Title:

Endosomal-lysosomal organellar assembly (ELYSA) structures coordinate lysosomal

degradation systems through mammalian oocyte-to-embryo transition

Authors:

Yuhkoh Satouh*[1], Takaki Tatebe[1], Isei Tanida[2], Junji Yamaguchi[2], Yasuo Uchiyama[2], Ken

Sato*[1]

Affiliations:

[1] Laboratory of Molecular Traffic, Institute for Molecular and Cellular Regulation, Gunma

University, Maebashi, Japan.

[2] Department of Cellular and Molecular Neuropathology, Juntendo University Graduate

School of Medicine, Tokyo, Japan.

*Correspondence to:

yuhkohs@gunma-u.ac.jp

sato-ken@gunma-u.ac.jp

Summary blurb

This study describes endosomal-lysosomal organellar assembly structures in mammalian oocytes, elucidating statistical alterations in their size, distribution, and correlation with lysosomal maturation.

Abstract

Mouse oocytes undergo drastic changes in organellar composition and their activities during maturation from the germinal vesicle (GV) to meiosis II (MII) stage. After fertilization, the embryo degrades parts of the maternal components via lysosomal degradation systems, including autophagy and endocytosis, as zygotic gene expression begins during embryogenesis. Here, we demonstrate that endosomal-lysosomal organelles form large spherical assembly structures, termed ELYSAs, in mouse oocytes. ELYSAs are observed in GV oocytes, attaining sizes up to 7–8 μm in diameter in MII oocytes. ELYSAs comprise tubular-vesicular structures containing endosomes and lysosomes along with cytosolic components. Most ELYSAs are also positive for an autophagy regulator, LC3. These characteristics of ELYSA resemble those of ELVA (endolysosomal vesicular assemblies) identified independently. The signals of V1-subunit of vacuolar ATPase tends to be detected on the periphery of ELYSAs in MII oocytes. After fertilization, the localization of the V1-subunit on endosomes and lysosomes increase as ELYSAs gradually disassemble at the 2-cell stage, leading to further acidification of endosomal-lysosomal organelles. These findings suggest that the ELYSA/ELVA maintain endosomal-lysosomal activity in a static state in oocytes for timely activation during early development.

Introduction

Fertilization triggers the dynamic remodeling of cellular components via zygotic gene expression as well as the degradation of cellular components from both maternal and paternal gametes[1]. After fertilization, maternal cytosolic proteins and RNAs are selectively degraded by the ubiquitin-proteasome system (UPS)[2] and specific RNA degradation systems[3, 4], respectively. Treatment with a proteasome inhibitor or knockdown of the zygote-specific proteasome assembly chaperone gene inhibits cytokinesis in zygotes, suggesting an essential role for the UPS in early development[5, 6]. Membrane components derived from each gamete are degraded by the lysosomal degradation system via autophagy and endocytosis[7, 8]. In mice, autophagic activity is relatively low in metaphase II (MII) oocytes, whereas it is rapidly upregulated approximately 4 h after fertilization, and LC3 (MAP1LC3: Microtubule-associated protein light chain 3)-positive structures appear throughout the cytoplasm. Autophagic activity decreases slightly before and after the 2-cell stage but remains high from the 4- to 8-cell stage[9]. Loss of Atg5 causes embryonic lethality at the 4- to 8-cell stages, suggesting that autophagy is essential for early mouse development. In contrast, some maternal plasma membrane (PM) proteins are internalized from the PM at the late 2-cell stage and are selectively degraded by lysosomes[10]. Endocytosis inhibitors prevent cell division in 2-cell embryos, suggesting that endocytosis plays a pivotal role in early development[10, 11]. Although lysosomal degradation systems are drastically upregulated after fertilization, their regulation remains unknown.

In *Caenorhabditis elegans*, the lysosomal activity in oocytes is upregulated by major sperm proteins secreted from sperms as the oocytes grow and mature[12]. In this process,

translational arrest of mRNA coding the V1-subunits of vacuolar ATPase in oocytes is released by the stimulation of sperm-derived factors. Consequently, the V1-subunits are recruited to the Vo-subunits in the lysosomal membrane, enhancing lysosomal acidification. This process is known as lysosomal switching[12]. In mouse embryos, LysoTracker staining analysis suggests that lysosomal acidification is prominent at the 2-cell stage but not in MII oocytes and zygotes[13], raising the possibility that lysosomal switching takes place during mouse embryogenesis.

In this study, we aimed to investigate the dynamics of the endolysosomal system during mouse oocyte maturation and embryogenesis. We found that endosomes and lysosomes assembled to form spherical structures in germinal vesicle (GV)-stage oocytes, which became larger as they matured into MII oocytes. Ultrastructural analysis showed that these structures were large spherical membrane assemblies, including endosomes, lysosomes, and tubulo-vesicular structures located in the periphery of the PM. We refer to this structure as the Endosomal-LYSosomal organellar Assembly (ELYSA). The ELYSA contains several cytosolic constituents. While ELYSAs are disassembled in late-stage zygotes, immature endosomes and lysosomes gradually appear in the cytoplasm of early 2-cell stage embryos. Live imaging analysis showed that the signal of V1-subunit of V-ATPase on the acidic compartment became stronger during embryogenesis, suggesting further acidification of lysosomes. These findings suggest that the ELYSA maintains low endosomal and lysosomal activities and acts as a waiting place for endolysosomal organelles to undergo fertilization-triggered activation toward embryogenesis.

Results

Endosomes and lysosomes coordinately form giant structures in MII oocytes

To examine the localization of endosomes and lysosomes in mouse MII oocytes, early or late endosomes were stained with anti-RAB5 or RAB7 antibody, respectively, and their localization was compared with that of lysosomes stained with anti-LAMP1 antibody. Giant structures harboring lysosomes with early and late endosomes were detected (**Fig. 1A, Supplementary Fig. 1, Video 1**). Although small punctate structures that are RAB5-positive but LAMP1-negative also spread over the cytosol, most giant structures were positive for RAB5 and LAMP1 (**Video 1**). Each giant structure in the MII oocytes was spherical, ellipsoidal, or concatenated with small spheres. These structures were located in the peripheral region just beneath the PM, excluding the metaphase plate region in proximity to the metaphase chromosomes. These structures were observed from the GV stage to the pronuclear-zygote (PN-zygote) stage but were mostly disassembled at the 2-cell stage. Dispersion and separation of RAB5 and LAMP1 signals after the first division indicated that these giant structures were transiently formed (**Fig. 1 B**). On the other hand, RAB5-positive and LAMP1-negative punctate structures tend to accumulate along with LAMP1-positive punctate structures larger than 1 μm at the late 2-cell stage.

Analyses of the number, size, and distribution of LAMP1-positive organelles were carried out by reconstituting three-dimensional objects using LAMP1-positive signals in immunocytochemically stained oocytes and embryos (**Supplementary Fig. 2**). Although the LAMP1 signals on the PM increased in the 2-cell stages, reconstitution of the objects was carried out only for the cytoplasmic fraction. The total number of the LAMP1-positive

objects was highest at the GV stage, and it decreased toward the late 2-cell stage, with a

significant difference compared to that in MII oocytes (**Fig. 2A**). Sorting of the objects based

on their sizes (indicated with the diameters when converting the volume of each object to the

corresponding sphere) revealed that the LAMP1-positive structures with diameters of 0.2–

0.8 μm, which were prominent in GV oocytes, significantly decreased toward the MII stage,

while the number of giant LAMP1-positive structures with diameters of 3–7 μm decreased

during the post-fertilization changes at the early and late 2-cell stage relative to that in MII

(**Fig. 2B**). The ratio of the number of the LAMP1-positive structures clearly indicated a

drastic increase in giant structures with 5–7 μm diameter during GV to MII transition (**Fig.**

2C). The transiently increased LAMP1-positive structures with 9–10 μm diameter at the

early 2-cell stage were observed as concatenated spheres that gathered upon cellular division;

concurrently, prominent signals on the PM appeared, especially at the cleavage furrow and

disappeared by the late 2-cell stage (**Fig. 2D, E**). The total volume of LAMP1-positive

objects was highest in MII, and it decreased over the early and late 2-cell stages (**Fig. 2F**).

LAMP1-positive structures with diameters of > 3 μm were also observed in GV oocytes, but

those larger than 5 μm in diameter were rarely detected, and the total volume of LAMP1-

positive structures larger than 3 μm in diameter in MII oocytes accounted for 51.3% of the

total volume. The large population of the LAMP1-positive structures with diameters larger

than 3 μm in MII transitioned to structures with diameters of 1–3 μm during the 2-cell stages,

as indicated by the percentage of the volume of each object size relative to the total volume

(**Fig. 2G**). The reconstitution of the objects above was conducted after the thresholding of

the fluorescent intensity. On the other hand, objects with diameters of 6–7 μm in MII oocytes

retained high intensity/volume, but those in GV oocytes or fertilized embryos revealed lower

intensities (**Supplementary Fig. 3**).

Furthermore, the analysis of object distribution for GV and MII oocytes indicated

that MII oocytes had a greater number of LAMP1-positive structures with > 4 μm diameter

in the cellular periphery, whereas GV oocytes had a greater number of these structures in the

cellular medial regions (**Fig. 2H, I**). These results suggest that the giant LAMP1-positive

structures are assembled and localized to the cellular periphery during oocyte maturation.

Endosomes, lysosomes, and tubulo-vesicular structures form an ELYSA

To clarify the internal structure of this giant structure containing lysosomes and

endosomes in the MII oocytes, we observed these in MII oocytes with in-resin correlative

light and electron microscopy (CLEM) using immunological reaction (immune in-resin

CLEM) with anti-RAB7 and -LAMP1 antibodies[14, 15]. RAB7- and LAMP1-signals were well

detected even in the ultrathin section (100 nm thickness) of the Epon-embedded MII oocyte,

fixed with the mixture of 4% paraformaldehyde and 0.025% glutaraldehyde (GA), which

showed similar distribution as that shown in **Fig. 1** and **Supplementary Fig. 4**. Toluidine

blue staining in adjacent ultrathin sections revealed that the RAB7/LAMP1-double positive

structures in the fluorescence observation were also positive for toluidine blue, with

interspaces often found in the center of spheres (**Fig. 3A**). Confocal fluorescent images of

the ultrathin sections clarified that RAB7-positive signals were frequently observed on the

periphery of these LAMP1-positive signals. Electron microscopic analysis of the

RAB7/LAMP1-double positive region in the ultrathin sections revealed that the

RAB7/LAMP1-double positive structures are assemblies of tubulo-vesicular structures, with

either vacuolated or filled interiors (**Fig. 3B**). The toluidine blue stain-positive structures

were confirmed to match in size and peripheral distribution with those identified via

RAB7/LAMP1 immunostaining, except for the metaphase plate, even under 2.5% GA

conditions for precise ultrastructural observation (**Fig. 3C, Supplementary Fig. 5, Video 2**).

Therefore, we examined the ultrastructural morphology of toluidine blue-positive giant

structures in MII oocytes fixed with 2.5% GA. The results showed that tubular membrane

vesicles with a short diameter of approximately 60.5 ± 15.9 nm assembled in the structure,

with gaps in the cytoplasm of approximately 24.0 ± 6.7 nm. Some of the vesicles also

harbored lysosome-like electron-dense contents, while others displayed multilamellar or

multivesicular body-like structures, and mitochondria appeared to be absent (**Fig. 3D**). Based

on these observations, we termed this membrane assembly the Endosomal-LYSosomal

organellar Assembly (ELYSA).

An autophagy regulator, LC3, also accumulated in ELYSA

We next examined whether autophagy-related proteins are included in ELYSAs. LC3 is a key

regulator of autophagy and often used as an autophagosome marker. It has been suggested

that the autophagic activity is decreased in MII oocytes since puncta formation of

transgenically-expressed a GFP fusion protein with LC3 disappeared at the MII stage[9]. Since

GFP fluorescence may be quenched under acidic conditions, we examined the localization of

LC3 in GV and MII oocytes by immunostaining with an anti-LC3 antibody. We found that

LC3 hardly formed puncta in the cytoplasm, but part of the LC3 signal overlapped with

ELYSA in both GV and MII oocytes (**Fig. 4A, B**), suggesting that LC3-positive membrane

structures were also present in the ELYSA. Further analysis of such double-positive

structures for the LC3 and LAMP1 antibodies indicated that their number decreased

drastically, while their total volume did not change during oocyte maturation

(**Supplementary Fig. 6A, B**). Analysis of double-positive structures indicated that their

average diameter significantly shifted from 2.36 ± 0.31 μm (GV) to 3.78 ± 0.20 μm (MII).

Moreover, LAMP1-positive structures smaller than 2 μm in diameter were rarely positive for

LC3 (**Supplementary Fig. 6C, D**). On the other hand, analysis of the distance from the

cellular geometric center indicated that they tended to be observed at the periphery of oocytes

in both stages (more than 80% in > 30 μm in the MII oocyte), and no clear tendency of double

positivity was observed along the distance (**Supplementary Fig. 6E, F**). A similar trend was

observed for RAB5 and LAMP1 double-positive structures, while the RAB5 signals tended

to overlap with LAMP1 signals more specifically at the cellular periphery than LC3, even in

the GV stage (**Supplementary Fig. 6G, H**).

ELYSA enlargement proceeds in an actin dependent manner

The localization of LAMP1-positive structures at the perinuclear (medial) region

before oocyte maturation and the specific localization at the cell periphery after maturation

(**Fig. 2H, I**) is similar to the redistribution of the endoplasmic reticulum (ER) and the

formation of cortical ER clusters. Fitzharris et al. showed that the inhibition of actin

cytoskeleton polymerization disturbs ER migration and cortical ER cluster formation[16]. Thus,

we examined the involvement of cytoskeletons in ELYSA formation by adding latrunculin A

(LatA) or cytochalasin B (CCB) as inhibitors of actin polymerization and nocodazole (Noco)

as an inhibitor of microtubule formation during *in vitro* maturation (IVM). While all the drugs

inhibited polar body extrusion and metaphase plate formation as reported previously[16], co-

immunostaining of LC3- or RAB5-antibody with LAMP1-antibody revealed their persistent

colocalization in ELYSAs (**Fig. 5A, Supplementary Fig. 7**).

Three-dimensional object analysis of the oocytes with or without cytoskeletal

inhibitor treatment indicated that the total number of LAMP1-positive structures was

increased only by CCB, and the total volume of the structures was unaffected

(**Supplementary Fig. 8**). We then classified LAMP1-positive structures by their diameters

and found that the number of structures with diameters of 0.8–2.0 μm increased when oocytes

were treated with LatA and CCB; very few structures with diameters of 4–7 μm that

corresponded to the enlarged ELYSAs were detected upon treatment with actin

polymerization inhibitors, not the tubulin inhibitor (**Fig. 5B**). Further examination of the

distance from the cell center of each LAMP1-positive structure with a diameter larger than 4

μm in IVM oocytes indicated that only the actin polymerization inhibitors reduced the

number of large structures at the cell periphery but not in the cellular medial region (**Fig. 5C,**

D).

V1 component of V-ATPase partially localizes to the periphery of ELYSAs in oocytes

We studied the relationship between ELYSA formation and endosomal/lysosomal

acidification in oocytes. V1A is a component of the V1 subunit of V-ATPase, which is

associated with the outer lysosomal membrane and drives endosomal/lysosomal acidification.

We used LAMP1-EGFP mRNA to observe ELYSA behaviors during oocyte maturation, with

ATP6V1A-mCherry mRNA (V1A-mCherry: a core component of V-ATPase fused with the

mCherry marker) to further examine the V1 subunit localization on ELYSAs. The mRNAs

were injected into GV oocytes, and the oocytes were applied to IVM.

Three-dimensional object analysis for the matured (MII) oocytes expressing

LAMP1-EGFP revealed a larger number of smaller LAMP1-positive structures (diameter of

0.2–0.4 μm) and a smaller number of structures with diameters of 0.6–1.0 μm, compared to

immunostaining (**Fig. 2**), but not significant difference in larger size (**Supplementary Fig.**

9). Live-cell imaging of LAMP1-EGFP fluorescence captured the dynamics of LAMP1-

positive organelles during oocyte maturation, notably demonstrating its stable expression

after germinal vesicle breakdown (GVBD, 2–3 h after initiation of IVM), and revealed that

the enlargement of ELYSAs larger than 5 μm in diameter occurred through the sequential

assembly of smaller ELYSAs that constantly migrated and contacted each other (**Fig. 6A,**

Video 3).

We then analyzed the localization of V1A-mCherry in the oocytes after IVM.

Although it was indistinguishable whether V1 subunit molecules or V1 subunit-positive

compartment were recruited to ELYSAs, V1A-mCherry signals were largely distributed on

the ELYSA surface and rarely observed within ELYSAs in deconvolved confocal images

(**Fig. 6B, Video 4**).

Acidification of lysosomes is facilitated as ELYSA disassembly proceeds in embryos

We examined the relationship between ELYSA disappearance and the emergence of

acidic membrane compartments during early development. LysoSensor probes exhibit

increased fluorescence in a pH-dependent manner upon acidification and enable the semi-

quantitative analysis of cellular acidic compartments, whereas LysoTracker probes exhibit

non-pH-dependent fluorescence. We first co-stained oocytes and embryos at various stages

with LysoTracker Red and LysoSensor Green (in the same medium drop). LysoTracker

showed a pronounced puncta-like signal after the 2-cell stage, as previously reported[13]. In

contrast, the ELYSAs in GV and MII oocytes were clearly identified by LysoSensor, whereas

LysoTracker only produced a very dim signal. Thus, the ratio of LysoTracker/LysoSensor

fluorescent intensities in the cytoplasm of each stage embryo indicated that ELYSAs were

present at a low ratio. In contrast, embryos at the 4-cell stage and beyond displayed small

punctate signals with a higher ratio (**Fig. 7**). A comparison between LysoTracker and V1A-

mCherry signals showed that these fluorescent signals colocalized well on small bright

vesicles but not on the ELYSA. Both LysoTracker and V1A-mCherry puncta were reduced

in embryos cultured in the presence of bafilomycin A_1, which inhibits the H^+ translocation of

V-ATPase (**Supplementary Fig. 10**). This suggests that LysoTracker detects membrane

compartment acidification. However, it may not sufficiently capture this process within the

ELYSA. In contrast, the results using LysoSensor suggest that lysosomes within the ELYSA

were not completely neutralized and remained acidic at a lower level, even though V1A was

difficult to localize.

When LAMP1-EGFP mRNA was injected into fertilized oocytes, it was difficult to

visualize the lysosomes due to the presence of a population that migrates onto the PM during

the 2-cell stage. Thus, to examine the spatiotemporal relationship between V1A localization

and lysosomal acidification during embryogenesis, we injected V1A-mCherry mRNA into

PN zygotes and cultured them in the presence of LysoSensor. It should be noted that stable

detection of LysoSensor and V1A-mCherry fluorescence required 4–8 h. In this process,

localization of V1A signal to lysosomes was rarely observed until ELYSA disappearance or

disassembly throughout the PN-zygote stage, but rapidly enhanced in the early 2-cell embryo

stage. LysoSensor-positive punctate structures, where V1A-mCherry signal colocalize,

increased in number during the transition from early to late 2-cell stage, indicating that

lysosomal acidification is promoted after ELYSA disappearance (**Fig. 8, Video 5**).

Low-level cathepsin-dependent proteolytic activity was maintained in ELYSA

Cathepsins are mainly responsible for the proteolytic activity in lysosomes. We

visualized the proteolytic activity of cathepsin B, the expression of which has been confirmed

in mouse embryos[13]. Mouse oocytes and embryos at various stages were simultaneously

collected and stained with Magic Red Cathepsin B staining solution to examine their

relationship with the ELYSA. Magic Red staining showed that proteolytic activity was

detected not only in the small, isolated membrane structures (mature lysosomes) but also in

the ELYSA in the GV and MII oocytes, whereas small punctate structures were

predominantly stained in other stages (**Fig. 9A, Supplementary Fig. 11A**). The intensity

plots of LysoSensor and Magic Red fluorescence revealed that the proteolytic activity

indicated by the Magic Red signal in the ELYSA was detected, but not as strong as that in

isolated punctate structures (lysosomes) in the GV and MII oocytes, and in developing

embryos (**Fig. 9B, Supplementary Fig. 11B, C**).

Discussion

In this study, we identified a large spherical structure, the ELYSA, consisting of endosomes, lysosomes, LC3-positive structures, and various vesicular-tubular structures in mouse oocytes. Small ELYSAs are present in GV oocytes and increase in size (3 to 7 μm in diameter) through assembly at the cell periphery in an actin-dependent manner during oocyte maturation and occupying more than half the volume of the LAMP1-positive fraction in MII oocytes. After fertilization, as ELYSAs gradually disassemble in zygotes, punctate structures positive for either RAB5- or LAMP1-signal alone appear in the cytoplasm of embryos at the early 2-cell stage. The number of membrane structures positive for V1A-mCherry increase upon ELYSA disassembly, indicating further acidification of the endosomal/lysosomal compartment. These results suggest that the ELYSA functions as a reservoir that accumulates endosomal and lysosomal compartments at the periphery of the cell cortex and maintains their activities in a static state until embryogenesis following fertilization.

We studied the dynamics and activity of endosomes and lysosomes in mouse oocytes and embryos and found that some oocyte PM proteins are internalized from the PM at the late 2-cell stage and are selectively degraded in lysosomes[10]. In this process, we noted that lysosomes were not well stained by LysoTracker until the early 2-cell stage, but their signals became stronger from the late 2-cell stage, suggesting the existence of a mechanism underlying developmentally regulated endosomal/lysosomal activation during embryogenesis[10, 13]. Notably, ELYSAs were stained with toluidine blue and LysoSensor, suggesting that these structures were acidified to some extent (**Fig. 3, 7**). In addition, Magic

Red, which reflects cathepsin B activity, weakly stained ELYSAs (**Fig. 9**). These

observations indicate that lysosomes in ELYSAs have some activity, albeit at low levels.

Mouse oocytes accumulate several maternal constituents, including proteins, lipids, and

mRNA, and grow to approximately 75 μm in diameter. In addition, oocytes must remain at

the GV and MII stages for an extended period to maintain their quality until fertilization. In

such situations, lysosomal degradation activity may be suppressed to a relatively low level

by the ELYSA, while nutrient uptake, storage, and protein synthesis are predominantly

upregulated in growing oocytes. Furthermore, we found that the V1-subunit signals were

largely detected on the periphery of ELYSAs but not in their interior in oocytes (**Fig. 6**). Thus,

it is possible that ELYSA formation prevents the localization of V1-subunits on the lysosomal

membrane inside the ELYSA.

Ultrastructural analysis revealed that the ELYSA consists of outer and inner

membrane assembly layers containing several endosomes and lysosomes, respectively (**Fig.**

1, 3). Considering the large number of RAB5-positive and LAMP1-negative punctate

structures in MII oocytes, these layers may also reflect the assembly mechanism of the

ELYSA. On the contrary, upon disassembly of the ELYSA, endosomes and lysosomes are

expected to be released from the ELYSA earlier and later, respectively. This may explain the

time lag between fertilization and lysosomal acidification beginning at the early 2-cell stage.

Since the ELYSA appears to contain cytosolic constituents, it may accommodate some

cytosolic constituents as a capsule to provide nutrition or signaling factors for development.

ELYSAs were distributed relatively widely in the GV oocyte, then specifically

located at the periphery of the oocyte cell cortex, except in the MII cell plate in the MII stage

(**Fig. 1, 2 Supplementary Fig. 6**). This change in distribution is reminiscent of that of the

ER and cortical ER clusters, whose distribution is regulated by the actomyosin system[16]. In

fact, the treatment with actin cytoskeleton polymerization inhibitors, but not a tubulin

polymerization inhibitor, prevented the redistribution of ELYSAs from the cellular medial

region to the cell periphery in oocytes (**Fig. 5**). The former treatment increased the number

of small LAMP1-positive organelles but reduced the size of ELYSAs in oocytes (**Fig. 5**).

These results suggest that actin cytoskeleton is involved in the assembly of smaller (< 2 μm

in diameter) LAMP1-positive organelles for the enlargement of ELYSAs in the cell periphery.

Our detailed imaging analysis showed that lysosomal acidification is promoted

simultaneously with the detection of the V1-subunits on lysosomes during ELYSA

disassembly in late zygotes and is further facilitated from the 2- to 4-cell stage and is

enhanced after the morula stage (**Fig. 8, Supplementary Fig. 10**). These findings suggest

that lysosomal acidification and maturation are linked to the ELYSA disassembly. Notably,

minor and major zygotic gene activation in mouse embryos occur at the 1- and 2-cell stages,

respectively, implying that lysosomal degradation of maternal components is coupled with

zygotic gene expression to promote embryogenesis[17]. Fertilization appears to trigger ELYSA

disassembly, but the downstream signaling pathways remain to be identified and warrant

further research.

Very recently, after sharing this ELYSA paper as a preprint in 2023[18], Zaffagnini et

al. reported the endolysosomal vesicular assemblies (ELVAs), which resemble ELYSAs, in

mouse oocytes[19]. The ELVA is defined as super-organelles containing endosomes, lysosomes,

autophagosomes, and proteasomes along with cytosolic proteins, which sequesters

detrimental protein aggregates inside and degrade them upon oocyte maturation[19].

Consistencies between the ELYSA and the ELVA in their contents and actin-dependent

dynamics during oocyte maturation/embryogenesis strongly suggest that ELYSAs and

ELVAs are identical structures[19], assuring the solidity of these findings in this paper.

Zaffagnini et al. also observed the accumulation of proteasomes and ubiquitinated proteins

in ELVAs and demonstrated that the depletion of a RUN and FYVE domain-containing

protein 1 (RUFY1) protein from oocytes by the Trim-Away method causes disappearance of

ELVAs, suggesting that RUFY1 is involved in the ELVA assembly. Analysis of *Rufy1*-

knockout ELVA/ELYSA free oocytes is awaited to elucidate the physiological role of ELVAs

or probably ELYSAs. They also suggested that ELVAs sequester and degrade specific

proteins that is known to be aggregate and cause diseases, including proto-oncogene receptor

tyrosine kinase KIT, upon oocyte maturation[19]. Complementary to their findings, our study

revealed that the recruitment of the V1-subunit is limited on ELYSAs in oocytes but enhanced

on the isolated LAMP1-positive vesicles after late 2-cell stage. These findings support a

hypothesis that ELYSA has a multimodal function, keeping endosomal and lysosomal

compartments at a relatively static state and supplying these membrane compartments as a

reservoir for embryogenesis.

Inhibition of lysosomal acidification by bafilomycin A_1 treatment arrests

embryogenesis at the 4–8-cell stage, while inhibition of cathepsin activity using a mixture of

E64d and pepstatin A blocks embryogenesis at the 8-cell or morula stage[13]. In contrast, the

inhibition of endocytosis by Pitstop2 treatment causes arrest at the 2-cell stage[10]. These

differences may imply that at the 2-cell stage, the endocytic activity is more important than

the lysosomal degradation activity, which is essential at later stages. However, insights from

this study shed light on the correlation between increased endocytic/autophagic activity and

lysosomal maturation through ELYSA disassembly and the potential significance of minor

lysosomal maturation in the early phase of embryogenesis.

The relationship between such endocytic/autophagic activities in early-stage

embryos and the embryonic developmental potential has been recently studied. Using GFP-

LC3 as an indicator of autophagic activity, Tsukamoto et al. found that aged oocytes from

14- to 15-month-old mice had lower autophagic activity than those from younger mice,

probably due to decreased lysosomal activity. Females transplanted with embryos with high

autophagic activity at the 4-cell stage have a significantly higher litter size ratio than that of

control females[20]. Analysis of ELYSA formation and disassembly in aged oocytes and

embryos may be a useful tool for determining oocyte quality by examining appropriate

regulation of endosomal/lysosomal activities.

Methods

Animal experiments and strains

Neither randomization nor blinding was used for animal selection (8–16-week-old). All

experimental protocols involving animals were approved and performed in accordance with

the guidelines of the Animal Care and Experimentation Committee of Gunma University

(Approval No. 22-076). Hybrid B6D2F1 mice were purchased from the Japan SLC

(Hamamatsu, Japan)

Antibodies and inhibitors

The following primary antibodies were used: rabbit monoclonal anti-RAB5 (C8B1; 3547; Cell Signaling Technology, Danvers, MA, USA), rabbit monoclonal anti-RAB7 (D95F2; 9367; Cell Signaling Technology), rat monoclonal anti-mouse LAMP1 (1D4B; sc-19992; Santa Cruz Biotechnology, Dallas, TX, USA), and mouse monoclonal anti-rat LC3 (Clones 4E12; M152-3; Medical and Biological Laboratories, Tokyo, Japan). The following secondary antibodies were used: goat anti-rabbit IgG (H+L) conjugated with Alexa Fluor 555 (A-21428; Thermo Fisher Scientific, Waltham, MA, USA), goat anti-rat IgG (H+L) cross-adsorbed secondary antibody conjugated with Alexa Fluor 488 and 647 (A-11006 and A-21247, respectively; Thermo Fisher Scientific), and goat anti-mouse IgG (H+L) cross-adsorbed secondary antibody conjugated with Alexa Fluor 546 (A-11030; Thermo Fisher Scientific). The following inhibitors or solutions were used for IVM or embryonic culture assays: dimethyl sulfoxide (DMSO) (13406-55; Nacalai Tesque, Kyoto, Japan); latrunculin A actin polymerization inhibitor that binds to monomer actin (428021; Sigma-Aldrich, St. Louis, MO, USA); cytochalasin B, an actin polymerization inhibitor that binds to the plus-end of polymerized actin (C6762; Sigma-Aldrich); and nocodazole, a tubulin polymerization inhibitor that binds to β-tubulin (M1404; Sigma-Aldrich). Stock solutions of latrunculin A, cytochalasin B, or nocodazole were prepared by solubilizing each in DMSO at 200× concentration for the assays. Bafilomycin A_1 was dissolved in DMSO (L266; Dojindo, Kumamoto, Japan) at a final concentration of 25 nM.

Oocyte preparation

GV oocytes were collected from the ovaries of female mice 46 h after injection with pregnant

mare serum gonadotropin or an anti-inhibin antibody (CARD HyperOva; Kyudo, Kumamoto,

Japan)[21]. Antral follicles were suspended in 30 µL of FHM medium drops (Sigma-Aldrich)

covered with liquid paraffin (Nacalai Tesque), punctured using 26G needles (Terumo, Tokyo,

Japan), and the GV oocytes were collected using a glass needle with a mouth pipette. MII

oocytes were collected from the oviducts of female mice 13 h after injection with CARD

HyperOva and 7.5 IU human chorionic gonadotropin (hCG; ASKA Pharmaceutical, Tokyo,

Japan) 48 h apart. The oviductal wall was punctured using a 26G needle to collect cumulus-

oocyte complexes in 20 µL potassium simplex optimization medium (KSOM; Kyudo) drops

covered with liquid paraffin.

DNA vector construction and mRNA synthesis

Complementary DNA for mouse *Lamp1* (accession number NM_001317353) or *Atp6v1a*

(accession number NM_001358204) was amplified using PCR with High-fidelity Taq

Polymerase, KOD FX Neo (Toyobo, Osaka, Japan), and a mouse ovary cDNA library as a

template (Genostaff, Tokyo, Japan). The amplicon was then subcloned into the entry vector

pDONR221 and transferred into the destination vectors pDEST-CMV-C-EGFP[22] (a gift from

Robin Ketteler (Addgene plasmid # 122844; http://n2t.net/addgene:122844; RRID:

Addgene_122844); for *Lamp1*) or pDest-mCherry-N1[23] (a gift from Robin Shaw (Addgene

plasmid # 31907; http://n2t.net/addgene:31907; RRID: Addgene_31907); for *Atp6v1a*) using

Gateway recombination cloning technology. The following primer combinations were used

for   cloning: 5′-

GCGCACAAGTTTGTACAAAAAAGCAGGCTGCCACCATGGCGGCCCCCGGCG-3′

and   5′-

GCGCACCACTTTGTACAAGAAAGCTGGGTTGATGGTCTGATAGCCGGCGTGACT

-3′   (for   *Lamp1*),   or   5′-

GCGCACAAGTTTGTACAAAAAAGCAGGCTGCCACCATGGATTTCTCCAAGCTAC

CC-3′   and   5′-

GCGCACCACTTTGTACAAGAAAGCTGGGTTGTCTTCAAGGCTACGGAATG-3′ (for

Atp6v1a). The cDNA encoding *Lamp1-EGFP* or *Atp6v1a-mCherry* were amplified using the

following primer combinations: 5′-CGCAAATGGGCGGTAGGCGTG-3′ and 5′-

TCCAGCAGGACCATGTGATCGC-3′   (for   *Lamp1-EGFP*),   or   5′-

CGCAAATGGGCGGTAGGCGTG-3′ and 5′-TTGGTCACCTTCAGCTTGG-3′ (for

Atp6v1a-mCherry). RNA synthesis using each PCR amplicon as a template, poly-A tail

addition, and mRNA purification were performed using an mMESSAGE mMACHINE T7

Ultra kit (Thermo Fisher Scientific) and MEGAclear transcription clean-up kit (Thermo

Fisher Scientific), according to the manufacturer's protocols. Synthesized mRNAs were

dissolved in RNase-free demineralized water at a concentration of 400 ng/μL and frozen in

aliquots until use.

In vitro maturation, fertilization, and embryonic culture

IVM for the GV oocytes was conducted by washing the oocytes twice and culturing in alpha-

modified Eagle's minimum essential medium (Thermo Fisher Scientific) containing 5%

(vol/vol) fetal bovine serum (FBS), 0.1 IU/mL follicle-stimulating hormone (MSD, Tokyo,

Japan), 1.2 IU/mL of hCG (ASKA Pharmaceutical), and 4 ng/mL of epidermal growth factor

(Thermo Fisher Scientific) for 17 h at 37 °C under a 5% CO_2 atmosphere[24]. For *in vitro*

fertilization, spermatozoa from a B6D2F1 epididymis were pre-incubated in a 100-μL

modified human tubal fluid medium (Kyudo) drop covered with liquid paraffin and

inseminated at a sperm concentration of 1.5×10^5 per mL for 1.5 h. Excess spermatozoa were

washed out, and the oocytes were incubated in a 20-μL KSOM drop (Kyudo) for further

development. IVM assays using latrunculin A and cytochalasin B were performed by

transferring oocytes to medium drops containing each inhibitor or DMSO 4 h after the start

of IVM (waiting for the completion of GVBD), as previously reported[16]. IVM media

containing DMSO, latrunculin A, or cytochalasin B was prepared by diluting the stock

solution (1:400), and the final concentration of the inhibitors was 10 μM. The embryonic

culture assay using bafilomycin A_1 was conducted by transferring the PN-zygote into a 20-

μL drop of KSOM containing 25 nM bafilomycin A_1 covered with liquid paraffin at 6 h post-

fertilization (hpf). Embryos were transferred at 24, 48, and 72 hpf to freshly prepared KSOM

containing bafilomycin A_1.

Immunostaining

Immunostaining of oocytes or embryos was performed as previously described[10]. Oocytes or

embryos were cultured to the indicated stages and fixed in 4% paraformaldehyde (PFA;

Nacalai Tesque) in phosphate-buffered saline (PBS) containing 0.1 mg/mL polyvinyl alcohol

(PVA; Sigma-Aldrich; PVA/PBS) for 15 min at 25–27 °C. After fixation, the cells were

permeabilized with PVA/PBS containing 0.1% Triton X-100 (Sigma-Aldrich) for 30 min at

25–27 °C and washed three times with PVA/PBS. After blocking with 10% FBS (Thermo

Fisher Scientific) in PBS for 1 h at room temperature, the cells were incubated with the

aforementioned primary antibodies (1:100 for all) overnight at 4 °C. Next, the cells were

washed three times in PVA/PBS and incubated with anti-mouse, rat, or rabbit IgG secondary

antibodies coupled with Alexa Fluor 488, 555, or 564 (1:200 for all; Thermo Fisher

Scientific) at 25–27 °C for 2–4 h. Thereafter, the cells were incubated with PVA/PBS

containing 10 μg/mL Hoechst 33342 (Dojindo) for 10 min and washed with fresh PVA/PBS

a few times. Subsequently, control and nonspecific staining were performed by processing

oocytes and embryos in the absence of primary antibodies. The oocytes and embryos were

imaged in PVA/PBS microdrops in 35-mm glass-bottomed dishes (MatTek Corporation,

Ashland, MA, USA). For each position, confocal images at 81 *z*-axis planes with 1-μm

increments were acquired on an IX-71 inverted microscope (Olympus Corporation, Tokyo,

Japan) equipped with a CSU W-1 confocal scanner unit (Yokogawa Electric Corporation)

using a 60× silicone immersion lens UPLSAPO60XS (Olympus Corporation).

LysoTracker/LysoSensor staining of oocytes and generation of ratio images

Acidic organelles in the oocytes were visualized using LysoTracker Red DND-99 and

LysoSensor Green DND-189 (Thermo Fisher Scientific) according to the manufacturer's

instructions. Briefly, the oocytes or embryos at target stages for one experimental round were

obtained simultaneously and incubated in a large drop of reaction mixture (KSOM containing

1/10,000 Magic Red stock solution dissolved in DMSO and 1/1,000 LysoSensor Green DND-

189 stock solution dissolved in DMSO) for 30 min at 37 °C under 5% CO_2 in a humidified

atmosphere in liquid paraffin in a 35-mm glass-bottom dish (MatTek Corporation).

Subsequently, confocal images at 41 z-axis planes with 2-μm increments were obtained using

a CSU W-1 spinning disk confocal microscope with a 30× silicone immersion lens

UPLSAPO30XS (Olympus Corporation). Acquisition conditions for fluorescence excited by

470-nm (LysoSensor) and 555-nm (LysoTracker) lasers were determined in preliminary

experiments to be almost 1:1 intensities between the two wavelengths for cytosolic punctate

structures in the 2-cell embryo. Further generation of ratio images for

LysoTracker/LysoSensor fluorescence were carried out using Fiji software[25]. Averaged

intensity projection was carried out, determining background intensity of each field by

subtracting the averaged intensities of four cellular blank regions from the whole image. Then,

32-bit ratio images for LysoTracker/LysoSensor fluorescence of cytosolic regions were

generated by processing images with "Math/Divide" function in Fiji, and "fire" look up table

images were converted to RGB images.

Magic Red/LysoSensor staining

Intracellular cathepsin activity was assayed in oocytes or embryos using Magic Red

Cathepsin B or Cathepsin L detection kits (ImmunoChemistry Technologies LLC, Davies,

CA, USA) according to the manufacturer's instructions. Briefly, the oocytes or embryos at

target stages for one experimental round were obtained simultaneously and incubated in a

large drop of reaction mixture (KSOM containing 1/250 Magic Red stock solution dissolved

in DMSO and 1/1,000 LysoSensor Green DND-189 stock solution dissolved in DMSO) for

25 min at 37 °C under 5% CO_2 in a humidified atmosphere in liquid paraffin in a 35-mm

glass-bottom dish (MatTek Corporation). Thereafter, Hoechst 33342 (10 μg/mL) was added

to the drop to a final concentration of 10 μg/mL, and incubated for 5 min. Subsequently,

oocytes or embryos at the same stage were collected, and confocal images at 41 z-axis planes

with 2-μm increments for both 470-nm (LysoSensor) and 555-nm (Magic Red) lasers were

obtained on a CSU W-1 spinning disk confocal microscopy using a 30× silicone immersion

lens UPLSAPO30XS (Olympus Corporation).

In-resin CLEM and electron microscopy

In-resin CLEM was performed as previously described with some modifications[14, 15]. Briefly,

oocytes were fixed in 4% PFA and 0.025% glutaraldehyde (GA). After permeabilization with

50 μg/mL digitonin (Nacalai Tesque) and 0.02% Triton X-100, oocytes were washed with

PVA/PBS three times and incubated with PBS containing 10% (w/v) FBS (Thermo Fisher

Scientific) for 1 h at room temperature. Thereafter, oocytes were incubated with PVA/PBS

containing primary antibody overnight at 4 °C. After incubation with a secondary antibody

in PVA/PBS for 2 h, oocytes were fixed in 4% PFA and 2.5% GA, embedded in 1.5% agarose,

stained with osmium tetroxide, dehydrated with graded ethanol solutions (QY1), and

embedded in epoxy resin (Oken Shoji, Tokyo, Japan). Ultrathin sections of Epon-embedded

oocytes (100-nm thickness) were prepared using a Leica UC6 ultramicrotome (Leica,

Nussloch, Germany) and placed on glass coverslips. Subsequently, fluorescence microscopy

images of the sections were obtained using a Nikon A1RHD25 confocal laser scanning

microscope with a NIS-Elements software (Nikon, Tokyo, Japan). Electron microscopic

images were observed using a Helios Nanolab 660 FIB-SEM instrument (FEI Company,

Hillsboro, USA).

For conventional electron microscopy, the oocytes were fixed in the presence of 2% PFA and

2.5% GA and stained with osmium tetroxide. After dehydration with ethanol, the oocytes

were embedded in epoxy resin, as described above. Sections were cut at a thickness of 100

nm using a UC6 ultramicrotome, mounted on glass coverslips, stained with uranyl acetate

and lead citrate, and observed under a Regulus8240 scanning electron microscope (Hitachi

High-Tech Corporation, Tokyo, Japan).

Serial semi-thin sections and 3D reconstruction of toluidine blue-positive structures

Serial semi-thin sections were cut at a thickness of 300 nm using a UC6 ultramicrotome and

mounted on glass slides. Next, the sections were stained with 1% toluidine blue/0.1 M PB,

and images were obtained using a BX50 light microscope (Olympus Corporation). Serial

toluidine blue-stained images were aligned using FIJI (National Institutes of Health, MD,

USA) with plugin-linear stack alignment using SIFT (LSAS), and 3D reconstruction was

performed using Amira software (Thermo Fisher Scientific).

Microinjection into the oocytes

Microinjection of mRNA into GV oocytes was performed as previously described[26,27]. Briefly,

antral follicles were harvested in FHM medium containing 250 μM of dbcAMP using 26G

needles (Terumo). After dissociating the cumulus cells by pipetting, the oocytes were

transferred to medium supplemented with 250 μM of dbcAMP and 10 μM of cytochalasin B

(Sigma-Aldrich). Thereafter, mRNA injection into the oocytes was performed using a piezo-

micromanipulator (Primetech, Tokyo, Japan) with a glass capillary needle. The mRNA

concentrations were 50 and 100 ng/µL for *V1A-mCherry* and *Lamp1-EGFP*, respectively.

The oocytes were washed three times and used for IVM. Microinjection of mRNA into zygote

stage embryo (at 3–4 hpf) was carried out as previously described[10]. The concentration of

mRNA was the same as that in the GV oocytes.

Time-lapse confocal imaging of oocytes/embryos

Low-invasive confocal fluorescence imaging of oocytes/embryos was performed as

previously described[28, 29, 30] with slight modifications. Briefly, on an inverted microscope IX-

71 (Olympus Corporation) equipped with a CSU W-1 confocal scanner unit (Yokogawa

Electric Corporation), an incubation chamber (Tokai Hit, Fujinomiya, Japan) was set at 37 °C

on the microscope stage, with a gas mixture of 5% CO_2 introduced into the chamber at 150

mL/min. Oocytes/embryos were placed in 8 µL of IVM medium or KSOM drops covered

with liquid paraffin (Nacalai Tesque) in 35-mm glass-bottomed dishes (MatTek Corporation).

Subsequently, confocal images at 41 *z*-axis planes with 2-µm increments for excitation lasers

of 470- and 555-nm from an LDI-7 Laser Diode Illuminator (89North, Williston, VT, USA)

were obtained every 30 or 60 min during IVM or embryonic culture observation, respectively,

through optimized band-pass filters using 30× silicone immersion lens UPLSAPO30XS

(Olympus Corporation).

Three-dimensional object analysis

Series of confocal images with 81 z-axis planes with 1-µm increments for LAMP1

immunostaining were used for 3D object analysis. Fields to analyze were cropped for each

oocyte/embryo, and the upper and lower end of LAMP1 fluorescent signals in the cytosol

were examined to determine the center plane (middle height) of the cell. The cellular region

in the center plane was used for masking and determination of the geometric center (as a 3D

center) of each oocyte. Deletion of the signals from the polar body or zona pellucida was

carried out manually. Objects were profiled using the 3D object counter plugin[31] of Fiji

software[25], to measure the number, volume, geometric center of each object, and mean

distance from the geometrical center of the object to its surface. Thresholding was carried

out after background subtraction using the Fiji software, manually applying intensities of

600–1200 as threshold to recognize objects separately. Signals on PM were eliminated both

via size filtering (0–5000 μm^3) of the 3D object counter plugin and manual erasing. The distal

distance of each object from the 3D center of the oocyte was used for distribution analysis to

avoid underestimation of the distance of large ELYSA. This was calculated by adding the

distance from the 3D center of the oocyte to the geometric center of each object and the mean

distance from the geometrical center of the object to its surface. Distance calculation and

generation of bar graphs were carried out using Excel 2016 software (Microsoft, Redmond,

WA, USA). Object-based colocalization analysis for the punctate structures were performed

using the JACoP plugin in the ImageJ Fiji software[31]. Confocal z-stack images of multiple

wave length for LAMP1 and LC3 or RAB5 antibodies were acquired for each oocyte as

above, and object-based colocalization was examined by identifying coincidence of

geometric centers of LC3 or RAB5 objects overlapping on LAMP1 objects. All of the

LAMP1-positive 3D objects were listed with the 3D object counter plugin applied with same

threshold, and the colocalized LAMP1-positive objects were identified and analyzed using

the Excel 2016 software.

Analysis of microscopic images and movies

Analysis of the acquired images, including the generation of a line plot of intensity,

adjustment of intensity, and generation of montage images at a uniform intensity, was

performed using Fiji software[25]. To enhance the images, a Gaussian Blur filter (at a sigma

radius of 1) and auto-adjustment of the intensity were applied to the stack of images. To

adjust the intensity, auto-adjustment of the intensity was applied to each image. The movies

were concatenated using Premiere Pro software (Adobe, San Jose, CA, USA).

Statistics and reproducibility

Sample sizes were estimated based on previous studies using similar experiments, and results

from preliminary experiments were examined to ensure statistical power more than 0.8 using

statistical power analysis in the G*Power software[32]. Statistical analyses and generation of

dot plot graphs were performed using GraphPad Prism 8 (GraphPad Software, Boston, MA,

USA). The significance of difference among groups were compared using one-way ANOVA

and Tukey's multiple comparison test. The experimenters were not blinded because of a

limitation in the availability of experienced personnel.

Data availability:

The data supporting the findings of this study are available from the corresponding authors

upon request.

Materials availability statement:

The materials supporting the findings of this manuscript are available from the corresponding

authors upon reasonable request.

Acknowledgments:

We would like to thank Editage (www.editage.jp) for English language editing.

Funding:

This study was supported by the Japan Society for the Promotion of Science KAKENHI

(Grant Numbers 19H05711, 20H00466, 24H02031 to K. Sato, 21H02435, 22H02872, and

22H04652 to I. Tanida, and 20K22744 to J. Yamaguchi), Takeda Science Foundation to K.

Sato (Grant Number 2024095741), the Mochida Memorial Foundation for Medical and

Pharmaceutical Research, Takeda Science Foundation (Grant Number 2022093867), and The

Cell Science Research Foundation to Y. Satouh. This work was partly supported by the

Project for Elucidating and Controlling Mechanisms of Aging and Longevity from the Japan

Agency for Medical Research and Development (AMED 21gm5010003 to Y. Uchiyama.,

22gm1710001 s0201 and 23gm1710001s0202 to I. Tanida) and by the MEXT-supported

Program for the Strategic Research Foundation at Private Universities (to Y. Uchiyama), and

the Center of Genomic and Regeneration Medicine, Juntendo University Graduate School of

Medicine (to Y. Uchiyama and I. Tanida).This work was partly supported by the joint

research program of the Institute for Molecular and Cellular Regulation, Gunma University

(24014 to I. Tanida and K. Sato).

Competing interests:

No competing interests declared.

Author contributions:

Y.S. and K.S. established the essential research concepts for this study. Y.S. and T.T.

performed immunostaining and assays using mouse oocytes/embryos. I. T., J. Y., and Y. U.

performed the CLEM experiments and analyses. Y. S. and K. S. prepared the manuscript.

References:

1.   Li L, Zheng P, Dean J. Maternal control of early mouse development. *Development*

137, 859-870 (2010).

2.   Verlhac MH, Terret ME, Pintard L. Control of the oocyte-to-embryo transition by

the ubiquitin-proteolytic system in mouse and *C. elegans*. *Curr Opin Cell Biol* **22**,

758-763 (2010).

3.   Jiang ZY, Fan HY. Five questions toward mRNA degradation in oocytes and

preimplantation embryos: when, who, to whom, how, and why? *Biol Reprod* **107**,

62-75 (2022).

4.   Schier AF. The maternal-zygotic transition: death and birth of RNAs. *Science* **316**,

406-407 (2007).

5.   Shin SW*, et al.* Mouse zygote-specific proteasome assembly chaperone important

for maternal-to-zygotic transition. *Biol Open* **2**, 170-182 (2013).

6.   Shin SW*, et al.* Inhibition of the ubiquitin-proteasome system leads to delay of the

onset of ZGA gene expression. *J Reprod Dev* **56**, 655-663 (2010).

7.   Sato K. Multiple roles of endocytosis and autophagy in intracellular remodeling

during oocyte-to-embryo transition. *Proc Jpn Acad Ser B Phys Biol Sci* **98**, 207-221

(2022).

8.   Birgisdottir AB, Johansen T. Autophagy and endocytosis - interconnections and

interdependencies. *J Cell Sci* **133**, (2020).

9.   Tsukamoto S, Kuma A, Murakami M, Kishi C, Yamamoto A, Mizushima N.

Autophagy is essential for preimplantation development of mouse embryos. *Science*

321, 117-120 (2008).

10.   Morita A, Satouh Y, Kosako H, Kobayashi H, Iwase A, Sato K. Clathrin-mediated

endocytosis is essential for the selective degradation of maternal membrane proteins

and preimplantation development. *Development* **148**, dev199461 (2021).

11.   Satouh Y, Sato K. Reorganization, specialization, and degradation of oocyte

maternal components for early development. *Reprod Med Biol* **22**, e12505 (2023).

12.   Bohnert KA, Kenyon C. A lysosomal switch triggers proteostasis renewal in the

immortal *C. elegans* germ lineage. *Nature* **551**, 629-633 (2017).

13.   Tsukamoto S*, et al.* Functional analysis of lysosomes during mouse preimplantation

embryo development. *J Reprod Dev* **59**, 33-39 (2013).

14.   Tanida I, Yamaguchi J, Suzuki C, Kakuta S, Uchiyama Y. Application of immuno-

and affinity labeling with fluorescent dyes to in-resin CLEM of Epon-embedded

cells. *Heliyon* **9**, e17394 (2023).

15.   Mitsui S, Yamaguchi J, Suzuki C, Uchiyama Y, Tanida I. TUNEL-positive structures

in activated microglia and SQSTM1/p62-positive structures in activated astrocytes

in the neurodegenerative brain of a CLN10 mouse model. *Glia* **71**, 2753-2769

(2023).

16.   FitzHarris G, Marangos P, Carroll J. Changes in endoplasmic reticulum structure

during mouse oocyte maturation are controlled by the cytoskeleton and cytoplasmic

dynein. *Dev Biol* **305**, 133-144 (2007).

17.   Schulz KN, Harrison MM. Mechanisms regulating zygotic genome activation. *Nat*

Rev Genet **20**, 221-234 (2019).

18.   Satouh Y, *et al.* Endosomal-lysosomal organellar assembly (ELYSA) structures

coordinate lysosomal degradation systems through mammalian oocyte-to-embryo

transition. [**Preprint**] *bioRxiv* 2023.12. 29.573616 (2023)

https://doi.org/10.1101/2023.12.29.573616

19.   Zaffagnini G*, et al.* Mouse oocytes sequester aggregated proteins in degradative

super-organelles. *Cell* **187**, 1109-1126 e1121 (2024).

20.   Tsukamoto S*, et al.* Fluorescence-based visualization of autophagic activity predicts

mouse embryo viability. *Sci Rep* **4**, 4533 (2014).

21.   Takeo T, Nakagata N. Superovulation using the combined administration of inhibin

antiserum and equine chorionic gonadotropin increases the number of ovulated

oocytes in C57BL/6 female mice. *PLoS One* **10**, e0128330 (2015).

22.   Agrotis A, Pengo N, Burden JJ, Ketteler R. Redundancy of human ATG4 protease

isoforms in autophagy and LC3/GABARAP processing revealed in cells.

Autophagy **15**, 976-997 (2019).

23.   Hong TT, *et al.* BIN1 localizes the L-type calcium channel to cardiac T-tubules.

PLoS Biol **8**, e1000312 (2010).

24.   Morohaku K, *et al.* Complete in vitro generation of fertile oocytes from mouse

primordial germ cells. *Proc Natl Acad Sci U S A* **113**, 9021-9026 (2016).

25.   Schindelin J, *et al.* Fiji: an open-source platform for biological-image analysis. *Nat*

Methods **9**, 676-682 (2012).

26.   Umeda R, *et al.* Structural insights into tetraspanin CD9 function. *Nat Commun* **11**,

1606 (2020).

27.   Kato K, *et al.* Structural and functional insights into IZUMO1 recognition by JUNO

in mammalian fertilization. *Nat Commun* **7**, 12198 (2016).

28.   Nozawa K, Satouh Y, Fujimoto T, Oji A, Ikawa M. Sperm-borne phospholipase C

zeta-1 ensures monospermic fertilization in mice. *Sci Rep* **8**, 1315 (2018).

29.   Satouh Y, Nozawa K, Yamagata K, Fujimoto T, Ikawa M. Viable offspring after

imaging of Ca^{2+} oscillations and visualization of the cortical reaction in mouse eggs.

Biol Reprod **96**, 563-575 (2017).

30.   Satouh Y, Inoue N, Ikawa M, Okabe M. Visualization of the moment of mouse

sperm-egg fusion and dynamic localization of IZUMO1. *J Cell Sci* **125**, 4985-4990

(2012).

31.   Bolte S, Cordelieres FP. A guided tour into subcellular colocalization analysis in

light microscopy. *J Microsc* **224**, 213-232 (2006).

32.   Faul F, Erdfelder E, Lang AG, Buchner A. G*Power 3: a flexible statistical power

analysis program for the social, behavioral, and biomedical sciences. *Behav Res*

Methods **39**, 175-191 (2007).

Figure legends

Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte

plasma membrane

(A) Meiosis II (MII) oocytes were fixed and co-stained with anti-RAB5 and anti-LAMP1

antibodies. The schematic diagram indicates the location of oocyte chromosomes and

metaphase plate in the MII oocytes. Arrowheads indicate the positions of oocyte

chromosomes. Magnified regions (right) are indicated by yellow boxes. (B) Embryos at

different stages were fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies.

Hour(s) post-fertilization is indicated as hpf. DNA was stained with Hoechst 33342.

Maximum intensity projection of confocal images at an axial scan range of 80 µm is shown

as Z-projection images.

Fig. 2. The giant structures are enlarged in the periphery of the MII oocyte PM during

oocyte maturation

Germinal vesicle (GV) oocyte, MII oocyte, early 2-cell (E2C: 24 hpf), and late 2-cell (L2C:

40 hpf) embryos were fixed and stained with anti-LAMP1 antibody. Reconstituted three-

dimensional objects for the LAMP1-positive organelles in 9 to 10 oocytes for each stage from

more than three independent experiments were further analyzed for number, size, and

distribution. (A) Averaged total number of LAMP1-positive objects per oocyte/embryo. (B)

Numbers of the LAMP1-positive objects sorted by size (indicated by the diameters of spheres

calculated from volumes). (C) Ratios of (B) for MII/GV to present changes during oocyte

maturation. (D) Ratios of (B) for E2C/MII or L2C/MII to present post-fertilization changes.

(E) A single confocal section image for E2C embryos, which have transient and concatenated

objects upon cellular division (arrowhead) and a prominent signal on the PM of cleavage

furrow (dotted circle). (F) Averaged total volume (μm^3) of the objects per oocyte/embryo. (F,

G) Total volumes of the objects sorted by size, or the percentage of each fraction relative to

the total LAMP1-positive object volume. (H, I) Numbers of large objects in GV or MII

oocytes at a distance of > 30 µm (cellular peripheral) or < 25 µm (medial) from the cellular

center. Groups with different letters are significantly different ($P < 0.05$, one-way ANOVA

and Tukey's multiple comparison test). Error bars indicate SEM. Dot plots show mean $\pm$

SEM and asterisks indicate significant difference between two groups.

Fig. 3. Endosomal-lysosomal organelles form assembly structures, ELYSA

(A) The internal structure of the giant structure in MII oocytes was analyzed by correlative

light and electron microscopy (CLEM) analysis using anti-RAB7 and anti-LAMP1

antibodies. A merged image of immunostaining and electron microscopy (EM) is indicated

(left), and a toluidine blue staining image of the adjacent thin section (right) is indicated.

Black arrowheads indicate the giant structures positive for RAB7/LAMP1/toluidine blue

staining. Note that RAB7/LAMP1 staining is indicated in a pseudo-color. (B) The fluorescent

distribution of RAB7/LAMP1 immunostaining was analyzed with deconvolution confocal

microscopy. Note that RAB7/LAMP1 staining is indicated in pseudo-color. (C) A series of

images of toluidine blue-stained serial semi-thin sections of the Epon-embedded MII oocyte

were reconstructed in 3D. Based on the images obtained, chromosomes were segmented blue,

toluidine blue-positive structures were segmented yellow, and the oocyte PM was segmented

white. (D) The internal structures of the toluidine blue-positive structure, ELYSA, were

observed using conventional EM of oocytes fixed with 2.5% glutaraldehyde (GA). Magnified

regions in a or b (right) are indicated by boxes (left). Arrows, arrowheads, or asterisks

indicate vesicles harboring high electron density contents and multilamellar or multivesicular

structures, respectively. ELYSA, endosomal-lysosomal organellar assembly.

Fig. 4. An autophagy regulator LC3 is detected in ELYSA

Germinal vesicle (GV) (A) and MII (B) oocytes were fixed and co-stained with anti-LC3 and

anti-LAMP1 antibodies. Magnified regions (right) are indicated by yellow boxes. DNA was

stained with Hoechst 33342. Arrowheads indicate the positions of oocyte chromosomes.

Maximum intensity projection of confocal images at an axial scan range of 80 µm is shown

as Z-projection images.

Fig. 5. Enlargement of ELYSA and its redistribution to the cell periphery occur in an

actin cytoskeleton-dependent manner

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without actin

polymerization inhibitors (10 µM latrunculin A or cytochalasin B) or a tubulin inhibitor (20

µM nocodazole), then fixed and co-stained with anti-LC3 and anti-LAMP1 antibodies. The

reconstituted objects for LAMP1-positive organelles in 10 oocytes for each treatment from

more than three independent experiments were analyzed for the number, size, and distribution.

(A) Maximum intensity projection of confocal images at an axial scan range of 80 µm is

shown as Z-projection images. DNA was stained with Hoechst 33342. Arrowheads indicate

the metaphase plate. (B) Average total LAMP1-positive object number per oocyte after

sorting the objects by size (indicated by the diameters of spheres calculated from volumes).

(C, D) Numbers of objects with diameters > 4 µm in oocytes at a distance of > 30 µm or <

25 µm from the cellular center. Groups with different letters are significantly different (P <

0.05, one-way ANOVA and Tukey's multiple comparison test). Error bars indicate SEM. Dot

plots are indicated with mean ± SEM.

Fig. 6. Assembly of ELYSA during migration and limited interaction with the

acidification machinery

(A) Snapshots of a time-lapse observation for EGFP fluorescence of GV oocytes injected

with *Lamp1-EGFP* and *V1A-mCherry* mRNA are indicated. Maximum intensity projection

of confocal images with a height of 40 μm (bottom half of the oocyte) are shown. Magnified

perinuclear regions (right) are indicated by yellow boxes. Arrowheads indicate assemblies

that adhere to each other in the subsequent frame. (B) Maximum intensity projection of

deconvolved confocal images at an axial scan range of 80 µm is shown as Z-projection

images. Magnified regions are indicated by yellow boxes (a, b) and the LAMP1-EGFP

organelle regions are indicated by yellow dotted line.

Fig. 7. Endosomes and lysosomes in the ELYSA are weakly acidified, but further

acidification occurs after the 4-cell stage

Oocytes and embryos at different stages were stained in the same drop for LysoSensor Green

and LysoTracker Red staining analysis, and confocal images were acquired. Averaged

intensity projection of confocal images at an axial scan range of 80 µm is shown as Z-

projection images. Magnified ratiometric image for LysoTracker/LysoSensor fluorescence of

cytosolic regions indicated by yellow boxes are presented as "fire" look up table images at

the bottom. Arrowheads indicate ELYSAs recognized with LysoSensor.

Fig. 8. ATP6V1A localization on acidic compartments increases as ELYSAs are

disassembled

Embryos injected with *V1A-mCherry* mRNA (at 3 hpf) were incubated to the blastocyst stage

in the presence of LysoSensor Green. Snapshots of a time-lapse observation of a

representative embryo are shown as maximum intensity projection of confocal images with

a height of 80 μm unless otherwise specified. Magnified perinuclear regions are indicated by

yellow boxes.

Fig. 9. Proteolytic activity of cathepsin B is maintained at a relatively low level in the

ELYSA

(A) GV and MII oocytes were stained in the same drop for LysoSensor Green and Magic Red

Cathepsin B staining analysis, and confocal images were acquired. (B) Fluorescent intensities

of LysoSensor and Magic Red on the yellow lines indicated in (A) are indicated by a line

plot. Small, filled dots indicate the origin (0 μm) in distance.

Supplementary Fig. 1. Endosomes and lysosomes form giant structures in the periphery

of the oocyte plasma membrane

MII oocytes were fixed and co-stained with anti-RAB7 and anti-LAMP1 antibodies. The

schematic diagram indicates the location of oocyte chromosomes and the metaphase plate in

the MII oocytes. Maximum intensity projection of confocal images at an axial scan range of

80 µm is shown as Z-projection images. Arrowheads indicate the positions of oocyte

chromosomes.

Supplementary Fig. 2. Schemes of the three-dimensional particle analysis in this study

Oocytes/embryos were fixed and stained, and three-dimensional images were acquired as

serial z-stack images using confocal microscopy. Fields to analyze were cropped for each

oocyte/embryo, and the upper and lower end of LAMP1 fluorescent signals in cytosol were

examined and used to determine the center plane of the cell. The cellular region in the center

plane was used for masking and determination of the geometric center (as a 3D center of the

oocyte). Deletion of the signals from the polar body or zona pellucida was carried out

manually. Signals on PM were eliminated by both size exclusion of 3D particle counter

plugin in Fiji software and by manual erasing. The distal distance of each particle from the

3D center of the oocyte was used for the distribution study to avoid underestimation of the

distance of large ELYSA particle, which was calculated by adding the 3D distance from the

center of cell to the geometric center of the particle and the mean diameter of the particle.

Supplementary Fig. 3. Large LAMP1-positive structures exhibit high fluorescent

intensity, especially in MII oocytes

Germinal vesicle (GV) oocytes, MII oocytes, early 2-cell (E2C: 24 hpf), and late 2-cell (L2C:

40 hpf) embryos were fixed and stained with anti-LAMP1 antibody. Reconstituted three-

dimensional objects for the LAMP1-positive objects in 9 to 10 oocytes for each stage from

more than three independent experiments were further analyzed for size and fluorescent

intensity. The sorting by size (indicated by the diameters of spheres calculated from volumes)

was carried out by thresholding the fluorescent intensity prior to measuring intensity. (A)

Total fluorescent intensity of LAMP1-positive objects with different sizes. (B)

Intensity/volume ratios of LAMP1-positive objects with different sizes. Groups with

different letters are significantly different ($P < 0.05$, one-way ANOVA and Tukey's multiple

comparison test). Error bars indicate SEM.

Supplementary Fig. 4. Giant structures in MII oocytes were immunostained with anti-

RAB7 and LAMP1 antibodies after glutaraldehyde (GA) fixation

MII oocytes were fixed with 0.025% GA and co-stained with anti-RAB5/RAB7 and anti-

LAMP1 antibodies. Single-section images from confocal observation are shown.

Supplementary Fig. 5. Toluidine blue-positive structures in MII oocytes show similar

localization patterns to the ELYSA

Aligned serial toluidine blue-stained light microscopic images were stacked and three-

dimensionally reconstructed by inversing the intensity and adjusting it to be transparent.

Supplementary Fig. 6. LC3 signals colocalize with LAMP1 signals in a size-dependent

manner

GV and MII oocytes stained with anti-LAMP1 antibody and anti-LC3 or anti-RAB5 antibody

were reconstituted three-dimensionally for their objects positive for the target proteins, and

3 to 4 oocytes for each stage were further analyzed for number, size, and distribution. (A, B)

Averaged total numbers or total volumes of the double-positive objects for LC3 and LAMP1

per oocyte/embryo, respectively. (C, D) Average numbers or percentage of total LAMP1-

positive objects of the double-positive objects for LC3 and LAMP1 after sorting of the

objects by size (indicated by the diameters of spheres calculated from volumes), respectively.

(E, F) The double-positive objects for LC3 and LAMP1 were sorted by distance from the

three-dimensional geometrical center of the oocyte. Average numbers or percentages of total

LAMP1-positive objects are indicated, respectively. (G, H) The double-positive objects for

RAB5 and LAMP1 were sorted by distance from the three-dimensional geometrical center

of the oocyte. Average numbers or percentages of total LAMP1-positive objects are indicated,

respectively. Significance was examined using a two-tailed Student's t-test, assuming equal

variances between groups; the significance is indicated with asterisks ($P < 0.05$). Error bars

indicate SEM.

Supplementary Fig. 7. An early endosome marker RAB5 is localized within the ELYSA

in the presence of inhibitors for cytoskeletons

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without inhibitors

of polymerized actin (10 μM latrunculin A or cytochalasin B) or tubulin (20 μM nocodazole),

and fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. Maximum intensity

projection of confocal images at an axial scan range of 80 µm is shown as Z-projection

images. Magnified regions are indicated by yellow boxes. DNA was stained with Hoechst

33342. Arrowhead and asterisk indicate the metaphase plate and first polar body, respectively.

Supplementary Fig. 8. Treatment with inhibitors for cytoskeltons affected the number

of LAMP1-positive objects but not their volume

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without actin

polymerization inhibitors (10 μM latrunculin A or cytochalasin B) or a tubulin inhibitor (20

μM nocodazole), then fixed and co-stained with anti-LC3 and anti-LAMP1 antibodies. The

reconstituted objects for LAMP1-positive objects in 10 oocytes for each treatment from more

than three independent experiments were analyzed for the number, size, and distribution. (A)

Average total object number per oocyte. (B) Total volumes of the objects. Groups with

different letters are significantly different (P < 0.05, one-way ANOVA and Tukey's multiple

comparison test). Error bars indicate SEM.

Supplementary Fig. 9. Number and distribution of large LAMP1-EGFP structures do

not significantly differ from intrinsic LAMP1 examined with immunofluorescence

After the *in vitro* maturation (IVM) for 18 h, LAMP1-EGFP fluorescence injected as mRNA

in the GV stage was compared with intrinsic LAMP1 detected with an anti-LAMP1 antibody.

The reconstituted objects for LAMP1-positive structures from 3 oocytes were compared with

those from immunofluorescent analysis in Figure 2 for the number and size. (A) Ratio of

average numbers of total LAMP1-positive to that of MII in Figure 2. (B) Ratios of numbers

of total LAMP1-positive structures sorted by size (indicated by the diameters of spheres

calculated from volumes). Significance was examined using a two-tailed Student's t-test,

assuming equal variances between groups; the significance is indicated with asterisks (P <

0.05). Error bars indicate SEM.

Supplementary Fig. 10. LysoTracker-positive punctate structures colocalized with V1A-

mCherry

Embryos were injected with V1A-mCherry mRNA (at 3 hpf) and cultured with or without

bafilomycin A_1 and stained with LysoTracker Green at the indicated time points. Maximum

intensity projection of confocal images at an axial scan range of 80 µm is shown as Z-

projection images.

Supplementary Fig. 11. Magic Red Cathepsin B staining in GV and MII oocytes showed

higher cytoplasmic signals but similar intensity in isolated punctate structures to

developing embryos

Oocytes and embryos at the indicated stages were collected and stained with Magic Red

Cathepsin B solution in the same drop of medium. (A) Averaged intensity projection of

confocal images at an axial scan range of 80 µm is shown as Z-projection images. (B) Early

2-cell and 4-cell embryos were stained in the same drop for LysoSensor Green and Magic

Red Cathepsin B staining analysis, and confocal images were acquired. (C) Fluorescent

intensities of LysoSensor and Magic Red on the yellow lines indicated in (B) are indicated

by a line plot. Small, filled dots indicate the origin (0 µm) in distance.

1000

1001 **Video 1.**

1002 MII oocytes were fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. A series

1003 of the confocal images with a height of 80 µm is shown from top to bottom.

1004

1005 **Video 2.**

1006 A series of continuous toluidine blue-stained light microscopic images of the same oocyte

1007 (top half) reconstructed to 3D. Chromosomes in the images obtained were segmented blue,

1008 toluidine blue-positive structures were segmented yellow, and oocyte PM was segmented

1009   white.

1010

Video 3.

GV oocytes were injected with the mixture mRNA solution for *Lamp1-EGFP* and *V1A-*

mCherry, and incubated for IVM. Time-lapse movie of EGFP fluorescence of a

representative oocyte is indicated as maximum intensity projection of confocal images with

a height of 80 μm. Magnified perinuclear regions in Fig. 5 are indicated by a box. The

timestamp indicates the time after IVM started.

1017

1018   **Video 4**

1019   GV oocytes were injected with V1A-mCherry mRNA and incubated for 18 h for IVM in the

1020   presence of LysoSensor green. Fluorescent images of a representative oocyte which matured

1021   to MII stage is presented. A series of the confocal images of 81 slices with a height of 80 μm

1022   is shown from top to bottom.

1023

1024   **Video 5.**

1025   Embryos were injected with V1A-mCherry mRNA at 3 hpf and incubated for further

1026   embryogenesis in the presence of LysoSensor Green during embryonic development. A time-

1027   lapse movie of a representative embryo is shown (maximum intensity projection of confocal

1028   images at a height of 80 μm).

1029

Figure 1

Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte plasma membrane
(A) Meiosis II (MII) oocytes were fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. The schematic diagram indicates the location of oocyte chromosomes and metaphase plate in the MII oocytes. Arrowheads indicate the positions of oocyte chromosomes. Magnified regions (right) are indicated by yellow boxes. (B) Embryos at different stages were fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. Hour(s) post-fertilization is indicated as hpf. DNA was stained with Hoechst 33342. Maximum intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images.

Figure 2

Fig. 2. The giant structures are enlarged in the periphery of the MII oocyte PM during oocyte maturation
Germinal vesicle (GV) oocyte, MII oocyte, early 2-cell (E2C: 24 hpf), and late 2-cell (L2C: 40 hpf) embryos were fixed and stained with anti-LAMP1 antibody. Reconstituted three-dimensional objects for the LAMP1-positive organelles in 9 to 10 oocytes for each stage from more than three independent experiments were further analyzed for number, size, and distribution. (A) Averaged total number of LAMP1-positive objects per oocyte/embryo. (B) Numbers of the LAMP1-positive objects sorted by size (indicated by the diameters of spheres calculated from volumes). (C) Ratios of (B) for MII/GV to present changes during oocyte maturation. (D) Ratios of (B) for E2C/MII or L2C/MII to present post-fertilization changes. (E) A single confocal section image for E2C embryos, which have transient and concatenated objects upon cellular division (arrowhead) and a prominent signal on the PM of cleavage furrow (dotted circle). (F) Averaged total volume (μm^3) of the objects per oocyte/embryo. (F, G) Total volumes of the objects sorted by size, or the percentage of each fraction relative to the total LAMP1-positive object volume. (H, I) Numbers of large objects in GV or MII oocytes at a distance of > 30 μm (cellular peripheral) or < 25 μm (medial) from the cellular center. Groups with different letters are significantly different ($P < 0.05$, one-way ANOVA and Tukey's multiple comparison test). Error bars indicate SEM. Dot plots show mean $\pm$ SEM and asterisks indicate significant difference between two groups.

Figure 3

A CLEM

RAB7
LAMP1

20 μm
0.025% GA

Toluidine blue
(adjacent section to CLEM)

B Merge   LAMP1   RAB7   EM

2 μm

C

D

2.5% GA

b

a

2 μm

a

200 nm

b

Fig. 3. Endosomal-lysosomal organelles form assembly structures, ELYSA

(A) The internal structure of the giant structure in MII oocytes was analyzed by correlative light and electron microscopy (CLEM) analysis using anti-RAB7 and anti-LAMP1 antibodies. A merged image of immunostaining and electron microscopy (EM) is indicated (left), and a toluidine blue staining image of the adjacent thin section (right) is indicated. Black arrowheads indicate the giant structures positive for RAB7/LAMP1/toluidine blue staining. Note that RAB7/LAMP1 staining is indicated in a pseudo-color. (B) The fluorescent distribution of RAB7/LAMP1 immunostaining was analyzed with deconvolution confocal microscopy. Note that RAB7/LAMP1 staining is indicated in pseudo-color. (C) A series of images of toluidine blue-stained serial semi-thin sections of the Epon-embedded MII oocyte were reconstructed in 3D. Based on the images obtained, chromosomes were segmented blue, toluidine blue-positive structures were segmented yellow, and the oocyte PM was segmented white. (D) The internal structures of the toluidine blue-positive structure, ELYSA, were observed using conventional EM of oocytes fixed with 2.5% glutaraldehyde (GA). Magnified regions in a or b (right) are indicated by boxes (left). Arrows, arrowheads, or asterisks indicate vesicles harboring high electron density contents and multilamellar or multivesicular structures, respectively. ELYSA, endosomal-lysosomal organellar assembly.

Figure 4

Fig. 4. An autophagy regulator LC3 is detected in ELYSA

Germinal vesicle (GV) (A) and MII (B) oocytes were fixed and co-stained with anti-LC3 and anti-LAMP1 antibodies. Magnified regions (right) are indicated by yellow boxes. DNA was stained with Hoechst 33342. Arrowheads indicate the positions of oocyte chromosomes. Maximum intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images.

Figure 5

Fig. 5. Enlargement of ELYSA and its redistribution to the cell periphery occur in an actin cytoskeleton-dependent manner

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without actin polymerization inhibitors (10 μM latrunculin A or cytochalasin B) or a tubulin inhibitor (20 μM nocodazole), then fixed and co-stained with anti-LC3 and anti-LAMP1 antibodies. The reconstituted objects for LAMP1-positive organelles in 10 oocytes for each treatment from more than three independent experiments were analyzed for the number, size, and distribution. (A) Maximum intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images. DNA was stained with Hoechst 33342. Arrowheads indicate the metaphase plate. (B) Average total LAMP1-positive object number per oocyte after sorting the objects by size (indicated by the diameters of spheres calculated from volumes). (C, D) Numbers of objects with diameters > 4 μm in oocytes at a distance of > 30 μm or < 25 μm from the cellular center. Groups with different letters are significantly different ($P < 0.05$, one-way ANOVA and Tukey's multiple comparison test). Error bars indicate SEM. Dot plots are indicated with mean $\pm$ SEM.

Figure 6

Fig. 6. Assembly of ELYSA during migration and limited interaction with the acidification machinery
(A) Snapshots of a time-lapse observation for EGFP fluorescence of GV oocytes injected with *Lamp1-EGFP* and *V1A-mCherry* mRNA are indicated. Maximum intensity projection of confocal images with a height of 40 µm (bottom half of the oocyte) are shown. Magnified perinuclear regions (right) are indicated by yellow boxes. Arrowheads indicate assemblies that adhere to each other in the subsequent frame. (B) Maximum intensity projection of deconvolved confocal images at an axial scan range of 80 µm is shown as Z-projection images. Magnified regions are indicated by yellow boxes (a, b) and the LAMP1-EGFP organelle regions are indicated by yellow dotted line.

Figure 7

Fig. 7. Endosomes and lysosomes in the ELYSA are weakly acidified, but further acidification occurs after the 4-cell stage

Oocytes and embryos at different stages were stained in the same drop for LysoSensor Green and LysoTracker Red staining analysis, and confocal images were acquired. Averaged intensity projection of confocal images at an axial scan range of 80 µm is shown as Z-projection images. Magnified ratiometric image for LysoTracker/LysoSensor fluorescence of cytosolic regions indicated by yellow boxes are presented as "fire" look up table images at the bottom. Arrowheads indicate ELYSAs recognized with LysoSensor.

Figure 8

Fig. 8. ATP6V1A localization on acidic compartments increases as ELYSAs are disassembled
Embryos injected with *V1A-mCherry* mRNA (at 3 hpf) were incubated to the blastocyst stage in the presence of LysoSensor Green. Snapshots of a time-lapse observation of a representative embryo are shown as maximum intensity projection of confocal images with a height of 80 μm unless otherwise specified. Magnified perinuclear regions are indicated by yellow boxes.

Figure 9

Fig. 9. Proteolytic activity of cathepsin B is maintained at a relatively low level in the ELYSA
(A) GV and MII oocytes were stained in the same drop for LysoSensor Green and Magic Red Cathepsin B staining analysis, and confocal images were acquired. (B) Fluorescent intensities of LysoSensor and Magic Red on the yellow lines indicated in (A) are indicated by a line plot. Small, filled dots indicate the origin (0 μm) in distance.

Supplementary Figure 1

Supplementary Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte plasma membrane

MII oocytes were fixed and co-stained with anti-RAB7 and anti-LAMP1 antibodies. The schematic diagram indicates the location of oocyte chromosomes and the metaphase plate in the MII oocytes. Maximum intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images. Arrowheads indicate the positions of oocyte chromosomes.

Supplementary Figure 2

Confocal planes (1μm pitch)

End of signal

End of signal

Center plane

Masking of cellular region and measurement of cellular geometric center (= three dimensional center of oocyte (XYZ))

- Background subtraction
- Erase signals from out side the oocyte/embryo

3D particle counter plugin (ImageJ: Fiji)
- Number of particles (after thresholding)
- Volume of each particle
- Geometric centers of each particle (XYZ)
- Mean distance from the geometrical center of each object to its surface

Analysis for volume and number count

+ analysis for distance (from center of cell)

1. 3D distance from center of oocyte to geometric center of each particle (calculated from XYZ positions)

2. Mean distance from particle geometric center of each object to its to surface

"Distance" in this study (=sum of 1 and 2): 3D distance of distal end of particle to the center of oocyte

Supplementary Fig. 2. Schemes of the three-dimensional particle analysis in this study
Oocytes/embryos were fixed and stained, and three-dimensional images were acquired as serial z-stack images using confocal microscopy. Fields to analyze were cropped for each oocyte/embryo, and the upper and lower end of LAMP1 fluorescent signals in cytosol were examined and used to determine the center plane of the cell. The cellular region in the center plane was used for masking and determination of the geometric center (as a 3D center of the oocyte). Deletion of the signals from the polar body or zona pellucida was carried out manually. Signals on PM were eliminated by both size exclusion of 3D particle counter plugin in Fiji software and by manual erasing. The distal distance of each particle from the 3D center of the oocyte was used for the distribution study to avoid underestimation of the distance of large ELYSA particle, which was calculated by adding the 3D distance from the center of cell to the geometric center of the particle and the mean diameter of the particle.

Supplementary Figure 3

A

B

Supplementary Fig. 3. Large LAMP1-positive structures exhibit high fluorescent intensity, specifically in MII oocytes

Germinal vesicle (GV) oocytes, MII oocytes, early 2-cell (E2C: 24 hpf), and late 2-cell (L2C: 40 hpf) embryos were fixed and stained with anti-LAMP1 antibody. Reconstituted three-dimensional objects for the LAMP1-positive objects in 9 to 10 oocytes for each stage from more than three independent experiments were further analyzed for size and fluorescent intensity. The sorting by size (indicated by the diameters of spheres calculated from volumes) was carried out by thresholding the fluorescent intensity prior to measuring intensity. (A) Total fluorescent intensity of LAMP1-positive objects with different sizes. (B) Intensity/volume ratios of LAMP1-positive objects with different sizes. Groups with different letters are significantly different (P < 0.05, one-way ANOVA and Tukey's multiple comparison test). Error bars indicate SEM.

Supplementary Figure 4

Supplementary Fig. 4. Giant structures in MII oocytes were immunostained with anti-RAB7 and LAMP1 antibodies after glutaraldehyde (GA) fixation
MII oocytes were fixed with 0.025% GA and co-stained with anti-RAB5/RAB7 and anti-LAMP1 antibodies. Single-section images from confocal observation are shown.

Supplementary Figure 5

Inversed (intact)	Inversed (transparent)

Supplementary Fig. 5. Toluidine blue-positive structures in MII oocytes show similar localization patterns to the ELYSA
Aligned serial toluidine blue-stained light microscopic images were stacked and three-dimensionally reconstructed by inversing the intensity and adjusting it to be transparent.

Supplementary Figure 6

Supplementary Fig. 6. LC3 signals colocalize with LAMP1 signals in a size-dependent manner

GV and MII oocytes stained with anti-LAMP1 antibody and anti-LC3 or anti-RAB5 antibody were reconstituted three-dimensionally for their objects positive for the target proteins, and 3 to 4 oocytes for each stage were further analyzed for number, size, and distribution. (A, B) Averaged total numbers or total volumes of the double-positive objects for LC3 and LAMP1 per oocyte/embryo, respectively. (C, D) Average numbers or percentage of total LAMP1-positive objects of the double-positive objects for LC3 and LAMP1 after sorting of the objects by size (indicated by the diameters of spheres calculated from volumes), respectively. (E, F) The double-positive objects for LC3 and LAMP1 were sorted by distance from the three-dimensional geometrical center of the oocyte. Average numbers or percentages of total LAMP1-positive objects are indicated, respectively. (G, H) The double-positive objects for RAB5 and LAMP1 were sorted by distance from the three-dimensional geometrical center of the oocyte. Average numbers or percentages of total LAMP1-positive objects are indicated, respectively. Significance was examined using a two-tailed Student's t-test, assuming equal variances between groups; the significance is indicated with asterisks ($P < 0.05$). Error bars indicate SEM.

Supplementary Figure 7

Supplementary Fig. 7. An early endosome marker RAB5 is localized within the ELYSA in the presence of inhibitors for cytoskeletons

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without inhibitors of polymerized actin (10 μM latrunculin A or cytochalasin B) or tubulin (20 μM nocodazole), and fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. Maximum intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images. Magnified regions are indicated by yellow boxes. DNA was stained with Hoechst 33342. Arrowhead and asterisk indicate the metaphase plate and first polar body, respectively.

Supplementary Figure 8

A

B

Supplementary Fig. 8. Treatment with inhibitors for cytoskeltons affected the number of LAMP1-positive objects but not their volume

GV oocytes were incubated for *in vitro* maturation (IVM) for 18 h with or without actin polymerization inhibitors (10 μM latrunculin A or cytochalasin B) or a tubulin inhibitor (20 μM nocodazole), then fixed and co-stained with anti-LC3 and anti-LAMP1 antibodies. The reconstituted objects for LAMP1-positive objects in 10 oocytes for each treatment from more than three independent experiments were analyzed for the number, size, and distribution. (A) Average total object number per oocyte. (B) Total volumes of the objects. Groups with different letters are significantly different (P < 0.05, one-way ANOVA and Tukey's multiple comparison test). Error bars indicate SEM.

Supplementary Figure 9

Supplementary Fig. 9. Number and distribution of large LAMP1-EGFP structures do not significantly differ from intrinsic LAMP1 examined with immunofluorescence

After the *in vitro* maturation (IVM) for 18 h, LAMP1-EGFP fluorescence injected as mRNA in the GV stage was compared with intrinsic LAMP1 detected with an anti-LAMP1 antibody. The reconstituted objects for LAMP1-positive structures from 3 oocytes were compared with those from immunofluorescent analysis in Figure 2 for the number and size. (A) Ratio of average numbers of total LAMP1-positive to that of MII in Figure 2. (B) Ratios of numbers of total LAMP1-positive structures sorted by size (indicated by the diameters of spheres calculated from volumes). Significance was examined using a two-tailed Student's t-test, assuming equal variances between groups; the significance is indicated with asterisks (P < 0.05). Error bars indicate SEM.

Supplementary Figure 10

Supplementary Fig. 10. LysoTracker-positive punctate structures colocalized with V1A-mCherry
Embryos were injected with V1A-mCherry mRNA (at 3 hpf) and cultured with or without bafilomycin
A$_1$ and stained with LysoTracker Green at the indicated time points. Maximum intensity projection of
confocal images at an axial scan range of 80 μm is shown as Z-projection images.

Supplementary Figure 11

A

| | GV | MII | Early 2-cell (24 hpf) | 4-cell (48 hpf) | Morula (72 hpf) | Blastocyst (96 hpf) |

BF

Magic Red B (averaged projection)

20 μm

LysoSensor (averaged projection)

B

Early 2-cell (24 hpf)

4-cell (48 hpf)

LysoSensor

Magic Red B

Merge

BF

20 μm

C

Early 2-cell

Isolated puncta

Cellular boundary

nuclear

LysoSensor
Magic-Red

4-cell

cortical

Isolated puncta

media

nuclear

LysoSensor
Magic-Red

Supplementary Fig. 11. Magic Red Cathepsin B staining in GV and MII oocytes showed higher cytoplasmic signals but similar intensity in isolated punctate structures to developing embryos
Oocytes and embryos at the indicated stages were collected and stained with Magic Red Cathepsin B solution in the same drop of medium. (A) Averaged intensity projection of confocal images at an axial scan range of 80 μm is shown as Z-projection images. (B) Early 2-cell and 4-cell embryos were stained in the same drop for LysoSensor Green and Magic Red Cathepsin B staining analysis, and confocal images were acquired. (C) Fluorescent intensities of LysoSensor and Magic Red on the yellow lines indicated in (B) are indicated by a line plot. Small, filled dots indicate the origin (0 μm) in distance.

Video 1

Video 1.
MII oocytes were fixed and co-stained with anti-RAB5 and anti-LAMP1 antibodies. A series of the confocal images with a height of 80 μm is shown from top to bottom.

Video 2

Video 2.
A series of continuous toluidine blue-stained light microscopic images of the same oocyte (top half) reconstructed to 3D. Chromosomes in the images obtained were segmented blue, toluidine blue-positive structures were segmented yellow, and oocyte PM was segmented white.

Video 3

Video 3.
GV oocytes were injected with the mixture mRNA solution for *Lamp1-EGFP* and *V1A-mCherry*, and incubated for IVM. Time-lapse movie of EGFP fluorescence of a representative oocyte is indicated as maximum intensity projection of confocal images with a height of 80 μm. Magnified perinuclear regions in Fig. 5 are indicated by a box. The timestamp indicates the time after IVM started

Video 4

Video 4
GV oocytes were injected with V1A-mCherry mRNA and incubated for 18 h for IVM in the presence of LysoSensor green. Fluorescent images of a representative oocyte which matured to MII stage is presented. A series of the confocal images of 81 slices with a height of 80 μm is shown from top to bottom.

Video 5

Video 5.
Embryos were injected with V1A-mCherry mRNA at 3 hpf and incubated for further embryogenesis in the presence of LysoSensor Green during embryonic development. A time-lapse movie of a representative embryo is shown (maximum intensity projection of confocal images at a height of 80 μm).