Title:

Endosomal-lysosomal organellar assembly (ELYSA) structures coordinate lysosomal

degradation systems through mammalian oocyte-to-embryo transition

Authors:

Yuhkoh Satouh*[1], Takaki Tatebe[1], Isei Tanida[2], Junji Yamaguchi[2], Yasuo Uchiyama[2], Ken

Sato*[1]

Affiliations:

1: Laboratory of Molecular Traffic, Institute for Molecular and Cellular Regulation, Gunma

University, Maebashi, Japan.

2: Department of Cellular and Molecular Neuropathology, Juntendo University Graduate

School of Medicine, Tokyo, Japan.

*Correspondence to:

yuhkohs@gunma-u.ac.jp

sato-ken@gunma-u.ac.jp

Abstract:

Mouse oocytes mature from the germinal vesicle (GV) to meiosis II (MII) stage and are

fertilized by a single sperm. After fertilization, the embryo degrades parts of the maternal

components via lysosomal degradation systems, including autophagy and endocytosis, as

zygotic gene expression begins during embryogenesis. Here, we demonstrate that

endosomal-lysosomal organelles form assembly structures (ELYSAs) in the periphery of

the oocyte plasma membrane. Small ELYSA structures appeared in GV oocytes and

combined to form large structures in MII oocytes. The ELYSA forms a large spherical

structure with tubular-vesicular structures approximately 2–10 μm in diameter and includes

cytosolic components at the MII stage. After fertilization, ELYSA structures are gradually

disassembled, and immature endosomes and lysosomes appear at the early 2-cell stages.

The V1-subunit of vacuolar ATPase is recruited to these endosomes and lysosomes and

causes further acidification of endosomal-lysosomal organelles, suggesting that the ELYSA

maintains endosomal-lysosomal activity in a static state in oocytes for timely activation

during early development.

Introduction:

Fertilization triggers the dynamic remodeling of cellular components via zygotic gene expression following the degradation of the cellular components of each maternal and paternal gamete. After fertilization, maternal cytosolic proteins and RNAs are selectively degraded by the ubiquitin-proteasome system (UPS)[1] and specific RNA degradation systems[2, 3], respectively. Treatment with a proteasome inhibitor or knockdown of the zygote-specific proteasome assembly chaperone gene inhibits cytokinesis in zygotes, suggesting an essential role for the UPS in early development[4, 5]. Membrane components derived from each gamete are degraded by the lysosomal degradation system via autophagy and endocytosis[6, 7]. In mice, autophagic activity is relatively low in meiosis II (MII) oocytes, whereas it is rapidly upregulated approximately 4 h after fertilization, and LC3-positive structures appear throughout the cytoplasm. Autophagic activity decreases slightly before and after the 2-cell stage but remains high from the 4- to 8-cell stage[8]. Loss of Atg5 causes embryonic lethality at the 4- to 8-cell stages, suggesting that autophagy is essential for early mouse development. In contrast, some maternal plasma membrane (PM) proteins are internalized from the PM at the late 2-cell stage and are selectively degraded by lysosomes[9]. Endocytosis inhibitors prevent cell division in 2-cell embryos, suggesting that endocytosis plays a pivotal role in early development[9, 10]. Although lysosomal degradation systems are drastically upregulated after fertilization, their regulation remains unknown.

In *Caenorhabditis elegans*, lysosomal activity in oocytes is upregulated by sperm-derived factors as the oocytes grow and mature. In this process, translational arrest of mRNA coding the V1-subunits of vacuolar ATPase in oocytes is released by the stimulation of

sperm-derived factors. As a result, the V1-subunits are recruited to the V0-subunits in the lysosomal membrane, enhancing lysosomal acidification. This process is known as lysosomal switching[11]. In mouse embryos, LysoTracker staining analysis suggests that lysosomal acidification is prominent at the 2-cell stage but not in MII oocytes and zygotes[12], raising the possibility that lysosomal switching takes place during mouse embryogenesis.

In this study, we aimed to investigate the dynamics of the endolysosomal system during mouse oocyte maturation and embryogenesis. We found that endosomes and lysosomes assembled to form spherical structures in germinal vesicle (GV)-stage oocytes, which became larger as they matured into MII oocytes. Ultrastructural analysis showed that these structures were large spherical membrane assemblies, including endosomes, lysosomes, and tubulo-vesicular structures located in the periphery of the PM. We refer to this structure as the Endosomal-LYSosomal organellar Assembly (ELYSA). The ELYSA contains several cytosolic constituents. While ELYSAs are disassembled in late-stage zygotes, immature endosomes and lysosomes gradually appear in the cytoplasm of early 2-cell stage embryos. Live imaging analysis showed that the V1-subunit of V-ATPase is targeted to the endolysosomal membrane, resulting in further acidification of lysosomal compartments during embryogenesis. These findings suggest that the ELYSA maintains low endosomal and lysosomal activities and acts as a waiting place for endolysosomal organelles to undergo fertilization-triggered activation toward embryogenesis.

Results

Endosomes and lysosomes coordinately form giant structures in MII oocytes

To examine the localization of endosomes and lysosomes in mouse MII oocytes,

early or late endosomes were stained with anti-RAB5 or -RAB7 antibody, respectively, and

their localization was compared with that of lysosomes stained with anti-LAMP1 antibody.

Giant structures harboring lysosomes with early and late endosomes were detected (**Fig. 1A,**

Supplementary Fig. 1, Supplementary Movie 1). Each giant structure in the MII oocytes

was spherical, ellipsoidal, or concatenated with small spheres with short and long diameters

of 3–5 μm and up to 10 μm, respectively. These structures were located in the peripheral

region just beneath the PM, excluding the metaphase plate region in proximity to the

metaphase chromosomes. These structures were observed until the pronuclear-zygote

(PN-zygote) stage but were mostly disassembled at the 2-cell stage. The RAB5 and

LAMP1 signals separated after the first division, indicating that these giant structures were

transiently formed (**Fig. 1 B**).

Endosomes, lysosomes, and tubulo-vesicular structures form an ELYSA

To clarify the internal structure of this giant structure containing lysosomes and

endosomes in the MII oocytes, we observed these in MII oocytes using correlative light and

electron microscopy (CLEM) using immunological reaction with anti-RAB7 and -LAMP

antibodies[13, 14]. RAB7- and LAMP1-signals were well detected even in the ultrathin section

(100 nm thickness) of the Epon-embedded MII oocyte, fixed with 0.025% glutaraldehyde

(GA), which showed similar distribution as shown in **Fig. 1** and **Supplementary Fig. 2**.

Toluidine blue staining in adjacent ultrathin section showed the RAB7/LAMP1-double

positive structures in the fluorescence observation were also positive for toluidine blue,

with interspaces were often found in the center of spheres (**Fig. 2A**). Confocal fluorescent

images in the ultrathin sections clarified that RAB7-positive signals were often observed on

the outside of these LAMP1-positive signals. Electron microscopic analysis of the

RAB7/LAMP-double positive region in the ultrathin section revealed that the

RAB7/LAMP1-dboule positive structures are assemblies of tubulo-vesicular structures, the

interior of that are either vacuolated or filled (**Fig. 2B**). The toluidine blue stain-positive

structures were confirmed to match in size and peripheral distribution with those identified

by RAB7/LAMP1 immunostaining, except for the metaphase plate, even under 2.5% GA

conditions for precise ultrastructural observation (**Supplementary Fig. 3, Supplementary**

Movie 2). Therefore, we examined the ultrastructural morphology of toluidine blue-positive

giant structures in MII oocytes fixed with 2.5% GA. The results showed that tubular

membrane vesicles with a short diameter of approximately 60.5 ± 15.9 nm assembled in the

structure, with gaps in the cytoplasm of approximately 24.0 ± 6.7 nm. Some of the vesicles

also harbored lysosome-like electron-dense contents, while others displayed multilamellar

or multivesicular body-like structures, whereas mitochondria appeared to be absent (**Fig.**

2C). Based on these observations, we termed this membrane assembly as the

Endosomal-LYSosomal organellar Assembly (ELYSA).

An autophagosome marker, LC3, also accumulated in ELYSA

Decreased autophagic activity in MII oocytes is suggested by the fact that puncta

formation of transgenically expressed GFP-LC3, an autophagosome marker, disappeared at

the MII stage[8]. Since GFP fluorescence may be quenched under acidic conditions, we

examined the localization of LC3 in MII oocytes by immunostaining with an anti-LC3

antibody. We found that LC3 hardly formed puncta in the cytoplasm, but part of the LC3

signal overlapped with the ELYSA in MII oocytes (**Fig. 3A**), suggesting that LC3-positive

membrane structures were also present in the ELYSA.

Immature ELYSA localizes in peripheral and perinuclear regions in preovulatory oocytes

We verified ELYSA formation in immature, preovulatory oocytes by performing

immunostaining for LAMP1 with RAB5 or LC3 using mouse GV oocytes. We observed

assemblies larger than 1 μm in GV oocytes, wherein fluorescent signals overlapped (**Fig.**

4A,B). The ELYSA at this stage was smaller than that in MII oocytes and was localized not

only in the peripheral region of the PM but also in the vicinity of the GV nucleus.

ELYSA maturation proceeds actin dependently and may prevent V1 component access

for vesicle acidification

Migration from the perinuclear to the peripheral region during oocyte maturation is

similar to the redistribution of the endoplasmic reticulum (ER) and the formation of cortical

ER clusters. Fitzharris et al. showed that the inhibition of actin cytoskeleton polymerization

disturbs ER migration and cortical ER cluster formation[15]. Therefore, we added latrunculin

A or cytochalasin B as inhibitors of actin polymerization and observed ELYSA formation

after *in vitro* maturation (IVM) by immunostaining. While both drugs inhibited polar body

extrusion and metaphase plate formation, as reported previously, these inhibitors reduced

ELYSA size, as indicated by LAMP1, LC3, and RAB5 immunostaining, and caused the

redistribution of LAMP1-, LC3-, and RAB5-positive signals from the cell periphery to the

cytoplasm compared with control GV oocytes (**Fig. 5A**).

147   We then observed GV oocytes injected with LAMP1-EGFP and

ATP6V1A-mCherry mRNA (V1A-mCherry: a core component of V-ATPase in the

cytoplasm, which is associated with the outer lysosomal membrane during lysosomal

acidification). Live-cell imaging of LAMP1-EGFP fluorescence clearly captured the

ELYSA maturation dynamics during IVM and revealed that ELYSA size enlargement was

accelerated after germinal vesicle breakdown (GVBD, 2–3 h after IVM started) and that

ELYSA–ELYSA assembly occurred as ELYSAs migrated to the peripheral region (**Fig. 5B,**

Supplementary Movie 3). We also examined the expression of V1A-mCherry in oocytes

after IVM. V1A-mCherry was partly detected on the surface of the ELYSA in MII oocytes

but not significantly within the ELYSA (**Fig. 5C**).

Acidification of lysosomes is initiated in coordination with ELYSA disassembly

159   We examined the relationship between ELYSA disappearance and the emergence

of acidic membrane compartments during early development. LysoSensor probes exhibit

increased fluorescence in a pH-dependent manner upon acidification and enable the

semi-quantitative analysis of cellular acidic compartments, whereas LysoTracker probes

exhibit non-pH-dependent fluorescence. Thus, we costained oocytes and embryos at

various stages with LysoTracker Red and LysoSensor Green (in the same medium drop).

LysoTracker showed a pronounced puncta-like signal after the 2-cell stage, as previously

reported[12]. In contrast, the ELYSAs in GV and MII oocytes were clearly identified by

LysoSensor, whereas LysoTracker only produced a very dim ELYSA signal (**Fig. 6A**).

A comparison of LysoTracker and V1A-mCherry showed that these fluorescent

signals colocalized well on small bright vesicles but not on the ELYSA, and both

LysoTracker and V1A-mCherry puncta were reduced in embryos cultured in the presence

of bafilomycin A_1, which inhibits the H^+ translocation of V-ATPase (**Supplementary Fig.**

4). This suggests that LysoTracker detects membrane compartment acidification, but not

that in the ELYSA. In contrast, the LysoSensor results suggested that lysosomes in the

ELYSA were not neutralized and remained acidic at a lower level, even though V1A was

difficult to localize.

Live-cell imaging of embryos expressing V1A-mCherry in the presence of

LysoSensor was performed to examine the spatiotemporal relationship between V1A

localization and lysosomal acidification. Although the fluorescence of both signals was not

stably detected until the first 2–3 frames after the start of imaging, this method revealed that

the ELYSA disappeared or disassembled throughout the PN-zygote stage until the first

mitotic division. In addition, V1A accumulation was rarely observed in the PN-zygote stage

but was rapidly enhanced in the early 2-cell embryo stage. LysoSensor-positive punctate

structures, where V1A-mCherry accumulated, increased in number during the 2-cell stage

(**Fig. 6B, Supplementary Movie 4**).

Low-level cathepsin-dependent proteolytic activity was maintained in ELYSA

Cathepsins are mainly responsible for the proteolytic activity in lysosomes. We

visualized the proteolytic activity of cathepsin B, the expression of which has been

confirmed in mouse embryos[12]. Mouse oocytes and embryos at various stages were

simultaneously collected and stained with Magic Red cathepsin B staining solution to

examine their relationship with the ELYSA. Magic Red staining showed that proteolytic

activity was detected not only in the small, isolated membrane structures (mature

lysosomes) but also in the ELYSA in the GV and MII oocytes, whereas small punctate

structures were predominantly stained in other stages (**Supplementary Fig. 5, Fig. 7A**).

The intensity plots of LysoSensor and Magic Red fluorescence revealed that the proteolytic

activity indicated by the Magic Red signal in the ELYSA was not as strong as that in

isolated punctate structures (lysosomes) in the GV and MII oocytes (**Fig. 7B**).

Discussions:

In this study, we identified a large spherical structure, the ELYSA, consisting of

endosomes, lysosomes, LC3-positive vesicles, and various vesicular-tubular structures in

mouse oocytes. ELYSAs appear in GV oocytes and become larger through assembly during

oocyte maturation. After fertilization, as ELYSAs gradually disassemble in zygotes, RAB5-,

RAB7-, or LAMP1-positive punctate structures appear in the cytoplasm of embryos at the

early 2-cell stage. The V1-subunit of V-ATPase is targeted to these structures, leading to

further acidification of the endosomal/lysosomal compartments. These results suggest that

the ELYSA functions as a reservoir that accumulates endosomal and lysosomal

compartments in the periphery of the cell cortex and maintains their activities in a static

state until embryogenesis following fertilization.

We previously found that some oocyte PM proteins are internalized from the PM at

the late 2-cell stage and are selectively degraded in lysosomes [9]. In this process, we noted

that lysosomes were not well stained by LysoTracker until the early 2-cell stage, but their

signals became stronger from the late 2-cell stage, suggesting the existence of a mechanism

underlying developmentally regulated endosomal/lysosomal activation during

embryogenesis [9, 12]. Notably, ELYSAs were stained with toluidine blue and LysoSensor,

suggesting that these structures were acidified to some extent. In addition, Magic Red,

which reflects cathepsin B activity, weakly stained ELYSAs. These observations indicate

that lysosomes in ELYSAs have some activity, albeit at low levels. Mouse oocytes

accumulate several maternal constituents, including proteins, lipids, and mRNA, and grow

to approximately 75 μm in diameter. In addition, oocytes must remain at the GV and MII

stages for an extended period to maintain their quality until fertilization. In such situations,

lysosomal degradation activity may be suppressed to a relatively low level by the ELYSA,

whereas nutrient uptake, storage, and protein synthesis are predominantly upregulated in

growing oocytes. Notably, the V1-subunits of V-ATPase were detected on the periphery of

ELYSAs but not in their interior. Thus, ELYSA formation may prevent V1-subunits from

targeting the lysosomal membrane inside the ELYSA.

Ultrastructural analysis revealed that the ELYSA consists of outer and inner

membrane assembly layers that contain several endosomes and lysosomes, respectively

(**Fig. 1, 2**). Upon disassembly of the ELYSA, endosomes in the outer layer and lysosomes

in the inner layer are expected to be released from the ELISA earlier and later, respectively.

This may explain the time lag between fertilization and lysosomal acidification beginning

at the early-2-cell stage. Since the ELYSA seems to contain cytosolic constituents, it may

accommodate some cytosolic constituents as a capsule to provide nutrition or signaling

factors for development.

Since the distance between membrane structures in the ELYSA appears to be

relatively uniform, cytoplasmic components in the ELYSA may contain factors that

promote membrane structure tethering or assembly. In fact, we observed that the cytoplasm

between the membrane structures inside the ELYSA had a slightly higher electron density.

In the future, identifying the cytoplasmic molecules tethering these membrane structures to

the ELYSA will be necessary.

We also found that ELYSAs were gradually distributed from the perinuclear region

to the periphery of the oocyte cell cortex, except in the MII cell plate, from the GV to the

MII stage. This movement is reminiscent of that of the ER and cortical ER clusters, whose

distribution is regulated by the actomyosin system[15]. In fact, actin cytoskeleton

polymerization inhibitors resulted in a reduced size of ELYSAs and their redistribution

from the cell cortex to the cytoplasm (**Fig. 5**), suggesting that the actin cytoskeleton is

involved in the formation and distribution of ELYSAs in oocytes.

Autophagy and endocytosis are upregulated in zygotes and 2-cell stage embryos,

respectively, and their substrates are delivered to the lysosomes for degradation. Our study

showed that lysosomal acidification was enhanced from the 2- to 4-cell stage and after the

morula stage. The first lysosomal acidification occurred when ELYSAs were disassembled

in late zygotes. These findings suggest that the timing of degradative substrate delivery to

lysosomes is developmentally linked to lysosomal acidification and maturation. Notably,

minor and major zygotic gene activation occurs at the 1- and 2-cell stages, respectively,

during mouse development, implying that lysosomal degradation of maternal components

is coupled with zygotic gene expression to promote embryogenesis[16]. Fertilization appears

to trigger ELYSA disassembly, but the downstream signaling pathway remains to be

identified and needs to be studied in the future.

Recently, it has been reported that appropriate lysosomal activity is important for

mammalian development[10]. Bafilomycin A_1 treatment arrests embryonic development at the

4–8-cell stage[12]. Inhibition of cathepsin activity using a mixture of E64d and pepstatin A

blocks the development of the 8-cell or morula stage[12]. However, autophagic activity in

fertilized eggs and embryonic developmental potential are correlated. Using GFP-LC3 as

an indicator of autophagic activity, Tsukamoto et al. found that aged eggs from 14- to

15-month-old mice had lower autophagic activity than those from younger mice, probably

due to decreased lysosomal activity. Females transplanted with embryos with high

autophagic activity have a significantly higher litter size ratio than that of control females[17].

Analysis of ELYSA formation and disassembly in aged oocytes and embryos may be a

useful tool for determining oocyte quality by examining whether appropriate regulation of

endosomal/lysosomal activities occurs.

Methods:

Animal experiments and strains

Neither randomization nor blinding was used for animal selection (8–16-week-old). All

experimental protocols involving animals were approved and performed in accordance with

the guidelines of the Animal Care and Experimentation Committee of Gunma University

(Approval No. 22-076). Hybrid B6D2F1 mice were purchased from the Japan SLC

(Hamamatsu, Japan)

Antibodies and inhibitors

The following primary antibodies were used: rabbit monoclonal anti-RAB5 (C8B1; 3547;

Cell Signaling Technology, Danvers, MA, USA), rabbit monoclonal anti-RAB7 (D95F2;

9367; Cell Signaling Technology), rat monoclonal anti-mouse LAMP1 (1D4B; sc-19992;

Santa Cruz Biotechnology, Dallas, TX, USA), and mouse monoclonal anti-rat LC3 (Clones

4E12; M152-3; Medical and Biological Laboratories, Tokyo, Japan). The following

secondary antibodies were used: goat anti-rabbit IgG (H+L) conjugated with Alexa Fluor

555 (A-21428; Thermo Fisher Scientific, Waltham, MA, USA), goat anti-rat IgG (H+L)

cross-adsorbed secondary antibody conjugated with Alexa Fluor 488 and 647 (A-11006 and

A-21247, respectively; Thermo Fisher Scientific), and goat anti-mouse IgG (H+L)

cross-adsorbed secondary antibody conjugated with Alexa Fluor 546 (A-11030; Thermo

Fisher Scientific). The following inhibitors or solutions were used for IVM or embryonic

culture assays: dimethyl sulfoxide (DMSO) (13406-55; Nacalai Tesque, Kyoto, Japan);

latrunculin A actin polymerization inhibitor that binds to monomer actin (428021;

Sigma-Aldrich, St. Louis, MO, USA); and cytochalasin B, an actin polymerization inhibitor

that binds to the plus-end of polymerized actin (C6762; Sigma-Aldrich). Stock solutions of

latrunculin A or cytochalasin B were prepared by solubilizing each in DMSO at 400×

concentration for assay. Bafilomycin A_1 was dissolved in DMSO (L266; Dojindo,

Kumamoto, Japan) at a final concentration of 25 nM.

Oocyte preparation

GV oocytes were collected from the ovaries of female mice 46 h after injection with pregnant mare serum gonadotropin or an anti-inhibin antibody (CARD HyperOva; Kyudo, Kumamoto, Japan). Antral follicles were suspended in 30 μL of FHM medium drops (Sigma-Aldrich) covered with liquid paraffin (Nacalai Tesque), punctured using 26G needles (Terumo, Tokyo, Japan), and the GV oocytes were collected using a glass needle with a mouth pipette. MII oocytes were collected from the oviducts of female mice 13 h after injection with CARD HyperOva and 7.5 IU human chorionic gonadotropin (hCG; ASKA Pharmaceutical, Tokyo, Japan) 48 h apart. The oviductal wall was punctured using a 26G needle to collect cumulus-oocyte complexes in 20 μL potassium simplex optimization medium (KSOM; Kyudo) drops covered with liquid paraffin.

DNA vector construction and mRNA synthesis

Complementary DNA for mouse *Lamp1* (accession number NM_001317353) or *Atp6v1a* (accession number NM_001358204) was amplified using PCR with High-fidelity Taq Polymerase, KOD FX Neo (Toyobo, Osaka, Japan), and a mouse ovary cDNA library as a template (Genostaff, Tokyo, Japan). The amplicon was then subcloned into the entry vector pDONR221, and transferred into the destination vectors pDEST-CMV-C-EGFP (#122844; for *Lamp1*; Addgene, Watertown, MA, USA) or pDest-mCherry-N1 (#31907; for *Atp6v1a*; Addgene) using Gateway recombination cloning technology. The following primer combinations were used for the constructions:

5′-GCGCACAAGTTTGTACAAAAAAGCAGGCTGCCACCATGGCGGCCCCCGGCG-

3′   and

5′-GCGCACCACTTTGTACAAGAAAGCTGGGTTGTCTTCAAGGCTACGGAATG-3′

(for   *Lamp1*),   or

5′-GCGCACCACTTTGTACAAGAAAGCTGGGTTGATGGTCTGATAGCCGGCGTGA

CT-3′   and

5′-GCGCACCACTTTGTACAAGAAAGCTGGGTTGTCTTCAAGGCTACGGAATG-3′

(for *Atp6v1a*). The cDNA encoding *Lamp1-EGFP* or *Atp6v1a-mCherry* were amplified

using the following primer combinations: 5′-CGCAAATGGGCGGTAGGCGTG-3′ and

5′-TCCAGCAGGACCATGTGATCGC-3′   (for   *Lamp1-EGFP*),   or

5′-CGCAAATGGGCGGTAGGCGTG-3′ and 5′-TTGGTCACCTTCAGCTTGG-3′ (for

Atp6v1a-mCherry). RNA synthesis using each PCR amplicon as a template, poly-A tail

addition, and mRNA purification were performed, using an mMESSAGE mMACHINE T7

Ultra kit (Thermo Fisher Scientific) and MEGAclear transcription clean-up kit (Thermo

Fisher Scientific), according to the manufacturer's protocols. Synthesized mRNAs were

dissolved in RNase-free demineralized water at a concentration of 400 ng/μL and frozen in

aliquots until use.

In vitro *maturation, fertilization, and embryonic culture*

IVM for the GV oocytes was conducted by washing the oocytes twice and culturing in

alpha-modified Eagle's minimum essential medium (Thermo Fisher Scientific) containing

5% (vol/vol) fetal bovine serum (FBS), 0.1 of IU/mL follicle-stimulating hormone (MSD,

Tokyo, Japan), 1.2 IU/mL of hCG (ASKA Pharmaceutical), and 4 ng/mL of epidermal

growth factor (Thermo Fisher Scientific) for 17 h at 37 °C under a 5% CO_2 atmosphere[18].

For *in vitro* fertilization, spermatozoa from a B6D2F1 epididymis were pre-incubated in a

100-μL modified human tubal fluid medium (Kyudo) drop covered with liquid paraffin and

inseminated at a sperm concentration of 1.5×10^5 per mL for 1.5 h. Excess spermatozoa

were washed out, and the oocytes were incubated in a 20-μL KSOM drop (Kyudo) for

further development. IVM assays using latrunculin A and cytochalasin B were performed

by transferring oocytes to medium drops containing each inhibitor or DMSO 4 h after the

start of IVM (waiting for the completion of GVBD), as previously reported[15]. IVM media

containing DMSO, latrunculin A, or cytochalasin B was prepared by diluting the stock

solution (1:400), and the final concentration of the inhibitors was 10 μM. The embryonic

culture assay using bafilomycin A_1 was conducted by transferring the PN-zygote into a

20-μL drop of KSOM containing 25 nM bafilomycin A_1 covered with liquid paraffin at 6 h

post-fertilization (hpf). Embryos were transferred at 24, 48, and 72 hpf to freshly prepared

KSOM containing bafilomycin A_1.

Immunostaining

Immunostaining of oocytes or embryos was performed as previously described [9]. Oocytes

or embryos were cultured to the indicated stages and fixed in 4% paraformaldehyde (PFA;

Nacalai Tesque) in phosphate-buffered saline (PBS) containing 0.1 mg/mL polyvinyl

alcohol (PVA; Sigma-Aldrich) (PVA/PBS) for 15 min at 25–27 °C. After fixation, the cells

were permeabilized with PVA/PBS containing 0.1% Triton X-100 (Sigma-Aldrich) for 30

min at 25–27 °C and washed thrice with PVA/PBS. After blocking with 10% FBS (Thermo

Fisher Scientific) in PBS for 1 h at room temperature, the cells were incubated with the

aforementioned primary antibodies (1:100 for all) overnight at 4 °C. Next, the cells were

washed thrice in PVA/PBS, and incubated with anti-mouse, rat, or rabbit IgG secondary

antibodies coupled with Alexa Fluor 488, 555, or 564 (1:200 for all; Thermo Fisher

Scientific) at 25–27 °C for 2–4 h. Thereafter, the cells were incubated with PVA/PBS

containing 10 µg/mL Hoechst 33342 (Dojindo) for 10 min and washed with fresh PVA/PBS

a few times. Subsequently, control and nonspecific staining were performed by processing

oocytes and embryos in the absence of primary antibodies. The oocytes and embryos were

imaged in PVA/PBS microdrops in 35-mm glass-bottomed dishes (MatTek Corporation,

Ashland, MA, USA). For each position, confocal images at 81 z-axis planes with 1-µm

increments were acquired on an IX-71 inverted microscope (Olympus Corporation, Tokyo,

Japan) equipped with a CSU W-1 confocal scanner unit (Yokogawa Electric Corporation)

using a 60× silicone immersion lens (Olympus Corporation).

LysoTracker/LysoSensor staining of oocytes at various stages

Acidic organelles in the oocytes were visualized using LysoTracker Red DND-99 and

LysoSensor Green DND-189 (Thermo Fisher Scientific) according to the manufacturer's

instructions. Briefly, the oocytes or embryos at target stages for one experimental round

were obtained simultaneously and incubated in a large drop of reaction mixture (KSOM

containing 1/10,000 Magic Red stock solution dissolved in DMSO and 1/1,000 LysoSensor

Green DND-189 stock solution dissolved in DMSO) for 30 min at 37 °C under 5% CO_2 in

a humidified atmosphere in liquid paraffin in a 35-mm glass-bottom dish (MatTek

Corporation). Subsequently, confocal images at 41 z-axis planes with 2-μm increments

were obtained using a CSU W-1 spinning disk confocal microscope with a 30× silicone

immersion lens.

Magic Red/LysoSensor staining

Intracellular cathepsin activity was assayed in oocytes or embryos using Magic Red

Cathepsin B or Cathepsin L detection kits (ImmunoChemistry Technologies LLC, Davies,

CA, USA) according to the manufacturer's instructions. Briefly, the oocytes or embryos at

target stages for one experimental round were obtained simultaneously and incubated in a

large drop of reaction mixture (KSOM containing 1/250 Magic Red stock solution

dissolved in DMSO and 1/1,000 LysoSensor Green DND-189 stock solution dissolved in

DMSO) for 25 min at 37 °C under 5% CO_2 in a humidified atmosphere in liquid paraffin in

a 35-mm glass-bottom dish (MatTek Corporation). Afterward, Hoechst 33342 (10 μg/mL)

was added to the drop to be a final concentration of 10 μg/mL, and incubated for a further 5

min. Subsequently, oocytes or embryos at the same stage were collected, and confocal

images at 41 z-axis planes with 2-μm increments for both 470-nm (LysoSensor) and

555-nm (Magic Red) lasers were obtained on a CSU W-1 spinning disk confocal

microscopy using a 30× silicone immersion lens.

In-resin CLEM and electron microscopy

In-resin CLEM was performed as previously described with some modifications[13, 14].

Briefly, oocytes were fixed in 4% PFA and 0.025% glutaraldehyde (GA). After

permeabilization with 50 μg/mL digitonin (Nacalai Tesque) and 0.02% Triton X-100,

oocytes were washed with PVA/PBS three times and incubated with PBS containing 10%

(w/v) FBS (Thermo Fisher Scientific) for 1 h at room temperature. Thereafter, oocytes were

incubated with PVA/PBS containing primary antibody overnight at 4 °C. After incubation

with a secondary antibody in PVA/PBS for 2 h, oocytes were fixed in 4% PFA and 2.5%

GA, embedded in 1.5% agarose, stained with osmium tetroxide, dehydrated with graded

ethanol solutions (QY1), and embedded in epoxy resin (Oken Shoji, Tokyo, Japan).

Ultrathin sections of Epon-embedded oocytes (100-nm thickness) were prepared using a

Leica UC6 ultramicrotome (Leica, Nussloch, Germany) and placed on glass coverslips.

Subsequently, fluorescence microscopy images of the sections were obtained using a Nikon

A1RHD25 confocal laser scanning microscope with a NIS-Elements software (Nikon,

Tokyo, Japan). Electron microscopic images were observed using a Helios Nanolab 660

FIB-SEM instrument (FEI Company, Hillsboro, USA).

For conventional electron microscopy, the oocytes were fixed in the presence of 2% PFA

and 2.5% GA and stained with osmium tetroxide. After dehydration with ethanol, the

oocytes were embedded in epoxy resin, as described above. Sections were cut at a thickness

of 100 nm using a UC6 ultramicrotome, mounted on glass coverslips, stained with uranyl

acetate and lead citrate, and observed under a Regulus8240 scanning electron microscope

(Hitachi High-Tech Corporation, Tokyo, Japan).

Serial semi-thin sections and 3D reconstruction of toluidine blue-positive structures

Serial semi-thin sections were cut at a thickness of 300 nm using a UC6 ultramicrotome

and mounted on glass slides. Next, the sections were stained with 1% toluidine blue/0.1 M

PB, and images were obtained using a BX50 light microscope (Olympus Corporation).

Serial toluidine blue-stained images were aligned using FIJI (National Institutes of Health,

MD, USA) with plugin-linear stack alignment using SIFT (LSAS), and 3D reconstruction

was performed using Amira software (Thermo Fisher Scientific).

Microinjection into the oocytes

Microinjection of mRNA into GV oocytes was performed as previously described[19, 20].

Briefly, antral follicles were harvested in FHM medium containing 250 μM of dbcAMP

using 26G needles (Terumo). After dissociating the cumulus cells by pipetting, the oocytes

were transferred to medium supplemented with 250 μM of dbcAMP and 10 μM of

cytochalasin B (Sigma-Aldrich). Thereafter, mRNA injection into the oocytes was

performed using a piezo-micromanipulator (Primetech, Tokyo, Japan) with a glass capillary

needle. The mRNA concentrations were 50 and 100 ng/μL for *V1A-mCherry* and

Lamp1-EGFP, respectively. The oocytes were washed thrice and used for IVM. mRNA was

microinjected into MII oocytes as previously described. The concentration of mRNA was

the same as that in the GV oocytes.

Time-lapse confocal imaging of oocytes/embryos

Low-invasive confocal fluorescence imaging of oocytes/embryos was performed as

previously described[21, 22, 23] with slight modifications. Briefly, on an inverted microscope

IX-71 (Olympus Corporation) equipped with a CSU W-1 confocal scanner unit (Yokogawa

Electric Corporation), an incubation chamber (Tokai Hit, Fujinomiya, Japan) was set at

37 °C on the microscope stage, with a gas mixture of 5% CO_2 introduced into the chamber

at 150 mL/min. Oocytes/embryos were placed in 8 μL of IVM medium or KSOM drops

covered with liquid paraffin (Nacalai Tesque) in 35-mm glass-bottomed dishes (MatTek

Corporation). Subsequently, confocal images at 41 *z*-axis planes with 2-μm increments for

excitation lasers of 470 and 555 nm from an LDI-7 Laser Diode Illuminator (89North,

Williston, VT, USA) were obtained every 30 or 60 min during IVM or embryonic culture

observation, respectively, through optimized band-pass filters using 30× silicone immersion

lens.

Imaging analysis

Analysis of the acquired images, including the generation of a line plot of intensity,

adjustment of intensity, and generation of montage images at a uniform intensity, was

performed using ImageJ software (National Institutes of Health). To enhance the images, a

Gaussian Blur filter (at a sigma radius of 1) and auto-adjustment of the intensity were

applied to the stack of images. To adjust the intensity, auto-adjustment of the intensity was

applied to each image. The movies were concatenated using Premiere Pro software (Adobe,

San Jose, CA, USA).

Data availability:

The data supporting the findings of this study are available from the corresponding authors

upon request.

Materials availability statement:

The materials supporting the findings of this manuscript are available from the

corresponding authors upon reasonable request.

Acknowledgments:

We would like to thank Editage (www.editage.jp) for the English language editing.

Funding:

This study was supported by the Japan Society for the Promotion of Science KAKENHI

(Grant Numbers 19H05711 and 20H00466 to K. Sato, 21H02435, 22H02872, and

22H04652 to I. Tanida, and 20K22744 to J. Yamaguchi), the Mochida Memorial

Foundation for Medical and Pharmaceutical Research, Takeda Science Foundation (Grant

Number 2022093867), and The Cell Science Research Foundation to Y. Satouh. This work

was partly supported by the Project for Elucidating and Controlling Mechanisms of Aging

and Longevity from the Japan Agency for Medical Research and Development (AMED

21gm5010003 to Y. Uchiyama., 22gm1710001 s0201 and 23gm1710001s0202 to I. Tanida)

and by the MEXT-supported Program for the Strategic Research Foundation at Private

Universities (to Y. Uchiyama), and the Center of Genomic and Regeneration Medicine,

Juntendo University Graduate School of Medicine (to Y. Uchiyama and I. Tanida).

Competing interests:

No competing interests declared.

Author contributions:

Y.S. and K.S. established the essential research concepts for this study. Y.S. and T.T.

performed the immunostaining and assays using mouse oocytes/embryos. I. T., J. Y., and Y.

U. performed the CLEM experiments and analyses. Y. S. and K. S. mainly prepared the

manuscript.

References:

1. Verlhac MH, Terret ME, Pintard L. Control of the oocyte-to-embryo transition by

the ubiquitin-proteolytic system in mouse and *C. elegans*. *Curr Opin Cell Biol* **22**,

758-763 (2010).

2. Jiang ZY, Fan HY. Five questions toward mRNA degradation in oocytes and

preimplantation embryos: when, who, to whom, how, and why? *Biol Reprod* **107**,

62-75 (2022).

3. Schier AF. The maternal-zygotic transition: death and birth of RNAs. *Science* **316**,

406-407 (2007).

4. Shin SW*, et al.* Mouse zygote-specific proteasome assembly chaperone important

for maternal-to-zygotic transition. *Biol Open* **2**, 170-182 (2013).

5.   Shin SW*, et al.* Inhibition of the ubiquitin-proteasome system leads to delay of the

onset of ZGA gene expression. *J Reprod Dev* **56**, 655-663 (2010).

6.   Sato K. Multiple roles of endocytosis and autophagy in intracellular remodeling

during oocyte-to-embryo transition. *Proc Jpn Acad Ser B Phys Biol Sci* **98**, 207-221

(2022).

7.   Birgisdottir AB, Johansen T. Autophagy and endocytosis - interconnections and

interdependencies. *J Cell Sci* **133**, (2020).

8.   Tsukamoto S, Kuma A, Murakami M, Kishi C, Yamamoto A, Mizushima N.

Autophagy is essential for preimplantation development of mouse embryos. *Science*

321, 117-120 (2008).

9.   Morita A, Satouh Y, Kosako H, Kobayashi H, Iwase A, Sato K. Clathrin-mediated

endocytosis is essential for the selective degradation of maternal membrane proteins

and preimplantation development. *Development* **148**, dev199461 (2021).

10.   Satouh Y, Sato K. Reorganization, specialization, and degradation of oocyte

maternal components for early development. *Reprod Med Biol* **22**, e12505 (2023).

11.   Bohnert KA, Kenyon C. A lysosomal switch triggers proteostasis renewal in the immortal *C. elegans* germ lineage. *Nature* **551**, 629-633 (2017).

12.   Tsukamoto S*, et al.* Functional analysis of lysosomes during mouse preimplantation embryo development. *J Reprod Dev* **59**, 33-39 (2013).

13.   Tanida I, Yamaguchi J, Suzuki C, Kakuta S, Uchiyama Y. Application of immuno- and affinity labeling with fluorescent dyes to in-resin CLEM of Epon-embedded cells. *Heliyon* **9**, e17394 (2023).

14.   Mitsui S, Yamaguchi J, Suzuki C, Uchiyama Y, Tanida I. TUNEL-positive structures in activated microglia and SQSTM1/p62-positive structures in activated astrocytes in the neurodegenerative brain of a CLN10 mouse model. *Glia* **71**, 2753-2769 (2023).

15.   FitzHarris G, Marangos P, Carroll J. Changes in endoplasmic reticulum structure during mouse oocyte maturation are controlled by the cytoskeleton and cytoplasmic dynein. *Dev Biol* **305**, 133-144 (2007).

16.   Schulz KN, Harrison MM. Mechanisms regulating zygotic genome activation. *Nat Rev Genet* **20**, 221-234 (2019).

564   17.   Tsukamoto S*, et al.* Fluorescence-based visualization of autophagic activity predicts

565   mouse embryo viability. *Sci Rep* **4**, 4533 (2014).

567   18.   Morohaku K*, et al.* Complete in vitro generation of fertile oocytes from mouse

568   primordial germ cells. *Proc Natl Acad Sci U S A* **113**, 9021-9026 (2016).

570   19.   Umeda R*, et al.* Structural insights into tetraspanin CD9 function. *Nat Commun* **11**,

571   1606 (2020).

573   20.   Kato K*, et al.* Structural and functional insights into IZUMO1 recognition by JUNO

574   in mammalian fertilization. *Nat Commun* **7**, 12198 (2016).

576   21.   Nozawa K, Satouh Y, Fujimoto T, Oji A, Ikawa M. Sperm-borne phospholipase C

577   zeta-1 ensures monospermic fertilization in mice. *Sci Rep* **8**, 1315 (2018).

579   22.   Satouh Y, Nozawa K, Yamagata K, Fujimoto T, Ikawa M. Viable offspring after

580   imaging of Ca2+ oscillations and visualization of the cortical reaction in mouse

581   eggs. *Biol Reprod* **96**, 563-575 (2017).

583   23.   Satouh Y, Inoue N, Ikawa M, Okabe M. Visualization of the moment of mouse

584   sperm-egg fusion and dynamic localization of IZUMO1. *J Cell Sci* **125**, 4985-4990

(2012).

Figure legends:

Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte

plasma membrane.

(A) Meiosis II (MII) oocytes were fixed and costained with anti-RAB5 and anti-LAMP1

antibodies. The schematic diagram indicates the location of oocyte chromosomes and

metaphase plate in the MII oocytes. Maximum intensity projection of confocal images of

80-µm in height are shown as Z-projection images. Arrowheads indicate the positions of

oocyte chromosomes. Magnified regions (right) are indicated by yellow boxes. DNA was

stained with Hoechst 33342. (B) Embryos at different stages were fixed and costained with

anti-RAB5 and anti-LAMP1 antibodies. Hour(s) post-fertilization is indicated as hpf.

Fig. 2. Endosomal-lysosomal organelles form assembly structures (ELYSA).

(A) The internal structure of the giant structure in MII oocytes was analyzed by correlative

light and electron microscopy (CLEM) analysis using anti-RAB7 and anti-LAMP1

antibodies. A merged image of immunostaining and electron microscopy (EM) is indicated

(left), and a toluidine blue staining image of the adjacent thin section (right) is indicated.

Black arrowheads indicate the giant structures positive for the RAB7/LAMP1/toluidine

blue stainings. Note that RAB7/LAMP1 staining is indicated in a pseudo-color. (B) The

fluorescent distribution of RAB7/LAMP1 immunostaining was analyzed with

deconvolution confocal microscopy. Note that RAB7/LAMP1 staining is indicated in

pseudo-color. (C) The internal structures of the toluidine blue-positive structure, ELYSA,

were observed using conventional electron microscopy of oocytes fixed with 2.5%

glutaraldehyde. Magnified regions in a or b (right) are indicated by boxes (left). Arrows,

arrowheads, or asterisks indicate vesicles harboring high electron density contents and

multilamellar or multivesicular structures, respectively. ELYSA, endosomal lysosomal

organellar assembly.

Fig. 3. An autophagy marker LC3 is also detected in ELYSA.

MII oocytes were fixed and costained with anti-LC3 and anti-LAMP1 antibodies. The

maximum intensity projection of the confocal images of 80-μm in height is shown as a

Z-projection image. Magnified regions (right) are indicated by yellow boxes. DNA was

stained with Hoechst 33342. Arrowheads indicate the positions of oocyte chromosomes.

Fig. 4. ELYSA grows and migrates to the peripheral region during oocyte maturation.

Germinal vesicle oocytes were fixed and costained with anti-RAB5 and anti-LAMP1

antibodies (A) or anti-LC3 and anti-LAMP1 antibodies (B). The maximum intensity

projection of the confocal images of 80-μm in height is shown as a Z-projection image.

Magnified perinuclear regions (right) are indicated by yellow boxes. DNA was stained with

Hoechst 33342.

Fig. 5. Enlargement of ELYSA and its redistribution to the cell periphery occur in an

actin cytoskeleton-dependent manner.

(A) GV oocytes, after *in vitro* maturation (IVM) treatment with or without actin polymerization inhibitors (10 μM latrunculin A or cytochalasin B), were fixed and costained with anti-LC3 and anti-LAMP1 antibodies. Maximum intensity projection of confocal images of 80-μm in height are shown as Z-projection images. DNA was stained with Hoechst 33342. Arrowheads or asterisks indicate the metaphase plate or first polar body, respectively. (B) EGFP fluorescence of GV oocytes injected with *Lamp1-EGFP* and *V1A-mCherry* mRNA was analyzed using time-lapse imaging. Maximum intensity projection of confocal images of 40-μm in height (bottom half of the oocyte) are shown. Magnified perinuclear regions (right) are indicated by yellow boxes. Arrowheads indicate assemblies that adhere to each other in the subsequent frame. (C) The same oocyte in (B) after IVM. Maximum intensity projection of confocal images for both EGFP and mCherry fluorescence of 80-μm in height are shown. Magnified regions in a or b (right) are indicated by yellow boxes.

Fig. 6. Endosomes and lysosomes in the ELYSA are acidified weakly, but further acidification occurs after ELYSA disassembly.

(A) Oocytes and embryos at different stages were stained in the same drop for LysoSensor Green and LysoTracker Red staining analysis, and confocal images were acquired. (B) Green and red fluorescence of pronuclear oocytes injected with *V1A-mCherry* mRNA was analyzed using time-lapse imaging in the presence of LysoSensor Green to the blastocyst stage. Magnified perinuclear regions are indicated by yellow boxes. Maximum intensity projection of confocal images of 80-μm in height are shown unless otherwise stated.

Fig. 7. Proteolytic activity of cathepsin B is maintained relatively at a low level in the ELYSA.

(A) GV and MII oocytes were stained in the same drop for LysoSensor Green and Magic Red cathepsin B staining analysis, and confocal images were acquired. (B) Fluorescent intensities of LysoSensor and Magic Red on the yellow lines indicated in (A) are indicated by a line plot. Small, filled dots indicate the origin (0 μm) in distance.

Supplementary Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte plasma membrane.

(A) MII oocytes were fixed and costained with anti-RAB7 and anti-LAMP1 antibodies. The schematic diagram indicates the location of oocyte chromosomes and the metaphase plate in the MII oocytes. Maximum intensity projection of confocal images of 80-μm in height are shown as Z-projection images. Arrowheads indicate the positions of oocyte chromosomes.

Supplementary Fig. 2. Giant structures in MII oocytes were immunostained with anti-RAB7 and LAMP1 antibodies after glutaraldehyde (GA) fixation.

(A) MII oocytes were fixed with 0.025% GA and costained with anti-RAB5/RAB7 and anti-LAMP1 antibodies. Single-section images from confocal observation are indicated.

Supplementary Fig. 3. Toluidine blue-positive structures in MII oocytes show similar

localization patterns to the ELYSA.

(A) Aligned serial toluidine blue-stained light microscopic images were stacked and

three-dimensionally reconstructed by inversing the intensity and adjusting it to be

transparent. (B) A series of images of toluidine blue-stained serial semi-thin sections of the

Epon-embedded MII oocyte were reconstructed to 3D. Based on the images obtained,

chromosomes were segmented blue, toluidine blue-positive structures were segmented

yellow, and oocyte PM was segmented white.

Supplementary Fig. 4. LysoTracker-positive punctate structures colocalized with

V1A-mCherry.

Green and red fluorescence images of embryos expressing V1A-mCherry (mRNA injected

at 3 h post-fertilization [hpf]) and stained with LysoTracker Green at the indicated time

points were acquired. The maximum intensity projection of confocal images of 80-μm in

height is indicated.

Supplementary Fig. 5. Magic Red cathepsin B staining gave signals in the cytoplasm

as well as the ELYSA in GV and MII oocytes..

Fluorescence images of oocytes and embryos at the indicated stages were obtained and

stained with Magic Red Cathepsin B solution (and LysoTracker Green; not shown) in the

same drop of medium. The average intensity projections of confocal images of 80-μm in

height are shown.

Supplementary Movie 1.

MII oocytes were fixed and costained with anti-RAB5 and anti-LAMP1 antibodies. A series

of the confocal images of 81 slices of 80-μm in height is shown from top to bottom.

Supplementary Movie 2.

A series of continuous toluidine blue-stained light microscopic images of same oocyte (top

half) reconstructed to 3D. Chromosomes in the images obtained were segmented blue,

toluidine blue-positive structures were segmented yellow, and oocyte PM was segmented

white.

Supplementary Movie 3.

Time-lapse movie of EGFP fluorescence in the GV oocytes injected with *Lamp1-EGFP* and

V1A-mCherry mRNA during IVM with a maximum intensity Z-projection of confocal

images of 80-μm in height. Magnified perinuclear regions in Fig. 5 are indicated by a box.

The timestamp indicates the time after IVM started.

Supplementary Movie 4.

Time-lapse movie of fertilized oocytes expressing V1A-mCherry mRNA in the presence of

LysoSensor Green during embryonic development. The maximum intensity projection of

confocal images of 80-μm in height is shown.

Figure 1

Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte plasma membrane.
(A) Meiosis II (MII) oocytes were fixed and costained with anti-RAB5 and anti-LAMP1 antibodies. The schematic diagram indicates the location of oocyte chromosomes and metaphase plate in the MII oocytes. Maximum intensity projection of confocal images of 80-μm in height are shown as Z-projection images. Arrowheads indicate the positions of oocyte chromosomes. Magnified regions (right) are indicated by yellow boxes. DNA was stained with Hoechst 33342. (B) Embryos at different stages were fixed and costained with anti-RAB5 and anti-LAMP1 antibodies. Hour(s) post-fertilization is indicated as hpf.

Figure 2

A

CLEM

RAB7
LAMP1

20 μm

0.025% GA

Toluidine blue
(adjacent section to CLEM)

B

Merge

LAMP1

RAB7

EM

2 μm

C

2.5% GA

b

a

2 μm

a

200 nm

b

Fig. 2. Endosomal-lysosomal organelles form assembly structures (ELYSA).
(A) The internal structure of the giant structure in MII oocytes was analyzed by correlative light and electron microscopy (CLEM) analysis using anti-RAB7 and anti-LAMP1 antibodies. A merged image of immunostaining and electron microscopy (EM) is indicated (left), and a toluidine blue staining image of the adjacent thin section (right) is indicated. Black arrowheads indicate the giant structures positive for the RAB7/LAMP1/toluidine blue stainings. Note that RAB7/LAMP1 staining is indicated in a pseudo-color. (B) The fluorescent distribution of RAB7/LAMP1 immunostaining was analyzed with deconvolution confocal microscopy. Note that RAB7/LAMP1 staining is indicated in pseudo-color. (C) The internal structures of the toluidine blue-positive structure, ELYSA, were observed using conventional electron microscopy of oocytes fixed with 2.5% glutaraldehyde. Magnified regions in a or b (right) are indicated by boxes (left). Arrows, arrowheads, or asterisks indicate vesicles harboring high electron density contents and multilamellar or multivesicular structures, respectively. ELYSA, endosomal lysosomal organellar assembly.

Figure 3

Fig. 3. An autophagy marker LC3 is also detected in ELYSA.
MII oocytes were fixed and costained with anti-LC3 and anti-LAMP1 antibodies. The maximum intensity projection of the confocal images of 80-µm in height is shown as a Z-projection image. Magnified regions (right) are indicated by yellow boxes. DNA was stained with Hoechst 33342. Arrowheads indicate the positions of oocyte chromosomes.

Figure 4

Fig. 4. ELYSA grows and migrates to the peripheral region during oocyte maturation.
Germinal vesicle oocytes were fixed and costained with anti-RAB5 and anti-LAMP1 antibodies (A) or anti-LC3 and anti-LAMP1 antibodies (B). The maximum intensity projection of the confocal images of 80-μm in height is shown as a Z-projection image. Magnified perinuclear regions (right) are indicated by yellow boxes. DNA was stained with Hoechst 33342.

Figure 5

Fig. 5. Enlargement of ELYSA and its redistribution to the cell periphery occur in an actin cytoskeleton-dependent manner.

(A) GV oocytes, after *in vitro* maturation (IVM) treatment with or without actin polymerization inhibitors (10 µM latrunculin A or cytochalasin B), were fixed and costained with anti-LC3 and anti-LAMP1 antibodies. Maximum intensity projection of confocal images of 80-µm in height are shown as Z-projection images. DNA was stained with Hoechst 33342. Arrowheads or asterisks indicate the metaphase plate or first polar body, respectively. (B) EGFP fluorescence of GV oocytes injected with *Lamp1-EGFP* and *V1A-mCherry* mRNA was analyzed using time-lapse imaging. Maximum intensity projection of confocal images of 40-µm in height (bottom half of the oocyte) are shown. Magnified perinuclear regions (right) are indicated by yellow boxes. Arrowheads indicate assemblies that adhere to each other in the subsequent frame. (C) The same oocyte in (B) after IVM. Maximum intensity projection of confocal images for both EGFP and mCherry fluorescence of 80-µm in height are shown. Magnified regions in a or b (right) are indicated by yellow boxes.

Figure 6

A

	GV	MII	2-cell (24 hpf)	4-cell (48 hpf)	Morula (72 hpf)	Blastocyst (96 hpf)

BF — 40 µm

LysoSensor

LysoTracker

B

PN-zygote	PN-zygote	Early 2-cell	Late 2-cell	4-cell	Morula	Blastocyst

LysoSensor

Adjusted

Single slice (enlarged/enhanced)

V1A-mCherry

BF   20 µm

06:00   16:00   22:00   35:00   45:00   70:00   95:00

Fig. 6. Endosomes and lysosomes in the ELYSA are acidified weakly, but further acidification occurs after ELYSA disassembly.

(A) Oocytes and embryos at different stages were stained in the same drop for LysoSensor Green and LysoTracker Red staining analysis, and confocal images were acquired. (B) Green and red fluorescence of pronuclear oocytes injected with *V1A-mCherry* mRNA was analyzed using time-lapse imaging in the presence of LysoSensor Green to the blastocyst stage. Magnified perinuclear regions are indicated by yellow boxes. Maximum intensity projection of confocal images of 80-μm in height are shown unless otherwise stated.

Figure 7

Fig. 7. Proteolytic activity of cathepsin B is maintained relatively at a low level in the ELYSA.
(A) GV and MII oocytes were stained in the same drop for LysoSensor Green and Magic Red cathepsin B staining analysis, and confocal images were acquired. (B) Fluorescent intensities of LysoSensor and Magic Red on the yellow lines indicated in (A) are indicated by a line plot. Small, filled dots indicate the origin (0 μm) in distance.

Supplementary Figure 1

Supplementary Fig. 1. Endosomes and lysosomes form giant structures in the periphery of the oocyte plasma membrane.
(A) MII oocytes were fixed and costained with anti-RAB7 and anti-LAMP1 antibodies. The schematic diagram indicates the location of oocyte chromosomes and the metaphase plate in the MII oocytes. Maximum intensity projection of confocal images of 80-μm in height are shown as Z-projection images. Arrowheads indicate the positions of oocyte chromosomes.

Supplementary Figure 2

Supplementary Fig. 2. Giant structures in MII oocytes were immunostained with anti-RAB7 and LAMP1 antibodies after glutaraldehyde (GA) fixation.
(A) MII oocytes were fixed with 0.025% GA and costained with anti-RAB5/RAB7 and anti-LAMP1 antibodies. Single-section images from confocal observation are indicated.

Supplementary Figure 3

A

Inversed (intact)   Inversed (transparent)

B

Supplementary Fig. 3. Toluidine blue-positive structures in MII oocytes show similar localization patterns to the ELYSA.

(A) Aligned serial toluidine blue-stained light microscopic images were stacked and three-dimensionally reconstructed by inversing the intensity and adjusting it to be transparent. (B) A series of images of toluidine blue-stained serial semi-thin sections of the Epon-embedded MII oocyte were reconstructed to 3D. Based on the images obtained, chromosomes were segmented blue, toluidine blue-positive structures were segmented yellow, and oocyte PM was segmented white.

Supplementary Figure 4

Supplementary Fig. 4. LysoTracker-positive punctate structures colocalized with V1A-mCherry.
Green and red fluorescence images of embryos expressing V1A-mCherry (mRNA injected at 3 h post-fertilization [hpf]) and stained with LysoTracker Green at the indicated time points were acquired. The maximum intensity projection of confocal images of 80-μm in height is indicated.

Supplementary Figure 5

Supplementary Fig. 5. Magic Red cathepsin B staining gave signals in the cytoplasm as well as the ELYSA in GV and MII oocytes..

Fluorescence images of oocytes and embryos at the indicated stages were obtained and stained with Magic Red Cathepsin B solution (and LysoTracker Green; not shown) in the same drop of medium. The average intensity projections of confocal images of 80-μm in height are shown.

Supplementary Movie 1

Supplementary Movie 1.
MII oocytes were fixed and costained with anti-RAB5 and anti-LAMP1 antibodies. A series of the confocal images of 81 slices of 80-μm in height is shown from top to bottom.

Supplementary Movie 2

Supplementary Movie 2.
A series of continuous toluidine blue-stained light microscopic images of same oocyte (top half) reconstructed to 3D. Chromosomes in the images obtained were segmented blue, toluidine blue-positive structures were segmented yellow, and oocyte PM was segmented white.

Supplementary Movie 3

Supplementary Movie 3.
Time-lapse movie of EGFP fluorescence in the GV oocytes injected with *Lamp1-EGFP* and *V1A-mCherry* mRNA during IVM with a maximum intensity Z-projection of confocal images of 80-μm in height. Magnified perinuclear regions in Fig. 5 are indicated by a box. The timestamp indicates the time after IVM started.

Supplementary Movie 4

Supplementary Movie 4.
Time-lapse movie of fertilized oocytes expressing V1A-mCherry mRNA in the presence of LysoSensor Green during embryonic development. The maximum intensity projection of confocal images of 80-μm in height is shown.