Auditory training alters the cortical representation of both learned and task irrelevant sound

features

Huriye Atilgan[1,2], Kerry M Walker[2], Andrew J. King[2], Jan W. Schnupp[2,3] and Jennifer K. Bizley[1,2]

[1] The Ear Institute, University College London, UK

[2] Department of Physiology, Anatomy and Genetics, University of Oxford, UK

[3] City University of Hong Kong, China

Corresponding Author: Jennifer Bizley j.bizley@ucl.ac.uk

UCL Ear Institute, 332 Gray's Inn Road, London, WC1X 8EE.

22
23

Abstract

Auditory learning is supported by long-term changes in the neural processing of sound. We mapped

neural sensitivity to timbre, pitch and location in animals trained to discriminate the identity of

artificial vowels based on their spectral timbre in a two-alternative forced choice (T2AFC, n=3,

female ferrets) or to detect changes in fundamental frequency or timbre of repeating artificial

vowels in a go/no-go task (n=2 female ferrets). Neural responses were recorded under anaesthesia

in two primary cortical fields and two tonotopically organised non-primary fields. Responses were

compared these data to that of naïve control animals. We observed that in both groups of trained

animals the overall sensitivity to sound timbre was reduced across three cortical fields but enhanced

in non-primary field PSF. Neural responses in trained animals were able to discriminate vowels that

differed in either their first or second formant frequency unlike control animals whose sensitivity

was mostly driven by changes in the second formant. Neural responses in the T2AFC animals, who

were required to generalise across pitch when discriminating timbre, became less modulated by

fundamental frequency, while those in the go/no-go animals were unchanged relative to controls.

Finally, both trained groups showed increased spatial sensitivity and altered tuning. Trained animals

showed an enhanced representation of the midline, where the speaker was located in the

experimental chamber. Overall, these results demonstrate training elicited widespread changes in

the way in which auditory cortical neurons represent complex sounds with changes in how both task

relevant and task-irrelevant features were represented.

Introduction

Sensory discrimination tasks are known to drive cortical plasticity, and increases in map area have

been proposed as providing the structural substrate for learning in auditory cortex (Rutkowski and

Weinberger, 2005; Schreiner and Polley, 2014). However, recent studies have questioned the

functional role of map plasticity suggesting that it may be a temporary phenomena associated with

learning that does not persist once a task is well-learned (Reed et al., 2011), and demonstrating that

auditory learning can occur in the absence of map plasticity (Galindo-Leon et al., 2009; Shepard et

al., 2016). Training animals to 'forage' for sounds with particular features leads to diverse changes

within primary auditory cortex independently of any changes in the tonotopic representation (Bao et

al., 2004; Whitton et al., 2014). Since one role for auditory cortex is thought to be in the formation

of behaviourally meaningful sound categories (Bizley and Cohen, 2013) a question of great interest is

to understand the cortical representation is changed in order to support the categorization of

spectrally overlapping naturalistic sounds.

Identifying auditory 'objects' requires that animal and human listeners are capable of discriminating

sounds along a given perceptual dimension while generalizing across variability in other dimensions

(Griffiths and Warren, 2004; Bizley and Cohen, 2013). At the level of the single neuron this requires

that neuronal responses are both selective for one sound feature but tolerant (or invariant) across

others (Ison and Quiroga, 2008; Bizley et al., 2009). While the consequences of behavioural training

on neural invariance is unknown, appropriate environmental exposure during development can

shape auditory cortical responses to complex sound features: In the auditory cortical neurons of

animals reared in complex acoustic environments fewer neurons respond to any single sound but

responses were more selective for particular spectro-temporal features and can tolerate greater

acoustic variability(Bao et al., 2013).

In this study we recorded from the auditory cortex of animals trained in one of two behavioural

paradigms that required that animals discriminate perceptual features of spectrally overlapping

artificial vowels. One set of animals was trained to discriminate the timbre of artificial vowels (Bizley

et al., 2013; Town et al., 2018) and did so across a range of fundamental frequencies (F0s). A second

set of animals were trained to detect changes in pitch and timbre of an ongoing sound sequence

(Walker et al., 2017).

After behavioral training was complete electrophysiological recordings were made from four

tonotopic auditory cortical fields. The perceptual features of complex sounds, such as their location

in space or spectral timbre, are distributed rather than systematically mapped within or across

auditory cortical fields (Bizley and Walker, 2010; Recanzone and Cohen, 2010; King and

Middlebrooks, 2011). Given this, and the broadband nature of the stimuli, it seemed unlikely that

learning could be supported by an expansion of the tonotopic map. We therefore sought to

determine how training altered (i) single neuron response sensitivity and selectivity to both learned

and task-irrelevant sound features, and (ii) whether training impacted on the distribution of neural

tuning across specific auditory cortical fields. We tested two hypotheses: Firstly, that training would

increase the sensitivity to trained features and the tolerance for changes in untrained features.

Secondly, consistent with hierarchical encoding theories, we hypothesized that training effects

would be strongest in non-primary areas. Our data demonstrate that the representation of both

trained and task-irrelevant features was changed by training, but that the direction of these changes

varied between cortical fields: increase in selectivity to the trained features occurred specifically in

the secondary posterior pseudosylvian field (PPF) whereas neurons in other fields became less

sensitive.

Materials & Methods

Animals

All animal procedures were approved by the local animal welfare and ethical review committee at

the University of Oxford and performed under license from the UK Home Office in accordance with

the Animal (Scientific Procedures) Act 1986. Five adult, female, pigmented ferrets (*Mustela putorius*)

were used in this study. Three of these animals experienced 1-2 years of training on a two

alternative forced-choice timbre discrimination task which required they report the identity of an

artificial vowel(Fig. 1C, T-2AFC; for the details of behavioral training see {Bizley, 2013 #925}). Two

animals were trained to perform a Go/No-go change detection task, where animals were presented

with a sequence of artificial vowels and had to report (in different sessions) either changes in F0 or

changes in vowel identity (Fig.1D, T/P-GNG; see {Walker, 2011 #93}).

Ferrets were housed in groups of either two or three, with free access to high-protein food pellets

and water bottles. On the day before behavioural training, water bottles were removed from the

home cages and were replaced on the last day of a training run. Training runs lasted for 5 days or

less, with at least 2 days between each run. On training days, ferrets received drinking water as

positive reinforcement while performing a sound discrimination task. Water consumption during

training was measured and supplemented as wet food in home cages at the end of the day to ensure

that each ferret received at least 60 ml of water per kilogram of body weight daily. Once behavioural

training was complete electrophysiological recordings were made under non-recovery anesthesia.

Recording under aneasthsia was necessary for the large scale mapping of neurons across cortical

fields, and in order to directly compare the resulting responses with data from control animals

previously collected under the same aneasthetic regime. Electrophysiological control data was

provided from 5 additional ferrets that formed a previously published dataset {Walker, 2011

#1485;Bizley, 2009 #607}. All animals were routinely monitored to ensure their ears were clean and

disease free throughout the study.

Stimuli

Acoustic stimuli for both behavioral testing and electrophysiology were artificial vowel sounds. For electrophysiological testing, sounds were all possible combinations of four F0 values: F0 = 200, 336, 565, and 951 Hz, four spectral timbres: /a/ (F1–F4 at 936, 1551, 2815, and 4290 Hz); /ɛ/ (730, 2058, 2979, and 4294 Hz); /u/ (460, 1105, 2735, and 4115 Hz); and /i/ (437, 2761, 3372, and 4352 Hz) and four spatial locations presented in virtual acoustic space: -45°, -15°, 15°, and 45° azimuth, at 0° elevation; giving a total of 64 sounds, each of which was 150 ms in duration. Additionally, noise bursts and pure tones were used to characterize individual units and to determine tonotopic gradients in order to confirm the cortical field in which any given recording was made {Bizley, 2005 #602}.

For animals trained in the 2AFC timbre identification task stimuli were /u/ and /ɛ/. Animals were trained initially with an F0 of 200 Hz, but tested across a range of values from 150 Hz – 500 Hz. For animals trained in the T/P GNG task reference sounds were the vowel /a/ at 200 Hz with F0 targets being the vowel /a/ with F0 values of 336, 565, and 951 Hz and timbre targets being the vowels /i/, /u/ and /ɛ/ presented with an F0 of 200 Hz.

Electrophysiological Recordings

Experimental methods were identical to those used in Bizley et al., 2009. Recordings were made under medetomidine/ketamine anaesthesia in the left auditory cortex and targeted at primary and non-primary tonotopic areas: primary auditory cortex (A1) and the anterior auditory field, (AAF), on the middle ectosylvian gyrus and the posterior pseudosylvian and posterior suprasylvian fields (PPF and PSF) located on the posterior ectosylvian gyrus (Fig. 2A). Recordings were made with silicon probe electrodes (Neuronexus Technologies, USA) either in a 16 x2 configuration (16 active sites spaced at 100μm intervals on each of two probes), a 32 x 1 configurations (50 μm spacing) or, in one animal, an 8 x 4 configuration (100μm spacing in depth, 200μm between shanks). Voltage signals were bandpass filtered (500-5000Hz), amplified (<20,000 times), and digitized at 25 kHz. Data acquisition and stimulus generation were performed using BrainWare (Tucker-Davis Technologies). A minimum of 80 units were recorded in each field in each dataset (mean = 118 units ± 28.32, Table 1). Data were combined across animals (control or trained) to make composite tonotopic maps (Fig. 1B). The distribution of CFs were not significantly different between trained and naïve animals. (2 way mixed ANOVA (2 groups x 4 fields), group: $F_{(1, 612)} = 0.02$, p=0.8792; field: $F_{(3,612)} = 39.79$, p<0.001; group×field: $F_{(3,612)} = 4.67$, p =0.08).

Behavioural Testing

Full details of the training apparatus and procedure for shaping animals can be obtained in {Bizley, 2013 #925;Walker, 2011 #93}. Briefly, water-restricted ferrets were positively conditioned to report the identity of a vowel sound (either /e/ or /i/) in a two-alternative forced choice task, or trained detect a change in the pitch or timbre of a repeating artificial vowel on a go/no-go task. In each case, the animal initiated each trial by inserting its nose in a poke hole situated at the center of the sound-isolated testing chamber. For the 2AFC task this resulted in the presentation of two repetitions of one of the vowel sounds and animals were rewarded for correctly responding at the side that was associated with that vowel. In the GNG task ferrets heard a sequence of artificial vowels which could change in identity or pitch at the third to seventh vowel in the sequence, and if ferrets withdrew from the nose poke hole during presentation of such a deviant, they were rewarded with water. Failures to withdraw to a deviant (within a 550 ms time window following deviant onset) resulted in a 12 s time out. In both tasks sounds were presented from a speaker located above the central 'go' spout at the animals midline.

Neural Data Analysis

Neural data was analysed using a variance decomposition approach developed in {Bizley, 2009 #607}. We first calculated spike counts for each of the 64 stimuli, averaged over repeated presentations of the same sound and binned with 20 ms resolution over the 300 ms immediately preceding stimulus onset. We then performed a 4-way ANOVA on the spike counts, where the 3 stimulus parameters (azimuth, pitch, and timbre) plus the time bin served as factors. To quantify the relative strength with which one of the three stimulus dimensions influenced the firing of a particular unit, we calculated the proportion of variance explained by each of azimuth, pitch, and timbre, Var_{stim}, as:

$$\text{Var}_{\text{stim}} = \frac{\text{SS}_{\text{stim.bin}} - \text{SS}_{\text{error}} \cdot \text{df}_{\text{stim.bin}}}{\text{SS}_{\text{total}} - \text{SS}_{\text{bin}}}$$

where "stim" refers to the stimulus parameter of interest (pitch, timbre, or azimuth), $\text{SS}_{\text{stim.bin}}$ is the sum of squares for the interaction of the stimulus parameter and time bin, SS_{error} is the sum of squares of the error term, $\text{df}_{\text{stim.bin}}$ refers to the degrees of freedom for the stimulus x time bin interaction, SS_{total} is the total sum of squares, and SS_{bin} is the sum of squares for the time bin factor. A significant SS_{bin} reflects the fact that the response rate was not flat over the duration of the 300 ms response window and by examining the stimulus-by- time-bin interactions, we were able to test the statistical significance of the influence a given stimulus parameter had on the temporal discharge

186 pattern of the response. Subtracting the $SS_{error} . df_{stim .bin}$ from the $SS_{stim .bin}$ term allows us to

187 calculate the proportion of response variance attributable to each of the stimuli, taking into account

188 the additional variance explained simply by adding extra parameters to the model. As in our previous

189 work, we considered a main effect or interaction term in the ANOVA to be statistically significant if it

190 exceeded $p < 0.001$.

Experimental Design and Data Analysis

191
192

Data from 5 control animals and 5 experimental animals were used in this experiment. This allowed

us to have a sufficient sample size of recordings from each cortical field (see Table 1) and to match the

frequency tuning distribution of our samples.

For statistical comparison of neural tuning measures (derived using the variance decomposition

approach described above) we used Analysis of Variance, with the specific analysis for each test being

reported in the Results section. Generalised Linear Mixed Models were fit using maximum likelihood

approaches with model selection performed using Akaike Information Criteria values. Where

appropriate Bonferoni corrected post-hoc comparisons were applied (p<0.05). The specific analysis

for each experimental question is detailed in full in the Results section.

Code and data availability

202
203

204 All code is available on Github: https://github.com/huriyeatg/trainingInducedPlasticity

205 Data are available on the following repository [link added on publication].

206 **Table 1**: Total number of recordings (probe placements and units) in each field for 5 control animals
207 and 5 trained animals.

	A1	AAF	PPF	PSF
Recordings from Control Animals				
Probe Placement(n)	10	7	11	7
Units (n)	189	101	152	96
Recordings from Trained Animals				
Probe Placement(n)	14	9	7	11
Units (n)	133	80	115	145

Results

Three ferrets were trained in a two alternative forced choice timbre discrimination (T-2AFC) task to

discriminate /u/ from /ɛ/, across a range of F0s (Fig.1C). Two ferrets were trained in a go/no-go task

to detect changes in timbre or F0 (T/P-GNG) from a repeating reference vowel (/a/ F0 = 200 Hz, Fig.1D).

Once behavioural training and testing was complete we recorded neural activity under

medetomidine/ketamine anesthesia, which allowed us to map neural responses across the cortical

surface of multiple cortical fields in each animal, and directly compare these data to those obtained

in naïve animals in a previous study that constitute the control data for this investigation. The

responses of 713 units (459 single neurons, 254 small clusters of units) which were responsive (paired

t-test on sound-evoked and spontaneous firing rates, *p<0.05*) to vowels were recorded from four

tonotopic auditory cortical fields (see Table 1). There were no systematic differences between the

response properties of single units and small unit clusters therefore the term 'units' will be used to

refer to both except when explicitly stated.

Figure 1:

A, Location of ferret auditory cortical fields and their tonotopic organisation (adapted from (Nelken

et al., 2004)). Recordings in this study targeted A1, AAF, PPF and PSF. Field boundaries are marked with dotted lines, and the pseudosylvian sulcus (pss) and suprasylvian sulcus (sss) are drawn as solid lines. **B**, Voronoi tessellation map showing the CFs of all unit recordings made in control animals (478 units, 5 animals) and trained animals (456 units, 5 animals). Tiles represent a recording site and are coloured according to the characteristic frequency (CF) of the unit recorded there. **C**, Schematic illustration of the timbre two-alternative forced choice paradigm (C, T-2AFC) and F0 / timbre change detection paradigm (D, T/P-GNG).

Response modulation by timbre, pitch and azimuth

In order to test our first hypothesis, that we would see an increase in both sensitivity and invariance in control animals, we determined the proportion of units whose responses were significantly modulated by variation in stimulus location, pitch (determined by fundamental frequency, F0) and timbre, using the variance decomposition approach used in Bizley et al. (2009). Since all five animals trained in a timbre discrimination task, we predicted a greater number of neurons might convey timbre information, and that we might observe fewer neurons that were additionally sensitive to untrained stimulus features (pitch in the 2AFC animals and azimuth in 2AFC and T/P GNG animals). However, contrary to these predictions, the number of units that showed timbre sensitivity was equivalent between trained and control groups (391/713 units and 293/538 units respectively, X^2 = 0.005, p=0.95). In both datasets units with joint stimulus sensitivity outnumbered those with sensitivity to only a single parameter, but the distribution of sensitivity to zero, one, two or three stimulus parameters was significantly different between the groups (X^2 =86.9, p<0.001). In the trained dataset many more units showed significant modulation by all three stimulus parameters, or by none of the stimulus parameters, than in the control dataset where most units were modulated by one or two stimulus dimensions. Taken together this suggests that training resulted in a relative increase in the number of units either insensitive to modulations in pitch, timbre or space, or sensitive to all three dimensions. We now consider sensitivity to each feature in turn, asking how training changes the responses of single units, and the cortical distribution of sensitivity.

Table 2: Percentages of recordings modulated by stimuli dimensions (azimuth, F0 and timbre) for naïve, T-2AFC trained and T/P-GNG trained animals

	Control Animals	T-2AFC trained	T/P-GNG trained
One Stimulus	**24% (120)**	**16% (103)**	**14% (11)**
Only Azimuth	*5% (29)*	*4% (24)*	*3% (3)*
Only F0	*6% (30)*	*7% (47)*	*6% (6)*
Only Timbre	*13% (71)*	*5% (32)*	*5% (5)*
Two Stimuli	**38% (203)**	**16% (92)**	**31% (32)**
Azimuth-F0	*2% (10)*	*6% (30)*	*22% (23)*
Azimuth-Timbre	*7% (35)*	*5% (26)*	*6% (6)*
F0-Timbre	*29% (158)*	*6% (36)*	*3% (3)*
Three Stimuli	**30% (159)**	**44% (272)**	**11% (11)**
Not significant	**13% (70)**	**24% (146)**	**34% (36)**
Total Units	538	607	106

Neural sensitivity to trained stimulus features: Timbre

For each unit we determined what proportion of the response variance was attributable to each of
the three stimulus dimensions, and their combinations. We start by considering sensitivity to timbre,
as both behavioural tasks required that animals discriminated this feature (Fig. 2A, T-2AFC animals,
Fig. 2B, T-GNG). Figure 2C shows the distribution of sensitivity to timbre (i.e. the proportion of variance
explained by timbre) across the four cortical fields examined and illustrates that while the proportion
of units with significant timbre sensitivity is equivalent across trained and control groups, the
magnitude of this sensitivity is substantially *lower* in the trained animals in three out of the four
cortical fields examined. This observation was confirmed with a two-way ANOVA (factors: group
(control, T-2AFC, T/P-GNG), and cortical field (A1, AAF, PPF, PSF), dependent variable: proportion of
variance explained by timbre) which showed a significant effect of group ($F_{(2, 1224)} = 74.48$, $p < 0.001$)
and field ($F_{(3, 1224)} = 7.17$, $p < 0.001$) and a significant group $\times$ field interaction ($F_{(6, 1224)} = 11.17$,
$p < 0.001$). In control animals, sensitivity to timbre was highest in the primary fields A1 and AAF. In
contrast in the T-2AFC animals timbre sensitivity was substantially higher in PPF than in any of the
other three areas. Tukey–Kramer post hoc comparisons between trained and untrained animals (*p*
<0.05) revealed lower timbre sensitivity in the primary areas (A1 and AAF) in T-2AFC and T/P-GNG
trained animals than in controls, but higher sensitivity in PPF in T-2AFC trained animals (Fig.2C,D).
Therefore, we conclude that long-term training in a spectral timbre discrimination task leads to a field-
specific increase in timbre sensitivity in field PPF which is accompanied by a more general decrease
timbre sensitivity in A1, AAF and PSF.

Figure 2: Timbre sensitivity is reduced overall, but enhanced in field PPF, of trained animals.

A, performance of three ferrets discriminating timbre across changes in F0 (chance = 50%). **B** performance of two animals detecting changes in timbre (chance = 25%). **C**, cortical distribution of sensitivity to timbre measured using the Proportion of variance explained metric (see methods). Each tile represents an electrode penetration, with individual sites averaged. **D**, swarm plots showing the distribution of timbre sensitivity across fields and groups. Each datapoint is a unit. The dot indicates a significant main effect of field (p<0.001), the star indicates a significant main effect of training (p<0.001), and a significant effect of field*training interaction is indicated with both symbols together (p<0.001). **E**, first and second formant frequencies for the vowels used in the study, **F**, difference values in F1 or F2 space for given vowel pairs **G,** Neural discriminability for vowel pairs estimated by calculating the proportion of neural response variance attributable to changes in timbre for subsets of the data comprising only those vowels. Boxplots illustrate the median and upper and lower quartile values of the data. **H**, as G, but broken down by cortical field.

Formant cues are reweighted in timbre-trained animals

Contrary to our expectation, long-term training on a spectral timbre discrimination task apparently
led to an overall decrease in neural sensitivity to timbre. However, the stimuli with which we recorded
neural responses comprised both trained and untrained sounds. To contrast responses to both types
of vowels, we therefore repeated the variance decomposition analysis using the neural responses to
pairs of vowels (i.e. subsets of 50% of the data comprising the responses to stimulus combinations of
two vowels, four F0s and four locations, 32 in total) with the proportion of variance explained by
timbre in each case now providing an estimate of how well neuronal responses differentiate a given
pair of vowels across variation in F0 and space. We hypothesized that if training led to enhanced
selectivity for the target vowels we would observed the highest timbre sensitivity measures for these:
for the T2AFC animals /u/ versus /e/ should yield the greatest timbre sensitivity, whereas for the T/P-
GNG animals, where /a/ was the reference vowel, we might expect the /a/-/i/, /a/-/e/, and /a/-/u/
contrasts to yield higher sensitivity measures than the pairs of vowels that did not include /a/.

Figure 2E shows the proportion of variance explained by timbre for the six possible vowel
combinations, organised according to the magnitude of the difference in second formant frequency
(ΔF2). In control animals the neural sensitivity to spectral timbre was well predicted by the size of the
difference in F2 frequency. In contrast, the pattern of timbre sensitivity in the T-2AFC animals was
neither dominated by the F2 difference, nor by the identity of the trained vowel pair. Instead the pairs
with lowest discriminability were those with the smallest difference in the frequency of the first
formants (sitting to the left of the space defined by ΔF1 and ΔF2 in Fig.2F).

To better understand how training group, cortical field and change in first and second formant
frequencies determined the proportion of the neural response variance was attributed to a change in
timbre we ran GLMM which predicted the neural response variance for every pair of vowels with
factors field, training group, delta F1 and delta F2 (where delta F1 and F2 were calculated as the
difference in first and second formant frequencies for the relevant vowel pair), with unit ID as a
random effect. The model was fitted using maximum likelihood estimation and the AIC was used to
determine the best model. Within this model both delta F1 and delta F2 were significant predictors,
and there was a significant interaction between training group and delta-F1 (but not training group
and delta F2) consistent with the idea that training led to an increased integration of both formant
frequencies (see Table 1 for full model). From this we conclude that training not only redistributed
timbre sensitivity within auditory cortex, but caused a reweighting of spectral integration from being
F2 dominated to incorporating differences in both F1 and F2.

Sensitivity to fundamental frequency

T-2AFC animals were able to identify trained vowels across a range of F0 values (Fig 3A) and we

therefore predicted that we might see a decreased sensitivity (i.e. increased tolerance) across changes

in F0. In contrast, the T/P-GNG animals were never required to discriminate one stimulus dimension

across changes in the other, but were trained to detect changes in either F0 (Fig 3B) or spectral timbre

(Fig 2B). When sensitivity to F0 is plotted across the cortical surface, or broken down by cortical field

it is apparent that compared to control data both trained groups show decreased F0 sensitivity. A

two-way ANOVA was performed determine the effect of group and cortical field. There was a

significant effect of group (F (2,1224) = 27.76, p<0.001) and a significant training x field interaction (F

(6,1224)= 5.02, p<0.001). Tukey–Kramer post hoc comparisons (*p* <0.05) on the responses recorded

in each of the cortical field across groups revealed lower F0 sensitivity in A1 of T-2AFC trained animals

compared to control animals (Fig.3C,D), with T/P-GNG animals being statistically indistinguishable

from controls.

Figure 3: Sensitivity to F0 is decreased in trained animals

A Behavioural performance for discriminating timbre across fundamental frequency (F0). Symbols show individual animals. **B** F0 change detection performance for two animals. **C** Voronoi tessellation maps and box plots plotting the proportion of variance explained by F0 for control and trained animals**.** Conventions as figure 2 C. **D** Swarm plots showing the distribution of F0 sensitivity across fields and training groups. Symbols (black star and dot) show ANOVA results idicating significant main effect of field (p<0.001), training (p<0.001) and a significant effect of field*training interaction.

Sensitivity to task-irrelevant features: Space

Finally we examined how training impacted sensitivity to variation in sound source location. In our control dataset we observed that the spatial tuning elicited by these vowel stimuli presented in virtual acoustic space was modest, and as anticipated, predominantly contralateral (Bizley et al., 2009). When the same neurons were tested with spatially modulated broadband noise (also in VAS) we observed considerably greater spatial modulation leading us to suggest that the low spatial sensitivity was a product of the stimuli rather than the neurons we were recording or the VAS technique. We had speculated that exposure to these sounds might enhance spatial sensitivity measured in auditory cortex using these stimuli. As Figure 4 show spatial sensitivity was indeed higher in trained animals: whether visualized as a cortical map (Fig.4A) or broken down by cortical field (Fig.4B), it is clear that in the non-primary areas spatial sensitivity was higher in the T-2AFC animals than in controls. A two-way ANOVA on the proportion of variance explained by space, with group and cortical field as factors, determined significant effects of group (F(2,1224) = 10.63, p<001) and field (F(3,1224)= 3.45, p=0.01) and a significant training × field interaction (F(6,1224)= 4.22, p<0.001). Tukey–Kramer post hoc comparisons (*p* <0.05) of the trained and control data in specific auditory fields revealed higher azimuth sensitivity in the non-primary areas (PPF and PSF) in trained animals compared to control animals.

Figure 4: Effect of training on the neural encoding of task-irrelevant features

A, Voronoi tessellation maps plotting the proportion of variance explained by azimuth for control
and trained animals. *B*, Swarm plots showing the proportion of variance explained by azimuth across
fields and training groups. The dot indicates a significant main effect of field (p<0.001), the star
indicates a significant main effect of training (p<0.001), and a significant effect of field*training
interaction is indicated with both symbols together (p<0.001). *C-E,* Population spatial tuning
functions (mean ± SEM for all responsive units) for control, T2AFC animals and T/P GNG animals. F
beta coefficients for the impact of model parameters on spike rate.
397

We next asked whether the increased spatial sensitivity in trained animals reflected altered spatial

tuning, or simply enhanced the gain of spatial receptive fields. To assess this we exploited the typical

contralateral spatial preference of most auditory cortical neurons to derive population tuning

functions. We calculated a normalised spatial response function for each neuron, taken as the mean

sound-evoked spike rate at each location across all pitch and timbre combinations, and then averaged

across all recorded units (Fig.4C-E). As expected, in the control animals (Fig.4C) this yielded a

monotonically increasing function with the most contralateral (+45°) location eliciting the strongest

firing rates. In contrast, in both trained groups of animals, spatial tuning was non-monotonic and

showed a peak at +15°. To quantify these effects we ran a GLMM predicting spike rates with group

(T-2AFC or T/P-GNG relative to reference category control), field (reference category A1), spatial

position (categorical predictor, relative to -45) with unit as a random effect. This confirmed significant

main effects of spatial position (-45 and +45 significantly different, reflecting contralateral tuning), and

significant field*group*position interactions showing for fields PPF and PSF with both trained groups.

The coefficients for these effects (Fig.4F) were negative for +45 and positive for ±15, confirming the

changes visible in the spatial tuning functions plotted in (Fig.4C-E). Thus despite the sound source

never varying in its spatial position, and not being relevant to the task, the location from which the

sounds were presented (i.e. central space) was substantially enhanced in the trained animals.

Discussion

In this study we trained two groups of ferrets to discriminate perceptual attributes of artificial vowels.

One group categorized vowels according to their identity across changes in F0 (T-2AFC animals), while

the other detected changes in either the F0 or timbre in a sequence of on-going vowel sounds (T/P

GNG animals). We predicted that the 2AFC animals would show enhanced sensitivity for timbre and

increased tolerance (i.e. decreased sensitivity) to other sound features. In fact, what we observed was

more complex: sensitivity to timbre decreased markedly in three cortical areas but was enhanced in

field PPF. Moreover, sensitivity to vowel identity became less contingent only on changes in the

frequency of the second formant, and instead was dependent on both changes in first and second

formant frequency. Sensitivity to F0, which the animals were required to generalize across, decreased. In contrast, sensitivity to space, which had no task-relevance, was enhanced, with spatial receptive fields shifted towards the midline, from where target sounds originated. In the T/P GNG animals our conclusions are somewhat limited by a smaller sample size, but there was also a decrease in timbre sensitivity relative to controls and an increase in spatial sensitivity, with F0 encoding matching that of control animals.

Previous studies investigating the impact of training on neural tuning in auditory cortex have focused on map plasticity in A1. However, higher auditory cortical fields are thought to become increasingly specialized for processing spatial or non-spatial stimulus attributes (Rauschecker and Tian, 2000; Bizley and Cohen, 2013; Elgueda et al., 2019) and neurons in higher cortical fields show enhanced attention related changes during behavior (Mesgarani and Chang, 2012; Atiani et al., 2014; Elgueda et al., 2019). It is currently unknown how these attention-related changes relate to more 'hard-wired' changes in neural sensitivity. Our data bridge this gap by suggesting that the areas that show larger attention related changes may also be those in which receptive fields are optimized through learning to process task-relevant stimuli.

The observation of a marked decrease in sensitivity to task-relevant features in the primary auditory cortical fields A1 and AAF is a potentially surprising finding. However, a number of studies report that engagement in a behavioural task causes suppression of neural responses in auditory cortex (Otazu et al., 2009; Town et al., 2018). One possibility is that training leads to the integration of diverse non-sensory inputs into auditory cortex that ultimately underlie the observation of choice or motor related activity, but that these inputs come at the cost of presumably redundant feed-forward stimulus evoked activity. Cooling primary auditory cortex in ferrets does not lead to an impairment in a vowel discrimination in silence in a task analogous to the one here (Town et al., 2023). Our neural data raise the testable prediction that inactivation of field PPF should cause a timbre identification deficit whereas cooling A1, AAF or PSF would have a more modest effect on behavior.

Calculating discriminability measures for pairs of vowels allowed us to observe that training on a spectral timbre task altered the way in which neuronal responses were sensitive to spectral timbre. In the naïve control animals the most discriminable stimuli were those in which there was a large difference in second formant frequency. This finding is mirrored behaviourally; when first and second formant cues are placed in conflict, animals tend to weight the position of the second formant over the first, with behaviour being best predicted by either F2 position or the position of the spectral centroid (Town et al., 2015). Nonetheless, the animals whose neural data were recorded for this study were also tested behaviourally with single-formant stimuli and these animals accurately classify F1 for

/u/ and F2 for /e/ (Bizley et al., 2013). The finding that neural responses in trained animals are

explained by both changes first and second formants for both trained and novel vowels suggest that

learning results in an enhanced integration of the cues that define the spectral envelope.

We also considered the impact of training on the neural representation of two task-irrelevant

features. In the T-2AFC task animals were required to generalise across F0 while discriminating timbre,

and neural responses showed an overall decrease in sensitivity to F0. In contrast, the T/P-GNG animals

were required, in separate sessions, to detect changes in timbre or F0 of a sequence of vowels and

showed F0 sensitivity that was statistically indistinguishable from controls. It remains possible that, as

with timbre in the 2AFC animals, this might reflect a reorganization such that there is a general

decrease in sensitivity that is accompanied by a local increase in F0 sensitivity in another cortical area

in one or both sets of animals. This possibility requires further investigation, possibly with high-density

recordings targeting the low frequency border of primary and posterior fields where specialization for

pitch might occur (Walker et al., in prep).

In contrast to the decreased sensitivity to non-spatial sound features we observed a marked increase

in spatial sensitivity in trained animals despite the task having no spatial component to it. These

changes occurred principally in the non-primary fields PPF and PSF and were not simply gain changes

but rather (presumably) adaptive shifts in tuning: the population tuning function shifted from

monotonically increasing for more contralateral sounds to peaking 15° contralateral to the midline.

Our spatial receptive fields were very coarsely measured with stimuli at ±15° and ±45° but if the

change in tuning observed in the recorded hemisphere were mirrored across the midline we would

expect to see that the representation of the midline, where stimuli were presented during

behavioural testing, was enhanced. Therefore it seems that repeated exposure to behaviourally

relevant sounds from this location led to an enhancement in non-primary auditory cortex (fields PPF

and PSF). Engaging in a sound discrimination task has been shown to refine spatial tuning in primary

auditory cortex, with changes occurring for both localisation tasks, and – more modestly – for non-

spatial tasks (Lee and Middlebrooks, 2011) but to our knowledge this is the first report of enhanced

location coding after repeated exposure to behaviourally relevant stimuli. Learning triggers

widespread changes in gene expression in auditory cortex (Graham et al., 2023). Future work can

seek to unpicking the specific molecular mechanisms which ultimately support the changes in

auditory cortical function, and ultimately auditory memory, that we observe here.

The stimuli that the animals were processing were relatively low frequency (<4 kHz). Our sampling

yielded balanced samples of neurons across the frequency axis in both groups of animals (Fig.1) but

we did not perform the high-resolution mapping that would probably be required to unequivocally

argue that there was no difference in the tonotopic organisation in these animals. Nonetheless, we

did not see any evidence for map reorganisation in these animals and our sampling of the cortical

surface did not yield a dominance of low frequency recording sites. Given evidence that map

plasticity may be a temporary phase of learning (Reed et al., 2011), and that learning of natural

sounds can occur in the absence of any map plasticity (Shepard et al., 2016) the lack of evidence for

tonotopic map plasticity here is perhaps not surprising and likely additionally reflects the broad band

nature of the sounds that the animals were exposed to.

A caveat of our study is that it is performed under anaesthesia. This was essential for performing

mapping across multiple cortical fields. Methods for recording in awake animals currently allow only

high density sampling from a small area of cortex, or sparse sampling across multiple fields.

Recording under anaesthesia does, however, allow us to separate out effects of sensitivity to

stimulus features from attention and allows us to measure static receptive field features to which

attentional and task related effects are likely added during active listening.

In summary, training causes diverse effects in auditory cortex: sensitivity to trained stimulus features

which is broadly distributed in naïve animals becomes localised to one distinct auditory field after

training. In contrast to control animals which strongly weight second formant frequency, receptive

fields integrate information about both first and second formant frequency. Finally, sensitivity to

task-orthogonal features – here auditory space – is enhanced when stimuli consistently originated

from a single location.

Acknowledgments

This work was supported by a RNID (formally Action on Hearing Loss) PhD studentship to H.A.,

grants from the Biotechnology and Biological Sciences Research Council (to J.W.H.S., J.K.B.,

and A.J.K. and to J.K.B. BB/H016813/1), a Royal Society Dorothy Hodgkin Fellowship (J.K.B.), a

Sir Henry Dale Fellowship (J.K.B. WT098418MA) and a Wellcome Trust Principal Research

Fellowship (A.J.K.).

References

Atiani S, David SV, Elgueda D, Locastro M, Radtke-Schuller S, Shamma SA, Fritz JB (2014) Emergent
selectivity for task-relevant stimuli in higher-order auditory cortex. Neuron 82:486–499.

Bao S, Chang EF, Teng CL, Heiser MA, Merzenich MM (2013) Emergent categorical representation of
natural, complex sounds resulting from the early post-natal sound environment.
Neuroscience 248:30–42.

Bao S, Chang EF, Woods J, Merzenich MM (2004) Temporal plasticity in the primary auditory cortex
induced by operant perceptual learning. Nat Neurosci 7:974–981.

Bizley JK, Cohen YE (2013) The what, where and how of auditory-object perception. Nat Rev
Neurosci 14:693–707.

Bizley JK, Walker KM (2010) Sensitivity and selectivity of neurons in auditory cortex to the pitch,
timbre, and location of sounds. Neuroscientist 16:453–469.

Bizley JK, Walker KM, King AJ, Schnupp JW (2013) Spectral timbre perception in ferrets:
discrimination of artificial vowels under different listening conditions. J Acoust Soc Am
133:365–376.

Bizley JK, Walker KM, Silverman BW, King AJ, Schnupp JW (2009) Interdependent encoding of pitch,
timbre, and spatial location in auditory cortex. J Neurosci 29:2064–2075.

Elgueda D, Duque D, Radtke-Schuller S, Yin P, David SV, Shamma SA, Fritz JB (2019) State-dependent
encoding of sound and behavioral meaning in a tertiary region of the ferret auditory cortex.
Nat Neurosci.

Galindo-Leon EE, Lin FG, Liu RC (2009) Inhibitory plasticity in a lateral band improves cortical
detection of natural vocalizations. Neuron 62:705–716.

Graham G, Chimenti MS, Knudtson KL, Grenard DN, Co L, Sumner M, Tchou T, Bieszczad KM (2023)
Learning induces unique transcriptional landscapes in the auditory cortex. Hear Res
438:108878.

Griffiths TD, Warren JD (2004) What is an auditory object? Nat Rev Neurosci 5:887–892.

Ison MJ, Quiroga RQ (2008) Selectivity and invariance for visual object perception. Front Biosci
13:4889–4903.

King AJ, Middlebrooks JC (2011) Cortical Representation of Auditory Space. In: The Auditory Cortex,
pp 329–341. Springer.

Lee CC, Middlebrooks JC (2011) Auditory cortex spatial sensitivity sharpens during task performance.
Nat Neurosci 14:108–114.

Mesgarani N, Chang EF (2012) Selective cortical representation of attended speaker in multi-talker
speech perception. Nature 485:233–236.

Nelken I, Bizley JK, Nodal FR, Ahmed B, Schnupp JW, King AJ (2004) Large-scale organization of ferret
auditory cortex revealed using continuous acquisition of intrinsic optical signals. J
Neurophysiol 92:2574–2588.

Otazu GH, Tai LH, Yang Y, Zador AM (2009) Engaging in an auditory task suppresses responses in
auditory cortex. Nat Neurosci 12:646–654.

Rauschecker JP, Tian B (2000) Mechanisms and streams for processing of "what" and "where" in
auditory cortex. Proc Natl Acad Sci U S A 97:11800–11806.

Recanzone GH, Cohen YE (2010) Serial and parallel processing in the primate auditory cortex
revisited. Behav Brain Res 206:1–7.

Reed A, Riley J, Carraway R, Carrasco A, Perez C, Jakkamsetti V, Kilgard MP (2011) Cortical map
plasticity improves learning but is not necessary for improved performance. Neuron 70:121–
131.

Rutkowski RG, Weinberger NM (2005) Encoding of learned importance of sound by magnitude of
representational area in primary auditory cortex. Proc Natl Acad Sci U S A 102:13664–13669.

Schreiner CE, Polley DB (2014) Auditory map plasticity: diversity in causes and consequences. Curr
Opin Neurobiol 24:143–156.

Shepard KN, Chong KK, Liu RC (2016) Contrast Enhancement without Transient Map Expansion for
Species-Specific Vocalizations in Core Auditory Cortex during Learning. eNeuro 3.

Town SM, Atilgan H, Wood KC, Bizley JK (2015) The role of spectral cues in timbre discrimination by
ferrets and humans. J Acoust Soc Am 137:2870–2883.

Town SM, Poole KC, Wood KC, Bizley JK (2023) Reversible Inactivation of Ferret Auditory Cortex
Impairs Spatial and Nonspatial Hearing. J Neurosci 43:749–763.

Town SM, Wood KC, Bizley JK (2018) Sound identity is represented robustly in auditory cortex during
perceptual constancy. Nature communications 9:4786.

Walker KM, Davies A, Bizley JK, Schnupp JW, King AJ (2017) Pitch discrimination performance of
ferrets and humans on a go/no-go task. Animal Behavior and Cognition. Available at:
http://biorxiv.org/lookup/doi/10.1101/165852 [Accessed December 29, 2023].

Whitton JP, Hancock KE, Polley DB (2014) Immersive audiomotor game play enhances neural and
perceptual salience of weak signals in noise. Proc Natl Acad Sci U S A 111:E2606-15.