1 **Title:**
2
3 **Pannexin 1 Channels Control Cardiomyocyte Metabolism and Neutrophil Recruitment During**
4 **Non-Ischemic Heart Failure**
5
6

7 Caitlin M. Pavelec[1,3], Alexander P. Young[2,3], Hannah L. Luviano[1], Emily E. Orrell[1], Nabin Poudel[4],
8 Abigail G. Wolpe[5], Samantha H. Thomas[1], Scott Yeudall[1], Clint M. Upchurch[1], Mark D. Okusa[4],
9 Brant E. Isakson[3,7], Matthew J. Wolf[2,3], Norbert Leitinger[1,3]
10

Affiliations
11
1. Department of Pharmacology, University of Virginia School of Medicine, Charlottesville,
VA
2. Cardiovascular Medicine, Department of Medicine, University of Virginia School of
Medicine, Charlottesville, VA
3. Robert M. Berne Cardiovascular Research Center, University of Virginia School of
Medicine, Charlottesville, VA
4. Division of Nephrology and Center for Immunity, Inflammation and Regenerative
Medicine, University of Virginia School of Medicine, Charlottesville, VA
5. Department of Cell Biology, University of Virginia School of Medicine, Charlottesville, VA
6. Department of Molecular Physiology and Biophysics, University of Virginia School of
Medicine, Charlottesville, VA

Funding:
This work was supported by NIH P01 HL120840 (to N.L., B.E.I.), NIH T32 HL007284 (to C.M.P.), NIH
F31HL165918-01A1 (to C.M.P.), NIH F31 HL149221 to (to A.G.W.), NIH R01 HL137112 (to B.E.I.),
NIH F30 HL154554-01 (to S.Y.), NIH T32 GM007055 (to S.Y., C.M.U.), NIH T32 GM007267 (to S.Y.),
NIH R01 HL162925 (to M.J.W.),NIH R01 DK085259 (to M.D.O.), NIH R01 DK123248 (to M.D.O.),
the summer research internship program at the University of Virginia (to S.H.T), University of
Virginia Office of Citizen Scholar Development Double Hoo Award (to H.L.L., to C.M.P.).
30

Acknowledgements:
We thank the members of the Pannexin Interest Group for their input and expertise throughout
the project. We thank the Cardiovascular Research Center histology core, and the Flow Cytometry
Core Facility at the University of Virginia School of Medicine for their expertise and resources.

<u>Abstract:</u>

Pannexin 1 (PANX1), a ubiquitously expressed ATP release membrane channel, has been

shown to play a role in inflammation, blood pressure regulation, and myocardial infarction.

However, a possible role of PANX1 in cardiomyocytes in the progression of heart failure has not

yet been investigated. We generated a novel mouse line with constitutive deletion of PANX1 in

cardiomyocytes (Panx1^{MyHC6}). PANX1 deletion in cardiomyocytes had no effect on unstressed

heart function but increased the glycolytic metabolism both *in vivo* and *in vitro*. *In vitro*, treatment

of H9c2 cardiomyocytes with isoproterenol led to PANX1-dependent release of ATP and Yo-Pro-1

uptake, as assessed by pharmacological blockade with spironolactone and siRNA-mediated

knock-down of PANX1. To investigate non-ischemic heart failure and the preceding cardiac

hypertrophy we administered isoproterenol, and we demonstrate that Panx1^{MyHC6} mice were

protected from systolic and diastolic left ventricle volume increases and cardiomyocyte

hypertrophy. Moreover, we found that Panx1^{MyHC6} mice showed decreased isoproterenol-induced

recruitment of immune cells (CD45$^+$), particularly neutrophils (CD11b$^+$, Ly6g$^+$), to the

myocardium. Together these data demonstrate that PANX1 deficiency in cardiomyocytes impacts

glycolytic metabolism and protects against cardiac hypertrophy in non-ischemic heart failure at

least in part by reducing immune cell recruitment. Our study implies PANX1 channel inhibition as

a therapeutic approach to ameliorate cardiac dysfunction in heart failure patients.

    <u>Introduction</u>

Heart failure, which affects five million Americans each year and has a five-year survival

rate of less than 50 percent[1], can be divided in to two sub-pathologies: heart failure with reduced

ejection fraction (HFrEF) and heart failure with preserved ejection fraction (HFpEF); HFpEF is the

more predominant form and its prevalence is on the rise[1,2]. While several effective therapies are

available for patients with HFrEF, HFpEF has limited treatment options underscoring the need to

identify new therapeutic strategies[3]. Pathological features of heart failure include cardiac

hypertrophy, chronic inflammation, and interstitial fibrosis of the heart muscle, which together

result in reduced cardiac contractile function[4]. Current pharmacologic interventions for HFrEF

include treatments with beta-blockers, loop diuretics, angiotensin converting enzyme (ACE)

inhibitors, angiotensin receptor blockers, and sodium glucose transporter 2 (SGLT2) inhibitors[4–6].

These therapies target the renin-angiotensin/ aldosterone systems and the adrenergic signaling

axes, providing more symptom management than improving function of the cardiac muscle[5].

Moreover, previous clinical trials that have targeted TNF-α signaling, modulated the immune

system using intravenous immunoglobulins[7], or decreased fibroblast activation with pirfenidone[8]

(an antifibrotic which down regulates growth factors and procollagens I and II), have not lead to

significant improvements in patient outcomes. To prevent cardiac muscle decline, it is necessary

to devise novel therapeutic approaches which directly target the mechanisms of heart failure

progression.

Proper contractile function of the heart requires, at the level of the cardiomyocyte,

sufficient cellular energy in the form of accessible ATP[9,10]. Adult cardiomyocytes rely on fatty acids

as their main fuel source to generate ATP, but they are metabolically flexible and can use glucose,

branched-chain amino acids, or ketones as needed[11]. During heart failure, cardiomyocytes

demonstrate an increased demand for ATP due to more frequent but less effective contraction[12].

In order to meet this increased energy demand, cardiomyocytes shift their metabolism from

utilizing fatty acids to utilizing glucose, which is accompanied by increased glycolysis[13], which

resembles the reliance on glucose and glycolysis in fetal cardiomyocytes[11,14]. Moreover, recent

work has demonstrated that decreased reliance on fatty acid oxidation confers greater resistance

to ischemia reperfusion injury and promotes cardiomyocyte proliferation in mice deficient in the

mitochondrial fatty acid transporter *Cpt1b*[15]. These findings suggest that a metabolic shift to

glucose utilization by cardiomyocytes protects against cardiac injury.

In response to local hypoxia or chronic beta-adrenergic stimulation, two conditions known

to cause cardiac hypertrophy and heart failure, cardiomyocytes release damage signals such as

ATP[16] and secrete pro-inflammatory cytokines, which leads to the recruitment of macrophages

and neutrophils and efferocytosis of dying cardiomyocytes[7,17]. Initially, recruited leukocytes

promote the repair of damaged myocardium by stimulating resident fibroblasts to secrete

extracellular matrix, and by inducing angiogenesis to provide needed oxygenation and nutrients

to the healing cardiac muscle[18,19]. While this acute inflammatory response is necessary to activate

fibroblasts and prevent excessive cardiomyocyte death, chronic activation of both the innate and

adaptive immune responses has been shown to exacerbate heart failure progression[7,20–22]. The

immune cell cascade and timeline for resolution of inflammation after ischemic myocardial

infarction has been well characterized, however, it is less well understood how immune cells are

recruited to the injured myocardium in chronically developing non-ischemic heart failure[23].

Pannexin 1 (PANX1) is a ubiquitously expressed transmembrane channel, through which

stressed or damaged cells release ATP and other small metabolites that are smaller than 1 kDa.

PANX1-dependent ATP release from apoptotic cells serves as a "find me" signal for efferocytosis

by immune cells[24–26]. PANX1 channel opening can be activated via multiple mechanisms including

α_{1D}[27]- or β-adrenergic receptor stimulation[28,29], and caspase-3-mediated cleavage[30]. Several

pharmacologic compounds including probenecid, carbenoxolone, and the FDA-approved heart

failure therapeutic spironolactone have been identified as PANX1 channel blockers[31–33]. Release

of ATP from cardiomyocytes through PANX1, which caused fibroblast activation *in vitro,* has

previously been described[16]. Moreover, global PANX1 deletion in mice decreased ischemic area

in a model of cardiac ischemia reperfusion injury[34], while endothelial cell-specific deletion of

PANX1 in mice decreased immune cell infiltration in a model of myocardial infarction[35]. While

these prior studies have strongly implied PANX1 as a therapeutic target for cardiovascular

diseases in general[36], the role for PANX1 in cardiomyocytes specifically in the regulation of

chronic cardiac pathologies, including heart failure, remains to be elucidated.

Here we report that PANX1 channels can be activated in cardiomyocytes through beta-

adrenergic receptor stimulation. Using mice with a cardiomyocyte-specific PANX1 deletion

(Panx1[MyHC6]), we demonstrate a crucial role for PANX1 channels in the development of non-

ischemic heart failure. Panx1[MyHC6] mice were protected against isoproterenol-induced [37,38] cardiac

hypertrophy, systolic dysfunction, and neutrophil infiltration into the myocardium. Furthermore,

we find that PANX1 deletion facilitates glucose utilization and increases glycolytic and decreases

inflammatory gene expression. Taken together our findings implicate PANX1 channels as

promising therapeutic targets in non-ischemic heart failure and cardiac hypertrophy.

Results:

Isoproterenol Activates Pannexin 1 channels in Cardiomyocytes:

Previous studies have demonstrated that Pannexin 1 (PANX1) channels are activated in response to β3 and α_{1D}-adrenergic receptor stimuli[27,28]. A recent study demonstrates beta-adrenergic receptor 1-dependent activation of Panx1 channels in cardiomyocytes via cAMP/PKA-mediated phosphorylation on Serine 206[29]. To activate Panx1 channels we used isoproterenol, a pan β-adrenergic receptor agonist, in H9c2 rat myoblasts which endogenously express all β-adrenergic receptor family members and Panx1[39–41]. We tested Panx1 channel opening by measuring both ATP release into the culture media and dye (Yo-Pro-1) uptake into cells. After stimulation with 2 or 20μM isoproterenol, H9c2 cells released significantly more ATP into the extracellular media compared to vehicle stimulation (**Figure 1A**). Isoproterenol-induced ATP release was abolished by either siRNA mediated knock-down of Panx1 (successful knock-down is demonstrated by significantly decreased mRNA and protein levels (S1A,B)), or pharmacological channel inhibition with spironolactone[33] (**Figure 1A,C**). Cell viability was not affected at either dose of isoproterenol (S1C).

Additionally, using a flow-cytometry based approach, we found that stimulation with 20μM isoproterenol caused a significant increase in Yo-Pro-1 positive, Annexin V negative H9c2 cells compared with vehicle stimulation (**Figure 1B**, representative gating S1D). Isoproterenol-induced Yo-Pro-1 uptake was significantly decreased after siRNA-mediated Panx1 knock-down (**Figure 1B,** S1D), and blunted with PANX1 channel blockade by spironolactone, carbenoxolone, or probenecid (**Figure 1D**). Taken together, these data demonstrate that PANX1 channel opening in cardiac myoblasts as a result of β-adrenergic receptor stimulation leads to the release of ATP.

Generation and characterization of a mouse with Pannexin 1 deletion specifically in cardiomyocytes

To investigate the role of PANX1 channels in cardiomyocytes *in vivo*, we generated a cardiomyocyte-specific PANX1 null mouse by crossing Panx1 floxed mice[42] with mice expressing Cre recombinase under the control of the alpha myosin heavy chain promoter (MyHC6). Confirmation of successful recombination was demonstrated by the presence of a null band on PCR analysis of genomic DNA isolated from the hearts of Panx1[MyHC6] mice, which was not present in the hearts of Panx1[fl/fl] mice or other tested organs (liver, kidney) from either mouse genotype (S2A). Levels of *Panx1* mRNA (by qPCR) and PANX1 protein (by immunofluorescence) were significantly decreased in isolated cardiomyocytes and cardiac troponin t (cTnnt) positive cells, respectively, from Panx1[MyHC6] mice (**Figure 2A**, S2B). Panx1[MyHC6] mice had no obvious gross developmental defects when compared to their Panx1[fl/fl] littermates as examined by body mass, organ mass, or tibial length at 12 weeks of age (S2C-F). Furthermore, assessment of cardiac function via 2D-echocardiography showed no significant differences in end-systolic or end-diastolic volume, wall thickness, or ejection fraction at baseline (**Figure 2B-E,** S2K). Finally, blood pressure measurements by radiotelemetry in 12-week-old male mice demonstrated no differences in mean arterial pressure, systolic or diastolic pressure, or heart rate in Panx1[MyHC6] mice when compared to littermate Panx1[fl/fl] mice (S2G-J).

Pannexin 1 deletion shifts cardiomyocyte metabolism from oxidative phosphorylation to glycolysis.

To explore effects of PANX1 deficiency on the cardiomyocyte transcriptome *in vivo*, we performed bulk RNA-sequencing on cardiomyocytes isolated from hearts of Panx1[MyHC6] and

Panx1[fl/fl] mice, which identified 1281 and 727 uniquely expressed genes, respectively (**Figure 3A**).

There were 1069 significantly differentially expressed genes between the genotypes, with 793

down-regulated and 276 up-regulated (**Figure 3B**). Gene ontology (GO) Term and KEGG pathway

analysis revealed that Panx1 deletion mainly affected pathways involved in intercellular

communication, cell adhesion and gap junction function, namely "cell-cell" and "Wnt signaling",

"microtubule-based movement", "ECM-receptor interactions", "leukocyte trans endothelial

migration", and "cell adhesion molecules", among others involved in inflammation and

cytoskeletal organization (S3A, B). KEGG Pathway analysis was corroborated by significantly lower

expression of transcripts of the claudin and cadherin family and their receptors (*Cdh1, Cldn7,*

Cdhr3, Cdhr4), as well as increased expression of Transport and Golgi Organization protein 2

Homolog (*Tango2*) , which has been associated with altered mitochondrial energy metabolism in

humans[43,44] (**Figure 3B**).

Pannexin channels have been demonstrated to control cellular metabolism, specifically

the regulation of glucose homeostasis and glycolytic metabolism[45–48]. A previous study by

Adamson et al. demonstrated that Panx1 channel function is required for insulin-induced glucose

metabolism in white adipocytes[49]. To assess whether loss of PANX1 impacted these pathways in

cardiomyocytes, we analyzed the genes associated with key glycolytic enzymes in PANX1-

deficient cardiomyocytes. Our data demonstrate increased expression of glycolytic pathway

genes in cardiomyocytes isolated from Panx1[MyHC6] mice compared to their floxed controls. Of

note, we found a significantly higher levels of the glucose transporter *Slc2a4* (Glut4), which is the

predominant isoform in cardiomyocytes (**Figure 3C**).

To determine if the observed transcriptional increase in genes involved in glycolysis

impacts cellular metabolism and glucose utilization, we first performed a glycolytic stress test

(GST) using a Seahorse extracellular flux analyzer on H9c2 cells that had been treated with either

Panx1 siRNA or control siRNA. PANX1 knock-down alone resulted in significantly increased basal

and maximal extracellular acidification rate (ECAR). The magnitude of increase was even further

increased by treatment with isoproterenol (**Figures 3D, E,** representative trace S3C).

Next, we performed glucose uptake assays to determine if the increase in glycolytic rate

shown in the glycolytic stress test could be attributed to an increase in glucose present in the

cardiomyocyte. Intracellular 2NBD-glucose levels were significantly increased after stimulation of

H9c2 cells with isoproterenol, which was abrogated after siRNA-mediated knock down of PANX1

(**Figure 3F**). Though cardiomyocyte metabolism was significantly changed by PANX1 deletion *in*

vitro, intracellular ATP content is not significantly different (S3D). Taken together these data

demonstrate a key role for PANX1 channels in regulating glucose uptake and glycolytic rate in

H9c2 cardiomyocytes.

Panx1 deficiency in cardiomyocytes protects against isoproterenol-induced cardiac

hypertrophy and cardiac dysfunction in non-ischemic heart failure

To examine the role of Panx1 channels in regulating cardiomyocyte function of the heart

under stress, we induced cardiac hypertrophy preceding non-ischemic heart failure using an

established protocol where we repeatedly administered isoproterenol via intraperitoneal

injection for 14 days at a dose of 15mg/kg/day[37,50]. We established this timepoint as an early

measure of disease progression, where cardiac hypertrophy is the predominate phenotype. We

performed 2D-echocardiography prior to the beginning of the study to collect baseline

measurements and then again after 14 days at the model endpoint (**Figure 4A**). As shown

previously, before the start of the treatment Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice displayed no overt

differences in cardiac function (Figure 2B-E) or weight (S4A). Repeated isoproterenol

administration resulted in significantly increased heart weight in Panx1$^{fl/fl}$ mice, but not in

Panx1^{MyHC6} mice (**Figure 4B,** S4B). Cardiomyocyte hypertrophy (as measured by cardiomyocyte

cross sectional area by wheat germ agglutinin (WGA) staining) was significantly increased in

hearts of in Panx1$^{fl/fl}$ mice compared with Panx1^{MyHC6} mice after isoproterenol treatment (**Figure

4C**).

Furthermore, treatment with isoproterenol caused significant increases in both end-

systolic and end-diastolic volume of the left ventricle in Panx1$^{fl/fl}$ mice as measured by

echocardiography (**Figure 4D-F,** S4G). Panx1^{MyHC6} mice did not have significant increases in either

metric compared to their baseline and had significant less change compared to Panx1$^{fl/fl}$ age and

sex-matched controls (**Figure 4D-F,** S4G). We determined that this disparity was likely caused by

heart-intrinsic functional defects and not systemic vascular or renal damage, as we observed no

differences in blood pressure or serum creatinine between Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice after

treatment with isoproterenol (S4C-F).

Additionally, to assess cardiac fibrosis found at later stages of non-ischemic heart failure

and to examine if Panx1 deficiency would also protect against fibrotic changes, we induced non-

ischemic heart failure by administration of isoproterenol for 28 days (15/mg/kg/day) by

subcutaneous implantation of osmotic pump (S5A). Prolonged isoproterenol treatment caused a

significantly greater increase in Collagen 1 (COL1A1) content in the myocardium of Panx1$^{fl/fl}$ mice

compared to Panx1^{MyHC6} mice, as determined by immunofluorescence staining (S5B).

These data demonstrate that cardiomyocyte-specific deletion of PANX1 is

cardioprotective against cardiac hypertrophy and interstitial fibrosis during the progression of

non-ischemic heart failure.

Pathways involved in the immune response and fatty-acid metabolism are down regulated in

the hearts of Panx1^{MyHC6} mice during non-ischemic heart failure

To elucidate potential mechanisms that are involved in the protection from cardiac

hypertrophy and dysfunction observed in Panx1^{MyHC6} mice, we performed bulk RNA-sequencing

on whole hearts of Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice after 14-day administration of isoproterenol.

We identified 1592 genes which were significantly differentially expressed between the two

genotypes, with 594 genes downregulated and 998 upregulated in Panx1^{MyHC6} mice compared to

Panx1$^{fl/fl}$ controls (**Figure 5A**).

KEGG pathway and GO Term analysis on the differentially expressed genes identified

regulatory pathway families including "hematopoietic cell lineage", "cell adhesion molecules", "B-

cell receptor signaling pathway", and "Rheumatoid arthritis", all of which suggested significant

changes in molecules associated with immune cell recruitment and infiltration. Moreover, we

identified "fatty acid degradation", "fatty acid metabolism", and "cardiac muscle contraction",

implicating a metabolic shift in the absence of PANX1 channels. Furthermore, identified GO Terms

indicated changes in plasma membrane proteins, specifically transporters and ion channels, as

well as "oxidoreductase activity", "scavenger receptor activity", and "hydrolase activity",

implicating the possible shift in the oxidative stress state.

Further analysis of genes associated with fatty acid beta-oxidation (as identified through

WikiPathways) demonstrated overall decreased expression of this group of genes in hearts from

Panx1^MyHC6 mice with non-ischemic heart failure, while some genes such as *Lipc* (encoding the

lipase C hepatic type), were expressed at higher levels (**Figure 5B**). Interestingly, a lowered

reliance on fatty acid beta-oxidation during ischemic stress has been previously associated with

increased cardiomyocyte viability[15].

Examination of genes involved in the inflammatory response identified lower expression

of *Cd40*, *Il2ra*, and *Il2* in hearts of Panx1^MyHC6 mice compared to their Panx1^fl/fl counterparts

(**Figure 5C**), suggesting a lower immune cell load in the hearts as CD40 and IL-2 expression are

associated with a wide range of leukocyte populations[51]. Additionally, we observed lower

expression of *Lamb1* (encoding Laminin subunit beta 1) and *Lama5* (encoding laminin subunit

alpha 5) as well as *Vtn* (encoding Vitronectin) suggestive of decreased cell-cell adhesion and

fibrosis (**Figure 5C**).

Since PANX1 channel function has been associated with activation of a variety of

purinergic receptors[29,52–54], we examined the expression of purinergic signaling receptors and

other associated molecular regulators in our RNA-Seq dataset (WikiPathway "Purinergic

signaling"). We found lower expression of *P2yr1*, *P2rx1*, *P2ry14,* and *Adora2a* in hearts of

Panx1^MyHC6 mice (**Figure 5D**) compared to Panx1^fl/fl controls. This finding is particularly interesting

as *P2rx1, P2yr14*, and *Adora2a* have all been previously associated with chemotaxis and activation

of immune cells, particularly neutrophils[55].

Together these data identify that PANX1 deficiency in cardiomyocytes causes significant

transcriptomic changes in genes associated with inflammation, immune cell recruitment,

metabolism and oxidative phosphorylation, all of which may contribute to the protection of

Panx1^{MyHC6} mice from cardiac hypertrophy and progressive heart failure in a model of

isoproterenol induced non-ischemic heart failure.

Neutrophil infiltration is significantly blunted in Panx1^{MyHC6} mice during heart failure induced

by isoproterenol

To determine the contribution of cardiomyocytes to the inflammatory state in

isoproterenol-injured hearts, we performed bulk RNA-Seq on the isolated cardiomyocytes from

Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice. We identified 1439 differentially expressed genes with the majority

being downregulated in Panx1-deficient cardiomyocytes (1037 down, 402 up) (S6A). Further

analysis of genes associated with "inflammation response" (WikiPathways) revealed decreased

expression of many of the genes identified as being regulatory in inflammation including Zap70,

Lack, Fn1, Col1a2, and multiple cytokine receptors (**Figure 6A**). This was corroborated by the

finding of the GO Terms "immune response" and "immune system process", which were

significantly enriched with genes downregulated in the isolated cardiomyocyte fraction of

Panx1^{MyHC6} mice compared to Panx1$^{fl/fl}$ mice (**Figure 6A**).

To determine whether the observed inflammatory gene regulation patterns was reflective

of perturbed leukocyte recruitment to the hearts of Panx1^{MyHC6} mice, we isolated the non-

myocyte fractions from the hearts of Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice after 14-day treatment with

isoproterenol and examined leukocyte populations by flow cytometry (representative gating S6B).

The number of total leukocytes (CD45$^+$) as well as neutrophils (Ly6g$^+$, CD11b$^+$) were significantly

lower in the hearts of Panx1^{MyHC6} mice (**Figure 6B, C,** S6B). These findings were confirmed by

immunofluorescence imaging of the left ventricle of hearts from Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice.

In accordance with the flow cytometry data, the number of Ly6g$^+$ puncta per field in the hearts

of Panx1^{MyHC6} mice was significantly lower compared to hearts from Panx1$^{fl/fl}$ mice (**Figure 6D,**

S6C).

Of note, there was no significant difference in the total number of macrophages (F4/80$^+$,

CD11b$^+$) present in Panx1^{MyHC6} mouse hearts during non-ischemic heart failure (**Figure 6E,** S6D).

Further analysis of macrophage subsets showed no differences in macrophages with the co-

stimulatory macrophage activation marker CD86 (F4/80$^+$, CD11b$^+$, CD86$^+$) or macrophages with

the anti-inflammatory marker CD163 (F4/80$^+$, CD11b$^+$, CD163$^+$/CD86$^-$) (**Figure 6E,** S6D). Finally,

we observed no differences in dendritic cells (CD11b$^+$, CD24$^+$) (**Figure 6E,** S6D).

These data demonstrate a critical role for PANX1 expressed in cardiomyocytes in recruiting

immune cells, specifically neutrophils, to the myocardium in response to chronic beta-adrenergic

stimulation. This finding is in line with previous work which demonstrates that PANX1-dependent

ATP release recruits neutrophils to the endothelium during acute inflammatory disease[56].

Discussion:

The role of ATP, adenosine, and other purinergic signaling molecules as both autocrine

and paracrine signals in the myocardium has long been understood and exploited clinically, using

compounds such as clopidogrel and adenosine[57]. However, the underlying mechanisms for

release of purinergic compounds in non-ischemic heart failure still need to be investigated[58,59].

To examine the role of PANX1 in cardiomyocytes, we generated a novel mouse with

cardiomyocyte-specific deletion of PANX1 (Panx1^{MyHC6}). We found that Panx1^{MyHC6} mice shift the

baseline metabolic state of their cardiomyocytes toward a more glycolytic phenotype, concurrent

with increased glucose uptake and GLUT4 transporter expression. After isoproterenol-induced

non-ischemic heart failure, PANX1[MyHC6] mice are protected from pathological cardiac hypertrophy

and dysfunction. We also find that in Panx1[MyHC6] mouse hearts during isoproterenol-induced

cardiac hypertrophy there is a decrease in gene expression of fatty-acid beta oxidation- associated

genes. Finally, PANX1 deficiency in cardiomyocytes blunts neutrophil recruitment to the

myocardium during isoproterenol-induced non-ischemic heart failure.

Two previous studies by Adamson *et al.* and Senthivinayagam *et al.* found that deletion of

PANX1 in either white or brown adipose tissue exacerbated diet-induced obesity, specifically by

increasing susceptibility to insulin resistance, establishing that PANX1 plays a critical role in

glucose homeostasis in the adipose tissue[28,49]. Since cardiomyocytes are among the most energy-

demanding cell types requiring high levels of ATP needed for contraction, we examined the role

for PANX1 in control of cardiomyocyte metabolism. We demonstrate that PANX1 deletion induces

a metabolic shift in the cardiomyocyte by priming the cardiomyocyte toward a glycolytic state at

baseline. Consequently, PANX1-deficient cardiomyocytes are less responsive to stress induced by

beta-adrenergic receptor activation. We found a significant increase in *Slc2a4* expression in

isolated primary murine cardiomyocytes of Panx1[MyHC6] mice, which is matched by increased

glucose uptake and glycolytic rate in PANX1 knock-down H9c2 myoblasts *in vitro*[14]. This glycolytic

phenotype has previously been observed in fetal cardiomyocytes and is associated with increased

proliferative capacity[11,14]. Furthermore, increased glucose availability and glycolytic rate in

unstressed situations leads to better outcomes and is associated with increased regenerative

capacity after ischemic/ reperfusion injury[15,60].

RNA-Sequencing of isolated cardiomyocytes from Panx1[MyHC6] and Panx1[flfl] mice with non-

ischemic heart failure revealed significantly decreased expression of genes associated with

immune system function. Previous work established that PANX1 deletion in endothelial cells

decreases the immune cell recruitment to the myocardium after a myocardial infarction, without

further specifying a particular immune cell population[35]. We find that deletion of PANX1 in

cardiomyocytes decreases overall immune cell recruitment, which was driven specifically by a

significant decrease of neutrophils. Immune cell recruitment to the myocardium has been shown

to exacerbate cardiomyocyte damage and impairment of myocardial function[7,61]. Therefore, the

decrease in immune cells in the myocardium of mice with PANX1-deficient cardiomyocytes

suggests that PANX1 may serve to propagate or initiate the sterile immune response in the

myocardium during times of chronic cardiac stress[22]. The changes in gene expression reveal an

overall decreased chronic inflammatory state of the myocardium in Panx1[MyHC6] mice after

isoproterenol induced heart failure.

While our observations of neutrophil recruitment by PANX1 are in the context of

isoproterenol-induced non-ischemic heart failure, PANX1 activation in cardiomyocytes may also

be important in ischemic heart failure or after ischemia-reperfusion injury. Future investigations

will elucidate if the effects on neutrophil infiltration attributed to cardiomyocyte-specific deletion

of PANX1 vary by injury type, which will provide opportunities to develop therapies that are

specific to the type of cardiac injury.

ATP is known to serve as recruitment or "find me" signal for immune cells during

inflammation[62,63] and apoptosis[24,25]. While we do not identify the responsible receptors on

neutrophils, our RNA sequencing data demonstrate that expression of key purinergic receptors,

namely P2X1, P2Y1, P2Y14, and ADORA2A, is decreased in the whole heart tissue of Panx1[MyHC6]

mice during isoproterenol-induced heart failure. Of these P2Y1, P2Y14, and ADORA2A can be

associated with decreased neutrophil recruitment[64]. Moreover, we cannot rule out other

purinergic molecules that could be reseased through PANX1 potentially driving the neutrophil

recruitment. P2Y1 and ADORA2A bind ADP and adenosine, respectively, which could be

generated from local exonucleases acting on the ATP released via PANX1 or could be released

from PANX1 itself. Additionally, P2Y14 on neutrophils binds UDP-glucose[65], which can be released

through PANX1[25,64]. Therefore, a combination of signaling molecules released by PANX1, including

ATP and UDP-glucose, may be necessary for the infiltration of neutrophils into the myocardium

during non-ischemic heart failure.

Historically, treatment of heart failure in patients has focused on therapies which treat the

symptoms of the disease, such as edema and dyspnea, rather than directly targeting cardiac

function or preventing functional decline of the cardiac muscle. Previously, clinical trials have

identified that spironolactone, a mineralocorticoid receptor antagonist (MRA) and PANX1 blocker,

improves patient outcomes in both all-cause mortality and sudden cardiac death in patients with

established heart failure over a placebo[33,66,67]. However, it is is not completely clear why this

therapy is more beneficial than others which also lower blood pressure. Based on our findings we

speculate that this can be explained by the additional action of spironolactone as a PANX1

channel blocker. Systemic PANX1 channel blockade may improve patient outcomes by preventing

immune cell recruitment, as has been shown in multiple mouse models[24,35,53]. Furthermore,

modulating cardiomyocyte metabolism to increase glucose uptake may prevent an energy

imbalance which is thought to be a driving factor in the functional decline of cardiac muscle.

Taken together our study illustrates how PANX1 deletion in cardiomyocytes protects

against the progression of non-ischemic heart failure. We demonstrate that PANX1 channel

opening can be induced by beta-adrenergic receptor stimulation in cardiomyocytes.

Cardiomyocyte-specific deletion in mice (Panx1^{MyHC6}) impacts their cardiac glycolytic metabolism,

which we attribute to a higher expression of *Slc2a4* and increased glucose uptake. Panx1^{MyHC6}

mice are protected from isoproterenol-induced non-ischemic heart failure and cardiac

hypertrophy, which we demonstrate to be at least in part due to a decrease in neutrophil

recruitment. Our findings identify PANX1 in cardiomyocytes as a therapeutic target to prevent

severe cardiac tissue damage in non-ischemic heart failure, which may also be relevant for

developing therapies targeted at PANX1 for other forms of heart failure such as HFpEF.

Methods

Animals: Cardiomyocyte-specific Panx1-deficient mice were generated by crossing *Panx1$^{loxP/loxP}$*

(Panx1$^{fl/fl}$) mice[42] with B6.FVB-Tg(Myh6-cre)2182Mds/J (MyHC6-Cre)[68] (strain #011038) obtained

from the Jackson Laboratory. For breeding purposes, mice hemizygous for the MyHC6 Cre were

cross with Panx1$^{fl/fl}$ mice to give littermate mice of Panx1$^{loxP/loxP}$ MyHC6$^{wt/wt}$ (Panx1$^{fl/fl}$) and

Panx1$^{loxP/loxP}$ MyHC6 $^{Cre/wt}$ (Panx1^{MyHC6}). All animals were maintained at the University of Virginia

Center for Comparative Medicine in a pathogen-free vivarium facility. Mice were maintained on

a 12-hour light/dark cycle and provided ad libitum access to standard rodent chow (Teklad 7912)

and water. All experiments performed were approved by the University of Virginia Animal Care

and Use Committee (protocol #3444). All experiments were performed using age-matched

littermate controls. Genotyping was performed via PCR from tail clip biopsies after proteinase K

digestion in DirectPCR tail lysis buffer (Viagen Biotech 102-T). PCR of genomic DNA was

performed using Apex Taq Master Mix and allele-specific primers as previously described[69].

Isoproterenol induction of heart failure: 8–10-week-old male mice were treated daily with

isoproterenol or saline vehicle for 14 days to induce heart failure as previously described.

Isoproterenol (medchem: HYB0468) was freshly prepared daily in sterile saline prior to injection

(15 mg/kg/day) and administered by intraperitoneal injection. Mice were weighed daily to

monitor for excessive weight loss and appropriate dose administration.

Isoproterenol progression of heart failure: For longer-term administration of isoproterenol,

osmotic pumps (Alzet 2004) were filled with isoproterenol in sterile saline plus 0.02% acetic acid

and mice were weighed prior to implantation to administer 15mg/kg/day according to pump rate

specifications. Pumps were equilibrated overnight in sterile saline prior to implantation. 8–10-

week-old male mice were anesthetized using isoflurane and implantation sites were exposed after

hair removal with Nair hair removal lotion with aloe and lanolin. Mice were administered

bupivicane analgesic and sites were sterilized with three washes of betadine and ethanol wipes.

Pumps were implanted in a subcutaneous pocket on the back of the mouse and surgical clips

were placed for wound closure. Mice were placed on a heating-pad and isoflurane anesthesia

was maintained for the length of the procedure. Surgical clips were removed 10-14 days after

placement.

Immunofluorescence staining: Hearts were harvested and fixed for 4 hours in 10% zinc-buffered

formalin prior to fixation in 50% ethanol. Tissues were paraffin embedded and cryosectioned to

7μm thickness and mounted. Slides were deparaffinized. Briefly, sections were submerged in

Histo-Clear (Electron Microscopy Sciences 64110-01) (3x10 minutes), 100% ethanol (2x3

minutes), 95% ethanol (2x3 minutes), 70% ethanol (1x3 minutes), and PBS (1x5 minutes). Antigen

retrieval was performed using citrate-based solution (Vector Laboratories H-330) where slides

were submerged in antigen retrieval solution and heated to boiling for 20 minutes, slides were

then cooled for 10 minutes at room temperature and 10 minutes at 4°C. Tissue sections were then

blocked for 1 hour in antibody blocking buffer (0.2% TritonX, 1mL donkey serum) at room

temperature. Antibody blocking buffer was removed and replaced with antibody blocking buffer

containing primary antibody (1:100) overnight at 4°C. Sections were then washed (3xPBS for 5

minutes) and incubated in antibody blocking buffer containing secondary antibody (1:500) for 1

hour at room temperature protected from light. Sections were washed (3xPBS) and

counterstained with DAPI (Thermo Fisher Scientific D3571) before mounting. Sections were

imaged on an Olympus Fluoview 1000 and are representative images of composite z-stacks.

Analysis (thresholding, manual counting, and fluorescence intensity quantification) were

performed in ImageJ.

Quantification of Cardiomyocyte cross-sectional area: Hearts were prepared as indicated in

immunofluorescence staining. Sections were stained with cTnnt (1:100 - Abcam ab8295) during

primary incubation and DAPI (1:500 - Thermo Fisher D3571) and Wheat-Germ-Agglutinin-Alexa

Fluor 647 (5 µg/m – Thermo Fisher W32466) during secondary incubation. Sections were imaged

on an Olympus Fluoview 1000 confocal microscope with five images per mouse taken in the left

ventricle across two sections. 50 round nucleated cardiomyocytes per mouse were outlined

across the images and were analyzed in ImageJ.

Flow cytometry of infiltrating cardiac leukocytes: Primary infiltrating leukocytes were isolated

from cardiac tissue by non-Langendorff perfusion previously described[70]. Briefly, hearts were

sequentially perfused with EDTA buffer, perfusion buffer, and digestion buffer containing

collagenase until hearts appeared soft. Hearts were then mechanically dissociated with scissors

and pipetted up and down. Cells were washed through a 100µm cell strainer to create a single

cell suspension. Myocytes were isolated by centrifugation at 120 x g for 5 minutes. Supernatants

(containing non-myocyte cells) were transferred to clean tubes and pelleted by centrifugation at

300 x g for 5 minutes. Pellets were treated with ACK lysis buffer for 1 minutes and then

resuspended in 1mL of FACS buffer (PBS, 2mM EDTA, and 1%BSA). Splenocytes were isolated by

grinding through a 100µm cell strainer and washing with PBS containing 2mM EDTA. Samples

were then incubated with ACK lysis buffer for 2 minutes and spun at 400 x g for 5 minutes. All

samples were then counted and resuspended at 100µL per 100,000 cells in PBS. Cells were

stained with LIVE/DEAD Yellow (1:1000) (Thermo Scientific L34967) for 30 minutes on ice. Fc-

receptors were then blocked for 15 minutes using FcBlock (BioRad – BUF041A). Cells were

pelleted by centrifugation (300 x g,5 minutes) and resuspended in FACS buffer (100μL per 100,000

cells) and stained with fluorophore-conjugated antibodies (1:100) for 1 hour on ice. See tables for

antibodies included below. Cells were washed in PBS (x3) after staining. Flow cytometry collection

and deconvolution was performed on an Cytek Northern Lights 3 laser Spectral Flow Cytometer.

Automatic deconvolution was performed using single stains generated from splenocytes. Gating

and post-hoc analysis was performed with FCS-Express 7.18. Representative gating is presented

for all flow cytometry experiments in supplemental figures.

RNA-Sequencing: Hearts of mice were removed after whole body perfusion with 10mL of PBS,

weighed, and snap frozen at -80^C prior to analysis. For some experiments, isolated

cardiomyocytes were suspended in 1mL of TRIzol^{TM} (Invitrogen) and stored at -80^{oC} for further

analysis[70]. After chloroform incubation and centrifugation per manufacture instructions RNA-

containing phase was diluted with 70% ethanol 1:1 and RNA isolation was performed using

RNeasy Mini Kit (Qiagen). For tissues, Total RNA was isolated using RNeasy Mini Kit (Qiagen) after

lysis in RLT buffer with bead homogenization. Total RNA quantity and quality were then assessed

on a NanoDrop 2000 spectrophotometer (Thermo Fisher Scientific). Library preparation including

quality metrics, sequencing, and clustering were performed by Novogene (Sacramento, CA).

Sample integrity was assessed prior to library preparation using the Bioanalyzer 2110 system with

an RNA Nano 6000 Assay Kit (Agilent Technologies). One hundred fifty-base paired-end

sequencing was performed using an Illumina NovaSeq platform according to manufacture

instructions. Fastq files were processed using in-house perl scripts from Novogene to obtain clean

reads. Clean reads were aligned to the mouse reference genome using Hisat2 v2.0.5.

FeatureCounts was used to calculate read numbers for each gene and fragments of per kilobase

of transcript sequence per millions base pairs sequenced (FPKM) were calculated. Differential

expression analysis was performed using DESeq2. Differentially expressed genes (DEG) were

assigned by criteria of P-value adjusted < 0.05 (-Log(p-value) > 1.3). KEGG pathway analysis and

Gene Ontology pathway analysis was performed on DEG using clusterProfiler.

Echocardiography: 8–16-week-old male mice were anesthetized using isoflurane. The chest was

exposed after hair removal and a four-lead electrocardiogram (ECG) was collected during the

procedure. Images were collected using a VevoView 1100 system. Images were collected in the

long-axis (B-mode) and short axis (M-mode) on mice serially throughout the experimental period.

Serial collection began prior to disease induction and occurred every two-weeks during disease

progression. Images were analyzed using Vevo Lab. Left-ventricle (LV) end-diastolic, end-systolic

volume, and wall thickness were assessed, and ejection fraction calculated.

Blood Pressure Assessments using Radiotelemetry: Blood pressure was measured using

telemetry equipment (Data Sciences International, DSI) as previously described[27]. Mice were

surgically implanted with radiotelemetry units (PA-C10 or HD-X10). Briefly, while under

isoflurane anesthesia, the catheter of a radiotelemetry unit was placed in the left carotid artery

and positioned such that the probe reached the aortic arch. The radio-transmitter was placed in

a subcutaneous pouch at the right flank. Buprenorphine was used as an analgesic. Mice were

allowed to recover for seven days prior to the initiation of recordings. Baseline blood pressure

measurements, including systolic pressure, diastolic pressure, mean arterial pressure (MAP) and

heart rate, were recorded every minute for a continuous period of 5 days using Dataquest A.R.T.

20 software (DSI). Diurnal (inactive period) MAP was measured during animal's light cycle (6:00

a.m. to 5:59 p.m.) and nocturnal (active period) MAP was measured during the animal's dark

cycle (6:00 p.m. to 5:59 a.m.).

Antibodies: Antibodies used in these studies were rabbit monoclonal anti-Panx1 (Cell Signaling -

#91137), rabbit monoclonal anti-CCR2 (Abcam – ab273050), rat monoclonal anti-Ly6g (R&D

Systems – MAB1037), mouse monoclonal anti-cardiac troponin T (Abcam – ab8295), and wheat

germ agglutinin (Invitrogen – W32466). Flow cytometry antibody used include CD45 rat anti-

mouse APC eFluor 780 (Invitrogen - 47-0451-80), CD11b rat anti-mouse FITC (Invitrogen - 11-

0112-82), CD163 rat anti-mouse PE (Invitrogen - 12-1631-80), CD86 rat anti-mouse SuperBright

436 (Invitrogen - 62-0869-41), LIVE/DEAD Yellow Cell Stain for 405 (L34967), Ly6g rat anti-mouse

PE-Cyanine5 (Invitrogen - 15-9668-80), F4/80 anti-mouse APC (eBioscience 17-4801-82), CD64

anti mouse PerCP-710 (Invitrogen – 46-0641-80), CD24 anti mouse PE-Cyanine7 (Invitrogen 25-

0242-82).

Cell culture: H9c2 rat myoblasts were obtained from ATTC and maintained in high-glucose DMEM

supplemented with 10% heat-inactivated FBS, 15 mM sodium bicarbonate (Glibco), 1mM sodium

pyruvate (Glibco), and 1% Pen-Strep. Cells were passaged when they reached 70% confluency on

tissue culture (TC) treated dishes using 0.25% trypsin (Glibco).

siRNA-Mediated Knock-Down of Panx1: H9c2 cells in 12 or 24 well TC treated plates or 10cm TC

treated dishes were washed in sterile PBS and then incubated with OptiMEM (Glibco). Cells were

treated with RNAiMax Lipofectamine and siRNA against Panx1 or negative control silencer #1

(Ambion - 197003) per the manufacturer's instructions. Experiments were performed 72 hours

after transfection with the siRNA.

RNA isolation and quantitative reverse transcription PCR: RNA was isolated from approximately

75,000 H9c2 rat myoblasts or isolated cardiomyocytes from one mouse heart and lysed in RLT

lysis buffer. RNA was isolated using the RNeasy Mini Kit (QIAGEN) according to the manufacturer's

instructions. RNA was quantified using a NanoDrop spectrophotometer (Thermo Fisher

Scientific), and complementary DNA (cDNA) libraries were generated using the iScript cDNA

Synthesis Kit (Bio-Rad). Quantitative reverse transcription PCR was performed using TaqManFast

Advanced Master Mix (Thermo Fisher Scientific) and TaqMan primers (mouse *Panx1* mm0045091

– Invitrogen) (Rat *Panx1* Rn01447976_m1 – Thermo Fisher Scientific) for genes of interest on a

CFX Connect real-time PCR instrument (Bio-Rad). Relative gene expression was determined using

18S (4333760 – Thermo Fisher Scientific) or *Hprt* (mm01545399 – Invitrogen) as a housekeeping

gene.

ATP release measurements: H9c2 cells were plated at 80,000 cells per well in a 48 well dish and

put in serum free media overnight prior to experimental treatment. Cells were incubated with

Panx1 inhibitors (spironolactone 50 µM in serum free media) or vehicle control for 1 hour prior

to stimulation. ARL (Tocris 1283) (300 µM) was added for 30 minutes prior to stimulation. Cells

were then stimulated with vehicle (PBS), isoproterenol (2 or 20 µM in sterile PBS), or

phenylephrine positive control (10 µM) for 15 minutes. One-third of the media in the well was

carefully collected with care taken to not bump cells in the bottom of the well and placed on ice.

ATP concentrations were assessed by CellTiterGlo (Promega G7571) and compared to standard

curve prepared by combining serum free H9c2 media with ATP disodium salt (Tocris 3245).

Dye Uptake Assay: H9c2 cells were plated at 1000,000 cells per well in a 12 well dish and

incubated with Panx1 inhibitors (spironolactone 50 µM, carbenoxolone 50 µM, or probenecid

1mM) or vehicle control for 1 hour prior to stimulation. Cells were stimulated with vehicle (PBS),

isoproterenol (2 or 20 μM in sterile PBS), or phenylephrine positive control (10 μM) for 1 hour at

the same time as Yo-Pro-1 dye (Thermo Fisher Scientific Y3603) (1μM) was added to the media.

Media was removed and cells were washed prior to detachment with trypsin (0.25%). Cells were

spun at 400 x g for 5 minutes and resuspended in PBS with LIVE/DEAD Yellow (1:1000) for 30

minutes. Cells were pelleted and washed in Annexin V binding buffer three times and then

resuspended in Annexin V AlexaFluor 568 (Thermo Fisher Scientific A13202) (1:500) (AV) for 10

minutes. Cells were washed and resuspended in FACS buffer as described above. Cells were

analyzed by flow cytometry on an Attune NxT. Single stain controls were performed using H9c2

cells for initial intensity. FMO were collected and used for deconvolution prior to gating. Cells

were gated for LIVE/DEAD Yellow and AV negative and the percentage of cells positive for Yo-Pro-

1 was assessed as compared to vehicle stimulation.

Glycolytic Stress Test: After treatment with control or Panx1-targeting siRNA as described above,

H9c2 rat myoblasts were plated (40,000 cells per well) in complete medium in XFe96 cell culture

microplate and allowed to settle for one hour at room temperature before overnight culture. The

following day cells were treated with ISO (20μM) for 1 hour in complete medium. Immediately

prior to the assay the media was changed to glycolytic stress test medium (Sigma Aldrich D5030)

supplemented with 143 mM NaCl and 2mM L-glutamine (Gibco 25030 – 081). Glycolytic activity

was assessed by measurement of extracellular acidification rate on a Seahorse XFe96 instrument

(Agilent Technologies). The rate of pH change was measured every 13 minutes for a 3-minute

interval before sequential challenge with 1) 20mM glucose (Sigma Aldrich D9434), 2) 1 μM

oligomycin (Sigma Aldrich 75351), and 3) 80 mM 2-deoxyglucose (Sigma Aldrich D8375). Basal

ECAR was measured as post-glucose ECAR minus post 2-DG ECAR, and maximal ECAR was

measured as post-oligomycin ECAR minus post-2DG ECAR.

2-NDB Glucose Uptake: H9c2 rat myoblasts were plated in a black, clear bottom 96 well plate and

allowed to adhere overnight in complete media. Cells had been cultured with either control or

Panx1 siRNA for 48 hours in OptiMem media prior to plating. The following day cells were treated

with ISO (20µM) for 1 hour or vehicle control. After stimulation media was removed and replaced

with glucose free DMEM as described above in the presence of 2-NDB glucose for 4 hours. The

uptake of fluorescent 2-NBDG was assessed according to manufacturer's instructions (Cayman

Chemicals - Glucose Uptake Cell-Based Assay Kit) on a Synergy HTX plate reader (BioTek).

CytoTox Cell Death Assay: H9c2 rat myoblasts were plated in a clear bottom 96 well plate and

allowed to adhere overnight in complete phenol red free media. Cells had been cultured with

either control or Panx1 siRNA for 48 hours in OptiMem media prior to plating. The following day

cells were treated with ISO (2,20µM) or phenylephrine (10 µM) for 15 minutes or vehicle control.

After stimulation media was removed and LDH presence in the media was assessed according to

manufacturer's instructions (Promega- CytoTox Non-radioactive cytotoxicity assay) on a Synergy

HTX plate reader (BioTek).

Serum Creatinine: Whole blood was collected from mice via cardiac puncture and placed into

serum separating tubes (BD Microtainer – 365967) for 30 minutes at room temperature. Serum

was separated by centrifugation (1500 x g, 5 minutes) and snap frozen on dry ice. Serum

creatinine was measured at 520 nm after addition of sodium picrate according to manufacturer's

instructions (Pointe Scientific - Creatinine Reagent Set C75391250) using a known standard

(Pointe Scientific C75391250).

Statistics: Statistical analyses were performed in Prism 9 (GraphPad). The presence of statistical

outliers was assessed by a ROUT test (2%), and identified outliers were excluded from analysis.

Statistical outliers were identified for data show in Figure 4E, F, which were excluded from

analysis. Statistical tests were performed with two-tailed analysis. Comparisons between two

groups was conducted by Welch's or Student's t-test, while comparisons between more than two

groups were made by one-way analysis of variance or mixed-effects analysis. Sidak's multiple

comparisons test was performed as post-hoc analysis where appropriate. Mixed effects analysis

was used for paired data with more than two groups. A p-value of less than 0.05 was considered

statistically significant. Data are represented as mean ± the standard error of the mean. Image

analysis was performed on 3-5 averaged images per mouse across a minimum of two cardiac

sections. Data points shown for imaging are representative of the average across a mouse, with

each data point representative of individual mice.

Data Availability: Sequencing data reported in this manuscript are deposited in the NCBI Gene

expression Omnibus. All data needed to evaluate the conclusions of this manuscript are present

in the manuscript, present in the supplementary figures, or available upon request to the

corresponding author. Graphics were prepared using BioRender. Panx1^{MyHC6} mice can be provided

by N.L. upon completion of a materials transfer agreement with the University of Virginia School

of Medicine. Requests may be submitted to N.L. (nl2q@virginia.edu).

References:

1. Tsao, C. W. *et al.* Heart Disease and Stroke Statistics—2023 Update: A Report From the American Heart Association. *Circulation* CIR.0000000000001123 (2023) doi:10.1161/CIR.0000000000001123.

2. Schiattarella, G. G. & Hill, J. A. Cardiometabolic HFpEF: Mechanisms and Therapies. *CardioMetabolic Syndr. J.* **1**, 117 (2021).

3. Simmonds, S. J., Cuijpers, I., Heymans, S. & Jones, E. A. V. Cellular and Molecular Differences between HFpEF and HFrEF: A Step Ahead in an Improved Pathological Understanding. *Cells* **9**, (2020).

4. Berthiaume, J. M. *et al.* Chapter 8 - Pathophysiology of Heart Failure and an Overview of Therapies. in *Cardiovascular Pathology (Fourth Edition)* (eds. Buja, L. M. & Butany, J.) 271–339 (Academic Press, 2016). doi:10.1016/B978-0-12-420219-1.00008-2.

5. Chavey, W. E., Hogikyan, R. V., Harrison, R. V. & Nicklas, J. M. Heart Failure Due to Reduced Ejection Fraction: Medical Management. *Am. Fam. Physician* **95**, 13–20 (2017).

6. McMurray, J. J. V. & Packer, M. How Should We Sequence the Treatments for Heart Failure and a Reduced Ejection Fraction? A Redefinition of Evidence-Based Medicine. *Circulation* CIRCULATIONAHA.120.052926 (2020) doi:10.1161/CIRCULATIONAHA.120.052926.

7. Zhang, Y., Bauersachs, J. & Langer, H. F. Immune mechanisms in heart failure. *Eur. J. Heart Fail.* **19**, 1379–1389 (2017).

8. Travers, J. G., Tharp, C. A., Rubino, M. & McKinsey, T. A. Therapeutic targets for cardiac fibrosis: from old school to next-gen. *J. Clin. Invest.* **132**, e148554 (2022).

9. Fearnley, C. J., Roderick, H. L. & Bootman, M. D. Calcium Signaling in Cardiac Myocytes. *Cold*

Spring Harb. Perspect. Biol. **3**, (2011).

10. Williams, G. S. B., Boyman, L. & Lederer, W. J. Mitochondrial calcium and the regulation of

metabolism in the heart. *J. Mol. Cell. Cardiol.* **78**, 35–45 (2015).

11. Lopaschuk, G. D. & Jaswal, J. S. Energy Metabolic Phenotype of the Cardiomyocyte During

Development, Differentiation, and Postnatal Maturation. *J. Cardiovasc. Pharmacol.* **56**, 130–

140 (2010).

12. Neubauer, S. The Failing Heart — An Engine Out of Fuel. *N. Engl. J. Med.* **356**, 1140–1151

(2007).

13. Fragasso, G. Deranged Cardiac Metabolism and the Pathogenesis of Heart Failure. *Card. Fail.*

Rev. **2**, 8–13 (2016).

14. Razeghi, P. *et al.* Metabolic Gene Expression in Fetal and Failing Human Heart. *Circulation*

104, 2923–2931 (2001).

15. Li, X. *et al.* Inhibition of fatty acid oxidation enables heart regeneration in adult mice. *Nature*

622, 619–626 (2023).

16. Dolmatova, E. *et al.* Cardiomyocyte ATP release through pannexin 1 aids in early fibroblast

activation. *Am. J. Physiol. - Heart Circ. Physiol.* **303**, H1208–H1218 (2012).

17. Sager, H. B. *et al.* Proliferation and Recruitment Contribute to Myocardial Macrophage

Expansion in Chronic Heart Failure. *Circ. Res.* **119**, 853–864 (2016).

18. Benjamin, I. J. *et al.* Isoproterenol-induced myocardial fibrosis in relation to myocyte

necrosis. *Circ. Res.* **65**, 657–670 (1989).

19. Revelo, X. *et al.* Cardiac Resident Macrophages Prevent Fibrosis and Stimulate Angiogenesis.

Circ. Res. CIRCRESAHA.121.319737 (2021) doi:10.1161/CIRCRESAHA.121.319737.

20. Blanton, R. M., Carrillo-Salinas, F. J. & Alcaide, P. T-cell recruitment to the heart: friendly

guests or unwelcome visitors? *Am. J. Physiol.-Heart Circ. Physiol.* **317**, H124–H140 (2019).

21. Lavine, K. J. *et al.* The Macrophage in Cardiac Homeostasis and Disease. *J. Am. Coll. Cardiol.*

72, 2213–2230 (2018).

22. Frieler, R. A. & Mortensen, R. M. Immune Cell and Other Non-Cardiomyocyte Regulation of

Cardiac Hypertrophy and Remodeling. *Circulation* **131**, 1019–1030 (2015).

23. Hartman, M. H. T., Groot, H. E., Leach, I. M., Karper, J. C. & van der Harst, P. Translational

overview of cytokine inhibition in acute myocardial infarction and chronic heart failure.

Trends Cardiovasc. Med. **28**, 369–379 (2018).

24. Chekeni, F. B. *et al.* Pannexin 1 channels mediate 'find-me' signal release and membrane

permeability during apoptosis. *Nature* **467**, 863–867 (2010).

25. Medina, C. B. *et al.* Metabolites released from apoptotic cells act as tissue messengers.

Nature **580**, 130–135 (2020).

26. Narahari, A. K. *et al.* ATP and large signaling metabolites flux through caspase-activated

Pannexin 1 channels. *eLife* **10**, e64787 (2021).

27. Billaud, M. *et al.* A molecular signature in the pannexin1 intracellular loop confers channel

activation by the α1 adrenoreceptor in smooth muscle cells. *Sci. Signal.* **8**, ra17–ra17 (2015).

28. Senthivinayagam, S. *et al.* Adaptive thermogenesis in brown adipose tissue involves

activation of pannexin-1 channels. *Mol. Metab.* **44**, 101130 (2021).

29. Zhang, Y. *et al.* Role of cAMP in Cardiomyocyte Viability: Beneficial or Detrimental? *Circ. Res.*

133, 902–923 (2023).

30. Chiu, Y.-H. *et al.* A quantized mechanism for activation of pannexin channels. *Nat. Commun.*

8, 14324 (2017).

31. Michalski, K. & Kawate, T. Carbenoxolone inhibits Pannexin1 channels through interactions

in the first extracellular loop. *J. Gen. Physiol.* **147**, 165–174 (2016).

32. Silverman, W., Locovei, S. & Dahl, G. Probenecid, a gout remedy, inhibits pannexin 1

channels. *Am. J. Physiol. Cell Physiol.* **295**, C761-767 (2008).

33. Good, M. E. *et al.* Pannexin 1 Channels as an Unexpected New Target of the Anti-

Hypertensive Drug Spironolactone. *Circ. Res.* **122**, 606–615 (2018).

34. Rusiecka, O. M. *et al.* Mitochondrial pannexin1 controls cardiac sensitivity to

ischaemia/reperfusion injury. *Cardiovasc. Res.* **119**, 2342–2354 (2023).

35. Good Miranda E *et al.* Endothelial Pannexin 1 Regulates Cardiac Response to Myocardial

Infarction. *Circ. Res.* **0**,.

36. Li, L., He, L., Wu, D., Chen, L. & Jiang, Z. Pannexin-1 channels and their emerging functions in

cardiovascular diseases. *Acta Biochim. Biophys. Sin.* **47**, 391–396 (2015).

37. Oudit, G. Y. *et al.* Phosphoinositide 3-Kinase γ–Deficient Mice Are Protected From

Isoproterenol-Induced Heart Failure. *Circulation* **108**, 2147–2152 (2003).

38. Balakumar, P., Singh, A. P. & Singh, M. Rodent models of heart failure. *J. Pharmacol. Toxicol.*

Methods **56**, 1–10 (2007).

39. Yang, Y. *et al.* Apelin-13/APJ induces cardiomyocyte hypertrophy by activating the Pannexin-

1/P2X7 axis and FAM134B-dependent reticulophagy. *J. Cell. Physiol.* **237**, 2230–2248 (2022).

40. Valentine, C. D. & Haggie, P. M. Confinement of β $_1$ - and β $_2$ -adrenergic receptors in the

plasma membrane of cardiomyocyte-like H9c2 cells is mediated by selective interactions

with PDZ domain and A-kinase anchoring proteins but not caveolae. *Mol. Biol. Cell* **22**,

2970–2982 (2011).

41. Branco, A. F. *et al.* Isoproterenol Cytotoxicity is Dependent on the Differentiation State of

the Cardiomyoblast H9c2 Cell Line. *Cardiovasc. Toxicol.* **11**, 191–203 (2011).

42. Poon, I. K. H. *et al.* Unexpected link between an antibiotic, pannexin channels and

apoptosis. *Nature* **507**, 329–334 (2014).

43. Heiman, P. *et al.* Mitochondrial dysfunction associated with TANGO2 deficiency. *Sci. Rep.* **12**,

3045 (2022).

44. Lujan, A. L. *et al.* Defects in lipid homeostasis reflect the function of TANGO2 in phospholipid

and neutral lipid metabolism. *eLife* **12**, e85345 (2023).

45. Lee, V. R. *et al.* Pannexin 1 regulates adipose stromal cell differentiation and fat

accumulation. *Sci. Rep.* **8**, 16166 (2018).

46. Sayedyahossein, S. *et al.* Pannexin 1 binds β-catenin to modulate melanoma cell growth and

metabolism. *J. Biol. Chem.* **296**, 100478 (2021).

47. Jorquera, G. *et al.* High extracellular ATP levels released through pannexin-1 channels

mediate inflammation and insulin resistance in skeletal muscle fibres of diet-induced obese

mice. *Diabetologia* **64**, 1389–1401 (2021).

48. Banuelos, A. *et al.* Pannexin-1 channels promotes CD8+ T cell effector and memory

responses through distinct metabolic pathways. *J. Immunol.* **210**, 226.15-226.15 (2023).

49. Adamson, S. E. *et al.* Pannexin 1 is required for full activation of insulin-stimulated glucose

uptake in adipocytes. *Mol. Metab.* **4**, 610–618 (2015).

50. Sun, J. *et al.* Human Relaxin-2 Fusion Protein Treatment Prevents and Reverses

Isoproterenol-Induced Hypertrophy and Fibrosis in Mouse Heart. *J. Am. Heart Assoc.* **8**,

e013465 (2019).

51. Von Hundelshausen, P. & Weber, C. Platelets as Immune Cells: Bridging Inflammation and

Cardiovascular Disease. *Circ. Res.* **100**, 27–40 (2007).

52. Filiberto, A. C. *et al.* Endothelial pannexin-1 channels modulate macrophage and smooth

muscle cell activation in abdominal aortic aneurysm formation. *Nat. Commun.* **13**, 1521

(2022).

53. Metz, L. M. *et al.* Platelet pannexin-1 channels modulate neutrophil activation and migration

but not the progression of abdominal aortic aneurysm. *Front. Mol. Biosci.* **10**, 1111108

(2023).

54. Bisht, K. *et al.* Capillary-associated microglia regulate vascular structure and function

through PANX1-P2RY12 coupling in mice. *Nat. Commun.* **12**, 5289 (2021).

55. Burnstock, G. & Boeynaems, J.-M. Purinergic signalling and immune cells. *Purinergic Signal.*

10, 529–564 (2014).

56. Lohman, A. W. *et al.* Pannexin 1 channels regulate leukocyte emigration through the venous

endothelium during acute inflammation. *Nat. Commun.* **6**, 7965 (2015).

57. Burnstock, G. Purinergic Signalling: Therapeutic Developments. *Front. Pharmacol.* **8**, 661

(2017).

58. Birkenfeld, A. L., Jordan, J., Dworak, M., Merkel, T. & Burnstock, G. Myocardial metabolism

in heart failure: Purinergic signalling and other metabolic concepts. *Pharmacol. Ther.* **194**,

132–144 (2019).

59. Burnstock, G. & Pelleg, A. Cardiac purinergic signalling in health and disease. *Purinergic*

Signal. **11**, 1–46 (2015).

60. Tran Diem H. & Wang Zhao V. Glucose Metabolism in Cardiac Hypertrophy and Heart

Failure. *J. Am. Heart Assoc.* **8**, e012673 (2019).

61. Martini Elisa *et al.* Single-Cell Sequencing of Mouse Heart Immune Infiltrate in Pressure

Overload–Driven Heart Failure Reveals Extent of Immune Activation. *Circulation* **140**, 2089–

2107 (2019).

62. Li, X. *et al.* Systemic Adenosine Triphosphate Impairs Neutrophil Chemotaxis and Host

Defense in Sepsis. *Crit. Care Med.* **45**, e97–e104 (2017).

63. Bao, Y., Chen, Y., Ledderose, C., Li, L. & Junger, W. G. Pannexin 1 Channels Link

Chemoattractant Receptor Signaling to Local Excitation and Global Inhibition Responses at

the Front and Back of Polarized Neutrophils. *J. Biol. Chem.* **288**, 22650–22657 (2013).

64. Wang, X. & Chen, D. Purinergic Regulation of Neutrophil Function. *Front. Immunol.* **9**, 399

(2018).

65. Sesma, J. I. *et al.* The UDP-sugar-sensing P2Y $_{14}$ receptor promotes Rho-mediated signaling

and chemotaxis in human neutrophils. *Am. J. Physiol.-Cell Physiol.* **303**, C490–C498 (2012).

66. Pitt, B. *et al.* The Effect of Spironolactone on Morbidity and Mortality in Patients with Severe

Heart Failure. *N. Engl. J. Med.* **341**, 709–717 (1999).

67. Ferreira, J. P. *et al.* Proteomic and Mechanistic Analysis of Spironolactone in Patients at Risk

for HF. *JACC Heart Fail.* (2021) doi:10.1016/j.jchf.2020.11.010.

68. Agah, R. *et al.* Gene recombination in postmitotic cells. Targeted expression of Cre

recombinase provokes cardiac-restricted, site-specific rearrangement in adult ventricular

muscle in vivo. https://www.jci.org/articles/view/119509/pdf (1997)

doi:10.1172/JCI119509.

69. Poon, I. K. H. *et al.* Unexpected link between an antibiotic, pannexin channels and

apoptosis. *Nature* **507**, 329–334 (2014).

70. Ackers-Johnson Matthew *et al.* A Simplified, Langendorff-Free Method for Concomitant

Isolation of Viable Cardiac Myocytes and Nonmyocytes From the Adult Mouse Heart. *Circ.*

Res. **119**, 909–920 (2016).

Figure 1: Beta-adrenergic receptor stimulation activates Pannexin 1 channels in cardiomyocytes *in vitro*

A) H9c2 were transfected with either control or Panx1-targeting siRNA for 72 hours and extracellular concentration of ATP was measured after 15 minutes of stimulation with isoproterenol (ISO) (2 and 20μM) or vehicle control. (N=4)

B) Flow cytometric analysis of Yo-Pro-1 dye uptake by control of Panx1-targeting siRNA treated H9c2 cells after stimulation with 20μM ISO (gated for live, annexin V⁻ cells). (N=4)

C) H9c2 cells were pretreated with spironolactone (50μM) or vehicle control and extracellular ATP levels were measured after 15 minutes of stimulation with isoproterenol (ISO) (2 and 20μM). (N=4)

D) Yo-Pro-1 dye uptake by H9c2 cells treated with isoproterenol (20µM) after pretreatment with spironolactone, carbenoxolone (CBX), or probenecid at the listed concentrations. (N=4)

Data are represented as mean $\pm$ standard error of the mean (SEM). Data shown are representatives of 2 independent experiments. Significance determined using One-way ANOVA (A, B, C, D) with Sidak's multiple comparison test for post-hoc analysis for comparisons between individual groups. ns = not significant See also S1 in reference to figure 1.

Figure 2: Generation and Characterization of a novel Pannexin 1 cardiomyocyte specific deletion mouse

A) Confocal micrographs of cardiomyocyte-rich whole heart sections from 12-week-old male Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice. Knockout of PANX1 in cardiomyocytes was assessed using immunofluorescent staining. Cardiac troponin T (cTnnt) was used to stain cardiomyocytes; DAPI was used as a nuclear stain. Left, representative micrographs with insets showing PANX1 punctate; middle, average Panx1 fluorescence intensity for individual mice; right, fluorescence intensities for individual micrographs showing demonstrating of PANX1 signal. (Panx1$^{fl/fl}$ N=4 mice N=13 images, Panx1^{MyHC6} N=3 mice N=9 images, 3-5 images per mouse averaged).

B) End-diastolic volume C) End-systolic volume D) Ejection fraction and E) Wall thickness

were measured in anesthetized 8- to 10-week-old male Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice

using *in vivo* 2D-echocardiography. (Panx1$^{fl/fl}$ N= 9 mice, Panx1^{MyHC6} N=8)

Data represented as mean $\pm$ SEM. Significance was determined by Welch's t-test (A-E).

ns = not significant. See also S2 in reference to figure 2.

Figure 3: Pannexin 1 deletion shifts cardiomyocyte metabolism from oxidative phosphorylation to glycolysis both *in vivo* and *in vitro*

A) Quantification of uniquely expressed genes from bulk RNA-Sequencing of isolated cardiomyocytes from untreated 12-week-old male Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice.

B) Volcano plot of differentially expressed genes with a cut off of p < 0.05. Individual genes are indicated with gene name and arrow.

C) Heatmap showing relative gene expression of glycolysis associated genes from isolated cardiomyocytes of untreated Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice plotted as the Log$_2$ (fold change) from Panx1$^{fl/fl}$ mouse. Purple heatmap represents -log(p-value) of two-tailed t-tests comparing fold change of Panx1$^{fl/fl}$ and Panx1^{MyHC6}. (Panx1$^{fl/fl}$ N=3 mice, Panx1^{MyHC6} N=3 mice)

D) Quantification of Basal extracellular acidification rate (ECAR) and

E) Maximal ECAR of control of Panx1-targeted siRNA treated H9c2 cells after stimulation with either vehicle or ISO (20µM) for 1 hour. (N=5)

F) Glucose uptake as measured by 2NBD-glucose fluorescence intensity of control or Panx1-targeted siRNA treated after 1 hour ISO (20µM) stimulation. (N=4)

Data represented as mean $\pm$ SEM. Significance determined using One-way ANOVA (D, E, F) with Sidak's multiple comparison test for post-hoc analysis for comparisons between individual groups.   ns = not significant. See also S3 in reference to figure 3.

Figure 4: Isoproterenol induced cardiac hypertrophy and dysfunction is abrogated in Panx1^MyHC6 mice

A) Schematic of experimental model of isoproterenol induced non-ischemic heart failure.

B) Normalized heart mass from 10–12-week-old Panx1^fl/fl or Panx1^MyHC6 mice treated for 14 days with isoproterenol (15 mg/kg/day), or saline control normalized to tibial length. (Saline: Panx1^fl/fl N=9, Panx1^MyHC6 N= 8; Isoproterenol Panx1^fl/fl N=11, Panx1^MyHC6 N= 9)

C) Representative confocal micrographs of wheat germ agglutinin (WGA) staining at 20X (Left) and 60X (Right) from Panx1^fl/fl or Panx1^MyHC6 mice treated with isoproterenol for

14 days. Cardiomyocyte cross sectional area was calculated from WGA. (Panx1$^{fl/fl}$ N=4 mice, Panx1^{MyHC6} N=4, data points represent average of 50 cells per mouse).

D) Representative M-mode images of Panx1$^{fl/fl}$ mice or Panx1^{MyHC6} after 14 days of isoproterenol treatment, visualized by 2D echocardiography.

E) Left, paired analysis of left ventricle end-systolic volume (µL) pre- and post-isoproterenol treatment; right, percent change from baseline after isoproterenol treatment. (Panx1$^{fl/fl}$ N=8, Panx1^{MyHC6} N=9)

F) Left, paired analysis of left ventricle end-diastolic volume (µL) pre- and post-isoproterenol treatment; right, percent change from baseline after isoproterenol treatment. (Panx1$^{fl/fl}$ N=8, Panx1^{MyHC6} N=9)

Data represented as mean $\pm$ SEM. Significance determined using One-way ANOVA (B with Sidak's multiple comparison test for post-hoc analysis for comparisons between individual groups; Student's t-test (C) and Welch's t-test (E, F) was used for comparison between two groups. Significance of paired data was determined using Mixed effects analysis (E, F) with Sidak's multiple comparison test for post-hoc analysis for comparisons between timepoints.   ns = not significant. See also S4, S5 in reference to figure 4.

Figure 5: Cardiomyocyte PANX1 knockout leads to transcriptional downregulation of fatty acid metabolism and immune activation pathways during non-ischemic heart failure

A) Quantification of differentially expressed genes identified in bulk RNA-Sequencing between whole hearts of Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice after 14 days of isoproterenol treatment. Differentially expressed gene determined by cut off of $p < 0.05$. Significantly changed, as determined by cut off of $p < 0.05$, KEGG Pathways and Gene Ontology (GO) Terms identified from differentially expressed genes plotted as ordered by -log(p-value).

B) Heatmap demonstrating genes identified in WikiPathways "Inflammatory response".

C) Heatmap demonstrating genes identified in WikiPathways "Fatty acid oxidation and Fatty acid beta-oxidation".

D) Heatmap demonstrating genes identified in WikiPathways "Purinergic signaling".

Gradient plotted as Log$_2$(Fold Change) of Panx1^{MyHC6} compared to Panx1$^{fl/fl}$. (Panx1$^{fl/fl}$

N=5, Panx1^{MyHC6} N=4)

Figure 6: Neutrophil infiltration is significantly blunted in Panx1^MyHC6 mice during heart failure induced by isoproterenol

A) Heatmap demonstrating genes identified in WikiPathway "Inflammatory response". Gradient plotted as Log_2(Fold Change) of Panx1^{MyHC6} isolated cardiomyocytes after 14-days of isoproterenol treatment compared to Panx1$^{fl/fl}$. Significantly decreased, as determined by cut off of $p < 0.05$, GO terms in Panx1^{MyHC6} cardiomyocytes compared with Panx1$^{fl/fl}$ cardiomyocytes. (Panx1$^{fl/fl}$ N=4, Panx1^{MyHC6} N=4 mice).

B) Flow cytometric analysis of CD45$^+$ cells as a percentage of total live cells isolated from Panx1$^{fl/fl}$ and Panx1^{MyHC6} hearts after 14-days of isoproterenol treatment as identified. Representative histograms of the CD45 populations from live parent gate. (Panx1$^{fl/fl}$ N=8, Panx1^{MyHC6} N=8 mice)

C) Flow cytometric analysis of neutrophils (CD11b$^+$, Ly6g$^+$) as a percentage of the CD45$^+$ cells in Panx1$^{fl/fl}$ and Panx1^{MyHC6} hearts after 14-days of isoproterenol treatment. Representative plots of the identified neutrophils from Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice hearts. (Panx1$^{fl/fl}$ N=8, Panx1^{MyHC6} N=8 mice)

D) Representative confocal micrographs of Ly6g staining at 60X from Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice treated with isoproterenol for 14 days. Ly6g was used to stain neutrophils; DAPI was used as a nuclear stain. Quantification of number of neutrophils, Ly6g$^+$ puncta, per 60X field in whole heart tissue from Panx1$^{fl/fl}$ and Panx1^{MyHC6} mice after 14-days of isoproterenol treatment. (Panx1$^{fl/fl}$ N=8, Panx1^{MyHC6} N=9 mice, 3-5 images per mouse averaged)

E) Flow cytometric analysis of macrophages (CD11b$^+$, F4/80$^+$) as a percentage of CD45$^+$ cells from Panx1$^{fl/fl}$ and Panx1^{MyHC6} hearts after 14-days of isoproterenol treatment. Macrophages expressing the co-stimulatory activation marker (CD86$^+$) and macrophages

expressing the anti-inflammatory marker (CD163$^+$, CD86$^-$) as a percentage of the macrophages (CD11b$^+$, F4/80$^+$). Flow cytometric analysis of dendritic cells (DC) (CD11b$^+$, CD24$^+$) as a percentage of CD45$^+$ cells from Panx1$^{fl/fl}$ and Panx1^{MyHC6} hearts after 14-days of isoproterenol treatment. (Panx1$^{fl/fl}$ N=6, Panx1^{MyHC6} N=6 mice)

Data represented as mean $\pm$ SEM. Significance determined using Welch's t-test (B, C, D, E) was used for comparison between two groups. ns = not significant. See also S6 in reference to figure 6.