maGENEgerZ: An Efficient AI-Based Framework Can Extract More Expressed Genes and Biological Insights Underlying Breast Cancer Drug Response Mechanism

Turki Turki[a*] and Y-h. Taguchi[b*]

[a] King Abdulaziz University, Department of Computer Science, Jeddah 21589, Saudi Arabia. Email: tturki@kau.edu.sa

[b] Chuo University, Department of Physics, Tokyo 112-8551, Japan. Email: tag@granular.com

Corresponding Authors. Turki Turki (tturki@kau.edu.sa) and Y-h. Taguchi (tag@granular.com)

Abstract

Understanding breast cancer drug response mechanism can play a crucial role in improving the treatment outcomes and survival rates. Existing bioinformatics-based approaches are far from perfect and do not adopt computational methods based on advanced artificial intelligence concepts. Therefore, we introduce a novel computational framework based on an efficient version of support vector machines (esvm) working as follows. First, we downloaded and processed three gene expression datasets related to breast cancer responding and non-responding to the treatments from the gene expression omnibus (GEO)

according to the following GEO accession numbers: GSE130787, GSE140494, and GSE196093. Our method esvm is formulated as a constrained optimization problem in the dual form as a function of λ. We recover the importance of each gene as a function of $\lambda, y,$ and x. Then, we select p genes out of n, provided as input to enrichment analysis tools, Enrichr and Metascape. Compared to existing baseline methods, results demonstrate superiority and efficiency of esvm achieving high performance results and having more expressed genes in (1) well-established breast cancer cell lines including MD-MB231, MCF7, and HS578T; and (2) breast tissues. Moreover, esvm is able to identify (1) various drugs including clinically approved ones (e.g., tamoxifen and erlotinib); (2) seventy-four unique genes (including tumor suppression genes such as TP53 and BRCA1); and (3) thirty-six unique TFs (including SP1 and RELA). These results have been reported to be linked to breast cancer drug response mechanism, progression, and metastasizing. Our method is available publicly in the maGENEgerZ web server at https://aibio.shinyapps.io/maGENEgerZ/.

Keywords: Breast cancer; drug response; gene expression; machine learning; AI application in cancer clinical trials

1. Introduction

The ability to elucidate various mechanisms underlying breast cancer drug response and resistance is a critical part in the clinical decision-making process, not only aiding in finding out the potential effectiveness of a drug compound but also spanning to (1) reducing the search space for candidate compounds; (2) having a greater awareness and management of probable adverse reactions before conducting clinical trials [1]; and (3) identifying potential drug targets associated with drug compounds. Studies have been conducted to analyze gene expression data obtained from biological experiments pertaining to breast cancer drug response to unveil various molecular mechanisms. Du et al. [2] utilized a bioinformatics approach to identify important genes that play a key role in overcoming breast cancer drug

resistance working as follows. First, two gene expression datasets were downloaded from the gene expression omnibus (GEO) database based on GEO accession numbers GSE28694 (Miller & Payne grades 4 and 5) and GSE28826 (Miller & Payne grades 1 and 2). The GSE28694 dataset had 13 samples treated as drug-sensitive group while the GSE28826 dataset had 28 samples treated as drug-resistant group. Both processed datasets were provided as input to limma to identify 255 differentially expressed genes (DEGs) with $p < 0.05$, assigned to enrichment analysis tool ClusterProfiler. The protein-protein interaction with the use of random walk identified three genes (i.e., PRC1, GGTLC1, and IRS1) that are involved in immune pathways and involved in the breast cancer drug resistance mechanism. Further validation of the importance of these three genes was performed using additional datasets from the GEO and TCGA databases.

Wu et al. [3] utilized a bioinformatics approach to identify a gene signature that aid in predicting neoadjuvant chemotherapy response for breast cancer patients. They downloaded a gene expression dataset from the GEO database according to the GEO accession number GSE25066. The processed dataset had 508 samples in which 16 samples were excluded because of missing data, resulting in 492 samples. To perform differential gene expression analysis, limma was applied to drug-resistant tumor samples against drug-sensitive tumor samples identifying 347 DEGs. Then, applying limma within the drug-resistant cell line samples against wild type cell samples to identify 296 DEGs. Then, identifying 36 genes common between the 347 and 296 DEGs. The 36 genes were provided as input to enrichment analysis finding out 12 hub genes considered as a gene-signature (HJURP, IFI27, RAD51AP1, EZH2, DNMT3B, SLC7A5, DBF4, USP18, ELOVL5, PTGER3, KIAA1324, and CYBRD1) from the PPI that has been validated to assess its discriminative power using lasso in which the same GSE25066 dataset was divided into training and validation sets.

Freitas et al. [4] performed a bioinformatics analysis to identify reliable biomarkers for adding carboplatin to the standard anthracycline/taxane treatment, which can aid in identifying triple negative

breast cancer (TNBC) patients achieving pathological complete response to neoadjuvant chemotherapy (NAC). Therefore, TNBC patients with expected poor clinical outcomes can be provided another treatment options. The processed gene expression data was for 66 patients in which 33 were treated with carboplatin + paclitaxel (composed of 19 having RD and 14 achieving PCR) while the remaining 33 were treated with paclitaxel (composed of 23 having RD and 10 achieving PCR). In the 33 patients treated with carboplatin + paclitaxel, they applied limma to identify 37 DEGs between RD and PCR while identifying 27 DEGs between RD and PCR patients treated only with paclitaxel. Moreover, identifying 24 DEGs between RD and PCR patients for the 66 patients. Then, selecting 10 statistically significant genes (BNIP3, ZBTB16, KCNB1, HAS1, HEMK1, TFF1, PLA2G4F, SNAI1, C5orf38 and GRIN2A) out of the 37 and 27 DEGs, and 3 statistically significant genes (ALDH1A1, MCM2, and CXADR) out of the 27 DEGs. These 13 genes acted as gene signatures and reported results demonstrated their feasibility to discriminate between patients having RD and those achieving PCR.

Stevens et al. [5] aimed to unveil the molecular mechanism behind inducing a chemotherapy-resistant in inflammatory breast cancer (IBC) patients. They had a dataset of 131 samples between IBC and non-IBC patients derived from several profiling methods distributed based on 14 samples using ChIP-seq profiling, 84 samples using RNA-seq profiling, 3 samples using single-cell RNA-seq profiling, and 30 samples using RNA-seq II profiling. The dataset was deposited into the GEO database with accession number GSE163397. Bioinformatics analysis coupled with enrichment analysis revealed that JAK2/STAT3 signaling is a key player in driving a chemo-resistance in IBC. Therefore, inhibition of JAK2/STAT3 coupled with the use of paclitaxel drug can overcome therapeutic resistance in IBC patients. Debets et al. [6] performed a bioinformatics analysis coupled with enrichment analysis to identify a molecular signature (ER2, HER4, ER, IGF1R, and Kalirin) predictive of treatment response and resistance in HER2-positive breast cancer patients. Other studies aimed to unveil various mechanisms leading to drug resistance in breast cancer patients [7-10].

Although current advancing in breast cancer drug mechanism within clinical testing mainly depends on bioinformatics-driven computational approaches with existing off-the-shelf tools, AI-driven computational frameworks are needed to unveil vast biological insights and to properly promote the use of AI in real clinical settings. The availability of such AI tools can help clinical oncologists to early avoid therapeutic targets associated with poor treatment responses and thereby advancing clinical understanding and reducing the search space for potential drugs with adverse effects.

Contributions. Our major contributions are outlined as follows:

(1) To unveil various molecular mechanisms pertaining to breast cancer drug response, we introduce an AI-driven computational approach consisting of an efficient variant of support vector machines (esvm) combined with enrichment analysis tools, Enrichr and Metascape [11-13].

(2) From the gene expression omnibus (GEO), we downloaded three gene expression datasets concerning breast cancer drug response according to the following GEO accession numbers: GSE130787, GSE140494, and GSE196093). Then, we provided each processed dataset as input to our method esvm and compared with four other baseline methods, including linear model for microarray (limma) [14], significance analysis of microarrays (SAM), *t*-test [15], and least absolute shrinkage and selection operator (lasso) for the identification of breast cancer cell lines and tissue identification. Enrichment analysis results demonstrate the superiority of esvm obtaining more expressed genes within breast tissue and breast cancer cell lines such as MD-MB231, MCF7, and HS578T. Additionally, performance results show that esvm has the highest area under the ROC curve obtained by esvm when compared to lasso and is 6.74 times more computationally efficient on average than the baseline SVM as reported in the conducted simulated study. Computational efficiency in esvm allows experimental biologists and computational scientists to study the effect of integrating various profiling data and to obtain results with a reasonable amount of time.

(3) Additional results from enrichment analysis report the following biological insights. We identified various drugs for breast cancer including FDA-approved ones such as Gemcitabine (Gemzar) and

tamoxifen (Nolvadex). Moreover, identifying 74 unique genes including tumor suppression genes such as TP53, PTEN, BRCA1, and RB1. Reporting 36 unique TFs including SP1, NFKB1, and RELA. All of which have been reported to play a key role in breast cancer drug response and resistance mechanism.

(4) For reproducibility of the analysis in this study, we made a publicly available implementation for our method esvm within the maGENEgerZ web server at https://aibio.shinyapps.io/maGENEgerZ/. Moreover, we included the processed datasets within the Supplementary Datasets folder. We also provided a maGENEgerZ_Screenshots.docx file within the Supplementary Information to show the use for our web tool.

2. Materials and Methods

2.1 Gene Expression Profiles

In this work, we retrieved three datasets from different gene expression experiments of different GEO accession numbers [16].

2.1.1 GSE130787: Dataset1

For this dataset derived from the gene expression experiment at the GEO database, we had 89 samples and 5267 genes. As a result, we encoded the dataset as an 89×5268 matrix including drug responses as a column vector. The 89 samples were distributed in terms of treatment into three groups as follows. Twenty-six samples for patients treated with docetaxel, carboplatin, and trastuzumab (TCH). Thirty-eight samples for patients treated with docetaxel, carboplatin, trastuzumab and lapatinib (TCHTy). Twenty-five samples for patients with docetaxel, carboplatin and lapatinib (TCTy). In terms of the distribution of drug responses, thirty-eight BC patients achieved pathological complete response (PCR) while fifty-one BC achieved residual disease (RD). The gene expression experiment was performed using the microarray

platform, Agilent-014850 Whole Human Genome Microarray 4x44K G4112F (Probe Name version). This dataset is referred to as Dataset1.

2.1.2 GSE140494: Dataset2

For this second dataset derived from the performed gene expression experiment, We had 91 samples and 5313 genes, which were approved as protein-coding genes (PCGs) by domain experts from HUGO Gene Nomenclature Committee (HGNC) Biomart at https://biomart.genenames.org/ [17, 18]. Thus, we encoded the dataset as a 91×5314 matrix, including a column vector for drug responses. The drug responses were distributed as follows. Nineteen BC patients achieved resistant response while seventy-two were sensitive to the treatment (i.e., docetaxel, followed by 5-fluorouracil, epirubicin, and cyclophosphamide (TFEC). The gene expression experiment was performed using the microarray platform, Affymetrix Human Genome U133 Plus 2.0 Array. This dataset is referred to as Dataset2.

2.1.3 GSE196093: Dataset3

For this third dataset, we had 736 samples and 118 genes. Consequently, we encoded the dataset as a 736×119 matrix, including a column vector for drug responses in which 256 BC patients achieved complete response (CR) while 480 had a failed complete response (FCR). We had 11 treatments in which the 736 samples were distributed accodingly as follows. Paclitaxel (169), Paclitaxel + ABT 888 + Carboplatin (63), Paclitaxel + AMG-386 (110), Paclitaxel + AMG-386 + Trastuzumab (18), Paclitaxel + MK-2206 (56), Paclitaxel + MK-2206 + Trastuzumab (31), Paclitaxel + Neratinib (105), Paclitaxel + Pembrolizumab (67), Paclitaxel + Pertuzumab + Trastuzumab (43), Paclitaxel + Trastuzumab (25), and T-DM1 + Pertuzumab (49). The gene expression data was performed using Reverse Phase Protein Array (RPPA) microarray at George Mason University. This dataset is referred to as Dataset3. Table 1 provides an overview for the three-studied datasets.

Table 1: Overview of the three studied breast cancer drug response datasets downloaded from the gene expression omnibus database.

Dataset	Samples	Responder	Non-responder	Genes	Platform	Organism	Experiment Type
GSE130787	89	38	51	5267	GPL6480	Homo sapiens	Expression profiling by array
GSE140494	91	72	19	5313	GPL570	Homo sapiens	Expression profiling by array
GSE196093	736	256	480	118	GPL28470	Homo sapiens	Protein profiling by protein array

2.2 Computational Framework

In Figure 1, we outline the main steps pertaining to our computational framework. In terms of the preprocessing part, biopsy samples were obtained from breast cancer patients. Then, collected samples were prepared and provided to a biological technology, measuring the gene expression levels [19]. In the machine learning part, the input data corresponds to a gene expression dataset, where x_i represents the ith sample and y_i is the associated drug response. In our study, y_i is a binary class label (e.g., {pathological complete response (PCR), residual disease (RD)}). The entire samples x_i (for $i = 1..m$) were encoded as an $m \times n$ matrix in which m and n are the number of samples and genes, respectively. All drug responses y_i (for $i = 1..m$) encoded as a $1 \times m$ column vector. To identify p important genes out of the n genes in which $p \ll n$, we used to find arguments (i.e., $w = [w_1 \ldots w_n]$ and $b \in R$) that minimize the objective function in Equation 1 subject to the linear constraints as in [20, 21]. After solving the optimization problem in Equation 1, weights in w correspond to the importance of genes in which higher weights indicate the more important these genes are. However, the main issue is as follows.

Figure 1: Flowchart of the introduced AI-based framework identifying drugs, drug targets, critical genes, and transcription factors in breast cancer. Figure created with BioRender.com.

The optimization problem in Equation 1 depends on w and *b,* where |W| *is* equal to *n,* which is way larger than *m* (i.e., $m \ll n$) in genomic sciences [22]. That makes the solution for the optimization problem in

Equation 1 computationally expensive, where the number of genes (n) is typically larger than the number of samples (m).

$$\min_{\mathbf{w},b} \frac{\|\mathbf{w}\|^2}{2} + C\sum_{i=1}^{m} \xi_i^k$$
$$\text{subject to } y_i(\mathbf{w} \cdot \mathbf{x}_i + b) \geq 1 - \xi_i, \ i = 1, 2, ..., m \tag{1}$$
$$\xi_i \geq 0$$

Therefore, we seek to solve the dual form of SVM (See Equation 2)[23]. It can be seen that the optimization problem now depends on finding the Lagrange multiplier (λ) in which each x_i is associated with λ_i. This is way faster than finding w in Equation 1.

$$\max_{\lambda} \sum_{i=1}^{m} \lambda_i - \frac{1}{2}\sum_{i=1}^{m}\sum_{j=1}^{m} \lambda_i\lambda_j y_i y_j K(x_i, x_j)$$
$$\text{subject to } 0 \leq \lambda_i \leq C \text{ and } \sum_{i=1}^{m} \lambda_i y_i, \ i = 1, 2, ..., m \tag{2}$$

.

In Equation 3, we recover w from λ in Equation 2 as follows:

$$w = \sum_{i=1}^{m} \lambda_i y_i x_i \tag{3}$$

For any x_i associated with $0 < \lambda_j < C$, b is recovered as

$$b = y_j - \sum_{i=1}^{m} \lambda_i y_i K(x_i, x_j) \tag{4}$$

where $K(x_i, x_j)$ is a similarity measure K (usually called kernel) of x_i and x_j.

A testing example z is predicted as

$$y^{'} = \text{sign}(\sum_{i=1}^{m} \lambda_i y_i K(x_i, z) + b) \tag{5}$$

where sign() is an indicator function mapping to 1 (corresponding to RD) if its argument is greater than or equal to 0. Otherwise, it is mapped to -1 (corresponding to PCR). In Equations 6-10, we show 5 popular kernels used with SVM.

Linear: $\quad K(x_i, z) = (x_i.z)$ $\hfill (6)$

Polynomial: $\quad K(x_i, z) = (x_i.z + 1)^d$ $\hfill (7)$

Gaussian: $\quad \exp^{-\sigma(\|x_i - z\|^2)}$ $\hfill (8)$

Sigmoid: $\quad \tanh(s(x_i.z) + a)$ $\hfill (9)$

Laplacian: $\quad \exp^{-\sigma(\|x_i - z\|)}$ $\hfill (10)$

When a linear kernel is used, the prediction model becomes as

$$y^{'} = \text{sign}(\sum_{i=1}^{m} \lambda_i y_i (x_i.z) + b) \tag{11}$$

We used CVXR in R to solve the dual form of SVM in Equation 2 and to find λ [24]. In terms of the enrichment analysis part, we uploaded the p genes as input to Enricher and Metascape, where these p genes are weighted with the top p weights ($|w_1| > |w_2| > ... > |w_p|$). Then, interpreting and identifying

biologically related terms. Including key expressed genes, drugs, drug targets, transcription factors, and others are provided in the next section.

3 Experiments and Results

3.1 Experimental Methodology

We compared our method esvm against the following baseline methods: significance analysis of microarrays (SAM), linear models for microarray data (limma) [14], significance analysis of microarrays (SAM), Student's t-test (t-test), and least absolute shrinkage and selection operator (lasso) [15]. The input to the 5 studied methods is a labeled gene expression data. Because the prediction in our efficient svm-based model (esvm) is defined as the $\text{sign}(\sum_{i=1}^{m} \lambda_i K(x_i, z) + b)$, we had to recover w as $\sum_{i=1}^{m} \lambda_i y_i x_i$ and then selecting p genes associated with the top p weights (i.e., w = $[|w_1|\ldots|w_p|]$), corresponding to the top p important genes. For lasso, as the model is expressed as $\beta_0 + \beta\text{x}$, we selected the p genes associated with top p coefficients (i.e., $\beta = [|\beta_1|\ldots|\beta_p|]$) (excluding genes associated with zero coefficients). For limma, SAM, and t-test baseline methods, genes were selected based on according to significantly adjusted p-values < 0.01.

To perform enrichment analysis and evaluate the results from biological perspective, we uploaded genes obtained from each method to Enrichr (https://maayanlab.cloud/Enrichr/) and Metascape (https://metascape.org/gp/index.html) [11, 13]. When retrieved terms are related to breast cancer, then a method that has terms associated with more genes is considered the superior method. Furthermore, we assessed the performance from the classification perspective against lasso as a baseline, reporting area under ROC curve (AUC) as a performance measure, followed by conducting an statistical significance test and reporting the running time. In this study, we utilized R to run the experiments [25]. Specifically, we used the CVXR package in R to aid in solving the formulated optimization problem [24]. We employed siggenes package to run SAM [26] and we employed the limma package in R to run limma using the the two functions, lmFit and eBayes [14]. In terms of the t-test, we employed the t.test function

within the stats package [25]. To run lasso, we used the glmnet package [27]. For SAM, limma, and *t*-test, to compute adjusted *p*-values, we employed the p.adjust function with the "BH", setting $p < 0.01$ as in [28, 29].

For Dataset1, the gene expression experiment is to analyse and evaluate neoadjuvant docetaxel and carboplatin plus trastuzumab and/or lapatinib for patients with HER2+ breast cancer, obtained from the school of medicine at department at the University of California, Los Angeles, USA. We retrieved gene expression profiles of patients responding to the treatment (labelled as pathological complete response (PCR)) and those not completely responding to the treatment (labelled as residual disease (RD)) from the gene expression omnibus (GEO) database (https://www.ncbi.nlm.nih.gov/geo/) with GEO accession number GSE130787. We utilized getGEO() function within GEOquery package [30] to download and get the gene expression data. We utilized the fData() function from the Biobase package [31] to process and get gene names, pData() function within the Biobase package was used to get drug responses (i.e., PCR and RD) associated with each sample, exprs() function within the Biobase package to get expression values, missForest() function within missForest package to impute missing values [32, 33]. We selected 5267 protein coding genes (PCGs) according to Biomart domain experts at HUGO Gene Nomenclature Committee (See subsection GSE130787: Dataset1).

For Dataset2, the gene expression experiment is for predicting neodjuvant chemotherapy response in early breast cancer, obtained from leibniz research centre for working environment and human factors, located at Dortmund, Germany. We retrieved gene expression profiles of breast cancer patients responding (labelled as PCR and pathological partial response (PPR)) and those not responding to treatment (labelled as pathological no change (PNC)) from the GEO database NIH under GEO accession number GSE140494. As in Dataset1, we used the five functions (i.e getGEO(), fData(), pData(), exprs(), and missForest()) to download, prepare and impute Dataset2. Moreover, we selected 5313 PCGs according to BioMart domain experts at HUGO Gene Nomenclature Committee (See subsection GSE140494: Dataset2).

For Dataset3, the gene expression experiment is for I-SPY 2 neoadjuvant chemotherapy/targeted-therapy trial for early-stage breast cancer patients with high risk, obtained from University of California, San Francisco, USA. Gene expression profiles pertaining to breast cancer patients responding to treatment (labelled as complete response) against those not responding properly to the treatment (labelled as failed complete response) were obtained from the GEO database under GEO Accession number GSE196093. As in Dataset2, we used the five functions (i.e getGEO(), fData(), pData(), exprs(), and missForest()) to download, prepare and impute Dataset3.

3.2 Results

3.2.1 Dataset1

In Table 2, we report terms obtained from Enrichr based on input genes provided via each method. Terms in Table 2(a) show expressed genes within breast tissues according to each method. The more expressed genes, the better the computational method is. Table 2(a) demonstrates that esvm outperformed all compared methods, obtaining 28 expressed genes (SLC22A3, FKBP10, GRP, STAB2, LPL, GLI3, ADCY5, OBP2A, SOSTDC1, APOD, HMGCS2, GABRE, CCL21, TMEM61, BBOX1, GFRA1, IGF1, BMP6, APLN, CAPN13, NR4A1, C1ORF116, ANGPTL7, KCNS3, SSPN, PGR, FGFR2, and LTF) within the breast tissue. The second-best method is limma with 14 expressed genes (TMEM86B, LRP1, MEOX1, PLCZ1, OBP2A, PPP1R1B, LRIG1, ECHDC3, MAFK, MZF1, A2M, RNF186, FMOD, and KANK3) within the breast tissue. The worst-performing method is lasso in which 8 expressed genes (DGAT1, RNASE7, FXYD1, VEGFB, ROR2, FGF1, HSPA12A, and PTPN14) are expressed in the breast tissue. In Table 2(b), we show retrieved terms (i.e., cancer cell lines) and expressed genes within NCI-60 Cancer Cell Lines. Our method, esvm, performed better than other methods, obtaining a total of 8 expressed genes within the retrieved breast cancer cell lines. Specifically, three genes (PNMT, COL13A1, BCAS1) were expressed within MD-MB231, three genes (MYO5C, GFRA1, FGFR2) were expressed

within MCF7, and two genes (COL13A1 and FAR2) were expressed within HS578T. The second-best method is limma (tied with sam), both having 4 expressed genes within two breast cancer cell lines. In terms of limma, one gene (ECHDC3) was expressed within MD-MB231, whereas three genes (RBM8A, USP18 and PI4K2A) were expressed within MCF7. For sam, one gene (CLSPN) was expressed within MD-MB231, three genes (TFAP2C, GFRA1, and CNNM3) were expressed within MCF7. The worst-performing method is lasso, having two expressed genes within breast cancer cell lines. These results demonstrate the superiority for our method (i.e., esvm) in identifying breast tissue as well as breast cancer cell lines used in cancer research. In Supplementary Table1_A and Table1_B, we include enrichment analysis results of ARCHS4 Tissues and NCI-60 Cancer Cell Lines, respectively.

In Figure 2(a), we provide a visualization of intersected genes produced by all computational methods. It can be seen that esvm has unique 94 genes out of 100 when compared to all other methods. This implies that each method generated a different list of genes. For example, limma, sam, ttest, and lasso had unique genes of 96, 95, 94, and 38, respectively. It can also be seen that the number of common genes between any pair of methods is at most 2. This indicates that each method generated a different gene set and similarity among methods is minimal. In Supplementary DataSheet1_B, we list the genes according to the UpSet plot in Figure 2(a).

Table 2: Enriched terms obtained from a) ARCHS4 Tissues and b) NCI-60 Cancer Cell Lines via Enrichr according to produced genes by each method when using Dataset1. The best result is shown in bold.

a) ARCHS4 Tissues					
Method	**Rank**	**Term**	**Overlap**	*P*-**value**	**Adjusted *P*-value**
esvm	10	BREAST (BULK TISSUE)	**28**/2316	5.92×10^{-6}	5.81×10^{-5}
limma	18	BREAST (BULK TISSUE)	14/2316	2.65×10^{-1}	9.99×10^{-1}
sam	19	BREAST (BULK TISSUE)	13/2316	3.72×10^{-1}	9.99×10^{-1}
t-test	49	BREAST (BULK TISSUE)	9/2316	8.32×10^{-1}	9.99×10^{-1}
lasso	1	BREAST (BULK TISSUE)	8/2316	9.50×10^{-2}	9.87×10^{-1}

b) NCI-60 Cancer Cell Lines					
Method	**Rank**	**Term**	**Overlap**	*P*-**value**	**Adjusted *P*-value**
esvm	12	MD-MB231	**3**/150	3.94×10^{-2}	2.50×10^{-1}
	33	MCF7	**3**/397	3.19×10^{-1}	7.21×10^{-1}
	24	HS578T	**2**/176	2.19×10^{-1}	6.63×10^{-1}
limma	45	MD-MB231	1/150	5.29×10^{-1}	7.74×10^{-1}
	29	MCF7	**3**/397	3.19×10^{-1}	6.74×10^{-1}
	-	HS578T	-	-	-
sam	46	MD-MB231	1/150	5.29×10^{-1}	8.85×10^{-1}
	29	MCF7	**3**/397	3.19×10^{-1}	7.83×10^{-1}
	-	HS578T	-	-	-
t-test	-	MD-MB231	-	-	-
	46	MCF7	2/397	5.93×10^{-1}	8.72×10^{-1}
	45	HS578T	1/176	5.87×10^{-1}	8.72×10^{-1}
lasso	26	MD-MB231	1/150	2.65×10^{-1}	5.11×10^{-1}
	-	MCF7	-	-	-
	29	HS578T	1/176	3.04×10^{-1}	5.24×10^{-1}

Figure 2: (a) UpSet plot of gene lists provided by the computational methods when using Dataset1. (b) Nine clusters in a protein-protein interaction based on genes of esvm coupled with Metascape. (c) Twelve transcription factors according to Metascape when coupled with genes from esvm. (d) Process and pathway enrichment analysis provided by Metascape according to the 100 genes of esvm.

As esvm demonstrated superiority among the other computational methods, we provide the 100 genes produced via esvm to Metascape for further enrichment analysis. Figure 2(b) displays the following 44 genes obtained from protein-protein interaction: GRP, TRH, LTF, TTR, APOD, CSSC4, CAPN9, CAPN13, S100A6, S100A2, CARTPT, DUSP13B, FKBP10, GGH, ERP27, POF1B, SOSTDC1, CEACAM5, BMP6, CHRDL2, GSTA1, QDPR, NDRG1, NR4A1, SOCs4, PPBP, CCL21, APLN, TRIM63, ADCY5, CALML4, MYO5C, RAB38, AQP3, TUBA3C, PTP4A3, FOS, MS4A2, MMP1, MMP3, MMP12, FGFR2, IGF1, and PGR. The enrichment analysis showed that 10 genes (BMP6, IGF1, MMP1, MMP3, MMP12, PPBP, S100A2, S100A6, CCL21, CHRDL2) were related to NABA MATRISOME ASSOCIATED, 7 genes (BMP6, IGF1, PPBP, S100A2, S100A6, CCL21, CHRDL2) were related to NABA SECRETED FACTORS, and 6 genes (BMP6, FGFR2, NR4A1, IGF1, MMP12, APLN) were related to positive regulation of epithelial cell proliferation. It has been reported that NABA MATRISOME ASSOCIATED and NABA SECRETED FACTORS pathways play a key role in breast cancer metastasis through its involvement with extracellular matrix proteins [34, 35]. Also, genes linked to positive regulation of epithelial cell proliferation biological process are related to induction of metastasis and inhibition of breast cancer cells apoptosis through the promotion of epithelial cell proliferation via estrogen [36, 37]. These results demonstrate the effectiveness of our computational method in unveiling important molecular mechanisms pertaining to breast cancer pathogenesis and metastasis.

Figure 2(c) reports 12 transcription factors (TFs): NFKBIA, FOS, RELA, STAT3, JUN, BRCA1, SP1, USF1, ETS1, CREB1, NFKB1, and MYC. BRCA1 is known to play a key role in various biological processes in breast cancer [38-42]. TFs such as ETS1 and STAT3 have been reported as potential therapeutic targets in breast cancer [43]. Suppression of NFKBIA and CREB1 have been reported to be related to the inhibition of breast cancer progression [44, 45]. These 12 TFs can (1) aid in understanding breast cancer molecular mechanism; and (2) act as potential therapeutic targets for breast cancer treatment. In Figure 2(d), genes provided via esvm are related to various biological processes and

pathways in breast cancer progression. The top enriched term is vitamin D receptor pathway, which ameliorates breast cancer via contributing to the growth regulation of breast cancer cells [46, 47]. Other enriched terms such as NABA MATRISOME ASSOCIATED, mammary gland development, and extracellular matrix organization have been reported to play a key role in various biological processes to breast cancer progression and treatment [48-52]. In Supplementary Enirchment_Dataset1, we list Metascape enrichment analysis results related to Dataset1.

In Table 3, we report terms (i.e., drugs) and expressed genes (i.e., drug targets) within IDG Drug Targets 2022. Tamoxifen and Fulvestrant are antiestrogen inhibitors (hormone therapy) that have been approved for breast cancer treatment (See Figure 3) [53-55]. Both drugs identified two drug targets, PGR, ATP1A2. Cisplatin is a chemotherapy used for breast cancer treatment [56, 57]. ATP1A2 was a drug target reported in association with cisplatin. The obtained biological knowledge contributes to more understanding of breast cancer progression and treatment. In Supplementary Table1_C, we provide enrichment analysis results pertaining to IDG Drug Targets 2022.

Table 3: Retrieved enriched terms from IDG Drug Targets 2022 via Enrichr according to uploaded genes from esvm.

Rank	Term	Class	Genes
35	Tamoxifen	Antiestrogen	PGR;ATP1A2
37	Fulvestrant	Estrogen Receptor Antagonist	PGR
64	Cisplatin	Platinum Coordination Complex	ATP1A2

Figure 3: Tamoxifen and fulvestrant anticancer drugs inhibiting estrogen, causing DNA damage and cell death. SERM is selective estrogen receptor modulator. SERD is selective estrogen degrader.

3.2.2 Dataset2

In Table 4(a), we report retrieved breast tissue terms associated with expressed genes while Table 4(b) reports breast cancer cell line terms with expressed genes after uploading genes produced via all computational methods. It can be seen from Table 4(a) that our method (esvm) performed better than its competing baseline methods. Particularly, esvm had breast tissue term with 20 expressed genes (COL15A1, PFKFB3, VCAM1, TAT, TNC, PLAT, LAMC2, ACTG2, NR4A1, CYP2A6, KRT19, KRT18, COL5A1, DUOXA1, SCNN1A, FOSB, KCNN4, CD300LG, LTF, and MPZL2) out of 2316. The second-best method was sam with 13 expressed genes (DSP, IGFBP5, ODF3, TACSTD2, KLK8, SLC5A6, EFEMP1, FABP4, PER3, SLPI, CRTAC1, STAB1, and DACT2) out of 2316. Limma and *t*-test were tied for third-best performing methods with both having 11 expressed genes within the breast tissue. Limma had the expressed genes, namely CAMSAP3, FABP4, JUP, TRAF4, SLFNL1, TRIM29, WNT9A, A2M, PCDH1, LOXL1, and SLC9A1 while t-test had these following genes: SLC22A23, MUCL1, GIPC3, PACS2, FZD7, PADI2, CLEC4F, CFB, KIAA0040, SYT7, and EGFR. The worst-performing method was lasso with 7 expressed genes (RECQL4, NR4A1, SLC44A4, GJD3, CLDN7, SHB, and LZTR1) out of 2316. Table 4(b) demonstrates enriched breast cancer cell lines terms from NCI-60 Cancer Cell Lines obtained via Enrichr. The best-performing method is esvm with a total of 7

expressed genes within MCF7 and HS578T breast cancer cell lines, distributed as follows. Six expressed genes (DCTN5, KRT19, DAAM1, KYNU, TRIM37, and MPZL2) out of 397 within MCF7 breast cancer cell line while 1 expressed gene (ACTG2) within the HS578T breast cancer cell line. In terms of MD-MB231 breast cancer cell line, esvm had no retrieved results. Therefore, results were designated as "-". The second-best performing method was sam, resulted in 5 expressed genes within MD-MB231 and MCF7 breast cancer cell lines. Two expressed genes (RHEB and GARNL3) out of 150 were expressed within MD-MB231 breast cancer cell line, whereas 3 genes (PIAS3, DCTN5, and CTCF) out of 397 were expressed within MCF7 breast cancer cell line**.** For HS578T breast cancer cell line, no retrieved results were reported for sam (See "-"). The worst-performing method was lasso with 3 expressed genes within MD-MB231, MCF7, and HS578T breast cancer cell lines. These results demonstrate the good performance results of esvm in identifying breast tissues and cancer cell lines. In Supplementary DataSheet2_A, we include genes obtained from computational methods, provided to Enrichr to derive enrichment analysis results. Additionally, enrichment analysis results related to Table 4 are included in Supplementary Table2_A and Table2_B. Figure 4(a) displays the UpSet plot in terms of intersection lists of genes of all computational methods. Ninety-one unique genes were attributed to esvm. Sam, limma, ttest, and lasso had 94, 93, 92, and 33 unique genes, respectively. These results indicate that each method incorporated different computational steps resulted in different lists of genes. The number of intersected genes between each pair of computational methods is upper bounded by 4. In Supplementary DataSheet2_B, we include gene lists of computational methods related to Figure 4(a).

Table 4: Enriched terms obtained from a) ARCHS4 Tissues and b) NCI-60 Cancer Cell Lines via Enrichr according to produced genes by each method when using Dataset2. The best result is shown in bold.

a) ARCHS4 Tissues					
Method	**Rank**	**Term**	**Overlap**	*P*-value	**Adjusted *P*-value**
esvm	21	BREAST (BULK TISSUE)	**20**/2316	1.00×10^{-2}	4.91×10^{-2}
limma	26	BREAST (BULK TISSUE)	11/2316	6.18×10^{-1}	9.99×10^{-1}
sam	11	BREAST (BULK TISSUE)	13/2316	3.72×10^{-1}	9.99×10^{-1}
t-test	17	BREAST (BULK TISSUE)	11/2316	6.18×10^{-1}	9.99×10^{-1}
lasso	6	BREAST (BULK TISSUE)	7/2316	1.03×10^{-1}	8.76×10^{-1}
b) NCI-60 Cancer Cell Lines					
Method	**Rank**	**Term**	**Overlap**	*P*-value	**Adjusted *P*-value**
esvm	-	MD-MB231	-	-	-
	4	MCF7	6/397	1.47×10^{-2}	2.75×10^{-1}
	58	HS578T	1/176	5.87×10^{-1}	7.72×10^{-1}
limma	-	MD-MB231	-	-	-
	43	MCF7	2/397	5.93×10^{-1}	8.36×10^{-1}
	-	HS578T	-	-	-
sam	32	MD-MB231	2/150	1.72×10^{-1}	4.48×10^{-1}
	41	MCF7	3/397	3.19×10^{-1}	6.46×10^{-1}
	-	HS578T	-	-	-
t-test	-	MD-MB231	-	-	-
	32	MCF7	3/397	3.19×10^{-1}	6.80×10^{-1}
	49	HS578T	1/176	5.87×10^{-1}	8.43×10^{-1}
lasso	14	MD-MB231	1/150	2.31×10^{-1}	6.54×10^{-1}
	33	MCF7	1/397	5.04×10^{-1}	6.54×10^{-1}
	15	HS578T	1/176	2.66×10^{-1}	6.54×10^{-1}

Figure 4: (a) UpSet plot of gene lists provided by the computational methods when using Dataset2. (b) Three clusters in a protein-protein interaction based on genes of esvm coupled with Metascape. (c) Eighteen transcription factors according to Metascape when coupled with genes from esvm. (d) Process and pathway enrichment analysis provided by Metascape according to the 100 genes of esvm.

As esvm performed better than baseline computational methods, we provided genes obtained from esvm to Metascape to unveil biological insights within breast cancer drug response. Figure 4(b) reports the following 12 genes obtained from PPI network: ACTG2, PRKAR2B, ALDH2, PRKACB, KRT19, KRT18, WASF3, DLG5, TRIM37, GSTM3, GSTA1, and CYP2A6. Four genes (KRT18, KRT19, PLCB4, and PRKACB) were related to Estrogen signaling pathway, which is reported to play a key role in breast cancer progression and treatment [58-60]. Three genes (CYP2A6, GSTA1, and GSTM3) were linked to chemical carcinogenesis - DNA adducts pathway, involved in cancer development [61, 62]. In Figure 4(c), we report the following 18 transcription factors: SP1, RELA, NFKB1, HIF1A, STAT3, ZEB1, FOXO3, NFE2L2, JUN, NRC1, SP3, EP300, E2F1, CEBPB, USF1, HDAC1, ETS1, and STAT1. STAT3 is involved in breast cancer progression [63]. E2F1 and EP300 have been reported to be involved in breast cancer development and metastasis [64-66]. These results demonstrate the importance of these TFs and such mutations or alterations can affect gene regulation and thereby contribute to breast cancer development. In Figure 4(d), the top enriched term was nuclear receptors meta-pathway. Ten genes (CYP2A6, GSTA1, GSTM3, HMOX1, ME1, S100P, SCNN1A, ABCC4, PLK2, and B3GNT5) were linked to nuclear receptors meta-pathway, which has been related to breast cancer cell growth via nuclear receptors such as estrogen receptors [67]. Moreover, the 13 genes (AGT, BCL6, CDKN2C, FGFR3, GJA1, TNC, MT1X, S100A8, S100A9, SYT1, SOCS2, SEMA3C, and CHPT1) were related to regulation of growth process, which is linked to breast cancer cell growth. Eleven genes (AGT, BIRC5, CCND2, FGFR3, GSTA1, GSTM3, HMOX1, CXCL8, LAMC2, PLCB4, and PRKACB) were related to Pathways in cancer, which is linked to breast cancer development and metastasis [68]. Six genes (BCL6, CCND2, CDKN2C, CXCL8, PLAT, and PROM1) were related to transcriptional misregulation in cancer, which is linked to mutations and altered gene expression in breast cancer [69]. Enrichment analysis results obtained from Metascape are provided in Supplementary Enrichment_Dataset2.

Table 5 shows drugs and drug targets within IDG Drug Targets 2022. Hydroxycarbamide and gemcitabine are ribonucleotide reductase enzyme (RNR) inhibitors that have been used for breast cancer

treatment (See Figure 5) [70, 71]. The two drugs are associated with RRM2 as a drug target [72]. Daunorubicin inhibits DNA replication and Cyclophosphamide causing damage to the DNA of cancer cells and thereby causing cancer cells to die. TOP2A was a drug target for Daunorubicin, whereas RRM2 was a drug target for Cyclophosphamide. In Supplementary Table2_C, we report enrichment analysis results for IDG Drug Targets 2022.

Table 5: Retrieved enriched terms from IDG Drug Targets 2022 via Enrichr according to uploaded genes from esvm.

Rank	Term	Class	Genes
5	Hydroxycarbamide	Antimetabolite	RRM2
13	Daunorubicin	Anthracycline	TOP2A
19	Cyclophosphamide	Alkylating Agent	RRM2
20	gemcitabine	Antimetabolite	RRM2

Figure 5: Hydroxycarbamide and gemcitabine inhibiting ribonucleotide reductase (RNR) enzyme, which inhibits the DNA replication during the cell cycle and inducing apoptosis.

3.2.3 Dataset3

After uploading produced genes via each method to Enrichr, we show retrieved breast tissue terms within ARCHS4 Tissues (See Table 6(a)) and breast cancer cell lines terms within NCI-60 Cancer Cell Lines (See Table 6(b)). From Table 6(a), we can see that esvm generated the best results. Particularly, esvm had 6 expressed genes (CSF1, CCND1, ERBB3, IRS1, ERBB2, and ESR1) out of 2316 within breast tissue, followed by sam and t-test, both having 4 common expressed genes (RET, CCND1, IRS1, and ERBB2) out of 2316 within breast tissue. Lasso had 2 (ESR1 and EGFR) expressed genes out of 2316 within breast tissue while limma was the worst-performing method having 1 expressed gene (ABL1) out of 2316 within breast tissue. In Table 6(b), esvm is also the best-performing method by having 3 expressed genes out of 397 within MCF7 breast cancer cell line. No results were associated with MD-MB231 and HS578T breast cancer cell lines. Therefore, we indicated results by "-". Both sam and t-test were tied by having 2 expressed genes out of 397 within MCF7 breast cancer cell line. As esvm, no reported results were found for the other two breast cancer cell lines (i.e., MD-MB231 and HS578T). The worst-performing method was limma where no reported results were found for the three breast cancer cell lines. These results demonstrate the superiority of esvm when identifying breast cancer tissue and cell lines. We include enrichment analysis results regarding Table 6 in Supplementary Table3_A and Table3_B.

Table 6: Enriched terms obtained from a) ARCHS4 Tissues and b) NCI-60 Cancer Cell Lines via Enrichr according to produced genes by each method when using Dataset3. The best result is shown in bold.

a) ARCHS4 Tissues					
Method	**Rank**	**Term**	**Overlap**	*P*-value	**Adjusted *P*-value**
esvm	3	BREAST (BULK TISSUE)	**6**/2316	5.28×10^{-1}	9.97×10^{-1}
limma	40	BREAST (BULK TISSUE)	1/2316	9.78×10^{-1}	9.78×10^{-1}
sam	4	BREAST (BULK TISSUE)	4/2316	2.99×10^{-1}	9.47×10^{-1}
t-test	4	BREAST (BULK TISSUE)	4/2316	2.99×10^{-1}	9.47×10^{-1}
lasso	9	BREAST (BULK TISSUE)	2/2316	4.12×10^{-1}	7.71×10^{-1}

b) NCI-60 Cancer Cell Lines					
Method	**Rank**	**Term**	**Overlap**	*P*-value	**Adjusted *P* value**
esvm	-	MD-MB231	-	-	-
	3	MCF7	3/397	7.68×10^{-2}	5.88×10^{-1}
	-	HS578T	-	-	-
limma	-	MD-MB231	-	-	-
	-	MCF7	-	-	-
	-	HS578T	-	-	-
sam	-	MD-MB231	-	-	-
	3	MCF7	2/397	8.14×10^{-2}	4.41×10^{-1}
	-	HS578T	-	-	-
t-test	-	MD-MB231	-	-	-
	3	MCF7	2/397	8.14×10^{-2}	4.41×10^{-1}
	-	HS578T	-	-	-
lasso	-	MD-MB231	-	-	-
	18	MCF7	1/397	2.13×10^{-1}	3.08×10^{-1}
	-	HS578T	-	-	-

In Figure 6(a), we show the UpSet plot showing intersection of produced genes among all methods when Dataset3 is used. From the leftmost, it appears that esvm differs from all other methods by having unique 20 unique genes. Limma and lasso have 17 and 6 unique genes, respectively. Esvm, sam and *t*-test

share 13 genes. Limma and esvm share 12 genes. Sam and t-test share 8 genes. Lasso and esvm share 3 genes. Sam, $\underline{t}$-ttest, and lasso share 2 genes. It can also be seen that the number of common genes between the remaining intersection of methods is 1. These results demonstrate that our method is different from remaining methods, attributed to different computational steps involved in the computations of esvm. In Supplementary DataSheet3_B, we report genes related to UpSet plot. Figure 6(b) reports the following 19 genes obtained from PPI network: AR, BIRC5, CCND1, RB1, STAT1, STAT3, ESR1, IRS1, PTEN, ERBB2, ERBB3, ALK, AKT1, MET, JAK2, IGFIR, EGFR, TP53, and MTOR. Thirteen genes (AKT1, ARAF, CCND1, EGFR, ERBB2, ERBB3, MTOR, IGF1R, JAK2, MET, PTEN, RAF1, and STAT3) were linked to EGFR tyrosine kinase inhibitor resistance, involved in resistance mechanism of EGFR inhibitor, and thereby breast cancer progression [73]. Twelve genes (AKT1, CCND1, ERBB2, ESR1, MTOR, IRS1, JAK2, NOS3, PTEN, RAF1, STAT1, and STAT3) were involved in leptin signaling pathway, which is related to breast cancer malignancy [74, 75]. Five genes (BIRC5, CCND1, ESR1, RB1, and CHEK2) were involved in PID FOXM1 PATHWAY, which has been reported as a crucial oncogenic transcription factor that promotes breast cancer progression and growth [76, 77].

In Figure 6(c), we report the following 20 transcription factors (TFs): TP53, BRCA1, HDAC1, RELA, STAT3, E2F1, SP1, NFKB1, YBX1, ESR1, NKX3-1, AR, VHL, PAX5, CTNNB1, PPARG, PGR, JUN, KDM4B, and DNMT1. Interestingly, TP53 mutation has been reported to be the most frequently occurring in breast cancer [78, 79]. Patients with BRCA1 mutations are at higher risk of developing breast cancer and thereby considered as an important biomarker [80, 81]. HDAC1 has been reported to be related to breast cancer cell proliferation [82]. These results demonstrate the importance of these TFs and can aim in developing therapeutic strategies for breast cancer treatment. Figure 6(d) reports top enriched terms regarding processes and pathways. Twenty-one genes (AKT1, ALK, BIRC5, AR, ARAF, CCND1, CASP7, EGFR, ERBB2, ESR1, MTOR, IGF1R, JAK2, MET, PTEN, RAF1, RB1, STAT1, STAT3, TP53, and FADD) were linked to Pathways in cancer (hsa05200), coinciding with recently reported results as one of the top enriched pathways in breast cancer [83].

a

b

c

d

Figure 6: (a) UpSet plot of gene lists provided by the computational methods when using Dataset3. (b) Three clusters in a protein-protein interaction based on genes of esvm coupled with Metascape. (c) Twenty transcription factors according to Metascape when coupled with genes from esvm. (d) Process and pathway enrichment analysis provided by Metascape according to the genes of esvm.

Thirteen genes (AKT1, ARAF, CCND1, EGFR, ERBB2, ERBB3, MTOR, IGF1R, JAK2, MET, PTEN, RAF1, and STAT3) were related to EGFR tyrosine kinase inhibitor resistance (WP4806), which was

reported as significant pathway associated with breast cancer [69, 84]. Twelve genes (AKT1, CCND1, ERBB2, ESR1, MTOR, IRS1, JAK2, NOS3, PTEN, RAF1, STAT1, and STAT3) were linked to Leptin signaling pathway (WP2034), reported to have a key role in breast cancer tumorigenesis [85]. Nine genes (AKT1, EGFR, ERBB2, ERBB3, ESR1, MTOR, IRS1, MET, and PTEN) were linked to PI3K/AKT Signaling in Cancer (R-HSA-2219528) in which its inactivation suppressing the proliferation of breast cancer cells and thereby inducing apoptosis [86]. Seven genes (AKT1, BIRC5, ATM, CASP7, RAF1, TP53, and FADD) were related to apoptosis (hsa04210), which plays a key role in controlling the excessive proliferation of breast cancer cells [87]. In Supplementary Enrichment_Dataset3, we include enrichment analysis results obtained from Metascape pertaining to Dataset3.

Table 7 reports drug terms and drug targets within IDG Drug Targets 2022. Ceritinib aims to inhibit anaplastic lymphoma kinase (ALK) enzyme and thereby blocking the ability of tumors to grow and promoting apoptosis (See Figure 7) [88]. Erlotinib inhibits the effect of tyrosine kinase enzyme on epidermal growth factor receptor (EGFR) and thereby preventing proliferation of cancer cells and inducing apoptosis. In Supplementary Table3_C, we report enrichment analysis results regarding drug terms within IDG Drug Targets 2022.

Table 7: Retrieved enriched terms from IDG Drug Targets 2022 via Enrichr according to uploaded genes using Dataset3, showing genes (column: Genes) associated with drugs (column: Term). Rank column shows the order of terms when retrieved.

Rank	Term	Class	Genes
1	Ceritinib	Tyrosine Kinase Inhibitor	ALK;JAK2;MET;IGF1R
2	Erlotinib	Tyrosine Kinase Inhibitor	ALK;ERBB3;ERBB2;JAK2;MET
3	Entrectinib	Kinase Inhibitor	ALK;JAK2;IGF1R

Figure 7: Ceritinib and erlotinib are anaplastic lymphoma kinase (ALK) and epidermal growth factor receptor (EGFR) inhibitors, receptively.

3.3 Models Introspection

Dataset1

In Figure 8, we aim to get computational insights pertaining to studied learning-based models applied in the Results section. Figure 8(a) demonstrated that our method esvm leads to non-zero weights while lasso in Figure 8(b) leading to a sparser representation and thereby many zero coefficients, attributed to the L1 penalty as in [20]. In Figure 8(c), it can be seen that lasso is ~ 6.14 times faster than esvm. Figures 8(d and e for esvm and lasso, respectively) demonstrate that prediction differences between breast cancer patients achieving pathological complete response against those having residual disease were statistically significant (p-value of all models $<2.2 \times 10^{-16}$, obtained from a t-test).

Figure 8: Classification models comparison for predicting drug response of TCH, TCHTy, and TCTy in breast cancer (BC) patients when Dataset1 is used. Gene importance when esvm (a) and lasso (b) are applied. Computational running time (c) for esvm and lasso. Boxplot and strip chart of drug sensitivity prediction for BC patients using esvm (d) and lasso (e). ROC curve (f) demonstrating the prediction performance. TCH is docetaxel, carboplatin, and trastuzumab. TCHTy is docetaxel, carboplatin, trastuzumab and lapatinib. TCTy is docetaxel, carboplatin and lapatinib. PCR is pathological complete response. RD is residual disease.

These results show that both induced models using Dataset1 are expected to be general predictors for

drugs with PCR and RD responses. For the ROC curves of esvm and lasso in Figure 8(f), both models

achieved an area under the ROC curve (AUC) of 1.00.

Dataset2

As shown in Figure 9, different aspects related to models are reported as follows. Figure 9(a) shows that

esvm leading to non-sparse representation in which the weight vector w consisting of non-zero weights.

On the other hand, Figure 9(b) demonstrates that lasso leading to a sparse representation in which many

of the coefficients β are zeros. Figure 9(c) displays that lasso is 44 times faster than esvm.

Figure 9: Classification models comparison for predicting TFEC drug response of in breast cancer (BC) patients when Dataset2 is used. Gene importance when esvm (a) and lasso (b) are applied. Computational running time (c) for esvm and lasso. Boxplot and strip chart of drug sensitivity prediction for BC patients sensitive against those resistant to the drug treatment. ROC curve (f) demonstrating the prediction performance. TFEC is docetaxel, 5-fluorouracil, epirubicin, and cyclophosphamide.

Figure 9 (d) for esvm demonstrates that prediction differences between sensitive breast cancer patients against those having resistant drug response were statistically significant (p-value $< 2.2 \times 10^{-16}$, obtained from a t-test). For lasso in Figure 9 (e), prediction differences between the two groups (i.e., BC patients sensitive to a drug against those having resistant response) were not statistically significant (p-value $= 1$, obtained from a t-test). These obtained results show that esvm is expected to be a general predictor for drug sensitivity and resistance. Lasso tends to be a specific predictor. Figure 9(f) displays the ROC curves for esvm and lasso, where the former has an AUC of 1.00 and the latter has an AUC of 0.947.

Dataset3

Figure 10 reports various computational aspects for studied models from classification perspective. It can be easily noticed that esvm doesn't shrink weights w to zero (Figure 11(a)) while lasso has sparser representation attributed to the L1 regularization shrinking coefficients β to zero (See Figure 10(b)). In terms of efficiency, Figure 10(c) shows that that lasso is 1974 times faster than esvm. In terms of generalization, Figure 10(d and e) demonstrates that prediction differences of esvm and lasso between breast cancer patients achieving complete response (CR) against those of failed complete response (FCR) were statistically significant (p-value $<2.2 \times 10^{-16}$, obtained from a t-test). However, esvm had a better AUC of 0.722 while lasso had a lower AUC of 0.555 (See Figure 11(f)).

Figure 10: Classification models comparison for predicting multiple drug combinations response in breast cancer (BC) patients when Dataset3 is used. Gene importance when esvm (a) and lasso (b) are applied. Computational running time (c) for esvm and lasso. Boxplot and strip chart of drug sensitivity prediction for BC patients sensitive against those resistant to the drug treatment. ROC curve (f) demonstrating the prediction performance.

These performance results indicate that prediction performance difference between esvm and lasso is 16.7% when AUC is considered.

Scalability

In Figure 11(a-j), we report the computational running time for increased dimensionality starting from 50000 dimensions to 500000 dimensions and fixing the number of rows to 100. The generation of X was done according to the uniform distribution $U(0,1)$, Y was generated in which we balanced the label distribution. When the number of dimensions is 50000 (See Figure 11(a)), lasso was 27 and 121.5 times faster than esvm and svm, respectively. Our method esvm was 4.5 faster than the baseline svm. The average running times for esvm, svm, and lasso spanning over all results in Figure 11(a-j) are 3.54 seconds, 23.88, and 0.275, respectively. That means our method, esvm, on average is 6.74 times faster than svm while lasso was 12.87 and 86.83 times faster than esvm and svm, receptively. These results demonstrate the computational efficiency of our method over the svm implementation using e1071 package in R . In Supplementary Running_Time, we include all running time results related to Figure 11(a-j).

Figure 11: Computational running time for our model esvm against baseline methods (svm and lasso) using simulated data of increased dimensionality.

4 Discussion

Identifying critical genes, drugs, drug targets, and transcription factors play a key role in unveiling the underlying drug response mechanism of breast cancer. Therefore, we introduced an AI-based computational framework functioning as follows. First, because gene expression datasets consist of many genes compared to the number of samples, the optimization problem formulation of svm in terms of finding weight vector w and bias term *b* is impractical (See Equation 1). Therefore, we employ the dual form of SVM optimization problem formulated in terms of finding lambda λ as shown in Equation 2. Then, we recover the weight vector w using Equation 3. Our method, esvm, takes as input gene expression data along with breast cancer drug responses, obtained from the GEO database. The output is a list of selected 100 genes according to the top 100 corresponding weights in w. Then, we performed enrichment analysis via providing the output genes to two enrichment analysis tools, Enrichr and Metascape. When Enrichr is considered, our method outperformed the baseline methods via having more expressed gene in breast cancer cell lines, including MD-MB231, MCF7, and HS578T. Moreover, our method esvm had more expressed genes in breast tissue. For Metascape, our method identified important genes including tumor suppressor genes (e.g., TP53 and BRCA1), TFs, drugs, drug targets, pathways and biological processes that play a key role in understanding breast cancer drug response mechanism.

As computational running time plays a key role in the gene selection process, our method esvm was way faster than the baseline svm. Although lasso was the fastest method, esvm had more expressed genes in breast cancer cell lines as well as breast tissues when compared to lasso. Therefore, the improvements in esvm are attributed to (1) superiority of computational efficiency when compared to the baseline svm; and (2) high performance results measured using the AUC when compared to lasso. Another advantage of esvm attributed to the computational efficiency is the computational feasibility to explicitly change the data representation and applying esvm to identify important genes. As a results, esvm can aid in analyzing gene expression data coupled with clinical data, and other profiling datasets.

Drug responses in this study were classified as follows. For Dataset1, the treated groups were classified during phase II trial as (1) pathological complete response referring to the absence of invasive cancer in axilla and breast; and (2) residual disease referring to the presence of invasive cancer in breast and axilla. In terms of Dataset2, treated groups after neoadjuvant therapy within Phase II multicenter trial were classified as (1) sensitive referring to patients completely responding to the treatment; and (2) resistant referring to not completely responding to the treatment. Regarding Dataset3, the treated groups were categorized during neoadjuvant I-SPY2 trial as (1) complete response when responding to the treatment; and (2) failed complete response when not completely responding to the treatment.

We specify parameters of methods in our study as follows. esvm($C = 2$), lasso($\lambda=0.05$), and svm (setting parameters associated with the linear kernel to their default values). Each method producing a gene list in which the probability of each method coinciding with our method esvm is estimated as follows. For Dataset1, Dataset2 and Dataset3, the probabilities are equal to $\frac{1}{\binom{5267}{100}}, \frac{1}{\binom{5313}{100}}$ and $\frac{1}{\binom{118}{100}}$, respectively; $\binom{n}{k}$ indicates the binomial coefficient in which n is the total number of genes in the considered dataset and k is to the total number of genes produced by esvm. Therefore, the odds of having a method producing results like ours are unlikely to occur.

For enrichment analysis results, the ranking results show the order of retrieved terms according to input genes expressed within terms. The overlab results take the form of X/Y, where X is the total number of genes produced by computational method out of total number of expressed genes in the considered term (e.g., MD-MB231). The higher the X, the better the method is. P-values and adjusted p-values indicate the significance of input genes provided to the enrichment analysis tool and are calculated according to Fisher exact test and Fisher exact test with corrected with Benjamini-Hochberg method, respectively [12]. It is worth noting that we considered the following terms (i.e., breast cancer cell lines) within NCI-60 Cancer Cell Lines category: MD-MB231, MCF7, and HS578T. These breast cancer cell lines are well-established in biology and medicine when analysing breast cancer as in [89-94].

5 Conclusion and Future Work

In this paper, we present a computational framework based on machine learning and enrichment analysis, unveiling critical genes, drugs, drug targets and other biological knowledge underlying breast cancer drug response mechanism. Our framework receives as an input a gene expression dataset pertaining to breast cancer patients responding and not responding to a treatment; we downloaded three different gene expression datasets from the GEO database according to the following accession numbers: GSE130787, GSE140494, and GSE196093. As $n \gg m$ arises challenge in computational genomics in which n and m are the number of genes and samples, respectively, we formulate our method according to the dual form in which we solve the optimization problem as a function of λ associated with m (instead of the formulation as a function of w associated with n). Then, we perform mathematical calculations to efficiently recover w as a function of three inputs $(\lambda, \mathrm{y}, \mathrm{x})$ and then identifying p important genes out of n, provided to enrichment analysis tools Enrichr and Metascape. In addition to (1) significantly achieving the highest performance results in terms of the area under curve; and (2) being more computationally efficient than the baseline SVM, results demonstrate that our method esvm outperformed existing baseline methods in (1) breast cancer cell line identification, showing more expressed genes; and (2) identifying breast tissue. Moreover, we reported several drugs (including tamoxifen, cisplatin, and erlotinib), 36 unique TFs (e.g., SP1, NFKB1, RELA), and 74 unique genes (including tumor suppression genes such as TP53, BRCA1, and RB1) that have been reported to be connected to drug response and resistance mechanism, progression, and metastasis of breast cancer. We made our computational method available publicly on the maGENEgerZ web server at https://aibio.shinyapps.io/maGENEgerZ/.

Future work includes (1) utilizing our framework to unveil various biological knowledge behind drug response mechanisms, related to different cancer types such as pancreatic cancer, liver cancer, and

multiple myeloma; (2) collaborating with clinical research physicians to apply our tool to analyze drug response mechanism in the neoadjuvant setting; and (3) integrating different profiling data related to cancer drug response and performing an assessment from biological perspective.

CRediT authorship contribution statement

Turki Turki: Conceptualization, Formal analysis, Methodology, Data curation, Software, Supervision, Writing - original draft, Visualization, Investigation. **Y-h. Taguchi:** Conceptualization, Formal analysis, Methodology, Data curation, Validation, Writing - original draft.

Conflict of interest

None declared.

Funding

This work is supported in part by funds from the Chuo University (TOKUTEI KADAI KENKYU).

References

1.   Tang, J., et al., *Drug target commons: a community effort to build a consensus knowledge base for drug-target interactions.* Cell chemical biology, 2018. **25**(2): p. 224-229. e2.
2.   Du, Y., et al., *Identification of Immune-Related Breast Cancer Chemotherapy Resistance Genes via Bioinformatics Approaches.* Frontiers in Oncology, 2022. **12**: p. 772723.
3.   Wu, J., et al., *A novel twelve-gene signature to predict neoadjuvant chemotherapy response and prognosis in breast cancer.* Frontiers in Immunology, 2022. **13**: p. 1035667.
4.   Freitas, A.J.A., et al., *Gene expression alterations predict the pathological complete response in triple-negative breast cancer exploratory analysis of the NACATRINE trial.* Scientific Reports, 2023. **13**(1): p. 21411.
5.   Stevens, L.E., et al., *JAK–STAT Signaling in Inflammatory Breast Cancer Enables Chemotherapy-Resistant Cell States.* Cancer Research, 2023. **83**(2): p. 264-284.
6.   Debets, D.O., et al., *Deep (phospho) proteomics profiling of pre-treatment needle biopsies identifies signatures of treatment resistance in HER2+ breast cancer.* Cell Reports Medicine, 2023. **4**(10).
7.   Jin, X., et al., *Chromatin-modifying protein 4C (CHMP4C) affects breast cancer cell growth and doxorubicin resistance as a potential breast cancer therapeutic target.* The Journal of Antibiotics, 2023: p. 1-9.

8.   Das, S., et al., *A novel computational predictive biological approach distinguishes Integrin β1 as a salient biomarker for breast cancer chemoresistance.* Biochimica et Biophysica Acta (BBA)-Molecular Basis of Disease, 2023. **1869**(6): p. 166702.

9.   Wu, M., et al., *Identification of chemoresistance-associated microRNAs and hub genes in breast cancer using bioinformatics analysis.* Investigational New Drugs, 2021. **39**(3): p. 705-712.

10.   Hoogstraat, M., et al., *Comprehensive characterization of pre-and post-treatment samples of breast cancer reveal potential mechanisms of chemotherapy resistance.* NPJ Breast Cancer, 2022. **8**(1): p. 60.

11.   Kuleshov, M.V., et al., *Enrichr: a comprehensive gene set enrichment analysis web server 2016 update.* Nucleic acids research, 2016. **44**(W1): p. W90-W97.

12.   Xie, Z., et al., *Gene set knowledge discovery with enrichr.* Current protocols, 2021. **1**(3): p. e90.

13.   Zhou, Y., et al., *Metascape provides a biologist-oriented resource for the analysis of systems-level datasets.* Nature communications, 2019. **10**(1): p. 1-10.

14.   Ritchie, M.E., et al., *limma powers differential expression analyses for RNA-sequencing and microarray studies.* Nucleic acids research, 2015. **43**(7): p. e47-e47.

15.   Tusher, V.G., R. Tibshirani, and G. Chu, *Significance analysis of microarrays applied to the ionizing radiation response.* Proceedings of the National Academy of Sciences, 2001. **98**(9): p. 5116-5121.

16.   Barrett, T., et al., *NCBI GEO: archive for functional genomics data sets—update.* Nucleic Acids Research, 2012. **41**(D1): p. D991-D995.

17.   Povey, S., et al., *The HUGO gene nomenclature committee (HGNC).* Human genetics, 2001. **109**: p. 678-680.

18.   Durinck, S., et al., *Mapping identifiers for the integration of genomic datasets with the R/Bioconductor package biomaRt.* Nature protocols, 2009. **4**(8): p. 1184-1191.

19.   Bi, M., et al., *Enhancer reprogramming driven by high-order assemblies of transcription factors promotes phenotypic plasticity and breast cancer endocrine resistance.* Nature cell biology, 2020. **22**(6): p. 701-715.

20.   Turki, T. and Y.h. Taguchi, *GENEvaRX: A novel AI-driven method and web tool can identify critical genes and effective drugs for Lichen Planus.* Engineering Applications of Artificial Intelligence, 2023. **124**: p. 106607.

21.   Turki, T. and Y. Taguchi, *A new machine learning based computational framework identifies therapeutic targets and unveils influential genes in pancreatic islet cells.* Gene, 2022: p. 147038.

22.   Taguchi, Y. and T. Turki, *Adapted tensor decomposition and PCA based unsupervised feature extraction select more biologically reasonable differentially expressed genes than conventional methods.* Scientific Reports, 2022. **12**(1): p. 17438.

23.   Mohri, M., A. Rostamizadeh, and A. Talwalkar, *Foundations of machine learning.* 2018: MIT press.

24.   Fu, A., B. Narasimhan, and S. Boyd, *CVXR: An R Package for Disciplined Convex Optimization.* Journal of Statistical Software, 2020. **94**(14): p. 1 - 34.

25.   Team, R.C., *R: A Language and Environment for Statistical Computing.* Journal of Statistical Software, 2008. **25**(1).

26.   Schwender, H., *siggenes: Multiple testing using SAM and Efron's empirical Bayes approaches.* R package version, 2012. **1**(0): p. 1-70.

27.   Friedman, J., T. Hastie, and R. Tibshirani, *Regularization paths for generalized linear models via coordinate descent.* Journal of statistical software, 2010. **33**(1): p. 1.

28.   Taguchi, Y. and T. Turki, *A new advanced in silico drug discovery method for novel coronavirus (SARS-CoV-2) with tensor decomposition-based unsupervised feature extraction.* PloS one, 2020. **15**(9): p. e0238907.

29.   Turki, T. and Y.h. Taguchi, *A new machine learning based computational framework identifies therapeutic targets and unveils influential genes in pancreatic islet cells.* Gene, 2023. **853**: p. 147038.

30.   Davis, S. and P.S. Meltzer, *GEOquery: a bridge between the Gene Expression Omnibus (GEO) and BioConductor.* Bioinformatics, 2007. **23**(14): p. 1846-1847.

31.   Huber, W., et al., *Orchestrating high-throughput genomic analysis with Bioconductor.* Nature methods, 2015. **12**(2): p. 115-121.

32.   Stekhoven, D.J. and M.D.J. Stekhoven, *Package 'missForest'.* R package version, 2013. **1**: p. 21.

33.   Stekhoven, D.J. and P. Bühlmann, *MissForest—non-parametric missing value imputation for mixed-type data.* Bioinformatics, 2011. **28**(1): p. 112-118.

34.   Zhao, W., et al., *Identification of PIMREG as a novel prognostic signature in breast cancer via integrated bioinformatics analysis and experimental validation.* PeerJ, 2023. **11**: p. e15703.

35.   Dai, Y.-H., et al., *Gene-associated methylation status of ST14 as a predictor of survival and hormone receptor positivity in breast Cancer.* BMC cancer, 2021. **21**(1): p. 1-14.

36.   Furuminato, K., et al., *The role of mesenchymal estrogen receptor 1 in mouse uterus in response to estrogen.* Scientific Reports, 2023. **13**(1): p. 12293.

37.   Zhang, X., et al., *Missing link between tissue specific expressing pattern of ERβ and the clinical manifestations in LGBLEL.* Frontiers in Medicine, 2023. **10**.

38.   Li, J., et al., *S100A9-CXCL12 activation in BRCA1-mutant breast cancer promotes an immunosuppressive microenvironment associated with resistance to immunotherapy.* Nature Communications, 2022. **13**(1): p. 1481.

39.   Choi, E., et al., *BRCA1 deficiency in triple-negative breast cancer: Protein stability as a basis for therapy.* Biomedicine & Pharmacotherapy, 2023. **158**: p. 114090.

40.   Sachsenweger, J., et al., *ABRAXAS1 orchestrates BRCA1 activities to counter genome destabilizing repair pathways—lessons from breast cancer patients.* Cell Death & Disease, 2023. **14**(5): p. 328.

41.   Tournant, F., *Stromal cells drive tumorigenesis in BRCA1 mutation carriers.* Nature Reviews Cancer, 2023: p. 1-1.

42.   Gajda-Walczak, A., et al., *New, fast and cheap prediction tests for BRCA1 gene mutations identification in clinical samples.* Scientific Reports, 2023. **13**(1): p. 7316.

43.   Chakraborty, S. and S. Banerjee, *Multidimensional computational study to understand non-coding RNA interactions in breast cancer metastasis.* Scientific Reports, 2023. **13**(1): p. 15771.

44.   Yang, W., et al., *Elevated expression of the rhythm gene NFIL3 promotes the progression of TNBC by activating NF-κB signaling through suppression of NFKBIA transcription.* Journal of Experimental & Clinical Cancer Research, 2022. **41**(1): p. 67.

45.   Chen, S., et al., *Cullin-5 deficiency orchestrates the tumor microenvironment to promote mammary tumor development through CREB1-CCL2 signaling.* Science Advances, 2023. **9**(3): p. eabq1395.

46.   Huss, L., et al., *Vitamin D receptor expression in invasive breast tumors and breast cancer survival.* Breast Cancer Research, 2019. **21**(1): p. 84.

47.   Sannappa Gowda, N.G., et al., *Quercetin activates vitamin D receptor and ameliorates breast cancer induced hepatic inflammation and fibrosis.* Frontiers in Nutrition, 2023. **10**.

48.   Ray, A. and P.P. Provenzano, *Aligned forces: Origins and mechanisms of cancer dissemination guided by extracellular matrix architecture.* Current Opinion in Cell Biology, 2021. **72**: p. 63-71.

49.   Jones, C.E., et al., *A miniaturized screening platform to identify novel regulators of extracellular matrix alignment.* Cancer research communications, 2022. **2**(11): p. 1471-1486.

50.   Wiseman, B.S. and Z. Werb, *Stromal effects on mammary gland development and breast cancer.* Science, 2002. **296**(5570): p. 1046-1049.

51. Hannan, F.M., et al., *Hormonal regulation of mammary gland development and lactation.* Nature Reviews Endocrinology, 2023. **19**(1): p. 46-61.

52. Batbayar, G., et al., *ASH2L, a COMPASS core subunit, is involved in the cell invasion and migration of triple-negative breast cancer cells through the epigenetic control of histone H3 lysine 4 methylation.* Biochemical and Biophysical Research Communications, 2023. **669**: p. 19-29.

53. Bradley, R., et al., *Aromatase inhibitors versus tamoxifen in premenopausal women with oestrogen receptor-positive early-stage breast cancer treated with ovarian suppression: a patient-level meta-analysis of 7030 women from four randomised trials.* The Lancet Oncology, 2022. **23**(3): p. 382-392.

54. Wang, J., et al., *Clinical efficacy of fulvestrant versus exemestane as first-line therapies for Chinese postmenopausal oestrogen-receptor positive/human epidermal growth factor receptor 2-advanced breast cancer (FRIEND study).* European Journal of Cancer, 2023. **184**: p. 73-82.

55. Torrisi, R., et al., *Predictors of fulvestrant long-term benefit in hormone receptor-positive/HER2 negative advanced breast cancer.* Scientific Reports, 2022. **12**(1): p. 12789.

56. Sultan, M.H., et al., *Characterization of cisplatin-loaded chitosan nanoparticles and rituximab-linked surfaces as target-specific injectable nano-formulations for combating cancer.* Scientific reports, 2022. **12**(1): p. 468.

57. Fatehi, R., et al., *Metformin enhances anti-cancer properties of resveratrol in MCF-7 breast cancer cells via induction of apoptosis, autophagy and alteration in cell cycle distribution.* Biochemical and Biophysical Research Communications, 2023. **644**: p. 130-139.

58. Bu, J., et al., *Dalpiciclib partially abrogates ER signaling activation induced by pyrotinib in HER2+ HR+ breast cancer.* Elife, 2023. **12**: p. e85246.

59. Bjørklund, S.S., et al., *Subtype and cell type specific expression of lncRNAs provide insight into breast cancer.* Communications Biology, 2022. **5**(1): p. 834.

60. Kanyomse, Q., et al., *KLF15 suppresses tumor growth and metastasis in Triple-Negative Breast Cancer by downregulating CCL2 and CCL7.* Scientific Reports, 2022. **12**(1): p. 19026.

61. Chen, H., et al., *A radiomics model development via the associations with genomics features in predicting axillary lymph node metastasis of breast cancer: a study based on a public database and single-centre verification.* Clinical Radiology, 2023. **78**(3): p. e279-e287.

62. Liao, H., et al., *Expression of the prognostic marker IL 8 correlates with the immune signature and epithelial mesenchymal transition in breast cancer.* Journal of Clinical Laboratory Analysis, 2023. **37**(3): p. e24797.

63. Bi, J., et al., *TMEM25 inhibits monomeric EGFR-mediated STAT3 activation in basal state to suppress triple-negative breast cancer progression.* Nature Communications, 2023. **14**(1): p. 2342.

64. Huo, Y., et al., *Analysis of breast cancer based on the dysregulated network.* Frontiers in Genetics, 2022. **13**: p. 856075.

65. Ring, A., P. Kaur, and J.E. Lang, *EP300 knockdown reduces cancer stem cell phenotype, tumor growth and metastasis in triple negative breast cancer.* BMC cancer, 2020. **20**: p. 1-14.

66. Li, C.H., et al., *The activation of EP300 by F11R leads to EMT and acts as a prognostic factor in triple negative breast cancers.* The Journal of Pathology: Clinical Research, 2023. **9**(3): p. 165-181.

67. Ma, S., et al., *Transcriptional repression of estrogen receptor alpha by YAP reveals the Hippo pathway as therapeutic target for ER+ breast cancer.* Nature communications, 2022. **13**(1): p. 1061.

68. Mo, D., et al., *A tRNA fragment, 5′-tiRNAVal, suppresses the Wnt/β-catenin signaling pathway by targeting FZD3 in breast cancer.* Cancer letters, 2019. **457**: p. 60-73.

69. McBean, B.N., et al., *Molecular mechanisms of intrinsic radioresistance in breast cancer.* Cancer Research, 2023. **83**(7_Supplement): p. 2401-2401.

70. Zuo, Z., et al., *Ribonucleotide reductase M2 (RRM2): Regulation, function and targeting strategy in human cancer.* Genes & Diseases, 2024. **11**(1): p. 218-233.

71. Gordon, D., et al., *Activator protein-1 (AP-1) signaling inhibits the growth of Ewing sarcoma cells in response to DNA replication stress.* Cancer Research, 2023. **83**(7_Supplement): p. 3532-3532.

72. Rudd, S.G., et al., *Ribonucleotide reductase inhibitors suppress SAMHD 1 ara CTP ase activity enhancing cytarabine efficacy.* EMBO molecular medicine, 2020. **12**(3): p. e10419.

73. Li, X., et al., *Can EGFR be a therapeutic target in breast cancer?* Biochimica et Biophysica Acta (BBA)-Reviews on Cancer, 2022: p. 188789.

74. Wang, Y., et al., *Leptin and leptin receptor expression as biomarkers for breast cancer: a retrospective study.* BMC Cancer, 2023. **23**(1): p. 260.

75. Hu, Y., et al., *Cancer-cell-secreted miR-204-5p induces leptin signalling pathway in white adipose tissue to promote cancer-associated cachexia.* Nature Communications, 2023. **14**(1): p. 5179.

76. Katzenellenbogen, B.S., V.S. Guillen, and J.A. Katzenellenbogen, *Targeting the oncogenic transcription factor FOXM1 to improve outcomes in all subtypes of breast cancer.* Breast Cancer Research, 2023. **25**(1): p. 76.

77. Ziegler, Y., et al., *Suppression of FOXM1 activities and breast cancer growth in vitro and in vivo by a new class of compounds.* npj Breast Cancer, 2019. **5**(1): p. 45.

78. Shahbandi, A., H.D. Nguyen, and J.G. Jackson, *TP53 mutations and outcomes in breast cancer: reading beyond the headlines.* Trends in cancer, 2020. **6**(2): p. 98-110.

79. Mitri, Z.I., et al., *Impact of TP53 mutations in Triple Negative Breast Cancer.* npj Precision Oncology, 2022. **6**(1): p. 64.

80. Liu, F., et al., *Overexpression of tripartite motif-containing 47 (TRIM47) confers sensitivity to PARP inhibition via ubiquitylation of BRCA1 in triple negative breast cancer cells.* Oncogenesis, 2023. **12**(1): p. 13.

81. Mateos, M.R.-C., et al., *17P New therapeutic target in triple-negative breast cancer for enhancing PARP inhibitor efficacy and stimulating the anti-tumour immune response.* ESMO Open, 2023. **8**(1).

82. Hu, Z., et al., *Histone deacetylase inhibitors promote breast cancer metastasis by elevating NEDD9 expression.* Signal Transduction and Targeted Therapy, 2023. **8**(1): p. 11.

83. Khanal, P., et al., *Systems and in vitro pharmacology profiling of diosgenin against breast cancer.* Frontiers in Pharmacology, 2023. **13**: p. 1052849.

84. Deng, Z., et al., *Curcumin and its nano-formulations: Defining triple-negative breast cancer targets through network pharmacology, molecular docking, and experimental verification.* Frontiers in Pharmacology, 2022. **13**: p. 920514.

85. Kong, X., et al., *Isoform-specific disruption of the TP73 gene reveals a critical role for TAp73γ in tumorigenesis via leptin.* eLife, 2023. **12**: p. e82115.

86. Lin, X., et al., *Oxypalmatine regulates proliferation and apoptosis of breast cancer cells by inhibiting PI3K/AKT signaling and its efficacy against breast cancer organoids.* Phytomedicine, 2023. **114**: p. 154752.

87. Yuan, L., et al., *Promoting apoptosis, a promising way to treat breast cancer with natural products: A comprehensive review.* Frontiers in Pharmacology, 2022. **12**: p. 801662.

88. Dong, S., et al., *Ceritinib is a novel triple negative breast cancer therapeutic agent.* Molecular Cancer, 2022. **21**(1): p. 138.

89.   Jiang, W., et al., *A mitochondrial EglN1–AMPK axis drives breast cancer progression by enhancing metabolic adaptation to hypoxic stress.* The EMBO Journal, 2023. **42**(20): p. e113743.

90.   Chen, J., et al., *Solasodine suppress MCF7 breast cancer stem-like cells via targeting Hedgehog/Gli1.* Phytomedicine, 2022. **107**: p. 154448.

91.   Balogh, B., et al., *Urocortin stimulates ERK1/2 phosphorylation and proliferation but reduces ATP production of MCF7 breast cancer cells.* Molecular and Cellular Endocrinology, 2022. **547**: p. 111610.

92.   Janacova, L., et al., *Catechol-O-methyl transferase suppresses cell invasion and interplays with MET signaling in estrogen dependent breast cancer.* Scientific Reports, 2023. **13**(1): p. 1285.

93.   Park, J.D., et al., *The ELK3-DRP1 axis determines the chemosensitivity of triple-negative breast cancer cells to CDDP by regulating mitochondrial dynamics.* Cell death discovery, 2023. **9**(1): p. 237.

94.   Lei, M., et al., *Gankyrin inhibits ferroptosis through the p53/SLC7A11/GPX4 axis in triple-negative breast cancer cells.* Scientific Reports, 2023. **13**(1): p. 21916.

Turki Turki received a B.S. in computer science from King Abdulaziz University, an M.S. in computer science from NYU.POLY, and a Ph.D. in computer science from the New Jersey Institute of Technology. He is currently an associate professor with the Department of Computer Science, King Abdulaziz University, Saudi Arabia. His research interests include machine learning, data science and bioinformatics. His research has been published in journals such as *Scientific Reports, IEEE Journal on Selected Topics in Signal Processing*, *Frontiers in Genetics and Computers in Biology and Medicine*. He is an editorial board member of *Sustainable Computing: Informatics and Systems, BMC Medical Genomics, PLOS ONE,* and *Computers in Biology and Medicine*.

Y-h. Taguchi received a B.S. degree in physics from the Tokyo Institute of Technology and a Ph.D. degree in physics from the Tokyo Institute of Technology. He is currently a full professor with the Department of Physics, Chuo University, Japan. His works have been published in leading journals such as *Physical Review Letters*, *Bioinformatics*, and *Scientific Reports*. His research interests include bioinformatics, machine learning, and nonlinear physics. He is also an editorial board member of *PloS ONE*, *BMC Medical Genomics*, *Medicine* (Lippincott Williams & Wilkins journal), *BMC Research Notes*, and *IPSJ Transaction on Bioinformatics*.