Spatially Resolved Transcriptomic Signatures of Hippocampal Subregions and

Arc-Expressing Ensembles in Active Place Avoidance Memory

Authors: Isaac Vingan[1], Victoria Sook Keng Tung[2], Shwetha Phatarpekar[3],

A. Iván Hernández[1,4,6], Oleg V. Evgrafov[2,3,5], Juan Marcos Alarcon[1,4,6,#]

Affiliations:

[1]School of Graduates Studies, Program in Neural and Behavioral Sciences, State University of

New York, Downstate Health Sciences University, Brooklyn, NY, USA

[2]School of Graduates Studies, Program in Molecular and Cell Biology, State University of New

York, Downstate Health Sciences University, Brooklyn, NY, USA

[3]Institute of Genomic Health, State University of New York, Downstate Health Sciences

University, Brooklyn, NY, USA

[4]Department of Pathology, State University of New York, Downstate Health Sciences

University, Brooklyn, NY, USA

[5]Department of Cell Biology, State University of New York, Downstate Health Sciences

University, Brooklyn, NY, USA

[6]The Robert F. Furchgott Center for Neural & Behavioral Science, State University of New

York, Downstate Health Sciences University, Brooklyn, NY, USA

#Corresponding author: Juan Marcos Alarcon (juanmarcos.alarcon@downstate.edu)

KEYWORDS

Spatial Transcriptomics, Gene Expression, Memory, Hippocampus, CA1, CA3, Dentate Gyrus,

Immediate Early Gene, IEG-Expressing Ensemble, Memory-associated Neuronal Ensemble

ACKNOWLEDGEMENTS

Supported by Senior Vice President of Research Seed Grant (State University of New York,

Downstate Health Sciences University) and partial support from NINDS grant R21NS091830.

36   Year: 2023

ABSTRACT

The rodent hippocampus is a spatially organized neuronal network that supports the formation of spatial and episodic memories. We conducted bulk RNA sequencing and spatial transcriptomics experiments to measure gene expression changes in the dorsal hippocampus following recall of active place avoidance memory. Our analysis focused on two specific levels of spatial resolution: hippocampal subregions and Immediate Early Gene (IEG) expressing cellular assemblies. Through bulk RNA sequencing, we examined the gene expression changes following memory recall across the functionally distinct subregions of the dorsal hippocampus. We found that training induced differentially expressed genes (DEGs) in the CA1 and CA3 hippocampal subregions were enriched with genes involved in synaptic transmission and synaptic plasticity, while DEGs in the dentate gyrus (DG) were enriched with genes involved in energy balance and ribosomal function. Through spatial transcriptomics, we examined gene expression changes following memory recall in putative memory-associated neuronal ensembles marked by the expression of the IEGs *Arc*, *Egr1*, and *c-Jun*. Within samples from both trained and untrained mice, the subpopulations of spatial transcriptomic spots marked by these IEGs were functionally and spatially distinct from one another. In only the hippocampus of trained mice, DEGs detected between Arc+ and Arc- spots were enriched in several memory-related gene ontology terms, including "regulation of synaptic plasticity" and "memory." Our results suggest that memory recall is supported by region-specific gene expression changes and transcriptionally distinct IEG expressing ensembles of neurons in the hippocampus.

INTRODUCTION

The neural operations supporting memory require the involvement of various brain systems interconnected through neural networks [1]. The activity of these networks is fine-tuned by experience through the selective recruitment of ensembles of neurons ascribed to contain the bits of information associated with memory [2]. Neuronal ensemble activity is shaped by synaptic plasticity mechanisms that modulate the weight and efficacy of the ensemble's synaptic connections [3, 4]. Changes in gene expression are a key underlying mechanism in synaptic plasticity [5], and recent studies have identified multiple profiles of gene expression associated with memory [6-8].

In the rodent hippocampus, encoding of spatial memory information is supported by the diversity of synaptic computations within and across its sub-fields [9, 10]. Each hippocampal subregion (i.e. Dentate Gyrus, CA1, CA2, CA3) shows distinct patterns of activity to process spatial information and to form or retrieve memories [11-16]. At the regional level, patterns of activity are conferred through the combination of cyto-architecture, local synaptic circuitry, neuronal cell types and synaptic inputs unique to each region [7, 13, 17-26]. At the single-cell level, the functional state of hippocampal neurons is defined by connectivity, hippocampal subregional location, and experience-dependent recruitment into a memory ensemble [2, 27].

The investigation of memory across molecular and functional levels necessitates an approach that connects the scales of hippocampal organization between cellular mechanisms and networks of cells. While the functional contribution of individual genes that underly synaptic plasticity at the cellular level are well-established [4, 28-31], it remains unclear what changes in gene expression across hippocampal subregions and subpopulations of the memory-associated neuronal ensembles characterize memory in the hippocampus.

Memory-associated neuronal ensembles are a sparsely distributed population of neurons that

strongly activate together during the acquisition and recall of a memory, and are thought to

encode the particular bits of information relevant to that memory [20, 32, 33]. Numerous studies

have identified Immediate Early Genes (IEGs) such as *Arc*, *c-Fos*, *Egr1*, *c-Jun*, *Npas4*, *Bdnf*, and

Fmrp to characterize memory-associated neuronal ensembles [2, 34-36]. In this study, the

location of memory-associated neuronal ensembles is inferred from the expression of IEGs in

spatial transcriptomic spots. The expression of *Arc* mRNA has been used to tag and characterize

the properties of memory-associated neuronal ensembles [37-39]. This proxy for the memory-

associated neuronal ensemble was used to explore the transcriptomic changes supporting

memory recall in the hippocampus. We characterized changes in gene expression across the

major subregions of the transverse dorsal hippocampus using bulk RNA sequencing of micro-

dissected dorsal hippocampal subregions, and investigated the spatial distribution of IEG-

associated gene expression through a spatial transcriptomic study of coronal dorsal hippocampal

slices.

RESULTS

APA trained mice exhibit place avoidance memory

To investigate gene expression changes induced by active place avoidance (APA) training in the hippocampal network, we conducted bulk RNA-sequencing and spatial transcriptomics (see Figure 1A for experimental timeline). In the APA, mice learn to avoid a 60° shock zone on a rotating circular arena [40, 41]. Mice trained in the APA showed a decrease in the number of shocks delivered during the acquisition trials (Figure 1B) and longer re-entry times to the shock zone during the memory retention test trials (Figure 1C).

Bulk RNA-seq transcriptomic profiling DG, CA3 and CA1 separate trained from untrained cohorts

To explore the molecular mechanisms underlying memory recall across the dorsal hippocampus, bulk RNA-sequencing was performed on microdissected DG, CA3 and CA1 hippocampal subregions. Principal component analysis (PCA) from all 3 subregions combined reveals a separation of samples by behavioral cohort (Figure 2A), indicating that differences in training contributes substantially to the overall variability of gene expression in our samples. Separation between samples is further accentuated when PCA is performed separately on data from each subregion (Figure 2C-D). Consistently, hierarchical clustering of gene expression profiles reliably grouped samples by their behavioral cohort (Figure 2E).

Synaptic plasticity and learning and memory-related biological processes are regionally enriched

We detected 1663, 606, 315 and 350 differentially expressed genes (DEGs) (adjusted p-value < 0.05) between trained and untrained samples for the combined hippocampal subregions and

separated CA1, CA3, and DG subregions, respectively (Figure 3A-D). 966 genes were

upregulated and 697 were downregulated in all hippocampal subregions. In CA1 393 genes were

upregulated and 213 were downregulated, in CA3 273 genes were upregulated and 42 were

downregulated, and in DG 117 genes were upregulated and 223 were downregulated.

In each comparison the majority of highly significant DEGs (FDR < 0.005) exhibited a modest

effect size ($|\log_2 \text{fold}| < 1.0$). Among these DEGs in the combined subregions with a high

significance and modest effect size, there are the genes *Ywhag*, *Bsn*, and *Nup153*, which are

related to cell signaling and morphogenesis [42-44]. In the CA1, we found *Camk2n1*, *Shank3*,

Shank1, and *Snap* 25 which are related to synaptic signaling and plasticity [45-48]. We found

Prckg, Ttr, and *Snhg11* in the CA3 which are related to memory [49-51]. *Kit*, *Srrm1*, and *Syn2*,

found in the DG, are related to cell morphogenesis and synaptic signaling [52-54].

To further elucidate the distribution of DEGs across hippocampal subregions we investigated the

overlap of significant DEGs (FDR < 0.05). We found the largest overlap of upregulated DEGs

between the CA3 and CA1 subregions with a total of 127 intersecting genes (Figure 3E). The

genes *Snap25, Shank1, Shank 3, Bsn and Ywhag* were identified in the overlap between the

upregulated DEGs of CA3 and CA1 and are known for their involvement in synaptic plasticity.

Ywhag was also found to be upregulated in the overlap of all three subregions.

Next, we performed Gene Ontology (GO) enrichment analyses to annotate each subregion (i.e.,

DG, CA3 and CA1) with GO terms overrepresented amongst the DEGs detected in each

subregional analysis [55]. In all hippocampal regions combined, we found DEGs were

significantly enriched with genes involved in synaptic machinery (Figure 3F & Supplemental

Figure 1F,G). Upregulated DEGs detected in both the CA3 and CA1 were enriched with genes

involved in synaptic transmission. Upregulated DEGs detected in the DG were enriched with

genes involved in neurogenesis and ATP production. In the DG downregulated DEGs were

involved in transcriptional machinery and epigenetic modifications (Figure 3F). Downregulated

DEGs in the CA1 and CA3 were not significantly enriched with any biological processes. These

data reveal a spatial distribution of gene expression across the dorsal hippocampus, defined by a

greater degree of similarity between the CA1 and CA3 than between either of these subregions

and the DG.

Differences in the spatial distribution of hippocampal gene expression between trained and

untrained animals

To further investigate the spatial distribution of DEGs associated with APA memory across the

hippocampus, we performed spatial transcriptomics on coronal sections containing the dorsal

hippocampus from one trained and one untrained mouse. Computational analyses of integrated

capture spots in the hippocampus of trained and untrained samples reveals distinct clusters (seen

in a Uniform Manifold Approximation and Projection (UMAP) plot) which map along

anatomical boundaries in a spatial transcriptomic plot (Figure 4A,B). Spatial transcriptomic data

were integrated with the Allen Brain Atlas to computationally annotate each spot with the cell-

type most prominently detected [56] (Figure 4C,D).

We detected 352 DEGs in all hippocampal spots (205 upregulated and 147 downregulated genes)

between trained and untrained samples. 75 DEGs in the CA1 cell layer (55 upregulated and 20

downregulated genes), 72 DEGs in the CA3 cell layer (54 upregulated and 18 downregulated

genes), and 187 DEGs in the DG cell layer (129 upregulated and 58 downregulated genes)

(Figure 5 A-D). Genes *Cnih2, Syn1,* and *Camk2b,* were differentially expressed in the CA1,

CA3, and DG, respectively. These genes are involved in synaptic plasticity and learning and

memory [57-59].

In the comparison of DEGs between all hippocampal spots, GO enrichment analyses identified

biological processes related to synaptic plasticity and synaptic signaling (Figure 5F &

Supplemental Figure 2B,C). In the CA1 and CA3 subregions, we found upregulation of genes

involved in biological processes also related to synaptic plasticity and synaptic signaling (Figure

5F). In the DG upregulated DEGs were enriched in biological processes involved in protein

expression and ribosomal function (Figure 5F). CA1 was the only subregion where

downregulated DEGs show enrichment for biological processes. Our results show a high

similarity in the biological functions overrepresented in both the spatial transcriptomic and bulk

RNA sequencing differential gene expression analyses between trained and untrained samples.

Arc-expressing hippocampal spots exhibit changes in the expression of memory-associated

genes and biological processes

Our data suggest that differences in behavioral conditioning contributed to variations in the

spatial distribution of synaptic plasticity related gene expression across the hippocampal

network. Differences in behavioral conditioning affect the recruitment of Arc-expressing

hippocampal neurons which form the memory-associated neuronal ensemble [60-63]. We

studied the *Arc*-expressing spatial transcriptomics spots as a proxy for the putative IEG tagged

memory-associated neuronal ensemble to assess changes in biological processes following APA

memory recall. Out of the 542 spots in the hippocampus from the trained sample, 195 spots had

detectable expression of *Arc* mRNA (*Arc+*). In the untrained sample, 169 out of 568

hippocampal spots were *Arc+* (Supplemental Figure 3G). In both samples, spots with the highest

level of *Arc* expression were located in the CA1 cell layer (Figure 6A,B).

To test whether *Arc*-expressing neuronal ensembles exhibit changes in the expression of genes

related to synaptic plasticity, we performed differential gene expression analyses between both:

Arc+ spots across trained and untrained samples (TR^{Arc+} vs. UT^{Arc+}), and between *Arc+* and *Arc-*

spots within each sample (TR^{Arc+} vs. TR^{Arc-} & UT^{Arc+} vs. UT^{Arc-}) (Figure 6 C-F). Comparison of

TR^{Arc+} vs. UT^{Arc+} spots detected 174 significant DEGs (85 upregulated and 89 downregulated

genes), and comparison of TR^{Arc-} vs. UT^{Arc-} detected 353 significant DEGs (113 upregulated and

240 downregulated genes) (Figure 6 C,D). *Bc1* and *Rac1*, known for their involvement in

learning and memory [64, 65], were found to be down- and upregulated, respectively, in the

analysis across both *Arc+* and *Arc-* spots. GO term enrichment analysis of upregulated DEGs

detected across the TR^{Arc+} vs. UT^{Arc+} showed enrichment with biological processes related to

synaptic signaling and protein translation (Figure 6 H & Supplemental Figure 3C,E).

Upregulated DEGs detected across the TR^{Arc-} vs. UT^{Arc-} spots enriched for genes related to

synaptic transport and plasticity. Conversely, across both TR^{Arc+} vs. UT^{Arc+} and TR^{Arc-} vs. UT^{Arc-}

spots, functions related to energy balance and protein translation were downregulated.

101 DEGs (95 upregulated, 6 downregulated) were detected within TR^{Arc+} vs. TR^{Arc-}, and 78

DEGs (47 upregulated, 31 downregulated) were detected within UT^{Arc+} vs. UT^{Arc-} (Figure 6 E,F).

Dkk3, *Cck*, and *Neurod6*, all genes related to learning and memory, were detected in *Arc+* spots

within the hippocampus from both the trained and untrained mice [66-68]. *Homer1*, *Sv2b*, and

Scn3b were among the 17 significant DEGs detected in both the analysis of TR^{Arc+} vs. TR^{Arc-}

spots and the bulk comparison of all subregions. While DEGs detected within UT^{Arc+} vs. UT^{Arc-}

spots were not significantly enriched with genes involved in any biological processes, TR^{Arc+} vs.

TR^{Arc-} spots were significantly enriched with upregulated DEGs involved in synaptic plasticity

(Figure 6 G & Supplemental Figure 3D,F).

IEG-expressing hippocampal spots comprise distinct gene expression profiles within

trained and untrained samples

Subsets of the complete memory-associated neuronal ensemble can be marked through the

expression of different IEGs [69, 70]. To evaluate the functional relationship between these

subsets of the ensemble we correlated the expression of 22 known neuronal IEGs [71, 72] with

the expression of *Arc*. Pairwise correlations of the expression of these IEGs identified *Egr1* and

c-Jun as the IEGs most and least correlated with *Arc*, respectively (Supplemental Figure 3 A,B).

The correlation coefficients for *Arc* and *Egr1* are $r(21) = .598$, $p = 2.24 \times 10^{-56}$ in the untrained

sample and $r(21) = .764$, $p = 5.45 \times 10^{-105}$ in the trained sample. The correlation coefficients for

Arc and *c-Jun* are $r(21) = -.057$, $p = 0.17$ in the untrained sample and $r(21) = -.105$, $p = 0.014$ in

the trained sample. Because of this, we performed differential gene expression analyses on *Egr1*

and *c-Jun* expressing spots across and within behavioral conditions.

In total, 380 out of 542 spots in the hippocampus of the trained animal and 256 out of 568 in the

untrained were found to express detectable levels of *Egr1* (Supplemental Figure 3G). In both

samples, spots with the highest level of *Egr1* expression were located predominantly in the CA1

cell layer, following the spatial distribution of *Arc* expression. For the *Egr1*-expressing spots,

555 DEGs (509 upregulated, 46 downregulated) were identified the trained animal, and 196

DEGs (150 upregulated, 46 downregulated) in the untrained animals (Figure 7 D,F). In total, we

found detectable levels of *c-Jun* in 308 out of 542 spots in the hippocampal sample from the

trained animal and 175 out of 568 in the untrained animal (Supplemental Figure 3G). In both

samples, spots with the highest level of *c-Jun* expression were located predominantly in the DG

granule cell layer. For the *c-Jun*-expressing spots, 480 DEGs (440 upregulated, 40

downregulated) were identified within the trained animal, and 166 DEGs (138 upregulated, 28

downregulated) within the untrained animal (Figure 7 J,L).

Due to the high degree of correlation between the expression of *Arc* and *Egr1*, and the low

correlation between the expression of *Arc* and *c-Jun*, we predicted that *Arc* and *Egr1*-expressing

spots would appear more functionally similar to one another than they would to *c-Jun*-expressing

spots. However, GO enrichment analyses of all three IEG expressing spot populations across

trained and untrained conditions showed similar sets of enriched biological processes (Figure 7M

& Supplemental Figure 3C,E), namely those involved in gene expression and energy balance.

In contrast to the within sample analyses of *Arc+* spots showing overrepresentation of biological

process only in the hippocampus of the trained mouse (Figure 6G), analyses of *Egr1+* and *c-*

Jun+ spots in the hippocampal samples of *both* trained and untrained mice revealed DEGs that

were enriched in numerous biological processes. Biological processes enriched among DEGs in

the *Egr1+* spots within the hippocampus of trained and untrained mice were similar, and

involved in synaptic plasticity and transmission (Figure 7N & Supplemental Figure 3C,E).

Additionally, in the hippocampus of the trained mouse, *Egr1+* spots show downregulation of

biological processes related to protein translation and ribosomal function. A distinct collection of

biological processes were enriched among DEGs detected in analysis of *c-Jun+* spots within the

trained and untrained samples. Biological processes involved in axonal growth and

neurotransmitter release were overrepresented in *c-Jun+* spots in both the hippocampal sections

from trained and untrained mice.

RT-qPCR of DEGs detected in the overlap between bulk RNA sequenced hippocampal

subregions

To further validate the spatial distribution of DEGs across the hippocampal network, we

performed RT-qPCR comparisons of relative gene expression in micro-dissected DG, CA3, and

CA1 subregions (6 trained and 4 untrained animals). To support the regionalized differential

expression of the genes identified in the bulk RNA sequencing data, we measured the expression

of *Kit*, *Sst*, *Cdh24* (identified in the DG, CA3, CA1 subregions, respectively). *Gapdh* expression

was measured as an internal control. Expression of *Slc6a7* was also measured. *Slc6a7* was found

to be upregulated in all regional analyses (Figure 3E, center sector). Relative gene expression

demonstrate a significant upregulation of *Kit* in the DG of trained animals (p-adj = 0.024). *Sst*

was also found to be upregulated in the DG (Supplemental Figure 4A). We found no significant

differences for the measured genes in CA3 (Supplemental Figure 4B). In CA1, *Cdh24* is

increased in trained compared to untrained samples although the difference did not reach

statistical significance (Supplemental Figure 4C). And, *Slc6a7* was not increased in any

subregion.

DISCUSSION

In this study, we used a combination of bulk RNA sequencing and spatial transcriptomics to identify changes in gene expression within major subregions of the dorsal hippocampus and within IEG-expressing spots following APA memory recall. We hypothesized that forming a memory-associated neuronal ensemble following memory training and recall leads to the enrichment of unique biological processes across the dorsal hippocampus. To test this hypothesis, we used a combination of bulk RNA sequencing and spatial transcriptomics to identify changes in gene expression within each major subregion of the dorsal hippocampus and within IEG-expressing spots following memory recall. Vanrobaeys et al. used spatial transcriptomics to study consolidation of a spatial memory and found distinct yet overlapping transcriptomic signatures across hippocampal subregions, with the CA1 and DG exhibiting upregulation of genes related to transcriptional regulation and protein folding, respectively [73]. The findings from this study illustrate the importance exploring the spatial distribution of gene expression profiles when investigating the memory-associated neuronal ensemble and the functional changes therein.

Regionalization of molecular mechanisms following memory recall

Different brain regions are recruited depending on the type of memory training an animal is exposed to [27, 74]. We found evidence supporting the regionalization of differential gene expression and overrepresentation of GO terms (corresponding to specific biological processes) across the dorsal hippocampus following spatial memory recall. Bulk RNA sequencing analyses between trained and untrained samples revealed significant similarities in the DEGs detected in CA1 and CA3 fields, but not between either of these subregions and the DG. Similarly, the

biological processes overrepresented amongst DEGs in the CA1 and CA3 subregions shared

more similarities than with those in the DG. Spatial transcriptomics also revealed regional

differences in DEG detection and biological process overrepresentation. DEGs detected via

spatial transcriptomics did not show considerable overlap compared with those detected via bulk

RNA sequencing. Such differences might be accounted for by the technical differences between

the two transcriptomic methodologies (addressed below).

Cells in the CA1 and CA3 subregions share similar morphologies and ontology as compared to

the dentate gyrus [75-77], which could account for our regionalized findings in the bulk RNA

sequencing study. During memory recall, the functional coupling of the CA3 and CA1

subregions [78, 79] is also reflected by the gene expression changes observed in this study. Our

data show that the spots in the CA3 and CA1 subregions have upregulated biological processes

involved in post synaptic machinery and synaptic plasticity, while the spots in the DG subregion

have upregulated processes involved in protein expression and energy balance, and have

downregulated processes involved in synaptic transmission. The cellular and molecular

modifications occurring during memory recall are known to include: 1) increases in neurogenesis

and protein synthesis in the DG [80-83] and 2) increased synaptic efficacy (e.g., LTP) across the

Schaffer collateral between the CA3 and CA1 [84, 85]. The overrepresentation of GO terms

involved in ATP metabolism and energy balance in the DG could indicate an understudied role

of the differential regulation of metabolic pathways in support of memory storage and recall in

the hippocampus [86, 87]

Circuitry in the hippocampus is differentially recruited depending on the phase, cognitive

demand, and valance of a memory[88-90]. We believe that the functional changes observed

during the recall phase of memory are one snapshot of the spatio-temporal modifications

occurring over the life of a memory in the brain. More studies are required to evaluate sequential

changes in regionalized hippocampal function with memory over time.

Technical considerations of using spatial transcriptomics to study memory-associated

neuronal ensembles

We utilized a combination of bulk RNA-seq and spatial transcriptomics to measure the gene

expression changes in the brain following memory recall. Both methodologies provided unique

utility and insight into the transcriptomic changes in the brain following memory and have

proven to be highly compatible [73, 91]. We observed strong regionalized differences between

training conditions, despite the fact that the changes we observed did not align between the two

transcriptomic techniques. This discrepancy can be accounted for by some of the chemical and

technical differences between the two methodologies. First, 10x's Visium spatial transcriptomic

platform permits a granular resolution of neuronal systems with its 55μm diameter capture spots

[92]. The resolution of this platform allows us to analyze gene expression differences between

specifically annotated populations of neurons in the cell layers of each hippocampal subregion,

rather than the entirety of the bulk section. Second, relative to bulk sequencing, 10x's Visium

spatial transcriptomics requires more tissue processing before RNA extraction and has lower

RNA detection sensitivity [93]. Consequently, some DEGs with lower levels of expression in our

bulk study may have gone undetected in the spatial transcriptomic study, thus skewing which

genes were profiled. Our study is amongst the first to examine gene expression changes within

individual cell populations across a spatially distributed neuronal network. The exploratory

nature of the present study, together with more accessible and higher resolution spatial

transcriptomic methods, is expected to develop into deeper characterizations of the mechanistic

architecture of memory.

***Arc*-expressing spots are enriched with genes involved in synaptic plasticity**

Using spatial transcriptomics, we were able to detect the spatial distribution of immediate early

gene expression in the dorsal hippocampus. Spatial transcriptomic spots sample regions 55μm in

diameter, which can contain more than a dozen cell bodies (depending on the brain region). We

inferred that groups of IEG-expressing memory-associated neurons would be sampled by these

spots, which we predicted would be enriched with genes related to synaptic plasticity. To

accomplish this, we targeted the expression windows of both immediate early genes and late

response genes by sacrificing mice following the retention test during the peak of IEG expression

and the beginning of late response gene expression [94, 95].

Data from the *Arc*-expressing spots highlight the distinct gene expression profile of TR^{Arc+} spots.

Only the upregulated DEGs detected in the analysis of TR^{Arc+} vs. TR^{Arc-} were enriched with any

biological process. This pattern of enrichment indicates both that TR^{Arc+} neurons are functionally

distinct from the surrounding TR^{Arc-} cells and the UT^{Arc+} cells. Specifically, TR^{Arc+} cells are

defined by an upregulation and enrichment of biological processes related to synaptic plasticity,

a number of which were also upregulated and enriched in the bulk differential expression

analyses of all hippocampal regions combined. These data suggest that the functional changes

induced in the *Arc*-expressing spots might substantially contribute to the differences observed

between training conditions in the analysis of bulk hippocampal subregions. We also detected

DEGs and biological processes between the TR^{Arc+} vs. UT^{Arc+} spots. In addition to biological

processes related to synaptic signaling, an abundance of biological processes related to energy

metabolism and protein translation –ATP metabolic process, ATP synthesis coupled proton

transport and cytoplasmic translation – were enriched in the TR^{Arc+} vs. UT^{Arc+} comparison.

It is not yet known if the functional changes that distinguish cell in a memory-associated

neuronal ensemble from the surrounding cells are the same functional changes that distinguish

these cells across training conditions. Our data suggests that there are two defining features of

the *Arc+* spots. The first is the upregulation of synaptic plasticity mechanisms when compared to

Arc- spots. This finding is supported by a recent study that found synaptic plasticity mechanisms

as a defining feature of a neuron in a memory-associated neuronal ensemble [96]. The second is

the increased recruitment of cellular machinery involved in energy production when compared to

the same spots in the hippocampus of the untrained animal. This result is supported by increases

in energy production and consumption during periods of high cognitive demand on neuronal

systems [97] In the case of the APA, the trained avoidance behavior is considered higher

cognitive demand than the untrained context habituation behavior [40, 98]. Arc+ spatial spots in

the trained animal are different from Arc- spots in the same animal and Arc+ spots in the

untrained animal.

Egr1- and *c-Jun*-expressing spots exhibit distinct transcriptomic profiles

The neuronal ensemble tagged by Arc is well studied for its role in memory storage and

maintenance [99], but the *Arc*-tagged ensemble encompasses only a subset of the neurons that

were active during a memory experience[69, 99, 100]. Subpopulations of this memory-associated

neuronal ensemble across brain systems might encode for multiple memory traces relevant to a

behavioral experience [1, 27, 101, 102]. Separate subpopulations of these neuronal ensembles are

tagged by at least one of the two dozen IEGs, the expression of which is linked to one of the

many patterns of strong neuronal activity elicited during a memory experience [27, 71, 103-105].

With spatial transcriptomics, we assessed gene expression profiles of IEG positive spots in the

dorsal hippocampus after the retention test in trained and untrained animals. We expanded our

investigation of the gene expression profile of the memory-associated neuronal ensemble to the

spots tagged by the IEGs *Egr1* and *c-Jun.* The expression of *Egr1* was most correlated with the

expression of *Arc,* while *c-Jun* was the least correlated.

Sparing some overlap, the populations of *Arc+, Egr1+,* and *c-Jun+* spots within both samples

were each enriched with genes involved in distinct collections of biological processes. DEGs

detected in the analysis of *Egr1+* and *c-Jun+* spots in *both* samples from trained and untrained

mice were enriched in numerous biological processes. However, DEGs detected in the analysis

of *Arc+* spots within *only* the trained sample were enriched with any biological processes,

namely those related to synaptic plasticity and synaptic function. The *Egr1+* spots within both

training conditions detected upregulated genes involved in biological processes related to

synaptic plasticity and synaptic transmission. *c-Jun+* spots within both training conditions

detected upregulated genes involved in biological processes related to axonal growth and

neurotransmitter release. Unlike the *Arc*-expressing spots in the trained sample, biological

process were enriched among downregulated DEGs detected *Egr1-* and *c-Jun*-expressing spots

the trained sample. Additionally, the spatial distribution of *Arc* and *Egr1* shows high levels of

expression in the CA1 cell-layer of trained and untrained samples, while the expression of *c-Jun*

is high in the DG. Expression of each IEG might be linked to particular patterns of neuronal

activation [106-108] [109] and output circuit function [89], highlighting the functional and

spatial diversity amongst recruited cells within memory associated neuronal ensembles. *Egr1* and

c-Jun expression are associated with fear memory and acute stress, respectively [110-112].

Perhaps, the differences observed amongst the *Egr1+* and *c-Jun+* spots may not be specific to

the behavior of trained active place avoidance behavior, but rather, sensitive to the shared

emotional, temporal and contextual components inherit to both long term memory experiences of

trained and untrained animals in the APA apparatus.

In contrast to the diversity of biological processes enriched among the DEGs between IEG-

expressing spots *within* each training condition, we found very similar collections of biological

processes in the GO enrichment analysis of IEG+ spots *across* training conditions. These

differentially regulated processes involved in energy production suggest that the training

experience induces a greater energy demand on the memory associated neuronal ensemble,

relative to the untrained experience. These data further highlight the need for greater energy

metabolism in higher cognitive demand memory experience, like that of training in the APA [87,

97].

Investigating memory across multiple scales in the brain

In summary, we performed an investigation of the transcriptomic profile of spatial memory recall

in the hippocampus using a combination of bulk RNA sequencing and spatial transcriptomics.

We found a regionalized pattern of gene expression with the CA1 and CA3 subregions being

enriched with genes involved in synaptic plasticity and transmission, while the DG subregion

was enriched with genes involved in protein expression and energy production. We also

discovered a unique transcriptomic profile of *Arc* expressing spots in the dorsal hippocampus of

trained mice enriched for upregulated genes involved synaptic plasticity and transmission.

Additionally, the transcriptomic profile of these *Arc* expression spots differed from those other

IEG expressing spots (i.e. *Egr1* and *c-Jun*).

This is amongst the first steps toward bridging our understanding of memory *across* scales

ranging from systems to cells to molecules. In this study, we investigate the spatial distribution

of memory-associated molecular function following the reactivation of a memory experience.

Through the joint investigation of the effectors of memory – their neuronal identities, spatial

location and functional changes – we open the opportunity for future studies that seek to

understand how memories are stored in the brain and how they change over time. Through

similar transcriptomic approaches that focus on space as a biologically relevant variable, we will

begin to map the distribution of memory-associated neuronal ensembles and their respective

gene expression changes as they undulate across the hippocampal network over the lifetime of

particular memory experiences.

METHODS

Animals

A total of 18 adult ArcCreERT2::eYFPflx mice [113] with C57BL/6 genetic background aged 3-4 months were used in this study (Bulk RNA sequencing – 3 male and 3 female, spatial transcriptomics – 2 male, RT-qPCR – 10 male). Mice were bred in-house at the SUNY Downstate Health Sciences University vivarium (Brooklyn, NY, USA). Mice were housed in groups of two to five until the start of the experiment, at which point they were single-housed in shoebox cages in a sound attenuation cubicle (Med Associates). For every experimental run, a minimum of two mice —each undergoing the same behavioral conditioning— were simultaneously housed in the sound attenuation cubicle. Ad libitum food and water was provided. Mice were randomly assigned to behavioral cohorts before the start of the experiment. Mice were handled daily for 3 days prior to the start of the experiment to reduce anxiety and improve voluntary approach. All animal procedures proposed are approved by, and will be performed following, the Institutional Animal Care and Use Committee guidelines at SUNY Downstate Health Sciences University.

Behavior

All procedures were performed in compliance with the Institutional Animal Care and Use Committee of the State University of New York, Downstate Health Sciences University. ArcCreERT2::eYFPflx male mice were trained in a hippocampus-dependent two-frame active place avoidance task. The place avoidance system consisted of a 40-cm diameter arena with a parallel rod floor that could rotate at 1 rpm. The position of the animal was tracked using PC-based software (Tracker, Bio-Signal Group Corp., Brooklyn, NY) that analyzed 30-Hz digital

video images from an overhead camera. Mice in the trained condition learned the

"Room+Arena−" task variant. Place avoidance of a 60° zone was reinforced by a constant

current foot shock (60 Hz, 500ms, 0.2mA) that was scrambled (5-poles) across pairs of the floor

rods. Rotation of the arena would carry the mouse into the shock zone unless the animal actively

avoided the zone. Entering the shock zone for more than 500 ms triggered shock. Additional

shocks occurred every 1.5 seconds until the animal left the shock zone. Measures of place

avoidance were computed by TrackAnalysis software (Bio-Signal Group Corp., Brooklyn, NY).

On day 1, mice received a 30-min trial with the shock off to habituate to the rotating arena.

Across the next two days the animals experienced 4 training trials, with 2 30-minute trials a day

with the activated shock with a 40 min inter-trial-interval.. Control (untrained mice) experienced

identical training conditions, except for the shock always being off.

Memory performance was assessed 24 hr after the final training session in a 10 minute retention

test with the shock off.

Microdissected bulk RNA sequencing

60 minutes after retention test, mice were euthanized and brains were extracted, washed in ice

washed in ice cold artificial cerebrospinal fluid, blocked, and mounted on a vibratome stage.

Hippocampal subregions (DG, CA3, CA1) were microdissected from 400 μm thick coronal live

tissue sections. Subregions from the dorsal hippocampus were extracted and microdissected

using microsurgical tools. Microdissected pieces of tissue from each animal were pooled by

subregion in 500 μL of prechilled TRIzol in a 1.5mL microcentrifuge tubes and stored at -80ºC.

Total RNA was extracted using the Zymo RNA mini prep kit (Zymo Research, Irvine, CA,

United States) according to the manufacturer's protocol. RNA quality was assessed on an

Agilent 2200 TapeStation (Agilent Technologies, Palo Alto, CA, United States). Samples with a

RIN greater than 7 were processed for library preparation. For library preparation NEBnext

ULTRA II RNA Library Preparation Kit for Illumina (New England Biolabs, Ipswich, MA,

United States) was used to: enrich mRNA from total RNA through rRNA depletion, fragment

mRNA into short fragments using fragmentation buffer, reverse transcribe mRNA into cDNA

using random primers, synthesize second-strand cDNA with DNA polymerase I, RNase H,

dNTP, and buffer. Double stranded cDNA fragments were preapered into sequenceable libraries

through multiple steps of purification using SPRIselect beads, end repair, A-tailing and adaptor

ligation. The ligated library was size selected using SPRIselect beads, PCR amplified and

sequenced using the Illumina Novaseq6000.

Spatial transcriptomics

60 minutes after retention test, mice were euthanized and brains were extracted and immediately

prepared for snap freezing in cuvettes of Tissue-Tek Optimal Cutting Temperature compound

(Sakura Finetek USA, Torrance, CA, United States) floating in a bath of methyl-butane chilled

by liquid nitrogen. Brain blocks were cryosectioned at -20ºC [114] and 10 µm thick coronal

sections containing the dorsal portion of the hippocampus were obtained (bregma -1.7 and -2.2

mm [115]). Collected tissue sections are trimmed to fit within the 5 mm x 5 mm capture area on

the Visium gene expression slide. Sections were mounted on a prechilled Visium gene

expression slide or a Tissue Optimization slide. A single tissue block was use to collect all tissue

sections mounted on the tissue optimization slide. One section of tissue per sample was mounted

in the capture areas of the Visium gene expression slide.

Tissue sections were fixed in chilled methanol and stained according to the Visium Spatial Gene

Expression User Guide [116] or Visium Spatial Tissue Optimization User Guide [117].

Tissues were permeabilized on the gene expression slide for 18 minutes which was selected from

the results of our tissue optimization experiment. Images were taken according to the Visium

Spatial Gene Expression Imaging Guidelines[118]. Brightfield histology images for the Visium

Tissue Optimization and Gene Expression slides were taken using the Leica Aperio CS2 Slide

Scanner at 20x magnification. Tissue optimization fluorescent images were taken on a Zeiss

LSM 800 (555nm LED,75% intensity, and 200ms exposure).

Tissue mRNA was extracted and libraries were prepared following the Visium Spatial Gene

Expression User Guide [116]. They were loaded at 300 pM and sequenced on a NovaSeq 6000

System (Illumina) using a NovaSeq S4 Reagent Kit (200 cycles, catalog no. 20027466,

Illumina), at a sequencing depth of approximately $250\text{–}400 \times 10^6$ read-pairs per sample.

Sequencing was performed using the following read protocol: read 1: 28 cycles; i7 index read:

10 cycles; i5 index read: 10 cycles; and read 2: 91 cycles.

RNA sequencing data analysis

Sequencing reads are processed at the Downstate Human Genomics Advanced Computing

Facility. Sequencing fastq files were demultiplexed with salmon, filtered and mapped to the

mouse MM10 reference genome [119] using STAR Aligner [120] and transcript reads were

quantified for each sample.

Bulk RNA sequencing analyses

Differential Gene Expression

Differential expression analysis was performed through DESeq2 on RNA counts between trained

and untrained sample groups. Genes with an of FDR below 0.05 were considered differentially

expressed genes/transcripts.

PCA

We ran 4 PCA analyses on normalized counts for samples. One analysis on all the the subregions

combined and 1 one the samples from each hippocampal subregion, 3 total.

Heatmap

Genewise Z-scores were calculated for the counts of each gene in each sample. Z-scores are

plotted as a heat map, where each row is a gene, and each column is a sample. Rows and

columns are then sorted in a dendrogram using Euclidean distance as a measurement for

similarity between any two given columns or rows.

Gene Ontology Enrichment Analysis

We used clusterProfiler's Overrepresentation analysis for GO terms [121] which uses the

hypergeometric test to calculate the enrichment Biological Processes, Cellular Components and

Molecular Functions amongst a list of differentially expressed genes (FDR < 0.05) against a

background of all detected genes. GO terms enriched with an FDR < 0.01 were considered

significant for our analyses. This analysis was also used for DEGs detected in the spatial

transcriptomic analyses. Dot plot figures show the top 10 GO terms ordered by significance for

each analysis plotted in the figured.

Venn Diagram

Vennplex software[122]was utilized to plot overlaps in the differentially expressed gene lists

from our regional differential gene expression analyses in a venn diagram format. This software,

stratifies gene lists by up- and down-regulated genes to display how many genes in overlapping

sectors are upregulated, downregulated or counterregulated (same gene with opposite effect sizes

in two regions).

Gene overlap statistics

The R-language package "SuperExactTest" [123] was used to statistically test the overlap of

gene names between regional lists of DEGs. The package uses the hypergeometric Fisher's Exact

test, against a background of all genes successfully compared in each differential gene

expression analysis. The software package also outputs an upset-style plot to visualize degree of

overlap between lists of DEGs and the level of significance of the overlap.

Spatial transcriptomics analyses

Reads from each sample were quantified and alighted to a predetermined loci or capture spots on

the histological image of each sample using 10x's space ranger software which references the

fiducial frame surrounding each capture area (10x Genomics Space Ranger 1.1.0).

Gene expression data were analyzed using the software package Seurat in the R coding

environment [124]. Spatial gene expression data were cropped to only include the capture spots

within the dorsal hippocampus. Data from each sample were then normalized, transformed and

integrated with one another using the default tools in Seurat. The integrated data set were further

normalized and clustered in PCA space. Clustered data can be visualized through UMAP

projections of the capture spots or in the 2D space of the tissue section. Cell type annotation was

performed by integrating our data with a random subset of 10,000 hippocampal cells from the

Allen Brain Atlas Cortex and Hippocampus Single Cell Taxonomy[56]. Predicted cell-type

identities were calculated and optimized based on gene expression similarity and anatomical

location.

Differential Gene Expression

Capture spots were grouped along two different criteria to calculate differential gene expression

in our samples. First, capture spots belonging to the anatomical cell layers of the Dentate Gyrus,

CA3 and CA1 regions were compared across training conditions together (All Regions) and

separately (region-wise). Next, within each training condition, hippocampal capture spots were

separated into two groups based on the detectable (>0) expression of the immediate early gene

(e.g. *Arc+* and *Arc-* groups). IEG expressing and not expressing groups of spots were compared

both within training conditions and across training conditions. Differential gene expression was

calculated using the Wilcoxon rank sum test (the default method for Seurat). Genes with an of

FDR below 0.05 were considered differentially expressed genes/transcripts.

Pairwise Correlations

Counts data for 23 known IEGs were extracted from hippocampal cell layer capture spots. In

each sample, Pearson's correlation coefficients were calculated pairwise for each IEG using the

base statistics in R-language. P values for Pearson's correlations were calculated for IEG's using

the base R-language correlation test.

RT-qPCR

Samples were extracted similarly to as described for bulk sample preparation to extract the dorsal

hippocampal subregions CA1,CA3, and DG from coronal sections of brain tissue. RNA

extraction using guanidinium-phenol-cholorform extraction with ethanol precipitation was

performed and quantified using spectrophotometry. cDNA was reverse transcribed from sample

bulk RNA using Thermo Fisher's SuperScript IV Kit (Thermo Fisher Scientific, Waltham, MA,

United States). Relative gene expression was quantified using the SYBR green master mix

system [125]. 10 µL reactions were prepared in the wells of a 384 well plate, hippocampal

regions were each prepared on separated plates. Each well contained 2.5ng of template cDNA

and 0.5 µM each of forward and reverse primers for one gene per well for a total reaction volume

of 10µL (sequences and melting temperatures can be found in table 1). Each gene was loaded in

triplicate in each plate to account for pipetting errors. Plates were incubated in the Bio-Rad

CFX384 Real-Time PCR Detection System (Bio-Rad Life Sciences, Hercules, CA, United

States) using the following schedule: 50°C for 5 min (1 cycle), 95°C for 10 min (1 cycle), 60°C

for 1 min (1 cycle), and 95°C for 15 s followed by 60°C for 1 min (40 cycles).

Gene	Direction	Sequence	Tm (°C)
Gahpdh	F	AGG TCG GTG TGA ACG GAT TG	57.47
	R	TGT AGA CCA TGT AGT TGA GTC A	52.84
Sst	F	ACC GGG AAA CAG GAA CTG G	57.44
	R	TTG CTG GGT TCG AGT TGG C	58.45
Kit	F	GCC ACG TCT CAG CCA TCT G	58.51
	R	GTC GCC AGC TTC AAC TAT TAA CT	55.14
Cdh24	F	AGC CTG TCC TTA TTG GCA AGC	57.92
	R	TGG GCC AGC AGC ACA TAT TG	58.33
Slc6a7	F	ACC TGG ATG TAG ACT TCG CAG	56.47
	R	CGC CAG ACA TTT CCC AAG C	56.97

Table 1. RT-qPCR primer sequences

Following 40 cycles of amplification wells were heated in a gradient from 55oC to 90oC to build

dissociation curves of primers and their transcripts to assess primer efficiency and specificity. Ct

values for each reaction are then automatically determined (roughly 20% of the plateau value).

Data were analyzed for relative gene expression quantification using the $\Delta\Delta$Ct method, using the

expression of *Gapdh* as the internal control. Relative expression values were normalized to the

expression value of the untrained group and Student's t-test were performed on each primer

target across training conditions using the ΔCt value for each comparison. Benjamini-Hochberg

correction was performed to adjust p-values for multiple hypothesis testing.

FIGURES

Figure 1. Trained mice learned to avoid the location of an unmarked shock zone

A) Experimental timeline. Mice were trained in the active place avoidance paradigm over the

course of 4 days following 3 days of handling. On day 0 they were habituated to a rotating arena

with no active shock zone (HT). On day 1 and 2 trained mice received two 30-minute trials with

an active shock zone. Untrained mice were not exposed to an active shock zone. On day 3 all

mice received a 10-minute retention test trial (exposed to the rotating arena with no active shock

zone). Mice were sacrificed 60 minutes following the end of the retention test, brains were

collected for further processing.

B) Training performance measured as number of shocks triggered. Trained mice learned to avoid

the location of the shock zone indicated by the decreased number of shocked triggered.

Exponential fit learning curves included with shaded error bars (SE).

C) Memory performance measured as time to second entrance in the retention test trial. Trained

mice showed higher time to second entrance during the retention test. Trained mean 375s ± 70.4

(SE), Untrained mean 58.7s ± 8.51 (SE).

Figure 2. APA training results in substantial changes in hippocampal subregion gene expression

profiles

A-D) Left, principal component analyses of all hippocampal subregions (A), CA1 (B), CA3 (C),

DG (D) illustrate a clear separation between trained and untrained samples along the axis that

explains the most variance. Right, heatmap with top 20 contributing genes to the first two

principal components of each are reported in the heat map to the right of each plot. Sample size

for the CA1 was 3 trained and 3 untrained (B). Sample size for the CA3 was 3 trained and 2

untrained (C). And Sample size for the DG was 3 trained and 2 untrained (D).

E) Normalized gene expression of the top 2000 most significant DEGs. Hierarchical clustering of

gene expression profiles groups samples by behavioral cohort.

Figure 3. Bulk RNA-seq of hippocampal subregions reveals similar expression profiles in CA1

and CA3 regions in trained samples

A-D) Volcano plot of DEGs between trained and untrained in the CA1, CA3 and DG samples,

combined and individually. See Supplemental Figure 1A-D for full sized plots. E) Overlaps in

regional DEGs demonstrate greatest similarity between CA1 and CA3 subregions. DEGs from

each regional analysis were stratified by direction of fold change. Overlaps in DEGs between the

CA3 and CA1 subregions were significant (Fisher's Exact Test). See Supplemental Figure 1E for

further details. F) Biological processes overrepresented amongst DEGs detected between

samples from trained and untrained animals for CA1, CA3 and DG subregions (all combined and

separated). Dot color reflects the statistical significance ($-\log_{10}(FDR)$) of the biological process

enrichment. Dot size reflects the number of detected DEGs mapped to the genes involved in a

given biological process. See Supplemental Figure 1 for enrichment of Cellular Component and

Molecular Function GO terms.

Figure 4. Clustering and cell-type annotation reflect anatomical boundaries of the hippocampal

subregions

A,B) UMAP plot (left) and spatial transcriptomic maps of hippocampal spots (right) from trained

and untrained mouse samples. Color reflects clustering of hippocampal spots. C,D) UMAP plot

(left) and cell-type annotated spatial transcriptomic maps of hippocampal spots (right) in samples

from the trained and untrained mouse.

Figure 5. DEGs detected between trained and untrained animals within hippocampal subregions

using spatial transcriptomics are enriched with genes involved in similar biological processes to

those seen in bulk RNA-seq

A-D) Volcano plot of DEGs between all hippocampal (A), CA1 (B), CA3 (C), and DG (D) cell-

layer spots from trained and untrained mice. E) Overlap of regionally detected DEGs stratified

by direction of fold change. Overlaps were tested with Fisher's exact test. See Supplemental

Figure 2A for further details. F) Biological processes overrepresented amongst DEGs detected

between samples from trained and untrained animals for hippocampal samples or all

hippocampal spots, and spots from the separated CA1, CA3 and DG cell layers. Dot color

reflects the statistical significance ($-\log_{10}$(FDR)) of the biological process enrichment. Dot size

reflects the number of detected DEGs mapped to the genes involved in a given biological

process. See Supplemental Figure 2 for enrichment of Cellular Component and Molecular

Function GO terms.

Figure 6. Arc-positive spots in the trained sample are enriched with genes involved in synaptic

plasticity

A,B) UMAP plot (left) and spatial transcriptomic maps of hippocampal spots (right) in the

trained and untrained samples. Colors show normalized expression of the *Arc* gene in

hippocampal cell layer spots. C,D) Volcano plots of differential gene expression of *Arc* positive

(C) and negative (D) spots across behavioral conditions. E,F) Volcano plots of differential gene

expression between *Arc* positive and negative spots within trained (E) and untrained (F) samples.

G) Biological processes overrepresented amongst DEGs detected in the analysis of *Arc*-

expressing spots within trained and untrained samples. Dot color reflects the statistical

significance ($-\log_{10}$(FDR)) of the biological process enrichment. Dot size reflects the number of

detected DEGs mapped to the genes involved in a given biological process. H) Biological

processes overrepresented amongst DEGs detected in the analysis of *Arc*-expressing spots across

trained and untrained samples. Dot color reflects the statistical significance ($-\log_{10}$(FDR)) of the

biological process enrichment. Dot size reflects the number of detected DEGs mapped to the

genes involved in a given biological process.

Figure 7. Spatial transcriptomics reveals distinct collections of biological processes enriched

among *Arc*, *Egr1*, and *c-Jun*-expressing spots

A,B) UMAP plot (left) and spatial transcriptomic maps (right) of hippocampal spots in the

trained and untrained sample. Color reflects the normalized expression of Egr1 in the

hippocampal cell layer spots. C,D,E,F) Volcano plots of differential gene expression of Egr1

positive (C) and negative (D) spots across behavioral conditions and within trained (E) and

untrained (F) samples. G,H) UMAP plot (left) and spatial transcriptomic maps (right) of

hippocampal spots in the trained and untrained samples. Color reflects normalized expression of

c-Jun in the hippocampal cell layer spots. I,J,K,L) Volcano plots of differential gene expression

of c-Jun positive (I) and negative (K) spots across behavioral conditions and within trained (J)

and untrained (L) samples. M) Biological processes overrepresented amongst DEGs in Arc,

Egr1, and c-Jun-expressing spots between trained and untrained samples. Dot color reflects

statistical significance (-log10(FDR)) of biological process enrichment. Dot size reflects number

of detected DEGs mapped to the genes involved in a given biological process. N) Biological

processes overrepresented amongst DEGs in Arc, Egr1, and c-Jun-expressing spots across

trained and untrained samples. Dot color reflects statistical significance (-log10(FDR)) of

biological process enrichment. Dot size reflects the number of detected DEGs mapped to the

genes involved in a given biological process.

 SUPPLEMENTAL FIGURES

Supplemental Figure 1. Trained CA1 and CA3 subregions express similar DEGs and

postsynaptic cellular component GO terms

A-D) Volcano plot of DEGs between trained and untrained CA1, CA3 and DG samples,

combined and individually. Triangular points mark data out of the frame of the main text figure.

E) Upset plot of overlaps in regional DEGs demonstrates greatest similarity between CA1 and

CA3 samples. DEGs from each regional analysis were stratified by direction of fold change. In

the upset plot, each bar represents the number of genes in an intersection of lists of DEGs. The

intersecting lists is represented by the dots underneath the bar. The hashed line on each bar

represents the expected intersection of two identically sized random lists. Overlaps were tested

with Fisher's exact test using the list of all expressed genes as a background. The color of the

bar represents the -$\log_{10}(P)$ for the BH corrected P-value of each intersection. F,G) Cellular

compartments (F) and molecular functions (G)overrepresented amongst DEGs detected between

the hippocampal subregions from trained and untrained samples for all regions combined and

separated CA1, CA3 and DG samples. Dot color reflects the statistical significance (-

$\log_{10}(FDR)$) of the biological process enrichment. Dot size reflects the number of detected DEGs

mapped to the genes involved in a given biological process.

Supplemental Figure 2. Cellular components and molecular functions enriched among DEGs

detected using spatial transcriptomics

A) Upset plot of overlaps in regional DEGs demonstrates greatest similarity between CA1 and

CA3 subregions. DEGs from each regional analysis were stratified by direction of fold change.

In the upset plot, each bar represents the number of genes in an intersection of lists of DEGs. The

intersecting lists is represented by the dots underneath the bar. The hashed line on each bar

represents the expected intersection of two identically sized random lists. Overlaps were tested

with Fisher's exact test using the list of all expressed genes as a background. The color of the

bar represents the $-\log_{10}(P)$ for the BH corrected P-value of each intersection. B,C) Molecular

functions (B) and cellular compartments (C) overrepresented amongst DEGs detected between

the hippocampal samples from trained and untrained samples for all hippocampal spots and spots

in the cell layers of CA1, CA3 and DG subregions. Dot color reflects the statistical significance

$(-\log_{10}(FDR))$ of the biological process enrichment. Dot size reflects the number of detected

DEGs mapped to the genes involved in a given biological process.

Cohort	IEG	All Spots	Cell Layers	CA1	CA3	DG
Trained	Arc	195/542	56/146	34/49	20/43	2/54
	Egr1	380/542	130/146	48/49	41/43	41/54
	c-Jun	308/542	106/146	22/49	32/43	52/54
Untrained	Arc	169/568	58/155	34/51	19/42	5/62
	Egr1	256/568	113/155	51/51	33/42	29/62
	c-Jun	175/568	83/155	12/51	18/42	53/62

Supplemental Figure 3. The expression of the IEGs *Egr1* and *c-Jun* are most and least correlated

with the expression of *Arc*, respectively

A, B) Heatmaps reflecting the pairwise Pearson correlation coefficients between IEGs detected

in the Trained (A) and Untrained (B) sample. Pearson correlation between *Arc* and *Egr1* or *c-Jun*

are marked by the caret on the x-axis colored to reflect the value of the correlation coefficient.

The correlation coefficients for *Arc* and *Egr1* are $r(21) = .598$, $p = 2.24 \times 10^{-56}$ in the untrained

sample and $r(21) = .764$, $p = 5.45 \times 10^{-105}$ in the trained sample. The correlation coefficients for

Arc and *c-Jun* are $r(21) = -.057$, $p = 0.17$ in the untrained sample and $r(21) = -.105$, $p = 0.014$ in

the trained sample. C,E) Molecular functions (C) and cellular compartments (E) overrepresented

amongst DEGs detected across trained and untrained samples for spots expressionr the IEGs *Arc*,

Egr1, and *c-Jun*. Dot color reflects the statistical significance ($-\log_{10}(FDR)$) of the biological

process enrichment. Dot size reflects the number of detected DEGs mapping to the gene

belonging to a given biological process D,F) Molecular functions (D) and cellular compartments

(F) overrepresented amongst DEGs detected between IEG positive and negative spots within

either the trained or untrained sample. Dot color reflects the statistical significance ($-\log_{10}(FDR)$)

of the biological process enrichment. Dot size reflects the number of detected DEGs mapping to

the gene belonging to a given biological process G) Table total number of IEG expressing spots

relative to the total number of hippocampal spots for each area.

Supplemental Figure 4. Relative expression of select DEGs through RT-qPCR

A-C)Average normalized relative expression ($2^{-\Delta\Delta Ct}$) of genes in trained samples (red) as

compared to untrained samples (blue) in dentate gyrus (A), CA3 (B) and CA1 (C) subregions. n=

6 trained and n=4 untrained animals. Student t-test was performed between ΔCt values from

trained and untrained samples for each gene.

REFERENCES

1. Kitamura, T., et al., *Engrams and circuits crucial for systems consolidation of a memory.* Science, 2017. **356**(6333): p. 73-78.
2. Josselyn, S.A. and S. Tonegawa, *Memory engrams: Recalling the past and imagining the future.* Science, 2020. **367**(6473).
3. Takeuchi, T., A.J. Duszkiewicz, and R.G. Morris, *The synaptic plasticity and memory hypothesis: encoding, storage and persistence.* Philos Trans R Soc Lond B Biol Sci, 2014. **369**(1633): p. 20130288.
4. Malenka, R.C. and M.F. Bear, *LTP and LTD: an embarrassment of riches.* Neuron, 2004. **44**(1): p. 5-21.
5. Mayford, M., S.A. Siegelbaum, and E.R. Kandel, *Synapses and memory storage.* Cold Spring Harb Perspect Biol, 2012. **4**(6).
6. Marco, A., et al., *Mapping the epigenomic and transcriptomic interplay during memory formation and recall in the hippocampal engram ensemble.* Nat Neurosci, 2020. **23**(12): p. 1606-1617.
7. Rao-Ruiz, P., et al., *Engram-specific transcriptome profiling of contextual memory consolidation.* Nat Commun, 2019. **10**(1): p. 2232.
8. Reshetnikov, V.V., et al., *Genes associated with cognitive performance in the Morris water maze: an RNA-seq study.* Sci Rep, 2020. **10**(1): p. 22078.
9. Poo, M.M., et al., *What is memory? The present state of the engram.* BMC Biol, 2016. **14**: p. 40.
10. Mayford, M., *The search for a hippocampal engram.* Philos Trans R Soc Lond B Biol Sci, 2014. **369**(1633): p. 20130161.
11. Middleton, S.J. and T.J. McHugh, *CA2: A Highly Connected Intrahippocampal Relay.* Annu Rev Neurosci, 2020. **43**: p. 55-72.
12. Chevaleyre, V. and S.A. Siegelbaum, *Strong CA2 pyramidal neuron synapses define a powerful disynaptic cortico-hippocampal loop.* Neuron, 2010. **66**(4): p. 560-572.
13. Senzai, Y., *Function of local circuits in the hippocampal dentate gyrus-CA3 system.* Neurosci Res, 2019. **140**: p. 43-52.
14. Basu, J. and S.A. Siegelbaum, *The Corticohippocampal Circuit, Synaptic Plasticity, and Memory.* Cold Spring Harb Perspect Biol, 2015. **7**(11).
15. Amaral, D.G. and M.P. Witter, *The three-dimensional organization of the hippocampal formation: a review of anatomical data.* Neuroscience, 1989. **31**(3): p. 571-91.
16. Sloviter, R.S. and T. Lomo, *Updating the lamellar hypothesis of hippocampal organization.* Front Neural Circuits, 2012. **6**: p. 102.
17. van Strien, N.M., N.L. Cappaert, and M.P. Witter, *The anatomy of memory: an interactive overview of the parahippocampal-hippocampal network.* Nat Rev Neurosci, 2009. **10**(4): p. 272-82.
18. Kesner, R.P., I. Lee, and P. Gilbert, *A behavioral assessment of hippocampal function based on a subregional analysis.* Rev Neurosci, 2004. **15**(5): p. 333-51.
19. Lisman, J.E., *Relating hippocampal circuitry to function: recall of memory sequences by reciprocal dentate-CA3 interactions.* Neuron, 1999. **22**(2): p. 233-42.
20. Ryan, T.J., et al., *Memory. Engram cells retain memory under retrograde amnesia.* Science, 2015. **348**(6238): p. 1007-13.
21. Hunsaker, M.R. and R.P. Kesner, *The operation of pattern separation and pattern completion processes associated with different attributes or domains of memory.* Neurosci Biobehav Rev, 2013. **37**(1): p. 36-58.
22. Molitor, R.J., et al., *Memory Reactivation during Learning Simultaneously Promotes Dentate Gyrus/CA$_{2,3}$ Pattern Differentiation and CA$_1$ Memory Integration.* The Journal of Neuroscience, 2021. **41**(4): p. 726-738.
23. Shpokayte, M., et al., *Hippocampal cells segregate positive and negative engrams.* Commun Biol, 2022. **5**(1): p. 1009.
24. Sullivan, K.E., R.M. Kendrick, and M.S. Cembrowski, *Elucidating memory in the brain via single-cell transcriptomics.* J Neurochem, 2021. **157**(4): p. 982-992.

25.   Shah, S., et al., *In Situ Transcription Profiling of Single Cells Reveals Spatial Organization of Cells*
in the Mouse Hippocampus. Neuron, 2016. **92**(2): p. 342-357.

26.   Cho, J.H., B.S. Huang, and J.M. Gray, *RNA sequencing from neural ensembles activated during*
fear conditioning in the mouse temporal association cortex. Sci Rep, 2016. **6**: p. 31753.

27.   Tonegawa, S., M.D. Morrissey, and T. Kitamura, *The role of engram cells in the systems*
consolidation of memory. Nat Rev Neurosci, 2018. **19**(8): p. 485-498.

28.   Sanes, J.R. and J.W. Lichtman, *Can molecules explain long-term potentiation?* Nat Neurosci, 1999.
2(7): p. 597-604.

29.   Alberini, C.M. and E.R. Kandel, *The regulation of transcription in memory consolidation.* Cold
Spring Harb Perspect Biol, 2014. **7**(1): p. a021741.

30.   Asok, A., et al., *Molecular Mechanisms of the Memory Trace.* Trends Neurosci, 2019. **42**(1): p. 14-
22.

31.   Magee, J.C. and C. Grienberger, *Synaptic Plasticity Forms and Functions.* Annual Review of
Neuroscience, 2020. **43**(1): p. 95-117.

32.   Guan, J.S., et al., *How Does the Sparse Memory "Engram" Neurons Encode the Memory of a*
Spatial-Temporal Event? Front Neural Circuits, 2016. **10**: p. 61.

33.   Chen, B.K., et al., *Artificially Enhancing and Suppressing Hippocampus-Mediated Memories.* Curr
Biol, 2019. **29**(11): p. 1885-1894 e4.

34.   Sauvage, M., T. Kitsukawa, and E. Atucha, *Single-cell memory trace imaging with immediate-*
early genes. J Neurosci Methods, 2019. **326**: p. 108368.

35.   Meenakshi, P., S. Kumar, and J. Balaji, *In vivo imaging of immediate early gene expression*
dynamics segregates neuronal ensemble of memories of dual events. Mol Brain, 2021. **14**(1): p.
102.

36.   Erwin, S.R., et al., *A Sparse, Spatially Biased Subtype of Mature Granule Cell Dominates*
Recruitment in Hippocampal-Associated Behaviors. Cell Rep, 2020. **31**(4): p. 107551.

37.   Guzowski, J.F., et al., *Environment-specific expression of the immediate-early gene Arc in*
hippocampal neuronal ensembles. Nat Neurosci, 1999. **2**(12): p. 1120-4.

38.   Ramirez-Amaya, V., et al., *Spatial exploration-induced Arc mRNA and protein expression:*
evidence for selective, network-specific reactivation. J Neurosci, 2005. **25**(7): p. 1761-8.

39.   Lee, B.H., et al., *Real-time visualization of mRNA synthesis during memory formation in live mice.*
Proc Natl Acad Sci U S A, 2022. **119**(27): p. e2117076119.

40.   Cimadevilla, J.M., et al., *Passive and active place avoidance as a tool of spatial memory research*
in rats. J Neurosci Methods, 2000. **102**(2): p. 155-64.

41.   Stuchlík, A., et al., *Place Avoidance Tasks as Tools in the Behavioral Neuroscience of Learning*
and Memory. Physiological Research, 2013: p. S1-S19.

42.   Wachi, T., B. Cornell, and K. Toyo-Oka, *Complete ablation of the 14-3-3epsilon protein results in*
multiple defects in neuropsychiatric behaviors. Behav Brain Res, 2017. **319**: p. 31-36.

43.   Annamneedi, A., et al., *The Presynaptic Scaffold Protein Bassoon in Forebrain Excitatory Neurons*
Mediates Hippocampal Circuit Maturation: Potential Involvement of TrkB Signalling. Int J Mol
Sci, 2021. **22**(15).

44.   Leone, L., et al., *Altered Nup153 Expression Impairs the Function of Cultured Hippocampal*
Neural Stem Cells Isolated from a Mouse Model of Alzheimer's Disease. Mol Neurobiol, 2019.
56(8): p. 5934-5949.

45.   Vigil, F.A., et al., *Prevention of long-term memory loss after retrieval by an endogenous CaMKII*
inhibitor. Sci Rep, 2017. **7**(1): p. 4040.

46.   Urrutia-Ruiz, C., et al., *Deletion of the Autism-Associated Protein SHANK3 Abolishes Structural*
Synaptic Plasticity after Brain Trauma. Int J Mol Sci, 2022. **23**(11).

47.   Hung, A.Y., et al., *Smaller dendritic spines, weaker synaptic transmission, but enhanced spatial*
learning in mice lacking Shank1. J Neurosci, 2008. **28**(7): p. 1697-708.

48.   Hou, Q., et al., *SNAP-25 in hippocampal CA1 region is involved in memory consolidation.* Eur J
Neurosci, 2004. **20**(6): p. 1593-603.

49. Gomis-Gonzalez, M., et al., *Protein Kinase C-Gamma Knockout Mice Show Impaired*
Hippocampal Short-Term Memory While Preserved Long-Term Memory. Mol Neurobiol, 2021.
58(2): p. 617-630.
50. Iqbal, J., *Transthyretin—A Key Gene Involved in Regulating Learning and Memory in Brain, and*
Providing Neuroprotection in Alzheimer Disease via Neuronal Synthesis of Transthyretin Protein.
Scientific Research Publishing, 2018. **8**(2).
51. Pratt, K.J.B., et al., *Loss of neuronal Tet2 enhances hippocampal-dependent cognitive function.*
Cell Rep, 2022. **41**(6): p. 111612.
52. Katafuchi, T., et al., *Impairment of spatial learning and hippocampal synaptic potentiation in c-kit*
mutant rats. Learn Mem, 2000. **7**(6): p. 383-92.
53. Ferreira, E., D.M. Shaw, and S. Oddo, *Identification of learning-induced changes in protein*
networks in the hippocampi of a mouse model of Alzheimer's disease. Transl Psychiatry, 2016. **6**(7):
p. e849.
54. Barbieri, R., et al., *Synapsin I and Synapsin II regulate neurogenesis in the dentate gyrus of adult*
mice. Oncotarget, 2018. **9**(27): p. 18760-18774.
55. The Gene Ontology, C., *The Gene Ontology Resource: 20 years and still GOing strong.* Nucleic
Acids Res, 2019. **47**(D1): p. D330-D338.
56. Yao, Z., et al., *A taxonomy of transcriptomic cell types across the isocortex and hippocampal*
formation. Cell, 2021. **184**(12): p. 3222-3241 e26.
57. Schober, D.A., et al., *Transmembrane AMPA receptor regulatory proteins and cornichon-2*
allosterically regulate AMPA receptor antagonists and potentiators. J Biol Chem, 2011. **286**(15):
p. 13134-42.
58. Lugara, E., et al., *Synapsin I Controls Synaptic Maturation of Long-Range Projections in the*
Lateral Amygdala in a Targeted Selective Fashion. Front Cell Neurosci, 2019. **13**: p. 220.
59. Nicole, O. and E. Pacary, *CaMKIIbeta in Neuronal Development and Plasticity: An Emerging*
Candidate in Brain Diseases. Int J Mol Sci, 2020. **21**(19).
60. Carrillo-Reid, L., *Neuronal ensembles in memory processes.* Semin Cell Dev Biol, 2022. **125**: p.
136-143.
61. Gouty-Colomer, L.A., et al., *Arc expression identifies the lateral amygdala fear memory trace.* Mol
Psychiatry, 2016. **21**(3): p. 364-75.
62. Hsieh, C., et al., *Persistent increases of PKMzeta in memory-activated neurons trace LTP*
maintenance during spatial long-term memory storage. Eur J Neurosci, 2021. **54**(8): p. 6795-6814.
63. Chia, C. and T. Otto, *Hippocampal Arc (Arg3.1) expression is induced by memory recall and*
required for memory reconsolidation in trace fear conditioning. Neurobiol Learn Mem, 2013. **106**:
p. 48-55.
64. Zhang, T., et al., *Expression of BC1 Impairs Spatial Learning and Memory in Alzheimer's Disease*
Via APP Translation. Mol Neurobiol, 2018. **55**(7): p. 6007-6020.
65. Liu, Y., et al., *Hippocampal Activation of Rac1 Regulates the Forgetting of Object Recognition*
Memory. Curr Biol, 2016. **26**(17): p. 2351-7.
66. Zhang, L., et al., *Dickkopf 3 (Dkk3) Improves Amyloid-beta Pathology, Cognitive Dysfunction, and*
Cerebral Glucose Metabolism in a Transgenic Mouse Model of Alzheimer's Disease. J Alzheimers
Dis, 2017. **60**(2): p. 733-746.
67. Hadjiivanova, C., S. Belcheva, and I. Belcheva, *Cholecystokinin and learning and memory*
processes. Acta Physiol Pharmacol Bulg, 2003. **27**(2-3): p. 83-8.
68. Tutukova, S., V. Tarabykin, and L.R. Hernandez-Miranda, *The Role of Neurod Genes in Brain*
Development, Function, and Disease. Front Mol Neurosci, 2021. **14**: p. 662774.
69. Sun, X., et al., *Functionally Distinct Neuronal Ensembles within the Memory Engram.* Cell, 2020.
181(2): p. 410-423 e17.
70. Nambu, M.F., et al., *What does engram encode?: Heterogeneous memory engrams for different*
aspects of experience. Curr Opin Neurobiol, 2022. **75**: p. 102568.

71.   Okuno, H., *Regulation and function of immediate-early genes in the brain: beyond neuronal*
activity markers. Neurosci Res, 2011. **69**(3): p. 175-86.
72.   Gallo, F.T., et al., *Immediate Early Genes, Memory and Psychiatric Disorders: Focus on c-Fos,*
Egr1 and Arc. Front Behav Neurosci, 2018. **12**: p. 79.
73.   Vanrobaeys, Y., et al., *Mapping the spatial transcriptomic signature of the hippocampus during*
memory consolidation. Nat Commun, 2023. **14**(1): p. 6100.
74.   Rolls, E.T., *Memory systems in the brain.* Annu Rev Psychol, 2000. **51**: p. 599-630.
75.   Stanfield, B.B. and W.M. Cowan, *The morphology of the hippocampus and dentate gyrus in normal*
and reeler mice. J Comp Neurol, 1979. **185**(3): p. 393-422.
76.   Khalaf-Nazzal, R. and F. Francis, *Hippocampal development - old and new findings.* Neuroscience,
2013. **248**: p. 225-42.
77.   Cembrowski, M.S. and N. Spruston, *Heterogeneity within classical cell types is the rule: lessons*
from hippocampal pyramidal neurons. Nat Rev Neurosci, 2019. **20**(4): p. 193-204.
78.   Montgomery, S.M. and G. Buzsaki, *Gamma oscillations dynamically couple hippocampal CA3 and*
CA1 regions during memory task performance. Proc Natl Acad Sci U S A, 2007. **104**(36): p. 14495-
500.
79.   Carr, M.F., M.P. Karlsson, and L.M. Frank, *Transient slow gamma synchrony underlies*
hippocampal memory replay. Neuron, 2012. **75**(4): p. 700-13.
80.   Alam, M.J., et al., *Adult Neurogenesis Conserves Hippocampal Memory Capacity.* J Neurosci,
2018. **38**(31): p. 6854-6863.
81.   Kitamura, T., et al., *Adult neurogenesis modulates the hippocampus-dependent period of*
associative fear memory. Cell, 2009. **139**(4): p. 814-27.
82.   Kitamura, T. and K. Inokuchi, *Role of adult neurogenesis in hippocampal-cortical memory*
consolidation. Mol Brain, 2014. **7**: p. 13.
83.   Deng, W., J.B. Aimone, and F.H. Gage, *New neurons and new memories: how does adult*
hippocampal neurogenesis affect learning and memory? Nat Rev Neurosci, 2010. **11**(5): p. 339-
50.
84.   Aksoy-Aksel, A. and D. Manahan-Vaughan, *The temporoammonic input to the hippocampal CA1*
region displays distinctly different synaptic plasticity compared to the Schaffer collateral input in
vivo: significance for synaptic information processing. Front Synaptic Neurosci, 2013. **5**: p. 5.
85.   Pavlowsky, A. and J.M. Alarcon, *Interaction between long-term potentiation and depression in*
CA1 synapses: temporal constrains, functional compartmentalization and protein synthesis. PLoS
One, 2012. **7**(1): p. e29865.
86.   Mendez-Lopez, M., et al., *Spatial short-term memory in rats: effects of learning trials on metabolic*
activity of limbic structures. Neurosci Lett, 2010. **483**(1): p. 32-5.
87.   Christie, S.T. and P. Schrater, *Cognitive cost as dynamic allocation of energetic resources.* Front
Neurosci, 2015. **9**: p. 289.
88.   Terranova, J.I., et al., *Hippocampal-amygdala memory circuits govern experience-dependent*
observational fear. Neuron, 2022. **110**(8): p. 1416-1431 e13.
89.   Roy, D.S., et al., *Distinct Neural Circuits for the Formation and Retrieval of Episodic Memories.*
Cell, 2017. **170**(5): p. 1000-1012 e19.
90.   Marks, W.D., et al., *Neuronal Ensembles Organize Activity to Generate Contextual Memory.* Front
Behav Neurosci, 2022. **16**: p. 805132.
91.   Li, X. and C.Y. Wang, *From bulk, single-cell to spatial RNA sequencing.* Int J Oral Sci, 2021.
13(1): p. 36.
92.   Stahl, P.L., et al., *Visualization and analysis of gene expression in tissue sections by spatial*
transcriptomics. Science, 2016. **353**(6294): p. 78-82.
93.   Asp, M., J. Bergenstrahle, and J. Lundeberg, *Spatially Resolved Transcriptomes-Next Generation*
Tools for Tissue Exploration. Bioessays, 2020. **42**(10): p. e1900221.
94.   Yap, E.L. and M.E. Greenberg, *Activity-Regulated Transcription: Bridging the Gap between*
Neural Activity and Behavior. Neuron, 2018. **100**(2): p. 330-348.

95.   Sheng, M. and M.E. Greenberg, *The regulation and function of c-fos and other immediate early*
genes in the nervous system. Neuron, 1990. **4**(4): p. 477-85.
96.   Jeong, Y., et al., *Synaptic plasticity-dependent competition rule influences memory formation.* Nat
Commun, 2021. **12**(1): p. 3915.
97.   Magistretti, P.J. and I. Allaman, *A cellular perspective on brain energy metabolism and functional*
imaging. Neuron, 2015. **86**(4): p. 883-901.
98.   Wesierska, M., C. Dockery, and A.A. Fenton, *Beyond memory, navigation, and inhibition:*
behavioral evidence for hippocampus-dependent cognitive coordination in the rat. J Neurosci,
2005. **25**(9): p. 2413-9.
99.   Lacagnina, A.F., et al., *Distinct hippocampal engrams control extinction and relapse of fear*
memory. Nat Neurosci, 2019. **22**(5): p. 753-761.
100.   Sweis, B.M., et al., *Dynamic and heterogeneous neural ensembles contribute to a memory engram.*
Curr Opin Neurobiol, 2021. **67**: p. 199-206.
101.   Tonegawa, S., et al., *Memory engram storage and retrieval.* Curr Opin Neurobiol, 2015. **35**: p. 101-
9.
102.   Terranova, J.I., et al., *Systems consolidation induces multiple memory engrams for a flexible recall*
strategy in observational fear memory in male mice. Nat Commun, 2023. **14**(1): p. 3976.
103.   Sun, X. and Y. Lin, *Npas4: Linking Neuronal Activity to Memory.* Trends Neurosci, 2016. **39**(4):
p. 264-275.
104.   Sheng, H.Z., R.D. Fields, and P.G. Nelson, *Specific regulation of immediate early genes by*
patterned neuronal activity. J Neurosci Res, 1993. **35**(5): p. 459-67.
105.   Guzowski, J.F., et al., *Experience-dependent gene expression in the rat hippocampus after spatial*
learning: a comparison of the immediate-early genes Arc, c-fos, and zif268. J Neurosci, 2001.
21(14): p. 5089-98.
106.   Lyons, M.R. and A.E. West, *Mechanisms of specificity in neuronal activity-regulated gene*
transcription. Prog Neurobiol, 2011. **94**(3): p. 259-95.
107.   Madabhushi, R. and T.K. Kim, *Emerging themes in neuronal activity-dependent gene expression.*
Mol Cell Neurosci, 2018. **87**: p. 27-34.
108.   Tyssowski, K.M., et al., *Different Neuronal Activity Patterns Induce Different Gene Expression*
Programs. Neuron, 2018. **98**(3): p. 530-546 e11.
109.   Flavell, S.W. and M.E. Greenberg, *Signaling mechanisms linking neuronal activity to gene*
expression and plasticity of the nervous system. Annu Rev Neurosci, 2008. **31**: p. 563-90.
110.   Cheval, H., et al., *Distinctive features of Egr transcription factor regulation and DNA binding*
activity in CA1 of the hippocampus in synaptic plasticity and consolidation and reconsolidation of
fear memory. Hippocampus, 2012. **22**(3): p. 631-42.
111.   Maddox, S.A., M.S. Monsey, and G.E. Schafe, *Early growth response gene 1 (Egr-1) is required*
for new and reactivated fear memories in the lateral amygdala. Learn Mem, 2011. **18**(1): p. 24-38.
112.   Sherrin, T., et al., *Hippocampal c-Jun-N-terminal kinases serve as negative regulators of*
associative learning. J Neurosci, 2010. **30**(40): p. 13348-61.
113.   Denny, C.A., et al., *Hippocampal memory traces are differentially modulated by experience, time,*
and adult neurogenesis. Neuron, 2014. **83**(1): p. 189-201.
114.   10x Genomics, *Visium Spatial Protocols – Tissue Preparation Guide.* 2023.
115.   Paxinos, G. and K.B.J. Franklin, *Paxinos and Franklin's the Mouse Brain in Stereotaxic*
Coordinates. 2019: Elsevier Science.
116.   Genomics, x., *Visium Spatial Gene Expression Reagent Kits User Guide.* 2023.
117.   Genomics, x., *Visium Spatial Gene Expression Reagent Kits - Tissue Optimization.* 2019.
118.   Genomics, x., *Visium Spatial Gene Expression Imaging Guidelines.* 2021.
119.   Consortium, G.R., *Genome Reference Consortium Mouse Build 38.* 2012.
120.   Dobin, A., et al., *STAR: ultrafast universal RNA-seq aligner.* Bioinformatics, 2013. **29**(1): p. 15-
21.

121. Wu, T., et al., *clusterProfiler 4.0: A universal enrichment tool for interpreting omics data.*
Innovation (Camb), 2021. **2**(3): p. 100141.
122. Cai, H., et al., *VennPlex–A Novel Venn Diagram Program for Comparing and Visualizing Datasets*
with Differentially Regulated Datapoints. PLoS ONE, 2013. **8**(1): p. e53388.
123. Wang, M., Y. Zhao, and B. Zhang, *Efficient Test and Visualization of Multi-Set Intersections.* Sci
Rep, 2015. **5**: p. 16923.
124. Butler, A., et al., *Integrating single-cell transcriptomic data across different conditions,*
technologies, and species. Nat Biotechnol, 2018. **36**(5): p. 411-420.
125. Ponchel, F., et al., *Real-time PCR based on SYBR-Green I fluorescence: an alternative to the*
TaqMan assay for a relative quantification of gene rearrangements, gene amplifications and micro
gene deletions. BMC Biotechnol, 2003. **3**: p. 18.
1036