Hypoxia impacts small intestinal organoid stemness and differentiation

Xi Lan[1], Ping Qiu[2], Chunfeng Mou[1]

[1]Department of Nuclear Medicine, Children's Hospital of Chongqing Medical University, Chongqing, 400014 China.
[2]Stem Cells and Regenerative Medicine Research Center, Sichuan Neo-life Stem Cell Biotech Inc., 15 Jinquan Road, Chengdu, 610036, China.

Summary

Comprised of several cell types functioning differently, the small intestinal epithelial cells perform their functions and synergistically maintain homeostasis in the first part of the mammalian intestine. Due to the uneven distribution of the vessel, the oxygen level exhibits a gradient decreasing pattern in the normal intestine and becomes aberrant in some intestinal diseases. In this work, we find certain levels of hypoxia simulated by cobaltous chloride ($CoCl_2$) caused an increase in the secretive cell types and a decrease in the absorptive cell types *in vitro* cultured mouse small intestinal organoids. Importantly, the intestinal stem cell amount is impacted which leads to attenuated epithelial regeneration. Our study highlights the cell-type-specific alterations under the hypoxia insult, which gives possible therapeutic hints for hypoxia-relevant gastrointestinal diseases.

Keywords

Hypoxia, intestinal stem cell, differentiation, organoid

Introduction

The human small intestinal epithelium is the fastest self-renewing tissue, which renews every 3 to 5 days, with the structure of bending cells forming the crypts and villi (1). Intestinal stem cells (ISCs) reside at the crypt base and differentiate into the transit-amplifying (TA) compartment which divides 4-5 times within 12 hours (2). After that, the TA cells move upwards and differentiate into multiple cell types including enterocytes, goblet cells and enteroendocrine cells. During normal homeostasis or some intestinal injuries, ISCs can undergo self-renewal and proliferate to replenish cells that are lost to shedding (3, 4). The self-renewal and lineage differentiation into mature enterocytes, goblet cells and enteroendocrine cells are under the control of Wnt/β-catenin, Notch and bone morphogenetic protein (BMP) pathway signalings (5-7). Among them, Wnt/β-catenin signaling is of critical significance, and its downstream regulated gene Leucine-rich repeat-containing G-protein coupled receptor 5 (Lgr5) is identified as the marker for proliferative ISCs (8, 9).

Mutations that activate the Wnt/β-catenin pathway can result in human colorectal cancer (10, 11). Studies suggested Wnt as an essential growth factor for the maintenance of intestinal proliferation and homeostasis (12, 13), and ectopic Wnt signaling has been

implicated in the initiation and progression of colorectal cancers (14, 15).

In a healthy human intestine, due to the counter-current blood and the existence of a large
number of gut bacteria, the oxygen pressure (pO2) level is lower than the normal oxygen
pressure (160 mmHg), therefore, it's described as "physiologic hypoxia" (16). The local pO2
level decreases along the crypt-villus axis and the oxygen gradient becomes the lowest at
the villus tip (17). This unique gradient physiologic oxygen pressure results in the
stabilization of the transcription factor hypoxia-inducible factor (HIF) and the activation of
hypoxia-relative signaling, which plays multiple roles in metabolism, mitochondrial
autophagy and inflammation (18). However, in pathological conditions, profound hypoxic or
anoxic are usually found in mucosal lesions (19). Hypoxia is also a feature of inflammatory
bowel disease (IBD), and it affects inflammation in some intestinal diseases (16, 20). Studies
have demonstrated the protective role of HIF on some intestinal diseases such as IBD and
necrotizing enterocolitis (NEC) upon *in vivo* hypoxia injury (21-23), and the effect of hypoxia
on intestinal cell response in intestinal cancer cells or upon radiotherapy and chemotherapy
in vitro (24-26). In both scenarios, the intestinal stem cells (ISCs) have been shown to react
by promoting intestinal recovery from injuries (27, 28). However, the mechanism of how
ectopic hypoxia affects ISC function and how it impacts intestinal cell lineage differentiation
is less clarified. Mazumdar et al. mentioned that HIF-1α regulates Wnt/β-catenin signaling in
hypoxic embryonic stem (ES) cells and neural stem cells (29). Rohwer et al. demonstrated
the role of HIF-1α in intestinal epithelial cells (IECs) during intestinal tumorgenesis,
controlling Wnt/β-catenin activity and regulating metabolic reprogramming and
inflammation (30). However, it remains to be elucidated how hypoxia impacts different cell
types in intestine with or without affecting Wnt signaling.

Intestinal organoids isolated from adult stem cells have been established as an ingenious
craft to study the human intestine (8). It's a 3D model of intestinal tissue embedded in 3D
matrices such as Matrigel and can be passaged and cultured for a long time. The organoids
display the characteristic villi-crypt organization by heterogeneous epithelial cell monolayer,
allowing it to recapitulate many features of normal gut epithelium with IECs comprising all
types of epithelial cells (31). In addition, the organoid system is applied to estimate IEC
functions *in vivo* such as barrier function and stem cell function (32, 33). Compared with the
in vivo mouse model and 2D cell lines (26, 34) to study the influence of hypoxia on intestinal
functions or tumorigenesis, intestinal organoids can be easily established from different
staging of endoscopic biopsies in patients, making it a better model that can closely
recapitulate the hypoxic intestine and a faster model to come up with therapeutic regimens.

Hypoxia exists in intestinal diseases such as IBD and NEC, wherein the mechanism by which
hypoxia impacts ISC stemness and differentiation remains poorly understood. In the present
study, we examined obvious changes of different cell lineages upon hypoxia treatment to
intestinal organoids, as well as the upstream transcription factors regulating the ISC
stemness. Our study further confirms organoids as the model to study intestinal diseases
that are concurrent with hypoxia. This cobaltous chloride (CoCl$_2$)-caused hypoxia-treated
organoid model makes it a good method to study HIF-1α regulated signaling in intestine,
helping the translation of basic science studies into IBD treatment.

Result

1. Hypoxia treatment affects intestinal organoid stem cell renewal function

It's known that the ability of IECs to form organoids and organoid regeneration after being
passaged depends mainly on ISC function (4, 8). To investigate the effect of hypoxia
exposure on intestine organoid stem cell regeneration capacity, we passaged the ISC-
containing organoid as an assay of intestinal regenerative potential (8). Increasing dosages
of $CoCl_2$ (0, 10, 50, 100, 200 μM) were used to induce cellular hypoxia (35, 36), and did not
impair the budding and morphology of organoids (Fig. 1A). The protein level of HIF-1α was
upregulated 24 hours after hypoxia exposure (Fig. 1B). The organoid regeneration, which
indicates stem cell function and generated from single intestinal organoid cells engendered
fewer progenies in a gradient decreasing manner, as showed in the clonogenicity assay (Fig
1C).
We further explored whether the reduction of the organoid regeneration function was
preceded by the reduction of stem cell trait. Indeed, the mRNA levels of Lgr5 which mark
the intestinal stem cells, were also down-regulated (Fig. 2A). Therefore, these data suggest
that hypoxia exposure in intestine organoids impairs ISC regeneration by reducing ISC
numbers and organoid-forming capacity.
105
2. Wnt signaling was down-regulated upon hypoxia treatment in intestinal organoids

Wnt signals play a crucial role in the regulation of intestinal stem cell proliferation (11, 37,
38). As the key stem cell regulator in intestine stem cells, Lgr5 is controlled by Wnt/β-
catenin signaling (4, 39) and exhibits a reduced mRNA level after 200 μM $CoCl_2$ treated
hypoxia treatment (Fig. 2A). To further clarify the effect of hypoxia on Wnt signaling in
intestinal organoid, multiple Wnt target genes were tested in hypoxia-exposed intestine
organoids. As the hypoxic condition gets severe, the protein level of β-catenin was down-
regulated in a dosage-dependent manner (Fig 2B). Also, the β-catenin target Cyclin-D1,
which is the gene involved in cell survival (40), exhibited significant protein level reduction.
Expression of another Wnt target, the receptor tyrosine kinase EphB2, which reduces from
the crypt base to the differentiated cells (41) and is highly enriched in ISCs (42), also shows a
reduced protein level after 24 hrs hypoxia treatment. Consistent with this, protein
expression of another Wnt/β-catenin target gene Axin2 (43) was also decreased. These
demonstrate that hypoxia treatment dampens the activity of Wnt/β-catenin signaling in
intestinal organoids.
121
3. Hypoxia treatment affects intestine differentiation and increases secretory cell types

Pathological conditions such as inflammation insults are usually accompanied by cellular
hypoxia, further disrupting intestinal epithelial barrier function and metabolism (18). The
barrier function of intestinal epithelium is largely composed of enterocytes to maintain
intact epithelium construction, and goblet cell secreted mucus helps to form the healthy
mucosa layer, making the epithelium devoid of bacteria (44).

Then we deciphered whether hypoxia drives different intestine cell lineage cell
differentiation, and if it impairs certain cell types executing their functions. As shown in the
mRNA levels, hypoxia exposure led to the increase of Lysozyme (Lyz), Chromogranin A (ChrA)
and Mucin 2 (Muc2), which mark Paneth cell, enteroendocrine cell and goblet cell,
respectively (Fig. 3A). At the same time, the protein expressions showed by the
immunofluorescence (IFC) signal indicated a greater number of Lysozyme positive Paneth
cells, Chromogranin A positive enteroendocrine cells, and Mucin2 positive goblet cells in
hypoxia treated intestinal organoids (Fig 3B). Paneth cells act as the source of Wnt signaling

and secrete Wnt legends. However, our data suggest that this increase in Paneth cells per
buds, though, does not compensate for the lower level of Wnt signaling upon hypoxia
treatment.

During active inflammation in Crohn's disease (CD) and ulcerative colitis, pro-inflammatory
signals like cytokines promote the stabilization of HIF proteins (45). Hypoxia contributes to
inflammation, and Paneth cells are considered to be the origin of intestinal inflammation
(46). We then test whether hypoxia induces the inflammatory response in intestinal
organoids. Our data showed that hypoxia-induced stabilization of HIF-1α promotes the
transcription of proinflammatory cytokines tumor necrosis factor alpha (TNF-α) in intestinal
organoids (Fig. 4A, left). This increased TNF-α is likely secreted by the Paneth cells and we
wonder if TNF-α would boost this cell lineage. Single intestinal organoid cells treated with a
gradient concentration of TNF-α (5-50 ng/ml) for 48 hrs showed increased clonogenic ability,
as well as organoid swelling phenotype, especially in the higher TNF-α concentration
treatment group (Fig. 4B, C). Interestingly, we notice atypical cell morphology at the bottom
of the organoid-Matrigel culture plate in the TNF-α treated cells, which show elongated cell
morphology with protrusions (Fig. 4B, red arrows). This boosted growth of organoids
probably leads to an adaptation behavior under inflammation assault.

Paneth cells have active endoplasmic reticulum (ER) functions, and Paneth cell
differentiation depends on unfolded protein response (UPR) signaling (47). Unresolved ER
stress is common in the IBD epithelial cells (48). Importantly, unresolved ER stress and UPR
activation within Paneth cells can be treated as the nidus for intestinal inflammation (46).
We detected that mRNA levels of UPR proteins Atf4 and Chop are elevated upon hypoxia
exposure (Fig. 4A, right). While X-box binding protein 1 (Xbp1), Grp78 and Perk are not
changed significantly (data not shown).

Accordingly, hypoxia treatment increased the secretory lineage cell differentiation in
intestinal organoids, further causing the Paneth cell-relative ER stress response.
162
4. Hypoxia treatment reduces the absorptive cell type in intestinal organoids

Enterocytes constitute the most abundant cells and the only absorptive lineage cells in of
intestine. In addition to being an element of the epithelial lining, the differentiated mature
enterocytes at the upper villi in small intestine play a critical role in digestive, metabolic, and
barrier physical integrity maintenance functions (49). They are the cells most susceptible to
hypoxia (50). We found that mRNA levels of alkaline phosphatase (Alpi) and fatty acid
binding proteins (Fabp1/2) which hallmark enterocytes decreased after the gradient of
$CoCl_2$ treatment (Fig. 5A). IFC staining also showed a reduced amount of aldolase B (Aldob,
marker of enterocyte) positive cells (Fig. 5B), suggesting that enterocyte differentiation is
specifically impaired.

Enterocytes are involved in nutrient metabolism including glycolysis, fatty acid and lipid
metabolism (51, 52). The adipogenic gene Peroxisome proliferator-activated receptor γ
(PPARγ) has a beneficial metabolic and anti-inflammatory effect in the GI tract (53). We
found that PPARγ, although expressed at a low level in small intestine, was down-regulated
after hypoxia treatment (Fig. 5C).

Cells that are exposed to inflammatory cytokines manifest increased glycolysis, which is
reminiscent of those observed in hypoxic cells (54, 55). Glucose is also a relevant energy

source for enterocytes, and pyruvate kinases are responsible for catalyzing the final step in
glycolysis (56). We detected the protein level of glycolysis gene pyruvate kinases M (PKM)
was reduced after hypoxia exposure (Fig. 5C).

In addition to the general structure of intestinal organoids as the barrier, which we detected
no change after hypoxia treatment (Fig. 1A), we wondered whether the intrinsic elements of
the epithelium barrier were affected. We detected hypoxia caused a gradient down-
regulation of the epithelial barrier protein E-cadherin (Fig. 5C), which might indicate higher
permeability in the gastrointestinal epithelium

Therefore, hypoxia exposure decreased the absorptive enterocytes, with reduced
metabolic-related proteins and might dampened intestinal barrier function.
190
Discussion

Hypoxia occurs in intestinal diseases such as IBD and NEC (22, 23). It is well established that
hypoxia-induced intestinal injuries and low oxygen-relative physiological responses (23).
Studies using murine model of colitis indicated that HIF-1α plays a protective role in
intestinal inflammatory mucosal disease, most of which suggested a barrier-protective
function of HIF-1α (16, 20, 22). As different intestinal lineage cells executive specific
functions, however, the mechanism of hypoxia caused various cellular or metabolic
responses failed to be elucidated at the cell-type specific level in IEC. In this study, we
showed that hypoxia exposure by hydroxylase inhibitor $CoCl_2$ significantly changes the small
intestinal organoid cell lineage differentiation and reduces the ISC function by down-
regulating Wnt signaling.

It's known that hypoxia contributes to colorectal cancer cell (CRC) and activates Wnt
signaling (57, 58). However, in normal intestine epithelium under homeostasis, it's less
clarified whether hypoxia impacts ISC function and alters Wnt signaling. Consistent with Liu
and Zu et al. showing that Wnt signaling is reduced in the hypoxia-induced NEC and
ischemia-reperfusion (I/R) murine model (59, 60), our results further demonstrate that
hypoxia-caused impaired ISC function was accompanied by reduced Wnt signaling.
This intestinal organoid-based hypoxia exposure model provides an opportunity with the
cell type-specific model which correlates HIF-1 and the target genes in inflammation or
metabolism dysfunctions.

Notch signaling plays a crucial role in regulating cell fate decisions in intestinal homeostasis
(61, 62). Upon binding to their ligands, Jagged (Jag) and Delta-like (Dll) proteins, the Notch
receptor was cleaved and the intracellular domain translocates to the nucleus, where it
activates the transcription of the target genes, such as hairy and enhancer split-1 (Hes1),
regulating enterocyte differentiation (63). In reverse, Hes1 represses the expression of
atonal homolog 1 (Atoh1), which regulates secretory cell lineage differentiation.

Studies have implied that hypoxia increases the expression of Notch direct downstream
genes in neuronal and myogenic cells (64). Enterocyte differentiation is impaired as a
consequence of reduced Notch signaling pathway in patients with CD which is accompanied
by hypoxia-featured inflammation (65). While Notch1 signaling is upregulated in colorectal
hypoxic Caco-2 cells (66). It is still controversial regarding the role of Notch signaling in
regulating hypoxia-relevant intestinal diseases. Yeung et al. reported that hypoxia increases

clonogenicity and maintains the stem-like phenotype of colorectal cell line-derived cancer
stem cells, preventing differentiation of enterocytes and goblet cells by activating Notch1
(67). Of note, it remains less reported about the relationship between hypoxia and Notch
signaling in mediating stemness or barrier function in intestine under homeostasis. Although
we found the absorptive enterocytes decrease after hypoxia treatment, its regulatory Notch
signal transcription factors such as Hes1 and Notch intracellular domain (NICD) should be
tested in the future to answer if hypoxia affects the Notch signal directly.

Cell cultures have been used to study intestinal epithelial permeability after
repair/restitution (68, 69), while intestinal organoids provide us better *in vitro* model to
study the mechanism of injuries such as hypoxia on intestinal epithelial barrier function
which can best mimic the *in vivo* intestine. In intestinal cell lines or *in vivo* hypoxia-exposed
models (70, 71), HIF-1α was reported to influence intestinal epithelial permeability (19, 72).
Liu et al. suggested that Notch1 signaling plays a role in maintaining the intestinal barrier
function by regulating TJ proteins (66). More experiments are expected in the future such as
testing the expression of TJs (Zo-1 and occludin) and intestinal mucosal permeability
alterations, to prove the regulation of Notch to TJ proteins and its role in maintaining barrier
function.

On the other hand, we detected a higher number of goblet cells with increasing Muc2 level
after hypoxia exposure, consistent with D. Ortiz-Masiá et al.'s finding with increased Muc2
in chronic CD patients (65). Goblet cells secret the mucins that form the physical epithelial
barrier, and Paneth cells secrete antimicrobial and some signaling factors to regulate the ISC
stemness (73). The differentiation of IECs from the ISCs into the terminal differentiated cells
is under the regulation of master transcription factors (74). The progenitor cells committed
to secretory lineage have high expression of Atoh1. GFI1 and Spdef specify goblet and
Paneth cells, Neurogenin3 and NeuroD1 mark enteroendocrine cell differentiation, and Elf3
and Klf4 regulate goblet cell differentiation. More investigation is expected to delineate
whether hypoxia shifts early differentiation toward the secretory lineage via changing the
upstream secretory transcription factors.

Cells undergo stress and increase the production of certain proteins in their endoplasmic
reticulum (ER) when they experience stressful conditions like hypoxia (75), which is known
as the unfolded protein response (UPR). Heijmans et al. showed ER stress plays an
important role in regulating normal intestinal epithelial stem cell differentiation (47). We
detected the UPR targets Chop and Atf4 are up-regulated after hypoxia exposure. As XBP1 is
a regulator of the unfolded protein response, it's required for ER expansion (Shaffer et al.,
2004) and adaptation of tumor cells to hypoxic conditions (48). XBP1 deletion induces
inflammation in intestine (48). Although XBP1 was reported to drive the triple-negative
breast cancer (TNBC) tumorigenicity by controlling the HIF1α pathway (76), however, we
observed the level of Xbp1 did not change obviously in hypoxia-exposes normal intestinal
organoids.

As hypoxia treatment dampens intestinal organoid clonogenicity, while TNF-α treatment
boosts it, they likely target different cell types in the intestinal organoids. Compared with
the absorptive cell types (i.e., enterocytes) as the most vulnerable cells under hypoxia threat,
Paneth cells being the major warriors in front of TNF-α might explain the opposite growth
trend under these two different insults. Because TNF-α treatment 2 days post single

organoid seeding has the opposite barrier function protein alteration compared with those
single-cell formed organoids (data not shown), further enhancing the possibility that TNF-α
might boost Paneth cells in the first two days of organoid growth (wherein stem cells are
the majority cells) while mainly kill the enterocyte cell at later days. These two different
insults targeting different intestinal cell types help to explain that each cell type performs its
function under homeostasis and pathologic conditions. Besides, as we also notice atypical
cell protrusions in TNF-α treated cells, further experiments such as hypoxia and TNF-α co-
treatment can be done to elucidate whether this inflammation-fueled cell type-specific
outcome helps intestinal cancer cell metastasis, and whether hypoxia enhanced the
secretive cell number can further promote this phenomenon.

Method

Animals and crypt isolation organoid culture

Small intestines were removed from sacrificed 4-16-week-old C57BL/6 mice, flushed in cold
PBS and cut open. Mesentery was removed and the tissue was washed three times before
being cut into pieces. Then the tissues were digested with 2 mM EDTA and stayed on ice for
45 min. The digested intestines were transferred into a tube with cold PBS and shaken
around 3 times to detach the villi and remove residual EDTA. The suspension was filtered
through 70-μm meshes (BD Falcon) to enrich crypts. The crypt-containing solution was
centrifuged at 150 g for 3 min. The crypt pellet was then resuspended with DMEM and
embedded in Matrigel (Corning, #356231) for solidification for 15 min in a 37° C incubator.
Then culture medium was added above the crypt-Matrigel mix with DMEM/F12 (Gibco)
medium supplemented with EGF (50 ng/ml, Invitrogen), Noggin (50 ng/ml, R&D SYSTEMS),
R-spondin 500 ng/ml (R&D SYSTEMS), 1 × N-2 (Invitrogen), 1 × B-27 (Invitrogen), HEPES (10
mM), 1 x GlutaMax (Gibco), penicillin and streptomycin (100 U/ml), N-Acetyl Cysteine (1 μM,
Sigma). Y27632 (10 μM, Selleck) was included in the culture medium during the first 2 days
after being passaged. The medium was changed every other day.

Hypoxia treatment

294 Organoids were treated with gradient concentrations of $CoCl_2$ (Sigma, #V900059) by 0, 10,
295 50, 100, and 200 μM for 24 hours. DMSO (concentration less than 0.1%) was used as the
296 solvent and control reagent.

Clonogenicity assay

298 The colony-forming experiment was measured by seeding 200–300 single-cell organoids in
299 Matrigel and calculating organoid number after 2 days. The organoids were cultured for five
300 to nine days, then the crypt number was collected each day. Single organoid cells were
301 obtained by TrypLE (ThermoFisher) treatment for 10 min at 37℃, followed by gentle
302 pipetting if cell clusters remained.

RNA Isolation and Reverse transcription-quantitative polymerase chain reaction (RT-qPCR)

Total cellular RNA was extracted using EZ-10 RNA Mini-Preps Kits Handbook, (#BS88583,
Markham ON, Canada). Around 250 ng RNA was used to do reverse transcription by
RevertAid First Strand cDNA Synthesis Kit (K1622, Thermo, Waltham, MA, USA). Primers are
shown in Table 1. RT-qPCR was performed using an SYBR qPCR Master Mix (Q321-03,
Vazyme, Nanjing, China) by the following conditions: pre-heat at 95℃ for 10 s, then

followed with 39 cycles at 95°C for 5/10 s and 60°C for 30 s (72°C for 30 sec). The relative
mRNA levels were analyzed using the $2^{-\Delta\Delta Ct}$ method. Experiments were repeated at least
three times. Then the mean value was used to calculate the significant alterations in
statistical analysis.

Table 1. Primers used for RT-qPCR.

Gene		Primer (5′–3′)
ACTB	Forward	GGCTGTATTCCCCTCCATCG
	Reverse	CCAGTTGGTAACAATGCCATGT
HIF-1α	Forward	TCTCGGCGAAGCAAAGAGTC
	Reverse	AGCCATCTAGGGCTTTCAGATAA
Lgr5	Forward	CCTACTCGAAGACTTACCCAGT
	Reverse	GCATTGGGGTGAATGATAGCA
Lyz	Forward	GAGACCGAAGCACCGACTATG
	Reverse	CGGTTTTGACATTGTGTTCGC
ChrA	Forward	ATCCTCTCTATCCTGCGACAC
	Reverse	GGGCTCTGGTTCTCAAACACT
Muc2	Forward	GAAGCCAGATCCCGAAACCA
	Reverse	GAATCGGTAGACATCGCCGT
Tnf-α	Forward	GCACAGAAAGCATGATCCGC
	Reverse	CCCCATCTTTTGGGGGAGTG
Alf4	Forward	GGACAGATTGGATGTTGGAGAAAATG
	Reverse	GGAGATGGCCAATTGGGTTCAC
Chop	Forward	CTCGCTCTCCAGATTCCAGTC
	Reverse	CTTCATGCGTTGCTTCCCA
Alpi	Forward	CCAGCAGTAACTCACCTCATGG
	Reverse	GAAGCCTTGTGGATTCCTGCTG
Fabp1	Forward	ATGAACTTCTCCGGCAAGTACC
	Reverse	CTGACACCCCCTTGATGTCC
Fabp2	Forward	TCCCTACAGTCTAGCAGACGGA
	Reverse	CCAGAAACCTCTCGGACAGCAA

Whole mount staining and Immunofluorescence analysis

Organoids were cultured in 6-well plate containing 2 ml medium. Four days after passaging
and then treated with CoCl$_2$ for 24h, organoids were collected in ice-cold culture medium
and Cell Recovery Solution (BD Biosciences) was added and incubated for 1 hour to remove
the Matrigel. After being washed 3 times by medium, the organoids were fixed by 2%
formaldehyde overnight at 4°C. On the second day, organoids were permeabilized in 0.1%
Tween 20/PBS for 30 min at room temperature. Blocking was done by 0.2% Bovine serum
albumin (BSA) for one hour at 4°C. Then organoids were incubated in 1st antibody at 4°C
overnight. On the third day, three times of washing were performed with ice-cold 0.1%
Tween 20/PBS, followed by organoid sedimentation at room temperature. Then secondary

rabbit or mouse anti-rat IgG (H+L) reagent was added to incubate for 1 hour, followed by
three times washing with 0.1% Triton X-100. Finally, the pelleted organoids were mounted
on the glass slides. Images were taken by Spinning Disk Confocal Microscopy and five
random fields were chosen to observe the 1st antibody (Lyz, ChrA, Muc2, and Aldob)
positive cells and measure their percentage. The experiment was repeated for at least three
times.

Western blotting

Organoids cultured in the 24-well plate were washed by cold PBS and moved into 1.5 ml
tubes. Supernatant with Matrigel was removed and 80 μl 4X loading buffer (with DTT) was
added and pipetted for mixing. Samples were denatured at 100°C for 10 min. Protein
samples were stored at -80°C if not used immediately. Protein samples (around 50 μg) were
loaded and for electrophoresis on 8% sodium dodecyl sulfate-polyacrylamide gels, followed
by transferring to a polyvinylidene fluoride (PVDF) membrane (Bio-Rad, CA, USA). Then the
blocking was carried in non-fat dry milk at 37°C for 1 h. First antibody incubation was done
at 4 degrees with the following antibodies (1:1000). The rabbit monoclonal anti-HIF-1α
(#36169T), anti-β-Catenin (#8480) and anti-Cyclin-D1 (#2978) were purchased from Cell
Signaling Technology (Devers, MA, USA). The rabbit monoclonal anti-Axin2 (#2151) was
obtained from Abcam (Cambridge, UK). The mouse monoclonal anti-E-cadherin was
purchased from BD Biosciences (610181, San Jose, CA, USA). The mouse monoclonal, anti-
Ephb2 (sc-130752), anti-PKM (sc-365684) and anti-PPARγ (sc-7273) were purchased from
Santa Cruz (Dallas, TX, US). After 1st antibody incubation, membranes were washed by Tris-
Buffered Saline Tween (TBST)-20 for 3 times. Incubation with the secondary goat anti-rabbit
or goat anti-mouse IgG (1:1000, ZSGB-BIO, Beijing, China) were performed at 37°C for 1 h.
Then the PVDF membranes were washed three times for 10 min each time. The blots were
developed using an enhanced chemiluminescence detection kit (Millipore, MA, USA). Image
J was used to do the densitometric analysis.

Statistical analysis

352 GraphPad Prism 9.0 was used to perform statistical analysis. one-way ANOVA was used to
353 test the differences among groups, which is followed by Tukey's multiple comparisons test.
354 Each experiment was repeated at least 3 times. The results were deemed significantly
355 different at $P < 0.05$.

Author contributions

357 Xi Lan designed the paper, Xi Lan, Qiu Ping and Mou Chunfeng did the experiments, and
358 Mou Chunfeng approved the version to be published. All authors approved this version for
359 publication.

Acknowledgment

361 May this tiny work find the sincerest passion and effort of those unrooted scientists, in
362 honor of their unseen sweat and industrious missing contribution.

Conflict of interest

364 The authors declare no conflict of interest.

References

1.   Snippert HJ, van der Flier LG, Sato T, van Es JH, van den Born M, Kroon-Veenboer
C, et al. Intestinal crypt homeostasis results from neutral competition between symmetrically
dividing Lgr5 stem cells. Cell. 2010;143(1):134-44.
2.   Koo BK, Clevers H. Stem cells marked by the R-spondin receptor LGR5.
Gastroenterology. 2014;147(2):289-302.
3.   Potten CS, Gandara R, Mahida YR, Loeffler M, Wright NA. The stem cells of small
intestinal crypts: where are they? Cell Prolif. 2009;42(6):731-50.
4.   Barker N, van Es JH, Kuipers J, Kujala P, van den Born M, Cozijnsen M, et al.
Identification of stem cells in small intestine and colon by marker gene Lgr5. Nature.
2007;449(7165):1003-7.
5.   Scoville DH, Sato T, He XC, Li L. Current view: intestinal stem cells and signaling.
Gastroenterology. 2008;134(3):849-64.
6.   van der Flier LG, Clevers H. Stem cells, self-renewal, and differentiation in the
intestinal epithelium. Annu Rev Physiol. 2009;71:241-60.
7.   Crosnier C, Stamataki D, Lewis J. Organizing cell renewal in the intestine: stem cells,
signals and combinatorial control. Nat Rev Genet. 2006;7(5):349-59.
8.   Sato T, Vries RG, Snippert HJ, van de Wetering M, Barker N, Stange DE, et al.
Single Lgr5 stem cells build crypt-villus structures in vitro without a mesenchymal niche.
Nature. 2009;459(7244):262-5.
9.   de Lau W, Barker N, Low TY, Koo BK, Li VS, Teunissen H, et al. Lgr5 homologues
associate with Wnt receptors and mediate R-spondin signalling. Nature. 2011;476(7360):293-
7.
10.   Fodde R, Smits R, Clevers H. APC, signal transduction and genetic instability in
colorectal cancer. Nat Rev Cancer. 2001;1(1):55-67.
11.   Kinzler KW, Vogelstein B. Lessons from hereditary colorectal cancer. Cell.
1996;87(2):159-70.
12.   Kuhnert F, Davis CR, Wang HT, Chu P, Lee M, Yuan J, et al. Essential requirement
for Wnt signaling in proliferation of adult small intestine and colon revealed by adenoviral
expression of Dickkopf-1. Proc Natl Acad Sci U S A. 2004;101(1):266-71.
13.   Valenta T, Degirmenci B, Moor AE, Herr P, Zimmerli D, Moor MB, et al. Wnt
Ligands Secreted by Subepithelial Mesenchymal Cells Are Essential for the Survival of
Intestinal Stem Cells and Gut Homeostasis. Cell Rep. 2016;15(5):911-8.
14.   Anastas JN, Moon RT. WNT signalling pathways as therapeutic targets in cancer. Nat
Rev Cancer. 2013;13(1):11-26.
15.   Voloshanenko O, Erdmann G, Dubash TD, Augustin I, Metzig M, Moffa G, et al.
Wnt secretion is required to maintain high levels of Wnt activity in colon cancer cells. Nat
Commun. 2013;4:2610.
16.   Karhausen J, Haase VH, Colgan SP. Inflammatory hypoxia: role of hypoxia-inducible
factor. Cell Cycle. 2005;4(2):256-8.
17.   Zheng L, Kelly CJ, Colgan SP. Physiologic hypoxia and oxygen homeostasis in the
healthy intestine. A Review in the Theme: Cellular Responses to Hypoxia. Am J Physiol Cell
Physiol. 2015;309(6):C350-60.
18.   Glover LE, Lee JS, Colgan SP. Oxygen metabolism and barrier regulation in the
intestinal mucosa. The Journal of clinical investigation. 2016;126(10):3680-8.
19.   Karhausen J, Furuta GT, Tomaszewski JE, Johnson RS, Colgan SP, Haase VH.
Epithelial hypoxia-inducible factor-1 is protective in murine experimental colitis. The Journal
of clinical investigation. 2004;114(8):1098-106.
20.   Robinson A, Keely S, Karhausen J, Gerich ME, Furuta GT, Colgan SP. Mucosal
protection by hypoxia-inducible factor prolyl hydroxylase inhibition. Gastroenterology.
2008;134(1):145-55.

21.   Kong T, Westerman KA, Faigle M, Eltzschig HK, Colgan SP. HIF-dependent
induction of adenosine A2B receptor in hypoxia. FASEB J. 2006;20(13):2242-50.
22.   Cummins EP, Seeballuck F, Keely SJ, Mangan NE, Callanan JJ, Fallon PG, et al. The
hydroxylase inhibitor dimethyloxalylglycine is protective in a murine model of colitis.
Gastroenterology. 2008;134(1):156-65.
23.   Baregamian N, Rychahou PG, Hawkins HK, Evers BM, Chung DH.
Phosphatidylinositol 3-kinase pathway regulates hypoxia-inducible factor-1 to protect from
intestinal injury during necrotizing enterocolitis. Surgery. 2007;142(2):295-302.
24.   Ravizza R, Molteni R, Gariboldi MB, Marras E, Perletti G, Monti E. Effect of HIF-1
modulation on the response of two- and three-dimensional cultures of human colon cancer
cells to 5-fluorouracil. Eur J Cancer. 2009;45(5):890-8.
25.   Xie A, Robles RJ, Mukherjee S, Zhang H, Feldbrugge L, Csizmadia E, et al. HIF-
1alpha-induced xenobiotic transporters promote Th17 responses in Crohn's disease. J
Autoimmun. 2018;94:122-33.
26.   Hongo K, Tsuno NH, Kawai K, Sasaki K, Kaneko M, Hiyoshi M, et al. Hypoxia
enhances colon cancer migration and invasion through promotion of epithelial-mesenchymal
transition. J Surg Res. 2013;182(1):75-84.
27.   Markel TA, Crisostomo PR, Lahm T, Novotny NM, Rescorla FJ, Tector J, et al. Stem
cells as a potential future treatment of pediatric intestinal disorders. J Pediatr Surg.
2008;43(11):1953-63.
28.   Gupta A, Dixit A, Sales KM, Winslet MC, Seifalian AM. Tissue engineering of small
intestine--current status. Biomacromolecules. 2006;7(10):2701-9.
29.   Mazumdar J, O'Brien WT, Johnson RS, LaManna JC, Chavez JC, Klein PS, et al. O2
regulates stem cells through Wnt/beta-catenin signalling. Nat Cell Biol. 2010;12(10):1007-13.
30.   Rohwer N, Jumpertz S, Erdem M, Egners A, Warzecha KT, Fragoulis A, et al. Non-
canonical HIF-1 stabilization contributes to intestinal tumorigenesis. Oncogene.
2019;38(28):5670-85.
31.   Barker N, van Oudenaarden A, Clevers H. Identifying the stem cell of the intestinal
crypt: strategies and pitfalls. Cell Stem Cell. 2012;11(4):452-60.
32.   Lindemans CA, Calafiore M, Mertelsmann AM, O'Connor MH, Dudakov JA, Jenq
RR, et al. Interleukin-22 promotes intestinal-stem-cell-mediated epithelial regeneration.
Nature. 2015;528(7583):560-4.
33.   Nalapareddy K, Nattamai KJ, Kumar RS, Karns R, Wikenheiser-Brokamp KA,
Sampson LL, et al. Canonical Wnt Signaling Ameliorates Aging of Intestinal Stem Cells.
Cell Rep. 2017;18(11):2608-21.
34.   Cosin-Roger J, Simmen S, Melhem H, Atrott K, Frey-Wagner I, Hausmann M, et al.
Hypoxia ameliorates intestinal inflammation through NLRP3/mTOR downregulation and
autophagy activation. Nat Commun. 2017;8(1):98.
35.   Xie C, Yagai T, Luo Y, Liang X, Chen T, Wang Q, et al. Activation of intestinal
hypoxia-inducible factor 2alpha during obesity contributes to hepatic steatosis. Nat Med.
2017;23(11):1298-308.
36.   Zhou X, Tu J, Li Q, Kolosov VP, Perelman JM. Hypoxia induces mucin expression
and secretion in human bronchial epithelial cells. Transl Res. 2012;160(6):419-27.
37.   Ghosh JC, Altieri DC. Activation of p53-dependent apoptosis by acute ablation of
glycogen synthase kinase-3beta in colorectal cancer cells. Clinical cancer research : an
official journal of the American Association for Cancer Research. 2005;11(12):4580-8.
38.   Booth C, Brady G, Potten CS. Crowd control in the crypt. Nat Med. 2002;8(12):1360-
1.

39.   Clevers H, Loh KM, Nusse R. Stem cell signaling. An integral program for tissue renewal and regeneration: Wnt signaling and stem cell control. Science. 2014;346(6205):1248012.

40.   Ayyaz A, Kumar S, Sangiorgi B, Ghoshal B, Gosio J, Ouladan S, et al. Single-cell transcriptomes of the regenerating intestine reveal a revival stem cell. Nature. 2019;569(7754):121-5.

41.   Batlle E, Henderson JT, Beghtel H, van den Born MM, Sancho E, Huls G, et al. Beta-catenin and TCF mediate cell positioning in the intestinal epithelium by controlling the expression of EphB/ephrinB. Cell. 2002;111(2):251-63.

42.   Merlos-Suarez A, Barriga FM, Jung P, Iglesias M, Cespedes MV, Rossell D, et al. The intestinal stem cell signature identifies colorectal cancer stem cells and predicts disease relapse. Cell Stem Cell. 2011;8(5):511-24.

43.   Lustig B, Jerchow B, Sachs M, Weiler S, Pietsch T, Karsten U, et al. Negative feedback loop of Wnt signaling through upregulation of conductin/axin2 in colorectal and liver tumors. Mol Cell Biol. 2002;22(4):1184-93.

44.   Pelaseyed T, Bergstrom JH, Gustafsson JK, Ermund A, Birchenough GM, Schutte A, et al. The mucus and mucins of the goblet cells and enterocytes provide the first defense line of the gastrointestinal tract and interact with the immune system. Immunol Rev. 2014;260(1):8-20.

45.   Peyssonnaux C, Datta V, Cramer T, Doedens A, Theodorakis EA, Gallo RL, et al. HIF-1alpha expression regulates the bactericidal capacity of phagocytes. The Journal of clinical investigation. 2005;115(7):1806-15.

46.   Adolph TE, Tomczak MF, Niederreiter L, Ko HJ, Bock J, Martinez-Naves E, et al. Paneth cells as a site of origin for intestinal inflammation. Nature. 2013;503(7475):272-6.

47.   Heijmans J, van Lidth de Jeude JF, Koo BK, Rosekrans SL, Wielenga MC, van de Wetering M, et al. ER stress causes rapid loss of intestinal epithelial stemness through activation of the unfolded protein response. Cell Rep. 2013;3(4):1128-39.

48.   Kaser A, Lee AH, Franke A, Glickman JN, Zeissig S, Tilg H, et al. XBP1 links ER stress to intestinal inflammation and confers genetic risk for human inflammatory bowel disease. Cell. 2008;134(5):743-56.

49.   Salvo Romero E, Alonso Cotoner C, Pardo Camacho C, Casado Bedmar M, Vicario M. The intestinal barrier function and its involvement in digestive disease. Rev Esp Enferm Dig. 2015;107(11):686-96.

50.   Blikslager AT. Life in the gut without oxygen: adaptive mechanisms and inflammatory bowel disease. Gastroenterology. 2008;134(1):346-8.

51.   Araujo JR, Tazi A, Burlen-Defranoux O, Vichier-Guerre S, Nigro G, Licandro H, et al. Fermentation Products of Commensal Bacteria Alter Enterocyte Lipid Metabolism. Cell Host Microbe. 2020;27(3):358-75 e7.

52.   Ramachandran D, Clara R, Fedele S, Michel L, Burkard J, Kaufman S, et al. Enhancing enterocyte fatty acid oxidation in mice affects glycemic control depending on dietary fat. Sci Rep. 2018;8(1):10818.

53.   Gutting T, Weber CA, Weidner P, Herweck F, Henn S, Friedrich T, et al. PPARgamma-activation increases intestinal M1 macrophages and mitigates formation of serrated adenomas in mutant KRAS mice. Oncoimmunology. 2018;7(5):e1423168.

54.   Bagby GJ, Lang CH, Giaimo ME, Spitzer JJ. Increased glucose metabolism by epitrochlearis muscle removed from endotoxin-treated rats. Circ Shock. 1986;20(3):171-9.

55.   L'Her E, Sebert P. A global approach to energy metabolism in an experimental model of sepsis. Am J Respir Crit Care Med. 2001;164(8 Pt 1):1444-7.

56. Sun X, Yao L, Liang H, Wang D, He Y, Wei Y, et al. Intestinal epithelial PKM2
serves as a safeguard against experimental colitis via activating beta-catenin signaling.
Mucosal Immunol. 2019;12(6):1280-90.
57. Vadde R, Vemula S, Jinka R, Merchant N, Bramhachari PV, Nagaraju GP. Role of
hypoxia-inducible factors (HIF) in the maintenance of stemness and malignancy of colorectal
cancer. Crit Rev Oncol Hematol. 2017;113:22-7.
58. Dong HJ, Jang GB, Lee HY, Park SR, Kim JY, Nam JS, et al. The Wnt/beta-catenin
signaling/Id2 cascade mediates the effects of hypoxia on the hierarchy of colorectal-cancer
stem cells. Sci Rep. 2016;6:22966.
59. Liu J, Zhu H, Li B, Robinson SC, Lee C, O'Connell JS, et al. Lactoferrin Reduces
Necrotizing Enterocolitis Severity by Upregulating Intestinal Epithelial Proliferation. Eur J
Pediatr Surg. 2020;30(1):90-5.
60. Zu G, Guo J, Che N, Zhou T, Zhang X, Wang G, et al. Protective effects of
ginsenoside Rg1 on intestinal ischemia/reperfusion injury-induced oxidative stress and
apoptosis via activation of the Wnt/beta-catenin pathway. Sci Rep. 2016;6:38480.
61. Obata Y, Takahashi D, Ebisawa M, Kakiguchi K, Yonemura S, Jinnohara T, et al.
Epithelial cell-intrinsic Notch signaling plays an essential role in the maintenance of gut
immune homeostasis. J Immunol. 2012;188(5):2427-36.
62. Nakamura T, Tsuchiya K, Watanabe M. Crosstalk between Wnt and Notch signaling
in intestinal epithelial cell fate decision. J Gastroenterol. 2007;42(9):705-10.
63. Takahashi T, Shiraishi A. Stem Cell Signaling Pathways in the Small Intestine.
International journal of molecular sciences. 2020;21(6).
64. Gustafsson MV, Zheng X, Pereira T, Gradin K, Jin S, Lundkvist J, et al. Hypoxia
requires notch signaling to maintain the undifferentiated cell state. Dev Cell. 2005;9(5):617-
28.
65. Ortiz-Masia D, Cosin-Roger J, Calatayud S, Hernandez C, Alos R, Hinojosa J, et al.
M1 Macrophages Activate Notch Signalling in Epithelial Cells: Relevance in Crohn's
Disease. J Crohns Colitis. 2016;10(5):582-92.
66. Liu Z, Li L, Chen W, Wang Q, Xiao W, Ma Y, et al. Aryl hydrocarbon receptor
activation maintained the intestinal epithelial barrier function through Notch1 dependent
signaling pathway. Int J Mol Med. 2018;41(3):1560-72.
67. Yeung TM, Gandhi SC, Bodmer WF. Hypoxia and lineage specification of cell line-
derived colorectal cancer stem cells. Proc Natl Acad Sci U S A. 2011;108(11):4382-7.
68. Nusrat A, Delp C, Madara JL. Intestinal epithelial restitution. Characterization of a
cell culture model and mapping of cytoskeletal elements in migrating cells. The Journal of
clinical investigation. 1992;89(5):1501-11.
69. Hidalgo IJ, Raub TJ, Borchardt RT. Characterization of the human colon carcinoma
cell line (Caco-2) as a model system for intestinal epithelial permeability. Gastroenterology.
1989;96(3):736-49.
70. Synnestvedt K, Furuta GT, Comerford KM, Louis N, Karhausen J, Eltzschig HK, et al.
Ecto-5'-nucleotidase (CD73) regulation by hypoxia-inducible factor-1 mediates permeability
changes in intestinal epithelia. The Journal of clinical investigation. 2002;110(7):993-1002.
71. Furuta GT, Turner JR, Taylor CT, Hershberg RM, Comerford K, Narravula S, et al.
Hypoxia-inducible factor 1-dependent induction of intestinal trefoil factor protects barrier
function during hypoxia. The Journal of experimental medicine. 2001;193(9):1027-34.
72. Colgan SP, Taylor CT. Hypoxia: an alarm signal during intestinal inflammation. Nat
Rev Gastroenterol Hepatol. 2010;7(5):281-7.
73. Sato T, van Es JH, Snippert HJ, Stange DE, Vries RG, van den Born M, et al. Paneth
cells constitute the niche for Lgr5 stem cells in intestinal crypts. Nature.
2011;469(7330):415-8.

74.　　Beumer J, Clevers H. Regulation and plasticity of intestinal stem cells during
homeostasis and regeneration. Development. 2016;143(20):3639-49.
75.　　Hamilton MJ, Stevens RL. Stress in the ER leads to inflammatory bowel disease.
Inflamm Bowel Dis. 2009;15(11):1765-6.
76.　　Chen X, Iliopoulos D, Zhang Q, Tang Q, Greenblatt MB, Hatziapostolou M, et al.
XBP1 promotes triple-negative breast cancer by controlling the HIF1alpha pathway. Nature.
2014;508(7494):103-7.

A

CoCl$_2$ conc.　　0　　　10 µM　　　50 µM　　　100 µM　　　200 µM

B

CoCl$_2$ (µM)　　0　10　50　100　200

HIF-1α　　　　　　　　　　　　　　　120 kDa

β-actin　　　　　　　　　　　　　　43 kDa

HIF-1α/β-actin　　0.04　0.05　0.31*　0.41*　0.76*

C

DMSO　　　　　　50 µM CoCl$_2$　　　　　200 µM CoCl$_2$

A

B

A

B

C