Gliogenesis from the subventricular zone modulates the extracellular matrix at the glial scar after brain ischemia

Maria Ardaya[1,2,3,9], Marie-Catherine Tiveron[4], Harold Cremer[4], Benjamin Dehay[5], Fernando Pérez-Cerdá[1,2,3], Carlos Matute[1,2,3], Federico N. Soria[1,2,3,6,*], Fabio Cavaliere[1,2,3,7,8,*]

[1]Achucarro Basque Center for Neuroscience, Leioa, Spain.

[2]Department of Neuroscience, University of the Basque Country (UPV/EHU), Leioa, Spain.

[3]CIBERNED, Madrid, Spain.

[4]Aix Marseille Univ, CNRS, IBDM, Campus de Luminy, Marseille, France.

[5]Univ. Bordeaux, CNRS, IMN, UMR 5293, F-33000 Bordeaux, France.

[6]Ikerbasque, Basque Foundation for Science, Bilbao, Spain.

[7]Basque Biomodel Platform for Human Research (BBioH), Leioa, Spain.

[8]Current Address: Instituto Biofisika (CSIC, UPV/EHU), Fundación Biofísica Bizkaia, Leioa, Spain

[9]Current Address: Donostia International Physics Center (DIPC), San Sebastian, Spain.

[*]Correspondence to fabio.cavaliere@ehu.eus (F.C.), federico.soria@achucarro.org (F.N.S.)

ABSTRACT

Activation of the subventricular zone (SVZ) following cerebral ischemia is one of the brain's early

responses to counteract neuron loss and minimize tissue damage. Impaired brain regions communicate

with the SVZ through various chemotactic signals that promote cell migration and differentiation,

primarily involving neural stem cells (NSC), neuroblasts, or glioblasts. However, the activation of

gliogenesis and the role of newly formed astrocytes in the post-ischemic scenario remain subjects of

debate. We have previously demonstrated that adenosine release after brain ischemia prompts the SVZ

to generate new astrocytes. Here, we used transient brain ischemia in mice to identify the cellular origin

of these astrocytes within the SVZ neurogenic niche and to investigate their role in the pathological

process. By combining immunofluorescence, BrdU-tracing, and genetic cell labeling, we tracked the

migration of newborn astrocytes, positive for the proteoglycan marker Thbs4, from the dorsal and

medial SVZ to the perilesional barrier surrounding the ischemic core, known as the "glial scar". We

found that these Thbs4-positive astrocytes modulate the dense extracellular matrix at the lesion border

by both synthesizing and degrading hyaluronan. We also show that while the accumulation of this

polymer at the lesion site is sufficient to recruit newborn astrocytes, its degradation at the SVZ correlates

with gliogenesis. These findings suggest that newborn astrocytes could be a promising pharmacological

target for modulating the glial scar after brain ischemia and facilitate tissue regeneration.

INTRODUCTION

Cell regeneration in the adult mammalian brain primarily occurs mainly in two neurogenic niches: the subventricular zone (SVZ) in the lateral ventricle and the subgranular zone (SGZ) in the dentate gyrus of the hippocampus (Bond et al., 2015). Adult brain stem cells are multipotent and originate from the embryonic radial glia, thus sharing **key** cell markers and features with resident astrocytes. In the adult mouse brain, neuroblasts from SVZ migrate through the rostral migratory stream (RMS) to the olfactory bulb (OB), where neurogenesis takes place (Luskin, 1993). Neuronal hyperactivity (e.g., during epilepsy), as well as pathological conditions (e.g., stroke) or natural (e.g., aging) processes, can impair the physiological functions of the neurogenic niches, altering their neurogenic and gliogenic potential (Chaker et al., 2016; Kralic et al., 2005; Sundholm-Peters et al., 2005). For instance, it is known that SVZ cells are activated *in vivo* after focal ischemia following a middle cerebral artery occlusion (MCAO) (Ceanga et al., 2021).

We have previously demonstrated in oxygen and glucose-deprived (OGD) organotypic brain slices that the ischemic region and the SVZ establish a biochemical interplay based on a gradient of toxic and protective factors released early after the insult (Cavaliere et al., 2006). Furthermore, others have demonstrated that brain stroke models in mice can activate astrogliogenesis at the expense of neurogenesis (Benner et al., 2013; Laug et al., 2019; Li et al., 2010). Benner and colleagues identified a SVZ-derived astrocyte population characterized by high levels of the membrane-bound proteoglycan thrombospondin 4 (Thbs4), a cell-cell and cell-matrix adhesion molecule that has been considered neuroprotective against ischemic damage (Benner et al., 2013; Laug et al., 2019). Ablation of Thbs4 in KO mice also produced abnormal ischemic scar formation and increased microvascular hemorrhage, suggesting a role of SVZ astrogliogenesis in brain damage and glial scar formation induced by ischemic stroke (Benner et al., 2013).

The perilesional barrier known as the *glial scar* assembles early after brain ischemia, isolating the damaged area and limiting both the spread of pathogens and potentially toxic damage-associated molecular patterns (DAMPs) into the surrounding area (Anderson et al., 2016; Bush et al., 1999;

Faulkner et al., 2004). Consisting of packed reactive astrocytes and a dense extracellular matrix

composed mostly of hyaluronic acid (HA), the glial scar represents a physical barrier for the SVZ-

generated newborn neurons that migrate into the ischemic core (Arvidsson et al., 2002; Yamashita et

al., 2006). Thus, modulating glial scar formation after brain ischemia may be a strategy to reduce tissue

damage and restore brain functionality. In ischemic stroke, the brain extracellular matrix (ECM) goes

through a significant reorganization due to injury and must undergo successful remodeling to promote

brain repair. ECM elements involved in the ischemic cascade induce activation of astrocytes and

microglia, which modulate ECM structure by releasing glycoproteins into the extracellular space

(Adams and Gallo, 2018; Fawcett, 2015). This crosstalk culminates in the formation of the glial scar,

which is created locally by reactive astrocytes and astrocytes-secreted ECM molecules such as HA and

chondroitin sulphate proteoglycans (Bush et al., 1999; Lau et al., 2013).

We have previously demonstrated that high extracellular adenosine levels, a hallmark of ischemic

insults, induce astrogliogenesis from the SVZ (Benito-Muñoz et al., 2016). Following our previous

results, here we characterize the SVZ response to brain ischemia and investigate the role of newly

generated astrocytes in the modulation of the glial scar after MCAO. We show that ischemic conditions

increase the number of newly generated astrocytes in the SVZ and their migration to the ischemic area.

We also propose the role of these astrocytes in modulating the extracellular matrix at the glial scar and

suggest this astrocyte population as a possible target for brain repair after brain ischemia.

RESULTS

Thbs4 is expressed in the neurogenic niche of the SVZ.

As suggested by others (Benner et al., 2013; Girard et al., 2014; Pous et al., 2020), Thbs4 is highly

expressed in astrocytes of the SVZ and RMS and might play a role in ischemia-induced neurogenesis.

To confirm these observations, we labeled the mouse SVZ by immunofluorescence for Thbs4. Thbs4

was mainly expressed in the SVZ, RMS, and OB but also in the cerebellum and ventral tegmental area

(Fig. 1A). In the SVZ, the staining was associated with cells with different morphologies (Fig. 1B), ranging from cells with small primary apical processes in contact with the ventricle, to rounded NSC-like cells in the dorsolateral SVZ horn. After quantification by direct count of Thbs4-positive cells, Thbs4-positive astrocytes represented 21.4% of the total SVZ cells (Fig. 1C), with more than 60% of them displaying a radial glia-like type-B morphology (Bond et al., 2015; Doetsch et al., 1999) (Fig. 1D). This is in accordance with other studies reporting Thbs4 expression in B-cells, either by immunohistochemistry (Beckervordersandforth et al., 2010; Benner et al., 2013) or single-cell RNA sequencing (Basak et al., 2018; Cebrian-Silla et al., 2021; Llorens-Bobadilla et al., 2015).

To validate the immunohistochemistry protocol for Thbs4, we labeled the SVZ cells with green fluorescent protein (GFP) under the control of the Thbs4 promoter (pThbs4-eGFP) by postnatal (P1) electroporation (Fig. 1E). Twenty-four hours after electroporation, almost 20% of total cells in the dorsal SVZ expressed GFP (Fig. 1F), with a greater expression in the rostral SVZ (Fig. 1G), confirming that Thbs4-positive cells were mainly localized in the neurogenic niche of the SVZ. This also suggests that Thbs4 is expressed in NSCs at early postnatal stages, generating astrocytes alongside development.

Thbs4 expression in the SVZ increases after brain ischemia

To investigate whether brain ischemia could activate astrogliogenesis in the SVZ, we analyzed cell proliferation and Thbs4 expression in the SVZ after MCAO. Sixty minutes of MCAO (Fig. S1A) caused neuronal damage in the cortex and striatum, as observed by cresyl violet staining (Fig. S1B) and NeuN immunofluorescence (Fig. S1C). The survival rate at 28 days post-lesion was around 50% (Fig. S1D). Moreover, motor deficits and animal weight of ischemic mice were monitored every day after MCAO for 30 days. Motor deficits improved overtime, while body weight dropped sharply the first 3 days after MCAO and was partially restored at 30 dpi (Fig. S1E-F).

To study cell proliferation in the SVZ, we administered 50 mg/kg BrdU i.p. every two hours (3 doses in total) the day before MCAO (Fig. S2A). We observed an increase in the number of total SVZ BrdU-

positive cells 24h after MCAO (Fig. S2B-C). Ki67 immunofluorescence of SVZ wholemount preparations (Fig. S2D) also revealed a more than tenfold increase of proliferative cells 24h after MCAO (Fig. S2E). In addition, we observed a significant increase in cleaved Caspase3-positive cells 24h post-lesion (Fig. S2F-G). These results suggest a fast and specific proliferative response in the ischemia-induced SVZ.

We next analyzed Thbs4 staining in combination with Nestin, an NSC marker, and Doublecortin (DCX), a marker of neuroblasts, before and after MCAO. SVZ was analyzed in sham and ischemic mice 7-, 15- and 30-days post-ischemia (dpi). The immunofluorescence analysis of the SVZ (Fig. 2A) showed a transient decrease in Nestin and DCX-positive cells but an increase of Thbs4-positive astrocytes up to 15 dpi (Fig. 2B-D) compared to sham animals. The increase of Thbs4 astrocytes was also confirmed by Western blot analysis using SVZ homogenates (Fig. 2E-F), evidencing a 2-fold increased expression of Thbs4 in the ischemic SVZ. Similarly to immunofluorescence, Western blot also showed decreased Nestin protein levels at 15 dpi (Fig. 2G). These results demonstrate that Thbs4 expression in the SVZ increases after ischemia, coinciding with a decrease in neuroblast markers, suggesting that the classical pro-neuroblast program of the SVZ is interrupted in favor of the generation of astrocytes after ischemia.

Characterization of ischemia-induced cell populations in the SVZ

To characterize the different cell populations of the SVZ activated after focal ischemia, we labeled the whole pool of proliferating cells with 1% BrdU in drinking water for 14 days before MCAO (Codega et al., 2014). One month after MCAO, BrdU in the SVZ primarily stained slow proliferative cells like type B cells, whereas in rapid transit-amplifying cells (type C), BrdU was diluted away. Therefore, we chose to label type C cells by intraperitoneal administration (50 mg/kg) of 5-Iodo-2′-deoxyuridine (IdU) 24h before euthanasia (Fig. 3A). To identify the cellular origin of newborn astrocytes, we analyzed the SVZ 30 dpi by immunofluorescence co-labeling Thbs4 and GFAP together with BrdU or IdU. We found a 2-fold increase in the total amount of proliferative cells (BrdU-positive) after MCAO with

respect to the sham group (Fig. 3B), whereas we did not observe a significant change in the number of IdU-positive cells (Fig. 3C). The number of slow proliferative NSC (Thbs4/GFAP/BrdU) increased linearly after MCAO (Fig. 3D, E) especially in the dorsal region of the SVZ (Fig. S3A), whereas the number of activated NSC (Thbs4/GFAP/IdU) did not change significantly, either in the whole SVZ (Fig. 3F) or in any subregion analyzed (Fig. S3B). These results suggest that, after ischemia, Thbs4-positive astrocytes derive from the slow proliferative type B cells.

Ischemia-induced newborn astrocytes migrate from the dorsal SVZ to ischemic regions

To confirm that Thbs4-positive astrocytes originate from type B cells of dorsal SVZ, we labeled NSCs of dorsal SVZ by electroporating the pCAGGSx-CRE plasmid in P1 Ai14 Rosa26-CAG-tdTomato transgenic mice (Fig. 4A). Plasmid electroporation induced the expression of tdTomato (tdTOM) by Cre recombination, enabling the spatial tracking and cell fate assessment of electroporated NCSs from the dorsal SVZ. Three to four months after electroporation, we performed the MCAO, and tdTOM fluorescence was analyzed at 7, 30, and 60 dpi. To confirm the efficient labeling of NSCs by the electroporation protocol, we followed the tdTOM fluorescence in the OB (Fig. S4A, see methods). Only those animals with tdTOM-positive cells in the OB were taken into account for later analysis. After quantification of tdTOM-positive NSCs 7, 30, and 60 dpi, we observed a significant decrease in tdTOM-positive cells in ischemic animals in all SVZ regions analyzed (Fig. 4B-C and Fig S4B), suggesting that they either left the area, or died. Conversely, there was a general increase of tdTOM-positive cells outside the SVZ (Fig. 4D-E), especially in the intermediate and caudal axis of the whole brain compared to sham animals (Fig. S4D-G), suggesting that electroporated cells had migrated away from the SVZ.

To identify the proportion of NSCs that differentiated into Thbs4-positive astrocytes following ischemia, we analyzed the co-expression of Thbs4 with tdTOM after MCAO. Despite the general decrease of tdTOM-positive cells in the SVZ, the fraction of Thbs4/tdTOM cells increased at 30 dpi after MCAO (Fig. 4F-G), reaching a maximum increase at 60 dpi (Fig. 4H), in particular in rostral and intermediate SVZ (Fig. S4C). Finally, we analyzed the Thbs4-positive astrocytes recruitment to the

164   ischemic regions. More than half of the total tdTOM-positive cells in the infarcted area expressed the

165   Thbs4 marker at 60 dpi (Fig. 4I-J). These results suggest that ischemia-induced astrogliogenesis in the

166   SVZ occurs in type B cells from the dorsal region and that these newborn Thbs4-positive astrocytes

167   migrate to the ischemic areas.

169   Thbs4-positive astrocytes accumulate at the glial scar after brain ischemia.

Besides its role in the neurogenic niches, Nestin is also rapidly and locally expressed by reactive

astrocytes and participates in glial scar formation (Frisén et al., 1995). Thus, we used Nestin

immunofluorescence as a reliable tool to identify the lesion border where the glial scar occurs (Fig. 5A).

Nestin staining of the scar was also compared with more classical methodologies like TTC staining

(Fig. 5B), where viable tissue is stained red and dead tissue is left unstained. The white-pale pink area

was then measured as infarct tissue by TTC staining (Fig. 5C). Unlike the TTC staining, which can only

be performed shortly after the lesion, the visualization of the glial scar with Nestin immunofluorescence

allowed us to analyze the lesion at later time-points. We observed a rapid formation of the glial scar as

early as 7 dpi (Fig. 5D). Analyzing the expression of the two astrocytic markers, Thbs4 and GFAP, as

a percentage of the glial scar region-of-interest (ROI), we observed different dynamics and localization.

GFAP expression increased rapidly after 7 days within the glial scar ROI, whereas Thbs4 was

upregulated at 15 dpi (Fig. 5E). This delay suggested that the glial scar harbors two different astrocytic

sub-populations: GFAP-only or GFAP/Thbs4. To analyze the spatial location of GFAP and Thbs4

markers within the Nestin border, we measured the fluorescence intensity profile of damaged area at 30

dpi. GFAP was found in both the scar and core areas, whereas Thbs4 fluorescence was found mostly at

the borders of the glial scar (Fig. 5F-G). Analysis and dynamics of Thbs4-positive astrocyte migration

were further performed in the corpus callosum (CC), striatum (Str), cortex (Ctx) and olfactory bulb

(OB) (Fig. S5A). We observed a linear increase of Thbs4-positive astrocytes from 7 to 60 dpi, both in

the Ctx and Str (Fig. S5B). Migration occurred mainly through the corpus callosum; among the total

GFAP-positive astrocytes, 36.5% were also positive for Thbs4, suggesting that more than one-third of

the astrocytes in the glial scar originated from the SVZ (Fig. S5C). Likewise, the number of DCX-

positive cells dramatically decreased 15 dpi in all OB layer (Fig. S5D-F) together with a significant

decrease of BrdU-positive cells (Fig. S5G) DCX/BrdU- positive neuroblasts in the RMS (Fig. S5H),

suggesting that ischemia induced a change in the neuroblasts' ectopic migratory pathway, depriving the

OB layers of SVZ newborn neurons.

196   Thbs4-positive astrocytes modulate extracellular matrix homeostasis at the glial scar

To investigate the role of Thbs4-positive astrocytes in the glial scar, we analyzed hyaluronic acid (HA),

the main component of the brain extracellular matrix (ECM), using the hyaluronic acid binding protein

(HABP) as a marker. Following ischemic damage, local astrocytes (GFAP+/Thbs4-) react by adopting

a fibrotic morphology, producing high-molecular weight HA (HMW-HA) to form the glial scar and

isolate the damaged area (Preston and Sherman, 2011). We then examined HA deposition in the corpus

callosum and the infarcted areas after MCAO. HABP immunofluorescence (Fig. 6A) and image

analysis revealed HA accumulation in the infarcted areas. Skeleton analysis, which quantifies the length

of HA chains and correlates with its molecular weight, identified progressively longer HA cable-like

structures at the glial scar as the infarct evolved (Fig. 6B). Fractal dimension analysis, which reflects

the interconnectivity of the extracellular matrix (Soria et al., 2020), further confirmed HA accumulation

and increased matrix complexity in infarcted areas, but not in corpus callosum (Fig. 6C), suggesting

that ECM accumulation is different in white and gray matter.

The proximity of Thbs4-positive astrocytes to HA in the ischemic scar (Fig. 6D-E) led us to investigate

if this population contributed to ECM production and, consequently, to scar formation. Since HA is

produced exclusively at the cell membrane (Sherman et al., 2015; Zimmermann and Dours-

Zimmermann, 2008), we analyzed HA labeling within 1 µm of the Thbs4-positive cell membrane in

high-resolution confocal images (Fig. 6F). Image analysis showed a significant increase in HABP area

at Thbs4-positive astrocyte-cell membranes after MCAO (Fig. 6G), suggesting a role of Thbs4-positive

astrocytes in HA production within the glial scar.

To explore the potential role of Thbs4-positive astrocytes in HA removal, we quantified intracellular

HABP spots in Thbs4-positive astrocytes at the glial scar. While HMW-HA cleavage occurs

extracellularly, HA fragments can be internalized for degradation by lysosomal hyaluronidases (Preston

and Sherman, 2011; Zimmermann and Dours-Zimmermann, 2008). Intracellular HABP spots were

compared between Thbs4-positive astrocytes and local astrocytes (GFAP-positive/Thbs4-negative)

(Fig. S6A). Thirty days after MCAO, Thbs4-positive astrocytes showed more intracellular HABP than

local astrocytes in the infarcted cortex (Fig. S6B). However, the number of HA spots internalized was

higher in local astrocytes compared with Thbs4 astrocytes (Fig. S6C), suggesting that Thbs4-positive

cells internalize HA in fewer vesicles. To confirm that astrocytes were degrading HA, we performed an

immunofluorescence analysis of CD44 (Fig. S6D). CD44 is a transmembrane glycoprotein receptor that

mediates HA internalization and degradation, and therefore ECM remodeling (Dzwonek and

Wilczynski, 2015). We detected that around 50% of the total Thbs4-positive astrocytes contained

internalized CD44 receptor, in the striatum 30 dpi (Fig. S6E), with higher numbers (60%) for the

infarcted corpus callosum. These findings suggest that Thbs4- positive astrocytes participate in both the

formation and degradation of the extracellular matrix at the glial scar.

To further investigate HA internalization by astrocytes, we conducted *in vitro* experiments using rat

SVZ neural stem cells cultures. An oxygen and glucose deprivation (OGD) protocol was applied as an

ischemia model and cells were fixed 7 days after OGD (Fig. S7A). HA was analyzed via HABP

immunofluorescence (Fig. S7B), both inside and outside the Thbs4-positive cells (Fig. S7C). Thbs4-

positive cells showed increased intracellular HA after OGD (Fig. S7D), while extracellular HA levels

decreased (Fig. S7E). However, this phenomenon was reversed after adding hyaluronidases in the

extracellular space (Fig. S7F-G), suggesting that Thbs4-positive cells from SVZ homogenates can

internalize or synthesize HA depending on external stimuli. HA internalization was also measured in

Iba1-positive cell cultures where we observed a reduced internalization compare to Thbs4-positive cells

(Fig. S7H). Lastly, metalloproteinase activity was found to be increased in Thbs4 cultures following

OGD (Fig. S7I). Altogether, these results suggest Thbs4-positive cells can internalize HA after ischemic

conditions but also synthetize it when ECM is degraded.

Hyaluronan accumulation is sufficient but not exclusive for the recruitment of Thbs4-positive astrocytes

Since Thbs4-positive astrocytes migrate to the glial scar but not to the ischemic core, we hypothesized

that HA, the primary component of the ischemic scar, may be the trigger signal for recruiting Thbs4-

positive astrocytes. To test this hypothesis, we simulated the ECM accumulation that occurs at the scar

by overexpressing the Hyaluronan synthase 2 (HAS2) from the naked mole rat (*Heterocephalus glaber*),

a tool often used in mice to produces HA of very high molecular weight (Tian et al., 2013; Zhang et al.,

2023). We used a viral vector (AAV2/9-CAG-HAS2-3xFlag) with a FLAG tag to visualize transduced

cells after intracerebral inoculation in the striatum and checked HA accumulation in FLAG-positive

areas using HABP staining (Fig. 7A). We overexpressed HAS2 (or GFP as control) unilaterally. In a

subset of animals that underwent MCAO, AAV-HAS2 was injected contralateral to the infarct, to

compare Thbs4-positive astrocytes recruitment by both stimuli, the inflammation from the ischemic

damage and the HA accumulation *per se* (Fig. 7B). FLAG-positive cells were quantified as a surrogate

marker of viral infection for both control and MCAO groups (Fig. 7C). The GFP control group did not

increase HA production *per se* (Fig. 7D). However, HA overexpression was observed in HAS2 groups

(both control and MCAO group) by HABP immunofluorescence (Fig. 7E-F).

HA accumulation was sufficient to recruit the Thbs4-positive astrocytes from the ipsilateral SVZ 45

days after AAV-HAS2 inoculation (Fig. 7G), with a similar time window to that observed after MCAO.

Interestingly, when we induced a lesion by MCAO and performed viral overexpression of HAS2 in the

contralateral hemisphere, we found that Thbs4-positive astrocytes preferentially migrated to the

ischemic glial scar (Fig. 7G). This migration of Thbs4-positive astrocytes to the site of ischemic damage

occurred despite no differences in HA overexpression between contralateral (HAS2 infected) and

ipsilateral (infarcted area) hemispheres (Fig. 7H). These results suggest that the accumulation of HMW-

HA alone is sufficient to activate and recruit Thbs4-positive astrocytes from the SVZ. However, the

preferential migration to the ischemic scar indicates that brain ischemia generates a more potent

biochemical signal that not only activates the SVZ but also recruits newborn astrocytes.

271   Ischemia-induced gliogenesis at the SVZ correlates with local hyaluronan degradation

It is known that the niche microenvironment has a unique ECM composition (Kjell et al., 2020) and

that the stiffness of the niche plays a crucial role in the proliferation and differentiation of progenitor

cells (Long and Huttner, 2019; Segel et al., 2019). After observing the close association between Thbs4-

positive astrocytes and the ECM at the glial scar, we decided to study the Thbs4/HA tandem in the SVZ

following MCAO (Fig. 8A). We did not observe any significant changes in the HA area across the

entire SVZ but noted a decrease at 7 and 15 dpi in the dorsal SVZ (Fig. 8B-D). Skeleton analysis showed

fragmentation of HA exclusively in the dorsal SVZ at 7 and 15 dpi (Fig. 8E-G), i.e., earlier than the

migration of these newborn astrocytes toward the lesion.

Next, we examined the expression of genes related to HA degradation (*Hyase 1, 2* and *3*) and synthesis

(*HAS 1* and *HAS 2*). We also analyzed *Thbs4* and hypoxia-inducible factor 1-alpha (*Hif1a*), which are

known to be upregulated in the SVZ following brain ischemia (Benito-Muñoz et al., 2016; Liu et al.,

2007). Fresh SVZ tissue was collected from sham, 15 and 30 dpi mice (Fig. 8H). We observed an

increased expression of all Hyases at 15 dpi and slightly decreased by 30 dpi, whereas synthases (HAS)

showed a gradual increase from 15 to 30 dpi (Fig. 8I). Additionally, we observed an upregulation of

Thbs4 and Hif1a both at 15 and 30 dpi (Fig. 8I). These results suggest that HA degradation in the SVZ

is correlated with the Thbs4 response to ischemia.

289   DISCUSSION

The effects of brain insult on adult neurogenesis have been extensively discussed, both in animal models

and humans. However, whether these injuries can trigger the generation of newborn astrocytes from the

neurogenic niches remains a topic of debate. Neural stem cells in the adult brain are a remnant of

embryonic radial glia. Some researchers suggest that depletion of neurogenesis in the adult brain is a

consequence of astrogliogenesis (Encinas et al., 2011), while others propose astrogliogenesis is part of

a neurorestorative program originating from the neurogenic niches (Bonzano et al., 2018; Sohn et al.,

2015). Despite these differing perspectives, direct and definitive evidence of the generation of newborn

astrocytes from the SVZ or the SGZ has yet to be presented, as well as the potential role of these

astrocytes in aging and pathological conditions. In this study, we confirm that a strong and acute

damage-associated stimulus in the brain, such as an ischemic infarct, prompts astrogliogenesis in the

SVZ. Furthermore, we demonstrate that these astrocytes, which are positive for Thbs4, migrate to the

glial scar and play a role in its modulation, being capable of synthesis and degradation of hyaluronan,

a key component of the ECM at the scar.

Our findings align with previous studies, where cortical injury initiated by photothrombotic/ischemia

was shown to trigger a substantial production of Thbs4-positive astrocytes from the SVZ, a process

modulated by Notch activation (Benner et al., 2013). In this work, the authors suggested a clear role for

Thbs4-positive astrocytes in modulating the ischemic scar. In fact, homozygotic deletion of the Thbs4

gene led to abnormal glial scar formation and increased microvascular hemorrhage following brain

ischemia. Similarly, our results demonstrate that the adult neurogenic niche at the SVZ responds early

after ischemic stroke, degrading local hyaluronan plausibly to modify niche stiffness and facilitate

proliferation and differentiation of NSCs. Further experiments using Thbs4-knockout models will help

confirm this hypothesis.

By labeling NSC via *in vivo* postnatal electroporation, we provided proof of the origin and migration

of newborn Thbs4-positive astrocytes from the SVZ to the ischemic scar in response to the injury.

Notably, Thbs4 is expressed in the newborn astrocytes that migrate to the scar, whereas local reactive

astrocytes at the site of injury express only GFAP. It is likely that SVZ-derived newborn astrocytes

require Thbs4 expression to facilitate their migration to the ischemic area, as Thbs4 deletion has been

shown to impair migration of SVZ-derived cells along the RMS (Girard et al., 2014).

It is noteworthy that while the local, "GFAP-only", reactive astrocytes at the scar are capable of forming the required ECM (Cregg et al., 2014; Wanner et al., 2013), we show that Thbs4-positive (Thbs4/GFAP) astrocytes are also recruited from the SVZ to modulate scar formation. Unlike local reactive astrocytes, which are hastily generated and present both at the core and periphery of the lesion, the newborn astrocytes from the SVZ arrive later to the injury. We demonstrate that these Thbs4-positive astrocytes migrate from the SVZ to the lesion and halt at the lesion border (precisely defined by Nestin staining), establishing residence at the scar but not beyond.

The molecular signals that trigger gliogenesis in the SVZ and recruit newborn astrocytes to the lesion remain unclear. We previously identified adenosine, a byproduct of ATP hydrolysis, as a strong candidate, as it stimulates astrogliogenesis through A1 receptors following OGD *in vitro* and ischemia *in vivo* (Benito-Muñoz et al., 2016). Here, we show that viral-mediated overproduction of HA, the primary component of the glial scar, is sufficient to activate the generation and migration of the Thbs4-positive astrocytes from the SVZ. However, our findings also show that the ischemic infarct, with its proinflammatory (cytokines, chemokines) and proteolytic (MMPs, ROS) signals (Jayaraj et al., 2019), serves as a more potent attractor than the ECM alone.

The glial response from the SVZ we propose in this study can have several functional implications. Thbs4-positive astrocytes generated from the SVZ might act as a second wave of reactivity, supporting local astrocytes in modulating ECM formation. Hyper-reactive astrocytes at the lesion border decrease over time, along with reductions in HA synthesis and CD44 activity, suggesting that the second wave of Thbs4-positive astrocytes from the SVZ could sustain high levels of HA production (Lindwall et al., 2013). Thus, Thbs4-positive astrocytes may be crucial during this delayed temporal window when glial scar remodeling is essential for neurological recovery (Cai et al., 2017; Dzyubenko et al., 2018). However, it's important to note that only about one-third of the astrocytes at the glial scar are derived from the SVZ. This suggests that while SVZ-derived astrocytes may be important for glial scar maintenance, local astrocytes remain the central players in this process.

Based on our observations of both HA production and internalization by Thbs4-positive astrocytes *in*
vivo within the ischemic region and *in vitro* following OGD, we propose a role for these astrocytes in
the modulation, and possibly maintenance, of the ECM at the scar. It is likely that ECM degradation at
the scar is also carried out by professional macrophages, such as microglia, which are known to be
recruited to the scar (Fawcett and Asher, 1999) and have been shown to degrade ECM through
phagocytosis (Nguyen et al., 2020; Soria et al., 2020). Understanding which cells within the scar
environment participate in its modulation, particularly its degradation, holds highly therapeutic value
for promoting regeneration (Dzyubenko et al., 2018; Silver and Miller, 2004). Here, we shed light on
the unknown role of SVZ-derived astrocytes activated by ischemic brain injury and propose them as
critical players in tissue regeneration after brain ischemia.

ACKNOWLEDGEMENTS

The authors thank Juan Carlos Chara, Zara Martínez-Paez, and Laura Escobar for their invaluable
technical support. We thank Laura Simon for proofreading the manuscript, and Abraham Martín,
Mónica Benito-Muñoz and Sol Beccari for training in MCAO in mice. We also thank Vera Gorbunova
(University of Rochester, USA) for providing the cDNA of HAS2 from the naked mole rat. We thank
Marie-Laure Thiolat and Dr. Nathalie Dutheil for AAV production. This study received financial
support from the French government in the framework of the University of Bordeaux's IdEx
"Investments for the Future" program/GPR BRAIN_2030 (B.D.). F.C. acknowledges funding from
BIOEF (BIO21/COV/012), Eukampus, Plan Complementario en Salud (PRTR-C17.I1) and the Basque
Government. F.N.S. was funded by the Spanish Research Agency (IJCI-2017-32114, PID2020-
115896RJ-I00 and RYC2021-032602-I co-funded by NextGeneration EU/PRTR). C.M. acknowledges
funding from the Basque Government (IT1203-19) and CIBERNED. M.A. was supported by a PhD
fellowship from the Basque Government.

AUTHOR CONTRIBUTIONS

M.A. conducted experiments and collected and analyzed data. M.C.T. and H.C. provided the Ai14-Rosa26-CAG-tdTomato$^{flx/flx}$ mice and technical input for electroporation. B.D. provided the AAV-HAS2 and conceived the idea of glial scar simulation. F.C., F.N.S. and C.M. secured funding. C.M. provided infrastructural support and scientific feedback. F.C. and F.N.S. conceived and supervised the study. F.P.C. supervised the study. M.A. and F.N.S. prepared the figures. M.A., F.N.S. and F.C. wrote the paper, with input from all authors.

METHODS

Animals

Male and female wild-type C57BL6/J mice and Ai14-Rosa26-CAG-tdTomato$^{flx/flx}$ transgenic mice bred

at the animal facility of UPV/EHU were used for *in vivo* experiments. Sprague Dawley rat pups from

P4 to P7 were used for *in vitro* experiments. All animals for *in vitro* and *in vivo* studies were handled

in accordance with the European Communities Council Directive (2010/63/EU). All possible efforts

were made to minimize animal suffering and the number of animals used. Animals were maintained

under standard laboratory conditions with food and water *ad libitum*.

Transient MCAO

Brain ischemia was induced in 3-4-month old C57BL/6 mice (approximately 25 g) by occluding 60

minutes of the lenticulo-striate arteries with a 10 mm length of silicone-coated 6-0 monofilament nylon

suture (602223PK10, Doccol Corporation) using a modification of the protocol described by

(Gelderblom et al., 2009). Arteries occlusion generated a specific lesion in the cortex and striatum that

was evaluated at different times after the MCAO (7, 15, 30 or 60 days after occlusion). Ischemic damage

was evaluated in the ipsilateral hemisphere by Cresyl violet staining, Nestin immunofluorescence or

hyaluronan staining. Sham animals, in which arteries were visualized but not disturbed, were used as

an experimental control. Both experimental groups, sham and ischemic, were anesthetized with a mix

of isoflurane and oxygen (induction of anesthesia with 4% isoflurane and maintenance for surgery at

1.5% isoflurane) and maintained under analgesia with buprenorphine 24 hours after ischemia (0.03

mg/Kg body weight i.p. every 12h). Animal wellness was evaluated daily, controlling the body weight

and the motor deficit of the animal by means of neurological scores and pole test. Motor deficit was

assessed in each animal 1h after MCAO and later every 24h. Animals without neurological deficits

were excluded from the study.

5-bromo-2'-deoxyuridine (BrdU) protocol

The total pool of proliferating cells was labelled with three injections of 50 mg/kg 5-bromo-2'-deoxyuridine (BrdU) the day before MCAO protocol. Animals were sacrificed the day after MCAO to analyse acute SVZ proliferation. Chronic BrdU protocol was performed for quiescent NSC identification (Codega et al., 2014). Briefly, 1% of BrdU was administered in the drinking water for 14 days. On day 15, BrdU was removed and animals were subjected to MCAO protocol. Mice were sacrificed and perfused at 3, 7, 15, and 30 days after MCAO, although only animals with a temporal window of 30 days between MCAO and perfusion were used for statistical analysis. Activated cells were labelled by i.p. injection of 50 mg/Kg 5-iodo-2′-deoxyuridine (IdU) 24 hours before euthanasia.

Electroporation

Ai14-Rosa26-CAG-tdTomato$^{flx/flx}$ transgenic mice and pCAGGS-CRE plasmid were kindly provided for Prof. Harold Cremer and Marie-Catherine Tiveron from Institut de Biologie du Développement de Marseille (IBDM). Postnatal electroporation was performed as described previously (Platel et al., 2019). Mice pups from P0-P2 were anesthetized by hypothermia. pCAGGS-CRE plasmid was used at a final concentration of 2.5 µg/ul and 2µl was injected in the lateral ventricle by expiratory pressure using a glass micropipette. Following injection, pup brains were placed between electrodes targeted the dorsal SVZ and subjected to five 95V electrical pulses (50 ms, 950 ms intervals). BTX ECM399 (Thermo Fisher Scientific) electroporator was used with 7 mm tweezer electrodes. Electroporated pups were reanimated at 37ºC before returning to the mother.

In order to analyze the Thbs4 positive cells in the SVZ, we subcloned the eGFP gene (from a pCMV-eGFP N1) into a PGL3-basic backbone containing mouse Thbs4 promoter, kindly provided by Stenina-Adognravi lab (Muppala et al., 2017). pThbs4-GFP was also electroporated in P0-P2 mice, and euthanized 1-day post-electroporation to corroborate Thbs4 expression in the SVZ at postnatal stages.

AAV vector production

AAV2/9-CAG-HAS2-3xFlag vector was produced by polyethylenimine (PEI) mediated triple transfection of low passage HEK-293 T/17 cells (ATCC; cat number CRL-11268). The respective AAV expression plasmid expression plasmid pNMRHAS2_N1 (provided by Dr Vera Gorbunova) was cotransfected with the adeno helper pAd Delta F6 plasmid (Penn Vector Core, cat # PL-F-PVADF6) and AAV Rep Cap pAAV2/9 plasmid (Penn Vector Core, cat # PL-TPV008). AAV vector was purified as previously described (Bourdenx et al., 2015). Cells are harvested 72 h post-transfection, resuspended in lysis buffer (150 mM NaCl, 50 mM Tris–HCl pH 8.5) and lysed by 3 freeze-thaw cycles (37°C/-80°C). The cell lysate is treated with 150 units/ml Benzonase (Sigma, St Louis, MO) for 1 h at 37°C, and the crude lysate is clarified by centrifugation. Vectors are purified by iodixanol step gradient centrifugation and concentrated and buffer-exchanged into Lactated Ringer's solution (Baxter, Deerfield, IL) using vivaspin20 100 kDa cut-off concentrator (Sartorius Stedim, Goettingen, Germany). Titrations were performed at the transcriptome core facility (Neurocentre Magendie, INSERM U862, Bordeaux, France). The genome-containing particle (gcp) titer was determined at a concentration of 1.10^{13} gcp/ml by quantitative real-time PCR using the Light Cycler 480 SYBR green master mix (Roche, cat # 04887352001) with primers specific for the AAV2 ITRs (fwd 50-GGAACCCCTAGTGATG GAGTT-3'; rev 50-CGGCCTCAGTGAGCGA-30) on a Light Cycler 480 instrument. The purity assessment of vector stocks was estimated by loading 10 µl of vector stock on 10 % SDS acrylamide gels; total proteins were visualized using the Krypton Infrared Protein Stain according to the manufacturer's instructions (Life Technologies).

In vivo AAV-HAS2 injections

Mice were anesthetized with isoflurane and fixed to a stereotaxic frame connected with a gas mask. We injected 1µl of adeno-associated virus (AAV2/9) CAG-HAS2-3xFlag (1x10^13 gc/µl) in the contralateral striatum (AP: +1.0; ML: -1.8; DV: -3.2) to simulate scar tissue hyaluronan. After two weeks, the ischemic group was subjected to the MCAO model in the ipsilateral hemisphere and perfused

30 days later. Only slices with representative lesions in both hemispheres were used for image and

statistical analysis.

Immunofluorescence

Mice were anesthetized with an intraperitoneal injection of pentobarbital (100 mg/kg). Brain tissues

were fixed by intracardiac perfusion using 4% paraformaldehyde. Fixed brains were sliced at 40 μm of

thickness using the Leica VT1200S vibratome (Leica Microsystems) and subjected to

immunofluorescence. For standard immunofluorescence protocol, tissue slices were permeabilized with

0.1% Triton and non-specific epitopes blocked with 1% Bovine Serum Albumin and 10% Donkey

serum in PBS. Primary antibodies (Supplementary Table 1) were incubated at different concentrations

(see Table 1 for specification) overnight at 4ºC and then washed three times with 0.1% Triton in PBS.

Secondary conjugated antibodies were incubated for 1 h at room temperature. After three washes with

0.1% Triton in PBS, cells were stained for 10 min with DAPI and further washed with PBS. Finally,

coverslips were mounted with Fluoromount (SouthernBiotech) and analyzed by fluorescence using the

confocal microscope Leica TCS SP8. For hyaluronic acid labeling, endogenous biotins were pre-

blocked using Streptavidin/Biotin blocking solution (Vector Labs, #SP-2002) following the

manufacturer's instructions and further blocked with PBS 1X 0.1% Saponin 1% BSA for one hour at

room temperature. Primary antibodies and biotinylated Hyaluronic Acid Binding Protein (HABP,

#385911 Millipore) were incubated for 72 hours at 4°C or 24 hours at room temperature. After three

washes with PBS, 1X 0.1% Saponin, secondary antibody and DAPI diluted in the same blocking buffer

were incubated for one hour at room temperature. Slices were mounted with Mowiol (Millipore

#475904) in superfrost slides (Thermo Fisher Scientific) and #1.5 coverslips.

Imaging and image analysis

Confocal images were acquired in a Leica TCS SP8 with 20x (0.8 NA) and 63x objectives (1.4 NA).

Images were taken with a pixel size of 90 nm. Widefield imaging was performed in a Zeiss AxioVision

fluorescence microscope and a 3DHISTECH panoramic MIDI II digital slide scanner. Settings were

adjusted for each individual experiment and kept for all conditions. Image processing was performed

using FIJI (Schindelin et al., 2012), CaseViewer and Hyugens (SVI, The Netherlands) software. For

Thbs4, Nestin and DCX immunofluorescence, wholemount SVZ preparations and AAV-HAS2

analysis, tile-scan was done using the navigator plug-in in the LasX software. For Nestin

immunofluorescence, images were deconvoluted using the Hyugens software.

SVZ images were reconstructed using the tile-scan tool (LasX software, Leica). SVZ region of interest

(ROI) was delimited manually and expression of each marker was measured in the SVZ ROI.

Quantification of all markers was done manually, except for HABP quantification, which was done

with custom FIJI scripts for in vitro (https://github.com/MariArdaya/Hyaluronan_v2) and in vivo

analysis (https://github.com/SoriaFN/Tools). The HABP signal was segmented with an automatic

threshold and colocalized with other markers such as CD44. This, combined with cell segmentation,

based on the Thbs4 mask, allowed for quantification of HABP area and number of spots inside cells,

outside cells or at the membrane (with a band ROI of 1 μm). The Analyze Skeleton and Fractal count

plugins were used on the skeletonized segmented HABP to quantify the longest shortest path and the

fractal dimension, respectively, to reveal the length and interconnectivity of the matrix network (Casey

et al., 2024; Soria et al., 2020).

Western blot

The SVZ was freshly dissected on ice and minced using a micro grinder. Total protein was extracted

after tissue lysis with RIPA buffer (Thermo Fisher Scientific #89900) and Protease and Phosphatase

Inhibitor Cocktail 100X (Thermo Fisher Scientific). Samples were lysed in ice for 10 minutes and

sonicated using an Ultrasonic Processor (Hielscher Ultrasound Technology). Sonicated tissues were

centrifuged at 1200 rpm for 10 min and 4°C to remove insoluble fragments. Protein concentration was

measured by chemioluminescence using RC DC™ Protein Assay (Bio-Rad). Forty micrograms of

protein samples were denatured in reducing loading buffer containing β-mercaptoethanol (M3148,

Sigma) and 0.0002% bromophenol blue at 95°C for 5 minutes and separated in pre-casted 12% or 7.5%

Tris-Glycine polyacrylamide gel using the Criterion cell system (Biorad). Electrophoresis was

conducted at 80 V in a Tris-Glycine buffer (25 mM Tris, 192 mM glycine, 0.1% SDS in dH2O, pH 8.3).

After complete separation, proteins were transferred to a nitrocellulose membrane (Biorad) using Trans-

Blot® TurboTM Transfer System (Bio-Rad). Nitrocellulose membranes were incubated in blocking

solution (Tris-Buffer Solution (TBS), 0.1% Tween-20 and 5% BSA) for one hour at room temperature.

Primary antibodies were incubated overnight at 4°C. Horseradish peroxidase-conjugated secondary

antibodies were incubated in blocking buffer for one hour at room temperature. Immunodetection was

performed by electrochemical luminescence in a Chemidoc MP (Biorad) and the intensity of the bands

was quantified using ImageLab (Bio-Rad) software corrected by Gaussian curves.

qRT-PCR

Fresh SVZ tissue from ipsilateral hemisphere was isolated as previously described from sham, 15 and

30 days after MCAO animals to analyze the expression of several genes (Supplementary Table 2). After

mechanical trituration, RNA was extracted from the tissue after mechanical trituration following the

manufacturer´s protocol (NZYTECH Total RNA Isolation kit protocol (NZYTECH, MB13402). RNA

concentration was measured by spectrophotometry using the Nanodrop 2000c spectrophotometer

(Thermo-Scientific, USA). Real time polymerase chain reaction (RT-PCR) from 10 ng of RNA was

performed using the One-step NZYTECH RT-qPCR Green kit according to manufacturer´s protocol

(NZYTECH, MB34302). RT-PCR was normalized using the housekeeping genes hypoxanthine

phosphoribosyl-transferase 1 (HPRT1, Qiagen QuantiTect Primer Assay, Barcelona, Spain). Specificity

of each amplification was confirmed by melting curve analysis. PCR reactions were performed in a

CFX96 Detection System (BioRad, Madrid, Spain). Semi-quantification was performed using the 2-

ΔΔCt algorithm. Unsupervised hierarchical distance matrix and distance matrix map was developed

using Orange3 software (Python v3.0).

Statistical analysis

Statistical analysis was done using GraphPad Prism software (version 8.0; GraphPad software). Data

are presented as mean ± SEM unless specified otherwise. Statistical analysis was performed using

unpaired Student´s t-test for independent conditions and paired Student´s t-test for dependent

conditions. In addition, one-way and two-way ANOVA were also performed to compare more than 2

groups among them. Tukey and Dunnett post-hoc tests were used. For HABP skeleton analysis,

Kruskal-Wallis analysis was performed to compare cumulative frequency curves in more than 2 groups.

P value was considered significant at $p < 0.05$. Sample number (*n*), *p* values and statistical test are

indicated in figure legends.

REFERENCES

Adams KL, Gallo V. 2018. The diversity and disparity of the glial scar. *Nat Neurosci* **21**:9–15.
doi:10.1038/s41593-017-0033-9
Anderson MA, Burda JE, Ren Y, Ao Y, O'Shea TM, Kawaguchi R, Coppola G, Khakh BS, Deming TJ,
Sofroniew MV. 2016. Astrocyte scar formation aids central nervous system axon
regeneration. *Nature* **532**:195–200. doi:10.1038/nature17623
Arvidsson A, Collin T, Kirik D, Kokaia Z, Lindvall O. 2002. Neuronal replacement from endogenous
precursors in the adult brain after stroke. *Nat Med* **8**:963–970. doi:10.1038/nm747
Basak O, Krieger TG, Muraro MJ, Wiebrands K, Stange DE, Frias-Aldeguer J, Rivron NC, van de
Wetering M, van Es JH, van Oudenaarden A, Simons BD, Clevers H. 2018. Troy+ brain stem
cells cycle through quiescence and regulate their number by sensing niche occupancy. *Proc*
Natl Acad Sci U S A **115**:E610–E619. doi:10.1073/pnas.1715911114
Beckervordersandforth R, Tripathi P, Ninkovic J, Bayam E, Lepier A, Stempfhuber B, Kirchhoff F,
Hirrlinger J, Haslinger A, Lie DC, Beckers J, Yoder B, Irmler M, Götz M. 2010. In vivo fate
mapping and expression analysis reveals molecular hallmarks of prospectively isolated adult
neural stem cells. *Cell Stem Cell* **7**:744–758. doi:10.1016/j.stem.2010.11.017
Benito-Muñoz M, Matute C, Cavaliere F. 2016. Adenosine A1 receptor inhibits postnatal
neurogenesis and sustains astrogliogenesis from the subventricular zone. *Glia* **64**:1465–
1478. doi:10.1002/glia.23010
Benner EJ, Luciano D, Jo R, Abdi K, Paez-Gonzalez P, Sheng H, Warner DS, Liu C, Eroglu C, Kuo CT.
2013. Protective astrogenesis from the SVZ niche after injury is controlled by Notch
modulator Thbs4. *Nature* **497**:369–373. doi:10.1038/nature12069

Bond AM, Ming G-L, Song H. 2015. Adult Mammalian Neural Stem Cells and Neurogenesis: Five
Decades Later. *Cell Stem Cell* **17**:385–395. doi:10.1016/j.stem.2015.09.003
Bonzano S, Crisci I, Podlesny-Drabiniok A, Rolando C, Krezel W, Studer M, De Marchis S. 2018.
Neuron-Astroglia Cell Fate Decision in the Adult Mouse Hippocampal Neurogenic Niche Is
Cell-Intrinsically Controlled by COUP-TFI In Vivo. *Cell Rep* **24**:329–341.
doi:10.1016/j.celrep.2018.06.044
Bourdenx M, Dovero S, Engeln M, Bido S, Bastide MF, Dutheil N, Vollenweider I, Baud L, Piron C,
Grouthier V, Boraud T, Porras G, Li Q, Baekelandt V, Scheller D, Michel A, Fernagut P-O,
Georges F, Courtine G, Bezard E, Dehay B. 2015. Lack of additive role of ageing in
nigrostriatal neurodegeneration triggered by α-synuclein overexpression. *Acta Neuropathol*
Commun **3**:46. doi:10.1186/s40478-015-0222-2
Bush TG, Puvanachandra N, Horner CH, Polito A, Ostenfeld T, Svendsen CN, Mucke L, Johnson MH,
Sofroniew MV. 1999. Leukocyte infiltration, neuronal degeneration, and neurite outgrowth
after ablation of scar-forming, reactive astrocytes in adult transgenic mice. *Neuron* **23**:297–
308. doi:10.1016/s0896-6273(00)80781-3
Cai H, Ma Y, Jiang L, Mu Z, Jiang Z, Chen X, Wang Y, Yang G-Y, Zhang Z. 2017. Hypoxia Response
Element-Regulated MMP-9 Promotes Neurological Recovery via Glial Scar Degradation and
Angiogenesis in Delayed Stroke. *Molecular Therapy* **25**:1448–1459.
doi:10.1016/j.ymthe.2017.03.020
Casey DT, Lahue KG, Mori V, Herrmann J, Hall JK, Suki B, Janssen-Heininger YMW, Bates JHT. 2024.
Local fractal dimension of collagen detects increased spatial complexity in fibrosis.
Histochem Cell Biol **161**:29–42. doi:10.1007/s00418-023-02248-8
Cavaliere F, Dinkel K, Reymann K. 2006. The subventricular zone releases factors which can be
protective in oxygen/glucose deprivation-induced cortical damage: an organotypic study.
Exp Neurol **201**:66–74. doi:10.1016/j.expneurol.2006.03.020
Ceanga M, Dahab M, Witte OW, Keiner S. 2021. Adult Neurogenesis and Stroke: A Tale of Two
Neurogenic Niches. *Frontiers in Neuroscience* **15**.
Cebrian-Silla A, Nascimento MA, Redmond SA, Mansky B, Wu D, Obernier K, Romero Rodriguez R,
Gonzalez-Granero S, García-Verdugo JM, Lim DA, Álvarez-Buylla A. 2021. Single-cell analysis
of the ventricular-subventricular zone reveals signatures of dorsal and ventral adult
neurogenesis. *eLife* **10**:e67436. doi:10.7554/eLife.67436
Chaker Z, Codega P, Doetsch F. 2016. A mosaic world: puzzles revealed by adult neural stem cell
heterogeneity. *WIREs Developmental Biology* **5**:640–658. doi:10.1002/wdev.248
Codega P, Silva-Vargas V, Paul A, Maldonado-Soto AR, DeLeo AM, Pastrana E, Doetsch F. 2014.
Prospective Identification and Purification of Quiescent Adult Neural Stem Cells from Their In
Vivo Niche. *Neuron* **82**:545–559. doi:10.1016/j.neuron.2014.02.039
Cregg JM, DePaul MA, Filous AR, Lang BT, Tran A, Silver J. 2014. Functional regeneration beyond the
glial scar. *Exp Neurol* **253**:197–207. doi:10.1016/j.expneurol.2013.12.024
Doetsch F, Caillé I, Lim DA, García-Verdugo JM, Alvarez-Buylla A. 1999. Subventricular zone
astrocytes are neural stem cells in the adult mammalian brain. *Cell* **97**:703–716.
doi:10.1016/s0092-8674(00)80783-7
Dzwonek J, Wilczynski GM. 2015. CD44: molecular interactions, signaling and functions in the
nervous system. *Front Cell Neurosci* **9**:175. doi:10.3389/fncel.2015.00175
Dzyubenko E, Manrique-Castano D, Kleinschnitz C, Faissner A, Hermann DM. 2018. Role of immune
responses for extracellular matrix remodeling in the ischemic brain. *Ther Adv Neurol Disord*
11:1756286418818092. doi:10.1177/1756286418818092
Encinas JM, Michurina TV, Peunova N, Park J-H, Tordo J, Peterson DA, Fishell G, Koulakov A,
Enikolopov G. 2011. Division-coupled astrocytic differentiation and age-related depletion of
neural stem cells in the adult hippocampus. *Cell Stem Cell* **8**:566–579.
doi:10.1016/j.stem.2011.03.010

Faulkner JR, Herrmann JE, Woo MJ, Tansey KE, Doan NB, Sofroniew MV. 2004. Reactive astrocytes
protect tissue and preserve function after spinal cord injury. *J Neurosci* **24**:2143–2155.
doi:10.1523/JNEUROSCI.3547-03.2004
Fawcett JW. 2015. The extracellular matrix in plasticity and regeneration after CNS injury and
neurodegenerative disease. *Prog Brain Res* **218**:213–226. doi:10.1016/bs.pbr.2015.02.001
Fawcett JW, Asher RA. 1999. The glial scar and central nervous system repair. *Brain Res Bull* **49**:377–
391. doi:10.1016/s0361-9230(99)00072-6
Frisén J, Johansson CB, Török C, Risling M, Lendahl U. 1995. Rapid, widespread, and longlasting
induction of nestin contributes to the generation of glial scar tissue after CNS injury. *Journal*
of Cell Biology **131**:453–464. doi:10.1083/jcb.131.2.453
Gelderblom M, Leypoldt F, Steinbach K, Behrens D, Choe C-U, Siler DA, Arumugam TV, Orthey E,
Gerloff C, Tolosa E, Magnus T. 2009. Temporal and spatial dynamics of cerebral immune cell
accumulation in stroke. *Stroke* **40**:1849–1857. doi:10.1161/STROKEAHA.108.534503
Girard F, Eichenberger S, Celio MR. 2014. Thrombospondin 4 deficiency in mouse impairs neuronal
migration in the early postnatal and adult brain. *Molecular and Cellular Neuroscience*
61:176–186. doi:10.1016/j.mcn.2014.06.010
Jayaraj RL, Azimullah S, Beiram R, Jalal FY, Rosenberg GA. 2019. Neuroinflammation: friend and foe
for ischemic stroke. *J Neuroinflammation* **16**:142. doi:10.1186/s12974-019-1516-2
Kjell J, Fischer-Sternjak J, Thompson AJ, Friess C, Sticco MJ, Salinas F, Cox J, Martinelli DC, Ninkovic J,
Franze K, Schiller HB, Götz M. 2020. Defining the Adult Neural Stem Cell Niche Proteome
Identifies Key Regulators of Adult Neurogenesis. *Cell Stem Cell* **26**:277-293.e8. doi:10/ghcp5v
Kralic JE, Ledergerber DA, Fritschy J-M. 2005. Disruption of the neurogenic potential of the dentate
gyrus in a mouse model of temporal lobe epilepsy with focal seizures. *European Journal of*
Neuroscience **22**:1916–1927. doi:10.1111/j.1460-9568.2005.04386.x
Lau LW, Cua R, Keough MB, Haylock-Jacobs S, Yong VW. 2013. Pathophysiology of the brain
extracellular matrix: a new target for remyelination. *Nat Rev Neurosci* **14**:722–729.
doi:10/f5b6w9
Laug D, Huang T-W, Huerta NAB, Huang AY-S, Sardar D, Ortiz-Guzman J, Carlson JC, Arenkiel BR, Kuo
CT, Mohila CA, Glasgow SM, Lee HK, Deneen B. 2019. Nuclear factor I-A regulates diverse
reactive astrocyte responses after CNS injury. *J Clin Invest* **129**:4408–4418.
doi:10.1172/JCI127492
Li L, Harms KM, Ventura PB, Lagace DC, Eisch AJ, Cunningham LA. 2010. Focal cerebral ischemia
induces a multilineage cytogenic response from adult subventricular zone that is
predominantly gliogenic. *Glia* **58**:1610–1619. doi:10.1002/glia.21033
Lindwall C, Olsson M, Osman AM, Kuhn HG, Curtis MA. 2013. Selective expression of hyaluronan and
receptor for hyaluronan mediated motility (Rhamm) in the adult mouse subventricular zone
and rostral migratory stream and in ischemic cortex. *Brain Res* **1503**:62–77.
doi:10.1016/j.brainres.2013.01.045
Liu XS, Zhang ZG, Zhang RL, Gregg S, Morris DC, Wang Y, Chopp M. 2007. Stroke induces gene profile
changes associated with neurogenesis and angiogenesis in adult subventricular zone
progenitor cells. *J Cereb Blood Flow Metab* **27**:564–574. doi:10.1038/sj.jcbfm.9600371
Llorens-Bobadilla E, Zhao S, Baser A, Saiz-Castro G, Zwadlo K, Martin-Villalba A. 2015. Single-Cell
Transcriptomics Reveals a Population of Dormant Neural Stem Cells that Become Activated
upon Brain Injury. *Cell Stem Cell* **17**:329–340. doi:10.1016/j.stem.2015.07.002
Long KR, Huttner WB. 2019. How the extracellular matrix shapes neural development. *Open Biol*
9:180216. doi:10/ghn8hs
Luskin MB. 1993. Restricted proliferation and migration of postnatally generated neurons derived
from the forebrain subventricular zone. *Neuron* **11**:173–189. doi:10.1016/0896-
6273(93)90281-u

Muppala S, Xiao R, Krukovets I, Verbovetsky D, Yendamuri R, Habib N, Raman P, Plow E, Stenina-Adognravi O. 2017. Thrombospondin-4 mediates TGF-β-induced angiogenesis. *Oncogene* **36**:5189–5198. doi:10.1038/onc.2017.140

Nguyen PT, Dorman LC, Pan S, Vainchtein ID, Han RT, Nakao-Inoue H, Taloma SE, Barron JJ, Molofsky AB, Kheirbek MA, Molofsky AV. 2020. Microglial Remodeling of the Extracellular Matrix Promotes Synapse Plasticity. *Cell* **182**:388-403.e15. doi:10/gg5cj3

Platel J-C, Angelova A, Bugeon S, Wallace J, Ganay T, Chudotvorova I, Deloulme J-C, Béclin C, Tiveron M-C, Coré N, Murthy VN, Cremer H. 2019. Neuronal integration in the adult mouse olfactory bulb is a non-selective addition process. *eLife* **8**:e44830. doi:10.7554/eLife.44830

Pous L, Deshpande SS, Nath S, Mezey S, Malik SC, Schildge S, Bohrer C, Topp K, Pfeifer D, Fernández-Klett F, Doostkam S, Galanakis DK, Taylor V, Akassoglou K, Schachtrup C. 2020. Fibrinogen induces neural stem cell differentiation into astrocytes in the subventricular zone via BMP signaling. *Nat Commun* **11**:630. doi:10.1038/s41467-020-14466-y

Preston M, Sherman LS. 2011. Neural Stem Cell Niches: Critical Roles for the Hyaluronan-Based Extracellular Matrix in Neural Stem Cell Proliferation and Differentiation. *Front Biosci (Schol Ed)* **3**:1165–1179.

Schindelin J, Arganda-Carreras I, Frise E, Kaynig V, Longair M, Pietzsch T, Preibisch S, Rueden C, Saalfeld S, Schmid B, Tinevez J-Y, White DJ, Hartenstein V, Eliceiri K, Tomancak P, Cardona A. 2012. Fiji: an open-source platform for biological-image analysis. *Nat Methods* **9**:676–682. doi:10/f34d7c

Segel M, Neumann B, Hill MFE, Weber IP, Viscomi C, Zhao C, Young A, Agley CC, Thompson AJ, Gonzalez GA, Sharma A, Holmqvist S, Rowitch DH, Franze K, Franklin RJM, Chalut KJ. 2019. Niche stiffness underlies the ageing of central nervous system progenitor cells. *Nature* **573**:130–134. doi:10/gf6j83

Sherman LS, Matsumoto S, Su W, Srivastava T, Back SA. 2015. Hyaluronan Synthesis, Catabolism, and Signaling in Neurodegenerative Diseases. *International Journal of Cell Biology* **2015**:e368584. doi:10/gccgm9

Silver J, Miller JH. 2004. Regeneration beyond the glial scar. *Nat Rev Neurosci* **5**:146–156. doi:10.1038/nrn1326

Sohn J, Orosco L, Guo F, Chung S-H, Bannerman P, Mills Ko E, Zarbalis K, Deng W, Pleasure D. 2015. The subventricular zone continues to generate corpus callosum and rostral migratory stream astroglia in normal adult mice. *J Neurosci* **35**:3756–3763. doi:10.1523/JNEUROSCI.3454-14.2015

Soria FN, Paviolo C, Doudnikoff E, Arotcarena M-L, Lee A, Danné N, Mandal AK, Gosset P, Dehay B, Groc L, Cognet L, Bezard E. 2020. Synucleinopathy alters nanoscale organization and diffusion in the brain extracellular space through hyaluronan remodeling. *Nat Commun* **11**:3440. doi:10.1038/s41467-020-17328-9

Sundholm-Peters NL, Yang HKC, Goings GE, Walker AS, Szele FG. 2005. Subventricular Zone Neuroblasts Emigrate Toward Cortical Lesions. *Journal of Neuropathology & Experimental Neurology* **64**:1089–1100. doi:10.1097/01.jnen.0000190066.13312.8f

Tian X, Azpurua J, Hine C, Vaidya A, Myakishev-Rempel M, Ablaeva J, Mao Z, Nevo E, Gorbunova V, Seluanov A. 2013. High-molecular-mass hyaluronan mediates the cancer resistance of the naked mole rat. *Nature* **499**:346–349. doi:10/ghcp25

Wanner IB, Anderson MA, Song B, Levine J, Fernandez A, Gray-Thompson Z, Ao Y, Sofroniew MV. 2013. Glial scar borders are formed by newly proliferated, elongated astrocytes that interact to corral inflammatory and fibrotic cells via STAT3-dependent mechanisms after spinal cord injury. *J Neurosci* **33**:12870–12886. doi:10.1523/JNEUROSCI.2121-13.2013

Yamashita T, Ninomiya M, Hernández Acosta P, García-Verdugo JM, Sunabori T, Sakaguchi M, Adachi K, Kojima T, Hirota Y, Kawase T, Araki N, Abe K, Okano H, Sawamoto K. 2006. Subventricular zone-derived neuroblasts migrate and differentiate into mature neurons in the post-stroke adult striatum. *J Neurosci* **26**:6627–6636. doi:10.1523/JNEUROSCI.0149-06.2006

Zhang Z, Tian X, Lu JY, Boit K, Ablaeva J, Zakusilo FT, Emmrich S, Firsanov D, Rydkina E, Biashad SA, Lu
Q, Tyshkovskiy A, Gladyshev VN, Horvath S, Seluanov A, Gorbunova V. 2023. Increased
hyaluronan by naked mole-rat Has2 improves healthspan in mice. *Nature*.
doi:10.1038/s41586-023-06463-0
Zimmermann DR, Dours-Zimmermann MT. 2008. Extracellular matrix of the central nervous system:
from neglect to challenge. *Histochem Cell Biol* **130**:635–653. doi:10.1007/s00418-008-0485-9
718

Figure 1 – Thbs4 labels type-B cells in the SVZ neurogenic niche. (a) Schematic drawing of the brain showing Thbs4 labeling throughout the mouse brain. Thbs4 was primarily found in the SVZ and RMS of the adult mouse telencephalon. Arrows indicate non-specific Thbs4 staining in cerebellum and ventral tegmental area. (b) Thbs4 positive cells in the SVZ displayed various morphologies. (c) Proportion of Thbs4-positive cells relative to the total NSCs population in the SVZ. (d) Classification of Thbs4-positive cells in the SVZ based to their NSC morphology. (e) Electroporation of pThbs4-GFP at P1 revealed Thbs4 expression in the postnatal dorsal SVZ. (f) Quantification of Thbs4-reporter expression showing the proportion of NSC expressing Thbs4 postnatally in the dorsal SVZ. (g) Thbs4-GFP reporter expression was upregulated in the rostral SVZ. BV: Blood vessel; LV: Lateral ventricle. Scale bar = 10 µm (b) and 50 µm (e). Bars represent mean ± SEM. $*p < 0.05$ and $**p < 0.01$, Tukey post hoc test (after one-way ANOVA was significant at $p < 0.05$).

Figure 2 – Thbs4 expression is upregulated in the SVZ after brain ischemia. (a) Representative images of Thbs4 (green), Nestin (red) and DCX (magenta) in the SVZ under physiological conditions (left) and 15 days post-injury (dpi) (right). (b, c) The number of actively proliferating NSCs and neuroblast decreases at 7 and 15 dpi in the SVZ, as shown by reduced Nestin (b) and DCX (c) levels, respectively. (d) Thbs4 expression increases in the SVZ over time following ischemic injury. (e) Representative western blot images of Nestin and Thbs4 in sham and 15 dpi SVZ. (f, g) Quantification of mean gray value (MGV) from (e), normalized to GAPDH, shows an increased in Thbs4 (f) and a decrease in Nestin (g) in the 15 dpi SVZ. Scale bar = 100 µm in (a). n = 6 (b, c, d) and 3 (e, f, g) per condition. Bars represent mean ± SEM. *$p < 0.05$ and **$p < 0.01$; two-tail Student´s t test (sham vs. 15ddMCAO) and Tukey post-hoc test (after one-way ANOVA was significant at $p < 0.05$). See also Supplementary Figures 1 and 2.

Figure 3 – Thbs4-positive cells derive from B-type NSCs in the SVZ after brain ischemia. (a) Experimental design:

chronic BrdU (1% in water) treatment was administered for two weeks. BrdU labeling persisted only in B-type NSCs 30 days

after treatment. Animals were injected with IdU (50 mg/kg) three times the day before euthanasia to label proliferating cells.

(b, c) BrdU (b) and IdU (c) positive cells in the SVZ. Only BrdU-positive cells showed a significant increase at 30 dpi. (d)

Representative confocal images showing Thbs4, GFAP and BrdU levels in sham (top) and 30 dpi mice (bottom). (e, f)

Thbs4/GFAP/BrdU positive cells (slow proliferative cells) increased in the SVZ at 30 dpi (e), while no changes were observed

in Thbs4/GFAP/IdU positive cells (fast proliferative cells). Scale bar = 10 µm (d). n = 4 (sham) and 3 (MCAO). Bars represent

mean ± SEM. *$p < 0.05$ by two-tail Student´s t-test (sham vs. 30dd MCAO). See also Supplementary Figure 3.

Figure 4 – Ischemia-induced Thbs4 astrocytes migrate from the SVZ to ischemic areas. (a) Experimental design: Ai14, ROSA26-CAG-tdTomato transgenic mice were electroporated in the dorsal SVZ at P1. The pCAGGSx-CRE plasmid induced tdTomato (tdTOM) expression in dorsal NSCs. MCAO was performed 3-4 months later in the same electroporated hemisphere. Mice were analyzed at 7, 30, and 60 dpi. (b) Quantification of tdTOM-positive cells shows a decrease in the dorsal SVZ following brain ischemia. (c) Representative images of tdTOM-positive cells in sham (top) and 60 dpi SVZ (bottom). (d) Quantification of tdTOM-positive cells in the whole brain shows a gradual increase outside the SVZ after brain ischemia. (e) Representative images of tdTOM expression in sham (left) and 60 dpi mice (right). (f) Representative image of Thbs4/tdTOM-positive cells in the dorsal SVZ. (g) Quantification shows an increase in Thbs4/tdTOM-positive cells in the SVZ at 30 and 60 dpi. (h) The proportion of Thbs4-positive cells within the tdTOM-positive pool increases in the SVZ at 30 and 60 dpi. (i) Around 50% of tdTOM-positive cells expressed Thbs4 in the infarcted tissue at 60 dpi. (j) Representative image of Thbs4 and tdTOM expression near the damaged area. Scale bar = 50 µm (c), 10 µm (f), and 100 µm (g). n = 6 animals per condition. Bars represent mean ± SEM. *$p < 0.05$, **$p < 0.01$, and ***$p < 0.001$ by Chi-square test (h) and Tukey post-hoc test (after one-way ANOVA was significant at $p < 0.05$). See also Supplementary Figure 4.

Figure 5 – Ischemia-induced Thbs4 astrocytes accumulate at the glial scar. (a) The glial scar could be observed at 30 dpi using Nestin immunofluorescence. (b) Classical TTC technique underestimated ischemic regions compared with Nestin immunofluorescence. (c) Representative images of TTC staining in infarcted brains. (d) Representative confocal images showing the time course of Nestin expression at 3, 7, 15 and 30 dpi. (e) The Nestin-positive glial scar was established by 15 dpi, while Thbs4 astrocytes arrived at the glial scar later, by 30 dpi. (f) Fluorescence profile of Thbs4, GFAP and Nestin marker from the core to penumbra areas shown Thbs4 was mainly located at the borders of the Nestin-positive glial scar. (g) Representative images of Thbs4 (green), GFAP (white) and Nestin (red) markers in sham (top) and 30 dpi mice (bottom). Scale bar = 50 µm (g). n = 3 animals per condition. Bars/lines represent mean ± SEM. *$p < 0.05$ and **$p < 0.01$ by two-tail Student´s t test (TTC vs. Nestin) and by Dunnett posthoc test (after two-way ANOVA was significant at $p < 0.05$). See also Supplementary Figure 5.

Figure 6 – Thbs4-positive astrocytes produce hyaluronan at the glial scar after MCAO. (a) Representative images of hyaluronic acid (HA) labeled with HABP in the contralateral (1) and ipsilateral hemisphere (2). Only perineuronal nets are clearly visible in the healthy tissue, while a dense interstitial matrix is observed in ischemic areas. (b) Cumulative distribution of skeletonized HA signal in sham and infarcted regions of 7, 15 and 30 dpi mice (p < 0.0001). (c) Quantification of fractal dimension showed increased extracellular matrix interconnectivity in the infarcted cortex and striatum (Ctx & Str) at 30 dpi. (d) Representative confocal images of Thbs4 (green) and HABP (red) in the cortex and striatum of sham and 30 dpi mice. Note the HA labelling on the surface of Thbs4-positive cells. (e) Fluorescence intensity profile of Thbs4, GFAP and HABP markers from the lesion core to penumbra areas. Thbs4 expression was mainly restricted to the borders of the HA-positive glial scar at 30 dpi. (f) Representative confocal images of Thbs4 and HABP markers in sham and 30 dpi mice (top). Examples of HA (red) occupying the membrane mask of Thbs4-positive astrocytes in sham and 30 dpi mice (bottom), showing increased HA synthesis after MCAO. (g) Quantification of membrane-bound HA showed increased HA along Thbs4-positive membrane after brain ischemia (corpus callosum: pink; infarcted cortex: blue; infarcted striatum: green). Scale bar = 10 µm (e). n = 6 animals per condition. *p < 0.05 and ****p < 0.0001 by Krustal Wallis test and two-way ANOVA (significant at *p* < 0.05). See also Supplementary Figures 6 and 7.

Figure 7 – Hyaluronan accumulation is sufficient but not exclusive to recruit Thbs4 astrocytes. (a) Representative images of HA accumulation caused by viral-mediated overexpression of HAS2 from naked mole rat. (b) Experimental design: HAS2 viral injections were performed 2 weeks before MCAO. Animals were sacrificed 30 days after MCAO. We used standard GFP virus as control. Two groups of HAS2-injected mice were studied, with only one subjected to MCAO. (c) FLAG-positive cells (HAS2 virus tag) did not show changes in either HAS2 control (HAS2Control) or HAS2-MCAO group (contralateral hemisphere). (d, e, f) Representative images of HABP, GFP and Thbs4 markers in GFP control virus (d), HAS2 control virus (e) and HAS2 MCAO group (f). (g) Thbs4 expression increased after HAS2 viral-mediated overexpression. However, Thbs4 was upregulated in the ischemic hemisphere when MCAO was performed together with the HAS2 viral infection (MCAO ipsi). GFP control virus did not induce increase of Thbs4. (h) HABP was used as a control for HA accumulation. Thbs4 astrocytes in HAS2 MCAO group increased in the ischemic hemisphere even though HA accumulation in both hemispheres (HAS2 and MCAO hemisphere), suggesting additional ischemia-induced signals recruit Thbs4 astrocytes to infarcted areas. Scale bar = 100 µm (a). n = 6 animals per condition. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ by two-tail Student´s t-test (HAS2Control vs. MCAO Contra) and Tukey posthoc test (after one-way ANOVA was significant at $p < 0.05$).

Figure 8 – HA degradation in the SVZ correlates with Thbs4 response to brain ischemia. (a) Representative image of Thbs4 and HA (HABP) markers in the SVZ at 30 dpi. (b) Representative images of Thbs4 and HA in the sham and 15, 30 dpi SVZ. (c, d) HA levels did not change in the entire SVZ after MCAO (c) but decreased significantly in the dorsal SVZ (d). Thbs4 expression increased throughout the SVZ after MCAO (c), with a marked rise at 30 dpi in the dorsal SVZ (d), coinciding with HABP reduction. (e) Representative confocal (top) and skeletonized (bottom) images of HABP in the dorsal SVZ of sham, 15 and 30 dpi mice. (f) Cumulative distribution of HA skeletons showed a reduction in the length of HA cable-like structures only in the dorsal SVZ at 7 and 15 dpi. (g) The interconnectivity of the extracellular matrix decreased by 7 dpi in the dorsal SVZ, as measured by fractal dimension. (h) Experimental design for qPCR: 3–4-month-old mice were subjected to MCAO, and fresh SVZ tissue was extracted 15 and 30 days after MCAO for RT-qPCR analysis. (i) Hyaluronan degradation genes (Hyal1, Hyal2 and Hyal3) increased by 15 and 30 dpi in the SVZ, while hyaluronan synthase genes (HAS1 and HAS2) were overexpressed later, at 30 dpi. Thbs4 and Hif1a were also upregulated by 15 and 30 dpi in the SVZ. Scale bar = 100 µm (a). n = 6 (c, d, e and g) and 3 (i). $*p < 0.05$, $**p < 0.01$ and $***p < 0.001$ by Krustal Wallis test and Tukey post-hoc test (after one-way ANOVA was significant at $p < 0.05$).

Supplementary Figure 1 – Characterization of the Middle Cerebral Artery Occlusion (MCAO) mouse model of brain

ischemia. Related to Figure 2. (a) Scheme of the procedure: 10 mm silicon-coated filament was inserted through an external

artery and re-directed to the internal one. Ischemia was induced after 60 minutes of occlusion. After that, the filament was

removed, and animals were observed every day before euthanasia. (b) Volume of infarction was observed by cresyl violet

staining. (c) Neuronal death was observed by NeuN immunofluorescence. (d) Around 50% of ischemic mice survived at 28

dpi. (e) Ischemic animals ameliorated neurological symptoms over time. (f) Weight was measured as controls for mice

healthcare. n = 17 (d, e, f). **$p < 0.01$ and ****$p < <0.0001$ by survival curve test and Tukey post-hoc test (after one-way

ANOVA was significant at $p < 0.05$).

Supplementary Figure 2 – Cell proliferation in the SVZ after brain ischemia. Related to Figure 2. (a) Experimental design: three BrdU (50mg/kg) injections were performed the day before MCAO protocol. Animals were sacrificed 24 hours after the brain stroke model. (b) BrdU-positive cells increased in the SVZ 24 hours after MCAO. (c) Representative image of BrdU positive cells in the ipsilateral (left) and contralateral (right) hemisphere 24 hours after MCAO. (d) Ki67 marker was also used to analyze cell proliferation in wholemount preparations of the SVZ. (e) Ki67 positive cells increased in the SVZ 24 hours after MCAO. (f) The Cleaved-Caspase 3 marker increased in the SVZ 24 hours after MCAO. (g) Cleaved-caspase 3 marker was used to assess apoptosis in the SVZ after brain ischemia. Scale bar = 15 µm (g). n = 5 (sham) and 4 (MCAO). Bars represent mean ± SEM. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ by two-tail Student´s t-test (sham vs. 24h MCAO).

846 **Supplementary Figure 3 – Thbs4 astrocytes derive from slow proliferative cells after brain ischemia.** Related to Figure

847 3. (a) Slow proliferative cells were quantified in the lateral (grey), ventral (orange), medial (yellow) and dorsal (blue) SVZ.

848 We only observed an increase of Thbs4/GFAP/BrdU positive cells in the dorsal SVZ 30 dpi. (b) Thbs4/GFAP/IdU positive

849 cells did not show changes in the lateral (grey), ventral (orange), medial (yellow) and dorsal (blue) SVZ after brain ischemia.

850 n = 4 (sham) and 3 (MCAO). **p < 0.01 by by two-tail Student´s t-test (sham vs. 30dd MCAO).

Supplementary Figure 4 – Ischemic-induced Thbs4 astrocytes migrate preferentially to caudal infarcted areas.

Related to Figure 4. (a) Representative image of tdTOM positive cells in the OB three weeks after electroporation (3wpi). (b) tdTOM-positive cells decreased in the rostral, intermediate and caudal SVZ 7, 30 and 60 dpi. (c) Thbs4/tdTOM positive cells increased in rostral and intermediate SVZ 30 and 60 dpi. (d) Expression of tdTomato increased in intermediate and caudal brain areas at 30 dpi. (e) Representative images of tdTOM expression in rostral (left), intermediate (middle) and caudal (right) brain areas of sham (top) and 60 dpi mice (bottom). (f) tdTOM positive cells increased in the infarcted cortex (Ctx) and striatum (Str) whereas decreased in the dorsal SVZ (SVZ). No changes were observed in the corpus callosum (CC). (g) tdTOM positive cells only increased in the infarcted cortex and striatum at intermediate and caudal brain regions. Scale bar = 100 µm (a) and 1000 µm (e). n = 6 animals per condition. Bars represent mean ± SEM. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ by Dunnett/Tukey post-hoc test (after two-way ANOVA was significant at $p < 0.05$).

Supplementary Figure 5 – SVZ NSC stop producing newborn neurons to send newborn astrocytes to the ischemic

areas. Related to Figure 5. (a) Thbs4 expression in the corpus callosum (cc), infarcted cortex (Ctx) and infarcted striatum

(Str) of sham and 7 and 30 dpi mice. (b) Thbs4 increased by 30 dpi in the corpus callosum (cc) and 60 dpi in the infarcted

cortex (Ctx) and infarcted striatum (Str). (c) Expression of Thbs4 increased in total GFAP population after 15, 30 and 60 dpi.

(d) Experimental design: chronic BrdU (1% in water) treatment was administered for two weeks. BrdU only persisted in

slow proliferative NSC (Codega et al., 2014). DCX and BrdU positive cells were measured in the OB layers 7, 15 and 30

dpi. (e) Representative images of BrdU and DCX marker in sham (left), 15 (middle) and 30 (right) dpi mice OB. (f) DCX

positive cells decreased 15 and 30 dpi in the RMS, granular OB layer (GrL) and glomerular OB layer (GL). (g) BrdU

positive cells only decreased 30 dpi in the glomerular OB layer (GL). Double DCX/BrdU positive cells significative

decreased in the RMS 30 dpi. In (f) and (g), values are normalized to total DAPI nuclei (100%). Scale bar = 500 µm (a). n =

6 animals per condition. Lines represent mean ± SEM. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ by Chi-square test and Tukey

post-hoc test (after two-way ANOVA was significant at $p < 0.05$).

Supplementary Figure 6 – Thbs4 astrocytes internalize HA in the infarcted areas. Related to Figure 6. (a) Representative
images of Thbs4/GFAP ("newborn astrocytes", top) and GFAP-only ("local astrocytes", bottom) cells in the infarcted areas at
30 dpi. (b) Thbs4-positive astrocytes internalized more HA compared to local astrocytes only labelled with GFAP marker. (c)
However, the number of HA spots internalized was higher in local astrocytes compared with Thbs4 astrocytes, suggesting that
Thbs4 internalize HA in fewer vesicles. (d) Representative images of HABP (red), Thbs4 (green) and CD44 (blue) at the lesion
border. Inset shows an orthogonal projection of Thbs4/CD44/HABP-positive intracellular vesicle. (e) Thbs4 positive
astrocytes expressed a high rate of CD44 receptors so much in corpus callosum (CC), infarcted cortex (Ctx) as in infarcted
striatum (Str). Scale bar = 10 µm (c, f). n = 5 animals per condition. Bars represent mean ± SEM. *$p < 0.05$ and **$p < 0.01$ by
Tukey posthoc test (after two-way ANOVA was significant at $p < 0.05$) and two-tail Student´s t-test (Thbs4 vs GFAP).

Supplementary Figure 7 – In vitro NSC-derived Thbs4 cells synthetize HA after a hypoxic-ischemic condition. Related to Figure 6. (a) Experimental design: P4 pup rats were sacrificed, and SVZ NSC were extracted and cultured in the proliferative medium for one week. After that, neurospheres were disaggregated and seeded isolated in a culture dish. The following day, oxygen and glucose deprivation protocol (OGD) was performed. Cells were fixed 7 days after culture them. (b) Representative images of in vitro Thbs4 positive cells (green) and HA (red). (c) HA spots were measured inside and outside Thbs4 positive cells. (d) HA spots increased inside Thbs4 positive astrocytes after OGD whereas (e) decreased in the extracellular space. However, when NSC were exposed to hyaluronidases in order to remove extracellular HA, Thbs4 positive cells decreased internalization of HA spots (f) and increased extracellular HA spots (g). (h) Neurosphere-derived NSC (NSPH1) internalized more HA spots compared with Iba1-positive cells. (i) Metalloproteinase activity was higher in OGD NSC cultures compared to control condition. Scale bar = 10 µm (b). n = 15 samples per condition. Bars represent mean ± SEM. *$p < 0.05$, **$p < 0.01$ and ***$p < 0.001$ by two-tail Student´s t-test (CTRL vs. OGD; NSPH1 vs. Iba1+).

Table 1. Primary antibodies used for in vivo immunofluorescence protocol

Antibody	Host	Isotope	Dilution	Reference	Company
Thbs4	Mouse	Monoclonal IgM kappa light chain	1/400	Sc-390734	Santa Cruz Biotechnology, USA.
Thbs4	Goat	Polyclonal IgG	1/200	AF2390	R&B systems, USA
S100β	Rabbit	Polyclonal	1/400	Z0311	Dako, Agilent Technologies, Inc.
S100β	Mouse	Monoclonal IgG2a	1/1000	MA5-12969	Invitrogen, Spain
GFAP	Mouse	Monoclonal IgG1	1/1000	MAB3402	Sigma, Spain
GFAP	Rabbit	Polyclonal	1/4000	Z0334	Dako, Agilent Technologies, Inc.
βIII tubulin	Rabbit	Polyclonal	1/400	ab18207	Abcam, UK
NeuN	Mouse	Monoclonal IgG1	1/1000	MAB377	Millipore, Germany
DCX	Rabbit	Polyclonal IgG	1/500	ab18723	Abcam, UK
Nestin	Chicken	Polyclonal IgY	1/1000	NES	Aves Lab, EEUU
Prominin 1	Rat	Monoclonal IgG1 kappa	1/300	14-1331-82	Invitrogen, Spain
EGFR	Mouse	Monoclonal	1/400	SAB4200809	Abcam, UK
BrdU	Rat	Monoclonal	1/400	MCA2060	Bio-Rad, USA.
BrdU	Mouse	Monoclonal IgG1, kappa	1/500	03-3900	Invitrogen, Spain
IdU	Mouse	Monoclonal IgG2b	1/100	MA5-24879	Invitrogen, Spain
Ki67	Rabbit	Polyclonal IgG	1/400	Ab15580	Abcam, UK
Cleaved-Caspase 3	Rabbit	Monoclonal IgG	1/300	#9664	Cell Signaling, Germany
CD44	Rat	Monoclonal IgG2b	1/500	MA517875	Invitrogen, Spain
FLAG	Mouse	Monoclonal IgG1	1/500	F1804	Sigma, Spain
Olig2	Mouse	Monoclonal IgG2aκ	1/1000	MABN50	Millipore, Germany

Table 2. Primer sequences for qPCR

Name	Sequence
Hyal1_FW	GTGCCAAGCCCTATGCTAATAAG
Hyal1_REV	GCATGTCCATTGCAAAGACTGA
Hyal2_FW	GTCCCACATACACCCGAGGA
Hyal2_REV	GGCACTCTCACCGATGGTAGA
Hyal3_Ffw	GGACGACCTGATGCAGACTATTG
Hyal3_REV	GGTCCCCCCAGAGTACCACT
Has1_FW	AGGGCTCTTAAAGGAGGAGTCC
Has1_REV	AGAAGGTAAACTGAGTCCCCAGAA
Has2_FW	CAAAGAGGTTCGTTCAAGTTCTGA
Has2_REV	TGTGTTTGTTTCCCACTAGCTCTC
Hif1a_FW	CATAAAGTCTGCAACATGGAAGGT
Hif1a_REV	ATTTGATGGGTGAGGAATGGGTT
HPRT1_FW	TGATAGATCCATTCCTATGACTGTAGA
HPRT1_REV	AAGACATTCTTTCCAGTTAAAGTTGAG