Genomic signatures associated with the evolutionary loss of egg yolk in parasitoid wasps

Xianxin Zhao[1], Yuanyuan Liu[1], Yi Yang[1], Chun He[1], Kevin C. Chan[2], Haiwei Lin[1], Qi Fang[1], Gongyin Ye[1, *], Xinhai Ye[3, *]

[1]State Key Laboratory of Rice Biology and Breeding & Ministry of Agricultural and Rural Affairs Key Laboratory of Molecular Biology of Crop Pathogens and Insects, Institute of Insect Sciences, Zhejiang University, Hangzhou 310058, China.

[2]Shanghai Institute for Advanced Study, Zhejiang University, Shanghai 201203, China.

[3]College of Advanced Agriculture Science, Zhejiang A&F University, Hangzhou 311300, China.

[*]Corresponding authors: yexinhai@zafu.edu.cn; chu@zju.edu.cn.

Abstract

Trait regression and loss have occurred repeatedly in numerous lineages throughout evolutionary history in response to changing environments. In parasitoid wasps, a mega-diverse group of hymenopteran insects, loss or reduction of yolk in the egg has been reported in many species. This phenotypic change likely evolved as a response to the shift from ectoparasitism to endoparasitism. However, the genetic basis of this trait and the impact of its loss on genome evolution remain poorly understood. Here, we performed a comparative genomic analysis of 64 hymenopteran insects. The conserved insect yolk protein gene *vitellogenin* (*Vg*) underwent five independent loss events in four families, involving 23 of the analyzed endoparasitoid species. Whole-genome alignment suggested that *Vg* loss occurred during genome rearrangement events. Analysis of *Vg* receptor gene (*VgR*) loss, selection, and structural variation in lineages

lacking *Vg* demonstrated functional biases in the patterns of gene loss.

The ectoparasitism to endoparasitism transition did not appear to be the

primary driver of *Vg* loss or the subsequent *VgR* evolution. A number of

parallel and convergent genomic changes were observed between *Vg*-loss

lineages, including gene family evolution and selection of genes related

to transport, development, and metabolism. These changes may have

facilitated embryonic development without the yolk in these lineages.

Together, these findings reveal the genomic basis underlying a unique

trait loss in parasitoid wasps. More broadly, this study enhances our

understanding of yolk loss evolution outside the class Mammalia,

highlighting a potential evolutionary trend arising from the availability of

an alternative nutrient source for embryonic development.

Key words: Trait loss evolution, Egg yolk loss, Gene loss, Vitellogenin,

Vitellogenin receptor, Parasitoid wasp, Parasitic strategy evolution

Introduction

A fundamental goal in the field of evolutionary biology is to understand

the genetic bases underlying phenotypic shifts and evolutionary novelties

throughout the tree of life. Trait regression or loss (the reduction or

complete loss, respectively, of an ancestral trait) are phenotypic shifts that

have occurred repeatedly throughout evolutionary history. These shifts

likely occur in response to alterations in the environment, habitat, or

species interaction network[1–3]. Multiple genome sequencing and

comparative genomic studies have been conducted to uncover the genetic

bases of evolutionary trait loss in animals, including eye and

pigmentation loss among animals living in darkness[4–8], tooth loss in

birds[9–11], limb loss in snakes[12], and tail loss in humans and apes[13]. Such

studies have highlighted the importance of gene loss and inactivating

mutations in the evolution of trait loss or regression. These investigations

have also demonstrated the power of comparative genomic approaches to

reveal the genetic mechanisms underlying complex trait evolution.

Trait loss and the associated genetic bases are topics of great research

interest in the context of parasite evolution, particularly because long-

term host dependence may lead to trait loss in processes such as nutrient

acquisition[14,15]. For example, many parasitic plants show a loss of genes

associated with photosynthesis, nutrient uptake, flowering regulation,

defense responses, and root development[16–19]. A large-scale genomic

analysis of parasitic nematodes and platyhelminths revealed the

widespread losses of metabolic pathways and differential pathway

coverage across different evolutionary clades[20]. Other studies have

revealed the loss of homeobox gene families[21] and cilium-related genes[22]

in parasitic worms, which may be related to their simplified morphology.

Insects are the largest and most diverse group of organisms on Earth.

Parasitoids, defined as parasitic species that eventually kill their hosts,

have evolved repeatedly and independently in several insect orders[23]. The

most common and representative parasitoid species are parasitoid wasps

of the order Hymenoptera; these are extremely biologically diverse, and a

rapid accumulation of genomic data has made them model organisms in

the field of evolutionary biology[24–31]. Comparative genomic analyses of

parasitoid wasps and their free-living relatives can reveal novel

evolutionary trait losses or regressions, providing new insights into the

evolution of parasitism.

Some parasitoid wasp species exhibit an intriguing trait loss with a

visible phenotype: loss of the egg yolk. Eggs with little or no yolk were

first recorded in the endoparasitoid *Chalcis sispes*, and are widespread

among parasitoid wasps from various lineages[33–37] (Supplementary Table

1). Such yolkless eggs have been termed "hydropic", referring to their

method of nutrient acquisition from host hemolymph, which distinguishes

them from anhydropic eggs[32]. Moreover, the phenomenon of radiolabeled

amino acids absorption from host hemolymph was found in *Microplitis*

croceipes, rather than protein taken[33], implying that yolkless insect eggs

have evolved to obtain the nutrients necessary for development directly

from the host. Indeed, yolk trait loss has been likely linked to the

evolution of parasitic strategies (e.g., the shift from ectoparasitism to

endoparasitism). Thus, yolk loss has likely evolved as a result of

embryonic development inside the host[23,38,39], although this has not been

proven.

To uncover the genetic basis of egg yolk loss in parasitoid wasps, we

here conducted a comparative genomic analysis using *vitellogenin* (*Vg*), a

conserved gene encoding an egg yolk protein precursor in insects[38,40].

Using a broad, representative selection of hymenopteran species, we

identified lineages with *Vg* loss events and established a putative loss

mechanism. Furthermore, we examined loss of a *Vg*-related gene to

assess co-evolution in multiple lineages. These findings enhance our

knowledge of yolk loss and parasitoid wasp genome evolution,

contributing more broadly to an improved understanding of the genomic

changes associated with shifts in parasitic strategies and embryonic

developmental conditions.

Results

***Vg* loss was recurrent in Hymenoptera but confined to**

endoparasitoid wasps

We first sought to identify *Vg*s and homologous genes in members of the

order Hymenoptera. Thus, a comparative genomic investigation was

conducted in 64 representative hymenopteran species, including the

major lineages: sawflies, wasps, ants, and bees (Fig. 1b). A

comprehensive homology-based search was followed by protein domain

screening and manual curation (see **Methods**), yielding a total of 237

genes. These genes were categorized into three groups based on the

presence or absence of specific functional domains: 64 complete classic

Vg genes (*Vg*s), 169 *Partial Vg* genes (*PVg*s), and four *Vg-like* genes

(*Vgl*s) (Fig. 1a; Supplementary Table 2). The classic *Vg*s encoded

proteins with three major domains[38,41], namely Vitellogenin_N or LPD_N

(PF01347), Vit_openb-sht (PF09172), and von Willebrand factor type D

(vWD) (PF00094); notably, we identified seven *Vg*s in seven species that

were incorrectly annotated in prior genome annotations (Supplementary

Fig. 1). The *PVg*s encoded other proteins containing the Vitellogenin_N

domain, whereas *Vgl*s encoded proteins without the Vitellogenin_N

domain but with the Vit_openb-sht and/or vWD domain.

To place our findings in an evolutionary context, a phylogenetic tree

was constructed for the 64 species using 1116 single-copy orthologous

genes. The *Vg* identification results were then mapped onto the resulting

tree. This revealed a great deal of variation in *Vg* copy number within the

order Hymenoptera. Interestingly, *Vg* was absent from the genomes of 24

parasitoid wasps in three superfamilies (Ichneumonoidea, Chalcidoidea

and Cynipoidea) that contain diverse parasitoid species (Fig. 1b;

Supplementary Table 2). In contrast, one to four copies of *Vg* were

conserved in all other non-parasitoid hymenopteran species, including

sawfly (*Athalia*), vespid wasps (*Vespa* and *Polistes*), ants (*Ooceraea*,

Atta, and *Solenopsis*), and bees (*Colletes*, *Megalopta*, *Apis*, and *Bombus*).

Vg loss was presumed in the *Drosophila* parasitoid *Asobara japonica*

(Braconidae), but this may have been an artifact due to the presence of a

PVg gene located at the end of the scaffold (Supplementary Fig. 2; Fig.

1b).

Vg loss appeared to have occurred independently in five different

parasitoid wasp lineages, leading to the observed lack of *Vg* in 23 species

(excluding *A. japonica*). Specifically, *Vg* was lost in the common ancestor

of two *Leptopilina* species in the superfamily Cynipoidea; the common

ancestor of two *Trichogramma* species in the superfamily Chalcidoidea;

the common ancestor of *Diadegma semiclausum* and *Hyposoter*

didymator in the family Ichneumonidae; the common ancestor of three

aphid parasitoids of the genera *Aphidius* and *Lysiphlebus* in the family

Braconidae; and the common ancestor of 14 braconid parasitoids from

the genera *Cotesia*, *Microplitis*, *Chelonus*, *Macrocentrus*, and

Eumacrocentrus in the family Braconidae (Fig. 1b). In addition, the

phylogenetic tree demonstrated extensive variations in *PVg* and *Vgl* gene

copy number across hymenopteran species, suggesting complex

evolutionary histories.

To understand the relationships between *Vg* loss and parasitic

strategies, the 64 species were divided into three categories based on the

location in which their eggs are laid and embryos develop:

ectoparasitoids (those that lay eggs on the external host surface),

endoparasitoids (those that lay eggs inside the host)[23], and unknown.

These categorizations were then mapped to the phylogenic tree (Fig. 1b).

Based on the key role of Vg proteins in yolk formation (vitellogenesis)[40],

we hypothesized that *Vg* losses would be restricted to endoparasitoids due

to the nutrient-rich environment (host hemolymph) [42,43] in which the eggs

reside. Indeed, all of the parasitoid wasps with *Vg* loss were

endoparasitoids. However, not all of the analyzed endoparasitoids

demonstrated *Vg* loss; 35% retained copies of *Vg*. This suggested that

endoparasitism may not have been the primary factor driving *Vg* loss in

parasitoid wasps.

PVg was not derived from _Vg_ disruption in _Vg_-loss species

We hypothesized that recently mutated *Vg*s retaining the Vitellogenin_N

domain (i.e., *PVg*s) may have retained some *Vg* functionality due to the

importance of the N-domain for oocyte uptake[44,45]. Thus, *Vg* loss may not

have directly resulted in a lack of vitellogenesis if *PVg* was present. To

investigate the evolutionary origins of *PVg*s and to test whether these

genes arose from recent *Vg* disruption among *Vg*-loss species, a

phylogenetic analysis was performed using all *Vg*s and *PVg*s identified in

this study. If the *PVg*s had primarily originated from *Vg* mutation, *PVg*s

would cluster together with *Vg*s. Instead, the Vg sequences showed a

distinct evolutionary history from the PVg sequences (Fig. 2a). Notably,

the N-terminal domain sequences of eight PVgs in seven species were

located in the Vg clade, suggesting that these specific PVgs may have

resulted from Vg disruption. However, none of the corresponding species

had undergone *Vg* loss events, with the exception of the ambiguous *Vg*

loss in *A. japonica* as noted above.

The functions of *PVg*s in insects have not yet been characterized. To

establish whether there was putative functional overlap between *Vg*s and

*PVg*s, we examined *Vg* and *PVg* expression in 10 hymenopteran species

for which there were publicly available RNA sequencing data generated

from adults of both sexes. *Vg* was generally expressed at higher levels in

females than in males, whereas the opposite was true of *PVg* (Fig. 2b;

Supplementary Table 3). These distinct expression patterns between *Vg*s

and *PVg*s in adult hymenopterans suggested that the genes have divergent

functionality. Overall, these results revealed dramatic differences between

*Vg*s and *PVg*s with respect to both evolutionary history and gene

expression. This supported the hypothesis that vitellogenesis was largely

arrested in endoparasitoid wasps with *Vg* loss, consistent with previous

reports of yolkless eggs[32,35] (Supplementary Table 1).

_Vg_ loss was likely caused by genome rearrangement

To explore the potential causes of *Vg* loss in endoparasitoid wasps, the

genomes of species with *Vg* loss were examined for regions that were

syntenic with *Vg* loci in their *Vg*-retaining relatives. Because *Vg* loss

events span large evolutionary distances across hymenopteran phylogeny,

syntenic regions containing *Vg* were examined only within each family or

superfamily. In Ichneumonidae, the genes encoding chondroitin sulfate

N-acetylgalactosaminyltransferase and alanine-glyoxylate

aminotransferase were identified as marker genes for the *Vg* region; these

two genes flanked the genomic region containing *Vg* in wasps of the *Vg*-

retaining genera *Ophion* and *Venturia*. A search for remnants of *Vg* (e.g.,

a pseudogene or inactive gene) in the corresponding region of *H.*

didymator, which had lost *Vg*, revealed no homologous sequences (Fig.

3a; see **Methods**). Similar searches did not reveal *Vg* remnants in other

Vg-loss species of the families/superfamilies Braconidae, Cynipoidea, or

Chalcidoidea (Fig. 3b, c; Supplementary Fig. 3). These results suggested

that *Vg* loss was not due to gene-inactivating mutations. However, it is

also possible that rapid sequence evolution prevented homologous

sequence detection.

We next asked whether genome rearrangements had contributed to *Vg*

losses. This was addressed through detection of genome rearrangement

breakpoints between *Vg*-loss species and their *Vg*-retaining relatives in three independent lineages (Cynipoidea, Chalcidoidea, and Braconidae) via whole-genome alignment (Fig. 3d; see **Methods**). Ichneumonid species were excluded from this analysis because the genome assemblies were highly fragmented, with over 13,000 scaffolds for *Vg*-loss species, which could have negatively affected the accuracy of genome-wide alignments. In Cynipoidea (Group 1), 2151 genome rearrangement breakpoints were identified between *Leptopilina boulardi* (*Vg* loss) and *Belonocnema kinseyi* (*Vg* retaining). Notably, the breakpoints significantly overlapped with the *Vg* locus in the *B. kinseyi* genome (Fig. 4e). Similar results were observed for species in Chalcidoidea and Braconidae (Fig. 4e; Observed breakpoints vs. a set of control regions by 10,000 times randomly shuffling breakpoints on the same chromosome, *p* = 0, one-tailed permutation test). Taken together, these findings indicated strong contributions of genome rearrangement to *Vg* loss events in endoparasitoid wasps.

Gene loss and relaxed selection on the *Vg* receptor gene (*VgR*) after *Vg* loss

Evolutionary patterns of gene loss are usually not stochastic, but rather display a clear bias with respect to gene function or genomic position[46–48]. This fact prompted us to investigate the mechanisms by which genes associated with *Vg* function evolved in *Vg*-loss lineages. We first focused on *VgR*, which is crucial for incorporation of Vg proteins into insect oocytes[38,40,41,49,50]. We hypothesized that *VgR*s may be absent or subject to distinct evolutionary pressure in *Vg*-loss species. To test this hypothesis, *VgR* genes were identified in the genomes of all 64 examined hymenopteran species. *VgR* was absent from 19 of the genomes and was

present as a single-copy gene in the other 45 (Fig. 4a). Based on *VgR*

presence or absence, we identified four independent *VgR* loss events

throughout hymenopteran evolution; these were strongly correlated with

Vg loss events (Fig. 4a). A high proportion of *Vg*-loss species, excluding

A. japonica, had also lost *VgR* (78%), with only five species retaining

VgR (namely two *Leptopilina* species, two *Trichogramma* species, and *H.*

didymator). Interestingly, only one species (*Telenomus remus* of the

superfamily Platygastroidea) retained *Vg* but lost *VgR*.

We next examined whether *VgR* was under relaxed selective pressure

after *Vg* was lost. Using the selection intensity parameter K, where K < 1

indicates relaxed selection and K > 1 indicates strong purifying selection,

we screened for signatures of relaxed selection on all test branches

leading to all *Vg*-loss *VgR*-retaining species, under the RELAX model[51]

in HyPhy[52]. As expected, the result revealed a significant relaxed

selection pattern on these *VgR*s (Fig. 4b; K = 0, *p* = 0). Independent tests

for each of the *Vg*-loss lineages indicated significant relaxed selection on

VgR in *Leptopilina* wasps (Fig. 4c; K = 0, *p* = 0) and in *H. didymator*

(Fig. 4e; K = 0.72, *p* = 0.004), but not in *Trichogramma* spp. (Fig. 4d; K

= 1.06, *p* = 0.916) or *A. japonica* (Fig. 4f; K = 1.21, *p* = 0.094). Overall,

these findings suggested that *Vg* loss in endoparasitoid wasps resulted in

subsequent loss or relaxed selection on *VgR*, a novel example of co-

elimination of functionally linked genes[46].

Structural variations in the conserved VgR domain of *Leptopilina*

may have led to functional loss

Relaxed selection often leads to amino acid substitutions that alter the

three-dimensional structures (and therefore functions) of the encoded

proteins. To determine whether this was true of VgRs in *Vg*-loss lineages,

we predicted the structures of all 45 VgRs identified in this study with

AlphaFold2. Based on the confidence scores (pLDDT), we focused only

on the three most well-structured regions, designated A, B, and C (Fig.

5b–d). Each region contained one YWTD repeat (six YWTD motifs) and

two epidermal growth factor (EGF)-like repeats (Fig. 5a). Comparative

analysis suggested that the structures of these three regions were

generally conserved between species (Fig. 5b–d). Notably, a portion of

region A in the two *Leptopilina* VgRs differed substantially from those of

all other VgRs (Fig. 5f); this region overlapped with the second EGF-like

repeat of region A. Previous studies have shown that mutations in the

EGF-like repeat of VgR severely disrupt the process of yolk formation[53]

and that this functional domain is essential for Vg dissociation under

acidic conditions[54]. We therefore hypothesized that structural variants in

this EGF-like repeat region may have led to loss of the original VgR

function in *Leptopilina* species, consistent with the results of a recent

study[55].

The transition from ectoparasitism to endoparasitism did not directly

cause *Vg*/*VgR* loss or relaxed selection

The transition from ectoparasitism to endoparasitism is considered a key

step and a major trend in the evolutionary history of parasitoid wasps[23,56].

As noted above, although *Vg* and *VgR* loss events were restricted to

endoparasitoids, a number of endoparasitoids retained one or both genes.

This implied that the transition from ectoparasitism to endoparasitism

was not the primary force driving these gene losses. To gain deeper

insights into the reasons for *Vg*/*VgR* loss, we next tested whether the *Vg*s

and *VgR*s in endoparasitoids evolved under relaxed purifying selection. If

the transition to endoparasitism drove *Vg* or *VgR* evolution, there should

be clear signals of relaxed selection on *Vg*s/*VgR*s in other *Vg*-retaining

endoparasitoids compared to both ectoparasitoids and free-living species.

However, the results showed no evidence of relaxed selection on *Vg*s or

*VgR*s in *Vg*-retaining endoparasitoids (K = 1.04, *p* = 0.561 for *Vg*; K =

1.21, *p* = 0 for *VgR*). We therefore concluded that the ectoparasitism to

endoparasitism transition was not the primary factor underlying either *Vg*

loss or the subsequent loss/relaxed selection on *VgR*s in parasitoid wasps.

Changes throughout the genome were consistent in *Vg*-loss lineages

Two genome-wide screens were next conducted to identify genetic

changes in *Vg*-loss species. These screens were designed to clarify

specific evolutionary patterns after *Vg* loss in different parasitoid wasp

lineages and to determine whether parallel or convergent genomic

changes had occurred. We first investigated changes in the sizes of gene

families, also known as orthogroups (OGs). Expanded and contracted

OGs were identified in the most ancient branches leading to the *Vg*-loss

lineages. In each of the five independent branches associated with *Vg*

loss, there were 39–462 expanded OGs and 120–713 contracted OGs

(Fig. 6a).

Comparative analysis showed that most changes in OG size were

branch-specific; only two expanded and 40 contracted OGs were likely to

have occurred in parallel (i.e., were shared between three or more

lineages) (Fig. 6b, c; Supplementary Table 4, 5). The expanded OGs

included a multidrug resistance-associated gene encoding a member of

the ATP-binding cassette (ABC) transporter family that transports

numerous substrates across the cellular membrane[57]. We inferred that

parallel expansion of this OG may have enhanced the capacity of

endoparasitoids to utilize host nutrients in the early development stage,

compensating for the lack of nutrients from the yolk. Parallel contracted

OGs included lipase, spaetzle, farnesol dehydrogenase, and trypsin.

Functional annotation was conducted for the expanded and contracted

OGs between the five lineages to search for signatures of convergent

functional evolution (see **Methods**). The results showed that

convergently expanded or contracted OGs (those that were shared by four

or all five lineages) were primarily associated with transport (e.g., organic

substance transport or monoatomic ion transport), metabolism (e.g.,

steroid, amine, or peptide metabolic processes), and development (e.g.,

embryo or cuticle development) (Fig. 6d, e; Supplementary Table 6, 7).

The second genome-wide screen entailed analyzing the 2652 protein-

coding one-to-one orthologous genes in the five ancestral branches

leading to *Vg*-loss lineages for signatures of positive or relaxed selection

(see **Methods**). One gene of unknown function was under positive

selection in all five branches; five genes were under positive selection in

four branches; and 27 genes were under positive selection in three

branches (Supplementary Table 8). These genes were annotated with

diverse functions, including "double-stranded RNA-specific editase",

"histone lysine N-methyltransferase", and "zinc finger protein". There

were 199 genes with relaxed selection signals in all five branches.

Functional enrichment analysis indicated that these genes were related to

nervous system development, embryo development, locomotion, and

stress responses (false discovery rate-adjusted $p < 0.05$; Supplementary

Table 9, 10; Supplementary Fig. 4). Together, these results demonstrated

parallel or convergent changes in specific gene families in addition to

shared gene selection events across five independent *Vg*-loss lineages.

These genomic changes may have been related to adaptation to the

endoparasitic lifestyle (Fig. 7).

Discussion

In this study, we used a comparative genomic approach to explore the genetic basis of egg yolk loss in endoparasitoid wasps, the origin and spread of which constitutes a long-standing puzzle in the field of evolutionary biology. We focused our analysis on *Vg*, an egg yolk precursor gene found in nearly all oviparous animals, including fish, birds, and insects[58]. To better understand *Vg* evolution, we annotated *Vg*s and homologs (*PVg*s and *Vgl*s) in 64 representative hymenopteran genomes. Although Vg structures and functions have been well characterized[38,41,50,59–62], the molecular functions of PVgs and Vgls in insects are largely unknown. Here, phylogenetic and gene expression analyses revealed distinct evolutionary histories and expression patterns of *Vg* compared to *PVg*, suggesting divergent functionality of these genes. Similarly, in *Apis mellifera* and *Formica* ants, *PVg* reportedly functions in inflammation and/or oxidative stress processes rather than vitellogenesis due to sub-functionalization or neofunctionalization[63,64]. Further studies will be required in other species to clarify the functionality and evolutionary patterns of *PVg*s.

We discovered notable *Vg* loss events in 23 species from five independent lineages of endoparasitoid wasps in three superfamilies (Ichneumonoidea, Chalcidoidea, and Cynipoidea). Of the species with *Vg* loss, 14 (namely *Aphidius ervi*, *Aphidius gifuensis*, *Cotesia congregata*, *Cotesia glomerata*, *Cotesia vestalis*, *Diachasma alloeum*, *Diadegma semiclausum*, *Fopius arisanus*, *Leptopilina boulardi*, *Lysiphlebus fabarum*, *Macrocentrus cingulum*, *Microplitis demolitor*, *Microplitis mediator*, and *Venturia canescens*) have previously been identified as having yolkless eggs (Supplementary Table 1); relevant studies of egg

morphology have not been conducted for the remaining 9 species. None

of the species containing *Vg* have been reported to have yolkless eggs,

demonstrating a strong correlation between *Vg* loss and egg yolk

disappearance in the species analyzed here. In contrast to the

endoparasitoids, all of the ectoparasitoids and free-living hymenopterans

were found to retain at least one copy of *Vg*, demonstrating restriction of

Vg loss to endoparasitoids.

Genome alignment and synteny analyses were conducted to identify

remnants of *Vg* genes, which could provide insights into the evolutionary

mechanism of *Vg* loss. Even using a low alignment threshold, *Vg*

pseudogenes (e.g., those with clearly disabling insertions, deletions,

frameshifts, or truncations) were not identified. This was likely due to the

relatively distant timing of *Vg* loss (estimated at ~24.32–124.58 million

years ago) and the high evolutionary rates of hymenopteran genomes[65,66].

Interestingly, in three independent lineages, *Vg* loci were frequently

found at genome rearrangement breakpoints, implying that a genome

rearrangement event in the *Vg* region was likely the cause of gene loss in

many of these species (Fig. 7). Comparison of a greater number of

genomes and identification of recent *Vg* loss events would provide more

robust evidence supporting this proposed mechanism of *Vg* loss.

Previous genomic studies have revealed apparent functional biases in

gene loss events[46], likely resulting from co-elimination of functionally

related genes. This may be caused by changes in selective pressure on a

specific trait due to adaption to a new living environment or to new

biological characteristics[46,67–69]. For example, eight skin-related genes

have been inactivated in both cetaceans and hippopotami over the course

of evolution, potentially due to aquatic skin adaptations[70]. Fruit bats have

lost functionality of three renal transporter genes (*SLC22A12*, *SLC2A9*,

and *SLC22A6*) and a renal ammonium secreting transporter gene

(*RHBG*), likely contributing to efficient excretion of the excess water

obtained through their unique diet[71]. Similarly, we here discovered co-

elimination of *Vg* and *VgR* in 21 endoparasitoid wasp species. Based on

current knowledge of *Vg* and *VgR* evolution in Hymenoptera, we propose

that loss or relaxed selection on *VgR* was likely a direct consequence of

Vg loss, which rendered *VgR* dispensable (Fig. 7).

Despite the widespread loss of *VgR* among endoparasitoids, five *Vg*-

loss species retained *VgR*, and the gene was under significantly relaxed

selection in two *Leptopilina* wasp species (*L. boulardi* and *L. heterotoma*)

and in *H. didymator*. The structure of one *VgR* functional domain was

altered in the two *Leptopilina* species, suggesting a loss of the original

VgR function. There are several hypotheses for *VgR* maintenance among

Vg-loss species. First, the relatively short evolutionary time may not have

been sufficient to produce inactivating mutations or complete gene loss.

Second, *VgR*s may have undergone neofunctionalization. Consistent with

the latter hypothesis, a recent study of *L. boulardi VgR*, which is under

relaxed selection, showed that this gene no longer performs its original

function in ovary development but is involved in host-searching and

mating behavior[55]. The genomic sequences of additional endoparasitoid

wasps from different families and superfamilies should be analyzed in the

future to further explore the coevolutionary histories of *Vg* and *VgR*.

The transition from ectoparasitism to endoparasitism may have led to

relaxed selection on genes associated with nutritional status during early

developmental stages (e.g., vitellogenesis genes). This is because

ectoparasitoid embryos have strict nutrient requirements of the eggs

(primarily the yolk proteins), whereas endoparasitoid embryos are

directly exposed to a nutrient-rich external environment (the insect host

hemolymph)[42,43]. Thus, the dispensability of yolk-related functions could

lead to the loss of related genes in endoparasitoids. However, we found

no evidence for direct selection or co-elimination of *Vg* and *VgR* as a

result of the ectoparasitism to endoparasitism transition. This was

supported by previous reports of functional Vg proteins in endoparasitoid

wasps such as *Pteromalus puparum*[72–75]. Nevertheless, we did find that

co-elimination of *Vg* and *VgR* was restricted to endoparasitoids. This may

have been because endoparasitism leads to an alternative method of

embryonic nutrient acquisition, which is a prerequisite for co-elimination

of *Vg* and *VgR*. These two genes are crucial for formation of the yolk,

which supports embryonic development in the absence of external

nutrient availability. Loss of these genes could therefore lead to

dependence on external nutrients, hindering a return to the ectoparasitoid

state due to the generally irreversible nature of gene loss[71].

Determination of the mechanism(s) underlying trait loss or regressive

trait evolution remains challenging[46]. Here, it proved difficult to assess

whether loss of the yolk protein in endoparasitoids had a neutral or

positive effect on fitness. The hypothesis for positive selection suggests

that yolk loss could save energy costs and increase fecundity[76], but this

hypothesis has not been directly tested. Even if loss of the yolk protein in

endoparasitoids was determined to be a neutral process, other genomic

changes, such as the transporter gene duplication identified through

comparative genomics, could be adaptive. Detailed functional studies of

closely related *Vg*-loss and *Vg*-retaining species could shed light on the

impacts of *Vg* loss on fitness. In mammals, loss of the gene encoding the

egg yolk protein (*VIT*) has previously been reported as coinciding with

the origin of lactation and placentation[77]. We therefore propose that loss

of the egg yolk is likely a consequence of regressive evolution, which

may occur repeatedly during animal evolution when an alternative

nutrient source becomes available during embryonic development. A

study of the model species *Caenorhabditis elegans* has shown that the

embryos can develop without a yolk[78]. Thus, future studies should also

investigate yolk loss in nematode species, especially endoparasitic

nematodes.

In conclusion, we here conducted an extensive comparative genomic

analysis of 64 hymenopteran species to study loss of the vitellogenesis

gene *Vg*. We found evidence of five independent *Vg* loss events

throughout the order Hymenoptera, primarily occurring in species with

endoparasitic strategies. *Vg* loci in species that retained the gene were

frequently found in regions corresponding to genome rearrangements in

Vg-loss species, suggesting that loss of the gene was due to genome

rearrangement. Furthermore, *Vg* loss often coincided with loss or relaxed

selection of *VgR*. Overall, these findings enhance our understanding of

genome evolution and embryonic development across an important order

of insects, revealing key genomic changes associated with yolk loss.

Materials and methods

Data collection

Genomic data (including genomic sequences and genome annotation

results) of 64 representative hymenopteran insects were mainly collected

from publicly database, including the National Center for Biotechnology

Information (NCBI)[79], InsectBase v2[80], and Darwin Tree of Life

Project[81]. For RNA-seq analysis, we also collected RNA-seq data of 10

species from NCBI Sequence Read Archive (SRA) database. Find details

about data collection in Supplementary Table 11.

Vg and *VgR* identification

We used Bitacora v1.4[82] to identify candidate *Vg*s and *VgR*s. Full mode who searches genes in genomic sequences and protein annotations was selected to identify gene candidates. Besides ncbi-blast v2.13.0+ [83] and hmmer v3.3.2[84], the implemented tool, GEMOMA v1.7.1[85], a homology-based gene prediction program that uses amino acid sequence and intron position conservation to reconstruct gene models from TBLASTN alignments was used to identify new genomic regions encoding gene families. We use 13 vitellogenin protein sequences of 11 species from Uniprot[86] or NCBI database as query reference (see Supplementary Table 12). And two refence sequences of vitellogenin receptor of *Drosophila melanogaster* and *Solenopsis invicta* were retrieved from Uniprot database (see Supplementary Table 13). The specific HMM profiles were generated with alignment files of these reference sequences using hmmbuild command of hmmer v3.3.2[84]. Finally, each candidate gene was manually inspected, and further verified by checking their complete open reading frames. Then, the gene was divided into subgroups according to the following rule: sequences with N-terminal domain (Vitellogenin_N or LPD_N) as classical *Vg* (*Vg*) and partial *Vg* (*PVg*), while the other homologous members without N-terminal domain were categorized as *Vg*-like sequence (*Vgl*). For the potential chimeric sequences (e.g., potential gene model errors), we extracted the upstream and downstream 2000 bp sequences of these genes for re-annotation, using Augusts web interface[87] (http://augustus.gobics.de/) or FGENESH+ service online[88] (http://www.softberry.com/).

Gene expression

We used STAR v2.7.10b[89] and RSEM v1.3.3[90] for transcriptomes

assembly and expression calculation, respectively. Fragments Per

Kilobase Million (FPKM) values were used to quantify gene expression.

Synteny analysis

Gene synteny of selected species in three family Ichneumonidae, family

Braconidae and superfamily Cynipoidea were performed with JCVI

toolkit (MCScan Python version 1.3.3) [91] based on the alignments of each

pair of species generated by comparisons from LAST[92]. And high-quality

blocks of the *Vg* gene and its flanking genes were extracted for

visualization using a tool implemented in the JCVI toolkit.

Genomic rearrangement

We used LASTZ v1.04.03[93] to perform the pairwise genome alignments

with genomes of *Vg*-loss and *Vg*-retaining wasps. The *netSyntenic* files

were identified with axtChain/chainNet/netSyntenic tools based on

resulting alignments, and as input to identify the genomic rearrangement

breakpoints followed the workflow mentioned by Liao et al[94]. We

conducted a shuffling test to determine whether the median distance

between the randomly shuffled breakpoints and the *Vg* locus/loci was

significantly different from what would be observed. The shuffled

breakpoints were produced using shuffle function of bedtools[95] with

constraint of breakpoints region on the same chromosome, and shuffling

was repeated 10,000 times. The *p* value was calculated according to the

method described by Sun et al.[96].

Gene Selection

HyPhy package v2.5.33[52] was used for detecting gene selection signals.

Indeed, aBSREL[97] was used for detecting positively selection genes while

RELAX model[51] was used for identifying genes evolving under relaxed

selection ($p < 0.05$, the Bonferroni-Holm corrected). In RELAX result, K

was introduced as a parameter to assess the degree of relaxed selection,

specifically K < 1 indicates that the selection was relaxed, while K > 1

indicates that the selection was strengthened. Whereas, in the result of

positive selection test, the value of dN/dS greater than 1 indicated that the

test branches were positively selected, otherwise they were under

purifying selection.

Briefly, we used RELAX for testing relaxed selection on *Vg*s and

*VgR*s. To test whether the *VgR*s of the *Vg* loss species were under relaxed

selection, we first performed a general test in which all branches leading

to the *Vg* loss species (excluding *A. japonica*) were set as test branches,

while the remaining branches were set as reference. In addition, we also

performed a separate test on each branch leading to the *Vg* loss species;

To test whether the ecto-to-endo transition is a major driving force to

relax the purifying selection on *Vg*s and *VgR*s and then eventually lead to

their losses, we compared branches leading to endoparasitoid wasps (test)

and non-endoparasitoid wasps (reference) for both *Vg*s and *VgR*s.

We also searched for common positively/relaxedly selected genes on

the branches leading to the *Vg* loss lineages, which may reflect the

parallel or convergent genomic changes in the *Vg* loss lineages. In total,

2652 one-to-one orthologue protein-coding genes from 17 species were

involved in this selection analysis.

Orthology inference and phylogenetic reconstruction

We utilized OrthoFinder v2.5.5[98] to identify the orthologous and

paralogous genes, the longest transcripts of genes were kept to analysis.

The STAG (Species Tree Inference from All Genes) algorithm[99] that

integrated in OrthoFinder to identify Orthogroups (OGs) for phylogenetic

reconstruction. Based on the usage of "-M msa" parameter, genes from

1116 OGs which were developed to leverage not only strictly single-copy

orthologous genes but also the data from multi-copy gene families were

used for species tree inference. Multiple sequence alignments (MSA)

were inferred for each OG using MAFFT v7.520[100] and the species tree

was inferred from the concatenated MSAs using IQ-TREE v2.2.2.3[101]

with 1,000 replicates for ultrafast bootstrap analysis. The best-fitting

model of sequence evolution estimated by ModelFinder that integrated in

IQ-TREE v2.2.2.3[101] was used.

And we used r8s v1.81[102] to estimate the divergence times between

species or clade. The information of divergent time was previously

reported[103] and we used two time points as fix age, 281 million years ago

(mya) for Hymenoptera generated, and 8 mya for the divergent time of

Nasonia vitripennis and *Pteromalus puparum*, and other 12 to constrain

nodes marked by MCRA to calibrate the phylogenetic tree: Orussidae-

Pteromalinae: 211–289 mya, Apinae-Ampulicidae: 128–182 mya, *Apis*

mellifera-Bombus terrestris: 93–132 mya, Apinae-Microgastrinae: 203–

276 mya, Apinae-Gonatopodinae: 160–224 mya, Myrmicinae-Dorylinae:

65–127 mya, Vespinae-Polistinae: 34–71 mya, Cynipinae-Telenominae:

181–246 mya, Microgastrinae-Macrocentrinae: 82–147 mya,

Microgastrinae-Aphidiinae: 116–177 mya, Microgastrinae-

Campopleginae: 151–218 mya, Campopleginae-Cryptinae: 79–135 mya.

Gene family evolution

The gene family expansion and contraction were calculated using the

CAFE v5[104]. The content of each OG of 64 species generated from

OrthoFinder and the phylogenetic tree with divergence times inferred by

r8s in above were used as input files. Based on the results, the expanded

and contracted OGs of interested branches/clades that experienced Vg

loss events were specifically extracted, and the shared OGs and

associated function among these five branches/clades were inferred.

Gene ontology enrichment and functional similarity analysis

We annotated the amino acid sequences of each species using eggNOG-

mapper v.2.1.12 webserver[105] (http://eggnog-mapper.embl.de.) against

eggNOG5 database under default settings. And the biological process

category of gene ontology (GO) terms was mainly focused to explore the

most relevant functions in the interested OGs. In addition, GO

enrichment analyses among candidate genes were performed with

GOATOOLS[106] with the go-basic.obo database (http://geneontology.org/),

and p values for each GO terms were corrected for an FDR < 0.05

(Benjamini-Hochberg method of multiple testing correction). Using a

simple clustering algorithm in REVIGO v1.8.1[107], semantic relationships

of enriched GO terms were analyzed and representative subset of the

terms was selected for visualization.

To further unscramble the functional convergence in five Vg loss

linages, with the aid of the R package constellatoR[108,109]

(https://github.com/MetazoaPhylogenomicsLab/constellatoR), we firstly

assigned the interested OGs (expanded and contracted) onto GO terms

based on the functional similarity. And the function will be clustered with

an exemplar OG with the distance matrices which were transferred from

pairwise OG similarity matrices.

Protein structure analysis

We predicted the three-dimensional structures of VgR proteins from various species using LocalColabFold[110], which implements the AlphaFold algorithm[111]. The initial multiple sequence alignments (MSAs) for each protein were generated with the MMSeqs2 server[112]. We controlled the depth of the input MSAs by setting max_msa_clusters to 512 and max_extra_msa to 1024. All five neural networks were utilized for non-template-based predictions. The max_recycle parameter was set to 3 to balance accuracy and computational efficiency, without performing the post-prediction model refinements. From the predictions generated for each species, we selected the structure with the highest internal rank for subsequent analyses. Structural alignment was performed with MatchMaker[113] in ChimeraX[114]. A detailed list of sequences used for predictions is provided in the Supplementary Table 14. All figures were made with ChimeraX[114].

Data availability

All data used in this study were downloaded from public database and sources were listed in Supplementary Tables. The data supporting the findings of this study are available from the corresponding authors upon reasonable request.

Acknowledgements

This work was supported by Key Program of National Natural Science Foundation of China (NSFC) (Grant No. 32330085 to G.Y.Y.), Program of NSFC (Grant No. 32202376 to X.H.Y.; Grant No. 32302337 to X.X.Z.), the Young Elite Scientists Sponsorship Program by China

Association for Science and Technology (Grant No. 2022QNRC001 to

X.H.Y.), the China Postdoctoral Science Foundation (Grant No.

2022M722804 to X.X.Z.), the Scientific Research Startup Fund Project of

Zhejiang A&F University (Grant No. 2023LFR119 to X.H.Y.), the

Fundamental Research Funds for Central Universities (Grant No.

2021FZZX001-31 to G.Y.Y.).

Author contributions

X.H.Y. and G.Y.Y. conceived and designed the study. X.X.Z., Y.Y.L.,

C.H., Y.Y., and X.H.Y. participated in comparative genomic analysis,

gene selection, genome rearrangement. K.C.C., X.X.Z., and X.H.Y.

performed VgR protein structure predictions. H.W.L. and Q.F.

participated in data collection and discussion. X.X.Z. and X.H.Y. wrote

the manuscript with input from all of the authors. All authors read and

approved the manuscript.

Competing interests

686 The authors declare no competing interests.

References

1.   Ellers, J., Toby Kiers, E., Currie, C. R., McDonald, B. R. & Visser, B. Ecological

interactions drive evolutionary loss of traits. *Ecol. Lett.* **15**, 1071–1082 (2012).

2.   Lahti, D. C. *et al.* Relaxed selection in the wild. *Trends Ecol. Evol.* **24**, 487–496

(2009).

3.   Sifuentes-Romero, I. *et al.* Trait loss in evolution: What cavefish have taught us

about mechanisms underlying eye regression. *Integr. Comp. Biol.* **63**, 393–406 (2023).

4.   Rétaux, S. & Casane, D. Evolution of eye development in the darkness of caves:

Adaptation, drift, or both? *EvoDevo* **4**, 26 (2013).

5.   Leys, R., Cooper, S. J. B., Strecker, U. & Wilkens, H. Regressive evolution of an

eye pigment gene in independently evolved eyeless subterranean diving beetles. *Biol.*

Lett. **1**, 496–499 (2005).

6.   Protas, M. E., Trontelj, P. & Patel, N. H. Genetic basis of eye and pigment loss in

the cave crustacean, *Asellus aquaticus*. *Proc. Natl. Acad. Sci. U.S.A.* **108**, 5702–5707

(2011).

7.   Emerling, C. A. & Springer, M. S. Eyes underground: Regression of visual

protein networks in subterranean mammals. *Mol. Phylogenet. Evol.* **78**, 260–270

(2014).

8.   Protas, M. E. *et al.* Genetic analysis of cavefish reveals molecular convergence in

the evolution of albinism. *Nat. Genet.* **38**, 107–111 (2006).

9.   Louchart, A. & Viriot, L. From snout to beak: The loss of teeth in birds. *Trends*

Ecol. Evol. **26**, 663–673 (2011).

10. Sire, JY., Delgado, S. C. & Girondot, M. Hen's teeth with enamel cap: From

dream to impossibility. *BMC Evol. Biol.* **8**, 246 (2008).

11. Meredith, R. W., Zhang, G., Gilbert, M. T. P., Jarvis, E. D. & Springer, M. S.

Evidence for a single loss of mineralized teeth in the common avian ancestor. *Science*

346, 1254390 (2014).

12. Peng, C. *et al.* Large-scale snake genome analyses provide insights into

vertebrate development. *Cell* **186**, 3519 (2023).

13. Xia, B. *et al.* The genetic basis of tail-loss evolution in humans and apes. Preprint

at https://www.biorxiv.org/content/10.1101/2021.09.14.460388v1 (2021).

14. Poulin, R. & Randhawa, H. S. Evolution of parasitism along convergent lines:

From ecology to genomics. *Parasitology* **142**, S6–S15 (2015).

15. Visser, B. *et al.* Loss of lipid synthesis as an evolutionary consequence of a

parasitic lifestyle. *Proc. Natl. Acad. Sci. U.S.A.* **107**, 8677–8682 (2010).

16. Cai, L. *et al.* Deeply altered genome architecture in the endoparasitic flowering

plant *Sapria himalayana* Griff. (Rafflesiaceae). *Curr. Biol.* **31**, 1002−1011.e9 (2021).

17. Guo, X. *et al.* The *Sapria himalayana* genome provides new insights into the

lifestyle of endoparasitic plants. *BMC Biol.* **21**, 134 (2023).

18. Sun, G. *et al.* Large-scale gene losses underlie the genome evolution of parasitic

plant *Cuscuta australis*. *Nat. Commun.* **9**, 2683 (2018).

19. Chen, X. *et al. Balanophora* genomes display massively convergent evolution

with other extreme holoparasites and provide novel insights into parasite–host

interactions. *Nat. Plants* **9**, 1627–1642 (2023).

20. International Helminth Genomes Consortium. Comparative genomics of the

major parasitic worms. *Nat. Genet.* **51**, 163–174 (2019).

21. Tsai, I. *et al.* The genomes of four tapeworm species reveal adaptations to

parasitism. *Nature* **496**, 57–63 (2013).

22. Cunha, T. J., De Medeiros, B. A. S., Lord, A., Sørensen, M. V. & Giribet, G.

Rampant loss of universal metazoan genes revealed by a chromosome-level genome

assembly of the parasitic Nematomorpha. *Curr. Biol.* **33**, 3514–521.e4 (2023).

23. Pennacchio, F. & Strand, M. R. Evolution of developmental strategies in parasitic

Hymenoptera. *Annu. Rev. Entomol.* **51**, 233–258 (2006).

24. Werren, J. H. *et al.* Functional and evolutionary insights from the genomes of

three parasitoid *Nasonia* species. *Science* **327**, 343–348 (2010).

25. Xiao, JH. *et al.* Obligate mutualism within a host drives the extreme

specialization of a fig wasp genome. *Genome Biol.* **14**, R141 (2013).

26. Park, D. *et al.* Uncovering the novel characteristics of Asian honey bee, *Apis*

cerana, by whole genome sequencing. *BMC Genom.* **16**, 1 (2015).

27. Lindsey, A. R. I. *et al.* Comparative genomics of the miniature wasp and pest

control agent *Trichogramma pretiosum*. *BMC Biol.* **16**, 54 (2018).

28. Ye, X. *et al.* A chromosome-level genome assembly of the parasitoid wasp

Pteromalus puparum. *Mol. Ecol. Resour.* **20**, 1384–1402 (2020).

29. Xu, H. *et al.* Comparative genomics sheds light on the convergent evolution of

miniaturized wasps. *Mol. Biol. Evol.* **38**, 5539–5554 (2021).

30. Ye, X. *et al.* Genomic signatures associated with maintenance of genome stability

and venom turnover in two parasitoid wasps. *Nat. Commun.* **13**, 6417 (2022).

31. Yang, Y. *et al.* Genome of the pincer wasp *Gonatopus flavifemur* reveals unique

venom evolution and a dual adaptation to parasitism and predation. *BMC Biol.* **19**,

145 (2021).

32. Flanders, S. E. Oösorption and ovulation in relation to oviposition in the parasitic

Hymenoptera. *Ann. Entomol. Soc. Am.* **35**, 251–266 (1942).

33. Ferkovich, S. M. & Dillard, C. R. A study of uptake of radiolabeled host proteins

and protein synthesis during development of eggs of the endoparasitoid, *Microplitis*

croceipes (Cresson) (Braconidae). *Insect Biochem.* **16**, 337–345 (1986).

34. Ralec, A. Egg contents in relation to host-feeding in some parasitic Hymenoptera.

Entomophaga **40**, 87–93 (1995).

35. Grbic, M. & Strand, M. R. Shifts in the life history of parasitic wasps correlate

with pronounced alterations in early development. *Proc. Natl. Acad. Sci. U.S.A.* **95**,

1097–1101 (1998).

36. Donnell, D. M. Vitellogenin of the parasitoid wasp, *Encarsia formosa*

(Hymenoptera: Aphelinidae): Gene organization and differential use by members of

the genus. *Insect Biochem. Mol. Biol.* **34**, 951–961 (2004).

37. King, P. E., Ratcliffe, N. A. & Fordy, M. R. Oögenesis in a braconid, *Apanteles*

glomeratus (L.) possessing an hydropic type of egg. *Z. Zellforsch.* **119**, 43–57 (1971).

38. Tufail, M. & Takeda, M. Molecular characteristics of insect vitellogenins. *J.*

Insect Physiol. **54**, 1447–1458 (2008).

39. Fisher, R. C. Aspects of the physiology of endoparasitic Hymenoptera. *Biol. Rev.*

46, 243–278 (1971).

40. Raikhel, A. S. & Dhadialla, T. S. Accumulation of yolk proteins in insect oocytes.

Annu. Rev. Entomol. **37**, 217–51 (1992).

41. Wu, Z., Yang, L., He, Q. & Zhou, S. Regulatory mechanisms of vitellogenesis in

insects. *Front. Cell Dev. Biol.* **8**, 593613 (2021).

42.   Pennacchio, F., Caccia, S. & Digilio, M. C. Host regulation and nutritional

exploitation by parasitic wasps. *Curr. Opin. Insect. Sci.* **6**, 74–79 (2014).

43.   Caccia, S. *et al.* Nutrient absorption by *Aphidius ervi* larvae. *J. Insect Physiol.* **51**,

1183–1192 (2005).

44.   Anderson, T., Levitt, D. & Banaszak, L. The structural basis of lipid interactions

in lipovitellin, a soluble lipoprotein. *Structure* **6**, 895–909 (1998).

45.   Du, X. *et al.* Functional characterization of Vitellogenin_N domain, domain of

unknown function 1943, and von Willebrand factor type D domain in vitellogenin of

the non-bilaterian coral *Euphyllia ancora*: Implications for emergence of immune

activity of vitellogenin in basal metazoan. *Dev. Comp. Immunol.* **67**, 485–494 (2017).

46.   Albalat, R. & Cañestro, C. Evolution by gene loss. *Nat. Rev. Genet.* **17**, 379–391

(2016).

47.   Koludarov, I. *et al.* Domain loss enabled evolution of novel functions in the snake

three-finger toxin gene superfamily. *Nat. Commun.* **14**, 4861 (2023).

48.   Sharma, V., Hecker, N., Walther, F., Stuckas, H. & Hiller, M. Convergent losses

of *TLR5* suggest altered extracellular flagellin detection in four mammalian lineages.

Mol. Biol. Evol. **37**, 1847–1854 (2020).

49.   Schneider, W. J. Vitellogenin receptors: Oocyte-specific members of the low-

density lipoprotein receptor supergene family. *Int. Rev. Cytol.* **166**, 103–137 (1996).

50.   Sappington, T. W. & S. Raikhel, A. Molecular characteristics of insect

vitellogenins and vitellogenin receptors. *Insect Biochem. Mol. Biol.* **28**, 277–300

(1998).

51.   Wertheim, J. O., Murrell, B., Smith, M. D., Kosakovsky Pond, S. L. & Scheffler,

K. RELAX: Detecting relaxed selection in a phylogenetic framework. *Mol. Biol.*

Evol. **32**, 820–832 (2015).

52.   Kosakovsky Pond, S. L. *et al.* HyPhy 2.5—A customizable platform for

evolutionary hypothesis testing using phylogenies. *Mol. Biol. Evol.* **37**, 295–299

(2020).

53. Lin, Y. *et al.* Vitellogenin receptor mutation leads to the oogenesis mutant

phenotype "scanty vitellin" of the silkworm, *Bombyx mori*. *J. Biol. Chem.* **288**,

13345–13355 (2013).

54. Davis, C. G. *et al.* Acid-dependent ligand dissociation and recycling of LDL

receptor mediated by growth factor homology region. *Nature* **326**, 760–765 (1987).

55. Sheng, Y. *et al.* The vitellogenin receptor gene contributes to mating and host-

searching behaviors in parasitoid wasps. *iScience* **26**, 106298 (2023).

56. Sharanowski, B. J. *et al.* Phylogenomics of Ichneumonoidea (Hymenoptera) and

implications for evolution of mode of parasitism and viral endogenization. *Mol.*

Phylogenet. Evol. **156**, 107023 (2021).

57. Locher, K. P. Mechanistic diversity in ATP-binding cassette (ABC) transporters.

Nat. Struct. Mol. Biol. **23**, 487–493 (2016).

58. Chen, J.-S., Sappington, T. W. & Raikhel, A. S. Extensive sequence conservation

among insect, nematode, and vertebrate vitellogenins reveals ancient common

ancestry. *J. Mol. Evol.* **44**, 440–451 (1997).

59. Leipart, V. *et al.* Identification of 121 variants of honey bee vitellogenin protein

sequences with structural differences at functional sites. *Protein Sci.* **31**, e4369

(2022).

60. Leipart, V. *et al.* Resolving the zinc binding capacity of honey bee vitellogenin

and locating its putative binding sites. *Insect Mol. Biol.* **31**, 810–820 (2022).

61. Leipart, V. *et al.* Structure prediction of honey bee vitellogenin: A multi-domain

protein important for insect immunity. *FEBS Open Bio* **12**, 51–70 (2022).

62. Tufail, M., Nagaba, Y., Elgendy, A. M. & Takeda, M. Regulation of vitellogenin

genes in insects. *Entomol. Sci.* **17**, 268–282(2014)

63. Salmela, H. *et al.* Ancient duplications have led to functional divergence of

vitellogenin-like genes potentially involved in inflammation and oxidative stress in

honey bees. *Genome Biol. Evol.* **8**, 495–506 (2016).

64. Morandin, C. *et al.* Not only for egg yolk—functional and evolutionary insights

from expression, selection, and structural analyses of *Formica* ant vitellogenins. *Mol.*

Biol. Evol. **31**, 2181–2193 (2014).

65. Dam, M. H. V. *et al.* The easter egg weevil (*Pachyrhynchus*) genome reveals

syntenic patterns in Coleoptera across 200 million years of evolution. *PLoS Genet.* **17**,

e1009745 (2021).

66. Thomas, G. W. C. *et al.* Gene content evolution in the arthropods. *Genome Biol.*

21, 15 (2020).

67. Aravind, L., Watanabe, H., Lipman, D. J. & Koonin, E. V. Lineage-specific loss

and divergence of functionally linked genes in eukaryotes. *Proc. Natl. Acad. Sci.*

U.S.A. **97**, 11319–11324 (2000).

68. Blomme, T. *et al.* The gain and loss of genes during 600 million years of

vertebrate evolution. *Genome Biol.* **7**, R43 (2006).

69. Koonin, E. V. *et al.* A comprehensive evolutionary classification of proteins

encoded in complete eukaryotic genomes. *Genome Biol.* **5**, R7 (2004).

70. Springer, M. S. *et al.* Genomic and anatomical comparisons of skin support

independent adaptation to life in water by cetaceans and hippos. *Curr. Biol.* **31**, 2124–

2139.e3 (2021).

71. Sharma, V. *et al.* A genomics approach reveals insights into the importance of

gene losses for mammalian adaptations. *Nat. Commun.* **9**, 1215 (2018).

72. Dong, SZ. *et al.* Vitellin of *Pteromalus puparum* (Hymenoptera: Pteromalidae), a

pupal endoparasitoid of *Pieris rapae* (Lepidoptera: Pieridae): Biochemical

characterization, temporal patterns of production and degradation. *J. Insect Physiol.*

53, 468–477 (2007).

73. Guo, JY. *et al.* Oosorption in the endoparasitoid, *Pteromalus puparum*. *J. of*

Insect Sci. **11**, 90 (2011).

74. Ye, GY. *et al.* Molecular cloning and developmental expression of the

vitellogenin gene in the endoparasitoid, *Pteromalus puparum*. *Insect Mol. Biol.* **17**,

227–233 (2008).

75. Dong, SZ., Ye, GY., Guo, JY., Yu, XP. & Hu, C. Oogenesis and programmed cell

death of nurse cells in the endoparasitoid, *Pteromalus puparum*. *Microsc. Res. Tech.*

73, 673–680 (2010).

76. Heimpel, G. E. & Collier, T. R. The evolution of host-feeding behaviour in insect

parasitoids. *Biol. Rev.* **71**, 373–400 (1996).

77. Brawand, D., Wahli, W. & Kaessmann, H. Loss of egg yolk genes in mammals

and the origin of lactation and placentation. *PLoS Biol.* **6**, e63 (2008).

78. Van Rompay, L., Borghgraef, C., Beets, I., Caers, J. & Temmerman, L. New

genetic regulators question relevance of abundant yolk protein production in *C.*

elegans. *Sci. Rep.* **5**, 16381 (2015).

79. Sayers, E. W. *et al.* Database resources of the national center for biotechnology

information. *Nucleic Acids Res.* **50**, D20–D26 (2021).

80. Mei, Y. *et al.* InsectBase 2.0: A comprehensive gene resource for insects. *Nucleic*

Acids Res. **50**, D1040–D1045 (2022).

81. The Darwin Tree of Life Project Consortium. Sequence locally, think globally:

The Darwin tree of life project. *Proc. Natl. Acad. Sci. U.S.A.* **119**, e2115642118

(2022).

82. Vizueta, J., Sánchez-Gracia, A. & Rozas, J. Bitacora: A comprehensive tool for

the identification and annotation of gene families in genome assemblies. *Mol. Ecol.*

Resour. **20**, 1445–1452 (2020).

83. Camacho, C. *et al.* BLAST+: Architecture and applications. *BMC Bioinform.* **10**,

421 (2009).

84. Eddy, S. R. Accelerated profile HMM searches. *PLoS Comput. Biols* **7**, e1002195

(2011).

85. Keilwagen, J., Hartung, F., Grau, J. GeMoMa: Homology-based gene prediction

utilizing intron position conservation and RNA-seq data. In *Gene Prediction:*

Methods and Protocols, Methods in Molecular Biology. vol. 1962, Humana, New

York, NY. (ed Kollmar, M.). 161−177 (Springer Science+Business Media, 2019).

86. The UniProt Consortium. UniProt: The universal protein knowledgebase in 2023.

Nucleic Acids Res. **51**, D523–D531 (2023).

87. Stanke, M. & Morgenstern, B. AUGUSTUS: A web server for gene prediction in

eukaryotes that allows user-defined constraints. *Nucleic Acids Res.* **33**, W465–W467

(2005).

88. Salamov, A. A. & Solovyev, V. V. *Ab initio* gene finding in *Drosophila* genomic

DNA. *Genome Res.* **10**, 516–522 (2000).

89. Dobin, A. *et al.* STAR: Ultrafast universal RNA-seq aligner. *Bioinformatics* **29**,

15–21 (2013).

90. Li, B. & Dewey, C. N. RSEM: Accurate transcript quantification from RNA-Seq

data with or without a reference genome. *BMC Bioinform.* **12**, 323 (2011).

91. Tang, H. *et al.* Synteny and collinearity in plant genomes. *Science* **320**, 486–488

(2008).

92. Kiełbasa, S. M., Wan, R., Sato, K., Horton, P. & Frith, M. C. Adaptive seeds tame

genomic sequence comparison. *Genome Res.* **21**, 487–493 (2011).

93. Harris, R. S. Improved pairwise alignment of genomic DNA. (Pennsylvania State

University, 2007).

94. Liao, Y., Zhang, X., Chakraborty, M. & Emerson, J. J. Topologically associating

domains and their role in the evolution of genome structure and function in

Drosophila. *Genome Res.* **31**, 397–410 (2021).

95. Quinlan, A. R. & Hall, I. M. BEDTools: A flexible suite of utilities for comparing

genomic features. *Bioinformatics* **26**, 841–842 (2010).

96. Sun, Y. H. *et al.* Amniotes co-opt intrinsic genetic instability to protect germline

genome integrity. *Nat. Commun.* **14**, 812 (2023).

97. Smith, M. D. *et al.* Less is more: An adaptive branch-site random effects model

for efficient detection of episodic diversifying selection. *Mol. Biol. Evol.* **32**, 1342–

1353 (2015).

98. Emms, D. M. & Kelly, S. OrthoFinder: Phylogenetic orthology inference for

comparative genomics. *Genome Biol.* **20**, 238 (2019).

99. Emms, D. M. & Kelly, S. STAG: Species tree inference from all genes. Preprint

at https://www.biorxiv.org/content/10.1101/267914v1 (2018).

100. Katoh, K. & Standley, D. M. MAFFT multiple sequence alignment software

version 7: Improvements in performance and usability. *Mol. Biol. Evol.* **30**, 772–780

(2013).

101. Minh, B. Q. *et al.* IQ-TREE 2: New models and efficient methods for

phylogenetic inference in the genomic era. *Mol. Biol. Evol.* **37**, 1530–1534 (2020).

102. Sanderson, M. J. r8s: Inferring absolute rates of molecular evolution and

divergence times in the absence of a molecular clock. *Bioinformatics* **19**, 301–302

(2003).

103. Peters, R. S. *et al.* Evolutionary history of the Hymenoptera. *Curr. Biol.* **27**,

1013–1018 (2017).

104. Mendes, F. K., Vanderpool, D., Fulton, B. & Hahn, M. W. CAFE 5 models

variation in evolutionary rates among gene families. *Bioinformatics* **36**, 5516–5518

(2021).

105. Cantalapiedra, C. P., Hernández-Plaza, A., Letunic, I., Bork, P. & Huerta-

Cepas, J. EggNOG-mapper v2: Functional annotation, orthology assignments, and

domain prediction at the metagenomic scale. *Mol. Biol. Evol.* **38**, 5825–5829 (2021).

106. Klopfenstein, D. V. *et al.* GOATOOLS: A python library for gene ontology

analyses. *Sci. Rep.* **8**, 10872 (2018).

107. Supek, F., Bošnjak, M., Škunca, N. & Šmuc, T. REVIGO summarizes and

visualizes long lists of gene ontology terms. *PLoS ONE* **6**, e21800 (2011).

108. Balart-García, P. *et al*. Parallel and convergent genomic changes underlie

independent subterranean colonization across beetles. *Nat. Commun.* **14**, 3842 (2023).

109. Aristide, L. & Fernández, R. Genomic innovation, convergence and

parallelism facilitated the ecological transition to life out of the sea in molluscs.

Preprint at https://doi.org/10.21203/rs.3.rs-2382076/v1 (2022).

110. Mirdita, M. *et al.* ColabFold: Making protein folding accessible to all. *Nat.*

Methods **19**, 679–682 (2022).

111. Jumper, J. *et al.* Highly accurate protein structure prediction with AlphaFold.

Nature **596**, 583–589 (2021).

112. Steinegger, M. & Söding, J. MMseqs2 enables sensitive protein sequence

searching for the analysis of massive data sets. *Nat. Biotechnol.* **35**, 1026–1028

(2017).

113. Meng, E. C., Pettersen, E. F., Couch, G. S., Huang, C. C. & Ferrin, T. E. Tools for

integrated sequence-structure analysis with UCSF Chimera. *BMC Bioinform.* **7**, 339

(2006).

114. Meng, E. C. *et al.* UCSF ChimeraX: Tools for structure building and analysis.

Protein Sci. **32**, e4792 (2023).

Figures

Fig. 1 | Evolutionary analyses of vitellogenin (*Vg*) and *Vg*-related genes. (a)

Domain schematics of Vg and homologs, namely Partial Vg (PVg) and Vg-like (Vgl)

proteins. Vg proteins had three major domains: Vitellogenin_N (LPD_N) (green),

Vit_open_b-sht (purple), and von Willebrand factor type D (vWD) (gray). Other Vg-

related proteins were classified as PVgs or Vgls based on the presence or absence,

respectively, of the Vitellogenin_N domain. SP, signal peptide. (**b**) Phylogenic

analysis of hymenopteran insects. The tree suggested the existence of at least five

independent *Vg* loss events. Divergence times were estimated using 12 calibration

points. The colored solid circle on each tree node indicates the bootstrap value. Stars

to the left of the species name indicate the parasitic strategy of that species,

endoparasitism (yellow) or ectoparasitism (white). Question marks to the left of the

species name indicate an unknown parasitic strategy. The numbers of *Vg* and *Vg*-

related genes in each genome are indicated at right; the pink question mark for *Vg* in

Asobara japonica indicates ambiguity. This loss may have been an artifact from

genome assembly (Supplementary Fig. 2).

Fig. 2 | Phylogenetic and gene expression analyses of *Vg*s and *PVg*s. (a)
Phylogenetic tree of *Vg*s (red) and *PVg*s (blue) in hymenopteran species. All of the
*Vg*s clustered together with some of the *PVg*s (~48%); the remainder of the *PVg*s
clustered together as a monophyletic group. (**b**) Expression patterns of *Vg*s and *PVg*s
in male and female adults of 10 hymenopteran species. Columns indicate individual
samples; each row represents expression of a *Vg* or *PVg* (as noted at right).
Expression values are represented with color and were calculated in fragments per
kilobase of transcript per million mapped reads (FPKM). For experiments without
biological replicates, gene expression values are shown in numerical form. The
phylogeny of the 10 selected hymenopteran insects was extracted from the data
shown in **Fig. 1**.

Fig. 3 | *Vg* was likely lost from endoparasitoid Hymenoptera due to genome
rearrangement. (**a–c**) Synteny of *Vg* and the flanking genes among representative
Vg-loss and *Vg*-retaining species within the (**a**) family Ichneumonidae, (**b**) family
Braconidae, and (**c**) superfamily Cynipoidea. *Vg*s are indicated with red. Grey lines
indicate orthologous relationships. (**d**) Phylogenetic analyses of three independent
lineages with *Vg* loss. Group1, Cynipoidea; Group2, Chalcidoidea; Group3,
Braconidae. Pairwise genome alignments were performed within each group to
identify genomic rearrangement breakpoints. Species shown in pink and purple were
Vg-retaining (reference) and *Vg*-loss (query) species, respectively. (**e**) Distance
between *Vg* loci and observed genomic rearrangement breakpoints (orange points) or
randomly shuffled control bins (purple violins).

Fig. 4 | Relaxed selection and complete loss of genes encoding the Vg receptor (*VgR*) in many lineages with *Vg* loss. (**a**) Phylogenetic tree (from **Fig. 1b**) showing instances of *Vg* and *VgR* loss (pink and purple arrows, respectively). Branches shown in green were used in molecular selection analyses. (**b–f**) Selection signatures of *VgR*s from independent analyses of several *Vg*-loss branches or clades, namely (**b**) all *Vg*-loss branches or clades, excluding *Asobara japonica*; (**c**) *Leptopilina* spp.; (**d**) *Trichogramma* spp.; (**e**) *Hyposoter didymator*; and (**f**) *A. japonica*. Nonsynonymous to synonymous substitution rate (dN/dS) distributions across sites in the *VgR*s are shown for test (branches listed above in **b–f**, light purple) and reference (branches left from test branches, dark grey) branches. dN/dS values of 1, > 1, and < 1 represent neutral evolution, positive selection, and purifying selection, respectively. Arrows show the direction of selective pressure between the test and reference branches. K values < 1 or > 1 indicate relaxed or intensified selection, respectively. The Bonferroni-Holm corrected *p*-values were showed.

1017

Fig. 5 | Predicted three-dimensional structures of vitellogenin receptors (VgRs).
(**a**) Domain schematic of a classic VgR protein. The ligand binding domains (LBD1
and LBD2) and the EGF precursor domains (EGF precursor1 and EGF precursor2)
are indicated at left. Three regions with well-defined structures based on the
confidence (pLDDT) scores were designated regions A, B, and C (labeled at right).
(**b–d**) Predicted structures of the VgR regions A, B, and C with AlphaFold2. Each
structure is a superimposed representation based on 45 VgR sequences. pLDDT
scores for each residue are indicated with color. (**e**) A superimposed structure of
region A based on 43 VgRs (excluding *Leptopilina boulardi* and *L. heterotoma*)
showing the highly conserved structure of the second EGF-like repeat. (**f**) Structural
comparison showing variations in the second EGF-like repeat of region A in *L.
boulardi* and *L. heterotoma*. (**g**) Multiple sequence alignment of the second EGF-like
repeat of region A from all 45 VgRs, demonstrating the sequence variations in *L.
boulardi* and *L. heterotoma*.

1032

Fig. 6 | Parallel or convergent gene family evolution in the *Vg*-loss lineages. (a)
Expansion and contraction of gene families, also known as Orthogroups (OGs) in
branches leading to *Vg*-loss lineages. The phylogenetic tree was taken from **Fig. 1b**.
The five branches affected by *Vg* loss events are indicated with dashed lines and
marked a–e. (**b, c**) Unique and overlapping (**b**) expanded and (**c**) contracted OGs
between the five independent *Vg*-loss branches. (**d, e**) Functional convergence of (**d**)
expanded and (**e**) contracted OGs in the five independent *Vg*-loss branches (see
Methods). Each point represents an expanded or contracted OG shared by four or five
of the independent *Vg*-loss lineages. The centroid OGs were labeled with putative
functions in each polygon. Gene Ontology terms associated with transport,
development, and metabolism were functionally converged.

1044

Fig. 7 | Schematic of the evolution of *Vg* and *VgR* in parasitoid wasps. The
ancestral parasitoid wasp is thought to have been an ectoparasitoid that retained both
the *Vg* and *VgR* genes and laid yolk-rich eggs. Hymenopteran evolution involved
several independent shifts from ectoparasitism to endoparasitism. Some
endoparasitoid wasps (e.g., *Pteromalus puparum*) retain *Vg* and *VgR* genes and lay
yolk-rich eggs. Others have lost *Vg*, likely through genome rearrangement, leading to
loss or relaxed selection of *VgR* genes due to disruption of vitellogenesis. These
wasps often produce yolkless eggs (**Supplementary Table 1**). There were several
shared genomic changes related to transport, metabolism, development, sensory
systems, and stress resistance among the five independent *Vg*-loss lineages; these may
have been related to endoparasitoid adaptation and/or contributed to loss of the yolk
as an available nutrient source to the early embryo.

1057 **Supplementary Figures**

Species	Chimeric genes ID	Fixed genes	
Amblyteles armatorius	ENSAAYP00000011660	*AarmVg1*	*AarmVg2*
Anisopteromalus calandrae	Acal20629-TA	*AcalVg1*	*AcalVg2*
Ooceraea biroi	XP_019887836.2	*ObirVg1*	*ObirVg2*
Pachycrepoideus vindemmiae	Pvin01909-TA	*PvinVg1*	*PvinVg2*
Solenopsis invicta	XP_039305184.1	*SinvVg1*	*SinvVg2*
Trichomalopsis sarcophagae	OXU21803.1	*TsarVg1*	*TsarVg2*
Diadromus collaris	DCO15731-TA	*DcolPVg1*	*DcolPVg2*

1058

1059 **Supplementary Fig. 1 Schematic of fixation of chimeric genes.** For genes joined
1060 together but owned repeat domain structures of *Vg* were split into two genes which
1061 composed domains of *Vg* or *PVg*. These genes were re-annotate using Augusts or
1062 FGENESH+ webserver, and resulted genes were further validated from using BLAST
1063 and HMMER.

1064

Supplementary Fig. 2 Domain structure and location of *PVg* in *Asobara japonica*.
PVg (*Ajap023976*) of *A. japonica* was located at the start terminal on the scaffold
JADHZF10000379.1 from scanning 3,043bp while the LPD_N domain predicted
using SMART webserver was covered whole protein-coding sequences from 22 to
631 site with 1 to 16 was signal peptide. However, whether this *PVg* was a result of
genome misassemble or bad annotation still unknown.

1071

Supplementary Fig. 3. Gene synteny of *Vg*s and flanking genes among representative *Vg*-loss species and *Vg*-retaining wasps in Chalcidoidea.
Pteromalus puparum as the *Vg*-retaining species was showed in the middle and two *Vg*s were highlighted in red, while *Trichogramma brassicae* and *T. pretiosum* were showed above and below. Results showed the two *Vg*s in *Pteromalus puparum* have no homologous genes in this syntenic block region.

1078

Supplementary Fig. 4. Enrichment visualization of 199 gene under relaxed selection using REVIGO. The enrichment results generated from GOATOOLS were used as input, and value C was pre-set as 0.5 which corresponding to the "small" list size, then REVIGO will clustered listed GO terms and reduce redundancy of them. Based on the information in Supplementary Table 10, dispensability of GO terms less than 0.2 were labelled in the figure and some interested GO terms were marked in pink.