An atlas of genetic variation for linking pathogen-induced cellular traits to human disease

Authors:

Liuyang Wang[1], Kelly J. Pittman[1], Jeffrey R. Barker[1], Raul E. Salinas[1], Ian B. Stanaway[2], Graham D. Williams[1], Robert J. Carroll[3], Tom Balmat[4], Andy Ingham[5], Anusha M. Gopalakrishnan[1], Kyle D. Gibbs[1], Alejandro L. Antonia[1], The eMERGE Network, Joseph Heitman[1,6], Soo Chan Lee[7], Gail P. Jarvick[8], Joshua C. Denny[3], Stacy M. Horner[1,6], Mark R. DeLong[5], Raphael H. Valdivia[1], David R. Crosslin[2], and Dennis C. Ko[1,6,9*]

Affiliations

[1]Department of Molecular Genetics and Microbiology, School of Medicine, Duke University, Durham, NC 27710

[2]Department of Biomedical Informatics and Department of Medicine, School of Medicine, University of Washington, Seattle, WA 98195

[3]Department of Biomedical Informatics, School of Medicine, Vanderbilt University, Nashville, TN 37212

[4]Social Science Research Institute, Duke University, Durham, NC 27710

[5]Duke Research Computing, Duke University, Durham, NC 27710

[6]Division of Infectious Diseases, Department of Medicine, School of Medicine, Duke University, Durham, NC 27710

25 [7]South Texas Center for Emerging Infectious Diseases (STCEID), Department of Biology, College of Sciences, the University of Texas at San Antonio, San Antonio, TX 78249

[8]Department of Medicine, Division of Medical Genetics, School of Medicine, University of Washington, Seattle, WA 98195

[9]Lead contact

30 *To whom correspondence should be addressed: Dennis C. Ko, 0049 CARL Building Box 3053, 213 Research Drive, Durham, NC 27710. 919-684-5834. dennis.ko@duke.edu. @denniskoHiHOST

35 **Summary**

Genome-wide association studies (GWAS) have identified thousands of genetic variants associated with disease. To facilitate moving from associations to disease mechanisms, we leveraged the role of pathogens in shaping human evolution with the Hi-HOST Phenome Project (H2P2): a catalog of cellular GWAS comprised of 79 phenotypes in response to 8 pathogens in

40 528 lymphoblastoid cell lines. Seventeen loci surpass genome-wide significance ($p<5x10^{-8}$) for phenotypes ranging from pathogen replication to cytokine production. Combining H2P2 with clinical association data from the eMERGE Network and experimental validation revealed evidence for mechanisms of action and connections with diseases. We identified a SNP near *CXCL10* as a cis-cytokine-QTL and a new risk factor for inflammatory bowel disease. A SNP in

45 *ZBTB20* demonstrated pleiotropy, partially mediated through NFκB signaling, and was associated with viral hepatitis. Data are available in an H2P2 web portal to facilitate further interpreting human genome variation through the lens of cell biology.

Keywords: genome-wide association study, lymphoblastoid cell line, immune quantitative trait

50 locus, heritability, pleiotropy, phewas, electronic medical record, CXCL10, ZBTB20, eldelumab

Introduction

The human genome has been shaped by migration, drift, admixture, and natural selection (*1-3*). One of the strongest driving forces in natural selection has been pathogens (*4, 5*), as first exemplified with A. C. Allison's demonstration that sickle cell allele (rs334) conferred resistance to malaria (*6*). Red blood cells from individuals with this allele are resistant to *Plasmodium* infection (*7*). Similarly, human resistance to HIV infections afforded by the *CCR5Δ32* allele can also be seen at the level of individual T cells (*8-10*). Therefore, identification and characterization of human genetic differences that impact cellular traits can help mechanistically link human genetic variation to disease susceptibility.

Previous studies have examined the genetic basis of molecular traits in human populations. Expression quantitative trait loci (eQTL) studies in lymphoblastoid cell lines (LCLs) defined abundant associations between human SNPs and expression levels of nearby genes (*11, 12*). LCLs are EBV-transformed B cells that are highly similar to antigen-activated primary B cells (*13*). LCLs serve as a standardized resource for functional human genetic variation studies, as they have been densely genotyped by the International HapMap Project (*14, 15*) and the 1000 Genomes Project (*16*). As eQTLs are often shared across tissues (e.g. 88% of cis-eQTLs are shared among LCLs, fibroblasts, and primary T cells (*17*)), LCL eQTL studies have led to important insights not only in immunity-related diseases but also for disorders where B cells are not believed to be primary drivers of disease (*18*).

Using LCLs, we developed Hi-HOST (High-throughput Human in vitrO Susceptibility Testing) to identify human genetic differences in pathogen-induced cellular traits, serving as a cell biological link between eQTL studies and GWAS of human disease (*19, 20*). Hi-HOST uses live pathogens to examine variation in innate immune recognition, but also in cell biological and

75   signaling processes that can be quantified as phenotypes for genome-wide association. This work

therefore builds on a long tradition of using cellular microbiology to elucidate basic cell biology

(*21*) and expands that utility to interpret the human genome. Using Hi-HOST, we leveraged LCL

responses to *Salmonella enterica* to demonstrate that genetic variation in the methionine salvage

pathway regulates pyroptosis, as well as human susceptibility to sepsis (*22, 23*). Similarly, we

80   recently reported that a genetic variant in *VAC14* is associated with both increased *S.* Typhi

invasion into LCLs and risk of typhoid fever in a Vietnamese population (*24*).

Here, we present the Hi-HOST Phenome Project (H2P2) to globally explore the genetic

basis of cellular outcomes in response to infectious agents. Using 7 microbes and 1 bacterial

toxin, we carried out GWAS of 79 host-pathogen phenotypes that serve as cellular readouts for

85   processes such as endocytosis, endosomal trafficking, signal transduction, cell death, and

transcriptional regulation. We identified 17 loci that reached the generally accepted threshold for

genome-wide significance ($p<5x10^{-8}$). We integrated H2P2 data with experimental validation

and disease association data from the eMERGE Network PheWAS pipeline (*25*) to define new

functions for genes in disease and provide new clues to disease pathophysiology.

90   **Results**

Phenotypic variation in H2P2 traits reveals biologically meaningful clusters

We measured variation in cellular traits in 528 LCLs stimulated with 7 different microbes

and 1 bacterial toxin (Figure 1A). These LCLs have been genotyped (*16*) and consist of 4 human

populations: ESN (Esan in Nigeria), GWD (Gambians in Western Divisions in the Gambia), IBS

95   (Iberian Population in Spain), and KHV (Kinh in Ho Chi Minh City, Vietnam). The LCLs used

in this study were all from parent-offspring trios, allowing for heritability estimation by parent-

offspring regression and for carrying out combined GWAS analysis with protection from stratification through family-based association methods (*26, 27*).

100   The microbes and toxin we selected affect billions of people worldwide. Non-typhoidal *Salmonella* infections caused 150 million diarrheal illnesses (gastroenteritis) and 0.6 million cases of invasive enteric disease (bacteremia) in 2010 (*28, 29*). Approximately 20 million cases of typhoid fever are caused by *Salmonella enterica* serovar Typhi (*S.* Typhi) every year (*30*). *Chlamydia trachomatis,* the most common bacterial sexually transmitted infection in the world, causes 100 million cases of genital tract infection every year (*31*), and 1.3 million individuals are

105   blind due to ocular infection with *C. trachomatis* (*32*). *Staphylococcus aureus* is a common cause of skin and soft tissue infections, bacteremia, and infective endocarditis, and its alpha toxin, which is utilized in H2P2, is a key virulence determinant (*33*). *Candida albicans* and related fungal species are a frequent cause of genitourinary tract infection that can cause more severe disease in immunocompromised individuals and are the fifth most common cause of hospital-

110   associated infections (*34*). *Mucor circinelloides* is another fungal species that causes severe infections, mucormycosis, in immunocompromised individuals (reviewed in (*35*)) and has also recently been connected to food-borne infections (*36*). Finally, over 6 billion people in the world have been infected with the protozoal pathogen *Toxoplasma gondii*, which can be fatal in infants and immunocompromised individuals (*37*). Thus the microbes and toxin used in H2P2 are

115   important causes of human disease.

Beyond their role in human disease, the pathogens selected also exploit a wide range of host cellular processes to either kill the host cell or to create replicative niches within them. *C. trachomatis, S. enterica* serovar Typhi, serovar Typhimurium (wildtype and Δ*sifA* mutant, which escapes from the pathogen-containing vacuole at a greater rate into the host-cell cytosol (*38*)),

120   and *T. gondii* are intracellular pathogens that employ diverse lifestyles. These microbes were

engineered to express the green fluorescent protein (GFP) to allow quantitation of pathogen

invasion, survival and replication, intercellular spread, and concurrent measurement of cell death

by flow cytometry. B cells are known *in vivo* host cells for *Salmonella* invasion and replication

(*39-41*) and the *C. trachomatis* strain that we utilize (L2) causes lymphogranuloma venereum

125   (*42*), primarily an infection of the lymph nodes, where there is ample opportunity for interaction

with B cells. Cell death was also measured as the readout for *Staphylococcus aureus* alpha toxin,

a pore-forming toxin that causes cell lysis (*33*). Microbes were also tested for induction or

suppression of cytokines (selected based on pilot studies of 41 cytokines) in infected LCLs. *S.*

Typhimurium and *C. trachomatis*-infected cells were measured for 3 and 17 cytokines,

130   respectively. The fungal pathogens *M. circinelloides* and *C. albicans* were included for their

ability to induce FGF-2 in LCLs (*43*). Definitions for all 79 phenotypes as well as histograms for

phenotypes are provided in the supplemental data (Table S1 and Figure S1). Importantly, 76 of

79 H2P2 phenotypes showed significant experimental repeatability based on measurements on

LCLs from three different passages (Figure S2).

135   Hierarchical clustering of traits based on inter-individual variation confirmed the

robustness of our measurements (Figure 1B). Levels of three cytokines (CXCL10 (IP-10), IL-10,

and MDC) measured in uninfected cells with two different methods (ELISA at 24hrs and

Luminex at 70hrs) showed strong correlation (R = 0.78 for CXCL10, R= 0.46 for IL-10, R=0.90

for MDC; Spearman correlation). The clustering of responses to *Salmonella* infection is

140   consistent with previous findings: *S.* Typhimurium, Typhi, and Typhimurium Δ*sifA* cluster for

the phenotype of invasion, as all utilize a similar type-3 secretion system for entry (*44*)

(correlation to *S.* Typhimurium, R=0.69 for *S.* Typhi and R=0.90 for *S.* Typhimurium Δ*sifA*). In

contrast, the correlation is weaker for intracellular survival and replication phenotypes (correlation to *S.* Typhimurium replication from 3.5 hrs to 24 hrs, R=0.36 for *S.* Typhi and

145 R=0.34 for *S.* Typhimurium Δ*sifA*), reflecting the different replications niches for Δ*sifA* (host cell cytoplasm) and wild-type Typhimurium (membrane bound vacuole) (*38*) or the use of a different repertoire of effectors by *S.* Typhi (*45*). In contrast, we observed almost no correlation between EBV copy number (from (*46*) for 284 LCLs also used in H2P2) and H2P2 traits (Figure S3; Table S2), indicating these phenotypes are not being driven by the LCL immortalization

150 method. Thus, clustering based on phenotypic diversity verified the reliability of measurements and confirmed biological relatedness established by previous work from multiple groups.

H2P2 traits are heritable based on parent-offspring regression and SNP-based heritability

If genetic differences regulate variation in cellular phenotypes, then the additive contributions of these differences to phenotypic variance can be estimated. We measured

155 narrow-sense heritability (h^2) with two independent and complementary methods: parent-offspring regression and SNP-based h^2. Heritability based on parent-offspring regression is estimated as the slope of the regression line for offspring phenotypes vs. mid-parent phenotypes (*47*). We observed h^2 estimates from -0.06 to 0.85 (average h^2=0.33), with the largest h^2 observed for *S.* Typhimurium-induced levels of IL-10 (h^2=0.85±0.19, p=3.5x10^{-15}) (Figure 1C;

160 Figure S4; Table S3). The majority (64/79) of phenotypes showed significantly non-zero h^2 by this method (p<0.05). In contrast, SNP-based h^2, as implemented in the GCTA software package (*48*) with the Zaitlen modification for related individuals (*49*), calculates the proportion of variance that can be explained by all genotyped SNPs. With this method, we observed pedigree h^2 ranging from 0.04 to 0.76 (average h^2=0.36), and SNP-based h^2 ranging from 0.02 to 0.66

165 (average h^2=0.19) (Figure 1C; Table S3).

We observed high correlation between h^2 estimated using the two different methods (R=0.58; p=2.9x10^{-8}; Figure 1D). The strong correlation between the two estimates of h^2, based on distinct statistical frameworks, provides additional evidence that LCLs provide a robust system to identify human SNPs that contribute to the heritability of cellular phenotypes.

170   *H2P2 reveals 17 genome-wide significant associations and enrichment for genic SNPs and regions of active chromatin*

We performed a family-based GWAS in PLINK (*26, 27*) on 79 traits for 528 LCLs using dense genotyping information (15.5 million SNPs after imputation; see Methods). Across 79 traits, we observed 17 loci that reached a genome-wide significance threshold of p<5x10^{-8}

175   (Figure 2A; Table 1; Figure S5). H2P2 demonstrated enrichment of associated SNPs for functional genome annotations. We used the GARFIELD package to calculate and visualize fold-enrichment of SNPs associated in H2P2 at variable p-value thresholds with different genomic features (*50*). In regards to SNP location, the greatest enrichment was observed for exonic SNPs (Figure 2B). The fold-enrichment was highest at the most stringent p-value

180   threshold (p<1x10^{-8}) for H2P2 traits (9.2-fold enrichment; p=0.10 by Fisher's exact test) and was statistically significant when using a p<5x10^{-8} threshold for H2P2 traits (3.9 fold-enrichment; p=0.047). In contrast, there was depletion for intergenic SNPs at the p<5x10^{-8} threshold (0.66 fold-enrichment). H2P2-associated SNPs showed even greater enrichment for regions of active chromatin as annotated by DNase hypersensitivity peaks from the ENCODE project (*51*) (Figure

185   2C). Consistent with H2P2 being conducted in LCLs, the 2nd greatest enrichment was observed for DNAase hypersensitivity peaks measured in the LCL GM06990 out of 424 cell types measured (at p<5x10^{-8}, 5.7 fold-enrichment; p=3.8x10^{-3}). Even stronger enrichment was noted at this threshold for human Th2 cells (7.3 fold-enrichment; p=3.5x10^{-4}), and enrichment was

observed for cells derived from most tissues (Figure 2C), consistent with LCLs being a relevant

190   model for genetic analysis for multiple human cell types.

A large-effect cis-regulatory variant regulates CXCL10 levels

The strongest association in the H2P2 dataset was observed for rs2869462 with levels of

the chemokine CXCL10 (also known as IP-10) following *Chlamydia* infection (p=$2x10^{-9}$; Figure

2D, E). This SNP is located 7.5kb 3' of the *CXCL10* coding sequence (Figure 2D) in a region

195   that also encodes for two related chemokines, CXCL9 and CXCL11. These chemokines bind to

CXC-chemokine receptor 3 (CXCR3), a G-protein coupled receptor that mediates inflammation

by coordinating T-helper 1 (Th1) recruitment and activating effector cells during infection and

autoimmunity (reviewed in (*52*)). Many cell types produce CXCL10 during infection, including

activated B cells (*53-56*). Notably, the effect of this SNP is large: rs2869462 accounts for 13.7%

200   of the variance in CXCL10 protein levels. While no dataset is available to replicate this

association at the protein level, this SNP also demonstrated an association with *CXCL10* mRNA

(p=$7x10^{-7}$) in an independent set of 465 uninfected LCLs ((*57*); none of the LCLs overlap the

H2P2 LCLs) (Figure S6).

Although rs2869462 was identified using genome-wide association of all 528 LCLs in

205   our dataset, there were large differences in allele frequencies among the populations. The derived

allele of rs2869462 (G) is present at the highest frequencies in Europe (28% in IBS) and Asia

(28% in KHV) and is substantially lower in Africa (1.5% in ESN, 1% in GWD; Figure 2F).

Strikingly, all populations demonstrated the same directionality of effect for the rs2869462 allele

on CXCL10 levels (C > G) (Figure 2G).

210   *SNPs lead to pleiotropic effects on multiple pathogen-induced traits*

Different pathogens can target common signaling pathways to establish an intracellular niche or to modulate immune responses. Therefore, we examined whether genome-wide significant hits were associated with just one trait or if they were associated with multiple traits, using either a threshold of $p<0.05$ or with Bonferroni multiple-test correction ($p<6.3 \times 10^{-4}$). All

215   of the 17 genome-wide significant hits were associated with at least 4 H2P2 traits with the $p<0.05$ threshold (Figure 3A). However, several phenotypes are closely related and therefore these cross-phenotype associations do not reflect true pleiotropy, multiple unrelated effects due to the same gene ([58]). For example, rs2869462 had 5 cross-phenotype associations but 4 of these traits are based on CXCL10 levels (Figure 3B). Beyond the existence of pleiotropy, the pattern

220   of which traits shared genetic associations provided additional insight (Figure 3C). A circle plot of cross-phenotype associations showed most cross-phenotype associations in H2P2 connect invasion, establishment of an intracellular niche, and intercellular spread. Traits that had high phenotypic correlation (Figure 1B) were more likely to have cross-phenotypic associations as expected (Figure 3D).

225   The greatest number of associated traits at $p<0.05$ occurred for rs953897, a SNP in the gene encoding the transcriptional repressor ZBTB20. Based on GTEx data, rs953897 is associated with *ZBTB20-AS1* transcript abundance ($p=1 \times 10^{-9}$) ([59]). While rs953897 is most strongly associated with high *C. trachomatis* burden at 46 hrs ($p = 1.3 \times 10^{-8}$), this SNP was associated with 20 H2P2 traits ($p<0.05$) and 5 traits using a multiple-test corrected threshold of

230    $p<6.3 \times 10^{-4}$. A QQ plot comparing p-values for all traits for rs953897 confirmed the high degree of pleiotropy for this SNP, demonstrating strong deviation from neutrality towards lower p-values (Figure 3D). These associations even included other pathogens and biological processes, as the third most strongly associated trait was *S*. Typhimurium-induced pyroptosis ($p=7.5 \times 10^{-5}$;

Figure 3D). Overall, out findings indicate that genetic variation in *ZBTB20* is linked to pleiotropic effects on multiple host-pathogen traits.

Genetic variants impacting ZBTB20 expression affect the outcome of Chlamydia and Salmonella infections

The T allele of rs953897 was associated with both higher levels of *Chlamydia* replication (Figure 4A) and *Salmonella*-induced pyroptosis (Figure 4B). Reduction of *ZBTB20* expression by RNAi increased *Chlamydia* replication and pyroptosis, mimicking the effect of the T allele (Figure 4C, D). RNAi-mediated depletion of ZBTB20 also indicated that some phenotypes that did not reach statistically significant associations with rs953897 after multiple-test correction were nonetheless mediated by ZBT20. Specifically, expression of IL-6 after infection with *Chlamydia* (p=0.03), was *reduced* after *ZTB20* RNAi treatment, again mimicking the effect of the T allele (Figure 4E). The effect of ZBT20 depletion on IL-6 expression is not the result of decreased bacterial burden since *Chlamydia* replication was enhanced under these conditions. Thus, the association data and functional validation point to a role for ZBTB20 in the regulation of multiple infection-related phenotypes.

ZBTB20 has been characterized as a transcriptional repressor during prenatal development in the liver and the brain (*60-62*). Rare protein-coding mutations in *ZBTB20* are responsible for Primrose syndrome, which has features as disparate as mental retardation, ossified external ears, and distal muscle wasting (*63*). We hypothesized that one ZBTB20 target gene could regulate a pathway that impacted multiple biological processes related to pathogen immunity. An attractive target, IκB (*NFKBIA*), the canonical suppressor of NFκB signaling, is subject to ZBTB20 transcriptional repression (*64*). Reduction of ZBTB20 expression in LCLs by RNAi causes increased expression of IκB (Figure 4F). An increase of IκB should cause

inhibition of NFκB signaling, resulting in increased *Chlamydia* replication but decreased

expression of pro-inflammatory cytokines including IL-6. Consistent with this prediction,

depletion of IκB by RNAi decreased *Chlamydia* replication and increased IL-6 production

260   (Figure 4C, E). In contrast, depletion of IκB did not impact *Salmonella*-induced pyroptosis,

indicating an IκB-independent mechanism (Figure 4D). These observations point to multiple

roles for ZBTB20 in regulating cellular functions during infection, both through suppression of

regulators of signaling pathways, such as NFκB (Figure 4J), but also through regulation of other

unidentified targets.

265   *SNPs linked to CXCL10 expression are also associated with inflammatory bowel disease*

We determined if SNPs associated with cellular traits in H2P2 were associated with

human disease. Consistent with the rs2869462 C allele being associated with increased CXCL10

and inflammation, we discovered that this allele is a previously unrecognized inflammatory

bowel disease (IBD) risk allele. CXCL10 inhibitory antibodies have undergone phase II clinical

270   trials for both sub-types of IBD, Crohn's disease (CD) and ulcerative colitis (UC) (*65, 66*), based

on evidence from animal models (*67-69*) and of elevated CXCL10 in patients with CD (*70*) and

UC (*71*). Comparison to GWAS summary statistics from the IBD Genetics Consortium meta-

analysis of 12882 IBD cases and 21770 controls (*72*) demonstrated that rs2869462 is associated

with IBD (p=1.7x10^{-4}; OR=1.08), as well as with CD (p=1.9x10^{-3}; OR=1.09) and UC subtypes

275   (p=0.016; OR=1.06). The cases in the subtype analysis are exclusive of one another and

therefore demonstrate an association of rs2869462 with two different cohorts of IBD cases

(though most controls are shared). Furthermore, the direction of association is consistent with

high levels of CXCL10 (the C allele) being associated with greater risk of IBD.

The availability of GWAS summary statistics from this dataset allowed us to conduct a

280   formal colocalization analysis to determine whether the CXCL10 protein level signal was the

same as the IBD signal. We utilized the COLOC package which uses a Bayesian framework to

determine whether GWAS signals in the same region are likely due to the same causal variant

(*73*). The posterior probability that both CXCL10 protein level and IBD share the same causal

variant is as high as 0.80 for the region, with rs2869462 being singled out as the SNP with the

285   greatest posterior probability as being causal (Table S4). Comparison of regional plots of

association along with the colocalization analysis indicate that the same LD block is associated

with both CXCL10 levels and risk of IBD (Figure 5A).

We independently tested for this association using electronic medical record (EMR) data

from the eMERGE Network (*74*). The eMERGE dataset holds genotype-phenotype correlations

290   of >80,000 individuals with phenotypes assigned based on ICD-9 patient billing codes (*25*). We

tested for association with the codes for "inflammatory bowel disease and other gastroenteritis

and colitis" and the more restrictive code for "ulcerative colitis." rs2869462 was associated with

both phenotypes in the predicted direction (p=0.003; OR=1.12, C allele for IBD, p=0.02;

OR=1.12, C allele for UC) (Figure 5B). Therefore by first identifying a SNP as associated with

295   CXCL10 levels following an LCL model of *C. trachomatis* infection, we have discovered and

replicated a new IBD risk allele.

H2P2 SNPs are associated with disease in PheWAS of electronic medical record traits

Next, we systematically generated hypotheses regarding the effects of the H2P2 genome-

wide significant hits on human disease by employing a PheWAS (phenome-wide association

300   study) approach, looking for associations across a large set (1338) of clinical measurements and

diseases cataloged in eMERGE (*25*). We found 5 of 16 H2P2 genome-wide significant hits

surpassed the multiple test corrected significance threshold with at least 1 EMR phenotype (Figure 6A; Table S5; 1 H2P2 SNP was not in the eMERGE dataset and had no good proxy based on LD).

305   The SNP that had the greatest number of PheWAS associations was rs7566597. This SNP, associated with the H2P2 phenotype of *Chlamydia*-infected levels of the chemokine MIP-1β (macrophage inflammatory protein-1beta; CCL4), was associated with five clinical traits. The most significant association (p $=1.76 \times 10^{-6}$; p=0.0024 after Bonferroni; OR=1.40) was observed with otorrhea (Figure 6B). Otorrhea is ear drainage most commonly caused by an ear infection.

310   Notably, MIP-1β is elevated in fluid from patients with middle ear infections (*75*), mice with genetic predisposition to middle ear infections (*76*), and primary middle ear epithelial cultures infected *in vitro* with either influenza or *Streptococcus pneumoniae* (*77*). The directionality of this association (G allele associated with both higher MIP-1β levels and increased risk of otorrhea), and the fact that genotype is fixed prior to disease, lead to the hypothesis that the

315   rs7566597 G allele causes higher levels of MIP-1β to increase risk of otorrhea.

    Similarly, rs953897 in *ZBTB20* was associated with viral hepatitis (Figure 6C; p=3.17×10^{-5}; p=0.025 after Bonferroni; OR=1.20). Most of the 1654 viral hepatitis cases in this cohort are due to hepatitis C virus (HCV). To further test this association experimentally, we performed RNAi against *ZBTB20* in Huh7 human hepatocytes. Depletion of *ZBTB20* mRNA

320   (Figure 6D) increased the percentage of HCV infected cells over time (Figure 6E) and increased the amount of infectious virus by 7-fold (Figure 6F). Future mechanistic and clinical studies will be required to further validate the association and determine how genetic variation in *ZBTB20* affects risk of viral hepatitis. However, this example demonstrates that coupling H2P2 with

PheWAS of clinical traits can lead to hypotheses that can be quickly tested in the most clinically

325   relevant cell type for that particular disease.

Discussion

With H2P2, we have coupled the ability of pathogens to influence genetic diversity in

human populations to their use as cellular probes to elucidate mechanisms of disease. This

cellular GWAS approach provides: 1) biomolecules and proteins that could serve as possible

330   biomarkers and therapeutic targets and 2) a cellular model to validate and dissect mechanisms of

how the genetic variant regulates the cell biological process now connected to disease. We have

developed an H2P2 database and web interface to allow for exploration of this rich dataset by the

research community (http://h2p2.oit.duke.edu).

Cellular GWAS studies have also been performed on inter-individual variation in levels

335   of immune cell subtypes, cell surface protein expression, and cytokine levels (*78-80*). However,

Hi-HOST is unique in using live pathogens to induce complex cellular phenotypes, such as cell

death and invasion, providing surrogate phenotypes that are intermediate between molecular

phenotypes of gene/protein expression and human population studies of disease. Additionally,

H2P2 utilizes cells derived from multiple human populations. Notably, allele frequencies for

340   rs2869462 and rs953897 vary greatly across the globe. These differences may play an important

role in susceptibility of different populations in cellular responses and disease. Finally, our use of

parent-offspring trios allowed us to make estimates of h^2 through both parent-offspring

regression and SNP-based methods.

Our h^2 estimates are consistent with a large fraction of variation in cellular responses as

345   being genetically determined. For parent-offspring regression 64 of 79 traits had significantly

non-zero h^2 (p<0.05). While relatively small sample size led to large standard errors for SNP-based estimates of h^2, the two estimates correlated quite strongly (r=0.58; p=$2.9x10^{-8}$). Our estimates of h^2 are similar to other reports for immune-related traits. Orru *et al.* examined levels of 95 immune cell types and found a mean h^2 of 41% (range 3-87%) (*79*). Our estimates of h^2 for

350   cytokine levels as well as for more complex host-pathogen phenotypes such as *Chlamydia* intracellular replication (46hr % high GFP; 33% by parent-offspring regression and 20% by SNP-based heritability) are consistent with a strong genetic basic for variation in immune cell traits and host-pathogen interactions, though environment also has a large effect. Furthermore, our ability to identify SNPs strongly associated with these traits that can explain a sizable portion

355   of the h^2 further confirms the importance of common SNPs in contributing to the h^2 of molecular and cellular pathogen-induced traits.

Integrating H2P2 with human genetic association data from published studies and the eMERGE Network revealed how genetic variants impacting cellular traits also influenced human disease. For rs2869462, we found that this SNP associated with CXCL10 levels is also a

360   previously unrecognized risk factor for IBD. While over a hundred IBD risk alleles have been identified (*72*), the fact rs2869462 was associated with levels of CXCL10 in H2P2 may make genotyping of this SNP clinically actionable if coupled to anti-CXCL10 therapy. While anti-CXCL10 demonstrated some benefit in phase 2 trials, neither study met statistical significance for its primary endpoint (*65, 66*). We hypothesize that rs2869462 genotype might be a predictive

365   biomarker for identifying the genetic subtype of patients who show the greatest benefit. This example, spanning molecular phenotype, human disease, and clinical utility serves as a template for how we envision the H2P2 web portal being used to make similar discoveries.

For *CXCL10*, *ZBTB20*, and other genes implicated by H2P2, there are numerous associations that do not reach genome-wide significance but are undoubtedly true-positives based on highly related phenotypes or experimental evidence. Indeed, our lab has previously pursued non-genome-wide significant hits revealed by Hi-HOST, resulting in a new metabolite biomarker for sepsis (*23*) and a new potential therapeutic strategy for typhoid fever (*24*). However, to fully illuminate how genetic variation contributes to the pathophysiology of disease will require the engagement of the research community with the H2P2 web portal and other similar datasets. These users, already experts on particular genes and/or cellular pathways, would be well-equipped to then validate and discover the mechanisms underlying these associations. Thus, H2P2 provides a hypothesis-generating engine for identifying new biomarkers and therapeutic strategies. Future studies will expand the panel of stimuli and the cell types used to create a more complete picture of how cellular traits impact human health and disease, an important step towards a future of more personalized care.

Author Contributions

All authors critically reviewed the manuscript and contributed input to the final submission. DCK, LW, KJP, ALA, IBS, GDW, TB, SMH, RHV, and DRC wrote the manuscript. DCK, LW, KJP, ALA, JH, SCL, SMH, RHV, and DRC contributed to strategy and project planning. KJP,

385 DCK, JRB, RES, GDW, AMG, KDG, and SCL carried out experiments and analysis. LW, IBS, RJC, tEM, GPJ, JCD, DRC, and DCK planned and carried out computational analysis and provided datasets. TB, AI, LW, MRD, and DCK designed and implemented the H2P2 database and web portal.

Competing interests: Duke University has submitted a provisional patent application ("A

390 Companion Diagnostic for IBD Therapy") on behalf of DCK, LW, and ALA.

Acknowledgments: Funding: LW, KJP, RES, KDG, ALA, YC, and DCK were supported by NIH R01AI118903. LW, KJP, JRB, RES, AMG, RHV, and DCK were supported by the Ecopathogenomics of Sexually Transmitted Infections (EPSTI) Cooperative Research Center (NIH U19AI084044). DCK was also supported by a Duke University Whitehead Scholarship,

395 the Butler Pioneer Award, and a Duke MGM Pilot Award. GDW and SMH were supported by NIH R01AI125416 and R21AI124100 and the Burroughs Wellcome Fund. IBS, RJC, JCD, and DRC were supported by eMERGE Network (Phase III). This phase of the eMERGE Network was initiated and funded by the NHGRI through the following grants: U01HG8657 (Group Health Cooperative/University of Washington); U01HG8685 (Brigham and Women's Hospital);

400 U01HG8672 (Vanderbilt University Medical Center); U01HG8666 (Cincinnati Children's Hospital Medical Center); U01HG6379 (Mayo Clinic); U01HG8679 (Geisinger Clinic); U01HG8680 (Columbia University Health Sciences); U01HG8684 (Children's Hospital of Philadelphia); U01HG8673 (Northwestern University); U01HG8701 (Vanderbilt University

Medical Center serving as the Coordinating Center); U01HG8676 (Partners Healthcare/Broad Institute); and U01HG8664 (Baylor College of Medicine). U01HG004438 (CIDR) and U01HG004424 (the Broad Institute) served as eMERGE Genotyping Centers. SCL was supported by NIH/NIAID R03 AI11917 and UTSA Research funds. JH was supported by NIH/NIAID R37 MERIT Award AI39115-20, NIH/NIAID R01 AI50113-13, and an award from Astellas. TB, AI, and MRD were supported by Duke Research Computing. We thank Dr. Yanlu Cao for performing FGF-2 ELISAs. We thank Luke C. Glover for help in plotting association data. We thank Dr. Gregory D. Sempowski and the Duke Immune Reconstitution & Biomarker Analysis Shared Resource in performing pilot Luminex measurements. We thank Dr. Charles Rice for the HCV NS5A antibody and the Duke Functional Genomics Shared Resource for use of the Cellomics ArrayScan. We thank Drs. Shelton Bradrick, David Tobin, Ana-Maria Xet-Mull, Andrew Alspaugh, and Jennifer L. Trenor for pilot studies in characterizing infection of various pathogens into LCLs. DNA image from Figure 1 is from the National Human Genome Research Institute Image Gallery. The content of this manuscript is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or other funding sources.

References

1. L. L. Cavalli-Sforza, P. Menozzi, A. Piazza, *The history and geography of human genes*. (Princeton University Press, Princeton, N.J., 1994), pp. xi, 541, 518 p.

2. J. Z. Li *et al.*, Worldwide human relationships inferred from genome-wide patterns of variation. *Science* **319**, 1100-1104 (2008).

3. J. K. Pickrell, D. Reich, Toward a new history and geography of human genes informed by ancient DNA. *Trends Genet* **30**, 377-389 (2014).

4. M. Fumagalli *et al.*, Signatures of environmental genetic adaptation pinpoint pathogens as the main selective pressure through human evolution. *PLoS Genet* **7**, e1002355 (2011).

5. K. J. Pittman, L. C. Glover, L. Wang, D. C. Ko, The Legacy of Past Pandemics: Common Human Mutations That Protect against Infectious Disease. *PLoS Pathog* **12**, e1005680 (2016).

6. A. C. Allison, Protection afforded by sickle-cell trait against subtertian malareal infection. *Br Med J* **1**, 290-294 (1954).

7. M. J. Friedman, Erythrocytic mechanism of sickle cell resistance to malaria. *Proc Natl Acad Sci U S A* **75**, 1994-1997 (1978).

8. M. Dean *et al.*, Genetic restriction of HIV-1 infection and progression to AIDS by a deletion allele of the CKR5 structural gene. Hemophilia Growth and Development Study, Multicenter AIDS Cohort Study, Multicenter Hemophilia Cohort Study, San Francisco City Cohort, ALIVE Study. *Science* **273**, 1856-1862 (1996).

9. R. Liu *et al.*, Homozygous defect in HIV-1 coreceptor accounts for resistance of some multiply-exposed individuals to HIV-1 infection. *Cell* **86**, 367-377 (1996).

10. M. Samson *et al.*, Resistance to HIV-1 infection in caucasian individuals bearing mutant alleles of the CCR-5 chemokine receptor gene. *Nature* **382**, 722-725 (1996).

11. B. E. Stranger *et al.*, Patterns of cis regulatory variation in diverse human populations. *PLoS Genet* **8**, e1002639 (2012).

12. B. E. Stranger *et al.*, Population genomics of human gene expression. *Nat Genet* **39**, 1217-1224 (2007).

13. E. D. Cahir-McFarland *et al.*, Role of NF-kappa B in cell survival and transcription of latent membrane protein 1-expressing or Epstein-Barr virus latency III-infected cells. *Journal of virology* **78**, 4108-4119 (2004).

14. I. H. Consortium, A haplotype map of the human genome. *Nature* **437**, 1299-1320 (2005).

15. C. International HapMap *et al.*, Integrating common and rare genetic variation in diverse human populations. *Nature* **467**, 52-58 (2010).

16. G. P. Consortium, A map of human genome variation from population-scale sequencing. *Nature* **467**, 1061-1073 (2010).

17. T. Flutre, X. Wen, J. Pritchard, M. Stephens, A statistical framework for joint eQTL analysis in multiple tissues. *PLoS Genet* **9**, e1003486 (2013).

18. A. C. Nica, E. T. Dermitzakis, Expression quantitative trait loci: present and future. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences* **368**, 20120362 (2013).

19. D. C. Ko *et al.*, A genome-wide in vitro bacterial-infection screen reveals human variation in the host response associated with inflammatory disease. *Am J Hum Genet* **85**, 214-227 (2009).

20. D. C. Ko, T. J. Urban, Understanding Human Variation in Infectious Disease Susceptibility through Clinical and Cellular GWAS. *PLoS Pathog* **9**, e1003424 (2013).

21. P. Cossart, P. Boquet, S. Normark, R. Rappuoli, Cellular microbiology emerging. *Science* **271**, 315-316 (1996).

470 22. D. C. Ko *et al.*, Functional genetic screen of human diversity reveals that a methionine salvage enzyme regulates inflammatory cell death. *Proc Natl Acad Sci U S A* **109**, E2343-2352 (2012).

23. L. Wang *et al.*, Human genetic and metabolite variation reveal methylthioadenosine is a prognostic biomarker and inflammatory regulator in sepsis. *Science Advances* **3**, 475 e1602096 (2017).

24. M. I. Alvarez *et al.*, Human genetic variation in VAC14 regulates Salmonella invasion and typhoid fever through modulation of cholesterol. *Proc Natl Acad Sci U S A* **114**, E7746-E7755 (2017).

25. J. C. Denny *et al.*, Systematic comparison of phenome-wide association study of 480 electronic medical record data and genome-wide association study data. *Nature biotechnology* **31**, 1102-1110 (2013).

26. S. Purcell *et al.*, PLINK: a tool set for whole-genome association and population-based linkage analyses. *Am J Hum Genet* **81**, 559-575 (2007).

27. S. Purcell, P. Sham, M. J. Daly, Parental phenotypes in family-based association analysis. 485 *Am J Hum Genet* **76**, 249-259 (2005).

28. M. D. Kirk *et al.*, World Health Organization Estimates of the Global and Regional Disease Burden of 22 Foodborne Bacterial, Protozoal, and Viral Diseases, 2010: A Data Synthesis. *PLoS Med* **12**, e1001921 (2015).

29. S. E. Majowicz *et al.*, The global burden of nontyphoidal Salmonella gastroenteritis. 490 *Clinical infectious diseases : an official publication of the Infectious Diseases Society of America* **50**, 882-889 (2010).

30. G. Dougan, S. Baker, Salmonella enterica serovar Typhi and the pathogenesis of typhoid fever. *Annual review of microbiology* **68**, 317-336 (2014).

31. L. Newman *et al.*, Global Estimates of the Prevalence and Incidence of Four Curable 495 Sexually Transmitted Infections in 2012 Based on Systematic Review and Global Reporting. *PLoS One* **10**, e0143304 (2015).

32. M. J. Burton, D. C. Mabey, The global burden of trachoma: a review. *PLoS Negl Trop Dis* **3**, e460 (2009).

33. S. Y. Tong, J. S. Davis, E. Eichenberger, T. L. Holland, V. G. Fowler, Jr., 500 Staphylococcus aureus infections: epidemiology, pathophysiology, clinical manifestations, and management. *Clin Microbiol Rev* **28**, 603-661 (2015).

34. N. Yapar, Epidemiology and risk factors for invasive candidiasis. *Ther Clin Risk Manag* **10**, 95-105 (2014).

35. L. Mendoza *et al.*, Human Fungal Pathogens of Mucorales and Entomophthorales. *Cold* 505 *Spring Harb Perspect Med* **5**, (2014).

36. S. C. Lee *et al.*, Analysis of a food-borne fungal pathogen outbreak: virulence and genome of a Mucor circinelloides isolate from yogurt. *MBio* **5**, e01390-01314 (2014).

37. J. M. Furtado, J. R. Smith, R. Belfort, Jr., D. Gattey, K. L. Winthrop, Toxoplasmosis: a global threat. *J Glob Infect Dis* **3**, 281-284 (2011).

510 38. C. R. Beuzon *et al.*, Salmonella maintains the integrity of its intracellular vacuole through the action of SifA. *EMBO J* **19**, 3235-3249 (2000).

39. D. Castro-Eguiluz *et al.*, B cell precursors are targets for Salmonella infection. *Microb Pathog* **47**, 52-56 (2009).

40. R. Rosales-Reyes, C. Alpuche-Aranda, L. Ramirez-Aguilar Mde, A. D. Castro-Eguiluz, V. Ortiz-Navarrete, Survival of Salmonella enterica serovar Typhimurium within late endosomal-lysosomal compartments of B lymphocytes is associated with the inability to use the vacuolar alternative major histocompatibility complex class I antigen-processing pathway. *Infect Immun* **73**, 3937-3944 (2005).

41. R. Rosales-Reyes *et al.*, Salmonella infects B cells by macropinocytosis and formation of spacious phagosomes but does not induce pyroptosis in favor of its survival. *Microb Pathog* **52**, 367-374 (2012).

42. R. Ceovic, S. J. Gulin, Lymphogranuloma venereum: diagnostic and treatment challenges. *Infect Drug Resist* **8**, 39-47 (2015).

43. S. C. Lee *et al.*, Calcineurin orchestrates dimorphic transitions, antifungal drug responses and host-pathogen interactions of the pathogenic mucoralean fungus Mucor circinelloides. *Molecular microbiology* **97**, 844-865 (2015).

44. C. M. Collazo, J. E. Galan, The invasion-associated type-III protein secretion system in Salmonella - A review. *Gene* **192**, 51-59 (1997).

45. J. Parkhill *et al.*, Complete genome sequence of a multiple drug resistant Salmonella enterica serovar Typhi CT18. *Nature* **413**, 848-852 (2001).

46. R. Mandage *et al.*, Genetic factors affecting EBV copy number in lymphoblastoid cell lines derived from the 1000 Genome Project samples. *PLoS One* **12**, e0179446 (2017).

47. D. S. Falconer, T. F. C. Mackay, *Introduction to Quantitative Genetics*. (Longmans Green, Harlow, Essex, U.K., ed. 4, 1996).

48. J. Yang, S. H. Lee, M. E. Goddard, P. M. Visscher, GCTA: a tool for genome-wide complex trait analysis. *Am J Hum Genet* **88**, 76-82 (2011).

49. N. Zaitlen *et al.*, Using extended genealogy to estimate components of heritability for 23 quantitative and dichotomous traits. *PLoS Genet* **9**, e1003520 (2013).

50. V. Iotchkova *et al.*, GARFIELD--GWAS Analysis of Regulatory or Functional Information Enrichment with LD correction. *bioRxiv* **doi: 10.1101/085738** (2016).

51. E. P. Consortium *et al.*, An integrated encyclopedia of DNA elements in the human genome. *Nature* **489**, 57-74 (2012).

52. J. R. Groom, A. D. Luster, CXCR3 ligands: redundant, collaborative and antagonistic functions. *Immunol Cell Biol* **89**, 207-215 (2011).

53. C. Hennig *et al.*, High-content cytometry and transcriptomic biomarker profiling of human B-cell activation. *J Allergy Clin Immunol* **133**, 172-180 e171-110 (2014).

54. S. T. Hoff *et al.*, Human B cells produce chemokine CXCL10 in the presence of Mycobacterium tuberculosis specific T cells. *Tuberculosis (Edinb)* **95**, 40-47 (2015).

55. A. Kato *et al.*, CpG oligodeoxynucleotides directly induce CXCR3 chemokines in human B cells. *Biochemical and biophysical research communications* **320**, 1139-1147 (2004).

56. J. Vollmer *et al.*, CpG oligodeoxynucleotides stimulate IFN-gamma-inducible protein-10 production in human B cells. *J Endotoxin Res* **10**, 431-438 (2004).

57. T. Lappalainen *et al.*, Transcriptome and genome sequencing uncovers functional variation in humans. *Nature* **501**, 506-511 (2013).

58. N. Solovieff, C. Cotsapas, P. H. Lee, S. M. Purcell, J. W. Smoller, Pleiotropy in complex traits: challenges and strategies. *Nature reviews. Genetics* **14**, 483-495 (2013).

59. G. T. Consortium, Human genomics. The Genotype-Tissue Expression (GTEx) pilot analysis: multitissue gene regulation in humans. *Science* **348**, 648-660 (2015).

60. Z. Xie *et al.*, Zinc finger protein ZBTB20 is a key repressor of alpha-fetoprotein gene transcription in liver. *Proc Natl Acad Sci U S A* **105**, 10859-10864 (2008).

61. C. Mitchelmore *et al.*, Characterization of two novel nuclear BTB/POZ domain zinc finger isoforms. Association with differentiation of hippocampal neurons, cerebellar granule cells, and macroglia. *J Biol Chem* **277**, 7598-7609 (2002).

62. J. V. Nielsen, F. H. Nielsen, R. Ismail, J. Noraberg, N. A. Jensen, Hippocampus-like corticoneurogenesis induced by two isoforms of the BTB-zinc finger gene Zbtb20 in mice. *Development* **134**, 1133-1140 (2007).

63. V. Cordeddu *et al.*, Mutations in ZBTB20 cause Primrose syndrome. *Nat Genet* **46**, 815-817 (2014).

64. X. Liu *et al.*, Zinc finger protein ZBTB20 promotes Toll-like receptor-triggered innate immune responses by repressing IkappaBalpha gene transcription. *Proc Natl Acad Sci U S A* **110**, 11097-11102 (2013).

65. W. J. Sandborn *et al.*, Eldelumab [anti-interferon-gamma-inducible protein-10 antibody] Induction Therapy for Active Crohn's Disease: a Randomised, Double-blind, Placebo-controlled Phase IIa Study. *J Crohns Colitis*, (2017).

66. W. J. Sandborn *et al.*, Eldelumab [Anti-IP-10] Induction Therapy for Ulcerative Colitis: A Randomised, Placebo-Controlled, Phase 2b Study. *J Crohns Colitis* **10**, 418-428 (2016).

67. J. G. Hyun *et al.*, Anti-interferon-inducible chemokine, CXCL10, reduces colitis by impairing T helper-1 induction and recruitment in mice. *Inflammatory bowel diseases* **11**, 799-805 (2005).

68. S. Sasaki *et al.*, Blockade of CXCL10 protects mice from acute colitis and enhances crypt cell survival. *Eur J Immunol* **32**, 3197-3205 (2002).

69. U. P. Singh, S. Singh, D. D. Taub, J. W. Lillard, Jr., Inhibition of IFN-gamma-inducible protein-10 abrogates colitis in IL-10-/- mice. *J Immunol* **171**, 1401-1406 (2003).

70. A. E. Ostvik *et al.*, Enhanced expression of CXCL10 in inflammatory bowel disease: potential role of mucosal Toll-like receptor 3 stimulation. *Inflammatory bowel diseases* **19**, 265-274 (2013).

71. M. Uguccioni *et al.*, Increased expression of IP-10, IL-8, MCP-1, and MCP-3 in ulcerative colitis. *Am J Pathol* **155**, 331-336 (1999).

72. J. Z. Liu *et al.*, Association analyses identify 38 susceptibility loci for inflammatory bowel disease and highlight shared genetic risk across populations. *Nat Genet* **47**, 979-986 (2015).

73. C. Giambartolomei *et al.*, Bayesian test for colocalisation between pairs of genetic association studies using summary statistics. *PLoS Genet* **10**, e1004383 (2014).

74. C. A. McCarty *et al.*, The eMERGE Network: a consortium of biorepositories linked to electronic medical records data for conducting genomic studies. *BMC Med Genomics* **4**, 13 (2011).

75. R. Kaur, J. Casey, M. Pichichero, Cytokine, chemokine, and Toll-like receptor expression in middle ear fluids of children with acute otitis media. *Laryngoscope* **125**, E39-44 (2015).

76. F. Han *et al.*, Mutation in Phex gene predisposes BALB/c-Phex(Hyp-Duk)/Y mice to otitis media. *PLoS One* **7**, e43010 (2012).

77.   H. H. Tong, J. P. Long, P. A. Shannon, T. F. DeMaria, Expression of cytokine and chemokine genes by human middle ear epithelial cells induced by influenza A virus and Streptococcus pneumoniae opacity variants. *Infect Immun* **71**, 4289-4296 (2003).

78.   Y. Li *et al.*, A Functional Genomics Approach to Understand Variation in Cytokine Production in Humans. *Cell* **167**, 1099-1110 e1014 (2016).

79.   V. Orru *et al.*, Genetic Variants Regulating Immune Cell Levels in Health and Disease. *Cell* **155**, 242-256 (2013).

80.   M. Roederer *et al.*, The genetic architecture of the human immune system: a bioresource for autoimmunity and disease pathogenesis. *Cell* **161**, 387-403 (2015).

81.   J. H. Marcus, J. Novembre, Visualizing the geography of genetic variants. *Bioinformatics* **33**, 594-595 (2017).

82.   R. J. Bastidas, R. H. Valdivia, Emancipating Chlamydia: Advances in the Genetic Manipulation of a Recalcitrant Intracellular Pathogen. *Microbiol Mol Biol Rev* **80**, 411-427 (2016).

83.   M. I. G. Roncero, Enrichment method for the isolation of auxotrophic mutants of Mucor using the polyene antibiotic N-glycosyl-polifungin. *Carlsberg Research Communications* **49**, 685 (1984).

84.   F. C. Odds, A. J. Brown, N. A. Gow, Candida albicans genome sequence: a platform for genomics in the absence of genetics. *Genome biology* **5**, 230 (2004).

85.   J. P. Saeij, J. P. Boyle, M. E. Grigg, G. Arrizabalaga, J. C. Boothroyd, Bioluminescence imaging of Toxoplasma gondii infection in living mice reveals dramatic differences between strains. *Infect Immun* **73**, 695-702 (2005).

86.   C. C. Chang *et al.*, Second-generation PLINK: rising to the challenge of larger and richer datasets. *Gigascience* **4**, 7 (2015).

87.   R. J. Pruim *et al.*, LocusZoom: regional visualization of genome-wide association scan results. *Bioinformatics* **26**, 2336-2337 (2010).

88.   B. N. Howie, P. Donnelly, J. Marchini, A flexible and accurate genotype imputation method for the next generation of genome-wide association studies. *PLoS Genet* **5**, e1000529 (2009).

89.   O. Delaneau, J. F. Zagury, J. Marchini, Improved whole-chromosome phasing for disease and population genetic studies. *Nature methods* **10**, 5-6 (2013).

90.   H. Li *et al.*, The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078-2079 (2009).

91.   K. Wang, M. Li, H. Hakonarson, ANNOVAR: functional annotation of genetic variants from high-throughput sequencing data. *Nucleic acids research* **38**, e164 (2010).

92.   R. C. Team, R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL: http://www.R-project.org/. (2016).

93.   M. Aligeti, A. Roder, S. M. Horner, Cooperation between the Hepatitis C Virus p7 and NS5B Proteins Enhances Virion Infectivity. *Journal of virology* **89**, 11523-11533 (2015).

94.   H. A. Saka *et al.*, Quantitative proteomics reveals metabolic and pathogenic properties of Chlamydia trachomatis developmental forms. *Molecular microbiology* **82**, 1185-1203 (2011).

95.   C. Pujol, J. B. Bliska, The ability to replicate in macrophages is conserved between Yersinia pestis and Yersinia pseudotuberculosis. *Infect Immun* **71**, 5892-5899 (2003).

96. K. A. Datsenko, B. L. Wanner, One-step inactivation of chromosomal genes in Escherichia coli K-12 using PCR products. *Proc Natl Acad Sci U S A* **97**, 6640-6645 (2000).

650   97. C. H. Li *et al.*, Sporangiospore Size Dimorphism Is Linked to Virulence of Mucor circinelloides. *Plos Pathogens* **7**, (2011).

98. B. Howie, C. Fuchsberger, M. Stephens, J. Marchini, G. R. Abecasis, Fast and accurate genotype imputation in genome-wide association studies through pre-phasing. *Nat Genet* **44**, 955-959 (2012).

655   99. P. M. Visscher, W. G. Hill, N. R. Wray, Heritability in the genomics era--concepts and misconceptions. *Nature reviews. Genetics* **9**, 255-266 (2008).

Figures

Figure 1. Inter-individual variation in H2P2 traits reveals clustering of phenotypes and heritable variation in cellular responses to infection. (A) Diagram of the H2P2 workflow for connecting genetic variation to cell biology. Lymphoblastoid cell lines (LCLs) from 528 people

(parent-offspring trios) from 4 populations were exposed to 8 different stimuli for 79 phenotypes.

665 Over 10 million SNPs were tested for association with each phenotype using family-based

association implemented in PLINK (*26, 27*). (B) Clustering of H2P2 phenotypes reveals a map

of trait similarity. Heatmap and dendrograms of hierarchical clustering based on inter-individual

phenotypic variation. Spearman correlation is color-coded in the heatmap from blue (negative

correlation) to red (positive correlation). Phenotypes are color-coded by stimuli (outer band) and

670 biological category of phenotype (inner band). Several clusters of related traits are highlighted.

(C) Narrow-sense heritability (h^2) estimates for H2P2 phenotypes based on GREML vs. parent-

offspring regression. h^2 was estimated using genome-wide SNP data using the Zaitlen method of

GREML for family data in GCTA or by parent-offspring regression. Both methods utilized a

covariate for batch effects. The GREML method gives a SNP-based h^2 (green) as well as a total

675 h^2 (yellow; the sum of SNP-based h^2 (green) plus the non-SNP-based h^2). Both methods indicate

a large range of h^2 estimates for different H2P2 traits and are consistent with many traits having

a significant genetic component. (D) h^2 estimates from parent-offspring regression vs. GREML

SNP-based h^2 are well correlated. Linear regression for all 79 H2P2 traits demonstrate the two

methods give similar h^2 estimates.

Figure 2. GWA of H2P2 reveals 17 genome-wide significant loci including a cis-cytokine-QTL near *CXCL10*. (A) A meta-Manhattan plot for 79 traits shows 17 peaks (red) with p<5x10^{-8} (dotted line). –log(p-values) were calculated using QFAM-parents in PLINK. (B) GARFIELD

enrichment plot of SNP location demonstrates enrichment of SNPs associated with H2P2

685   phenotypes in exons, and 5' and 3' UTRs. SNPs associated with H2P2 traits at various p-value

thresholds were plotted in the indicated colors and the height of the peak within each category

indicates fold enrichment from 0 to 10. (C) GARFIELD enrichment plot of DNase

hypersensitivity peaks demonstrates enrichment of SNPs associated with H2P2 phenotypes in

active chromatin regions in multiple cell/tissue types. SNPs associated with H2P2 traits at

690   various p-value thresholds were plotted in the indicated colors and the height of the peak within

each category indicates fold enrichment from 0 to 14. (D) Regional plot around the *CXCL10*

gene demonstrates association of rs2869462 with CXCL10 levels from *C. trachomatis*-infected

cells. SNPs are plotted by position on chromosome 4 and –log(p-value) and color-coded by r^2

value to rs2869462 from 1000 Genomes European data. (E) Genotypic medians, first and third

695   quartiles (box), and maximum and minimum values (whiskers) for rs2869462 for CXCL10

levels from *C. trachomatis*-infected LCLs from all LCLs. (F) Map of rs2869462 allele

frequencies (C = orange; G = blue) from Geography of Genetic Variants Browser (*81*). (G)

Individual population genotypic median plots for rs2869462 for CXCL10 levels demonstrate C >

G in all populations. P-values are from QFAM-parents in PLINK.

Figure 3. Cross-phenotype associations and pleiotropy are abundant among cellular host-pathogen traits. (A) Histograms of the number of cross-phenotype associations for the 17 H2P2 genome-wide significant hits. For the most strongly associated SNP in each GWAS peak, the

number of traits with associations at $p<0.05$ and $p<6.33 \times 10^{-4}$ is shown. (B) QQ-plot and

705   PheWAS plot for the association of rs2869462 with the 79 H2P2 phenotypes shows deviation

from neutral expectation only for the 4 CXCL10 phenotypes. This is an example of cross-

phenotype associations but not pleiotropy. (C) Circle plot of 79 phenotypes by category and lines

connecting traits that share the same genome-wide significant hit at $p< 1 \times 10^{-5}$. (D) Plot of

pairwise trait phenotypic similarity (Spearman correlation) vs. similarity of shared SNPs (Jaccard

710   index). Traits that are more phenotypically similar have more shared SNPs with $p< 1 \times 10^{-3}$ for

both traits. (E) QQ-plot and PheWAS plot for the association of rs953897 with the 79 H2P2

phenotypes shows deviation from neutral expectation for dozens of phenotypes, including traits

in different biological categories in the PheWAS plot. This is an example of pleiotropy.

Figure 4. Genetic variation influencing *ZBTB20* regulates multiple host-pathogen traits. (A)

Regional plot for *ZBTB20* demonstrates an association of rs953897 with *C. trachomatis* high

GFP infected cells at 46 hrs (p=1.3x10⁻⁸). Genotypic medians plot of rs953897 with high GFP

infected cells at 46 hrs in IBS LCLs. (B) Regional plot for ZBTB20 demonstrates an association

of rs953897 with *S.* Typhimurium-induced pyroptosis (p=7.5x10⁻⁵). Genotypic medians plot of

rs953897 with *S.* Typhimurium-induced pyroptosis in all LCLs. (C-F) LCL GM1761 was treated

with non-targeting, ZBTB20 (53±9% knockdown), or IκB (83±2% knockdown) Accell RNAi for

three days prior to infection. All cellular phenotypes measured were normalized to non-targeting

values (treatment/non-targeting) prior to statistical analysis. (C) ZBTB20 and IκB regulate *C.*

trachomatis replication. By 46hrs, *C. trachomatis* replication has resulted in a high GFP+

population of cells with an enlarged GFP+ *Chlamydia*-containing vacuole. ZBTB20 knockdown

results in a greater percentage of high GFP+ cells, similar to what is seen with the T allele, while

IκB knockdown produces fewer GFP+ enlarged vacuoles. Mean (±SEM) percentage of high

GFP+ cells in non-targeting samples is 1.63% (±0.07%). (D) ZBTB20 regulates *Salmonella*-

induced pyroptosis independent of IκB. ZBTB20 knockdown results in a greater percentage of

pyroptotic cells, similar to what is seen with the T allele, while IκB knockdown shows no

significant change in pyroptosis. Percentage of pyroptotic cells in non-targeting samples is

35.1% (±1.4%). (E) Both the T allele and ZBTB20 knockdown result in reduced IL-6.

Genotypic median plot of rs953897 with *C. trachomatis*-induced IL-6 in all LCLs. ZBTB20

knockdown reduces IL-6 levels, while knockdown of IκB leads to increased levels measured at

70 hours. IL-6 levels from non-targeting LCLs were 186 pg/mL (±31.6 pg/mL). (F) ZBTB20

knockdown results in increased IκB mRNA. RNA was collected from nontargeting and ZBTB20

RNAi treated LCLs, cDNA was synthesized, and qPCR was conducted using TaqMan probes for

18s (control) and IκB (target). The ΔΔCT method was used to determine fold change of IκB. (G)

Proposed model for ZBTB20 effect on proinflammatory targets. C-F were from 8-12 biological

replicates from 2-4 experiments. p-values for C-E were generated from one-way ANOVA

analysis while F was calculated by an unpaired parametric t-test.

Figure 5. SNPs associated with high CXCL10 in H2P2 are also associated with increased risk of IBD. (A) Overlaid association plots of the CXCL10 region on chromosome 4

745   demonstrates colocalization of signals for *Chlamydia*-infected CXCL10 levels from H2P2 (grey) and IBD GWAS (blue) (*72*). rs2869462 from the two studies is highlighted in red and yellow. Colocalization analysis indicates an 80% probability that the peaks for CXCL10 level

and IBD are due to the same causal variant (Table S4). (B) OR plot for rs2869462 and IBD, CD and UC based on data generated in (*72*) and replication of the association with "IBD and other gastroenteritis and colitis" and UC from the eMERGE Network. The high CXCL10 allele (C) is associated with increased odds of IBD.

Figure 6. PheWAS of H2P2 hits reveals connections to human disease including an association of *ZBTB20* and viral hepatitis. (A) Chromosome landscape of H2P2 genome-wide significant hits that demonstrate significant associations with eMERGE phenotypes, after

Bonferroni correction for the number of eMERGE phenotypes. The five SNPs demonstrating

significant associations are described in Table S5. (B) (C) eMERGE PheWAS plots of

rs7566597 and rs953897. eMERGE phenotypes are grouped into functional categories. Y-axis is

–log(p-value), with the red line indicating Bonferroni corrected p-value = 0.05. (D) Huh7 cells

760   were transfected with non-targeting or ZBTB20 siRNA for two days prior to infection. qPCR

demonstrated significant knockdown of ZBTB20. (E) ZBTB20 suppresses HCV infection. The

percentage of cells infected with HCV was assessed at 24 and 48 hpi by immunofluorescence for

HCV protein NS5A (green) and staining with DAPI (blue) for total cells, with automated

quantification by Cellomics. ZBTB20 depletion causes the percentage of HCV-infected cells to

765   increase more rapidly over time. (F) ZBTB20 suppresses HCV production. Supernatants

collected at 72 hpi were used in a HCV focus forming assay to determine the concentration of

productive HCV particles. ZBTB20 depletion causes a 7-fold increase in HCV focus forming

units (FFUs/ml). For D-F, mean (±SEM) are shown and p-values are from a paired t-tests from

three separate experiments.

Table 1. Genome-wide-significant H2P2 SNPs. A single SNP with the lowest p-value is listed for each peak. SNPs described in detail in the text are in bold.

SNP ID	Chr	Position	Cellular trait	p-value	risk allele	gene*	Parent-offspring h^2 for trait	SNP-based h^2 for trait	variance explained by SNP (parental LCLs only)
rs7566597	**2**	**56289463**	**MIP1B_Chlamydia**	**4.67E-08**	**G**	**CCDC85A**	**58.2%**	**33.3%**	**4.4%**
rs9883818	3	337848	MDC_Uninfected_ELISA	1.70E-08	A	CHL1	28.6%	9.8%	1.2%
rs150230900	3	38822664	Chlamydia_70hr_GFP	3.50E-08	-	near RP11-134J21.1	21.1%	12.9%	1.0%
rs953897	**3**	**114349113**	**Chlamydia_46hr_highGFP**	**1.30E-08**	**T**	**ZBTB20**	**33.0%**	**20.1%**	**2.4%**
rs11287866	4	39317796	IL4_Chlamydia	2.60E-08	TA	RFC1	36.8%	22.7%	1.6%
rs2869462	**4**	**76013566**	**IP10_Chlamydia**	**2.00E-09**	**C**	**ART3; near CXCL10**	**48.3%**	**25.2%**	**13.7%**
rs28540901	4	184492384	IL10_Uninfected_Luminex	1.60E-08	T	near IRF2	10.8%	9.2%	2.9%
rs11957501	5	149250737	MDC_S_typhimurium	3.25E-08	C	ABLIM3	43.3%	10.7%	4.9%
rs6555828	5	157066729	Chlamydia_70hr_median_GFP	1.60E-08	G	near HAVCR1	27.3%	22.6%	2.8%
rs139408032	8	8882324	FGF2_Mucor	1.65E-08	A	MFHAS1	38.2%	21.9%	2.7%
rs61836093	10	14235152	FGF2_Candida	3.80E-08	G	FRMD4A	43.7%	16.3%	2.0%
rs74142986	10	81018235	TNFb_Chlamydia	1.85E-08	T	-	35.6%	18.2%	2.5%
rs10750312	11	99526242	S_typhimurium_Intracellular_Replication_24_3_5hr_median_GFP	3.00E-09	G	CNTN5	22.7%	10.2%	3.2%
rs4905049	14	93121941	Chlamydia_46hr_highGFP	3.30E-08	G	near ITPK1	33.0%	20.1%	0.3%
rs4949082	16	63851836	IP10_S_typhimurium	3.70E-08	A	-	40.1%	9.1%	1.2%
rs16956501	17	48419912	Chlamydia_70hr_median_GFP	1.10E-08	C	SKAP1	27.3%	22.6%	0.2%
rs4121804	18	59790252	Chlamydia_46hr_highGFP	4.11E-08	G	-	33.0%	20.1%	0.6%

*gene the SNP is located in or "near" (within 20kb)

Methods

KEY RESOURCES TABLE

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
Bacterial and Virus Strains		
S. enterica Typhi (strain Ty2) +pMMB67GFP	This paper	N/A
S. enterica Typhimurium (strain 14028s) +pMMB67GFP	Dennis Ko	(*19*)
S. enterica Typhimurium (strain 14028S *ΔsifA*) +pMMB67GFP	This paper	N/A
Chlamydia trachomatis LGV-L2, Rif^R + pGFP::SW2	Raphael Valdivia	(*82*)
Mucor circinelloides f. *lusitanicus* R7B (*leuA-*)	Soo Chan Lee	(*83*)
Candida albicans SC5314	Joseph Heitman	(*84*)
Toxoplasma gondii strain RHgfpluc	John Boothroyd	(*85*)
Biological Samples		
Chemicals, Peptides, and Recombinant Proteins		
alpha-hemolysin (Staph. aureus alpha toxin)	Sigma	H9395-.5MG
7-AAD	Enzo Life Sciences	BML-AP400-0001
Critical Commercial Assays		
Human FGF basic DuoSet	R&D Systems	DY233
Human IL-10 DuoSet	R&D Systems	DY217B
Human CXCL10 DuoSet	R&D Systems	DY266
Human MDC Duoset	R&D Systems	DY336
Human IL-6 Duoset	R&D Systems	DY206
Milliplex MAP Human Cytokine Custom 17-plex panel	Millipore EMD	HCYTOMAG

Deposited Data		
Illumina HumanOmni 2.5M array	1000 Genomes	ftp://ftp.1000genomes.ebi.ac.uk/vol1/ftp/release/20130502/supporting/hd_genotype_chip/
H2P2 database and web portal	This paper	h2p2.oit.duke.edu
Experimental Models: Cell Lines		
1000 Genomes LCLs	Coriell	GWD, ESN, IBD, KHV
Experimental Models: Organisms/Strains		
Oligonucleotides		
Accell siRNA for non-targeting #1, *ZBTB20*, *NFKBIA*	Dharmacon	D-001910-01-50; E-020529-00-0010; E-004765-00-0010
Taqman human gene expression assays for *ZBTB20*, *NFKBIA, 18S RNA,*	ThermoFisher	Hs00210321_m1; Hs00355671_g1; Hs03928990_g1
Recombinant DNA		
Software and Algorithms		
PLINK v1.9	(*86*)	https://www.cog-genomics.org/plink/1.9/
LocusZoom	(*87*)	http://locuszoom.sph.umich.edu/
IMPUTE v2.3.2	(*88*)	http://mathgen.stats.ox.ac.uk/impute/impute_v2.html
SHAPEIT v2.r790	(*89*)	https://mathgen.stats.ox.ac.uk/genetics_software/shapeit/shapeit.html

Samtools v1.5	(*90*)	http://samtools.sourceforge.net/
GCTA v1.26	(*48*)	http://cnsgenomics.com/software/gcta
GARFIELD	(*50*)	http://www.ebi.ac.uk/birney-srv/GARFIELD
CIRCOS v0.69	(Krzywinski et al. 2009)	http://circos.ca/software/download/circos/
ANNOVAR v2016FEB01	(*91*)	http://annovar.openbioinformatics.org/
GGV Browser	(*81*)	http://popgen.uchicago.edu/ggv/
R v3.3.2	(*92*)	https://cran.r-project.org/
ggplot2	The R Foundation	https://cran.r-project.org/web/packages/ggplot2/index.html
data.table	The R Foundation	https://cran.r-project.org/web/packages/data.table/index.html
dplyr	The R Foundation	https://cran.r-project.org/web/packages/dplyr/index.html
biomaRt	Bioconductor	https://bioconductor.org/packages/release/bioc/html/biomaRt.html
stringr	The R Foundation	https://cran.r-project.org/web/packages/stringr/index.html
RColorBrewer	The R Foundation	https://cran.r-project.org/web/packages/RColorBrewer/index.html
reshape2	The R Foundation	https://cran.r-project.org/web/packages/reshape2/index.html
GenomicRanges	Bioconductor	https://bioconductor.org/packages/release/bioc/html/GenomicRanges.html
PheWAS	The R Foundation	https://phewascatalog.org/

Other		

CONTACT FOR REAGENT AND RESOURCE SHARING

780   Further information and requests for reagents should be directed to and will be fulfilled by the

Lead Contact, Dennis Ko (dennis.ko@duke.edu).

EXPERIMENTAL MODEL AND SUBJECT DETAILS

Cells

785   1000 Genomes LCLs (528; all trios) from ESN (Esan in Nigeria), GWD (Gambians in Western

Divisions in the Gambia), IBS (Iberian Population in Spain), and KHV (Kinh in Ho Chi Minh

City, Vietnam) populations were purchased from the Coriell Institute. LCLs were maintained at

37˚C in a 5% CO_2 atmosphere and were grown in RPMI 1640 media (Invitrogen) supplemented

with 10% fetal bovine serum (FBS), 2 mM glutamine, 100 U/ml penicillin-G, and 100 mg/ml

790   streptomycin.

Human hepatoma Huh7 cells were grown in DMEM (Mediatech) supplemented with 10% fetal

bovine serum (HyClone), 2.5 mM HEPES, and 1× non-essential amino acids (complete,

cDMEM; Thermo Fisher Scientific). The identity of the Huh7 cell line was verified using the

Promega GenePrint STR kit (DNA Analysis Facility, Duke University), and cells were verified

795   as mycoplasma free by the LookOut Mycoplasma PCR detection kit (Sigma). Infectious stocks

of a cell culture-adapted strain of genotype 2A JFH1 HCV were generated and titrated by focus-

forming assay (FFA), as described (*93*).

METHOD DETAILS

LCL screening

LCLs were received from Coriell and cultured for 8 days prior to assays. LCLs were counted with a Guava Easycyte Plus flow cytometer (Millipore). LCLs were washed once with RPMI 1% FBS and then plated out in RPMI 10% FBS at 200,000 cells/200µl for Salmonellae, 100,000 cells/100µl for fungi, and 40,000 cells/100µl for *S. aureus* alpha toxin, *C. trachomatis,* and *T. gondii*. Cells were passaged at 150,000/ml in 20ml total volume for three days.

C. trachomatis **infection**

C. trachomatis LGV-L2 Rif^R pGFP::SW2 was grown and purified as previously described (*94*). *C. trachomatis* was added at MOI 5 in 100µl assay media, mixed by multichannel pipetting, and centrifuged onto cells at 3000 RPM for 30min at 4ºC. At 27, 46, and 70hrs, cells were mixed and 25µl was taken for flow cytometry measurement (4000 cells). 25µl of supernatant at 70hrs was measured by Luminex assay for 17 cytokines.

Salmonella **infection**

Salmonellae were tagged with an inducible GFP plasmid [pMMB67GFP from (*95*)]. *sifA* deletion mutants was constructed with lambda red (*96*) and verified by PCR. Assaying LCLs for Salmonellae infection was conducted as previously described (*19*). Overnight bacterial cultures were subcultured with a 1:33 dilution and grown for 2 hr 40 min at 37˚C. Invasion was conducted for 1hr at a multiplicity of infection (MOI) of 10 for *S.* Typhi and MOI 30, followed by addition of gentamicin (50µg/ml) for 1hr, and then culture was split into two separate cultures of 60µl of cells with 140µl of media to dilute gentamicin (15µg/ml) and allow for collection at two timepoints. IPTG (1.4mM) was added to turn on GFP expression for 75 min prior to 3.5hr and 24hr timepoints. For the 3.5hr timepoint, 150µl of cells were stained with 7-AAD (7-

aminoactinomycin D; Enzo Life Sciences) and green and red fluorescence of 7000 cells was measured on a Guava Easycyte Plus flow cytometer (Millipore). For the 24hr timepoint, cells were spun down and 2 aliquots of 55ul of supernatant was removed and stored at -80C for

825   subsequence IL10 (25µl), CXCL10 (25µl), and MDC (4µl) ELISAs (R&D Systems). 55µl of cells were stained with 7-AAD and measured by flow cytometry.

Fungal infection

The *Mucor circinelloides* f. *lusitanicus* R7B (*leuA-*) (*83*) strain and *Candida albicans* SC5314 (*84*) strain were used as wild-type to test the expression of FGF-2 from the LCLs. Leucine

830   autotropism (*leuA⁻*) was found not to impact virulence (*97*). To prepare *Mucor* spores, potato dextrose agar (PDA, 4 g potato starch, 20g dextrose, and 15 g agar per liter) was inoculated and incubated for 4 days at 26°C in the light. To collect spores, sterile deionized distilled water was added onto the plates and spores were released by gently scraping the colonies with a cell spreader. Spores were counted by using a hemocytometer. To prepare *C. albicans* yeast, yeast

835   dextrose broth (10 g yeast extract, 20 g peptone, 20 g glucose per liter) was inoculated and incubated at 30°C by shaking at 250 rpm overnight. The yeast cells were quantified by using a hemocytometer. To co-culture with LCLs, all fungal cells were washed with sterile PBS twice. Fungi were added at MOI 1 in 10µl and incubated for 24 hrs. Culture supernatant was collected and stored at -80°C for later FGF-2 ELISA analysis (R&D Systems).

840   ***S. aureus* toxin treatment**

LCLs were treated with alpha-hemolysin (Sigma) at 1µg/ml for 23hrs. Cells were mixed and cell death quantified by 7-AAD staining (concentration) and flow cytometry.

***T. gondii* infection**

T. gondii strain RHgfpluc was grown on confluent human foreskin fibroblast cells. The infected

845   cells were then scraped and transferred to a 50ml polystyrene tube and centrifuged at 500 x g for

10 minutes at 4°C. Pellet was resuspended in 3 ml of PBS and the suspension was aspirated three

times using a 20g needle attached to a 10ml syringe. 30 ml of PBS was added and centrifuged at

500 x g for 10 min at 4°C. Supernatant was removed, and pellet resuspended in 5ml of PBS.

Concentration of a 1:200 dilution was determined by flow cytometry and added at MOI 2 to

850   cells. At 5, 30, and 48 hrs infection, cells were mixed, and 25µl taken for measuring 4000 cells

by flow cytometry.

LCL RNAi experiments

LCLs ($2x10^5$ cells) were treated for three days in 500µl of Accell media (Dharmacon) with either

non-targeting Accell siRNA #1 or an Accell SmartPool directed against human *ZBTB20* or

855   *NFKBIA* (1µM total siRNA; Dharmacon). Prior to infection, cells were plated at $1x10^5$ in 100µl

RPMI complete media (without antibiotics) in 96-well plates. Infections were conducted as

described above.

HCV infection experiments

Huh7 cells were seeded in 12-well plates at a density of $1x10^5$ cells per-well in cDMEM and

860   transfected the next day using 9µL RNAiMAX (Thermo) and 3µL of the indicated 10µM siRNA

(siGenome Smartpool (Dharmacon)), in Optimem (Thermo). Four hours post-transfection, the

transfection mixture was removed and 1ml fresh cDMEM was added. HCV infections were

performed at an MOI of 0.3 for 24, 48, or 72 hr. For each condition, duplicate wells were either

infected or mock-infected, RNA was harvested from one well and the other utilized for

865   visualization of infected cells. Supernatant was collected from both wells for virus titration.

HCV focus forming assay. Serial dilutions of supernatants collected from non-targeting or ZBTB20-targeting siRNA treated cells collected 72hpi were used to infect naive Huh7.5 cells in triplicate wells of a 48-well plate. At 48 hpi, cells were fixed, permeabilized, and immunostained with HCV NS5A antibody (1:500; gift of Charles Rice, Rockefeller University). Following

870 binding of horseradish peroxidase (HRP)-conjugated secondary antibody (1:500; Jackson ImmunoResearch), infected foci were visualized with the VIP Peroxidase Substrate Kit (Vector Laboratories) and counted at 40× magnification.

Visualization of HCV infected cells. 48 hours post-transfection, Huh7 cells treated with non-targeting or ZBTB20-targeting siRNA were infected with HCV (MOI 0.3) or mock-infected. 24

875 or 48 hours post-infection, cells were fixed in 4% paraformaldehyde in PBS, permeabilized with 0.2% Triton X-100 in PBS, and blocked with 3% BSA in PBS, then immunostained with HCV NS5A antibody (1:1,000), washed 3x in PBS-Tween, then visualized with AlexaFluor- 488 Donkey anti-Mouse secondary antibody (1:1000, Thermo). Cell nuclei were stained with DAPI in the first of three PBS-Tween washes following the addition of secondary antibody. Two-color

880 images were collected with the Cellomics ArrayScan VTI HCS (Thermo), at 20x magnification in the Duke Functional Genomics Shared Resource. 10 fields of per-well, per-condition were acquired and the percentage of identified nuclei with detectable NS5A staining was quantified using VHSview software (Thermo).

885 **QUANTIFICATION AND STATISTICAL ANALYSIS**

Phenotype repeatability

Repeatability of each cellular trait was calculated to measure the variation among 3 independent experiments. The inter- and within-individual component of variance was calculated by fitting to

one-way ANOVA. The estimated within-individual component of variance gave the repeatability

890   coefficient.

Testing effect of EBV copy number on H2P2 cellular phenotypes

EBV relative copy numbers were retrieved for 1753 LCLs (*46*), of which 284 cell lines

overlapped with H2P2 samples (73 ESN; 105 GWD; 106 IBS). Prior to analysis, H2P2 cell

phenotypes were averaged from the three experiment replicates and then transformed to Z-scores

895   within each experimental batch. EBV loads were standardized to Z-score within each population.

Correlation between H2P2 cellular traits and EBV loads were tested using linear regression with

population as a covariate.

Genotype and imputation

Genotypes for 1000 Genome LCLs (*16*) were from Illumina HumanOmni 2.5M array (905,788

900   SNPs; see details in STAR resource table). Genome-wide imputation of autosomal genotypes

with 1000 genome Phase 3 haplotype as reference panel were performed through two steps, a

pre-phasing step using SHAPEIT2 (*89*) and an imputation step using IMPUTE2 (*98*). Imputed

genotype was further filtered by imputation accuracy score (IMPUTE's INFO) < 0.9 and minor

allele frequency < 0.01. A total of 339 samples overlap with 1000 genome Phase 3 individuals.

905   We merged those direct sequenced genotypes from 1000 genome Phase 3 project into our

imputed genotypes. We eventually obtained 15,581,278 SNPs (8,817,925 SNPs have minor

allele frequency ≥ 0.05). The human genome reference assemble (GRCH37/hg19) was used for

all analysis.

Phenotype- and SNP-based heritability analysis

910   Two different methods were applied to estimate heritability. The parent-offspring (PO)

regression method estimated additive heritability exclusively using phenotypic values. Linear

regression of child against average of parents was performed, and the slope was used as a heritability estimator. Batch was incorporated as a covariate. Genotype-based heritability was estimated using the GCTA GREML method (Yang et al. 2012). Autosomal SNPs with minor allele frequency filtering of 0.05 were used to create a genetic relationship matrix (GRM). Zaitlen and colleagues developed a method, "big K/small K", which enables to precise estimate heritability by jointly using closely related and unrelated individuals (*49*). Following Zaitlen's method, variance explained by genome-wide SNPs (σ_g^2) was then estimated for each cellular trait. The Zaitlen modification provides joint estimates of 1) h^2 based on pedigree relatedness and 2) h^2 based on inferred relatedness from genome-wide SNPs.

While the standard error for SNP-based h^2 estimates were quite large, we nonetheless observed very strong correlation between these estimates and the parent-offspring h^2 estimates (see Figure 1D). We estimated h^2 based on the analysis of LCLs from all populations in H2P2 to increase the precision of our estimates by including more individuals. Although h^2 is a population-specific parameter, h^2 are often quite similar across different populations and even species (*99*). Nonetheless, we include estimates and standard errors for the combined analysis as well as individual populations in Table S2.

Genome-wide association analysis

Genome-wide association analysis was conducted with PLINK v1.9 (*26*). Analysis was carried out using the QFAM-parents approach with adaptive permutation and a maximum of 10^9 permutations. The QFAM procedures implemented in PLINK use linear regression to test for association while employing permutation of within- and between-family components separately to control for family structure (*27*).

Enrichment analysis

935 Enrichment analyses were carried out using GARFIELD (*50*). GARFIELD has predefined a total of 1005 features from ENCODE and the NIH Roadmap project, and applies generalized linear regression models while accounting for the effects of linkage disequilibrium (LD), minor allele frequency, and local gene density. The GWAS summary statistics were used to quantify fold-enrichment against predefined annotation features at different GWAS p-value thresholds.

940 **PheWAS Analysis**

Testing for association of H2P2 genome-wide hits with clinical phenotypes was performed with the eMERGE biobank dataset of 83,717 individuals from 12 contributing medical centers (*1-4*) with ICD-9 derived PheWAS codes (*5, 6*). A merged set of unified variant genotypes across 78 batches of samples with different genotype platforms (e.g. various Illumina and Affymetrix arrays) was produced by imputation using the Michigan Imputation Server (MIS) with the

945 HRC1.1 haplotype reference set (*7-9*). Sixteen variants were selected for PheWAS based on association in H2P2 and their inclusion in the imputed eMERGE biobank dataset. The PheWAS codes were defined by query of the ICD-9 electronic medical record datasets of the contributing medical centers. Two types of PheWAS code phenotypes were used in the association to ascertain more chronic versus singleton diagnoses: the minimum code count of one (mcc1) ICD-

950 9 code to define an individual as a PheWAS code case, and minimum code count of two (mcc2) instances to define an individual as a chronically represented case. In the mcc2 cases, individuals were excluded from analysis if they only had one instance of the ICD-9 code. If there were less than 500 cases we did not include the ICD-9 derived PheWAS code in analysis because it would

955 likely be underpowered and impact the multiple testing correction. We also did not include medical centers that had low ascertainment of the ICD-9 by excluding medical centers which had less than 10 cases. This resulted in 1,338 for mcc1 and 788 for mcc2 phenotypes being included

in the analysis. We used the PLINK1.9 identity by descent genome file to find the set of unrelated individuals to bring forward for analysis. PheWAS association was implemented in the

960   R *glm()* logistic regression of the case-control data and plotting was carried out using the PheWAS R package (*5*). The covariates of gender and the PLINK1.9 computed 1 and 2 principal components from the pruned (>5% minor allele frequency, genotype and sample missingness > 0.1 and LD r-square<0.7.) genome wide imputation variant genotypes were included in the regressions. The p.adjust() R function with Bonferroni methods was used to adjust p-values of

965   the tested PheWAS codes within a particular SNVs sets of tests for multiple comparisons. Bonferroni of less than 0.05 was used as a significance threshold.

Colocalization analysis

Colocalization analysis was performed using R "coloc" v2.3.1 package (available at http://cran.r-project.org/web/packages/coloc). This software applies a Bayesian framework to estimate the

970   posterior probability of genomic variants affecting both cellular trait and disease based on pre-computed GWAS p values, odds ratios, and minor allele frequencies. We ran colocalization on a 400-kb region centered on the focus SNP rs2869462 using default COLOC parameters $(P1=P2=1x10^{-4}; P12=1x10^{-5})$. Summary statistics of IBD GWAS (*72*) were obtained from www.ibdgenetics.org.

975   **Gene expression analysis on 1000 genome RNAseq project**

Gene expression data of 465 individuals (*57*) were obtained from the EBI website (https://www.ebi.ac.uk/Tools/geuvadis-das/). The rs2869462 genotype data was downloaded from the 1000 genome project (*16*). Effects of rs2869462 on *CXCL10* gene expression were tested by linear regression on both combined populations and individual population.

980   **Descriptive statistics and visualization**

Descriptive statistics were performed with GraphPad Prism 6 (GraphPad Software, US) and with

R (*92*). QQ plots were plotted using quantile-quantile function in R. Regional Manhattan plot

were made using LocusZoom (*87*). Circos v0.69 was used to visualize the shared SNPs among

different groups. The size of each study or number of replicates, along with the statistical tests

985   performed can be found in Figure Legends. All numerical data are presented as the mean ± SEM

(standard error of mean).

DATA AND SOFTWARE AVAILABILITY

All H2P2 data is available for browsing and download at http://h2p2.oit.duke.edu

990   The H2P2 application server is running Red Hat Enterprise Linux Server v7.4, Apache v2.4.6,

Shiny Server v1.5.3.838, R v3.4.1, and Microsoft ODBC Driver 13 for SQL Server.

Implemented R packages include shiny v1.0.3, RODBC v1.3-15, ggplot2 v2.2.1, d3heatmap

v0.6.11, and DT v0.2. The H2P2 database server is running MS Windows Server 2016 and MS

SQL Server 2016. Large volume tables and indexes (2.5 billion GWAS observations and 8.5

995   billion genotype observations) were partitioned for improved query performance. Parallelized

query implementation was also used to improve performance.

Supplemental Information

Table S1. Phenotype definitions and representative flow cytometry plots.

Figure S1. Histograms of 79 H2P2 traits. For flow cytometric data, raw phenotype values are used. For cytokine data, concentrations are log2 transformed.

Figure S2. Repeatability of 79 H2P2 traits. Repeatability was calculated as the inter-individual component of variance from ANOVA of 3 measurements of LCLs taken on sequential cell passages. Phenotypes are color-coded by stimuli. All traits have significant repeatability ($p < 0.05$ marked by asterisk), except for three of the toxoplasma traits.

Figure S3. Minimal correlation between EBV copy number and H2P2 traits. The regression slopes of EBV copy number (from (*46*)) vs. the phenotypic values of each H2P2 trait (transformed into a Z-score and with population as a covariate) were determined. For all H2P2 traits grouped by biological category, slopes were rank ordered and plotted with the 95% confidence interval. Only two traits (IL12P70_Chlamydia and S_typhi_3_5hr_median_GFP) had slopes that were significantly non-zero at a nominal $p < 0.05$ threshold, and even for these traits, EBV copy number accounted for only a small proportion of total variance ($r^2 = 0.028$ and 0.025).

Table S2. Linear regression of EBV copy number and H2P2 traits.

Figure S4. Heritability of 79 H2P2 traits by parent-offspring regression. Individual plots for all H2P2 traits are shown with parent-offspring regression lines with (blue) and without (red) batch covariate. Estimated h^2 and p-value for parent-offspring regression with batch covariate are listed on each plot.

Table S3. Heritability of 79 H2P2 traits by parent-offspring regression and SNP-based heritability.

Figure S5. QQ plots of 79 H2P2 traits. All SNPs with MAF > 5% are shown.

1020 **Figure S6.** Association of rs2869462 with *CXCL10* mRNA. rs2869462 is associated with expression of the *CXCL10* mRNA (p=7x10^{-7}) from the RNAseq data in (*57*)) in the same direction as the H2P2 *CXCL10* protein data.

Table S4. COLOC analysis of the CXCL10 region.

Table S5. eMERGE PheWAS results for 16 H2P2 hits. All eMERGE phenotypes with

1025 Bonferroni corrected p < 0.05 are shown.