Phase-tuned neuronal firing encodes human contextual representations for navigational goals

Andrew J Watrous[1], Jonathan Miller[1], Salman E Qasim[1], Itzhak Fried[2], Joshua Jacobs[1]

1. Department of Biomedical Engineering, Columbia University, New York, NY 10027
2. Department of Neurosurgery, David Geffen School of Medicine and Semel Institute for Neuroscience and Human Behavior, University of California, Los Angeles, Los Angeles, CA 90095, USA

Abstract

We previously demonstrated that the phase of oscillations modulates neural activity representing categorical information using human intracranial recordings and high-frequency activity from local field potentials (Watrous et al., 2015b). We extend these findings here using human single-neuron recordings during a navigation task. Cells with firing rate modulations were observed primarily in entorhinal and frontal cortices. Using a novel oscillation detection algorithm, we identify phase-locked neural firing that encodes information about a person's prospective navigational goal. These results provide evidence for contextual accounts of human MTL function at the single-neuron level and identify phase-coded neuronal firing as a component of the human neural code.

Introduction

Single-neuron firing forms a fundamental basis of the neural code during perception and memory. In addition to the well-established role for behavior-related changes in neuronal firing rates, converging evidence across species and behaviors suggests that interactions between single-neuron spike timing and network oscillations observed in the local field potential (LFP) also contribute to the neural code (Hyman et al., 2005; Huxter et al., 2003; Rutishauser et al., 2010; Belitski et al., 2008; Ng et al., 2013; Kayser et al., 2009; Siegel et al., 2009). For instance, rodent hippocampal cells show phase precession relative to theta oscillations during navigation (O'Keefe & Recce, 1993; Terada et al., 2017), in which the theta phase of neuronal firing represents information about a rat's position (Jensen & Lisman, 2000). Synthesizing these findings in Spectro-Contextual Encoding and Retrieval Theory (SCERT), we have hypothesized that frequency-specific and phase-locked neuronal firing at different phases (i.e. phase coding) also forms a basis of the human neural code (Watrous & Ekstrom 2014; Watrous et al., 2015a). We previously reported evidence for SCERT (Watrous et al., 2015b) using high-frequency activity in the LFP as a proxy for single-cell spiking (Crone et al., 1998; Manning et al., 2009; Miller et al., 2014). However, given the uncertain relationship (Ekstrom et al., 2007; Rey et al., 2014) between single neurons and high-frequency activity in the human medial temporal lobe (MTL), it is unclear whether phase coding manifests in MTL neurons. We clarify this issue here by testing new aspects of SCERT, seeking to extend our previous findings of phase coding (Watrous et al., 2015b) to the single-neuron level.

Several lines of evidence indicate that the human MTL forms active representations of spatial context (Ranganath & Ritchey, 2012) such as navigational goals (Watrous et al., 2011; Brown et al., 2016), yet how such representations are instantiated at the single-neuron level remains largely unknown. Drawing upon SCERT, we hypothesized that phase-coding in single neurons also supports spatial contextual representations for prospective goals.

We analyzed a dataset that simultaneously measured human single-neuron and
oscillatory activity from MTL, frontal, and lateral temporal regions during a goal-directed
navigation task (Jacobs et al., 2010; Miller et al., 2015). Following the analytic strategy
from our previous work (Watrous et al., 2015b), we first tested for frequency-specific
phase locking and then directly tested for phase coding, which would appear as
individual neurons that spiked at different phases according to the prospective goal. We
examined these patterns first in the medial temporal lobe and then extratemporal areas.
Our results confirmed the existence of rate and phase coding for navigational goals in
individual neurons, thus providing the first evidence for the oscillatory phase coding of
spatial contextual information in the human MTL.
62
Results
Slow theta oscillations (3Hz) in the MTL during virtual navigation
Our primary aim was to test if human MTL neurons encode behavioral
information by modulating their spiking based on the phase of slow oscillations.
Examining this hypothesis required that we accurately identify the presence and phase
of slow oscillations, particularly because human MTL oscillations are lower frequency
and less stationary compared to the stable theta oscillations observed in rodents
(Watrous et al., 2013; Vass et al., 2016). We developed a novel method, the Multiple
Oscillations Detection Algorithm ("MODAL"; Figure 1A-C), to detect and characterize
neural oscillations in adaptively identified band(s) whose frequency ranges are
customized for each recording site according to its spectral properties. MODAL
identifies narrow-band oscillations exceeding the background 1/f spectrum (Figure 1A)
and calculates the instantaneous phase and frequency of oscillations in each band (see
Methods) while excluding timepoints without oscillations or that exhibited epileptogenic
activity (Gelinas et al., 2016). Thus, MODAL allowed us to test for phase coding of
spikes during the presence of narrowband oscillations.

79
Figure 1 Multiple Oscillation Detection Algorithm ("MODAL")
A-C) Key steps in the algorithm, shown for an example electrode from the right hippocampus of
patient 9. A) Mean log power averaged over time (black) and a fit line of the 1/f background
spectrum (gray). A slow theta band (blue) and a beta band (green) are identified as contiguous
frequencies exceeding the fit line.   B) Example output from MODAL depicting a raw trace
example of the LFP (upper) with the detected oscillations in each band (lower). The
instantaneous frequency of the detected oscillation in each band is overlaid on a spectrogram
and gray portions of the spectrogram indicate power values exceeding a local fit (similar to A but
using a 10s epoch). C) Accumulating detections over time reveals the prevalence of oscillations
at each frequency on this electrode (black). Blue and green bars indicate the overall prevalence
of oscillations in each frequency, independent of the exact frequency within a band. D)
Population data demonstrating low frequency oscillations. Grey line indicates the percent of LFP

channels with a detected band as a function of frequency. Of those channels with a detected
band, the black line indicates the average amount of time each frequency was detected. Slow
theta oscillations (below 5Hz) are observed using both metrics.
95
MODAL reliably identified oscillations at multiple frequencies that were visible in
the raw trace (Figure 1B-C). Analyzing each of 385 LFP signals across the entire task
period using MODAL, we found that most signals showed a band of activity centered at
"slow theta" (~3Hz; 93% of signals; Figure 1D, gray line). Analyzing the overall amount
of time each frequency was detected on these electrodes, we found that slow theta was
detected most often (Figure 1D, black line). These results are consistent with previous
work showing the prevalence of slow theta in the human MTL (Watrous et al., 2011;
Watrous et al., 2013; Vass et al., 2016, Jacobs, 2014; Bohbot et al., 2017). We
subsequently restricted our analysis to the low-frequency band (1–10 Hz) in order to
mirror the approach from our previous work (Watrous et al., 2015b).
106
Phase-locked neuronal firing
We leveraged MODAL's ability to precisely track the instantaneous phase during
oscillations to probe how phase coordinates the activity of individual neurons. Focusing
first on the MTL, we analyzed 441 (83%) neurons that each had a simultaneously
recorded LFP with an oscillation at 1–10 Hz. In many cells we observed significant
phase-locking, an overall tendency for firing to increase at particular phases of the LFP
oscillation (Jacobs et al., 2007; Rey et al., 2014). Phase locking is evident by examining
the LFP phase distribution for all spikes which occurred during oscillations from a given
cell (Figure 2A upper, Rayleigh p<.005). Across our population of recordings, we
identified phase-locked neural firing in 119 neurons (111/441, 25%, Rayleigh test,
p<.005), a proportion significantly above chance (Binomial p<.00001). We observed that
phase locked neural firing was clustered just after the trough of the oscillation for these
cells (Figure 2B, Rayleigh test p<.00001) and most phase locking occurred to slow-theta
oscillations below 5 Hz (Figure 2C).

121
Figure 2 Phase-Locked Neural Firing
A) Spike-triggered average of a phase-locked neuron from the right hippocampus of Patient 1
(left). Red tick mark denotes a spike. Circular histograms (right) show phases at which spikes
occurred relative to two detected bands. Spiking was phase-locked to the ascending phase in the
1.5-5 Hz band (red) but not in the 7.5-9 Hz band (Rayleigh test, p=.004 and p=.34,
respectively). B) Population data: Pooling over frequencies, mean spike phases were
significantly clustered near the initial ascending phase of the oscillation. C) Population scatter
plot of the mean phase of firing and maximally detected frequency within the band for each of 119
phase-locked neurons.
131

The LFPs associated with 48 neurons displayed oscillations at two distinct
frequency bands in the 1–10Hz range. We next tested if the spike–LFP phase locking
was specific to an individual frequency band or present for both bands. 12.5% of these
cells (6/48) showed frequency-specific phase locking, showing phase-locked firing in
only one LFP frequency band (Figure 2a; p<.005 in one band, p>.1 in all other
bands). Extending previous findings (Jacobs et al., 2007) by examining phase-locking to
adaptively-identified narrowband signals, we find that human neuronal firing is
modulated by the phase of low-frequency oscillations in a band and frequency-specific
manner, as predicted by SCERT (Watrous & Ekstrom, 2014).
141
LFP-spike phase coding of goal information
Previous work has identified single neurons responsive to navigational goals
(Ekstrom et al., 2003). To understand the behavioral relevance of phase-tuned neuronal
activity, we tested whether neurons also used phase-tuned neural firing to encode
contextual information about the patient's prospective navigational goal, analogous to
the phase coding for location in the rodent hippocampus (O'Keefe & Recce, 1993). We
identified 160 goal cells (36%) whose firing rates were significantly modulated by the
patient's navigational goal (Figure 3A-B, all $\chi2(5)>$, p<.0001). These cells were present
in 11 of 12 patients. This result replicates previous studies implicating the MTL in the
representation of navigational goals (Ekstrom et al., 2003; Watrous et al., 2011).

152
Figure 3 Example cells showing goal coding by firing rate and spike-LFP phase
A) Example neuron from patient 4 whose firing rate was significantly modulated by navigational
goal (chi-square test, p<.00001) but not by spike-LFP phase (decoding p>.05, not shown). Firing
rate is plotted as a function of each navigational goal. B) Another example neuron showing firing
rate modulation by goal from patient 11. C) Example neuron from patient 1 showing significant
spike-LFP phase coding for goal 4 (difference score (DS) = 2) compared to goals 5 and 6.
Circular histograms show spike counts separately for different goals, and only goals with a
difference score greater than zero are plotted for clarity. Black line at center of each plot shows
the resultant vector and the colored arc indicates the 95th percentile confidence interval of the
circular mean. D) Example cell from patient 6 showing phase coding for goal 6. Each cell in
Figure 3 is unique and from a different patient. LEC/REC: Left/Right entorhinal cortex; LH/RH:
Left/Right hippocampus
165
We then asked if neurons additionally represent information about the
prospective goal via phase coding. We first examined the LFP phase distribution for

each cell's spiking using a difference score (DS) approach, in which we use circular
statistics to compare distributions of spike phases between individual goals (Watrous et
al., 2015b). This analysis revealed cells that fired at significantly different LFP phases
(p<.0001) according to the patient's goal (Figure 3C-D). For instance, Figure 3C shows
the spike–phase distribution from a right hippocampal neuron that fired preferentially
near the oscillatory peak when the patient was seeking goal #4 and near the trough for
goals 5 and 6. To more systematically quantify phase coding and probe whether this
phenomenon is distinct from rate coding, we examined the 158 neurons whose firing
rates were not goal-modulated (p>.05). Of these, we identified 28 neurons (17%) with
significantly different spike phases for different goals (DS>0 for at least one goal), a
proportion significantly above chance (Binomial test, p<.000001, chance = .237 cells).
Thus, independent information about the patient's prospective goal could be recovered
by considering the LFP phase at which these neurons fired.
To verify this interpretation and further ensure that these phase differences were
robust, we used a decoding approach (Watrous et al., 2015) to test whether the patient's
prospective goal could be predicted from the phase of neuronal spiking for cells that did
not demonstrate rate coding. We observed significant decoding of goal information from
spike phase in 19 of 158 (12%) neurons for at least one band (Binomial test, p<.00014,
chance = 7.9 cells). Cells that exhibited phase coding of goal information were present in
7 of 12 patients. We observed a similar proportion of phase-coding neurons (67/441,
15%) when considering all neurons in our dataset, indicating that our exclusion of rate-
modulated cells did not bias our results.
Finally, we explored the anatomical distribution of rate and phase-coding cells
across our dataset (see Methods). We found that rate-coding cells were differentially
clustered in particular regions ($\chi^2(5)$=70.5, p <10^{-12}). The entorhinal cortex (58% of 162
cells) and frontal cortex (44% of 355 cells) had the largest proportions of cells with firing
rate modulations for goals. In contrast, phase coding cells were not significantly
clustered by brain region ($\chi^2(5)$=7.3, p=.19). Together, these results extend our previous
findings (Watrous et al., 2015b) to single neurons, providing the first evidence for single-
neuron phase coding during navigation in humans.
198
General Discussion
Analyzing recordings from epilepsy patients performing a goal-directed
navigation task, we expand our previous observation of phase-coding with high-
frequency LFPs (Watrous et al., 2015b) to the domain of single neuron spiking. While we
replicated the earlier finding of firing-rate coding of goal representations in human single-
cell activity (Ekstrom et al., 2003), we also found a distinct population of cells in which
spike-LFP phase coding contributed to representations in the absence of significant
changes in firing rate (Rutishauser et al., 2010). Furthermore, we found neurons that
were phase-locked to frequency-specific narrowband oscillations primarily in the slow-
theta band. Together, these findings provide new, stronger evidence for the SCERT
model at the single-neuron level.
Our analyses benefited from employing the MODAL algorithm, which combines
features of earlier algorithms (Whitten et al., 2011; Lega et al., 2012; Cohen 2014) to
identify oscillatory bands in a manner that is customized for each recording site. We
believe MODAL is an improvement on these methods because it adaptively identifies
oscillatory band(s) without introducing experimenter bias regarding bands of interest,
excludes periods when phase is noisy because oscillations are absent, and provides
exactly one estimate of phase and frequency per band.
Our findings provide the first evidence of phase coding during human navigation
and provide a theoretically important link to other model systems where phase coding is

present (Siegel et al., 2009; Kayser et al., 2009; Ng et al., 2013), such as phase-
precession (O'Keefe and Recce, 1993; Terada et al., 2017). However, we found
prominent phase-locking and phase-coding to slower frequency oscillations below 5 Hz,
suggesting that phase coding exists beyond the canonical 8-Hz theta signal seen in rats.
These findings thus lend further credence to findings indicating that (virtual) navigation-
related theta occurs at a slower frequency in humans (Watrous et al., 2013; Jacobs,
2014; Bohbot et al., 2017) and demonstrates that these oscillations modulate neuronal
spiking.
These results align with work implicating the human MTL in spatial contextual
representation (Ranganath & Ritchey, 2012) of navigational goals (Ekstrom et al., 2003;
Watrous et al., 2011; Brown et al., 2016) in support of ongoing behavior (Warren et al.,
2011; Yee et al., 2014) and provide further evidence that the timing of MTL activity is
critical for behavior (Reber et al., 2017; Rey et al., 2014). Combined with previous
human studies (Kraskov et al., 2007; Lopour et al., 2013; Watrous et al., 2015b; ten
Oever & Sack, 2015), our work indicates that both firing rate and the precise timing of
activity relative to LFP phase are general coding mechanisms in the human MTL across
behaviors and tasks, suggesting that other types of contextual information may also be
encoded using LFP phase. Future studies can build off these findings to directly assess
phase coding of other types of contextual information in humans, such as phase-
precession to space or time.
239
Methods
Neural Recordings and behavioral task
We analyzed data from 12 patients with drug-resistant epilepsy undergoing
seizure monitoring (surgeries performed by I.F.). The Medical Institutional Review Board
at the University of California-Los Angeles approved this study. Patients were implanted
with microwire depth electrodes (Fried et al., 1999) targeting the medial temporal lobe
and medial frontal lobe sites. Groups were formed for recordings in hippocampus,
entorhinal cortex, parahippocampal gyrus, amygdala, frontal cortex (orbitofrontal,
cingulate, motor), and lateral temporal cortices (n=214,162,65,212,355,95 neurons,
respectively). Our primary analyses of 441 neurons focused on signals from
hippocampal, entorhinal, and parahippocampal regions. Microwire signals were
recorded at 28-32 kHz and captured LFPs and action potentials, which were spike-
sorted using *wave_clus* (Quiroga et al., 2004). Signals were then downsampled to 2
kHz.
We examined data from a total of 31 recording sessions in which patients
performed a virtual-taxi driver game in a circular environment. Patients were instructed
to drive passengers to one of 6 goal stores in the virtual environment. The recordings
and behavioral task have been detailed in prior publications that have characterized the
spatial-tuning of neurons using firing rate alone (Jacobs et al., 2010; Miller et al., 2015).
Here, our primary analyses in this study focused on how contextual information about
navigational goals may be encoded based on firing rates and spike-LFP interactions.
261
Detection and Rejection of Epileptogenic signals
We implemented an automated algorithm to detect and exclude epochs of signal
likely resulting from epileptic activity following prior work (Gelinas et al., 2016). We first
low-pass filtered (4th order Butterworth) the signal below 80 Hz to remove any spike-
contamination at high frequencies. Epochs were marked for rejection if the envelope of
the unfiltered signal was 4 standard deviations above the baseline or if the envelope of
the 25-80Hz bandpass filtered signal (after rectification) was 4 standard deviations
above the baseline. In some cases, we noted short "bad data" epochs lasting less than

one second were not detected. We conservatively elected to exclude these epochs by
marking any "good data" epoch lasting less than one second as "bad". Bad data epochs
were excluded from all analyses.
273
274   *Multiple Oscillations Detection Algorithm ("MODAL")*
275   Numerous factors contribute to the presence and characteristics of band-limited
neural oscillations, broadly including neuroanatomy, behavioral state, and recording
equipment (Buzsaki et al., 2012). We developed an algorithm to adaptively detect and
characterize neural oscillations in bands exceeding the background 1/f spectrum
motivated by rodent studies that exclude periods of low amplitude theta oscillations
when assessing phase coding (Lenck-Santini & Holmes, 2008). To this end, we
modified the "frequency sliding" algorithm (Cohen 2014), which provides the
instantaneous phase and frequency of oscillations in a band, in two important ways.
283   First, rather than calculating frequency sliding in *a priori* bands, we defined bands
for subsequent analysis on each electrode as those frequencies exceeding the
background 1/f spectrum. We calculated power values in .5Hz steps from 1 to 50 Hz
using 6 cycle Morlet wavelet convolution. We then created a power spectrum by
averaging values over time (and excluding bad data epochs), and fit a line to this
spectrum in log-log space using *robustfit* in Matlab. Similar approaches have been used
previously (Lega et al., 2012; Podvalny et al., 2015). Frequency band edges were
defined as the lowest and highest frequencies in a contiguous set of frequencies which
had values exceeding this fit; several bands could be detected on each electrode. We
then calculated the instantaneous frequency and phase in each detected band using the
"frequency sliding" algorithm (Cohen 2014).
294   Second, frequency sliding provides a frequency and phase estimate at every
moment in time, regardless of the presence or absence of an oscillation. We ensured
that phase & frequency estimates were only obtained during time periods where there
was increased power in the band of interest. We recomputed the power spectrum in 10
second, non-overlapping windows and recomputed the fit line as described above. We
excluded phase and frequency estimates at time points 1) in which the power was below
the fit line or, 2) were during bad data epochs. Finally, we also excluded noisy
frequency estimates outside of the band, which can occur based on "phase slips"
(Cohen 2014). All analyses were conducted in Matlab using custom code which is
available upon request.
304
Statistical Analyses We used Rayleigh tests to identify phase-locked neural firing,
extracting the phase of the LFP during each spike in each detected frequency band. All
analyses were done considering each band separately and statistical thresholding was
set at $p<.005$ for each cell. This was chosen to be stricter than $p<.05$ Bonferroni-
correction across the number of bands detected in the 1-10Hz range. We identified cells
with firing rate modulated by navigational goal using chi-square tests. Under the null-
assumption of Poisson-spiking, which is independent of navigational goal, we derived
expected spike counts for each goal by multiplying total spike count by the proportion of
time the goal occurred throughout the task session.
314   Difference scores were calculated identically to our previous work (Watrous et
al., 2015b) and used the Watson-Williams test to compare phases during spikes that
occurred for each goal. We again used $p<.0001$ for statistical thresholding, as it
corresponded to Bonferroni-correction ($p<.05$) for the 15 pairwise combinations of 6
goals. We then used a decoding-based approach to validate our findings, employing a
linear decoder with fivefold cross-validation to predict the behavioral goal from the phase
of the LFP during neural spiking. We first computed the sine and cosine of the phase

values before classification following previous work (Lopour et al., 2013; Watrous et al.,
2015b). Chance performance varies across cells because we classified goal information
associated with the LFP phase for each spike and the distribution of spikes across goals
varied between cells. We accounted for this using a permutation procedure, re-running
our classification 500 times per cell using shuffled goal information (*circshift* in Matlab to
maintain the temporal structure of the session) to get a surrogate distribution of
classification accuracies per cell. We then obtained a p-value for classification by
ranking our observed classification accuracy to the surrogate distribution; p-values less
than .05 were considered significant.
To analyze the regional specificity of rate and phase coding, we expanded our
analyses to our entire dataset of neurons. We used chi square tests to assess if rate
coding or phase coding cells were differentially prevalent in each region.
333

Acknowledgements
335 We wish to thank the patients for their participation in this study. This work was
336 supported by National Institutes of Health grants NS033221 and NS084017 (I.F.),
337 MH061975 (J.J.), and National Science Foundation GRFP (S.E.Q).

References
340 Belitski, A., A. Gretton, C. Magri, Y. Murayama, M. A. Montemurro, N. K. Logothetis and
341 S. Panzeri (2008). "Low-frequency local field potentials and spikes in primary visual
342 cortex convey independent visual information." J Neurosci 28(22): 5696-5709.
343
344 Bohbot, V. M.S. Copara, J. Gotman, A.D. Ekstrom (2017). "Low-frequency theta
345 oscillations in the human hippocampus during real-world and virtual navigation." Nature
346 Communications 8: 14415.
347
348 Brown, T.I., V.A. Carr,K.F. LaRocque, S.E. Favila, A.M. Gordon, B. Bowles, , J.N.
349 Bailenson, A.D. Wagner (2016). Prospective representation of navigational goals in the
350 human hippocampus. Science, 352:1323-1326.
351
352 Buzsaki, G., C. A. Anastassiou and C. Koch (2012). "The origin of extracellular fields and
353 currents--EEG, ECoG, LFP and spikes." Nat Rev Neurosci 13(6): 407-420.
354
355 Buzsaki, G. and E. I. Moser (2013). "Memory, navigation and theta rhythm in the
356 hippocampal-entorhinal system." Nat Neurosci 16(2): 130-138.
357
358 Cohen, M. X. (2014). "Fluctuations in oscillation frequency control spike timing and
359 coordinate neural networks." J Neurosci 34(27): 8988-8998.
360
361 Crone, N.E., D.L. Miglioretti, B. Gordon, and R.P. Lesser (1998). "Functional mapping of
362 human sensorimotor cortex with electrocorticographic spectral analysis. II. Event-related
363 synchronization in the gamma band". Brain, *121*(12), 2301-2315.
364
Ekstrom, A. D., M. J. Kahana, J. B. Caplan, T. A. Fields, E. A. Isham, E. L. Newman and
I. Fried (2003). "Cellular networks underlying human spatial navigation." Nature
425(6954): 184-188.

Ekstrom, A.D., I. Viskontas, M. Kahana, J. Jacobs, K. Upchurch, S. Bookheimer, , and I.
Fried (2007). "Contrasting roles of neural firing rate and local field potentials in human
memory." Hippocampus, *17*(8): 606-617.
372
373 Fried, I., C.L. Wilson, N.T. Maidment, J. Engel Jr,E. Behnke, T.A. Fields, K.A.
374 Macdonald, J.W. Morrow, and L. Ackerson (1999). "Cerebral microdialysis combined
375 with single-neuron and electroencephalographic recording in neurosurgical patients". J
376 Neurosurgery 91(4): 697-705.
377
378 Gelinas, J. N., D. Khodagholy, T. Thesen, O. Devinsky and G. Buzsaki (2016). "Interictal
379 epileptiform discharges induce hippocampal-cortical coupling in temporal lobe epilepsy."
380 J Neurosci 22(6): 641-648.
381
382 Huxter, J., N. Burgess and J. O'Keefe (2003). "Independent rate and temporal coding in
383 hippocampal pyramidal cells." Nature 425(6960): 828-832.
384
385 Hyman, J. M., E. A. Zilli, A. M. Paley and M. E. Hasselmo (2005). "Medial prefrontal
386 cortex cells show dynamic modulation with the hippocampal theta rhythm dependent on
387 behavior." Hippocampus 15(6): 739-749.
388
389 Jacobs, J., M. J. Kahana, A. D. Ekstrom and I. Fried (2007). "Brain oscillations control
390 timing of single-neuron activity in humans." J Neurosci 27(14): 3839-3844.
391
392 Jacobs, J., M. J. Kahana, A. D. Ekstrom, M. V. Mollison and I. Fried (2010). "A sense of
393 direction in human entorhinal cortex." Proc Natl Acad Sci U S A 107(14): 6487-6492.
394
395 Jacobs, J. (2014). "Hippocampal theta oscillations are slower in humans than in rodents:
396 implications for models of spatial navigation and memory." Philos Trans R Soc Lond B
397 Biol Sci 369(1635): 20130304.
398
399 Jensen, O. and J. E. Lisman (2000). "Position reconstruction from an ensemble of
400 hippocampal place cells: contribution of theta phase coding." J Neurophysiol 83(5):
401 2602-2609.
402
403 Kayser, C., M. A. Montemurro, N. K. Logothetis and S. Panzeri (2009). "Spike-phase
404 coding boosts and stabilizes information carried by spatial and temporal spike patterns."
405 Neuron 61(4): 597-608.
406
407 Kraskov, A., R. Q. Quiroga, L. Reddy, I. Fried and C. Koch (2007). "Local field potentials
408 and spikes in the human medial temporal lobe are selective to image category." J Cogn
409 Neurosci 19(3): 479-492.
410
411 Lega, B. C., J. Jacobs and M. Kahana (2012). "Human hippocampal theta oscillations
412 and the formation of episodic memories." Hippocampus 22(4): 748-761.
413
414 Lenck-Santini P.P., G.L. Holmes (2008). "Altered phase precession and compression of
415 temporal sequences by place cells in epileptic rats." J Neurosci 28(19): 5053-5062.
416
417 Lopour, B. A., A. Tavassoli, I. Fried and D. L. Ringach (2013). "Coding of information in
418 the phase of local field potentials within human medial temporal lobe." Neuron 79(3):
419 594-606.
420

421 Manning J.R., J. Jacobs, I. Fried, M.J. Kahana (2009). "Broadband shifts in local field
422 potential power spectra are correlated with single-neuron spiking in humans." J
423 Neuroscience 29:13613-13620.
424
425 Miller, J. F., I. Fried, N. Suthana and J. Jacobs (2015). "Repeating spatial activations in
426 human entorhinal cortex." Curr Biol 25(8): 1080-1085.
427
428 Miller K.J., C.J. Honey, D. Hermes, R. Rao, M. denNijs, J.G. Ojemann (2014).
429 "Broadband changes in the cortical surface potential track activation of functionally
430 diverse neuronal populations". Neuroimage 85:711-720.
431
432 Ng, B. S., N. K. Logothetis and C. Kayser (2013). "EEG phase patterns reflect the
433 selectivity of neural firing." Cereb Cortex 23(2): 389-398.
434
435 O'Keefe, J. and M. L. Recce (1993). "Phase relationship between hippocampal place
436 units and the EEG theta rhythm." Hippocampus 3(3): 317-330.
437
438 Podvalny, E., N. Noy, M. Harel, S. Bickel, G. Chechik, C. E. Schroeder, A. D. Mehta, M.
439 Tsodyks and R. Malach (2015). "A unifying principle underlying the extracellular field
440 potential spectral responses in the human cortex." J Neurophysiol 114(1): 505-519.
441
442 Quiroga, R. Q., Z. Nadasdy and Y. Ben-Shaul (2004). "Unsupervised spike detection
443 and sorting with wavelets and superparamagnetic clustering." Neural Comput 16(8):
444 1661-1687.
445
446 Ranganath, C. and M. Ritchey (2012). "Two cortical systems for memory-guided
447 behaviour." Nat Rev Neurosci 13(10): 713-726.
448
449
450 Reber, T.P., J. Faber, J. Niediek, J Boström, C.E. Elger, and F. Mormann, (2017).
451 "Single-Neuron Correlates of Conscious Perception in the Human Medial Temporal
452 Lobe". Current Biology.
453
454 Rey, H. G., I. Fried and R. Quian Quiroga (2014). "Timing of single-neuron and local field
455 potential responses in the human medial temporal lobe." Curr Biol 24(3): 299-304.
456
457 Rutishauser, U., I. B. Ross, A. N. Mamelak and E. M. Schuman (2010). "Human memory
458 strength is predicted by theta-frequency phase-locking of single neurons." Nature
459 464(7290): 903-907.
460
461 Siegel, M., M.R. Warden, and E.K. Miller (2009) "Phase-dependent neuronal coding of
462 objects in short-term memory" PNAS 106(50) 21341-21346.
463
464 ten Oever, S. and A.T. Sack (2015). "Oscillatory phase shapes syllable
465 perception". Proceedings of the National Academy of Sciences, 112(52), 15833-15837.
466
467 Terada, S., Y. Sakurai, H. Nakahara and S. Fujisawa (2017). "Temporal and Rate
468 Coding for Discrete Event Sequences in the Hippocampus." Neuron 94(6): 1248-
469 1262.e1244.
470
471 Vass, L. K., M. S. Copara, M. Seyal, K. Shahlaie, S. T. Farias, P. Y. Shen and A. D.
472 Ekstrom (2016). "Oscillations Go the Distance: Low-Frequency Human Hippocampal

Oscillations Code Spatial Distance in the Absence of Sensory Cues during Teleportation." Neuron 89(6): 1180-1186.

Warren, D. E., M. C. Duff, U. Jensen, D. Tranel and N. J. Cohen (2012). "Hiding in plain view: lesions of the medial temporal lobe impair online representation." Hippocampus 22(7): 1577-1588.

Watrous, A. J., J. Fell, A. D. Ekstrom and N. Axmacher (2015a). "More than spikes: common oscillatory mechanisms for content specific neural representations during perception and memory." Curr Opin Neurobiol 31: 33-39.

Watrous, A. J., L. Deuker, J. Fell and N. Axmacher (2015b). "Phase-amplitude coupling supports phase coding in human ECoG." Elife 4.

Watrous, A. J. and A. D. Ekstrom (2014). "The spectro-contextual encoding and retrieval theory of episodic memory." Front Hum Neurosci 8: 75.

Watrous, A. J., I. Fried and A. D. Ekstrom (2011). "Behavioral correlates of human hippocampal delta and theta oscillations during navigation." J Neurophysiol 105(4): 1747-1755.

Watrous, A. J., D. J. Lee, A. Izadi, G. G. Gurkoff, K. Shahlaie and A. D. Ekstrom (2013). "A comparative study of human and rat hippocampal low-frequency oscillations during spatial navigation." Hippocampus 23(8): 656-661.

Whitten, T.A., A.M. Hughes, C.T. Dickson, J.R. Caplan (2011). "A better oscillation detection method robustly extracts EEG rhythms across brain state changes: the human alpha rhythm as a test case." Neuroimage 54(2)860-874.

Yee, L. T., D. E. Warren, J. L. Voss, M. C. Duff, D. Tranel and N. J. Cohen (2014). "The hippocampus uses information just encountered to guide efficient ongoing behavior." Hippocampus 24(2): 154-164.