Transoceanic pathogen circulation in the age of sail and steam

Elizabeth N. Blackmore[1, 2] and James O. Lloyd-Smith[1 ✉]

[1]Department of Ecology & Evolutionary Biology, University of California, Los Angeles , CA 90095, USA
[2]Department of Ecology and Evolutionary Biology, Yale University, New Haven, CT 06520
✉jlloydsmith@ucla.edu

Abstract | In the centuries following Christopher Columbus's 1492 journey to the Americas, transoceanic voyages opened unprecedented pathways for global pathogen circulation. Yet no biological transfer is a single, discrete event. We use mathematical modeling to quantify historical risk of shipborne pathogen introduction, exploring the respective contributions of journey time, ship size, population susceptibility, transmission intensity, density dependence, and pathogen biology. We contextualize our results using port arrivals data from San Francisco, 1850–1852, and from a selection of historically significant voyages, 1492–1918. We offer numerical estimates of introduction risk across historically-realistic journey times and ship population sizes, and show that steam travel and shipping regimes that involved frequent, large-scale movement of people both substantially increased risk of transoceanic pathogen circulation.

Introduction

In the centuries following Christopher Columbus's 1492 journey to the Americas, transoceanic voyages opened unprecedented pathways in global pathogen circulation. Fifty years ago, environmental historian and geographer Alfred Crosby coined the term "Columbian Exchange" to describe the ecological shifts that followed: a pivotal, irreversible transfer of plants, pathogens, and people between the "Old" world and the "New" [18]. Yet while "Columbian Exchange" narratives rightly characterize modern shipping as an important ecological force, no biological transfer is a single, discrete event. The Columbian Exchange was and is a centuries-long process.

Since the initial publication of Crosby's work in 1972, a broad range of ecological scholarship has demonstrated the power of human mobility to shape pathogen ecology, evolution, and epidemiology [38,48,61,63,67,88]. Contact structures have well-documented impacts on the dynamics of influenza [6,11,83], measles [39,93], Covid-19 [10,78], cholera [33,55], malaria [87] and a host of other diseases. This raises a historical question. How quickly, and how uniformly, did transoceanic shipping lead to globalized pathogen ecosystems — and with what consequences along the way for pathogen ecology, pathogen evolution, and human health?

We contend that transoceanic pathogen transfer was slower, less uniform, and less inevitable than traditional "Columbian Exchange" narratives imply. In particular, we argue that acute respiratory pathogens such as smallpox, measles, and influenza rarely survived early colonial voyages. Between

short generation times, lengthy periods at sea, and intense shipboard transmission, these "crowd diseases" could rapidly exhaust all susceptible people on board long before a vessel reached port — leaving no pathogen to introduce.

The idea that onboard pathogens often went extinct among passengers and crew before a ship's arrival is qualitatively intuitive. Historians have made the point repeatedly [45,49,58]. However, a quantitative and theoretical investigation of shipboard outbreak duration offers sharper insight into the historical contours of transoceanic pathogen circulation.

Previously, historical geographers Andrew Cliff and Peter Haggett [14] have argued that technological innovation between the eighteenth and twentieth centuries — above all, the emergence of transoceanic steam travel — transformed global pathogen circulation by substantially reducing ship journey times. Paterson et al. [64] applied stochastic epidemiological modeling to assess the plausibility of pre-1850 measles transfer from the United Kingdom to Australia, arguing that shorter journey times and rising ship population susceptibility (in particular, greater numbers of child emigrants) were both necessary for pathogen introduction across this route. We offer a broader and more general assessment of the risks of shipborne pathogen introduction using the toolkit of contemporary theoretical ecology.

We present a stochastic SEIR (Susceptible, Exposed, Infected, Recovered) model which quantifies the probability of an outbreak lasting a given duration across a range of shipboard conditions. We investigate the respective contributions of pathogen natural history, onboard transmission intensity and density dependence, ship population size, and ship susceptibility to shaping overall outbreak duration. Finally, we use port arrivals data from Gold Rush-era San Francisco, California, 1850-1852, to explore the potential real-world implications of variation in journey length, ship size, and natural history for transoceanic pathogen circulation.

Our results indicate that shipborne pathogen introductions were neither trivial nor inevitable. Ships were not simple pathogen vectors: they were populations. As in all populations, the extinction and survival dynamics of pathogens on ships were complex and highly contingent on population size, composition, and mixing patterns. Thus, the history of transoceanic disease introduction is a story both of fundamental pathogen biology, and of human economics, technology, and behavior. Theoretical modelling can reveal how these forces interacted to shape global disease transmission.

Results

Basic Dynamics

Transoceanic pathogen introduction requires a chain of infections that lasts at least as long as a ship's journey time. To investigate the basic dynamics of shipboard outbreak duration, we simulate outbreaks in a fully susceptible population ($N = 100$) using a hypothetical pathogen which has characteristics typical of acute respiratory viruses (mean incubation and infectiousness periods of 5 days each) (Figure 1). We define outbreak duration as the time until nobody on board ship is infected with

Fig. 1. Basic Dynamics. (**A**) Outbreak duration, (**B**) outbreak size and (**C**) number of transmission generations by R_0, assuming a fully-susceptible population of $N = 100$ and a theoretical pathogen with $\mu_E = \mu_I = 5$ days and $k_E = k_I = 3$. Solid black lines show median outbreak duration, outbreak size, and number of generations. Top and bottom dashed lines respectively show 95^{th} and 5^{th} percentile outbreak duration, outbreak size, and number of generations. (**D**) Probability of at least one person in state E or state I ("introduction risk") for any given journey time by R_0, using the same population and pathogen parameters. (**E**) Outbreak length distribution by R_0 in a fully susceptible population of $N = 100$ for influenza, measles and smallpox, using epidemiological parameters detailed in Table S1.

the pathogen, that is, until $E = I = 0$.

Historical accounts indicate that transmission on board ships was substantially more intense than transmission in typical land settings (Text S1). Thus, we explore a broad range of transmission intensities. These are summarized by the epidemiological parameter R_0, or the average number of infections that a single person will produce in a fully-susceptible population.

We observe three outbreak duration regimes, all of which depend heavily on transmission intensity. Under strongly subcritical transmission ($R_0 \lesssim 0.8$), the majority of simulations result in no transmission beyond the index case (Figure 1A). These "single-generation" outbreaks last only as long as the course of infection in a single person, in this case an average of 10 days.

Under strongly supercritical transmission ($R_0 \gtrsim 5$), outbreaks are large and almost universally reach or exceed the threshold for ship herd immunity, $\frac{S}{N} < \frac{1}{R_0}$ (Figure 1B). This reliably results in 35-55 day outbreaks for the modeled scenario, with duration steadily decreasing as R_0 increases. Occasionally, simulations under strongly supercritical R_0 also result in short single- or two-generation outbreaks (Figure 1A, 1C); these are consistent with the occurrence of minor outbreaks due to random extinction in stochastic systems [9,90].

Values of R_0 near criticality ($0.8 \lesssim R_0 \lesssim 5$), produce the longest outbreaks, with median duration peaking around $R_0 = 2$. These are made possible by extended, multi-generational transmission chains (Figure 1C). Yet while near-critical conditions give rise to the longest outbreaks, they also result in the widest range of outbreak durations. An R_0 of 1 may result in outbreaks that last 150 days or more, but median outbreak duration under these conditions is just 14 days.

Transmission intensity modulates a pathogen's overall introduction risk — here defined as the net probability that at least one passenger is carrying the pathogen (i.e. in state E or state I) upon arrival, for any given journey length. Under strongly subcritical transmission (e.g. $R_0 = 0.5$), introduction risk decays rapidly with journey time, with 50% probability of introduction at 10 days and 25% probability at 17 days (Figure 1D). Under strongly supercritical transmission ($R_0 = 8$), pathogen introduction risk is sigmoidal: introduction is near certain ($\geq 95\%$) for journeys of 33 days or less, then falls rapidly for journey times exceeding this threshold. Introduction is least predictable for weakly supercritical values of R_0 ($R_0 = 2$). Here, many outbreaks end quickly due to random extinction (Figure 1A). Yet past this threshold, risk broadly plateaus until much longer journey times (~ 60 days), then declines with a long tail. Thus, the relative introduction risk of weakly versus strongly supercritical transmission depends on journey length. Strongly supercritical transmission is significantly more likely to result in pathogen introduction for journeys of 33 days or less, since intense transmission carries minimal risk of early extinction. But across journeys of 40 days or more, pathogen introduction is most likely under weakly supercritical transmission.

For real pathogens, introduction thresholds are governed by pathogen-specific natural history, above all by the durations of a pathogen's latent and infectious periods (Figure 1E). We explore outbreak length for influenza, measles, and smallpox at subcritical, near-critical, and strongly supercritical R_0 (Table S1). The results demonstrate that relative introduction risk can be often inferred from pathogen natural history, even in the absence of shipboard R_0 estimates. At any R_0, smallpox

typically survives longer on board a ship than measles, which in turn typically survives longer than influenza. Natural history also indicates some general introduction thresholds, which hold regardless of transmission intensity. For example, for a ship with 100 people on board, influenza introduction is extremely unlikely for journeys lasting longer than 100 days, regardless of R_0.

Incorporating Population Size and Susceptibility

Next, we expand our analysis beyond the unlikely scenario of one ship with $N = 100$ and 100% population-level susceptibility to consider the combined effects of ship population size, N, and initial proportion susceptible, $\frac{S(0)}{N}$, on ship outbreak duration.

In populations with some initial immunity to infection (i.e. where $\frac{S(0)}{N} < 1$), transmission intensity is most meaningfully measured as a pathogen's "effective" reproduction number, R_e. Because population immunity levels change over the course of an outbreak, this is commonly expressed as a function of time, i.e. $R_e(t)$. Notably, $R_e(t)$ is a linear function of a pathogen's basic reproduction number, R_0. Hence, $R_e(t) = \frac{S(t)}{N} R_0$, with critical transmission occurring at the threshold $R_e(t) = 1$. We consider shipboard transmission at $t = 0$, where $R_e(0) = \frac{S(0)}{N} R_0$.

First, we vary the total number of people who are initially susceptible, $S(0)$, while holding $R_e(0)$ constant. We do so by fixing $N = 1001$, choosing $S(0)$, and back-calculating R_0 to maintain the same effective rate of transmission. This results in a roughly log-linear relationship between initial susceptible population size and outbreak duration at critical and supercritical values of $R_e(0)$ (Figure 2A). At $R_e(0) = 1$, increasing $S(0)$ has little influence on median outbreak duration but substantially increases 95th percentile outbreak duration. At $R_e(0) = 2$, increasing $S(0)$ increases both median and 95th percentile outbreak times. Finally, at $R_e(0) = 8$, increasing $S(0)$ dependably increases median, 5^{th} percentile, and 95^{th} percentile outbreak times.

Next, we vary N as well as $S(0)$. This opens the question of what relationship we expect between N, S and R_0 in the unique environment on board a historical ship. Records from the time indicate that many vessels suffered from inadequate ventilation and extreme levels of crowding (Text S1). On land, these conditions generally give rise to "density-dependent" patterns of transmission, where contact rates scale linearly with population size ($R_0 \propto N$). Yet ships were also famously structured and compartmentalized environments, which typically align with assumptions of "frequency-dependent" transmission (Text S1); here contact rates are assumed to remain constant, regardless of total population size ($R_0 \perp\!\!\!\perp N$).

In practice, we expect that effective density dependence varied substantially according to ship layout and construction, social norms, and pathogen-side biology. Thus we consider three density dependence scenarios: classical density dependence ($R_0 \propto N$), classical frequency dependence ($R_0 \perp\!\!\!\perp N$), and an intermediate degree of density dependence ($R_0 \propto N^{0.5}$). Under each scenario, we explore the effect of initial ship susceptibility, $\frac{S(0)}{N}$, on median outbreak duration across several total population sizes, N.

In all circumstances, larger and more susceptible populations generally present greater risks of

Fig. 2. Effect of susceptible population size. (**A**) Outbreak duration by initial susceptible population ($S(0)$) and effective reproduction number ($R_e(0)$). We fix $N = 1001$ and back-calculate R_0 for each value of $S(0)$ to maintain constant $R_e(0)$. As above, we base simulations on a theoretical pathogen with $\mu_E = \mu_I = 5$ days. Solid black lines show median outbreak duration; top and bottom dashed lines show 95[th] and 5[th] percentile durations respectively. Colors match those used in Figure 1A-C and denote single-generation outbreaks (grey), outbreaks that reach herd immunity (red), and outbreaks that terminate before herd immunity is achieved (blue). (**B**) Median outbreak duration by density dependence, transmission intensity, ship size (N) and initial proportion susceptible ($S(0)/N$). Dashed vertical lines show $R_e(0) = 1$ by ship size. *Top row:* classical frequency dependence ($R_0 = \mu_I \beta_{fd}$), with β_{fd} values of 0.04 (left), 1 (center) and 2 (right) and $\mu_I = 5$. Dashed black vertical lines show $R_e(0) = 1$ for all N values. *Middle row:* intermediate density dependence ($R_0 = \mu_I(\beta_{fd}\beta_{dd}N)^{0.5}$), fixing $\beta_{fd} = 1$ and with β_{dd} of 0.01 (left), 0.02 (center) and 0.04 (right). *Bottom row:* classical density-dependent transmission ($R_0 = \beta_{dd}N$) with with β_{dd} of 0.01 (left), 0.02 (center) and 0.04 (right).

pathogen introduction across any given journey (Figure 2B). Yet they do so in different ways, and for different reasons.

Under classical frequency dependence, $R_0 = \mu_I \beta_{fd}$, where μ_I represents the average duration of an individual's infectious period and where β_{fd} represents the average number of onward infections that a single infected person would generate, per day, in a fully susceptible population. Critical transmission, $R_e(0) = 1$, occurs at the constant threshold $\frac{S(0)}{N} = (\mu_I \beta_{fd})^{-1}$. The threshold value of $\frac{S(0)}{N}$ required for $R_e(0) = 1$ is independent of total ship population, N. However, for any given $\frac{S(0)}{N}$, ships with greater N must have a proportionally greater number of susceptibles, $S(0)$. Since ships with greater $S(0)$ experience longer outbreaks at a given supercritical value of $R_e(0)$ (Figure 2A, ships with greater total populations display longer median outbreak times at $R_e(0) > 1$ (Figure 2B, top row).

Under classical density dependence, $R_0 = \mu_I \beta_{dd} N$, where β_{dd} represents the average proportion of a given population, N, that a single infected person would infect per day in a fully-susceptible population. Critical transmission occurs at the threshold $\frac{S(0)}{N} = (\mu_I \beta_{dd} N)^{-1}$. Multiplying both sides by a factor of N reveals that critical transmission depends solely on initial susceptible population size: $S(0) = (\mu_I \beta_{dd})^{-1}$. When N is large, this threshold for $S(0)$ represents a smaller fraction of the total population. But, in contrast to frequency-dependent transmission, $S(0)$ is constant at any given $R_e(0)$, and so peak outbreak duration does not vary across ships of different sizes. Rather, larger ship populations give rise to near-critical and supercritical transmission at lower values of $\frac{S(0)}{N}$ (Figure 2B, bottom row).

Under intermediate transmission, $R_0 = \mu_I(\beta_{fd}\beta_{dd}N)^{0.5}$. Critical transmission occurs at the threshold $\frac{S(0)}{N} = \mu_I^{-1}(\beta_{fd}\beta_{dd}N)^{-0.5}$, and at the total susceptibility level $S(0) = N^{0.5}\mu_I^{-1}(\beta_{fd}\beta_{dd})^{-0.5}$. Under this model, larger ship populations reach critical transmission at slightly lower initial proportions of susceptibility, have a critical $S(0)$ that scales sublinearly with N, and hence display slightly higher outbreak length for any given $R_e(0)$ (Figure 2B, middle row).

Finally, we note that regardless of density dependence, ships with a higher rate of contact (represented either by β_{fd} or by β_{dd}) require lower initial susceptibility for critical transmission (compare across rows in Figure 2B). Thus, ships with higher rates of social mixing (e.g. more crowded ships) require fewer susceptible people to achieve supercritical transmission, regardless of total ship population size.

Thus, even in the absence of detailed reconstructions of ship transmission patterns, we can conclude that ships with larger, more crowded populations presented greater risks of pathogen introduction — be this by increasing total persistence times, decreasing the susceptibility fraction required for critical transmission, or both. In practice, the risk associated with larger ship populations was almost certainly boosted further by an increased chance of carrying at least one infected person on departure. We do not account for this difference, instead conditioning on the assumption that all ships depart with a single infected individual. Yet in cases with low infection prevalence at the port of origin, this elevated chance of having at least one infected person on board at the time of departure would have substantially increased net introduction risk. Thus, ships with larger populations were

both more likely to depart with infection on board and more likely to sustain this infection outbreak until arrival.

Historical Applications

Voyage characteristics such as journey time, population size, and population susceptibility varied substantially across different time periods, transit routes, and ship constructions. We explore some of this variation, and its implications, using port arrivals data for Gold Rush-era San Francisco, 1850-1852, originally collected by historian and genealogist Louis J. Rasmussen (Figure 3) [68–70].

By the mid-nineteenth century, acute respiratory infections such as smallpox and measles had only recently begun to arrive across the Pacific basin. California saw its first region-wide outbreaks of smallpox and measles in 1806 and 1838, respectively [16,79,80]. Smallpox was first introduced to Australia in 1788 but did not see second introduction until 1829, while measles appears to have arrived for the first time in 1850 [42,64]. Several Pacific islands saw initial introductions well into the late nineteenth century, including Hawai'i (Smallpox, 1853); Easter Island (Smallpox, 1863); Fiji (Measles, 1875); and Tonga (Measles, 1893) [42].

During the years 1850-1852, passengers journeyed to San Francisco from across East Asia, Australasia, South America, and Europe. In an era preceding reliable transcontinental rail, ocean travel also provided one of the fastest and safest routes from eastern North America to the newly-established state of California [24]. Median sailing times ranged from 7 days (from Oregon Territory) to 190 days (from Liverpool, England), with considerable variation within routes (Figure 3A-B, Table S2). Longer-range sail voyages displayed an especially broad range of transit times. For example, sail journeys from New York City could be as long as 283 days (on the *Primoguet*) or as short as 89 days (on the *Flying Cloud* — reportedly "the fastest [sail] voyage ever recorded") [69].

Steam travel represented a phase transition in transoceanic pathogen circulation for several reasons. First, in most cases, the technology dramatically reduced journey times. Median transit times from Panama were 63 days by sail but just 20 days by steam. Meanwhile, steam reduced median journey times from Oregon from 7 days to just 3 days (Table S2). These shorter journey times would have increased risk of shipborne pathogen introduction significantly.

Second, steam ships transported some of the greatest numbers of passengers (Figure 3C). Steamers from Panama carried a median of 196 passengers and as many as 1,050, compared with a median of 53 and a maximum of 287 by sail (Table S2). Oregon steamers carried a median of 28 passengers and as many as 157, compared with a median of 4 and a maximum of just 12 by sail. Finally, steam vessels from New York City carried a median of 111 passengers and a maximum of 743, compared to a median of 5 and a maximum of 160 by sail. The only sail route that could compete with steam travel on passenger numbers was the route from Hong Kong, which transported a median of 201 passengers and a maximum of 553. As demonstrated above, larger passenger numbers would have substantially increased ships' capacity for sustained pathogen circulation.

Finally, steam travel represented some of the most frequent voyages (Figure 3C), resulting in a

Fig. 3. San Francisco Arrivals, June 1850 – June 1852. (A) Map of arrivals into San Francisco harbor, June 6th 1850 - June 9th 1852, with median journey times by ship technology. **(B)** Journey time, **(C)** passenger number and **(D)** number of voyages by origin port and by ship technology. Data from Louis J. Rasmussen's *San Francisco Passenger Lists* [68–70]

greater cumulative risk of pathogen introduction across any given period. Particularly striking are the 116 steam journeys from Panama that arrived between June 1850 and June 1852.

Taken together, the speed, passenger numbers, and frequency of steam travel all indicate that this technology carried a substantially elevated risk of pathogen introduction. Indeed, Rasmussen's notes offer frequent evidence of disease on steamships, especially the route from Panama. One ship travelling this route, the *SS Panama*, arrived in San Francisco on June 1, 1851 having experienced "a good deal of sickness among the passengers during the voyage, principally fever and dysentery". Eleven days later a second steamer, the *SS Gold Hunter* arrived with 163 passengers and one case of smallpox [69].

To explore possible differences in introduction risk across each route and type of ship, we simulate influenza, measles, and smallpox outbreaks across the full range of ship populations and journey times represented in the San Francisco dataset (Figure 4A). Here, contours represent pathogen introduction risk by journey time and by total ship population, assuming 5% population-level susceptibility, intermediate density dependence, $\beta_{dd} = 0.05$, and β_{fd} calculated according to standard literature R_0 values (Table S1). We overplot individual journeys into San Francisco to assess pathogen introduction risk across each route (Figure 4A). We plot a selection of routes on each panel for visual clarity. However, the observations below concern introduction risk for all pathogens across all routes travelled. For select voyages we also provide numerical introduction risk estimates for each pathogen in Table 1.

Influenza's relatively low R_0 and extremely fast generation period result in a very low risk of introduction into San Francisco from any origin port except Oregon, Panama and Hawai'i. Even then, only the fastest voyages presented any significant risk of pathogen transfer. Had a person with influenza been present on board the *Columbia* steam ship (3 days, 74 passengers) at its time of departure from Oregon, we estimate a 68% risk of introduction into San Francisco (Table 1). By contrast, we estimate just a 11% risk for the *Tarquina* sail ship from Oregon (7 days, 11 passengers), and a < 1% risk on the *Columbus* steam ship from Panama (21 days, 225 passengers) and the *Baltimore* sail ship from Hawai'i (14 days, 29 passengers).

Measles, with longer latent and infectious periods, presents moderate introduction risks across all journey times $\lesssim$ 25 days (Figure 4A, Table 1) – consistent with this pathogen's range of durations for

Fig. 4. Historical Applications. Introduction risk for influenza, measles, and smallpox by journey time and by total ship population, N, assuming 5% initial population-level susceptibility, intermediate density dependence ($q = 0.5$), $\beta_{dd} = 0.05$, and μ_E, μ_I, and β_{fd} according to consensus natural history parameters (Table S1), with β_{fd} back-calculated as $1/\mu_I$ times a pathogen's typical R_0 on land. (**A**) overplots data on San Francisco Port arrivals, June 1850 – June 1852. Here, total population (N) represents only the passengers on board each ship, as crew data is not available. Introduction risks for all three pathogens are shown for 14 selected voyages in Table 1; these voyages are indicated with black crosses. (**B**) overplots selected historical journeys, 1492–1918, chosen to be indicative of the broad trends in transoceanic shipping. N represents the combined totals of passengers and crew. Sources and further data are available in Table S3. Numerical introduction risk estimates for (**B**) are provided in Table 2

.

single-generation outbreaks (Figure 1) This range includes the vast majority of journeys originating from Oregon (by steam or sail), Hawai'i (by steam or sail), and Panama (by steam). Additionally, we estimate plausible introductions from other ports of origin if ships transported large populations. Had the *Iowa* sail ship (54 days, 377 people) departed Hong Kong with a measles patient on board, we estimate a 61% chance of introduction despite the long journey (Table 1). Had the *Golden Gate* steam ship (65 days, 458 passengers) departed New York City with one infected passenger, we likewise estimate an 63% introduction risk.

Smallpox has a substantially longer generation period than either measles or influenza ($\mu_E = 12$ days; $\mu_I = 17.5$ days). Consequently, journeys of $\lesssim 40$ days present a moderate introduction risk at any ship population size. This expands the introduction range described above for measles to include several sail ships originating in Panama and Valparaíso. As before, we estimate higher introduction risks for ships with larger populations (Table 1). Yet since smallpox is less transmissible than measles, ships require larger population sizes to achieve a equivalent $R_e(0)$. Thus, in some cases highly-populated ships present a lower risk of introducing smallpox than they did measles. For instance, had the *Golden Gate* departed New York City with one infected patient on board, we estimate a 54% risk of smallpox introduction (Table 1).

Finally, we use these analyses to inform the plausibility of ship-borne pathogen transfer across a selection of historical voyages, chosen to reflect the variety of shipping routes, technologies, and practices between the 15$^{\text{th}}$ and 20$^{\text{th}}$ centuries (Figure 4B, Table S3). For these analyses we again assume a 5% rate of susceptibility, although in practice we expect this rate varied significantly by location, time period and ship population.

Under these assumptions, early transatlantic voyages of exploration could plausibly have introduced measles or smallpox to their ports of arrival (Figure 4B, Table 2). We estimate a 24% chance of measles introduction and 33% chance of smallpox introduction had Christopher Columbus's 1492 voyage on *Santa María* (35 days, 41 people) departed with one case on board, and a 2% and 7% introduction risk of smallpox and measles respectively on John Cabot's 1497 exploration on the *Matthew* (54 days, 20 people). However introduction risks for both pathogens were substantially higher on the transatlantic slave trade ship *Diana*, which carried 448 enslaved people and crew from Îles del Los, off the coast of West Africa, to Curaçao, in the Caribbean: an 67% risk for measles and a 61% risk for smallpox, had one person been infected at the time of departure.

Meanwhile, the lengthy journey times of the *Empress of China* trade ship (184 days), which traveled from New York City to present-day Macao and the *America* convict ship (180 days), which traveled from the United Kingdom to Australia, suggest a compelling explanation for the substantially later introduction of smallpox and measles to the South Pacific. Even with 175 passengers, a voyage such as the *America*'s is outside the range of plausible influenza and measles introduction, and extremely unlikely to introduce smallpox (0.2% risk).

Our analyses indicate that by far the greatest introduction risk of smallpox and measles – and the only plausible influenza introduction – came from fast, highly-populated ships such as the WW1 troop ship HMAT *Boonah* (1,095 passengers and crew), here undertaking a 17-day journey from South

Table 1. Numerical introduction risk estimates for influenza, measles, and smallpox across selected voyages to San Francisco, 1850-1852

| Route | Vessel | Type | Duration | N | Pathogen introduction risk | | |
					Influenza	Measles	Smallpox
New York	*Elsinore*	Sail	195 days	17	< 0.001	< 0.001	< 0.001
New York	*Golden Gate*	Steam	66 days	458	< 0.001	0.629	0.541
Valparaíso	*Aurora*	Sail	59 days	16	< 0.001	0.012	0.055
Valparaíso	*Huntress*	Sail	101 days	175	< 0.001	0.055	0.097
Panama	*Columbus*	Steam	21 days	225	0.001	0.697	0.739
Panama	*Margaret*	Sail	110 days	143	< 0.001	0.016	0.036
Oregon	*Columbia*	Steam	3 days	74	0.682	0.996	0.999
Oregon	*Tarquina*	Sail	7 days	11	0.110	0.936	0.980
Hawai'i	*Edgar*	Sail	40 days	142	< 0.001	0.421	0.426
Hawai'i	*Baltimore*	Sail	14 days	29	0.002	0.673	0.821
Sydney	*Cameo*	Sail	112 days	28	< 0.001	< 0.001	< 0.001
Sydney	*Edgar*	Sail	83 days	142	< 0.001	0.079	0.117
Hong Kong	*George Pollock*	Sail	56 days	47	< 0.001	0.050	0.104
Hong Kong	*Iowa*	Sail	54 days	377	< 0.001	0.609	0.521

Table 2. Numerical introduction risk estimates for influenza, measles, and smallpox across selected historical voyages, 1492-1918

| Year | Vessel | Type | Duration | N | Pathogen introduction risk | | |
					Influenza	Measles	Smallpox
1492	*Santa María*	Sail	35 days	41	< 0.001	0.238	0.334
1497-98	*Matthew*	Sail	54 days	20	< 0.001	0.022	0.079
1609	*Sea Venture*	Sail	53 days	150	< 0.001	0.331	0.330
1620	*Mayflower*	Sail	66 days	127	< 0.001	0.135	0.175
1778	*Diana*	Sail	41 days	413	< 0.001	0.673	0.617
1784	*Empress of China*	Sail	184 days	34	< 0.001	< 0.001	< 0.001
1832	*America*	Sail	180 days	175	< 0.001	< 0.001	0.002
1918	HMAT *Boonah*	Steam	17 days	1095	0.036	0.905	0.890

Africa to Australia. Had this ship departed with one infected person on board, it would have had a 4% risk of introducing influenza and around a 90% chance of introducing either measles or smallpox into its destination. Thus, such a combination of fast transit and extremely large passenger populations substantially increased both the magnitude of introduction risk for moderately fast-burning pathogens (measles and smallpox) and expanded the range of potential introduction to include pathogens (e.g. influenza) with much faster life cycles.

Discussion

Canonically, transoceanic pathogen transfer has been told as a story of colonial exploration. Our analysis indicates that introductions of smallpox and, to a lesser extent, measles via early transatlantic voyages of discovery were plausible, though by no means guaranteed. Depending on weather, these voyages could last just 5-10 weeks — a reasonable time frame for these pathogens to persist on board a ship [31]. Yet early transatlantic introductions of faster-burning pathogens such as influenza are unlikely, as are introductions across longer-range journeys such as voyages beyond the Atlantic Ocean. Moreover, unless early exploratory voyages had high levels of crew susceptibility, it is unlikely that

smaller ship populations such as the *Matthew*'s (18 people) or the *Santa María*'s (42 people), could have reached the supercritical values of $R_e(0)$ necessary for measles outbreaks longer than ~ 30 days and smallpox outbreaks longer than ~ 50 days.

More recently, transoceanic pathogen transfer has become a story of technological innovation. Our work supports and extends Cliff and Haggett [14]'s argument that steam technology transformed rates of transoceanic pathogen transfer. Steam ships travelled more quickly, could carry greater numbers of passengers and, in the case of Gold Rush-era San Francisco, made more frequent voyages. Under the right conditions, this could have increased both the rate and the geographic range of transoceanic pathogen transfer substantially (Figure 4A; Table 1).

However, steam travel was not the only force that could revolutionise global pathogen circulation. Our analysis indicates that historical processes involving large-scale movement of people were potential watersheds for transoceanic pathogen ecology. This mirrors a longstanding historical literature demonstrating links between war, colonialism, and infectious disease [27,75], as well as recent phylogenetic work indicating that the transatlantic slave trade introduced parvovirus and hepatitis-B virus to Mexico [41]. In the case of 1850s California, ship population size could easily have been the difference between plausible introduction and epidemiological isolation. In 1852, two ships sailed from Hong Kong, the *Catalpa* and the *Iowa*. Both displayed similar transit times into San Francisco: 60 days and 54 days, respectively [68,69]. Yet while the *Catalpa* carried "Chinese merchandise, rice, cordage, and assorted goods" – along with one solitary passenger – the *Iowa* brought "377 unidentified in steerage", likely Chinese people bound for California's gold fields [62]. As our analyses show, the presence of 377 people on board transformed the *Iowa*'s capacity to sustain outbreaks of smallpox and measles across the journey from Hong Kong to San Francisco (Figure 4A, Table 1).

Our study raises several questions that require further consideration. One concerns the mechanics of shipboard transmission. Little is known concerning either the density dependence or the intensity of transmission on board historical vessels – although our results suggest that crowded ships with larger and more susceptible populations presented greater risks regardless of the precise form of density dependence, and that rough introduction thresholds at any transmission intensity can be inferred from pathogen natural history (Figure 1, Figure 2).

A related and more challenging question concerns the extent of population structure on board historical ships. Highly compartmentalized populations could have prolonged ship outbreaks; to some extent, these effects can be approximated in our model by reducing R_e. Yet while ships were famously hierarchical environments, transmission patterns may not have followed directly from social structure; on a ship with poor ventilation, airborne transmission could plausibly have been homogeneous regardless of social behaviour. Thus, while population structure almost certainly shaped outbreak duration, incorporating these effects is challenging in the absence of high-resolution outbreak data from a given ship.

Our work sheds light on how shipboard transmission dynamics shape introduction risk, but reconstructing historically accurate circulation rates would require more information regarding pathogen dynamics in source populations. This matters both for inferring likely immunity rates in ship pop-

ulations, and for assessing the probability of at least one infected individual on board ship at the point of departure. Longitudinal mortality data exists for diseases such as smallpox and measles, especially in European and North American contexts, with particularly well-preserved time series in the London Bills of Mortality [50,52]. Reconstructing historical prevalence and immunity landscapes from these sources is difficult, but is critical for estimating realistic pathogen transfer rates pre-20th century contexts.

A related question concerns the contribution of partly-immune individuals to pathogen circulation within a given population. The ability of partly-immune people to be infected and transmit infection has long been recognized as an important driver of influenza [65] and smallpox [32] epidemiology. Partial immunity also provides a compelling explanation for recent resurgences in mumps [26] and pertussis [1,72]. Moreover, it is plausible that the contribution to transmission from partly-immune individuals was more significant on board a ship than it was on land, due to extended exposures or large infectious doses. This possibility – and the influence of partial immunity on outbreak duration more broadly – require further investigation.

Finally, our analysis specifically considers human-to-human direct transmission of infectious pathogens. Pathogens with significant food-, water- or fomite-borne transmission (e.g. cholera, *Salmonella*), with vector-borne transmission (e.g. malaria, yellow fever), with multi-species transmission (e.g. plague), or infecting only non-human animals (e.g. rinderpest) will require modified modeling approaches and likely additional historical data. This issue is also pertinent to smallpox, for which the extent of fomite transmission is unclear. Recent research indicates that orthopoxviruses can remain viable on surfaces for weeks [94]. The World Health Organization's smallpox eradication campaign found that fomites caused only a small minority of outbreaks [32], but in the context of historical pathogen circulation even rare introductions can be impactful.

Our model offers a general assessment of outbreak duration in a closed population, which holds significance beyond historical systems. Understanding infection persistence in discrete subpopulations is critical for studying pathogen circulation in any system with limited host connectivity, from wildlife populations [20,21] to agricultural biosecurity [5,13] to human populations ditributed across regions [37,83]

These findings carry important historical implications, connect to present-day disease dynamics, and may, some day, inform interplanetary risk of pathogen spread. Centuries before the present-day upheaval of air travel, technological advancements and large-scale human re-locations combined to transform global pathogen ecology [4,12,15,83]. Yet this process was almost certainly lengthy, geographically uneven, and contingent on complex interplays between technology, shipping practices, and specific pathogen biology. This presents a rich avenue for collaboration between ecologists, historians, and social scientists. How do global social, economic and technological forces combine to shape global pathogen ecology – and with what consequences along the way for the world's people, places, and pathogens?

Materials and Methods

Model Description

We simulate shipboard outbreaks using a stochastic SEIR model ext S3). We implement continuous-time stochastic simulations in R with the Gillespie Algorithm, using the package GillespieSSA [66]. All simulations assume a single index case in state E at the time of departure. We define outbreak duration as the time until both state E and state I contain zero individuals.

To achieve a more realistic depiction of the time course of infection, we use the Linear Chain Trick to make dwell times in state E and state I Erlang-distributed [44]. For all simulations, we use shapes $k_E = k_I = 3$ and rates k_E/μ_E and k_I/μ_I for states E and I respectively. This technique gives a unimodal distribution with a long right-hand tail, such that disease progression is relatively constrained in most individuals, but occasional individuals experience substantially longer periods of incubation or infectiousness [85]. We assume that state E is pre-symptomatic and that initially infected individuals could board ship at any point during this period, randomly assigning index cases across sub-states $E_1, E_2, ..., E_{k_E}$ at the point of departure.

Our model also tracks infection across pathogen generations, when needed. The I_n infectious individuals from generation n produce new exposed individuals E_{n+1}, which represent the $(n+1)^{\text{st}}$ generation of infections.

To account for uncertainty and variation in the density dependence of shipboard contact rates, our model uses a flexible depiction of density dependence encoded by the equation:

$$R_0 = \mu_I (\beta_{dd} N)^q (\beta_{fd})^{1-q}$$

R_0 is the pathogen's basic reproduction number on board a given ship. This represents the average number of infections that an infected person generates in a fully-susceptible population, where μ_I is the average period of infectiousness. The density dependence of transmission is adjusted with the parameter q, with $q = 1$ representing classical density-dependent transmission ($R_0 \propto N$), $q = 0$ representing classical frequency-dependent transmission ($R_0 \perp\!\!\!\perp N$), and $0 < q < 1$ representing intermediate density dependence ($R_0 \propto N^q$). The parameters β_{dd} and β_{fd} modulate the intensity of transmission under each density dependence pole — intuitively, the proportion (β_{dd}) and the raw number (β_{fd}) of people on board ship that a single infected individual will infect per day, on average, in a fully-susceptible population.

For these analyses we assume no formal relationship between β_{dd} and β_{fd}. Instead, we adjust each parameter explicitly to assess the effect of different combinations of q, β_{fd}, β_{dd} and N on outbreak duration. In cases where N is constant and where we do not explore the effect of density dependence, we set $q = 1$ and assume that $R_0 = \beta_{dd} N$; mathematically, this is equivalent to setting $q = 0$ with β_{fd} fixed at $\beta_{fd} = \beta_{dd} N$.

Historical Data

To provide real-life context for our theoretical results, we collected data on ship arrivals into the port of San Francisco between June 6 1850 and June 19 1852 from volumes I, II and III of genealogist and historian Louis. J. Rasmussen's reference book *San Francisco Ship Passenger Lists* [68–70]. We recorded the port of origin, the ship type, the journey time, and the number of passengers for ships originating from seven locations: Hawai'i; Hong Kong; Oregon Territory; New York City; Sydney, Australia; Valparaíso, Chile; and Liverpool, England. In the few cases where Rasmussen reports ships making multiple stops at one or more of these locations in the course of their voyage, we record both the journey time into San Francisco from a ship's port of origin and, where available, journey times into San Francisco from the intermediate port(s). We exclude ships where substantial numbers of people ($N > 10$) boarded during a voyage, as our model does not account for changes in population size subsequent to the initial point of departure.

For almost all ships, Rasmussen provides passenger numbers but not numbers of crew. We assume that in most cases, crew (i) represented a small proportion of a ship's total population; (ii) were, as professional sailors, more likely to possess immunity to common maritime infections, and so represented an even smaller proportion of a ship's susceptible people. Thus, in the absence of crew size data, analyses considering population size of vessels arriving into San Francisco approximate N as the total number of passengers on board each ship.

References

1. Benjamin M. Althouse and Samuel V. Scarpino. Asymptomatic transmission and the resurgence of Bordetella pertussis. *BMC Medicine*, 13(1):146, December 2015. doi: 10.1186/s12916-015-0382-8.

2. Anonymous. Another Instance of Pestilence Engendered in a Ship Crowded with Passengers from Ireland. *The Medical Repository (And Review Of American Publications On Medicine, Surgery And The Auxiliary Of Science)*, V(II):234–236, 1802.

3. Parham Azimi, Zahra Keshavarz, Jose Guillermo Cedeno Laurent, Brent Stephens, and Joseph G. Allen. Mechanistic transmission modeling of COVID-19 on the Diamond Princess cruise ship demonstrates the importance of aerosol transmission. *Proceedings of the National Academy of Sciences*, 118(8):e2015482118, February 2021. doi: 10.1073/pnas.2015482118.

4. Rachel E. Baker, Ayesha S. Mahmud, Ian F. Miller, Malavika Rajeev, Fidisoa Rasambainarivo, Benjamin L. Rice, Saki Takahashi, Andrew J. Tatem, Caroline E. Wagner, Lin-Fa Wang, Amy Wesolowski, and C. Jessica E. Metcalf. Infectious disease in an era of global change. *Nature Reviews Microbiology*, 20 (4):193–205, April 2022. doi: 10.1038/s41579-021-00639-z.

5. Lindsay M. Beck-Johnson, Erin E. Gorsich, Clayton Hallman, Michael J. Tildesley, Ryan S. Miller, and Colleen T. Webb. An exploration of within-herd dynamics of a transboundary livestock disease: A foot and mouth disease case study. *Epidemics*, 42:100668, March 2023. doi: 10.1016/j.epidem.2023.100668.

6. Trevor Bedford, Steven Riley, Ian G. Barr, Shobha Broor, Mandeep Chadha, Nancy J. Cox, Rodney S. Daniels, C. Palani Gunasekaran, Aeron C. Hurt, Anne Kelso, Alexander Klimov, Nicola S. Lewis, Xiyan Li, John W. McCauley, Takato Odagiri, Varsha Potdar, Andrew Rambaut, Yuelong Shu, Eugene Skepner, Derek J. Smith, Marc A. Suchard, Masato Tashiro, Dayan Wang, Xiyan Xu, Philippe Lemey, and Colin A. Russell. Global circulation patterns of seasonal influenza viruses vary with antigenic drift. *Nature*, 523 (7559):217–220, July 2015. doi: 10.1038/nature14460.

7. Matthew Biggerstaff, Simon Cauchemez, Carrie Reed, Manoj Gambhir, and Lyn Finelli. Estimates of the reproduction number for seasonal, pandemic, and zoonotic influenza: a systematic review of the literature. *BMC infectious diseases*, 14:480, September 2014. doi: 10.1186/1471-2334-14-480.

8. I. W. Brewer. Report of Epidemic of 'Spanish Influenza', Which Occured at Camp A. A. Humphreys, VA., During September and October, 1918. *Journal of Laboratory and Clinical Medicine*, 4:87–111, December 1918.

9. Tom Britton. Stochastic epidemic models: A survey. *Mathematical Biosciences*, 225(1):24–35, May 2010. doi: 10.1016/j.mbs.2010.01.006.

10. Serina Chang, Emma Pierson, Pang Wei Koh, Jaline Gerardin, Beth Redbird, David Grusky, and Jure Leskovec. Mobility network models of COVID-19 explain inequities and inform reopening. *Nature*, 589 (7840):82–87, January 2021. doi: 10.1038/s41586-020-2923-3.

11. Vivek Charu, Scott Zeger, Julia Gog, Ottar N. Bjørnstad, Stephen Kissler, Lone Simonsen, Bryan T. Grenfell, and Cécile Viboud. Human mobility and the spatial transmission of influenza in the United States. *PLOS Computational Biology*, 13(2):e1005382, February 2017. doi: 10.1371/journal.pcbi.1005382.

12. Matteo Chinazzi, Jessica T. Davis, Marco Ajelli, Corrado Gioannini, Maria Litvinova, Stefano Merler, Ana Pastore y Piontti, Kunpeng Mu, Luca Rossi, Kaiyuan Sun, Cécile Viboud, Xinyue Xiong, Hongjie Yu, M. Elizabeth Halloran, Ira M. Longini, and Alessandro Vespignani. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science*, 368(6489):395–400, April 2020. doi: 10.1126/science.aba9757.

13. Irina Chis Ster, Peter J. Dodd, and Neil M. Ferguson. Within-farm transmission dynamics of foot and mouth disease as revealed by the 2001 epidemic in Great Britain. *Epidemics*, 4(3):158–169, August 2012. doi: 10.1016/j.epidem.2012.07.002.

14. Andrew Cliff and Peter Haggett. Time, travel and infection. *British Medical Bulletin*, 69:87–99, 2004. doi: 10.1093/bmb/ldh011.

15. Vittoria Colizza, Alain Barrat, Marc Barthélemy, and Alessandro Vespignani. The role of the airline transportation network in the prediction and predictability of global epidemics. *Proceedings of the National Academy of Sciences*, 103(7):2015–2020, February 2006. doi: 10.1073/pnas.0510525103.

16. Sherburne Friend Cook. *The Conflict Between the California Indian and White Civilization*. University of California Press, Oakland, 1976.

17. Valentina Costantino, Mohana P Kunasekaran, Abrar A Chughtai, and Chandini R MacIntyre. How Valid Are Assumptions About Re-emerging Smallpox? A Systematic Review of Parameters Used in Smallpox Mathematical Models. *Military Medicine*, 183(7-8):e200–e207, July 2018. doi: 10.1093/milmed/usx092.

18. Alfred W. Crosby. *The Columbian Exchange: Biological and Cultural Consequences of 1492*. Greenwood Publishing Company, Westport, CT, 1972.

19. Eleanor R. Cross, Larry A. Hermansen, William M. Pugh, Martin R. White, Curtis Hayes, and Kenneth C. Hyams. Upper Respiratory Disease in Deployed U.S. Navy Shipboard Personnel. *Military Medicine*, 157(12):649–651, December 1992. doi: 10.1093/milmed/157.12.649.

20. Paul C. Cross, James O. Lloyd-Smith, Philip L. F. Johnson, and Wayne M. Getz. Duelling timescales of host movement and disease recovery determine invasion of disease in structured populations. *Ecology Letters*, 8(6):587–595, 2005. doi: 10.1111/j.1461-0248.2005.00760.x.

21. Paul C. Cross, Philip L. F. Johnson, James O. Lloyd-Smith, and Wayne M. Getz. Utility of R0 as a predictor of disease invasion in structured populations. *Journal of The Royal Society Interface*, 4(13):315–324, November 2006. doi: 10.1098/rsif.2006.0185.

22. J.H.L. Cumpston. *The History of Small-pox in Australia, 1788-1908*. Albert J. Mullett, Government Printer, Melbourne, 1914.

23. John Howard Lidgett Cumpston. *Influenza and Maritime Quarantine in Australia*. Issued under the Authority of the Minister for Trade and Customs, Albert J. Mullett, Government Printer, Melbourne, 1919.

24. James P. Delgado. *To California by Sea: a Maritime History of the California Gold Rush*. University of South Carolina Press, Columbia, 1990.

25. Kieran Doherty. *Sea Venture: Shipwreck, Survival, and the Salvation of Jamestown*. St. Martin's Publishing Group, New York, 2013.

26. Matthew Donahue, Blake Hendrickson, Derek Julian, Nicholas Hill, Julie Rother, Samir Koirala, Joshua L. Clayton, Thomas Safranek, and Bryan Buss. Multistate Mumps Outbreak Originating from Asymptomatic Transmission at a Nebraska Wedding — Six States, August–October 2019. *Morbidity and Mortality Weekly Report*, 69(22):666–669, June 2020. doi: 10.15585/mmwr.mm6922a2.

27. Jim Downs. *Maladies of Empire: How Colonialism, Slavery, and War Transformed Medicine*. Harvard University Press, September 2021.

28. Sheldon F. Dudley. The Biology of Epidemic Influenza, Illustrated by Naval Experience. *Proceedings of the Royal Society of Medicine*, 14:37–50, May 1921. doi: 10.1177/003591572101402304.

29. Kenneth C. Earhart, Christine Beadle, Larry K. Miller, Martin W. Pruss, Gregory C. Gray, Elizabeth K. Ledbetter, and Mark R. Wallace. Outbreak of influenza in highly vaccinated crew of U.S. Navy ship. *Emerging Infectious Diseases*, 7(3):463–465, 2001.

30. Peter Eicher. *Raising the Flag: America's First Envoys in Faraway Lands*. University of Nebraska Press, Lincoln, 2018.

31. J. H. Elliott. *Empires of the Atlantic world: Britain and Spain in America, 1492-1830*. Yale University Press, New Haven, 2006.

32. Frank Fenner, Donald A. Henderson, Isao Arita, Zdenek Jezek, Ivan Danilovich Ladnyi, and World Health Organization. Smallpox and its Eradication. Technical report, World Health Organization, 1988.

33. Flavio Finger, Tina Genolet, Lorenzo Mari, Guillaume Constantin de Magny, Noël Magloire Manga, Andrea Rinaldo, and Enrico Bertuzzo. Mobile phone data highlights the role of mass gatherings in the spreading of cholera outbreaks. *Proceedings of the National Academy of Sciences*, 113(23):6421–6426, June 2016. doi: 10.1073/pnas.1522305113.

34. Rebecca Fraser. *The Mayflower: The Families, the Voyage, and the Founding of America*. St. Martin's Publishing Group, New York, 2017.

35. Robert H. Fuson. *The Log of Christopher Columbus*. McGraw-Hill Companies, New York, 1987.

36. Albert L. Gihon. Naval Hygeine. In *A Reference Handbook of the Medical Sciences: Embracing the Entire Range of Scientific and Practical Medicine and Allied Science*, pages 766–773. William Wood & Company, New York, 1886.

37. Bryan T. Grenfell and Benjamin M. Bolker. Cities and villages: infection hierarchies in a measles metapopulation. *Ecology Letters*, 1(1):63–70, 1998. doi: 10.1046/j.1461-0248.1998.00016.x.

38. Bryan T. Grenfell and John Harwood. (Meta)population dynamics of infectious diseases. *Trends in Ecology & Evolution*, 12(10):395–399, October 1997. doi: 10.1016/S0169-5347(97)01174-9.

39. Bryan T. Grenfell, Ottar N. Bjørnstad, and J. Kappey. Travelling waves and spatial hierarchies in measles epidemics. *Nature*, 414(6865):716–723, December 2001. doi: 10.1038/414716a.

40. Fiona M Guerra, Shelly Bolotin, Gillian Lim, Jane Heffernan, Shelley L Deeks, Ye Li, and Natasha S Crowcroft. The basic reproduction number (R 0) of measles: a systematic review. *The Lancet Infectious Diseases*, 17(12):e420–e428, December 2017. doi: 10.1016/S1473-3099(17)30307-9.

41. Axel A Guzmán-Solís, Viridiana Villa-Islas, Miriam J Bravo-López, Marcela Sandoval-Velasco, Julie K Wesp, Jorge A Gómez-Valdés, María De La Luz Moreno-Cabrera, Alejandro Meraz, Gabriela Solís-Pichardo, Peter Schaaf, Benjamin R TenOever, Daniel Blanco-Melo, and María C Ávila Arcos. Ancient viral genomes reveal introduction of human pathogenic viruses into Mexico during the transatlantic slave trade. *eLife*, 10:e68612, August 2021. doi: 10.7554/eLife.68612.

42. Peter Haggett. The Invasion of Human Epidemic Diseases into Australia, New Zealand, and the Southwest Pacific: The Geographical Context. *New Zealand Geographer*, 49(2):40–47, 1993. doi: 10.1111/j.1745-7939.1993.tb02038.x.

43. Sophia Hatzianastasiou, Varvara A. Mouchtouri, Androula Pavli, Maria Tseroni, Spyros Sapounas, Charalampos Vasileiou, Katerina Dadouli, Maria Kyritsi, Michalis Koureas, Panagiotis Prezerakos, Matthaios Speletas, Georgios Panagiotakopoulos, Sotirios Tsiodras, and Christos Hadjichristodoulou. COVID-19 Outbreak on a Passenger Ship and Assessment of Response Measures, Greece, 2020. *Emerging Infectious Diseases*, 27(7):1927–1930, July 2021. doi: 10.3201/eid2707.210398.

44. Paul J. Hurtado and Adam S. Kirosingh. Generalizations of the 'Linear Chain Trick': incorporating more flexible dwell time distributions into mean field ODE models. *Journal of Mathematical Biology*, 79(5): 1831–1883, October 2019. doi: 10.1007/s00285-019-01412-w.

45. David Igler. *The Great Ocean: Pacific Worlds from Captain Cook to the Gold Rush*. Oxford University Press, Oxford, 2016.

46. Evan T. Jones, Margaret M. Condon, and John Cabot. *Cabot and Bristol's Age of Discovery: The Bristol Discovery Voyages 1480-1508*. University of Bristol Press, Bristol, United Kingdom, 2016.

47. Matthew R. Kasper, Jesse R. Geibe, Christine L. Sears, Asha J. Riegodedios, Tina Luse, Annette M. Von Thun, Michael B. McGinnis, Niels Olson, Daniel Houskamp, Robert Fenequito, Timothy H. Burgess, Adam W. Armstrong, Gerald DeLong, Robert J. Hawkins, and Bruce L. Gillingham. An Outbreak of Covid-19 on an Aircraft Carrier. *New England Journal of Medicine*, 383(25):2417–2426, December 2020. doi: 10.1056/NEJMoa2019375.

48. Matt J. Keeling, Ottar N. Bjørnstad, and Bryan T. Grenfell. Metapopulation Dynamics of Infectious Diseases. In Ilkka Hanski and Oscar E. Gaggiotti, editors, *Ecology, Genetics and Evolution of Metapopulations*, chapter 17, pages 415–445. Academic Press, Burlington, January 2004. doi: 10.1016/B978-012323448-3/50019-2.

49. Paul Kelton. *Cherokee Medicine, Colonial Germs: An Indigenous Nation's Fight against Smallpox, 1518–1824*. University of Oklahoma Press, Norman, 2015.

50. Olga Krylova and David J. D. Earn. Patterns of smallpox mortality in London, England, over three centuries. *PLOS Biology*, 18(12):e3000506, December 2020. doi: 10.1371/journal.pbio.3000506.

51. Lincoln L. H. Lau, Benjamin J. Cowling, Vicky J. Fang, Kwok-Hung Chan, Eric H. Y. Lau, Marc Lipsitch, Calvin K. Y. Cheng, Peter M. Houck, Timothy M. Uyeki, J. S. Malik Peiris, and Gabriel M. Leung. Viral Shedding and Clinical Illness in Naturally Acquired Influenza Virus Infections. *The Journal of Infectious Diseases*, 201(10):1509–1516, May 2010. doi: 10.1086/652241.

52. Hyeok Jun Lee. *Measles and Whooping Cough in London 1750-1900, and the Role of Immune Amnesia in Recurrent Epidemics*. Thesis, McMaster University, Ontario, 2023.

53. Justin Lessler, Nicholas G. Reich, Ron Brookmeyer, Trish M. Perl, Kenrad E. Nelson, and Derek AT Cummings. Incubation periods of acute respiratory viral infections: a systematic review. *The Lancet Infectious Diseases*, 9(5):291–300, May 2009. doi: 10.1016/S1473-3099(09)70069-6.

54. Haruka Maeda, Eiichiro Sando, Michiko Toizumi, Yuzo Arima, Tomoe Shimada, Takeshi Tanaka, Masato Tashiro, Ayumi Fujita, Katsunori Yanagihara, Hayato Takayama, Ikkoh Yasuda, Nobuyuki Kawachi, Yoshitaka Kohayagawa, Maiko Hasegawa, Katsuaki Motomura, Rie Fujita, Katsumi Nakata, Jiro Yasuda, Koichi Morita, Shigeru Kohno, Koichi Izumikawa, Motoi Suzuki, and Konosuke Morimoto. Epidemiology of Coronavirus Disease Outbreak among Crewmembers on Cruise Ship, Nagasaki City, Japan, April 2020. *Emerging Infectious Diseases*, 27(9), 2021. doi: 10.3201/eid2709.204596.

55. L. Mari, E. Bertuzzo, L. Righetto, R. Casagrandi, M. Gatto, I. Rodriguez-Iturbe, and A. Rinaldo. Modelling cholera epidemics: the role of waterways, human mobility and sanitation. *Journal of The Royal Society Interface*, 9(67):376–388, February 2012. doi: 10.1098/rsif.2011.0304.

56. Frank Marryat. *Mountains and Molehills, Or, Recollections of a Burnt Journal*. Longman, Brown, Green, and Longmans, London, 1855.

57. Aims McGuinness. *Path of Empire: Panama and the California Gold Rush*. Cornell University Press, December 2007.

58. John R. McNeill. Disease Environments in the Carribbean to 1850. In *Sea and Land: An Environmental History of the Caribbean*, pages 130–186. Oxford University Press, Oxford, 2022.

59. Joy M. Miller, Theresa W. S. Tam, Susan Maloney, Keiji Fukuda, Nancy Cox, James Hockin, Daniel Kertesz, Alexander Klimov, and Martin Cetron. Cruise Ships: High-Risk Passengers and the Global Spread of New Influenza Viruses. *Clinical Infectious Diseases*, 31(2):433–438, August 2000. doi: 10.1086/313974.

60. Arezou Minooee and Leland S. Rickman. Infectious Diseases on Cruise Ships. *Clinical Infectious Diseases*, 29(4):737–743, October 1999. doi: 10.1086/520426.

61. Joël Mossong, Niel Hens, Mark Jit, Philippe Beutels, Kari Auranen, Rafael Mikolajczyk, Marco Massari, Stefania Salmaso, Gianpaolo Scalia Tomba, Jacco Wallinga, Janneke Heijne, Malgorzata Sadkowska-Todys, Magdalena Rosinska, and W. John Edmunds. Social Contacts and Mixing Patterns Relevant to the Spread of Infectious Diseases. *PLOS Medicine*, 5(3):e74, March 2008. doi: 10.1371/journal.pmed.0050074.

62. M. Ngai. *Chinese Gold Miners and the "Chinese Question" in Nineteenth-Century California and Victoria*. W. W. Norton, New York, 2022.

63. Romualdo Pastor-Satorras, Claudio Castellano, Piet Van Mieghem, and Alessandro Vespignani. Epidemic processes in complex networks. *Reviews of Modern Physics*, 87(3):925–979, August 2015. doi: 10.1103/RevModPhys.87.925.

64. Beverley J. Paterson, Martyn D. Kirk, A. Scott Cameron, Catherine D'Este, and David N. Durrheim. Historical data and modern methods reveal insights in measles epidemiology: a retrospective closed cohort study. *BMJ Open*, 3(1):e002033, 2013. doi: 10.1136/bmjopen-2012-002033.

65. Velislava N. Petrova and Colin A. Russell. The evolution of seasonal influenza viruses. *Nature Reviews. Microbiology*, 16(1):60, January 2018. doi: 10.1038/nrmicro.2017.146.

66. Mario Pineda-Krch. GillespieSSA: Implementing the Gillespie Stochastic Simulation Algorithm in R. *Journal of Statistical Software*, 25:1–18, April 2008. ISSN 1548-7660. doi: 10.18637/jss.v025.i12.

67. Oliver G. Pybus, Andrew J. Tatem, and Philippe Lemey. Virus evolution and transmission in an ever more connected world. *Proceedings of the Royal Society B: Biological Sciences*, 282(1821), December 2015. doi: 10.1098/rspb.2014.2878.

68. Louis J. Rasmussen. *San Francisco Ship Passenger Lists*, volume 1. San Francisco historic record & genealogy bulletin, Colma, 1965.

69. Louis J. Rasmussen. *San Francisco Ship Passenger Lists*, volume 2. San Francisco historic record & genealogy bulletin, Colma, 1965.

70. Louis J. Rasmussen. *San Francisco Ship Passenger Lists*, volume 3. San Francisco historic record & genealogy bulletin, Colma, 1965.

71. Joanna J. Regan, J. Scott Vega, and Clive M. Brown. Infectious illnesses on cruise and cargo ships. In Eskild Petersen, Lin H. Chen, and Patricia Schlagenhauf-Lawlor, editors, *Infectious Diseases*, pages 35–44. Wiley, 1 edition, May 2017. doi: 10.1002/9781119085751.ch4.

72. Pejman Rohani, Xue Zhong, and Aaron A. King. Contact Network Structure Explains the Changing Epidemiology of Pertussis. *Science*, 330(6006):982–985, November 2010. doi: 10.1126/science.1194134.

73. Paul E Sampson. "The lungs of a ship": Ventilation, acclimatization, and labor in the maritime environment, 1740–1800. *History of Science*, 61(2), September 2021. doi: 10.1177/00732753211046449.

74. SlaveVoyages. Transatlantic Slave Trade Database: Voyage 92593, Diana, 1778, 2023. URL https://www.slavevoyages.org/voyage/92593/variables.

75. Matthew Smallman-Raynor and Andrew Cliff. *War Epidemics: An Historical Geography of Infectious Diseases in Military Conflict and Civil Strife, 1850-2000*. OUP Oxford, June 2004.

76. Elise Juzda Smith. 'Cleanse or Die': British Naval Hygiene in the Age of Steam, 1840–1900. *Medical History*, 62(2):177–198, April 2018.

77. Jessica E. Stockdale, Theodore Kypraios, and Philip D. O'Neill. Modelling and Bayesian analysis of the Abakaliki smallpox data. *Epidemics*, 19:13–23, June 2017. doi: 10.1016/j.epidem.2016.11.005.

78. Zachary Susswein, Eva C. Rest, and Shweta Bansal. Disentangling the rhythms of human activity in the built environment for airborne transmission risk: An analysis of large-scale mobility data. *eLife*, 12:e80466, April 2023.

79. R K Valle. Prevention of smallpox in Alta California during the Franciscan Mission Period (1769-1833). *California Medicine*, 119(1):73–77, July 1973.

80. Rosemary K. Valle. James Ohio Pattie and the 1827-1828 Alta California Measles Epidemic. *California Historical Quarterly*, 52(1):28–36, 1973. doi: 10.2307/25157415.

81. Thijs Veenstra, Patrick D. van Schelven, Yvonne M. ten Have, Corien M. Swaan, and Willem M. R. van den Akker. Extensive Spread of SARS-CoV-2 Delta Variant among Vaccinated Persons during 7-Day River Cruise, the Netherlands. *Emerging Infectious Diseases*, 29(4):734–741, April 2023. doi: 10.3201/eid2904.221433.

82. Delphis M. Vera, Ricardo A. Hora, Anarina Murillo, Juan F. Wong, Armando J. Torre, David Wang, Darbi Boulay, Kathy Hancock, Jacqueline M. Katz, Mariana Ramos, Luis Loayza, Jose Quispe, Erik J. Reaves, Daniel G. Bausch, Gerardo Chowell, and Joel M. Montgomery. Assessing the impact of public health interventions on the transmission of pandemic H1N1 influenza a virus aboard a Peruvian navy ship. *Influenza and Other Respiratory Viruses*, 8(3):353–359, 2014. doi: 10.1111/irv.12240.

83. Cécile Viboud, Ottar N. Bjørnstad, David L. Smith, Lone Simonsen, Mark A. Miller, and Bryan T. Grenfell. Synchrony, Waves, and Spatial Hierarchies in the Spread of Influenza. *Science*, 312(5772):447–451, April 2006. doi: 10.1126/science.1125237.

84. Emilia Vynnycky, Amy Trindall, and Punam Mangtani. Estimates of the reproduction numbers of Spanish influenza using morbidity data. *International Journal of Epidemiology*, 36(4):881–889, August 2007. doi: 10.1093/ije/dym071.

85. Helen J. Wearing, Pejman Rohani, and Matt J. Keeling. Appropriate Models for the Management of Infectious Diseases. *PLOS Medicine*, 2(7):e174, July 2005. doi: 10.1371/journal.pmed.0020174.

86. T.S. Wells. *The scale of medicines with which merchant vessels are to be furnished*. John Churchill, 1861.

87. Amy Wesolowski, Nathan Eagle, Andrew J. Tatem, David L. Smith, Abdisalan M. Noor, Robert W. Snow, and Caroline O. Buckee. Quantifying the Impact of Human Mobility on Malaria. *Science*, 338(6104):267–270, October 2012. doi: 10.1126/science.1223467.

88. Amy Wesolowski, Elisabeth zu Erbach-Schoenberg, Andrew J. Tatem, Christopher Lourenço, Cecile Viboud, Vivek Charu, Nathan Eagle, Kenth Engø-Monsen, Taimur Qureshi, Caroline O. Buckee, and C. J. E. Metcalf. Multinational patterns of seasonal asymmetry in human movement influence infectious disease dynamics. *Nature Communications*, 8(1):2069, December 2017. doi: 10.1038/s41467-017-02064-4.

89. Laura Forsberg White and Marcello Pagano. Transmissibility of the Influenza Virus in the 1918 Pandemic. *PLOS ONE*, 3(1):e1498, January 2008. doi: 10.1371/journal.pone.0001498.

90. P. Whittle. The outcome of a stochastic epidemic—a note on Bailey's paper. *Biometrika*, 42(1-2):116–122, 1955.

91. Kathryn S Willebrand, Lauren Pischel, Amyn A Malik, Samuel M Jenness, and Saad B Omer. A review of COVID-19 transmission dynamics and clinical outcomes on cruise ships worldwide, January to October 2020. *Eurosurveillance*, 27(1), January 2022. doi: 10.2807/1560-7917.ES.2022.27.1.2002113.

92. M. E. Willing. San Francisco Correspondence. *Sacramento Transcript*, page 2, 10 August, 1850.

93. Yingcun Xia, N. Bjørnstad, Ottar, and T. Grenfell, Bryan. Measles Metapopulation Dynamics: A Gravity Model for Epidemiological Coupling and Dynamics. *The American Naturalist*, 164(2):267–281, August 2004. doi: 10.1086/422341.

94. Claude Kwe Yinda, Dylan H. Morris, Robert J. Fischer, Shane Gallogly, Zachary A. Weishampel, Julia R. Port, Trenton Bushmaker, Jonathan E. Schulz, Kyle Bibby, Neeltje van Doremalen, James O. Lloyd-Smith, and Vincent J. Munster. Stability of Monkeypox Virus in Body Fluids and Wastewater. 29(10), 2023. doi: 10.3201/eid2910.230824.

Supplementary Material

Text S1. Transmission on board Historical Ships

Historical Evidence

Intensity. We expect that transmission on board historical ships was substantially more intense than transmission in land settings. In a study of pandemic 1918 influenza, Vynnycky et al. [84] estimated R_0 values of approximately 4-11 on board the troop ship His Majesty's Australian Transit (HMAT) *Boonah*, approximately 5-17 on board the troop ship HMAT *Devon*, and approximately 3.5-8 on board the troop ship HMAT *Medic*. These compare to an estimated range of roughly 1.5-4 in American and Scandinavian towns and cities. Similarly, White and Pagano [89] calculated an R_0 of roughly 4.98 for the same epidemics on board HMAT *Boonah* and HMAT *Medic*, compared with an R_0 of 1.34-3.21 in Maryland communities.

Similar analyses are not available for earlier time periods. However qualitative descriptions likewise indicate that conditions on board pre-1918 ships were highly conducive to intense pathogen transmission. One 1801 newspaper report describes an emigrant ship from Ireland to New York City so crowded that 'the space between decks, occupied by nearly 300 persons, became the receptacle of all excremental matters'. [2]. Half a century later, English author Frank Marryat recalled travelling from Panama to San Francisco on a boat "so crowded with passengers, that it was not until it was ascertained that there was scarcely standing-room for those on board that she tripped her anchor" [56]. Longer excerpts from both texts, printed below, offer vivid depictions of the conditions that passengers suffered during these voyages.

In addition to crowding, pre-1918 ships also faced substantial ventilation challenges. On sail ships, opening portholes risked losing heat and letting in water. Thus, below-deck spaces were sealed, resulting in notoriously poor air quality [73,76]. The steam revolution somewhat improved onboard environments, both by providing a source of heat and, in some cases, powering active ventilation systems. Yet steam ventilation still failed – sometimes catastrophically. In 1848, seventy-three people died from suffocation on board the *Londonderry* emigrant steamer when hatches were closed during a storm [73]. Six years later, fifty passengers out of a total of 204 died by suffocation in an unnamed ship carrying emigrants from Mauritius [86]. "Under no circumstance can a ship of any kind be made as healthful or as comfortable as the house on shore," wrote United States Navy surgeon Albert L. Gihon in 1886. Gihon added: "It is, practically, a floating box sealed against the admission of water, and, of course, also of air." [36]

Density Dependence. The nature of density dependence of pathogen transmission on board historical ships is unclear. On land, early-twentieth century studies demonstrate a clear and intuitive link between population density and rates of respiratory infection. Notably, Brewer [8] documented higher rates of pandemic influenza in military units training in more crowded environments in a 1918 study of Fort Humphreys, Virginia.

Some contemporaries clearly thought of ship transmission in what we would today consider density-dependent terms. In a report on 1918 pandemic influenza on board Royal Navy ships, surgeon-commander Sheldon F. Dudley argued that 'infective material must become so dense and diffused as to saturate the ship. That is to say, everyone on board receives a dose of the specific agent sufficient to cause influenza' [28]. Dudley's report is excerpted below.

Yet density dependence does not appear to describe every instance of shipboard transmission. In an analysis of smallpox outbreaks on board vessels bound into Australia, 1850-1908, quarantine director J. H. Cumpston found that most shipboard smallpox transmission was limited to close contacts of infected patients, such as family members, close colleagues, or those sharing beds and cabins [22]. Such a pattern typically argues for 'frequency-dependent' transmission, in which an infected individual on average transmits only to a fixed number of close contacts, regardless of a ship's total population size.

Contemporary Examples. Even with the benefits of modern hygiene, sophisticated air circulation, and more highly-regulated living conditions, infectious disease outbreaks are common on present-day cruise [59,60,91], cargo [71] and naval ships [19]. Attack rates are often high. Vera et al. [82] reported a 49.1% attack rate of 2009 H1N1 pandemic influenza in an unnamed Peruvian Navy ship carrying 355 people, with greatest risk in cadets assigned high-density living quarters. Earhart et al. [29] observed a 42% attack rate of H3N2 influenza across more than 500 people on board Navy vessel USS *Arkansas*, despite 95% vaccination coverage. Brotherton et al. (2000) reported a 37% attack rate of influenza-like illness on a cruise ship carrying over 1600 passengers and crew.

Recent reports of SARS-CoV-2 transmission on board passenger ships [3,43,54,81] and military ships [47] indicate high attack rates across both vaccinated and unvaccinated populations. Multiple studies suggest that outbreaks could spread broadly across ship populations. Using a mechanistic model, Azimi et al. [3] estimate similar transmission contributions from long-range aerosols and from short-range aerosols and droplets on board the *Diamond Princess* cruise ship. Meanwhile, in a study of an extensive outbreak on board a Dutch river cruise (60 of 132 passengers infected), Veenstra et al. [81] observed no apparent clustering by cabin layout or by meal-time seating arrangements.

Text S2. Qualitative Descriptions of Shipboard Infection and Transmission, 1801-1921

Three written descriptions of disease transmission on board nineteenth- and early twentieth-century ships are excerpted below. We do not intend these extracts to give a comprehensive view of shipboard transmission, but rather to offer a qualitative view of possible transmission scenarios. All spelling and grammar is original.

"Another instance of pestilence engendered in a ship crowded with passengers from Ireland." [2]

The ship Nancy, Capt. John Herron, was charterd by a commercial house at Sligo, to carry passengers from that port to New-York. She sailed from Sligo on the 12th July, 1801, and arrived, after a passage of 77 days, at the port of New York, on the 27th of September following. This ship, of the burthen of 202 tons, received on board 417 passengers, and was navigated by nine seamen. The provisions, mere refuse, put up by government-contractors with the view of saving expense, were of the worst kind: and the water, which was also of bad quality, from the unexpected length of the voyage, became extremely scanty before the arrival of the ship.

In order to receive so great a number of passengers on board of this ship, temporary cabbins [sic] were built on the quarter-deck, which were filled with eighty persons. Three hundred were crowded into the space between decks.

It will excite no surprise that a vessel thus crowded became sickly soon after sailing from Sligo. Typhous fever and dysentery began to prevail, and destroyed the lives of a large proportion of the passengers.

In addition to the wretchedness of being confined in so small a space, these unhappy emigrants suffered all the evils which their habits of uncleanliness could produce. Their bodies and clothes, covered and saturated with filth, exhuded poison all around them. Partly from the want of strength and assistance among the sick, and partly from the want of a sense of decency, the space between decks, occupied by nearly 300 persons, became the receptacle of all excremental matters, insomuch that they issued in streams from the scuppers. The filth on the upper deck was nearly over the shoes. The sides of the ship were daubed and incrusted [sic] with excrements; and even the rope for the support of such that wished to go on board were unfit to be handled. The stench was intolerably offensive.

In such condition arrived this unfortunate vessel at the place assigned for quarantine in the port of New York. Ninety persons had died on the passage; one hundred and eighty were sick. Scarcely a healthy countenance was to be seen on board of the ship; very few had escaped disease; and many had suffered from three to four relapses. About forty were taken ill after their arrival.

As soon as possible after their arrival the sick were brought ashore; stropped of their filthy and pestilential clothes; their bodies thoroughly washed and scoured with soap and water; and then wrapped up in clean blankets, and carried into the wards appointed for their reception in the Marine Hospital. The permanent buildings of the establishment were insufficient to receive so great a number; tents,

and other temporary accommodations, were provided for the remainder. Separation, ventilation, and cleanliness, as soon as they could be brought into action, accomplished every thing that could be expected. And only twenty-six have died since their arrival at this port.

Frank Marryat's description of a voyage from Panama to San Francisco, 1851 [56]

It seemed that we had brought the yellow fever with us to Panama, or rather it appeared at the time of our arrival, and it was now spreading with great rapidity. Cholera also broke out, and deaths from one or the other of these causes became very numerous.

The people being panic-struck, a great rush was made for the Californian boats, of which there happened, at this time, to be very few.

So soon as I was able to move, there was but one small screw steamer in port, and as the place was daily becoming more unhealthy, I secured, by great favour, a cabin in her.

Nothing could excuse the state in which this ship was put to sea, not even the panic; for she was not only ill-found in every respect, but was so crowded with passengers, that it was not until it was ascertained that there was scarcely standing-room for those on board that she tripped her anchor.

I had secured a dog-hole of a cabin, and was no sooner on board than my wife, worn out by fatigue and anxiety, was attacked by violent fever. There were two young doctors on board, but both were attacked shortly after we started. Then the epidemic (an aggravated intermittent fever) broke out among the passengers, who – crowded in the hold as thick as blacks in a slaver – gave way to fear, and could not be moved from the lower deck, and so lay weltering in their filth.

Sheldon F. Dudley's Report of 1918 pandemic influenza on board Royal Navy vessels, 1921 [28]

The density of susceptible persons in a ship must be very great as compared with an assemblage of susceptible persons on shore. If, for example, we contrast the sleeping accommodation in a ship with that of a big institution, we find that in the ship hammock hooks are less than 2 ft. apart, whereas institution bed centres are rarely less than 8ft.apart. Even when head to toe slinging is insisted on in a ship the men's heads must often be within 3ft. of each other. Now the volume of spray from a mouth at 3ft. is nearly twenty times that at 8ft. Therefore how much more readily will the man sleeping in a battleship's mess-deck get a requisite dose of infectious material than the man sleeping in an institution ashore? Again we also hear of many men in a ship not ill enough to go off duty, and, thus being immobilized, wandering about among their fellows.

When we consider these points, and at the same time realize that a modern battleship, with its tiers of lumbered decks, its cramped accommodation, and its crew of often over 1,000 men, covers an area of less than one-fiftieth of a square mile, I do not think it possible to doubt the infective material must become so dense and diffused as to saturate the ship. That is to say, everyone on board receives a dose of the specific agent sufficient to cause influenza, unless he happens to be highly immune at that time to the strain of organism responsible for the epidemic. In a ship, the density of susceptible

persons, the mass of infection, and the local migration are all so great that any diminution in one or more of these factors that may be produced by the use of sprays and gargles, by early isolation of cases and disinfection, is scarcely likely to diminish the rate of spread in a ship, once influenza has obtained a footing on board. And I think naval experience, where all these preventive measures have been vigorously employed, justifies this pessimism, as I have been unable to learn of any definite cases in which they did any good. In ships the outbreaks lasted ten days to three weeks; ashore the wave took about three months to pass over a locality. The longer wave period ashore was probably due to the lesser density of susceptible persons and infective sources, more time being required for the infection to hunt out all the susceptible individuals within its reach.

Text S3. Model Description and Equations

To achieve a more realistic depiction of the time course of infection, we make dwell times in state E and state I Erlang-distributed using the Linear Chain Trick [44]. This technique gives a unimodal distribution with a long right-hand tail, such that disease progression is relatively constrained in most individuals, but with occasional individuals experiencing substantially longer periods of incubation or infectiousness [85].

Individuals progress through k_E exposed states, $E_1, E_2, ..., E_{k_E}$, and through k_I infectious states, $I_1, I_2, ..., I_{k_I}$. We use $k_E = k_I = 3$ for all simulations.

The rate of progression from state E_e to state E_{e+1} and from state E_{k_E} to state I_1 is $\frac{k_E}{\mu_E}$, where μ_E represents the mean length of time that an individual spends in all exposed states. Similarly, the rate of progression from state I_i to state I_{i+1} and from state I_{k_I} to state R is is $\frac{k_I}{\mu_I}$, where μ_I represents the mean length of time that an individual spends in all infectious states.

We track infection across $g > 1$ transmission generations, where $E_{n,e}$ and $I_{n,i}$ denote n^{th}-generation individuals in states E_e and I_i respectively. The $\sum_{i=1}^{k_I} I_{n,i}$ infectious individuals from generation n produce new exposed individuals $E_{n+1,1}$, which represent the $(n + 1)^{\text{st}}$ generation of infections.

The number of first-generation individuals, $n = 1$, is fixed at $t = 0$. For all simulations, we assume a single first-generation individual. We randomly assign this person to a state from $E_{1,1}, E_{1,2}, ...E_{1,k_E}$ at the time of departure, $t = 0$.

To account for uncertainty and variation in the density dependence of shipboard contact rates, our model does not assume either classical density dependence or classical frequency dependence. Instead, we model the shipboard transmission with the equations:

$$\frac{dS}{dt} = -\beta_{fd}^{1-q}(\beta_{dd}N)^q S \sum_{n=1}^{g} \sum_{i=1}^{k_I} \frac{I_{n,i}}{N} \qquad\qquad 0 \leq q \leq 1$$

Here, q represents the degree of density dependence on board the ship, with $q = 0$ representing classical frequency dependence, $q = 1$ representing classical density dependence, and $0 < q < 1$ representing intermediate modes of transmission. The parameters β_{dd} and β_{fd} modulate the intensity of transmission under each density dependence pole — intuitively, the proportion (β_{dd}) and the raw number (β_{fd}) of people on board ship that a single infected individual will infect per day, on average, in a fully-susceptible population. Since susceptible people can be infected by infectious people in any generation and at any stage of infection, $\frac{dS}{dt}$ is proportional to the total number of infected people across all g transmission generations and all k_I infectious states.

The following equations show the deterministic analogue of our model. We implement continuous stochastic simulations in R using the Gillespie Algorithm, using the package GillespieSSA [66].

Model Equations

$$\frac{dS}{dt} = -\beta_{fd}^{1-q}(\beta_{dd}N)^q S \sum_{m=1}^{g} \sum_{j=1}^{k_I} \frac{I_{m,j}}{N}$$

$$\frac{dE_{1,1}}{dt} = -\frac{k_E}{\mu_E} E_{1,1}$$

$$\frac{dE_{n,1}}{dt} = \beta_{fd}^{1-q}(\beta_{dd}N)^q S \sum_{j=1}^{k_I} \frac{I_{n-1,j}}{N} - \frac{k_E}{\mu_E} E_{n,1} \qquad 2 \leq n \leq g$$

$$\frac{dE_{n,e}}{dt} = \frac{k_E}{\mu_E}(E_{n,e-1} - E_{n,e}) \qquad 1 < n \leq g; \; 2 \leq e \leq k_E$$

$$\frac{dI_{n,1}}{dt} = \frac{k_E}{\mu_E} E_{n,k_E} - \frac{k_I}{\mu_I} I_{n,1} \qquad 1 \leq n \leq g$$

$$\frac{dI_{n,i}}{dt} = \frac{k_I}{\mu_I}(I_{n,i-1} - I_{n,i}) \qquad 1 \leq n \leq g; \; 1 < i \leq k_I$$

$$\frac{dR}{dt} = \frac{k_I}{\mu_I} \sum_{m=1}^{g} I_{m,k_I}$$

Supplementary Tables

Table S1. Natural History Parameters

Pathogen	Mean latent period, μ_E	Mean infectious period, μ_I	Typical Land R_0	$\beta_{fd}^{\dagger}$	Reference
Influenza	2 days	3 days	1.5	0.5	[7,51,53]
Measles	12 days	8 days	15	1.875	[40,53]
Smallpox	12 days	17.5 days	7	0.4	[17,30,77]

$\dagger$ Calculated as $\beta_{fd} = \frac{R_0}{\mu_I}$

Table S2. San Francisco Port Arrivals Statistics, June 1850 – June 1852

Origin	Type	n	Journey Time (days)			Number of Passengers		
			Median	Range	Interdecile Range	Median	Range	Interdecile Range
Liverpool, UK	Sail	18	189.5	150-300	162-245.5	4.5	1-18	2-18
New York, USA*	Sail	92	155.5	89-283	113.1-219.8	5	0-160	1.1-20.0
	Steam	9	180	65-364	87.2-296	111	5-743	11.4-515.0
Valparaíso, Chile	Sail	100	57	23-190	45.0-74.2	7	0-175	1.0-39.0
Panama**	Sail	39	63	36-122	43.4-83.4	53	1-287	4.6-164.6
	Steam	116	20	12-58	15.7-28.0	196	11-1050	50.6-527.4
Oregon	Sail	48	7	3-16	3.4-11.3	4	0-12	1.0-6.0
	Steam	47	3	2-8	2.5-5.0	28	3-157	12.0-60.0
Hawai'i	Sail	76	22.5	14-80	16.0-31.5	5	0-142	2.0-22.0
	Steam	1	13	–	–	18	–	–
Sydney, Australia	Sail	20	91	67-188	70.0-111.2	15.5	2-142	3.9-57.0
Hong Kong	Sail	14	58.5	33-95	46.7-84.0	201	1-553	14.5-378.4

* For reasons that are unclear, four steam voyages originating in New York City had transit times longer than 200 days: the *SS New Orleans* (210 days), the *SS Sea Bird* (240 days), the *SS Goliah* (279 days) and the *SS Chesapeake* (364 days). The *SS Sea Bird* ran aground on San Martine on its way to California and had to be repaired [69]. Meanwhile, the extraordinarily long voyage of the *SS Chesapeake* astonished contemporary observers, with companies who had shipped goods on the vessel suing for damages on its arrival [92].

** Voyages from Panama predate the Panama Canal, which opened in 1914. Prior to its construction, travellers from Europe and the east coast took a ship to the eastern coast of Panama, crossed the Panama Isthmus by land, and took a second ship from the west coast of Panama to San Francisco [57].

Table S3. Selected Voyages, 1492-1918

Year	Vessel	Type	Purpose	Journey	Duration	Population	Ref.
1492	*Santa María*	Sail	Exploration	San Sebastian de la Gomera, Canary Islands to San Salvador, present-day Bahamas	35 days	41-60 crew (estimates vary; we use the more recent estimate of 41)	[35]
1497-98	*Matthew*	Sail	Exploration	Bristol, England to a disputed location in northeastern North America, possibly present-day Maine, Newfoundland, or Nova Scotia	35 days	20 crew	[46]
1609	*Sea Venture*	Sail	Emigration	Bristol, England to present-day Bermuda	54 days	150 passengers and crew	[25]
1620	*Mayflower*	Sail	Emigration	Leiden, Holland to near Cape Cod, Massachusetts	65 days	102 passengers, 20-30 crew (we assume 25 crew)	[34]
1784	*Empress of China*	Sail	Trade	New York City to Macao	184 days	34 crew	[30]
1778	*Diana*	Sail	Slave ship	Iles de Los, present-day Guinea to Curaçao	41 days	413 crew and enslaved people	[74]
1832	*America*	Sail	Convict transport	Unknown (likely United Kingdom to Australia)	180 days	175 crew and convicts	[64]
1918	HMAT *Boonah*	Steam	Troop ship	Durban, South Africa to Fremantle, Australia	17 days	164 crew, 931 troops	[23]