1

Control of ciliary transcriptional programs during spermatogenesis by antagonistic transcription factors

Weihua Wang[1,2,5]✉, Junqiao Xing[1,2,5], Xiqi Zhang[1,5], Hongni Liu[1,2], Xingyu Liu[3], Haochen Jiang[1,2], Cheng Xu[1,2], Xue Zhao[1,2], Zhangfeng Hu[1,2,4]✉

[1]Institute of Microalgae Synthetic Biology and Green Manufacturing, School of Life Sciences, Jianghan University, Wuhan, 430056 Hubei, China

[2]Hubei Engineering Research Center for Protection and Utilization of Special Biological Resources in the Hanjiang River Basin, School of Life Sciences, Jianghan University, Wuhan, 430056 Hubei, China

[3]NO.1 Middle School Affiliated to Central China Normal University, Wuhan, 430056 Hubei, China

[4]Hubei Key Laboratory of Environmental and Health Effects of Persistent Toxic Substances, Jianghan University, Wuhan, 430056 Hubei, China

[5]These authors contributed equally: Weihua Wang, Junqiao Xing, Xiqi Zhang

✉ e-mail: wangwh2019@163.com; huzhangfenghzf@163.com

Abstract

Existence of cilia in the last eukaryotic common ancestor (LECA) raises a fundamental question in biology: how the transcriptional regulation of ciliogenesis has evolved? One conceptual answer to this question is by an ancient transcription factor regulating ciliary gene expression in both unicellular and multicellular organisms, but examples of such transcription factors in eukaryotes are lacking. Previously, we showed that an ancient transcription factor XAP5 is required for flagellar assembly in *Chlamydomonas*. Here, we show that XAP5 and XAP5L are two conserved pairs of antagonistic transcription regulators that control ciliary transcriptional programs during spermatogenesis. Male mice lacking either XAP5 or XAP5L display infertility, as a result of meiotic prophase arrest and sperm flagella malformation, respectively. Mechanistically, XAP5 positively regulates the ciliary gene expression by activating the key regulators including FOXJ1 and RFX families during the early stage of spermatogenesis. In contrast, XAP5L negatively regulates the expression of ciliary genes via repressing these ciliary transcription factors during the spermiogenesis stage. Our results provide new insights into the mechanisms by which temporal and spatial transcription regulators are coordinated to control ciliary transcriptional programs during spermatogenesis.

Introduction

Cilia and flagella are evolutionarily conserved hair-like microtubule organelles that project from the surfaces of most eukaryotic cells(Derderian, Canales, & Reiter, 2023; Mitchell, 2017). They play essential roles in sensory reception, signal transduction and cell movement(Goetz & Anderson, 2010; Nachury & Mick, 2019; Singla & Reiter, 2006). The dysfunction of cilia in humans leads to an emerging class of genetic diseases called ciliopathies classified into first-order and second-order ciliopathies(Ishikawa & Marshall, 2011; Lovera & Lüders, 2021; Reiter & Leroux, 2017). First-order ciliopathies are caused by aberrations in ciliary proteins, and second-order ciliopathies occur due to defects in non-ciliary proteins, such as transcription factors, that are required for cilia formation and function(Reiter & Leroux, 2017).

Cilia are complex organelles and are dynamically regulated during development and the cell cycle(Eggenschwiler & Anderson, 2007; Kasahara & Inagaki, 2021; Santos & Reiter, 2008). The transcriptional regulation of ciliary genes must be precisely coordinated during ciliogenesis(Choksi, Lauter, Swoboda, & Roy, 2014; Collins, Ventrella, & Mitchell, 2021; Lewis & Stracker, 2021; Thomas et al., 2010). In metazoans, several transcription factors, including FOXJ1 and RFXs, have been shown to be involved in directing the expression of ciliary genes(Stubbs, Oishi, Izpisua Belmonte, & Kintner, 2008; Swoboda, Adler, & Thomas, 2000; X. Yu, Ng, Habacher, & Roy, 2008). These key ciliary transcription factors show dynamic expression patterns during development and differentiation(Stubbs et al., 2008; Swoboda et al., 2000; X. Yu et al.,

2008). However, how these key transcriptional programs are spatially and temporally controlled by cell type-specific transcription factors and signaling pathways in order to activate specific target ciliary genes and generate diverse cilia is largely unknown.

Despite transcriptional regulation of ciliary genes is required for ciliogenesis in both unicellular and multicellular organisms, the underlying mechanism mediating ciliary gene expression may be fundamentally different, as the two major transcription factors, FOXJ1 and RFXs, are absent from many unicellular organisms(Chu, Baillie, & Chen, 2010; Li et al., 2018; Piasecki, Burghoorn, & Swoboda, 2010; Vij et al., 2012). Cilia and ciliary genes have been highly conserved throughout evolution, suggesting that the regulation of ciliary genes could be programmed by yet undiscovered transcriptional mechanisms, which possibly coevolved with multicellularity. Recently, an ancient transcription factor, XAP5, is found to regulate ciliary gene expression in a unicellular organism, *Chlamydomonas reinhardtii*(Li et al., 2018). XAP5 is highly conserved among different species and is widely expressed across tissues(Martin-Tryon & Harmer, 2008; Mazzarella, Pengue, Yoon, Jones, & Schlessinger, 1997). In vertebrates, *XAP5-like* (*XAP5L*) with an intronless open reading frame originated in Therians via retrotransposition of the ancestral *XAP5*, and it is highly expressed during spermatogenesis(Sedlacek et al., 1999; A. Zhang et al., 2011). However, whether XAP5 and XAP5L play essential roles in cilia development in multicellular organisms remains unknown.

Here, we show that XAP5/XAP5L are antagonistic transcription factors required for

ciliary transcriptional programs during spermatogenesis. XAP5 is widely expressed in

different tissues, while XAP5L is exclusively present in the testes. To explore the roles of

XAP5/XAP5L in ciliogenesis, we generated a *XAP5L* knockout (KO) mouse line and

found that *XAP5L* KO male mice are infertile due to the malformation of sperm flagella.

Interestingly, male mice with the loss of XAP5 protein in germ cells also exhibited

infertile, due to the arrest of spermatogenesis in the meiotic prophase stage. These data

suggest that XAP5/XAP5L play crucial roles in sperm flagellar assembly in a

spatiotemporal manner. Mechanistically, XAP5 positively modulates transcriptional

rewiring of ciliary genes via activating the key regulators including FOXJ1 and RFXs

during the early stage of spermatogenesis. Conversely, XAP5L negatively regulates the

transcriptional program controlling ciliogenesis by repressing these ciliary transcription

factors during the spermiogenesis stage. Thus, ciliary transcriptional programs are

spatially and temporally controlled by XAP5/XAP5L antagonistic transcription factors

during spermatogenesis.

Results

XAP5 and XAP5L are indispensable for male fertility

To explore the potential functions of XAP5/XAP5L during ciliogenesis *in vivo*, we

performed Western blot analysis to investigate the spatiotemporal expression of

XAP5/XAP5L in various mouse tissues. We observed that XAP5 protein was widely

expressed in diverse tissues and present at all stages of postnatal testes maturation,

whereas XAP5L protein was mainly present in the testes and its level increased

dramatically from postnatal day 21 (P21) and continued into adulthood (Figures 1A, B).

Next, we identified the specific testicular cell types expressing XAP5/XAP5L using

published single-cell RNA-seq data(Jung et al., 2019). XAP5 mRNA was found

predominantly expressed in spermatogonia, while XAP5L mRNA was observed in

pachytene spermatocytes, and remained until elongating spermatids (Figure 1C).

Consistent with the mRNA localization patterns, immunofluorescence staining showed

that XAP5 protein was mainly detected in the nuclei of spermatogonia, whereas XAP5L

protein was present in the nuclei of pachytene spermatocytes and spermatids within the

seminiferous tubules, and neither of XAP5 or XAP5L was detected in mature sperm

(Figure 1D and Figure S1).

To investigate the physiological function of the uncharacterized XAP5/XAP5L protein,

we attempted to generate global KO mice using the CRISPR-Cas9 system. We

successfully established a *XAP5L* KO mouse line (Figures S2A-C) and found that while

all *XAP5L* KO mice survived to adulthood and appeared indistinguishable from the

wildtype (WT) mice, *XAP5L* KO males were sterile (Figures 1E, F). Notably, generating

mosaic F0 founders with *XAP5* mutant allele proved challenging, and no F1 generation

inherited the mutant alleles from the few F0 founders in over a year of breeding

experiments, underscoring the critical roles of XAP5 in mouse development.

Subsequently, we generated floxed-*XAP5* mice and crossed them with *Stra8-GFPCre*

knockin mice(Lin et al., 2017) to produce *XAP5* germline-specific KO mice (*XAP5* cKO)

(Figures S2D-F). Similar in appearance to controls ($XAP5^{fl/Y}$), all $XAP5$ cKO males were

also found to be sterile (Figures 1G, H). Collectively, these results suggest that

XAP5/XAP5L play essential roles in spermatogenesis.

XAP5 and XAP5L function at different stages of spermatogenesis

To uncover the cause of male infertility, we conducted gross and histological analyses of

the testes from WT and $XAP5L$ KO mice. There were no differences in the gross

appearance of testis or in the weight of mouse body and testis between WT and $XAP5L$

KO mice (Figure 2A), but the number of sperm collected from cauda epididymis of

$XAP5L$ KO mice was significantly lower than that from WT mice (Figure 2B), and sperm

motility was significantly reduced in KO mice (Figure 2C and Videos S1, 2).

Histological analysis did not reveal any severe abnormity in the seminiferous epithelial

cycle of XAP5L KO testes (Figure 2D). Further examination of the morphology and

structure of cauda epididymal sperm in $XAP5L$ KO mice revealed significantly higher

rates of abnormal flagella (Figures 2E-G), indicating that XAP5L is crucial for flagellar

assembly during spermiogenesis.

We also investigated the spermatogenic defects in the seminiferous tubules and

epididymis of $XAP5$ cKO male mice. Notably, the testes of $XAP5$ cKO males were

significantly smaller compared to WT controls at P16 and older (Figure 3A). Many

seminiferous tubules in P16 and older $XAP5$ cKO males were vacuolated due to the

absence of spermatocytes and spermatids, as indicated by the lack of XY body formation

and the absence of PNA signal, a marker of spermatids and sperm, in $XAP5$ cKO germ

cells (Figures 3B-D). As a consequence, no sperm were observed in the adult cauda

epididymides of *XAP5* cKO mice (Figure 3E). Taken together, these data indicate that

germ cell development in *XAP5* cKO males was arrested in meiotic prophase, and XAP5

is indispensable for meiotic progression during spermatogenesis.

XAP5 and XAP5L function as antagonistic regulators of ciliary transcriptional

regulatory networks via regulating transcription factors

To gain insight into the regulatory mechanisms of XAP5L during sperm flagellar

assembly, we employed RNA-seq analysis to examine mature sperm collected from the

cauda epididymis of WT and *XAP5L* KO adult mice. Differential expression analysis

identified 2093 upregulated and 267 downregulated genes in *XAP5L* KO sperm (Figure

S3A). Gene ontology analysis revealed that the upregulated genes were associated with

cilia assembly or function (Figure 4A and Figures S3B, C). When crossed with the

previously compiled ciliogenesis-related gene list(Nemajerova et al., 2016), the

upregulated genes were observed encompassing diverse structural and functional

components of the ciliogenic program (Table S1 and Figure S3D). Notably, several key

transcription factor (TF) regulators of ciliogenesis were among the upregulated genes,

including FOXJ1 and RFX families (Figures 4B, C and Figure S3E). Moreover, several

key regulators of spermatogenesis, particularly spermiogenesis, were identified, including

RFX2(Kistler et al., 2015; Wu et al., 2016), SOX families(Schartl et al., 2018; D. Zhang

et al., 2018), TAF7L(Zhou et al., 2013) and TBPL1(Martianov et al., 2001) (Figure 4B

and Figure S3E). Many core ciliary genes were also upregulated, such as

CFAP206(Beckers et al., 2020) and *IFT81*(Qu et al., 2020) which are critical for male

fertility (Figure S3E), consistent with the tight coregulation between spermiogenesis and

cilia-related genes during spermatogenesis. These findings suggest that the genes

identified in our study provide an excellent resource for candidates with novel ciliary or

spermatogenesis-related functions, and the newly identified testes-specific protein

TULP2(Oyama et al., 2022; Zheng et al., 2021) was selected for validation. We observed

that *TULP2* KO male mice were sterile due to malformation of sperm flagella (Figure

S4). Overall, our results clearly indicate that XAP5L acts as a central transcriptional

repressor of ciliogenesis during mouse spermatogenesis.

Furthermore, we performed RNA-seq analysis using P16 testes samples from WT and

XAP5 cKO mice to explore the underlying mechanism of XAP5-mediated germ cell loss.

Differential expression analysis identified 554 upregulated and 1587 downregulated

genes in *XAP5* cKO sperm (Figure S5A). We observed that the downregulated genes in

XAP5 cKO males were involved in cilium assembly or functions (Figure 4D and Figures

S5B-D). Strikingly, many of these downregulated genes were upregulated in *XAP5L* KO

sperm, including the key ciliogenesis regulators FOXJ1 and RFX families (Figures 4E, F,

Figure S5E and Table S1). In addition, meiosis initiation marker STRA8(Anderson et al.,

2008; Ferder et al., 2019), meiosis-specific genes SPO11(Romanienko & Camerini-

Otero, 2000) and DMC1(Bishop, Park, Xu, & Kleckner, 1992), key transcription

regulators of spermiogenesis RFX2(Kistler et al., 2015; Wu et al., 2016), SOX30(D.

Zhang et al., 2018) and CREM(Blendy, Kaestner, Weinbauer, Nieschlag, & Schutz, 1996;

Nantel et al., 1996) were also downregulated in *XAP5* cKO mice (Figures 4E and S5F),

consistent with the aforementioned result that loss of XAP5 induced spermatogenesis

arrest in meiotic stage.

To identify direct targets of XAP5/XAP5L, we conducted CUT&Tag analyses on P16

mouse testicular germ cells for XAP5, and on P56 mouse testicular germ cells for

XAP5L. Our results revealed 23143 peaks enriched for XAP5, compared to only 2290

peaks enriched for XAP5L (Table S1), and the limited number of enriched peaks for

XAP5L suggests that the antibody used for XAP5L in the CUT&Tag experiment might

have suboptimal performance. A substantial proportion of these peaks were located

within promoter regions (Figures S6A, B), indicating that XAP5/XAP5L primarily

function to regulate the expression of protein-coding genes. De novo motif analyses

showed that the binding motifs for XAP5 and XAP5L shared a conserved sequence

(CCCCGCCC/GGGCGGGG) (Figure S6C), suggesting that XAP5 may regulate a subset

of the same downstream genes as XAP5L.

To further elucidate the target genes of XAP5 and XAP5L, we integrated our

CUT&Tag data with RNA-seq datasets (Table S1). This analysis revealed that among the

genes downregulated in *XAP5* cKO germ cells, 891 were bound by XAP5 (Figure S6D).

Similarly, 75 genes upregulated in *XAP5L* KO sperm were occupied by XAP5L (Figure

S6E). GO analysis indicated that these genes are associated with cilium assembly and

related functions (Figures S6D-G). Notably, both *FOXJ1* and *RFX2* are regulated and

occupied by XAP5, with the conserved sequence present within the XAP5-bound regions

(Figure 4G). Furthermore, real-time PCR validation of the CUT&Tag experiments confirmed that *RFX2* is also occupied by XAP5L, despite the initial CUT&Tag data not showing enriched peaks for the RFX2 gene (Figure 4G). Unfortunately, due to the suboptimal performance of the XAP5L antibody, we did not detect the occupancy of XAP5L on the FOXJ1 promoter. Nevertheless, these findings suggest that XAP5 and XAP5L regulate ciliary gene expression primarily through direct binding to the conserved motif present in these genes.

Considering the evolutionary conservation of XAP5/XAP5L protein across different species and the expression patterns of XAP5, XAP5L, FOXJ1 and RFX factors during spermatogenesis (Figures 1C and S7), these results suggest that XAP5 and XAP5L have antagonistic effects, and they may function upstream of the transcription factor families, including FOXJ1 and RFX factors, to coordinate the ciliogenesis during spermatogenesis (Figure 4H).

Discussion

Cilia and flagella are ancient organelles present in the last eukaryotic common ancestor (LECA)(Mitchell, 2017). Cilia assembly and maintenance are under strict transcriptional regulation in both unicellular and multicellular organisms(Choksi et al., 2014; Collins et al., 2021; Lewis & Stracker, 2021; Thomas et al., 2010). Despite cilia and flagella having an ancient origin, the evolutionary history of ciliary gene regulation has remained an unsolved problem. The main reason is that the master ciliary transcription factors found

in multicellular organisms, including FOXJ1 and RFXs, are absent from unicellular

organisms(Choksi et al., 2014; Collins et al., 2021; Lewis & Stracker, 2021; Thomas et

al., 2010). Previously, we showed that an ancient transcription factor, XAP5, was

required for flagella assembly in the unicellular green algae *Chlamydomonas*(Li et al.,

2018). XAP5 proteins are evolutionarily conserved across diverse organisms(Li et al.,

2018; Martin-Tryon & Harmer, 2008), which offers the possibility to investigate the

conservation of the role of XAP5 in multicellular organisms. In the present study, we

report that XAP5 positively regulates transcriptional network of ciliary genes by

activating the key regulators including FOXJ1 and RFXs during spermatogenesis.

Cilia are dynamic organelles, and the expression profile of ciliary genes is dynamic

during the cell cycle and under certain physiological conditions(Eggenschwiler &

Anderson, 2007; Kasahara & Inagaki, 2021; Santos & Reiter, 2008). Cells have

developed regulatory mechanisms to generate functional cilia in a temporal and spatial

manner(Choksi et al., 2014; Collins et al., 2021; Lewis & Stracker, 2021; Thomas et al.,

2010). However, how ciliary transcription factors determine temporal and spatial ciliary

transcriptional programs is poorly understood. Sperm flagellar assembly is likewise

tightly regulated during a highly complex temporal process named

spermatogenesis(Holstein, Schulze, & Davidoff, 2003; Thomas et al., 2010). Several key

transcriptional regulators of spermatogenesis, such as FOXJ1 and RFX2, have been

identified(Chen, Knowles, Hebert, & Hackett, 1998; Kistler et al., 2015; Wu et al., 2016).

The flagellar formation of FOXJ1-null sperm is severely impaired, and the FOXJ1 target

genes are required for sperm motility(Beckers et al., 2020; Chen et al., 1998; Weidemann

et al., 2016). *RFX2* KO mice are completely sterile and RFX2 regulates the expression of

ciliary genes during spermiogenesis(Kistler et al., 2015; Wu et al., 2016). Ciliary

transcription factors (FOXJ1 and RFX2) are dynamically regulated during

spermatogenesis(Jung et al., 2019). Intriguingly, our results show that XAP5 and XAP5L

are two conserved pairs of antagonistic transcription factors that regulate these key

transcriptional regulators required for spermatogenesis.

By using Western blot analysis, we found that XAP5 was ubiquitously expressed in all

tissues examined, whereas XAP5L expression was restricted to the testes (Figure 1A).

Interestingly, previous research found that XAP5L was widely expressed in normal tissue

via using TCGA expression data(Thompson et al., 2021). Moreover, loss of

XAP5/XAP5L perturbs transcriptional programs in cancer cells. At present, as shown by

the recent data, we cannot rule out the possibility that XAP5L is dynamically expressed

in other tissues during development. Although XAP5 and XAP5L are highly expressed in

testes and critical for spermatogenesis, they are not present in mature sperm (Figure S1).

Thus, XAP5/XAP5L are not needed for sperm maintenance. Given the dynamic

expression of XAP5/XAP5L during spermatogenesis, it will be interesting to identify the

key signaling factors that regulate the timely and spatial expression of XAP5 and

XAP5L.

A role for XAP5 in human brain development is suggested by the association of X-

linked intellectual disability (XLID) with rare *XAP5* missense variants(Lee et al., 2020).

However, how defects in XAP5 lead to XLID syndrome is unknown. Although there is as

yet no evidence that XAP5 regulates ciliogenesis in any system besides *Chlamydomonas*

and mice, the function in cilia could help clarify the unexplained role of *XAP5* mutations

in causing XLID. The possibility that XAP5 could be a ciliary transcription factor in

humans would add XLID to the growing list of the second-order ciliopathies.

Materials and Methods

Mice

C57BL/6J strains (from Vital River Laboratories, Beijing, China) and *Stra8-GFPCre*

knockin mice (from Cyagen Biosciences) were used. Floxed-*XAP5* mice, *XAP5L* and

TULP2 knockout mice were generated on the C57BL/6J background using the

CRISPR/Cas9 system. The sgRNA sequences used to target *XAP5*, *XAP5L* and *TULP2*

were as follows: *XAP5L*, sgRNA1-GGC TAC CAG AAA CAG GGA CT, sgRNA2-GGG

AGT AAG GTC CCC AAA CT; *XAP5*, sgRNA3-CTA CAG GGC ACT TAT TAA TA,

sgRNA4-ATA GTA ATT CCC CCG TGC TT; *TULP2*, sgRNA5-TGA CTA ATT AGG

CCC GAG AG, sgRNA6-GGT TCT TAG AGA GTC AAC GT. Experimental protocols

were approved by the ethics committees of Jianghan University. Mice were maintained in

a pathogen-free environment with a room temperature of 23 ± 2 °C under a humidity

level of 30-70%. The light was maintained on a 12-hour day/night cycle.

Genome PCR

Mouse genomic DNA was isolated from the tail tip using One Step Mouse Genotyping Kit (Vazyme, PD101), and subjected to PCR with $2 \times$ Taq Plus Master Mix (Vazyme, P212) following the manufacturer's instructions. The PCR products were separated by 2% agarose gel electrophoresis. The primers used are listed in Table S2.

RNA isolation and real-time PCR

Total RNA was isolated using the TRIzol reagent (Invitrogen, 15596018) and then converted to cDNA with HiScript II 1st Strand cDNA Synthesis Kit (Vazyme, R212) following the manufacturer's instructions. Real-time PCR was conducted using ChamQ SYBR qPCR Master Mix (Vazyme, Q311) on an ABI StepOnePlus real-time PCR system (Applied Biosystems). Expression values were normalized using the ΔΔCt method with *Gapdh* as the internal control. The primer sequences are indicated in Table S2.

Western blot analysis

Cell lysates from mouse tissues were prepared in RIPA buffer (Beyotime, P0013B) containing Protease Inhibitor Cocktail (Roche, 11873580001) and 1mM PMSF. Subcellular lysates of testicular nuclear and cytoplasmic fractions were prepared using Nuclear and Cytoplasmic Protein Extraction Kit (Beyotime, P0027) following the manufacturer's instructions. The protein lysates were subjected to standard

immunoblotting analysis. Antibodies used were as follows: anti-β-actin (ABclonal,

AC026), anti-XAP5 (Prospertech, HU-412003), anti-XAP5 (Sigma, HPA003585), anti-

XAP5L (Prospertech, Hu-412001), anti-LMNB (Proteintech, 66095-1-Ig), anti-GAPDH

(ABclonal, AC002), anti-FOXJ1 (D360355, Sangon Biotech), anti-RFX2 (K110716P,

Solarbio) and anti-TULP2 (Prospertech, Hu-412002).

Immunofluorescence

Mouse testis were fixed in 4% paraformaldehyde in PBS at 4 °C overnight, dehydrated in

gradient ethanol, embedded in paraffin, and sectioned into 5 μm slices. After

deparaffinization and rehydration, testis sections were boiled in 1 mM EDTA, pH 8.0 for

15 min using a microwave oven for antigen retrieval, followed by cooling to room

temperature. The primary antibodies, including anti-XAP5 (Prospertech, HU-412003,

1:250 dilution), anti-XAP5 (Sigma, HPA003585, 1:250 dilution), anti-XAP5L

(Prospertech, Hu-412001, 1:250 dilution), anti-lectin PNA, Alexa Fluor 488 Conjugate

(ThermoFisher, L21409, 1:250 dilution), and anti-TULP2 (Prospertech, Hu-412002,

1:250 dilution), were then incubated at 4 °C overnight. Subsequently, the sections were

incubated with the secondary antibody donkey anti-rabbit Alexa Fluor Plus 488

(Invitrogen, A32790, 1:500 dilution) for 1 hour and DAPI for 5 minutes at room

temperature. Images were collected using a Zeiss Axio Vert A1 microscope.

Histological analysis

Testes and epididymides were fixed in Bouin's solution (sigma, HT10132) overnight at

4 °C, embedded in paraffin and sectioned into 5 μm slices. The sections were then

subjected to H&E staining using standard procedures and observed under a Zeiss Axio

Vert A1 microscope.

Assessment of sperm counts, motility and morphology

Cauda epididymides were collected from male mice and dissected in 37°C pre-warmed

human tubal fluid (HTF) culture medium (EasyCheck, M1130), followed by incubation

at 37°C for 15 min to release the sperm. To quantify sperm abundance, the sperm

suspension was fixed in 1% paraformaldehyde for 30 min. Sperm were then counted

under a Zeiss ×10 bright-field microscope. To assess sperm motility, the sperm

suspension was observed and recorded under a Leica total internal reflection fluorescence

microscope. Motility was defined as any movement of the sperm flagellum during a 10-

second observation cycle. After immobilization with 1% paraformaldehyde, sperm

morphology was observed by H&E staining using standard procedures. Images were

captured using a Zeiss Axio Vert A1 microscope equipped with a ×100 oil immersion

objective. Sperm without staining were also used to assess sperm morphology. For

ultrastructural observation, sperm samples were fixed in 3% glutaraldehyde in 0.1M

sodium phosphate buffer (pH7.4), and post-fixed with 1% (wt/vol) OsO_4. After

dehydration, the samples were placed in propylene oxide and embedded in a mixture of

Epon 812 and Araldite. Ultrathin sections obtained by a Leica EM UC7 ultramicrotome

were stained with uranyl acetate and lead citrate, then analyzed using a HT7700 TEM

(Hitachi).

Fertility test

Each of the adult male mice was mated with two females for 2 months. All the mice used

were aged 6-8 weeks. Vaginal plugs were checked every morning, and the number of

newborn pups was counted.

Chromosome spreads and immunofluorescence

Testicular chromosome spreads were prepared and immunolabeled as

described(Reinholdt, Ashley, Schimenti, & Shima, 2004). The primary antibodies used

were as follows: anti-γ H2AX (Millipore, 05-636, 1:300 dilution), anti-SYCP3

(Proteintech, 23024-1-AP, 1:300 dilution). Secondary antibodies used were Alexa Dye

(AlexaFluor Plus 488/594) conjugates (ThermoFisher) at 1:500 dilutions. Images were

collected with a Zeiss Axio Vert A1 microscope.

RNA-seq analysis

The mature sperm samples from P56 mice and testis samples from P16 mice were

washed three time in 1× PBS after harvest. Total RNA was extracted from the samples

using the TRIzol reagent (Invitrogen) and purified with the NEBNext Poly(A) mRNA

Magnetic Isolation Module (NEB, E7490). The quality of the RNA samples was

examined using the Agilent 2100 Bioanalyzer system (Agilent Technologies). Qualified

RNA samples were subjected to sequencing library construction using the NEBNext

Ultra II RNA Library Prep Kit for Illumina (NEB, E7775). Paired-end (2×150 bp)

sequencing was carried out on the Illumina NovaSeq6000 platform. The sequencing reads

were mapped to the Mus musculus reference genome (GRCm38/mm10) using HISAT2

v2.1.0(Kim, Langmead, & Salzberg, 2015), and StringTie v1.3.3b(Pertea et al., 2015)

was applied to assemble the mapped reads. Fragments Per Kilobase of transcript per

Million fragments mapped (FPKM) was used to measure the expression level of a gene.

Differential expression analysis was processed by DESeq2 v1.12.4(Subramanian et al.,

2005), and the significant differentially expressed genes were identified with a fold

change >2. Gene Ontology analysis was performed using the DAVID database.

CUT&Tag analysis

The testicular germ cells were isolated using a two-step enzymatic digestion process as

previously described (Liu et al., 2017). Briefly, after removing the tunica albuginea, the

testes were incubated in 1 mg/mL type IV collagenase (Sigma, C5138) at 37℃ for 15

min with gentle shaking every 5 min. The resulting cell suspension was centrifuged at

1000 rpm for 3 min, and the pellet was further digested with 0.25% Trypsin-EDTA

(Gibco, 25200072) at 37℃ for 10 min. The digestion was terminated by adding fetal

bovine serum (FBS), and the cell suspension was centrifuged at 1000 rpm for 3 min. The

cell pellet was resuspended in DMEM medium containing 10% FBS and cultured at 37℃

in a 5% CO2 atmosphere for 3 h. Floating and weakly adherent cells were collected as

germ cells. CUT&Tag and CUT&Tag real-time PCR were performed according to the

manufacturer's protocol using the Hyperactive Universal CUT&Tag Assay Kit for

Illumina Pro (Vazyme, TD904). Approximately 100,000 germ cells were used per sample.

For real-time PCR, IgG sera were used as a negative control for the XAP5 CUT&Tag,

while lysates from *XAP5L* KO testis were incubated with the XAP5L antibody as a

negative control for the XAP5L CUT&Tag. DNA spike in was used as the internal

control. The primer sequences are indicated in Table S2.

For analysis, quality control and read trimming were performed using FastQC (v0.11.9)

with default parameters and Trimmomatic (v0.39) with the following parameters:

ILLUMINACLIP:NexteraPE-PE.fa:2:30:10:8:true LEADING:3 TRAILING:3

SLIDINGWINDOW:4:15 MINLEN:8. Clean reads were aligned with bowtie2

(v2.4.4)(Langmead & Salzberg, 2012) against the Mus musculus (mm39) genome with

the parameters -X 1000 --very-sensitive. PCR duplicates were removed using picard's

MarkDuplicates with default parameters. Peak calling was conducted using using

MACS2 (v2.1.4) with the parameters -f BAMPE -B --SPMR --keep-dup all. Peaks were

annotated to the nearest feature using ChIPseeker (v1.32.0)(G. Yu, Wang, & He, 2015).

Motif finding and annotation were performed using Homer *findMotifsGenome.pl* and

annotatePeaks.pl functions, respectively(Heinz et al., 2010). Track screen shots were

produced in IGV (version 2.18.1)(Robinson, Thorvaldsdottir, Turner, & Mesirov, 2023).

Statistical analysis

Statistical analyses were conducted using GraphPad Prism 8.0.2 software. All data were presented as mean ± SEM. Values of $P < 0.05$ were considered statistically significant. Statistical significance between two groups was calculated using an unpaired, parametric, two-sided student's t test.

Materials and Data availability

The RNA-seq and CUT&Tag data generated in this study have been deposited in the Gene Expression Omnibus (GEO) database and are publicly available under accession numbers GSE236388 for RNA-seq and GSE279586 for CUT&Tag, respectively.

The authors declare that all the materials and the data supporting the findings of this study are available within the article and its supplementary materials or from the corresponding author upon reasonable request.

Acknowledgements

This work was supported by the National Key R&D Program of China (grant number: 2020YFA0907400), the National Natural Science Foundation of China (grant number: 32170702 and 82000828) and the Major Special Funding Program for First-class Discipline Construction of Jianghan University (grant number: 2023XKZ021). We thank Yuan He (Research Center for Medicine and Structural Biology, Wuhan University,

441 China) for the assistance during the experimental design of the transmission electron

442 microscope analysis.

Author Contributions

445 W.W., J.X., X.Z. and Z.H. designed research; W.W., J.X., X.Z., H.L., X.L., H.J., C.X. and

446 X.Z. performed research; W.W. and Z.H. contributed new reagents/analytic tools; W.W.,

447 J.X., X.Z. and Z.H. analyzed data; and W.W. and Z.H. wrote the paper.

Competing interests

450 The authors declare no competing interests.

Additional files

Table S1

454 The lists of differential expression genes, ciliogenesis-related genes, and enriched peaks

455 for XAP5/XAP5L.

Table S2

457 The lists of primers used in the Genome PCR and real-time PCR.

Video S1

459 Representative video of living sperm cells from WT mouse.

Video S2

461 Representative video of living sperm cells from *XAP5L* KO mouse.

References

Anderson, E. L., Baltus, A. E., Roepers-Gajadien, H. L., Hassold, T. J., de Rooij, D. G., van Pelt, A. M., & Page, D. C. (2008). Stra8 and its inducer, retinoic acid, regulate meiotic initiation in both spermatogenesis and oogenesis in mice. *Proc Natl Acad Sci U S A, 105*(39), 14976-14980. doi:10.1073/pnas.0807297105

Beckers, A., Adis, C., Schuster-Gossler, K., Tveriakhina, L., Ott, T., Fuhl, F., . . . Gossler, A. (2020). The FOXJ1 target Cfap206 is required for sperm motility, mucociliary clearance of the airways and brain development. *Development*. doi:10.1242/dev.188052

Bishop, D. K., Park, D., Xu, L., & Kleckner, N. (1992). DMC1: a meiosis-specific yeast homolog of E. coli recA required for recombination, synaptonemal complex formation, and cell cycle progression. *Cell, 69*(3), 439-456. doi:10.1016/0092-8674(92)90446-j

Blendy, J. A., Kaestner, K. H., Weinbauer, G. F., Nieschlag, E., & Schutz, G. (1996). Severe impairment of spermatogenesis in mice lacking the CREM gene. *Nature, 380*(6570), 162-165. doi:10.1038/380162a0

Chen, J., Knowles, H. J., Hebert, J. L., & Hackett, B. P. (1998). Mutation of the mouse hepatocyte nuclear factor/forkhead homologue 4 gene results in an absence of cilia and random left-right asymmetry. *J Clin Invest, 102*(6), 1077-1082. doi:10.1172/JCI4786

Choksi, S. P., Lauter, G., Swoboda, P., & Roy, S. (2014). Switching on cilia: transcriptional networks regulating ciliogenesis. *Development, 141*(7), 1427-1441. doi:10.1242/dev.074666

Chu, J. S., Baillie, D. L., & Chen, N. (2010). Convergent evolution of RFX transcription factors and ciliary genes predated the origin of metazoans. *BMC Evol Biol, 10*, 130. doi:10.1186/1471-2148-10-130

Collins, C., Ventrella, R., & Mitchell, B. J. (2021). Building a ciliated epithelium: Transcriptional regulation and radial intercalation of multiciliated cells. *Curr Top Dev Biol, 145*, 3-39. doi:10.1016/bs.ctdb.2020.08.001

Derderian, C., Canales, G. I., & Reiter, J. F. (2023). Seriously cilia: A tiny organelle illuminates evolution, disease, and intercellular communication. *Dev Cell, 58*(15), 1333-1349. doi:10.1016/j.devcel.2023.06.013

Eggenschwiler, J. T., & Anderson, K. V. (2007). Cilia and developmental signaling. *Annu Rev Cell Dev Biol, 23*, 345-373. doi:10.1146/annurev.cellbio.23.090506.123249

Ferder, I. C., Fung, L., Ohguchi, Y., Zhang, X., Lassen, K. G., Capen, D., . . . Wang, N. (2019). Meiotic gatekeeper STRA8 suppresses autophagy by repressing Nr1d1 expression during spermatogenesis in mice. *PLoS Genet, 15*(5), e1008084. doi:10.1371/journal.pgen.1008084

Goetz, S. C., & Anderson, K. V. (2010). The primary cilium: a signalling centre during vertebrate development. *Nat Rev Genet, 11*(5), 331-344. doi:10.1038/nrg2774

Heinz, S., Benner, C., Spann, N., Bertolino, E., Lin, Y. C., Laslo, P., . . . Glass, C. K. (2010). Simple combinations of lineage-determining transcription factors prime cis-regulatory elements required for macrophage and B cell identities. *Mol Cell, 38*(4), 576-589. doi:10.1016/j.molcel.2010.05.004

Holstein, A.-F., Schulze, W., & Davidoff, M. (2003). Understanding spermatogenesis is a prerequisite for

treatment. *Reproductive Biology and Endocrinology, 1*(1). doi:10.1186/1477-7827-1-107

Ishikawa, H., & Marshall, W. F. (2011). Ciliogenesis: building the cell's antenna. *Nat Rev Mol Cell Biol, 12*(4), 222-234. doi:10.1038/nrm3085

Jung, M., Wells, D., Rusch, J., Ahmad, S., Marchini, J., Myers, S. R., & Conrad, D. F. (2019). Unified single-cell analysis of testis gene regulation and pathology in five mouse strains. *Elife, 8*. doi:10.7554/eLife.43966

Kasahara, K., & Inagaki, M. (2021). Primary ciliary signaling: links with the cell cycle. *Trends Cell Biol, 31*(12), 954-964. doi:10.1016/j.tcb.2021.07.009

Kim, D., Langmead, B., & Salzberg, S. L. (2015). HISAT: a fast spliced aligner with low memory requirements. *Nat Methods, 12*(4), 357-360. doi:10.1038/nmeth.3317

Kistler, W. S., Baas, D., Lemeille, S., Paschaki, M., Seguin-Estevez, Q., Barras, E., . . . Reith, W. (2015). RFX2 Is a Major Transcriptional Regulator of Spermiogenesis. *PLoS Genet, 11*(7), e1005368. doi:10.1371/journal.pgen.1005368

Langmead, B., & Salzberg, S. L. (2012). Fast gapped-read alignment with Bowtie 2. *Nat Methods, 9*(4), 357-359. doi:10.1038/nmeth.1923

Lee, Y.-R., Khan, K., Armfield-Uhas, K., Srikanth, S., Thompson, N. A., Pardo, M., . . . Schwartz, C. E. (2020). Mutations in FAM50A suggest that Armfield XLID syndrome is a spliceosomopathy. *Nature Communications, 11*(1). doi:10.1038/s41467-020-17452-6

Lewis, M., & Stracker, T. H. (2021). Transcriptional regulation of multiciliated cell differentiation. *Semin Cell Dev Biol, 110*, 51-60. doi:10.1016/j.semcdb.2020.04.007

Li, L., Tian, G., Peng, H., Meng, D., Wang, L., Hu, X., . . . Hu, Z. (2018). New class of transcription factors controls flagellar assembly by recruiting RNA polymerase II in Chlamydomonas. *Proc Natl Acad Sci U S A, 115*(17), 4435-4440. doi:10.1073/pnas.1719206115

Lin, Z., Hsu, P. J., Xing, X., Fang, J., Lu, Z., Zou, Q., . . . Tong, M. H. (2017). Mettl3-/Mettl14-mediated mRNA N(6)-methyladenosine modulates murine spermatogenesis. *Cell Res, 27*(10), 1216-1230. doi:10.1038/cr.2017.117

Liu, W., Wang, F., Xu, Q., Shi, J., Zhang, X., Lu, X., . . . Li, L. (2017). BCAS2 is involved in alternative mRNA splicing in spermatogonia and the transition to meiosis. *Nat Commun, 8*, 14182. doi:10.1038/ncomms14182

Lovera, M., & Lüders, J. (2021). The ciliary impact of nonciliary gene mutations. *Trends Cell Biol, 31*(11), 876-887. doi:10.1016/j.tcb.2021.06.001

Martianov, I., Fimia, G. M., Dierich, A., Parvinen, M., Sassone-Corsi, P., & Davidson, I. (2001). Late arrest of spermiogenesis and germ cell apoptosis in mice lacking the TBP-like TLF/TRF2 gene. *Mol Cell, 7*(3), 509-515. doi:10.1016/s1097-2765(01)00198-8

Martin-Tryon, E. L., & Harmer, S. L. (2008). XAP5 CIRCADIAN TIMEKEEPER coordinates light signals for proper timing of photomorphogenesis and the circadian clock in Arabidopsis. *Plant Cell, 20*(5), 1244-1259. doi:10.1105/tpc.107.056655

Mazzarella, R., Pengue, G., Yoon, J., Jones, J., & Schlessinger, D. (1997). Differential expression of XAP5, a candidate disease gene. *Genomics, 45*(1), 216-219. doi:10.1006/geno.1997.4912

Mitchell, D. R. (2017). Evolution of Cilia. *Cold Spring Harb Perspect Biol, 9*(1). doi:10.1101/cshperspect.a028290

Nachury, M. V., & Mick, D. U. (2019). Establishing and regulating the composition of cilia for signal transduction. *Nat Rev Mol Cell Biol*. doi:10.1038/s41580-019-0116-4

Nantel, F., Monaco, L., Foulkes, N. S., Masquilier, D., LeMeur, M., Henriksen, K., . . . Sassone-Corsi, P. (1996). Spermiogenesis deficiency and germ-cell apoptosis in CREM-mutant mice. *Nature, 380*(6570), 159-162. doi:10.1038/380159a0

Nemajerova, A., Kramer, D., Siller, S. S., Herr, C., Shomroni, O., Pena, T., . . . Lize, M. (2016). TAp73 is a central transcriptional regulator of airway multiciliogenesis. *Genes Dev, 30*(11), 1300-1312. doi:10.1101/gad.279836.116

Oyama, Y., Miyata, H., Shimada, K., Larasati, T., Fujihara, Y., & Ikawa, M. (2022). TULP2 deletion mice exhibit abnormal outer dense fiber structure and male infertility. *Reprod Med Biol, 21*(1), e12467. doi:10.1002/rmb2.12467

Pertea, M., Pertea, G. M., Antonescu, C. M., Chang, T.-C., Mendell, J. T., & Salzberg, S. L. (2015). StringTie enables improved reconstruction of a transcriptome from RNA-seq reads. *Nature Biotechnology, 33*(3), 290-295. doi:10.1038/nbt.3122

Piasecki, B. P., Burghoorn, J., & Swoboda, P. (2010). Regulatory Factor X (RFX)-mediated transcriptional rewiring of ciliary genes in animals. *Proc Natl Acad Sci U S A, 107*(29), 12969-12974. doi:10.1073/pnas.0914241107

Qu, W., Yuan, S., Quan, C., Huang, Q., Zhou, Q., Yap, Y., . . . Zhang, Z. (2020). The essential role of intraflagellar transport protein IFT81 in male mice spermiogenesis and fertility. *Am J Physiol Cell Physiol, 318*(6), C1092-C1106. doi:10.1152/ajpcell.00450.2019

Reinholdt, L., Ashley, T., Schimenti, J., & Shima, N. (2004). Forward genetic screens for meiotic and mitotic recombination-defective mutants in mice. *Methods Mol Biol, 262*, 87-107. doi:10.1385/1-59259-761-0:087

Reiter, J. F., & Leroux, M. R. (2017). Genes and molecular pathways underpinning ciliopathies. *Nat Rev Mol Cell Biol, 18*(9), 533-547. doi:10.1038/nrm.2017.60

Robinson, J. T., Thorvaldsdottir, H., Turner, D., & Mesirov, J. P. (2023). igv.js: an embeddable JavaScript implementation of the Integrative Genomics Viewer (IGV). *Bioinformatics, 39*(1). doi:10.1093/bioinformatics/btac830

Romanienko, P. J., & Camerini-Otero, R. D. (2000). The mouse Spo11 gene is required for meiotic chromosome synapsis. *Mol Cell, 6*(5), 975-987. doi:10.1016/s1097-2765(00)00097-6

Santos, N., & Reiter, J. F. (2008). Building it up and taking it down: the regulation of vertebrate ciliogenesis. *Dev Dyn, 237*(8), 1972-1981. doi:10.1002/dvdy.21540

Schartl, M., Schories, S., Wakamatsu, Y., Nagao, Y., Hashimoto, H., Bertin, C., . . . Herpin, A. (2018). Sox5 is involved in germ-cell regulation and sex determination in medaka following co-option of nested transposable elements. *BMC Biol, 16*(1), 16. doi:10.1186/s12915-018-0485-8

Sedlacek, Z., Munstermann, E., Dhorne-Pollet, S., Otto, C., Bock, D., Schutz, G., & Poustka, A. (1999). Human and mouse XAP-5 and XAP-5-like (X5L) genes: identification of an ancient functional retroposon differentially expressed in testis. *Genomics, 61*(2), 125-132. doi:10.1006/geno.1999.5931

Singla, V., & Reiter, J. F. (2006). The primary cilium as the cell's antenna: signaling at a sensory organelle. *Science, 313*(5787), 629-633. doi:10.1126/science.1124534

Stubbs, J. L., Oishi, I., Izpisua Belmonte, J. C., & Kintner, C. (2008). The forkhead protein Foxj1 specifies
node-like cilia in Xenopus and zebrafish embryos. *Nat Genet, 40*(12), 1454-1460.
doi:10.1038/ng.267

Subramanian, A., Tamayo, P., Mootha, V. K., Mukherjee, S., Ebert, B. L., Gillette, M. A., . . . Mesirov, J. P.
(2005). Gene set enrichment analysis: A knowledge-based approach for interpreting genome-wide
expression profiles. *Proceedings of the National Academy of Sciences, 102*(43), 15545-15550.
doi:10.1073/pnas.0506580102

Swoboda, P., Adler, H. T., & Thomas, J. H. (2000). The RFX-type transcription factor DAF-19 regulates
sensory neuron cilium formation in C. elegans. *Mol Cell, 5*(3), 411-421. doi:10.1016/s1097-
2765(00)80436-0

Thomas, J., Morle, L., Soulavie, F., Laurencon, A., Sagnol, S., & Durand, B. (2010). Transcriptional control
of genes involved in ciliogenesis: a first step in making cilia. *Biol Cell, 102*(9), 499-513.
doi:10.1042/BC20100035

Thompson, N. A., Ranzani, M., van der Weyden, L., Iyer, V., Offord, V., Droop, A., . . . Adams, D. J.
(2021). Combinatorial CRISPR screen identifies fitness effects of gene paralogues. *Nature
Communications, 12*(1). doi:10.1038/s41467-021-21478-9

Vij, S., Rink, J. C., Ho, H. K., Babu, D., Eitel, M., Narasimhan, V., . . . Roy, S. (2012). Evolutionarily
ancient association of the FoxJ1 transcription factor with the motile ciliogenic program. *PLoS
Genet, 8*(11), e1003019. doi:10.1371/journal.pgen.1003019

Weidemann, M., Schuster-Gossler, K., Stauber, M., Wrede, C., Hegermann, J., Ott, T., . . . Gossler, A.
(2016). CFAP157 is a murine downstream effector of FOXJ1 that is specifically required for
flagellum morphogenesis and sperm motility. *Development, 143*(24), 4736-4748.
doi:10.1242/dev.139626

Wu, Y., Hu, X., Li, Z., Wang, M., Li, S., Wang, X., . . . Han, C. (2016). Transcription Factor RFX2 Is a Key
Regulator of Mouse Spermiogenesis. *Sci Rep, 6*, 20435. doi:10.1038/srep20435

Yu, G., Wang, L. G., & He, Q. Y. (2015). ChIPseeker: an R/Bioconductor package for ChIP peak
annotation, comparison and visualization. *Bioinformatics, 31*(14), 2382-2383.
doi:10.1093/bioinformatics/btv145

Yu, X., Ng, C. P., Habacher, H., & Roy, S. (2008). Foxj1 transcription factors are master regulators of the
motile ciliogenic program. *Nat Genet, 40*(12), 1445-1453. doi:10.1038/ng.263

Zhang, A., Skaar, D. A., Li, Y., Huang, D., Price, T. M., Murphy, S. K., & Jirtle, R. L. (2011). Novel
retrotransposed imprinted locus identified at human 6p25. *Nucleic Acids Res, 39*(13), 5388-5400.
doi:10.1093/nar/gkr108

Zhang, D., Xie, D., Lin, X., Ma, L., Chen, J., Zhang, D., . . . Han, C. (2018). The transcription factor
SOX30 is a key regulator of mouse spermiogenesis. *Development, 145*(11).
doi:10.1242/dev.164723

Zheng, M., Chen, X., Cui, Y., Li, W., Dai, H., Yue, Q., . . . Zhu, H. (2021). TULP2, a New RNA-Binding
Protein, Is Required for Mouse Spermatid Differentiation and Male Fertility. *Front Cell Dev Biol,
9*, 623738. doi:10.3389/fcell.2021.623738

Zhou, H., Grubisic, I., Zheng, K., He, Y., Wang, P. J., Kaplan, T., & Tjian, R. (2013). Taf7l cooperates with
Trf2 to regulate spermiogenesis. *Proc Natl Acad Sci U S A, 110*(42), 16886-16891.

Figures

Figure 1. Loss of XAP5 and XAP5L proteins results in male mice infertility.

(A) Detection of XAP5/XAP5L protein in different tissues of adult mice by Western blot

analysis. **(B)** Expression of XAP5/XAP5L protein during postnatal testicular

development. β-actin was used as control. **(C)** t-SNE plots displaying *XAP5*/*XAP5L* gene

expression levels in individual cell types of mouse testis visualized using published

single-cell data(Jung et al., 2019). Black arrow represents the developmental pseudotime

corresponding to the developmental ordering of each cell progressing through

spermatogenesis. Dot color intensity represents the expression level. Cells from different

stages are color-coded. **(D)** Immunofluorescence staining of XAP5/XAP5L in adult WT

testis. Representative spermatogonium (Sg), round spermatid (Rs) and elongating

spermatid (Es) are indicated. Scale bars: 20 μm (white), 5 μm (red). **(E)** Similar

development between WT and *XAP5L* KO mice. Scale bar: 1 cm. **(F)** Average litter size

of pups obtained by fertility test. Each male was caged with two females for 2 months, 5

males were used for each genotype. ***$P < 0.001$, n.s. stands for not significant. **(G)**

Similar appearance of WT and *XAP5* cKO mice. Scale bar: 1 cm. **(H)** Average litter size

of pups obtained by fertility test. Each male was caged with two WT females for 2

months, 4 males were used for each genotype. ***$P < 0.001$. Error bars depict means $\pm$

SEM. All *P*-values were calculated using an unpaired, two-tailed Student's t-test.

Figure 2. XAP5L is required for normal sperm formation.

(A) Similar gross testes morphology, body weight and testis weight between WT and *XAP5L* KO mice. Scale bar: 1 mm. n = 6, n.s. stands for not significant. **(B)** Significantly reduced caudal epididymal sperm counts in XAP5L KO mice compared with WT males. n = 5, **$P < 0.01$. **(C)** Significantly reduced sperm total motility of *XAP5L* KO mice. n =

3, **$P < 0.01$. **(D)** H&E staining showing similar histological structure of the testes from

WT and *XAP5L* KO mice. Scale bars: 100 μm. **(E)** H&E staining showing the

morphology of sperm in WT and *XAP5L* KO mice. Scale bars: 10 μm. **(F)** Significantly

increased abnormal sperm in *XAP5L* KO mice. n = 6, ***$P < 0.001$. **(G)** Electron

microscopic analysis displaying ultrastructural defects in *XAP5L* KO sperm flagella,

including cytoplasmic vacuoles enveloping mitochondrial sheath (as shown by red

arrowheads), a complete lack of mitochondrial sheath (as shown by red dotted circle),

two or more cross-sections of the same sperm flagellum enclosed within one cell

membrane (as shown by black arrowheads), partially formed outer dense fibers, and the

axonemal microtubules (as indicated by asterisks). Scale bars: 2 μm. Error bars depict

means ± SEM. All *P*-values were calculated using an unpaired, two-tailed Student's t-

test.

Figure 3. Ablation of XAP5 in germ cells results in arrested spermatogenesis and absence of sperm.

(A) Testes sizes, body weight and testis weight from WT and *XAP5* cKO mice at various ages were evaluated. Similar body weight was observed between WT and *XAP5* cKO mice. No significant difference in size and weight was detected between WT and *XAP5* cKO testes until P16. Scale bars: 1 mm. n = 5, ***P < 0.001, n.s. stands for not

significant. **(B)** Histology of testes from WT and *XAP5* cKO mice at various ages were

evaluated by H&E staining. No pachytene spermatocytes were found, and many

vacuolated tubules were observed in *XAP5* cKO testis since P16. Black arrowheads

indicate pachytene spermatocytes; black arrows indicate round spermatids. Vacuolated

seminiferous tubules are indicated by asterisks. Scale bars: 20 μm (black), 2.5 μm (red).

(C) Chromosome spreads staining showing pachytene spermatocytes from P16 WT male

mouse. Yellow dotted circle indicates the XY body. Scale bars: 10 μm. **(D)**

Immunofluorescence staining of PNA in adult testes indicating the absence of spermatids

in *XAP5* cKO mice. Scale bars: 20 μm. **(E)** H&E staining of adult epididymis displaying

absence of sperm in *XAP5* cKO mice. Scale bars: 50 μm. Error bars depict means ± SEM.

All *P*-values were calculated using an unpaired, two-tailed Student's t-test.

Figure 4. Aberrant ciliogenic and spermatogenic gene expression in *XAP5L* KO and

XAP5 cKO male germ cells.

(A) Gene ontology analysis showing top ten biological process terms significantly enriched among upregulated genes in *XAP5L* KO sperm. **(B)** Real-time PCR validation of genes involved in ciliogenesis and spermatogenesis. n = 3, **$P < 0.01$, ***$P < 0.001$, n.s. stands for not significant. **(C)** Detection of FOXJ1 and RFX2 in mature sperm of WT and *XAP5L* KO mice by Western blot analysis. GAPDH was used as a control. **(D)** Gene ontology analysis showing top ten biological process terms significantly enriched among downregulated genes in P16 *XAP5* cKO testes. **(E)** Real-time PCR validation of genes involved in ciliogenesis and spermatogenesis. n = 3, *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, n.s. stands for not significant. **(F)** Detection of FOXJ1 and RFX2 in P16 testes of WT and *XAP5* cKO mice by Western blot analysis. GAPDH was used as a control. **(G)** XAP5/XAP5L occupancies at *FOXJ1* and *RFX2* promoters assessed by CUT&Tag-seq (left) and CUT&Tag real-time PCR (middle and right). Red boxes indicate the binding regions of XAP5 and XAP5L. n = 3, *$P < 0.05$, ***$P < 0.001$, n.s. stands for not significant. **(H)** A model on the involvement of XAP5 and XAP5L in the regulation of ciliogenesis during spermatogenesis. Error bars depict means ± SEM. All *P*-values were calculated using an unpaired, two-tailed Student's t-test.

Supplementary Figures

Figure S1. Localization of XAP5/XAP5L in testicular cells.

(A) Western blot analysis showing no localization of XAP5/XAP5L in mature sperm. Western blot analysis showing the subcellular localization of endogenous XAP5 (**B**) and XAP5L (**C**). LMNB used as a nuclear protein marker, GAPDH used as a cytoplasmic protein marker. **(D)** Immunofluorescence staining of XAP5/XAP5L in adult epididymis. Scale bars: 50 μm.

Figure S2. Generation of *XAP5L* KO and *XAP5* cKO mice.

(A) Schematic illustration of the targeting strategy for generating a *XAP5L* knockout mouse line using CRISPR-Cas9 system. Upper panel showing genomic structure of the *XAP5L* gene. The location of two sgRNAs (sgRNA1 and sgRNA2) used and three primers (blue arrows) for genome PCR are indicated. Lower panel displaying nucleotide

sequence of WT allele in the vicinity of two sgRNAs (red color) and the corresponding

sequence of the *XAP5L* homozygous mutant (KO) allele. The deletion of *XAP5L* gene

was verified by Sanger sequencing. **(B)** Genome PCR analysis in WT and *XAP5L* KO

mice. **(C)** Detection of XAP5L in adult testes extracts of WT and *XAP5L* KO mice by

Western blot analysis. β-actin was used as a control. **(D)** Schematic illustration of the

targeting strategy for generating *XAP5* cKO mice. The location of LoxP sites inserted and

the primers (blue arrow) used for genome PCR are indicated. **(E)** Genome PCR analysis

in WT and *XAP5* cKO mice. **(F)** Detection of XAP5 in testes extracts of WT and *XAP5*

cKO mice by Western blot analysis. β-actin was used as a control.

Figure S3. XAP5L-mediated regulation of ciliary gene expression during mouse spermatogenesis.

(A) MA plot showing the numbers of differentially expressed genes in *XAP5L* KO sperm.

Gene ontology analysis showing top ten cellular component (**B**) and molecular function

(**C**) terms significantly enriched among upregulated genes in *XAP5L* KO sperm. **(D)** Venn

diagram representing total genes upregulated/downregulated by at least twofold in

XAP5L KO sperm and published cilia-related genes(Nemajerova et al., 2016). **(E)** Real-

time PCR validation of genes involved in ciliogenesis and spermatogenesis. n = 3, *P <

0.05, ***P < 0.001. Error bars depict means ± SEM. All *P*-values were calculated using

an unpaired, two-tailed Student's t-test.

Figure S4. *TULP2* KO male mice were infertile due to malformation of sperm.

(A) Detection of TULP2 protein in different tissues of adult mice by Western blot

analysis. **(B)** t-SNE plot displaying *TULP2* gene expression pattern during spermatogenesis visualized using published single-cell data(Jung et al., 2019). Black arrow represents the developmental pseudotime corresponding to the developmental ordering of each cell progressing through spermatogenesis. Dot color intensity represents the expression level. Cells from different stages are color-coded. **(C)** Immunofluorescence staining showing TULP2 expression in the cytoplasm of round spermatids (Rs) and elongating spermatids (Es). Scale bars: 5 μm. **(D)** Similar development of adult WT and *TULP2* KO mice. Scale bar: 1 cm. **(E)** Average litter size of pups obtained by fertility test. Each male was caged with two females for 2 months. ***$P < 0.001$, n.s. stands for not significant. **(F)** Similar gross testes morphology, body weight and testis weight between WT and *TULP2* KO mice. Scale bar: 1 mm. n = 4, n.s. stands for not significant. **(G)** Significantly reduced caudal epididymal sperm counts in *TULP2* KO mice compared with WT males. n = 7, ***$P < 0.001$. **(H)** Significantly reduced sperm total motility of *TULP2* KO mice. n = 3, ***$P < 0.001$. Significantly increased abnormal sperm in *TULP2* KO mice (**I** and **J**). Scale bars: 10 μm. n = 3, ***$P < 0.001$. **(K)** Electron microscopic analysis displaying ultrastructural defects in *TULP2* KO sperm flagella, including two cross-sections of the same sperm flagellum enclosed within one cell membrane (as shown by red arrowheads), disorganized outer dense fibers and the axonemal microtubules (as shown by red dotted circle). Scale bars: 8 μm (black), 2 μm (red). Error bars depict means ± SEM. All *P*-values were calculated using an unpaired, two-tailed Student's t-test.

Figure S5. XAP5-mediated regulation of ciliary gene expression during mouse

spermatogenesis.

(A) Heat map showing variably expressed genes in P16 *XAP5* cKO testes. Number of

related genes was indicated. Gene ontology analysis showing top ten cellular component

(**B**) and molecular function (**C**) terms significantly enriched among downregulated genes

in *XAP5* cKO testes. **(D)** Venn diagram representing total genes

upregulated/downregulated in *XAP5* cKO testes and published cilia-related

genes(Nemajerova et al., 2016). **(E)** Venn diagram representing total genes

downregulated in *XAP5* cKO testes, upregulated in *XAP5L* KO sperm by at least twofold

and published cilia-related genes(Nemajerova et al., 2016). **(F)** Real-time PCR validation

of genes involved in ciliogenesis and spermatogenesis. n = 3, *$P < 0.05$, **$P < 0.01$, ***P

< 0.001. Error bars depict means ± SEM. All *P*-values were calculated using an unpaired,

two-tailed   Student's   t-test.

Figure S6. Identification of XAP5/XAP5L-occupancy sites

Location of XAP5 (**A**) and XAP5L (**B**)-occupancy peaks relative to the nearest annotated

gene. (**C**) DNA sequence identified by Homer de novo motif analysis from

XAP5/XAP5L peaks. The P value of the motif comparison is indicated. Overlapped

sequences are shown in red boxes. (**D**) Left, overlapping genes identified by XAP5

CUT&Tag and RNA-seq analyses. Right, representative biological processes enriched

among the overlapping genes that are downregulated in *XAP5* cKO germ cells and bound

by XAP5 (as identified by CUT&Tag). (**E**) Left, overlapping genes identified by XAP5L

CUT&Tag and RNA-seq analyses. Right, representative biological processes enriched

among the overlapping genes that are upregulated in *XAP5* KO sperm and bound by

XAP5L (as identified by CUT&Tag). (**F**) XAP5/XAP5L occupancies at *SOX30* promoter

assessed by CUT&Tag-seq (top) and CUT&Tag real-time PCR (bottom). Red boxes

indicate the binding regions of XAP5/XAP5L. n = 3, *$P < 0.05$, **$P < 0.01$. (**G**)

Overlapping genes among XAP5/XAP5L target genes (the 891 genes in Figure S6D and

the 75 genes in Figure S6E) and published cilia-related genes(Nemajerova et al., 2016).

Error bars depict means ± SEM. All *P*-values were calculated using an unpaired, two-

tailed Student's t-test.

Figure S7. t-SNE plots displaying expression patterns of *FOXJ1* and *RFX* families in individual cell types of mouse testes.

The black arrow represents the developmental pseudotime corresponding to the developmental ordering of each cell progressing through spermatogenesis. Dot color intensity represents the relative expression level. Cells from different stages are color-coded. The gene expression was visualized using the published single-cell data(Jung et al., 2019).