Increased inflammatory signature in myeloid cells of non-small cell lung cancer

patients with high clonal hematopoiesis burden

Hyungtai Sim,[1]* Hyun Jung Park,[2,3]* Geun-Ho Park, [4,5]*, Yeon Jeong Kim,[2] Woong-

Yang Park,[2] Se-Hoon Lee, [4,5]** Murim Choi[1]**

*, ** These authors contributed equally

Affiliations:

[1]Department of Biomedical Sciences, Seoul National University College of Medicine,

Seoul, Republic of Korea

[2]Samsung Genome Institute, Samsung Medical Center, Seoul, Republic of Korea

[3]Research Institute for Veterinary Science, College of Veterinary Medicine, Seoul National

University, Seoul, Republic of Korea

[4]Division of Hematology-Oncology, Department of Medicine, Samsung Medical Center,

Sungkyunkwan University School of Medicine, Seoul, Republic of Korea

[5]Department of Health Sciences and Technology, Samsung Advanced Institute for

Health Science & Technology (SAIHST), Sungkyunkwan University, Seoul, Republic of

Korea

Corresponding authors:

Murim Choi, Department of Biomedical Sciences, Seoul National University College of

Medicine,103 Daehak-ro, Jongno-gu, Seoul 03080, Republic of Korea

Email: murimchoi@snu.ac.kr

Se-Hoon Lee, Division of Hematology-Oncology, Department of Medicine, Samsung

Medical Center, Sungkyunkwan University School of Medicine, 81 Irwon-ro, Gangnam-

gu, Seoul 06351, Republic of Korea

Email: sehoon.lee119@gmail.com

Running title: Increased inflammatory signature in NSCLC patients with CHIP

Keywords: Clonal Hematopoiesis, Immune checkpoint inhibitor, Inflammation, NSCLC

Abstract

Clonal hematopoiesis of indeterminate potential (CHIP) allows estimation of clonal

dynamics and documentation of somatic mutations in the hematopoietic system. Recent

studies utilizing large cohorts of the general population and patients have revealed

significant associations of CHIP burden with age and disease status, including in cancer

and chronic diseases. An increasing number of cancer patients are treated with immune

checkpoint inhibitors (ICI), but the association of ICI response in non-small cell lung

cancer (NSCLC) patients with CHIP burden remains to be determined. We collected

blood samples from 100 metastatic NSCLC patients before and after ICI for high-depth

sequencing of the CHIP panel and 63 samples for blood single-cell RNA

sequencing(scRNA-seq). Whole exome sequencing (WES) was performed in an

independent replication cohort of 180 patients. The impact of CHIP status on the

immunotherapy response was not significant. However, metastatic lung cancer patients

showed higher CHIP prevalence (44/100 for patients vs 5/42 for controls; P = 0.01). In

addition, lung squamous cell carcinoma patients showed increased burden of larger

clones compared to lung adenocarcinoma patients (8/43 for LUSC vs 2/50 for LUAD; P

= 0.04). Furthermore, single cell RNA-seq analysis of the matched patients showed

significant enrichment of inflammatory pathways mediated by NF-κB in myeloid clusters

of the severe CHIP group. Our findings suggest minimal involvement of CHIP mutation

and clonal dynamics during immunotherapy but a possible role of CHIP as an indicator

of immunologic response in NSCLC patients.

Introduction

Recent studies have revealed that somatic mutations detected in hematopoietic malignancies are also commonly observed with clonal expansion in the individuals with no hematological conditions (1). Among such individuals, asymptomatic cases with a variant allele frequency (VAF) of ≥ 2% are defined as clonal hematopoiesis of indeterminate potential (CHIP) (2). In general, CHIP is strongly associated with aging; and mutagenic anticancer therapies and smoking can also induce clonal expansion in specific genes (1,3). Longitudinal monitoring of CHIP mutations offers important clues into the clonal dynamics of blood (4).

Hematopoietic stem and progenitor cells (HSPCs) carrying CHIP mutations serve as a major source of expanded clones, and even a few cells are sufficient to influence clonal homeostasis (5). Representative CHIP mutation-carrying genes, such as *DNMT3A*, *ASXL1*, and *TET2*, impact the myeloid population, including monocytes and granulocytes, and have been linked to hematologic malignancies such as myelodysplastic syndrome and acute myeloid leukemia through inflammation mediated by the NF-kB and NLRP3 inflammasome (6-8). The prevalence of CH in cancer patients is also significantly higher than that in healthy populations, and high CH burden has been implicated in a negative effect on overall survival in multiple cancer types (9,10). CHIP is also associated with diseases beyond hematologic malignancies, such as atherosclerosis, type 2 diabetes, and chronic obstructive pulmonary disease (11-13). Multiple studies, including both population and pan-cancer cohorts, have explored the correlation of CHIP with solid tumors (9,10,14,15). However, establishing a causal

relationship has proven challenging due to confounding factors such as history of

smoking and anticancer therapy, both of which act as risk factors for CHIP (10).

Non-small cell lung cancer (NSCLC) stands out globally as the solid tumor with the

highest mortality burden (16). The difficulty in early detection of NSCLC significantly

contributes to its mortality rate being high compared to its incidence rate. Recent

studies examining CHIP within lung cancer cohorts and general population indicate a

possible association of lung cancer risk with CH (15,17,18). As mentioned above,

factors such as a patient's history of anticancer treatments and smoking serve as

common risk factors for CH; however, the interplay between lung cancer and CHIP is

not yet fully understood. Immune checkpoint inhibitors (ICI) targeting PD-(L)1 and

CTLA-4 have revolutionized treatment of advanced NSCLC, both as monotherapies and

in combination with chemotherapy (19). However, response rates to ICI treatments

remain relatively low. The predictive ability of markers like PD-1 expression and tumor

mutational burden (TMB) is insufficient, requiring the identification of additional markers.

Recent studies suggest a marginal effect of ICI on the clonal dynamics of CHIP (20,21),

but the current understanding of the role of CHIP in ICI-treated patients is not

comprehensive. Also, only limited analysis has been performed concerning the impact

of CHIP on cancer patients' blood. To understand the interplay between ICI and CHIP in

NSCLC patients, we conducted CHIP-targeted panel sequencing and single-cell RNA

sequencing (scRNA-seq) on blood samples collected before and 1-3 weeks after ICI

administration. Our results indicate a marginal interaction between ICI and CHIP, but

also validate that high CHIP burden manifests an inflammatory phenotype in metastatic

NSCLC patients' myeloid cells.

Materials and Methods

Patients

Description of the discovery cohort is previously reported (22). Briefly, 100 patients (50

LUAD, 43 LUSC and 7 others) with metastatic NSCLC stage IV who were treated with

anti-PD(L)1 (*e.g.*, Atezolizumab, Nivolumab, and Pembrolizumab) were recruited at

Samsung Medical Center (SMC) under the permission of SMC Institutional Review

Board (No. 2018-04-048, 2022-01-094). Whole blood sample was acquired before and

one to three weeks after the treatment and used for genome sequencing and scRNA-

seq. For replication, blood samples from additional 180 lung cancer patients (125 LUAD

and 55 LUSC) were used for genome analysis (23).

Targeted sequencing and whole exome sequencing (WES)

We designed a targeted sequencing panel comprising 167 selected genes associated

with putative drivers of CHIP at Twist Bioscience (South San Francisco, CA). For WES,

we utilized the SureSelect V5 panel from Agilent (Santa Clara, CA). CHIP-targeted

sequencing and WES were both conducted at Samsung Genome Institute, respectively

employing 50 ng and 200 ng of genomic DNA extracted from whole blood. The targeted

sequencing panel was captured and sequenced using NextSeq 2000, with paired-end

sequencing performed at a read length of 150 base pairs (bps).

Variant filtering and calling of CHIP variants

For somatic variant calling, the Sarek pipeline (version 3.2.1) implemented in Nextflow

was used (24). In brief, the pipeline employed GATK's standard data preprocessing

procedure and Mutect2 for somatic variant calling based on a BED file (25). All passed

calls were collected and annotated using Snpeff (version 4.3 and db version 105) and

VEP (version 108) (26,27). To filter CHIP variants, a stepwise approach based on the

following criteria was applied. First, variants should meet the following conditions: a)

variant allele frequency (VAF) 2-35%, b) variant depth >500x for the discovery set and

>40x for the WES replication set, c) 4 or more (2 or more for the WES replication set)

variant-covering read pairs in each forward and reverse direction, d) global allele

frequency from gnomAD <1e-5, or <1e-3 if a variant is in the COSMIC database (28))

allele count <5% in all processed samples, and f) no homopolymer signature found in 6

bp upstream or downstream of the candidate variant. Next, we called CHIP variants

based on a) protein sequence alterations, including exonic splicing variants, and b)

manual assessment of variant calls using IGV (29). Finally, we curated data from

previous studies and the COSMIC database, assigning variants reported more than 10

times or matched within a predefined list as CHIP with putative drivers (called as clonal

hematopoiesis of indeterminate potential with putative drivers, or CHIP-PD) (30).

Power estimation

The power of the Poisson binomial test was calculated as the probability that a variant

would be detected exclusively in one of two arbitrary groups. In brief, the most prevalent

CHIP variant, *DNMT3A* p.Arg882, constitutes approximately 10% of the *DNMT3A* CHIP

cases. When this variant is totally absent in the one group, a minimum of 16 *DNMT3A*

CHIP patients in the other group would be required to achieve a power of 0.8.

Considering the current prevalence of *DNMT3A* CHIP in our cohort (8/50 CHIP

samples), this suggests that a minimum of 100 CHIP samples per group is required for

determining the prevalence of any specific variant.

scRNA-seq pre-processing and analysis

The scRNA-seq dataset was generated using the 10x Genomics 5' single-cell kit and

was partially derived from a previous study (22). Reads were aligned to the hg38

reference using 10X CellRanger (ver. 7.0). Contaminant RNA barcodes were removed

with cellbender and sample demultiplexing was performed by demuxlet (31,32). After

filtering the cell expression matrix, we used Seurat (v 4.3.0) for downstream analysis

(33). At the sample level, we filtered out cells with mitochondrial RNA content >15%,

UMI counts <200 or >6,000, and calculated library-size corrected counts using

SCTransform (34). Then, we computed PCA based on the normalized SCTransform

values and integrated all samples using Harmony (35). Using the corrected PCA, we

performed unsupervised clustering via Louvain algorithms and UMAP. For each cluster,

we identified marker genes using the Wilcoxon rank-sum test with presto and validated

annotations using Azimuth (33).

Gene set enrichment analysis (GSEA)

GSEA was employed to compare the expression patterns according to CHIP status

based on their concordance to the hallmark pathways in the Molecular Signatures

Database (MSigDB) (36). For GSEA of scRNA-seq data, we used fgsea (37). Briefly,

the AUC rank of DEGs obtained with and without CHIP for each cluster was used as

input. Significant gene set enrichment was determined only for adjusted $P < 0.05$.

scRNA-seq weighted correlation network analysis (WGCNA)

We utilized WGCNA to construct a gene co-expression network from the gene

expression profiles of myeloid cells utilizing the R package hdWGCNA (38). In brief, the

co-expression network was constructed using a soft power of 12 for SCT-transformed

cell counts. Within the constructed co-expression network, we explored the type and

specificity of genes in each module, excluding unassigned genes. The functionality of

the black module was validated through GSEA and by examining its overlap with DEGs

identified between CHIP groups (38).

Gene regulatory network (GRN) inference

To further understand the altered gene modulation in major cell clusters, including B

cells, NK cells, CD4 T cells, CD8 T cells, and monocytes, we constructed a GRN using

pySCENIC (39). We created meta-cells for each sample using SEACells and employed

these meta-cells for GRN calculation to address the sparse nature of scRNA expression

and enhance resolution (40,41). The full transcriptome matrix and the GRN derived from

meta-cells were then used to analyze regulon signatures for each cell. Finally, we

mapped the regulon AUC values of each cell onto Seurat's UMAP and visualized the

results using Seurat.

Cell to cell communication analysis

To understand the interactions between myeloid and other cell types, we utilized the

Python package CellphoneDB (42), which curates human cell-cell interactions. The

counts matrix, processed with SCTransform, was divided according to CHIP status and

used as input for CellphoneDB. Significant interactions were then counted to determine

the number of cell-cell interactions, and myeloid-related interactions were compared

between samples of different CHIP status.

Statistical analysis

We present raw data if applicable. For scRNA-seq data, scaled and log-normalized raw

counts from cells are displayed in violin plots. To determine statistical significance

between high-burden, CHIP, and negative groups, the Wilcoxon test or Kruskal-Wallis

test was used for continuous variables and the Poisson binomial test or logistic

regression for categorical variables. Statistical metrics from GESA were calculated

using the fgsea package in R. All statistical analyses were conducted in R 4.2.1.

Data availability statement

Raw data for this study were generated at Samsung Genome Institute. Derived data

supporting the findings of this study are available from the corresponding author upon

request.

Results

CHIP profiles in metastatic NSCLC patients treated with ICI

To investigate the interplay between ICI treatment and CHIP in NSCLC, we collected

blood samples before and after ICI treatment from 100 metastatic NSCLC patients (Fig.

1, Supplementary Table S1). These samples were utilized for a CHIP targeted

sequencing panel (median depth ~600x) and scRNA-seq (**Fig. 1A** and Methods). At the

baseline, we identified 67 CHIP mutations in 100 patients, 26 of them in known putative

driver genes of hematopoietic cancer (CHIP-PD). After treatment, 68 CHIP and 32

CHIP-PD mutations were found in 91 patients who passed quality control assessment

(**Fig. 1A**, Supplementary Fig. S1-S3 and Supplementary Table S3).

Among the CHIP genes, *DNMT3A*, *PPM1D* and *TET2* were the most frequently mutated

(**Fig. 1B**). *PPM1D* truncating mutations were also common, and have previously

demonstrated association with patients' smoking habits and possible history with

chemotherapy (3,9). Variants involving the *DNMT3A* p.Arg882 residue and *TP53*, which

are frequent in CHIP, were not prevalent in our dataset, although our panel extensively

covered these regions and these loci were manually inspected (data not shown).

Overall, we observed ICI treatment to effect minimal changes in CHIP burden (**Fig. 1B**).

Also, VAF was highly correlated across all traced mutations and not dependent on

clinical responses (**Fig. 1C** and **D**). However, we noted increased number of CHIP

mutations with VAF >10% in patients with lung squamous cell carcinoma (LUSC). While

the specific mutated genes differed between patients with different histology or

responses, mutated gene burden did not significantly differ between groups.

Prevalence of high CHIP burden showed bias to LUSC patients

Next, we investigated the association of CHIP status with clinical parameters in the

cohort (**Fig. 2**). CHIP prevalence was significantly higher in the lung cancer cohort

compared to the control group (controls 5/42 vs. patients 44/100, age-adjusted logistic

regression $P = 0.01$; **Fig. 2A**, Supplementary Table S1 and S2). However, the number

of CHIP-positive patients did not differ before and after ICI treatment, nor in relation to

treatment response (**Fig. 2B** and **C**). As previously reported, CHIP burden is heavily

dependent on age (**Fig. 2D**). Smoking history may also impact CHIP prevalence, but we

could not find a significant contribution of smoking, as almost all NSCLC patients in our

study have a smoking history (CHIP prevalence of non-smoker 2/9 vs. smoker 42/91;

Supplementary Fig. S4).

Subsequently, we examined clonal size in the cohort, as that has been reported as a

risk factor of clinical outcome (43). Taking CHIP VAF as an indicator of clonal size, we

observed that high CHIP burden was prevalent in LUSC, while multiple concurrent CHIP

mutations were common in lung adenocarcinoma (LUAD) patients (**Fig. 2C** and

Supplementary Fig. S5). To determine if this finding would reproduce in an independent

cohort, we performed CHIP mutation calling using WES data of median ~80x depth from

180 additional peripheral blood mononuclear cell (PBMC) samples from an independent

cohort of NSCLC patients (**Fig. 2E-G**, Supplementary Fig. S6, Supplementary Table S1,

S3-S4 and Methods). The results showed similar patterns between the two cohorts, but

the replication cohort alone was not powerful enough to achieve significance for

difference in clonal sizes based on pathology (**Fig. 2E-G**).

High CHIP burden causes single-cell transcriptomic changes

Next, to test whether the presence of CHIP and ICI prognosis affects blood cell

transcriptomes, we analyzed PBMC scRNA-seq data from 63 samples of the discovery

cohort. After quality control processes, we retrieved 468,596 cells in a total of 26

clusters defined by systemic integration and clustering using the Louvain algorithm, then

validated using Azimuth (**Fig. 3A**, Supplementary Fig. S7A-B, Supplementary Table S5,

and Methods) (33). Cell composition analysis showed an increased proportion of NK

cells and a decreased CD4 TEM population after ICI administration, but high variability

between samples precluded systematic differences by CHIP status (Supplementary Fig.

S7C). Subsequently, DEG analysis using Wilcoxon's rank-sum test according to CHIP

VAF bins in each cluster (see Methods) highlighted genes important in immune

response and transcriptional activation (**Fig. 3C**, Supplementary Fig. S8). To further

implicate these DEGs functionally, a gene set enrichment test was performed using the

AUC statistics from DEGs and the hallmark pathway gene sets in msigDB (36). The

results indicated activation of the TNF signaling pathway via *NFKB* in the high CH

burden group for most cell lineages, including both classical dendritic cells (DC) and NK

cells (**Fig. 3D**, **3E**, Supplementary Fig. S9, and Supplementary Table S6; see Methods).

To understand possible interactions between ICI treatment response and CHIP burden,

we grouped samples by CHIP burden and ICI response (Supplementary Fig. S10A-B).

However, the interactions mostly depended on CH status only, and not on response to

ICI.

Implication of high CHIP burden in inflammatory signatures of myeloid cells

We performed a gene network analysis to comprehensively analyze the increased

expression of inflammatory signaling genes in patients with higher CHIP burden. First,

we used hdWGCNA to construct co-expression networks for genes in myeloid cells,

which yielded a black module that exhibited myeloid specificity and represented most of

the signaling-related DEGs identified via GSEA (Supplementary Fig. S6 and S11 and

Supplementary Table S7) (38). We determined gene ontology (GO) term enrichment in

this module and found the inflammatory and transcriptional activation pathways

observed in DEGs to be clearly identified (Supplementary Fig. S11C). Additionally,

further GSEA on the myeloid-specific module showed its function to resemble that of the

overall set of DEGs (Supplementary Fig. S11D).

In addition to WGCNA, GRN analysis using SCENIC (39) (Supplementary Fig. S12)

indicated that the regulatory activity of transcription factors in the AP-1 and NF-kB

pathways was particularly notable in the myeloid lineage of high-burden patients

(Supplementary Fig. S12A-B). Both the NF-kB regulon and the ATF3 regulon, which

represents the AP-1 pathway, were activated in the myeloid cluster of high-burden

patients (Supplementary Fig. S12C-E).

Recent studies have shown that expansion of HSPC-derived monocytes is the cause of

CHIP, and representative CHIP genes, including *DNMT3A* and *TET2*, exert immune-

regulatory effects through the NLRP3 inflammasome (5,8,44). Our network analysis

also points to this pathway (Supplementary Fig. S12). Notably, a VAF of 10% in

heterozygote mutations corresponds to approximately 20% of cells; thus, the presence

of such mutations in HSPC-derived monocytes of high-burden patients would have

resulted in an overall transcriptional profile change for blood monocytes.

In addition to conducting gene network analysis, we explored the effect of the

inflammatory signature on cell-cell interaction using CellPhoneDB (Supplementary Fig.

S13) (42). The number of significant cell-cell interactions did not differ significantly

between high-burden CHIP samples and negative samples, although the number of

cells in each group seemed likely to have an impact (Supplementary Fig. S13A-B).

However, when the interactions in the TNF and IL-1B pathways were analyzed

separately, monocytes and DCs in high-burden cases were activated as receptors for

TNF and Lymphotoxin A (Supplementary Fig. S13C). Collectively, our results suggest a

sensitized response of monocytes to inflammation in lung cancer patients with high

CHIP burden.

Effect of CHIP on clinical outcome

Finally, to understand whether clonal hematopoiesis status can affect the clinical

outcomes of lung cancer patients undergoing ICI treatment, we conducted a survival

analysis (Supplementary Fig. S14). Our results showed the progression-free survival

(PFS) of patients after ICI treatment to be reduced in the high-burden CHIP group (825

vs 404, $P = 0.056$), with no difference in ICI response rates between the negative and

high-burden CHIP groups. Previous studies have already observed a negative effect on

life expectancy in CHIP patients with solid tumors. Therefore, it is possible that these

results will be validated by analyzing additional samples.

Discussion

To address the influence of CH in lung cancer patients with differential prognosis following ICI, we sought to determine the following: (i) CH profiles from metastatic NSCLC patients, (ii) changes in PBMC gene expression by scRNA-seq, (iii) relationship of CH burden and transcriptome profile, and (iv) potential implication for the clinical outcomes. Here we report that CH prevalence was significantly higher in NSCLC patients and higher burden of mutations with VAF of greater than 10% was frequently observed in LUSC. Using scRNA-seq analysis, we found myeloid cells in NSCLC patients with high CH burden to exhibit an altered inflammatory milieu via increased NF-kB pathway expression. Our results provide possible explanations for heterogeneity in response to immunotherapy and suggest immune response to be altered due to CHIP in cancer patients.

Recent studies have employed mouse models and single-cell transcriptomics to understand the roles of specific mutations in CH and provided evidence of their implications for inflammatory response in the myeloid lineage (45,46). In particular, a study linking COVID-19 and CH showed an increased inflammatory response from monocytes in severe COVID-19 patients with CH, mediated via IFN-γ and TNF-α signaling signatures (47).Likewise, we observed that myeloid cells from NSCLC patients with CH tend to harbor increased inflammatory response, and this response is mediated by NF-kB pathway signaling (**Fig. 3**).

Our GSEA results specifically indicated the enrichment of TNF signaling and hypoxia pathways in most clusters of patients with severe CH (Supplementary Fig. S9). The

leading-edge genes from GSEA results showed core genes such as *FOS*, *DUSP1*, *JUN*,

and *PPP1R15A,* which are known to play critical roles in the inflammatory phenotypes

of immune cells, were shared between the TNF signaling and hypoxia pathways in all

significant clusters (Supplementary Fig. S15). Furthermore, gene regulatory network

analysis using SCENIC indicated a higher enrichment of inflammatory signatures in

myeloid lineages (Supplementary Fig. S9), highlighting the pronounced inflammatory

phenotype of CH clones in these cell lineages. To gain insight into the functional

implications of myeloid-driven inflammatory gene pathways within CH in NSCLC

patients, we conducted gene network analyses focusing on the myeloid lineage. The

"black module" from the WGCNA analysis (Supplementary Fig. S9) and the NFKB1 and

ATF3 GRN from SCENIC (Supplementary Fig. S11) revealed activation and adaptation

of myeloid-specific inflammatory pathways (38,39). Additionally, cell-cell interaction

analysis using cellphoneDB unveiled notable interactions between myeloid cells and

other lineages mediated through the TNF pathway (Supplementary Fig. S12) (42).

As ICIs modulate pathways involved in immune response rather than cell proliferation, it

is reasonable that ICIs do not exert a direct influence on CH development (21). Also,

our results did not support significant difference of ICI outcome based on CH burden

(**Fig. 2**). According to our power analysis, a minimum of 100 samples per group with

CHIP is required to achieve a statistical power of 0.8 from pair-wise comparisons

between each genes and variants (Methods for details). However, our findings do

suggest that elevated inflammatory activity in the bloodstream may impact long-term

patient survival pre- and post-ICI administration (Supplementary Fig. S14). Therefore, a

future study with a larger sample size may yet clarify a causal relationship of CH with

ICI outcome. Notably, involvement of the IL-1 pathway in the aggressiveness and

metastasis of lung cancer has been reported in experimental studies (48,49), and

inhibition of IL-1β has proven effective in reducing lung cancer-related mortality, as

shown in the CANTOS trial (50). Indeed, our analysis indicated a modest decrease in

PFS in the ICI-responder group with high-CH patients (Supplementary Fig. S14, $P =$

0.056). Our study design is potentially limited by the relatively short observation period

for defining ICI response Indeed, it is worth noting that more potent factors, such as

smoking and anticancer therapies, may have a more substantial impact than

immunotherapy (10). Consistent with this postulation, our analyses indicated minimal

changes in in CH-related signatures before and after ICI treatment (**Fig. 3**,

Supplementary S7, S9, and S10).

Also, the distinct characteristics of our cohort can be confounders for our results.

Compared to control patients, our cohort was biased to slightly older ages, a higher

prevalence of smoking habits, and with a higher proportion of males (mean age: 64.1 vs.

58.9; current smokers: 37/100 vs. 11/42; male/female: 91/9 vs. 18/24 Supplementary

Figures S1 and S3). However, previous studies have reported similar prevalence rates

of clonal hematopoiesis in NSCLC patients, aligned with our findings (9,51). Moreover,

our most prevalent CH mutations, including *DNMT3A*, *TET2*, and *PPM1D*, were

marginally affected by smoking, and this trend has been consistently observed in both

healthy populations and NSCLC patients (10,51,52).

Here we have illustrated increased inflammatory activation in myeloid cells of NSCLC

patients with high CH burden. Our scRNA-seq analysis has revealed that NF-kB

signaling mediated by the IL-1β and the TNF superfamilies may enhance interactions of

myeloid cells with other cell lineages. These findings propose the potential utility of CH

as a predictive marker for classifying NSCLC patients. Future investigations involving

ICI-treated patients and their clinical outcomes may shed light on the impact of CH,

offering a basis for the development of adjuvant anticancer strategies aimed at

modulating CH.

Authors' contributions

H. Sim: Data curation, formal analysis, validation, investigation, writing-original draft. **H.-J. Park**: Conceptualization, data curation, formal analysis, methodology, investigation. **G.-H. Park**: Data curation, formal analysis, methodology, investigation. **Y. J. Kim**: Data curation, methodology, investigation. **W.-Y. Park**: Conceptualization, investigation, methodology. **S. H. Lee**: Conceptualization, investigation, writing-original draft. **M. Choi**: Conceptualization, investigation, writing-original draft.

Acknowledgements:

We thank George Vassiliou for critical comments. The authors also acknowledge the Korea Research Environment Open Network (KREONET) service and the usage of the Global Science Experimental Data Hub Center (GSDC) provided by Korea Institute of Science and Technology Information (KISTI). This work was supported in part by the grants from the Korean Research Foundation (NRF-2021R1A2C3014067, NRF-RS-2023-00207857 to M. Choi, and NRF-2020R1A2C3006535 to S.-H. Lee), the Korea Health Technology R&D Project through the Korea Health Industry Development Institute (KHIDI) (HR20C0025 to S.-H. Lee), and Future Medicine 20*30 Project of the Samsung Medical Center (SMX1230041 to S.-H. Lee).

Conflict of interest:

Authors declare no conflict of interest.

Figures

Figure 1. Profile of CH in NSCLC patients treated with ICI. **A,** Overall study design. The

numbers indicate sample counts in each group. **B,** Frequency of CH variant detection in

each gene in NSCLC samples before (light blue) and after (blue) ICI treatment. **C,**

Comparison of variant allele frequency before and after ICI treatment, divided by

pathology type (LUAD, LUSC, other). **D,** Effect of ICI treatment on the clonal landscape

of all genes (left) and frequently detected CH genes.

Figure 2. Prevalence of CH in relation to various parameters. **A-D,** discovery cohort (*n*

= 100), **E-G,** replication cohort (*n* = 180; see Methods for description). **A**, **E,** CH

prevalence by pathology (LUSC and LUAD). **B,** Effect of ICI treatment on CH. **C**, **F,**

Effect of post-ICI prognosis to ICI on CH. **D**, **G,** Age distribution of the cohort, stratified

by CHIP allele frequency and pathology status.

Figure 3. scRNA-seq analysis detected myeloid-specific inflammatory signatures in

patients with high CH burden. **A,** UMAP plot of scRNA-seq data from the discovery

cohort ($n = 63$). **B,** Effect of pathology (left) and VAF (right) on cell composition. **C,**

Expression of selected genes in the NF-κB pathway from the scRNA-seq data. **D,**

GSEA of DEGs from myeloid populations. Color represents normalized effect score

(NES), and dot size represents adjusted *P*-values. **E,** GSEA plots of selected pathways

from **C**.

References

1. Jaiswal S, Fontanillas P, Flannick J, Manning A, Grauman PV, Mar BG, *et al.* Age-related clonal hematopoiesis associated with adverse outcomes. *N Engl J Med* 2014;**371**(26):2488-98 doi 10.1056/NEJMoa1408617.

2. Jaiswal S, Ebert BL. Clonal hematopoiesis in human aging and disease. *Science* 2019;**366**(6465):eaan4673 doi doi:10.1126/science.aan4673.

3. Hsu JI, Dayaram T, Tovy A, De Braekeleer E, Jeong M, Wang F, *et al.* PPM1D Mutations Drive Clonal Hematopoiesis in Response to Cytotoxic Chemotherapy. *Cell Stem Cell* 2018;**23**(5):700-13 e6 doi 10.1016/j.stem.2018.10.004.

4. Uddin MM, Zhou Y, Bick AG, Burugula BB, Jaiswal S, Desai P, *et al.* Longitudinal profiling of clonal hematopoiesis provides insight into clonal dynamics. *Immun Ageing* 2022;**19**(1):23 doi 10.1186/s12979-022-00278-9.

5. Welch JS, Ley TJ, Link DC, Miller CA, Larson DE, Koboldt DC, *et al.* The origin and evolution of mutations in acute myeloid leukemia. *Cell* 2012;**150**(2):264-78 doi 10.1016/j.cell.2012.06.023.

6. Muto T, Walker CS, Choi K, Hueneman K, Smith MA, Gul Z, *et al.* Adaptive response to inflammation contributes to sustained myelopoiesis and confers a competitive advantage in myelodysplastic syndrome HSCs. *Nat Immunol* 2020;**21**(5):535-45 doi 10.1038/s41590-020-0663-z.

7. Cai Z, Kotzin JJ, Ramdas B, Chen S, Nelanuthala S, Palam LR, *et al.* Inhibition of Inflammatory Signaling in Tet2 Mutant Preleukemic Cells Mitigates Stress-Induced Abnormalities and Clonal Hematopoiesis. *Cell Stem Cell* 2018;**23**(6):833-49 e5 doi 10.1016/j.stem.2018.10.013.

8. Abplanalp WT, Cremer S, John D, Hoffmann J, Schuhmacher B, Merten M, *et al.* Clonal Hematopoiesis-Driver DNMT3A Mutations Alter Immune Cells in Heart Failure. *Circ Res* 2021;**128**(2):216-28 doi 10.1161/CIRCRESAHA.120.317104.

9. Coombs CC, Zehir A, Devlin SM, Kishtagari A, Syed A, Jonsson P, *et al.* Therapy-Related Clonal Hematopoiesis in Patients with Non-hematologic Cancers Is Common and Associated with Adverse Clinical Outcomes. *Cell Stem Cell* 2017;**21**(3):374-82 e4 doi 10.1016/j.stem.2017.07.010.

10. Bolton KL, Ptashkin RN, Gao T, Braunstein L, Devlin SM, Kelly D, *et al.* Cancer therapy shapes the fitness landscape of clonal hematopoiesis. *Nat Genet* 2020;**52**(11):1219-26 doi 10.1038/s41588-020-00710-0.

11. Miller PG, Qiao D, Rojas-Quintero J, Honigberg MC, Sperling AS, Gibson CJ, *et al.* Association of clonal hematopoiesis with chronic obstructive pulmonary disease. *Blood* 2022;**139**(3):357-68 doi 10.1182/blood.2021013531.

12. Tobias DK, Manning AK, Wessel J, Raghavan S, Westerman KE, Bick AG, *et al.* Clonal Hematopoiesis of Indeterminate Potential (CHIP) and Incident Type 2 Diabetes Risk. *Diabetes Care* 2023;**46**(11):1978-85 doi 10.2337/dc23-0805.

13. Jaiswal S, Natarajan P, Silver AJ, Gibson CJ, Bick AG, Shvartz E, *et al.* Clonal Hematopoiesis and Risk of Atherosclerotic Cardiovascular Disease. *N Engl J Med* 2017;**377**(2):111-21 doi 10.1056/NEJMoa1701719.

14. Xie M, Lu C, Wang J, McLellan MD, Johnson KJ, Wendl MC, *et al.* Age-related mutations associated with clonal hematopoietic expansion and malignancies. *Nat Med* 2014;**20**(12):1472-8 doi 10.1038/nm.3733.

15. Stacey SN, Zink F, Halldorsson GH, Stefansdottir L, Gudjonsson SA, Einarsson G, *et al.* Genetics and epidemiology of mutational barcode-defined clonal hematopoiesis. *Nat Genet* 2023 doi 10.1038/s41588-023-01555-z.

16.   Leiter A, Veluswamy RR, Wisnivesky JP. The global burden of lung cancer: current status and
future trends. *Nat Rev Clin Oncol* 2023;**20**(9):624-39 doi 10.1038/s41571-023-00798-3.

17.   Hong W, Li A, Liu Y, Xiao X, Christiani DC, Hung RJ, *et al.* Clonal Hematopoiesis Mutations in
Patients with Lung Cancer Are Associated with Lung Cancer Risk Factors. *Cancer Research*
2022;**82**(2):199-209 doi 10.1158/0008-5472.Can-21-1903.

18.   Tian R, Wiley B, Liu J, Zong X, Truong B, Zhao S, *et al.* Clonal Hematopoiesis and Risk of Incident
Lung Cancer. *J Clin Oncol* 2023;**41**(7):1423-33 doi 10.1200/JCO.22.00857.

19.   Reck M, Rodriguez-Abreu D, Robinson AG, Hui R, Csoszi T, Fulop A, *et al.* Five-Year Outcomes
With Pembrolizumab Versus Chemotherapy for Metastatic Non-Small-Cell Lung Cancer With PD-
L1 Tumor Proportion Score >/= 50. *J Clin Oncol* 2021;**39**(21):2339-49 doi 10.1200/JCO.21.00174.

20.   Miller PG, Sperling AS, Brea EJ, Leick MB, Fell GG, Jan M, *et al.* Clonal hematopoiesis in patients
receiving chimeric antigen receptor T-cell therapy. *Blood Adv* 2021;**5**(15):2982-6 doi
10.1182/bloodadvances.2021004554.

21.   Miller PG, Gibson CJ, Mehta A, Sperling AS, Frederick DT, Manos MP, *et al.* Fitness Landscape of
Clonal Hematopoiesis Under Selective Pressure of Immune Checkpoint Blockade. *JCO Precis*
Oncol 2020;**4** doi 10.1200/PO.20.00186.

22.   Kim H, Park S, Han KY, Lee N, Kim H, Jung HA, *et al.* Clonal expansion of resident memory T cells
in peripheral blood of patients with non-small cell lung cancer during immune checkpoint
inhibitor treatment. *J Immunother Cancer* 2023;**11**(2) doi 10.1136/jitc-2022-005509.

23.   Litchfield K, Reading JL, Puttick C, Thakkar K, Abbosh C, Bentham R, *et al.* Meta-analysis of
tumor- and T cell-intrinsic mechanisms of sensitization to checkpoint inhibition. *Cell*
2021;**184**(3):596-614.e14 doi https://doi.org/10.1016/j.cell.2021.01.002.

24.   Garcia M, Juhos S, Larsson M, Olason PI, Martin M, Eisfeldt J, *et al.* Sarek: A portable workflow
for whole-genome sequencing analysis of germline and somatic variants. *F1000Res* 2020;**9**:63
doi 10.12688/f1000research.16665.2.

25.   McKenna A, Hanna M, Banks E, Sivachenko A, Cibulskis K, Kernytsky A, *et al.* The Genome
Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data.
Genome Res 2010;**20**(9):1297-303 doi 10.1101/gr.107524.110.

26.   Cingolani P, Platts A, Wang le L, Coon M, Nguyen T, Wang L, *et al.* A program for annotating and
predicting the effects of single nucleotide polymorphisms, SnpEff: SNPs in the genome of
Drosophila melanogaster strain w1118; iso-2; iso-3. *Fly (Austin)* 2012;**6**(2):80-92 doi
10.4161/fly.19695.

27.   McLaren W, Gil L, Hunt SE, Riat HS, Ritchie GR, Thormann A, *et al.* The Ensembl Variant Effect
Predictor. *Genome Biol* 2016;**17**(1):122 doi 10.1186/s13059-016-0974-4.

28.   Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, *et al.* COSMIC: the Catalogue Of
Somatic Mutations In Cancer. *Nucleic Acids Res* 2019;**47**(D1):D941-D7 doi 10.1093/nar/gky1015.

29.   Robinson JT, Thorvaldsdottir H, Wenger AM, Zehir A, Mesirov JP. Variant Review with the
Integrative Genomics Viewer. *Cancer Res* 2017;**77**(21):e31-e4 doi 10.1158/0008-5472.CAN-17-
0337.

30.   Niroula A, Sekar A, Murakami MA, Trinder M, Agrawal M, Wong WJ, *et al.* Distinction of
lymphoid and myeloid clonal hematopoiesis. *Nat Med* 2021;**27**(11):1921-7 doi 10.1038/s41591-
021-01521-4.

31.   Fleming SJ, Chaffin MD, Arduini A, Akkad AD, Banks E, Marioni JC, *et al.* Unsupervised removal of
systematic background noise from droplet-based single-cell experiments using CellBender. *Nat*
Methods 2023;**20**(9):1323-35 doi 10.1038/s41592-023-01943-7.

32.   Kang HM, Subramaniam M, Targ S, Nguyen M, Maliskova L, McCarthy E, *et al.* Multiplexed
531   droplet single-cell RNA-sequencing using natural genetic variation. *Nat Biotechnol*
532   2018;**36**(1):89-94 doi 10.1038/nbt.4042.

533   33.   Hao Y, Hao S, Andersen-Nissen E, Mauck WM, 3rd, Zheng S, Butler A, *et al.* Integrated analysis of
534   multimodal single-cell data. *Cell* 2021;**184**(13):3573-87 e29 doi 10.1016/j.cell.2021.04.048.

535   34.   Hafemeister C, Satija R. Normalization and variance stabilization of single-cell RNA-seq data
536   using regularized negative binomial regression. *Genome Biol* 2019;**20**(1):296 doi
537   10.1186/s13059-019-1874-1.

538   35.   Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, *et al.* Fast, sensitive and accurate
539   integration of single-cell data with Harmony. *Nat Methods* 2019;**16**(12):1289-96 doi
540   10.1038/s41592-019-0619-0.

541   36.   Liberzon A, Birger C, Thorvaldsdottir H, Ghandi M, Mesirov JP, Tamayo P. The Molecular
542   Signatures Database (MSigDB) hallmark gene set collection. *Cell Syst* 2015;**1**(6):417-25 doi
543   10.1016/j.cels.2015.12.004.

544   37.   Sergushichev AA. An algorithm for fast preranked gene set enrichment analysis using cumulative
545   statistic calculation. *bioRxiv* 2016:060012 doi 10.1101/060012.

546   38.   Morabito S, Reese F, Rahimzadeh N, Miyoshi E, Swarup V. hdWGCNA identifies co-expression
547   networks in high-dimensional transcriptomics data. *Cell Rep Methods* 2023;**3**(6):100498 doi
548   10.1016/j.crmeth.2023.100498.

549   39.   Van de Sande B, Flerin C, Davie K, De Waegeneer M, Hulselmans G, Aibar S, *et al.* A scalable
550   SCENIC workflow for single-cell gene regulatory network analysis. *Nat Protoc* 2020;**15**(7):2247-
551   76 doi 10.1038/s41596-020-0336-2.

552   40.   Bravo Gonzalez-Blas C, De Winter S, Hulselmans G, Hecker N, Matetovici I, Christiaens V, *et al.*
553   SCENIC+: single-cell multiomic inference of enhancers and gene regulatory networks. *Nat*
554   *Methods* 2023;**20**(9):1355-67 doi 10.1038/s41592-023-01938-4.

555   41.   Persad S, Choo ZN, Dien C, Sohail N, Masilionis I, Chaligne R, *et al.* SEACells infers transcriptional
556   and epigenomic cellular states from single-cell genomics data. *Nat Biotechnol* 2023 doi
557   10.1038/s41587-023-01716-9.

558   42.   Efremova M, Vento-Tormo M, Teichmann SA, Vento-Tormo R. CellPhoneDB: inferring cell-cell
559   communication from combined expression of multi-subunit ligand-receptor complexes. *Nat*
560   *Protoc* 2020;**15**(4):1484-506 doi 10.1038/s41596-020-0292-x.

561   43.   Kessler MD, Damask A, O'Keeffe S, Banerjee N, Li D, Watanabe K, *et al.* Common and rare
562   variant associations with clonal haematopoiesis phenotypes. *Nature* 2022;**612**(7939):301-9 doi
563   10.1038/s41586-022-05448-9.

564   44.   Sano S, Oshima K, Wang Y, MacLauchlan S, Katanasaka Y, Sano M, *et al.* Tet2-Mediated Clonal
565   Hematopoiesis Accelerates Heart Failure Through a Mechanism Involving the IL-1beta/NLRP3
566   Inflammasome. *J Am Coll Cardiol* 2018;**71**(8):875-86 doi 10.1016/j.jacc.2017.12.037.

567   45.   Nam AS, Dusaj N, Izzo F, Murali R, Myers RM, Mouhieddine TH, *et al.* Single-cell multi-omics of
568   human clonal hematopoiesis reveals that DNMT3A R882 mutations perturb early progenitor
569   states through selective hypomethylation. *Nat Genet* 2022;**54**(10):1514-26 doi 10.1038/s41588-
570   022-01179-9.

571   46.   Fuster JJ, MacLauchlan S, Zuriaga MA, Polackal MN, Ostriker AC, Chakraborty R, *et al.* Clonal
572   hematopoiesis associated with TET2 deficiency accelerates atherosclerosis development in mice.
573   *Science* 2017;**355**(6327):842-7 doi 10.1126/science.aag1381.

574   47.   Choi B, Kang CK, Park S, Lee D, Lee AJ, Ko Y, *et al.* Single-cell transcriptome analyses reveal
575   distinct gene expression signatures of severe COVID-19 in the presence of clonal hematopoiesis.
576   *Exp Mol Med* 2022;**54**(10):1756-65 doi 10.1038/s12276-022-00866-1.

48.   Li R, Ong SL, Tran LM, Jing Z, Liu B, Park SJ, *et al.* Chronic IL-1beta-induced inflammation
regulates epithelial-to-mesenchymal transition memory phenotypes via epigenetic
modifications in non-small cell lung cancer. *Sci Rep* 2020;**10**(1):377 doi 10.1038/s41598-019-
57285-y.
49.   Zhang J, Veeramachaneni N. Targeting interleukin-1beta and inflammation in lung cancer.
Biomark Res 2022;**10**(1):5 doi 10.1186/s40364-021-00341-5.
50.   Ridker PM, MacFadyen JG, Thuren T, Everett BM, Libby P, Glynn RJ, *et al.* Effect of interleukin-
1beta inhibition with canakinumab on incident lung cancer in patients with atherosclerosis:
exploratory results from a randomised, double-blind, placebo-controlled trial. *Lancet*
2017;**390**(10105):1833-42 doi 10.1016/S0140-6736(17)32247-X.
51.   Hong W, Li A, Liu Y, Xiao X, Christiani DC, Hung RJ, *et al.* Clonal Hematopoiesis Mutations in
Patients with Lung Cancer Are Associated with Lung Cancer Risk Factors. *Cancer Res*
2022;**82**(2):199-209 doi 10.1158/0008-5472.CAN-21-1903.
52.   Levin MG, Nakao T, Zekavat SM, Koyama S, Bick AG, Niroula A, *et al.* Genetics of smoking and
risk of clonal hematopoiesis. *Sci Rep* 2022;**12**(1):7248 doi 10.1038/s41598-022-09604-z.