Wing shape evolution is not constrained by ancestral genetic covariances in the invasive *Drosophila suzukii*

Antoine Fraimout[1†], Stéphane Chantepie[1], Nicolas Navarro[2], Céline Teplitsky[3] and Vincent Debat[1]

Short title: Quantitative genetics of *Drosophila suzukii* wing shape

[1]Institut de Systématique, Evolution, Biodiversité, ISYEB – CNRS, MNHN, UPMC, EPHE, Université des Antilles, Muséum National d'Histoire Naturelle, Sorbonne Universités, 57 rue Cuvier, 75005 Paris, France
[2]UMR CNRS 6282 Biogéosciences, Université de Bourgogne-Franche-Comté, 21000 Dijon, France
[3]Centre d'Ecologie Fonctionnelle et Evolutive, CEFE – UMR 5175 – CNRS, IRD, EPHE, Université de Montpellier, Université Paul Valéry, 34293 Montpellier, France

[†]Corresponding author, E-mail: fraimout.antoine@gmail.com

Abstract

The extent to which phenotypic evolution can be constrained by genetic correlations is an important question in evolutionary biology. To address this question, biological invasions are opportune models where derived, invasive populations can be compared to their extant ancestors, allowing to track the evolution of genetic correlations from the ancestor, throughout the invasion process. In this paper, we focused on the worldwide invasion of *Drosophila suzukii* (Matsumara, 1931)*,* and investigated the evolution of the genetic covariance matrix **G** of wing shape between ancestral native, and derived invasive populations. Leveraging demographic history resolved by population genetics approaches, we tested whether **G** remained stable during the invasion. Using a multivariate Q_{ST}-F_{ST} approach, we further tested whether or not the observed phenotypic divergence in wing shape aligned with a neutral scenario of evolution. Our results show moderate yet significant quantitative genetic differentiation of wing shape among *D. suzukii* populations and a relative stability in the structure of **G**, presenting a roughly spherical shape but slightly different volumes. These characteristics likely reflect the demographic history of populations and suggest a low level of genetic constraint on wing shape evolution. The divergence between populations was greater than expected under a purely neutral model of evolution, compatible with an effect of divergent selection among them. Overall, our study suggests that selection and drift, but not ancestral genetic constraints, affected the early stages of wing shape evolution during *D. suzukii* invasion.

Key words: *Drosophila suzukii*, **G**-matrix, geometric morphometrics, biological invasion, quantitative genetics, Q_{ST}-F_{ST}

Introduction

Biological invasions often involve stochastic demographic changes through founder effects and genetic bottlenecks (Golani *et al.*, 2007) or the shuffling of different alleles by genetic admixture between populations (Rius & Darling, 2014). Invasive populations can also be subjected to multiple and often novel selective pressures in the new environment and throughout the invasion process (Kilkenny & Galloway, 2013). Consequently, they tend to show marked genetic and phenotypic differences compared to their source populations, and distinguishing whether this differentiation results from adaptive or neutral evolution is challenging (Lee, 2002; Wares, 2005; Facon *et al.*, 2006; Hayes & Barry, 2008; Keller & Taylor, 2008; Prentis *et al.*, 2008; Bock *et al.*, 2015; Dlugosch *et al.*, 2015; Estoup *et al.*, 2016). Furthermore, these evolutionary changes can happen over relatively short time scales (Colautti & Lau, 2015). Hence, a topical question in invasion biology is how fast can invasive populations evolve in their new environment? Can we detect adaptive evolution occurring at the onset of the invasion (Szűcs *et al.*, 2017)?

In a new environment, the evolution of a population's mean phenotype towards a new fitness optimum (*i.e.* adaptation) is conditioned by the genetic variances and covariances of the traits under selection, summarized in **G**, the genetic variance-covariance matrix (Lande, 1976). For a set of traits measured in a given population, the **G** matrix informs about genetic variation for each individual trait (diagonal elements) and for the genetic covariation between traits (off-diagonal elements). Statistical analysis of **G** through an eigen decomposition can then identify the major axis of variation in the population, corresponding to the combination of traits with the most genetic variation, also known as **g**max (Schluter 1996). Identifying the axes of genetic covariance between traits is particularly important as theory predicts that they will influence the evolutionary response: adaptation will be facilitated if selection is aligned with **g**max, or constrained if it is in a direction where there is little, or no genetic variation for the combination of traits under selection (Arnold *et al.*, 2001, Agrawal & Stinchcombe, 2009; Fischer *et al.*, 2016). This is formalized in the multivariate breeder's equation (Lande, 1979):

$$R = G\beta$$

where the response to selection R - the change in average phenotype of the population in one generation - equals the product of **G** and the directional selection gradient β.

It has been proposed in the context of evolutionary diversification, that the **G** matrix of an ancestral population can constrain derived populations to diverge along certain directions of the phenotypic space (*i.e.* the lines of least evolutionary resistance; Schluter, 1996). For instance, early stages of adaptive phenotypic divergence could be constrained by the direction of the major axis of the ancestral **G** matrix (*i.e.* ancestral **g**max; Schluter, 1996; McGuigan, 2006; Walter *et al.*, 2018) and result in an alignment between **g**max and the leading vector of divergence of the among-populations divergence matrix (**D**). On the other hand, under a neutral model of phenotypic evolution, divergence among populations is also expected to take place along that same axis of greatest genetic variation (Lande 1979) and result in proportional **D** and **G** matrices (Chenoweth *et al.* 2010). In other words, a concordance between the major axes of **D** and **G** could be both suggestive of neutral evolution (Arnold *et al.* 2001; Bégin & Roff, 2003) or indicative of selection.

Furthermore, valid predictions of evolutionary trajectories depend on whether or not **G** remains stable across space and time (Jones *et al.*, 2004; Arnold *et al.*, 2008). The stability of **G** has received much attention (*e.g.* Roff, 2000; Steppan *et al.*, 2002; Bégin & Roff, 2003; Mezey & Houle, 2003; Björklund, 2004; Blows & Hoffmann, 2005; Arnold *et al.*, 2008; Walsh & Blows, 2009) and its sensitivity to selection, migration and drift has been investigated both theoretically (*e.g.* Jones *et al.*, 2004; Guillaume & Whitlock, 2007, Chantepie & Chevin, 2020), and empirically under natural and laboratory conditions (Merilä *et al.*, 1994; Phillips *et al.*, 2001; Conner *et al.*, 2003; Cano *et al.*, 2004; Eroukhmanoff. & Svensson, 2011; Björklund *et al.*, 2013; McGlothlin *et al.*, 2018; Walter 2023; Henry & Stinchcombe 2023; Chantepie *et al.* 2024) but no clear consensus has emerged. For instance, the recent work of Henry & Stinchcombe (2023) has shown that the **G** matrix of floral traits in the ivy (*Ipomoea hederacea*) remained stable across a large latitudinal gradient in continental USA, despite evidence of phenotypic divergence among populations. Conversely, Mallard *et al.* (2023) have shown using a large experimental evolution study that the orientation of the **G** matrix of locomotor traits in *C. elegans* can be altered by genetic drift, despite phenotypic stasis. Multiple additional lines of evidence suggest that **G** can be reshaped by selection (Jones *et al.*, 2003; Jones *et al.*, 2004; Dugand *et al.* 2021), drift (Phillips *et al.*, 2001) and environmental conditions (*e.g.* Sikkink *et al.*, 2015; Wood & Brodie, 2015) even over the span of a single generation (Siren *et al.*, 2017), but, there is also recent evidence for stable

G matrices across populations and continents (*e.g.*, Puentes *et al.*, 2016; Delahaie *et al.*, 2017, Henry & Stinchcombe 2023). Overall clear-cut predictions on the interplay between **G** and phenotypic divergence in response to selection or drift are difficult to formulate, particularly at a contemporary timescale. Therefore, the stability of **G** needs to be further explored empirically along with clear testing of the neutral expectation for among-population divergence against observed data (Merilä & Björklund 2004).

Here, we focus on the worldwide invasion of the spotted wing drosophila, *Drosophila suzukii* (Matsumura, 1931) to assess the stability of **G** between native and invasive populations across continents. *Drosophila suzukii* is a highly invasive pest of agricultural crops in Europe and North America where it was first detected almost simultaneously in 2008 (see Asplen *et al.*, 2015 for review). We reconstructed the detailed history of this worldwide invasion and showed the presence of multiple independent introductions from Asia involving population bottlenecks and genetic admixture events within invaded regions (Fraimout *et al.*, 2017). A consequence of this demographic history is that native and invasive populations show significant levels of neutral genetic differentiation as measured by F_{ST} estimated using microsatellite loci (Fraimout *et al.*, 2015; Fraimout *et al.*, 2017). At the phenotypic level, we showed using laboratory lines derived from samples of natural populations that native (Japanese) and invasive (French and North-American) *D. suzukii* display genetically based differences in wing shape and size (Fraimout *et al.*, 2018). However, whether this phenotypic divergence is pervasive across replicated populations, and whether it is associated with changes in the structure of the underlying **G** matrix is unknown. This invasion therefore provides an opportune context to address long-standing questions about the evolution of multivariate phenotypes and the structure of genetic variances and covariances in colonizing populations. Particularly the presence of multiple invasive populations within Europe and USA allows assessing the magnitude and direction of phenotypic evolution from the native ancestral populations. In other words, it is possible to estimate how much the mean phenotype of each invasive population has diverged from the ancestral (*i.e.* native) population and whether phenotypic evolution repeatedly occurred in the same direction or at random, in all directions of the phenotypic space. Given the availability of extant ancestral populations, this in turns allows to determine whether phenotypic divergence among invasive populations has been constrained by ancestral genetic covariances. Furthermore, comparing populations with known history could help identify the effects of demographic processes on the structure of

the **G** matrix. For instance, we expect that periods of bottlenecks or population admixture – all experienced by

invasive *D. suzukii* populations to a different extent (Fraimout *et al.*, 2017, Olazcuaga *et al.* 2020) – may have

altered the structure of **G** in invasive populations. Specifically, depletion of genetic variance due to bottlenecks

could result in a **G** matrix of smaller volume. Alternatively, new combinations of alleles produced by genetic

admixture, could increase the volume or **G**, or alter its shape, by changing the distribution of genetic variance

over the first axes of variation. Our main objective was thus to investigate whether or not **G** remained stable

throughout the invasion history of *D. suzukii*. Additionally, access to both molecular and quantitative measures

of genetic variation allows testing for a signal of selection in the observed phenotypic divergence. By

comparing F_{ST} calculated from molecular markers to its phenotypic equivalent Q_{ST}, it is possible to determine

whether phenotypic divergence exceeds or is equal to the divergence expected under genetic drift alone (Spitze

1993, Whitlock 2008). While the former ($Q_{ST} > F_{ST}$) would be indicative of divergent selection acting on the

study traits, the latter ($Q_{ST} = F_{ST}$) would suggest neutral phenotypic evolution.

We used an isofemale line breeding design (Hoffmann & Parsons, 1988; David *et al.*, 2005; Berger *et al.*,

2013) and a geometric morphometric approach (*e.g.* Klingenberg 2003) to estimate the quantitative genetic

divergence in wing shape between six natural populations of *D. suzukii* sampled in the native area (Japan) and

the two main invaded areas in Europe and the USA. We specifically addressed the following questions: i) how

did wing shape evolve across multiple invasive populations? ii) did the **G**-matrix remain stable during the

invasion**?** iii) has phenotypic divergence among invasive populations been constrained by ancestral genetic

covariances and iv) can it be attributed to selection acting on wing shape?

Material and methods

Samples, lines and data acquisition

Drosophila suzukii adults were sampled in 2014 using banana bait traps and net swiping at six localities

representative of their global distribution range in Japan (native range), USA and France (invasive range) (Fig.

1). Within each country, we sampled two localities representative of a northern and a southern latitude.

Repeated clinal evolution of wing morphology has been described in the related invasive species *Drosophila*

subobscura and proposed as an adaptation to different temperature optima at northern and southern latitudes

(Huey *et al.*, 2000; Gilchrist *et al.*, 2001; Gilchrist *et al.*, 2004). Thus, we hypothesized that sampling populations close to the extremes of their latitudinal distribution would maximize our chance to capture possibly adaptive phenotypic divergence in the invaded range. In USA and France, these localities were also representative of the invasion history, with the southern localities corresponding to the first reported occurrences of *D. suzukii* (Watsonville, USA and Montpellier, France; Hauser, 2011; Calabria *et al.*, 2012) and the northern localities representing the expansion front of the species in the area at the time of sampling (Dayton, Oregon, USA and Paris, France). It is worth stressing that although the most likely genetic source of Europe and USA invasions was identified as continental Asia (*i.e.* China, Fraimout *et al.*, 2017), *F* statistics and Bayesian clustering methods indicate weak genetic differentiation between these sources and our Japanese population samples, thereby suggesting that the use of Japanese populations as ancestral representatives is an acceptable approximation. For the Japanese flies, a northern population was sampled in Sapporo (Hokkaido Island) and a southern population was sampled in Tokyo (Honshu Island). In both Japanese localities, traps were placed in gardens inside the university campuses. In the invasive North American range, sampling habitats corresponded to vineyards in Dayton, and strawberry fields in Watsonville. French samples were collected in the botanical garden of the Natural History Museum in Paris and close to vineyards in Montpellier. 30 isofemale lines on average (see Table S1 for details and population codes) were established for each locality by placing randomly chosen wild-caught gravid females in separate rearing vials, and splitting F_1 offsprings in five replicate vials. Following the isofemale approach, we further expanded each line by letting individuals mate freely within tubes (*i.e.*, full-sib mating) and transferring their progeny to new rearing vials each generation. We repeated this for five generations and in the fifth generation, a single female per line was isolated in a new rearing tube to produce the focal offspring used in the study. At the end of the experiment, 5 to 10 full-sib males were randomly chosen from each line from each population for phenotyping. Rearing medium consisted of cornstarch and yeast with added anti-biotic (hydroxy-4 benzoate). Lines were randomly positioned in incubators and kept for the five generations at 22°C. A total of 1323 individuals distributed across 177 isofemale lines were phenotyped. Wings were carefully clipped from over-anesthetized flies using tweezers and mounted on glass slides in a mixture of ethanol and glycerin. Coverslips were sealed using nail polish and maintained with small weights to flatten the wing as much as possible. Images of the wings were acquired with a Leica DFC 420 digital camera mounted on a Leica Z6 APO microscope. Fifteen landmarks

were defined on the dorsal side of the right wing and digitized using a custom *ImageJ* plugin written by C. P.

Klingenberg (pers. comm.).

Wing shape differentiation – Geometric morphometric analyses

We focused our study on the analysis of wing shape variation among *D. suzukii* populations. Shape data have

been used in multiple quantitative genetics studies (Klingenberg & Leamy, 2001; Klingenberg, 2003; Monteiro

et al., 2003; Hansen & Houle 2004; Mezey & Houle, 2005; Myers *et al.*, 2006; Polly, 2008; Klingenberg,

2010; Adams, 2011; Leinonen *et al.*, 2011; Martínez-Abadías *et al.*, 2012) and are suitable for the estimation

of **G**, and relevant for the study of multivariate phenotypic evolution. We followed a standard geometric

morphometrics protocol (Bookstein 1996) to characterize wing shape: first, Procrustes generalized least

squares superimposition was used to extract shape information from the landmarks coordinates (GLS, *e.g.*

Dryden & Mardia, 1993). Briefly, this method allows to scale, rotate and translate the configuration of

landmarks to obtain shape information that is free of orientation or size differences. Second, a principal

component analysis (PCA) was applied to the set of 30 coordinates (15 landmarks in 2 dimensions x and y)

obtained after superimposition. From the PCA, 26 non-null components were conserved as input data

(corresponding to the 30 coordinates minus 4 degrees of freedom lost through the scaling, rotation and

translations (x and y)). We estimated the centroid size of the wing as the square root of the sum of the squared

distances of all landmarks from their centroid, and used this centroid size as a size variable in the subsequent

analyses. As divergence in wing shape first appeared to be weak on the PCA plot, we further performed a

between-group principal component analysis (*bgPCA*) to visualize the overall variation in wing shape among

populations (*e.g.* Mitteroecker & Bookstein, 2011). We then estimated the statistical significance of the shape

divergence among populations by applying a multivariate analysis of variance (MANOVA) using the 26 PCs

as response variables, and population of origin as a fixed effect. We further estimated the pairwise differences

among populations with permutational pairwise MANOVAs, using the *RVAideMemoire* R package (v.0.9-55,

Hervé, 2016). We used the Wilk's lambda statistics and set the number of permutations to 10000 while

correcting for multiple testing using the *fdr* method (Benjamini & Hochberg 1995). Finally, to account for any

allometric effect (i.e. a change in shape associated with a change in size,e.g. Klingenberg 2010), we also fitted

the MANOVAs using the residuals of a size/shape multivariate regression, corresponding to the allometry-free

components of wing shape variation.

Wing shape differentiation – phenotypic divergence vector analysis

To estimate the divergence in wing shape among populations of *D. suzukii*, particularly the magnitude and

direction of phenotypic evolution in the derived invasive populations from their Asian ancestors, we applied a

geometric approach to phenotypic evolution (*e.g.,* Stuart *et al.* 2017, Fraimout *et al.* 2022), by calculating

vectors of phenotypic divergence among populations. The phenotypic mean of each population was first

estimated from the MANOVA (see section above) using the 26 non-null PC representative of wing shape, and

accounting for differences in wing size by adding the wing centroid size as covariate. Then, we estimated the

differences in mean wing shape between each invasive population and the ancestral wing shape, assessed as

the average of the two Japanese populations (Fig. 2). We also estimated divergence at the country level, using

the three countries average wing shapes. In total, we obtained six phenotypic divergence vectors describing

the difference between the ancestral Japanese and each invasive population (4 vectors) and between the

ancestor and each country's mean wing shape (2 vectors). To test whether phenotypic divergence occurred

along similar or different directions, we assessed the similarity between these vectors of divergence by

comparing their angle to the distribution of angles between 10,000 pairs of random vectors of the same

dimensionality (Klingenberg and McIntyre, 1998). Angles between vectors were calculated as:

$$\alpha_1 = \cos^{-1}(\Delta z_{Di}, \Delta z^{T}_{Dj}) \ [1]$$

where z_D corresponds to a divergence vector of population *i* or *j* from the ancestor. An eigenanalysis of the

matrix containing the four population vectors was applied to synthesize the main axes of divergences.

*Matrices estimations – population-specific **G** matrices*

We estimated the six **G** matrices corresponding to the six sampled populations under a Bayesian framework

using the R-package *MCMCglmm* (Hadfield, 2010; Fig. 3). To this end, we partitioned the phenotypic variance

in wing shape for each population using a multivariate animal model (Kruuk, 2004) of the form:

$$y = \sum \quad X\beta + \sum \quad Z\gamma + \varepsilon \ [2]$$

where *y* is the multivariate vector of phenotypic values for wing shape corresponding to the 26 non-null

principal components (PCs) accounting for the totality of shape variance; β is the vector of fixed effect; γ is

the vector of random effects; ε is the vector of residual errors and X and Z the design matrices relating to the

fixed and random effects, respectively. We estimated **G** matrices separately for each population using model

[2] and appended the matrix of relatedness between individuals to the vector of random effects with the

pedigree option implemented in the *MCMCglmm* function. The pedigree corresponding to our crossing design

was coded in R using custom scripts (see *Data availability* statement). In all models, wing centroid size was

fitted as a fixed effect. We used a prior set with variance estimates following a 'broken-stick' model of decrease

of variance, appropriate for the use of PC scores representing decreasing proportion of explained variance

(Jolliffe, 2002), with null covariances and a degree of belief $nu = n + 0.002$ (where n is the number of traits,

Houle & Meyer, 2015). We ran each model for 1 200 000 iterations with a burnin period of 200 000 and a

thinning rate of 1000, resulting in 1000 Monte Carlo Markov Chain (MCMC) posterior samples. The

convergence of models was checked visually from plots of MCMC chains and posterior distributions as well

as autocorrelation level (<0.05 in all models). All analyses were performed in R (v.4.2.2; R Core Team, 2022).

*Test of **G**-matrix stability – Differences in the size and shape of **G***

One of our main objectives was to assess the stability of the **G** matrix in the face of the demographic changes

linked to the invasion. According to previous results (Fraimout *et al.* 2017) all invasive *D. suzukii* populations

have experienced genetic bottlenecks, and some populations show evidence of genetic admixture from

differentiated sources. Our hypothesis was that the variation in total genetic variance due to these demographic

events - namely, reduction of genetic variance by bottlenecks or increased genetic variance from admixture -

should be reflected in the structure of **G**. To test this hypothesis, we estimated three parameters pertaining to

the structure of **G** and performed pairwise comparisons among all six studied populations. Matrix size (*i.e.*,

volume), estimating the overall amount of genetic variance for wing shape in the population, was calculated

as the sum of the matrix eigenvalues. Matrix shape, estimating whether some directions of wing shape variation

present more genetic variance than others, was assessed by the matrix eccentricity (*i.e.* the ratio of the first to

the second eigenvalue; Jones *et al.*, 2003). Matrix shape was also estimated by the proportion of variance along

$\mathbf{g}_{max}$. Significance of metric differences between populations was assessed by estimating the ratio between each

population metrics and examining the overlap between the 95% highest posterior density (HPD) of these ratios

calculated from the MCMC posterior samples, with zero.

*Test of **G**-matrix stability – Genetic covariance tensor*

We further tested for differences among **G** matrices by following the approach of Hine *et al.* (2009) and

Aguirre *et al.* (2014) using a fourth-order genetic covariance tensor analysis (hereafter 'covariance tensors').

Briefly (see details in Hine *et al.* 2009; Aguirre *et al.* 2014), variances and covariances among all elements of

the original set of **G** matrices are calculated to construct the fourth-order covariance tensor Σ and the second-

order (*i.e.* matrix) tensor **S**. Eigenanalysis of the **S**-matrix allows for the construction of eigentensors which

represent independent vectors of differences in variance and covariance among **G** matrices (Fig. 4). Covariance

tensors provide a way to compare more than two genetic covariance matrices and describe independent axes

of divergence among them. In other words, covariance tensors allow identifying trait combinations explaining

most genetic variation among **G** matrices. We applied the covariance tensor analysis to our set of six **G**

matrices using the full 1000 MCMC samples for each matrix and compared the observed divergence to the

null expectation obtained from the analyses of random **G** matrices. We used an approach based on Morrissey

et al. (2019) to simulate the null distribution of **G** matrices. For each population, the breeding values of

founders were drawn from a multivariate distribution with zero mean and a variance-covariance matrix equal

to the **G**-matrix estimated from this population with the animal model [2]. Breeding values were then randomly

shuffled among the founders and the breeding values of all remaining individuals were estimated based on the

pedigree of each population. As our breeding design included five generations of inbreeding, we modeled a

pedigree to better approximate the actual relatedness of our measured individuals. The breeding values of an

individual were drawn from a multivariate normal distribution with a vector of means equal to the mean

breeding values of its parents and a deviation equal to the segregation variance (Walsh & Lynch, 2018). For

an individual, the segregation variance was estimated as half the product of the randomized **G** matrix of their

population (estimated on the new population's founders) and one minus the average inbreeding coefficient of

its parents (Walsh & Lynch, 2018, pp.557). The phenotypes were then recomposed by adding the random and

fixed effects estimated by the animal model following [2].

We repeated this sequence for each posterior sample of animal models and resulted in 1000 null datasets for

each population. The **G** matrix of null datasets was estimated following eq. [2]. To reduce computational burden, we sampled 200 posterior samples for each null animal model (Morrissey *et al.*, 2019) to compute a mean null distribution **G** matrix for each population that accounts for both the sampling effect and the uncertainty in the **G** matrix estimates. Each final null **G** matrix consisted of the average of the 1000 MCMC posterior for each of the 100 replicates and were used to estimate random genetic covariance tensor as described in Aguirre *et al.* (2014) using custom scripts in R (see *Data availability* section for source code),

*Test of genetic constraints: proportionality between **G** and **D***

In the case of populations deriving from a common ancestor, quantitative genetics theory predicts that genetic constraints imposed by the ancestral **G**-matrix can influence among-population divergence in a manner, leading to the proportionality of the divergence matrix **D** and the ancestral **G** under neutral or adaptive scenarios (Lande, 1979; Merilä & Björklund 1994, 2004). This proportionality induced by genetic constraints is expected under selection, where the evolutionary response is biased along the ancestral main directions of genetic variation (e.g. Schluter 1996, McGlothlin et al 2018), but also under neutral evolution, as genetic drift is expected to induce a global reduction in genetic variances, but no predictable change in the orientation of the divergence, leaving on average **G** and **D** proportional (Lande 1979; Phillips *et al* 2001).

The similarity between **D** and **G** was first investigated by computing the angle between a divergence vector **d** and the ancestral **G** matrix following Krzanowski (1979):

$$\theta = cos^{-1}\{(dLL^t d^t)^{-0.5}\}$$

where **L** is the set of first q eigenvectors of **G**, where q was set equal to half the shape space dimensionality ($q = (2k - 4)/2 = 13$; and k is the number of digitized landmarks) and **d** is a divergence vector..

We compared the observed angles to their random expectations by simulating 1000 random vectors v_{2k} projected in the tangent space according to

$$d = (I_{2k} - \mu\mu^t)v$$

where **μ** is the vectorised mean shape (Dryden and Mardia, 1998).

Test of selection: multivariate Q_{ST}-F_{ST} comparison

A proportionality between **G** and **D** is indicative of an effect of genetic constraints on phenotypic evolution,

but is insufficient to identify the evolutionary process at play, pointing at the need for a more stringent test to

decipher the adaptive and neutral scenarios (Martin *et al*. 2008). To test the hypothesis that phenotypic

evolution among invasive *D. suzukii* populations resulted from divergent selection on wing shape, we used the

two steps method proposed by Martin *et al*. (2008). Briefly, this method is a multivariate extension of the

classical Q_{ST}-F_{ST}, testing whether the observed phenotypic divergence is greater ($Q_{ST} > F_{ST}$), equal to ($Q_{ST} =$

F_{ST}), or smaller ($Q_{ST} < F_{ST}$) than expected under drift alone (see Martin *et al*. 2008 and Chapuis *et al*. 2008 for

more details). First, proportionality between **D** and **G** was tested using the MANOVA estimates of the

covariance matrices. Second, the coefficient of proportionality ρ_{ST} between the ancestral **G** matrix and the

divergence matrix **D** was estimated and compared to the predicted neutral divergence $2F_{ST}/(1-F_{ST})$ obtained

from neutral molecular markers (Rogers & Harpending 1983; Martin *et al*. 2008). While ρ_{ST} is not strictly

equivalent to Q_{ST}, the rationale behind its comparison to a measure of neutral differentiation (*i.e.* F_{ST}) is very

similar. ρ_{ST} is expressed as $1/n_f(\rho_{MS} - 1)$ (equation 11 in Martin *et al*. 2018; pp. 2139), where n_f is the number

of families sampled, and ρ_{MS} is the coefficient of proportionality between the among- and within-population

mean square matrices, estimated from phenotypic data. Comparing the confidence intervals of ρ_{ST} to that of

$2F_{ST}/(1-F_{ST})$ estimated from molecular markers provides a test of neutrality similar in rationale to the Q_{ST}-F_{ST}

comparison: in case of an overlap between the CIs, the neutrality hypothesis is not rejected, this is similar to

observing $Q_{ST} \approx F_{ST}$. Conversely, deviation from this overlap would be indicative of selection. To test for

proportionality between **D** and **G**, we used the Bartlett-adjusted test of proportionality between matrices as

recommended in Martin *et al*. (2008). We used the *k.prop* R function of the neutrality package from Martin *et*

al. (2008) to calculate ρ_{ST} and the *hierfstat* R package (v.0.5.11; Goudet & Jombart 2015) to get bootstrapped

estimates of F_{ST} from 25 neutral microsatellite loci (Fraimout *et al*. 2015, 2017). Inputs for both tests were

obtained by performing a MANOVA on the wing shape data and using the log-centroid size as covariate and

population of origin as fixed effect. Also following recommendation from Martin *et al*. (2008), we used the

first 11 PCs describing wing shape variation to maximize accuracy of the statistical tests so that $p < 2nb$ (where

p is the number of traits and *nb* is the number of populations sampled) while retaining PCs accounting for >

90% of variance.

Test of selection: solving the multivariate breeder's equation

It has been suggested that in Drosophila, flight performance in different thermal conditions might be under selection, favouring the evolution of latitudinal gradients in wing size and shape (e.g. Gilchrist *et al.* 2001). This hypothesis is indirectly supported by phenotypic plasticity analyses, showing better flight performance in cold air in flies reared at low temperatures, associated with different wing morphologies (Frazier *et al.* 2008). While we are mostly oblivious to the selective factors affecting *D. suzukii* invasive success (but see Camus *et al.* 2025), and specifically the evolution of its wing morphology, flight performance might play a role in its dispersive ability, or be under selection in the diverse thermal conditions encountered across its latitudinal ranges. We used data obtained in a previous study showing an association between wing shape and flight speed in *D. suzukii* (Fraimout *et al.* 2018), to explore the hypothesis that selection on flight speed might have driven wing shape divergence among invasive populations, using the multivariate breeder's equation (Lande 1979). First, an experimental selection differential was derived from Fraimout *et al*. (2018) data. In this previous study, we reconstructed 3D flight trajectories of flies reared under different developmental temperatures and measured flight parameters such as speed, acceleration, or sinuosity. Recorded individual wings were digitized and wing landmarks coordinates used to assess the relationship between flight parameters and wing shape (i.e. we identified wing shape variation associated with flight speed variation). Landmark data from this previous study were aligned onto the reference configuration of the present study to allow direct comparability. Solving the breeder's equation assumes that traits change is correlated with fitness. In the absence of fitness data, we hypothesized that flight performance might significantly contribute to fitness in Drosophila (e.g. Frazier *et al*. 2008) and we therefore used the shape change associated with the strongest flight speed difference as a hypothetical selection differential. Wing shape was regressed onto the log of flight speed using a multivariate regression (*e.g.* Mardia *et al.* 1979; Monteiro 2003) to estimate the vector of shape change associated with highest flight velocity s_{velocity}. The corresponding response to selection Δz was then predicted using the multivariate breeder's equation (Lande & Arnold 1983) and the estimated ancestral (*i.e.* Japanese) **G** and **P** matrices:

$$\Delta z = G_{jap} P_{jap}^{-} s_{velocity}$$

where $^{-}$ is the Moore-Penrose generalized inverse. This predicted response to velocity selection was compared

368   to the different vectors of population divergences (see above) by computing the angle between them following

$$\alpha2 = cos^{-1} \frac{\Delta z}{\|\Delta z\|} \cdot \frac{d^t}{\|d\|}$$

The observed angles were then compared to the expectation of the angle between Δz and 1000 random vectors

(see above). Our hypothesis here is that if a high flight speed has been favoured by selection throughout the

invasion, we should observe a match between the evolutionary shape changes (the phenotypic divergence

among populations) and the shape change resulting from the hypothetical response to selection.

Test of selection: percentage of genetic variance along divergence vectors

Finally, we reasoned that if selection was involved in the short-term evolution of wing shape, it should manifest

in a depletion of genetic variation in the selected directions (*e.g.* Agrawal & Stinchcombe 2009), corresponding

to the directions of divergence from the hypothetical common ancestor (Fig. 5). To test this prediction, **G**

matrices of each population were projected onto the shape space directions defined by the divergence vectors,

$d^t Gd$, and expressed in proportion of the total genetic variance. The amount of genetic variance in the

direction defined by *d* was then compared between the ancestral and derived **G** matrices. We predicted that a

reduction in this proportion of variance should be detected if selection was involved in the evolutionary

divergence, as depletion of genetic variance due to drift would randomly affect all directions. To test whether

genetic variance along the directions of divergence was lower than expected by chance, we compared it to the

variance estimated in random directions in the (tangent) shape space.

Results

Significant divergence of wing size and shape across populations

We observed differentiation in wing centroid size between American and all other populations, as well as

between the two American populations (Fig. S1): the American populations displayed smaller wing size, with

US-WAT displaying the smallest wings. Although the divergence for wing shape among populations was

weak, the *bgPCA* showed a signal of geographic differentiation, opposing the pairs of populations according

to their geographic origin (Fig. S2). This shape divergence was confirmed by the significant population effect

in the MANOVA (Wilk's lambda = 0.143, df = 5, $p < 0.001$). The MANOVA performed on the residuals of

the regression between shape and size also revealed a significant population effect (Wilk's lambda = 0.199, df = 5, $p < 0.001$), suggesting that the shape difference goes beyond a simple allometric effect. The pairwise permutational MANOVAs revealed significant shape differences between all pairs of populations ($p < 1e-04$). The analysis of the angles between the divergence vectors suggests that both French populations diverged from their common ancestor in the same direction of the phenotypic space: these two vectors are more similar than two random vectors ($\alpha = 31.9°$, $p < 10^{-4}$, Fig. 2, Table S2). For the American populations, the angle between the two divergence vectors was greater than that of the French populations ($\alpha = 49.9°$, $p = 5 \times 10^{-4}$, Fig 2, Table S2) and hint towards divergence along different directions. The average American and French direction of divergence from their common ancestor were strikingly different, and no more similar than random vectors ($\alpha = 71.9°$, $p = 0.11$, Table S2). In terms of amount of divergence based on the Procrustes distance, the American populations appeared to have slightly less diverge than French populations, from their hypothetical ancestor (5.72×10^{-3}, 7.46×10^{-3}, respectively; Table S3) and as observed from the *bgPCA* (Fig. S2).

*The **G**-matrix for wing shape is mostly stable across populations*

We found significant differences in the size of the **G** matrix among 3 pairs of populations, all involving the southern USA population US-WAT (Table 1). The **G** matrix of the US-WAT population had a significantly larger volume than those of US-DAY (northern USA) and of both French populations (FR-PAR and FR-MON). None of the other pairwise comparisons revealed any significant differences in the volumes of **G**.

Regarding the shape of the **G** matrices, none of the **G** matrices exhibited pronounced eccentricity (i.e. no single PC dominated the others, inducing an elongated, "cigar-shaped" matrix). Visual inspection of the first two principal components indicated that all **G** matrices were approximately circular in two-dimensional space (with the exception of FR-MON), and would likely appear relatively spherical if represented as three-dimensional ellipsoids. (Fig. 3). Eccentricity values calculated as the ratio of the first to the second eigenvalues of **G** were indeed low and close to 1 for all matrices (Fig. S3). In other words, genetic variation in each population was relatively equally distributed among combinations of traits. We did not find any significant differences in the shape of the **G** matrices as eccentricity and the amount of variance explained by g_{max} between any pairs of populations were not statistically different (Fig. S3). Similarly, no significant difference was found between the eigentensors of **S** calculated from the set of random and observed sets of **G** matrices based on the

overlapping 95% HPD intervals (Fig. 4), suggesting a global stability of the **G** matrices across populations.

*Proportionality of **D** and ancestral **G**: a constrained response to selection?*

Overall, the divergence matrix appeared fairly similar to the ancestral **G** matrix, its first two PCs and the

ancestral **G** matrix respectively forming narrow angles of 13.6 and 15.8°, smaller than expected for random

vectors (47.9°± 7.18). While the null hypothesis of proportionality between both matrices was not rejected

(Bartlett-adjusted X^2 test; *p* = 1), suggesting a close match, the coefficient of proportionality (ρ = 1.063; 95%

CI, 0.763 – 1.751) was in turn higher than the coefficient estimated for microsatellite markers ($2F_{ST}/(1-F_{ST})$ =

0.160; 95% CI, 0.128 – 0.198). Therefore, while **D** and **G** are proportional, their coefficient of proportionality

deviates from the neutral expectation, suggesting a role for selection in the observed divergence.

No reduction in genetic variance in the direction of divergence

We further projected each **G** matrix onto their respective direction of divergence from the hypothetical ancestor

and estimated the corresponding percentage of variance (Fig. 5). Overall, we found that amounts of genetic

variance in these directions were not statistically different among populations. Furthermore, we did not observe

a lower amount of genetic variance between the projection of each population **G** matrix and the projection of

the ancestral **G** matrix on their corresponding divergence vectors. This result goes against our expectation that

selection would result in a depletion of genetic variance along each population's divergence vector from the

ancestor.

The response to a theoretical selection does not match the observed divergences

To investigate the possible selective factors affecting wing shape evolution in *D. suzukii* invasion, we simulated

the evolutionary responses to a hypothetical selection differential, the wing shape difference between fast and

slow flying *D. suzukii* individuals, as reported in Fraimout *et al.* (2018; Fig. 6). The evolutionary response is

shown on Fig. 6, as the average response derived from the full posterior of MCMC samples. The shape

divergence of French and American populations from their common hypothetical ancestor are shown on Figure

6c and d (the two Japanese populations used to compute the ancestral **G** matrix were considered too close from

this ancestral state and were thus not considered). The pattern of shape change associated with $S_{velocity}$ and the

average response **Δz** were very similar (mean α = 22.74°, SD 6.03) confirming that **G** does not constrain the

response to selection. This result is in line with the relatively spherical shape of the ancestral **G** matrix, which

should not constrain the response to selection. While a similarity in the shift of the basal landmarks can be

found between the divergence of French populations and the theoretical response to selection, no obvious

similarity with American populations divergences is detected (Fig 6, Fig. S4). However, this congruence was

not reflected in the correlation between vectors when accounting for the full landmark configurations as the

evolution of the distal part of the wing did not match the expected response to selection (Fig. 6, Fig. S4).

Overall, the similarity between the divergence vectors from each population and the response to selection **Δz**

was low (Fig. S4) suggesting that the observed divergence in wing shape does not result from selection on

faster flight capacity.

Discussion

Genetic divergence of wing shape across populations of Drosophila suzukii

Geographical variation of wing morphology has been described in several *Drosophila* species (*e.g.* Azevedo

et al., 1998; Haas & Tolley, 1998; Hoffmann & Shirriffs, 2002), including introduced ones (Huey *et al.*, 2000).

In particular, geographical clines of wing size and shape have been detected along latitudinal and altitudinal

gradients (Huey *et al.* 2000; Gilchrist *et al.*, 2001; Gilchrist *et al.*, 2004; Pitchers *et al.*, 2013; Adrion *et al.*,

2015). These clines have typically been interpreted as adaptations to flight conditions, specifically temperature

and air pressure (Dillon & Frazier 2006; Pitchers *et al*. 2013), although direct evidence on the nature of

selection is lacking. In *D. suzukii*, clines in wing morphology have been reported along an altitudinal gradient

in Hawaii – the first and oldest invaded territory (Asplen *et al.*, 2015) – with flies from high altitude (ca.

2000m) displaying significantly larger wings than flies sampled at sea level (Koch *et al.*, 2020). Here, we

detected geographic variation of wing shape across all populations, but the differences among north and south

populations were different among continents. Gilchrist *et al.* (2000) reported differences in latitudinal wing

shape changes in *D. melanogaster* from three continents, and suggested the occurrence of specific local

selective pressures on wing shape. However, they also detected parallel clines in wing size across continents,

suggesting common adaptive effects. In our study, wing size was more stable: the only difference between

northern and southern populations was found in the USA. As for wing shape, although we found that

populations from the same continent have evolved in a relatively similar direction of the phenotypic space, we did not find any evidence for parallel latitudinal clines. At the time of sampling, the invasion of *D. suzukii* was initiated less than a decade ago in Europe and the USA and it is possible that adaptive geographical clines did not have enough time to evolve in these regions, contrary to the first invaded range in Hawaii (Koch *et al.* 2020). As a comparison, a latitudinal cline in *D. subobscura* following colonization of the New World, was reported after two decades, while no difference had been detected ten years after the start of the invasion (Huey *et al.*, 2000; Gilchrist *et al.*, 2004). At this point, it is unclear whether *D. suzukii* wing phenotypes are locally adapted, and additional work is warranted to decipher the interplay between wing plasticity, genetics and their association with fitness in natural *D. suzukii* populations. Nevertheless, by rearing flies in controlled laboratory conditions for a few generations, we showed that the phenotypic divergence observed among populations is genetic rather than plastic (see also Fraimout *et al* 2018) and that some evolutionary process independent of phenotypic plasticity have shaped the wings of *D. suzukii* in the wild.

G matrix of wing shape is stable across populations

Despite the wealth of theoretical and empirical research (e.g. Roff, 2000; Steppan *et al.*, 2002; Bégin & Roff, 2003; Jones *et al.*, 2003; Mezey & Houle, 2003; Björklund, 2004; Jones *et al.*, 2004; Blows & Hoffmann, 2005; Arnold *et al.*, 2008; Walsh & Blows, 2009; Jones *et al.* 2014; Wood & Brodie, 2015; Puentes *et al.*, 2016; Svensson *et al.*, 2021), predicting how natural selection or drift will affect the structure of **G** and at which pace **G** can evolve in the wild remains a difficult task. As stated by Jones *et al.* (2003): "[...] *the current state of analytical theory on G-matrix evolution is that existing theory cannot guarantee stability of the G-matrix (Turelli 1988), but also does not guarantee instability*." Following this assertion, simulation studies (Jones *et al.*, 2003; Jones *et al.*, 2004; Arnold *et al.*, 2008; Jones *et al.*, 2014; Björklund & Gustafsson, 2015; Chantepie & Chevin 2020) have identified how correlative selection, pleiotropic mutations, population size and other evolutionary forces can affect the stability of **G**'s structure and orientation. However, empirically and 20 years later, evidence for both stability (Delahaie *et al.*, 2017; McGlothlin *et al.*, 2018, Henry & Stinchcombe 2023) and lability (*e.g*., Uesugi *et al.*, 2017, Chantepie *et al*. 2024) have been reported in varying contexts.

Here we found that the **G** matrices were globally stable across populations, the differences between populations

mainly pertaining to the volume of **G** (i.e. the global amount of genetic variance). Specifically, French

populations seemed to display less genetic variance as indicated by their smaller volumes, while the US-WAT

population seemed more genetically variable than the rest of the populations. Reduction of genetic variance

and, therefore, of the volume of **G**, is expected under certain conditions such as increased genetic drift due to

reduced effective population size (Shaw *et al.*, 1995; Roff, 2000; Jones *et al.*, 2004) or increased strength of

stabilizing selection (Jones *et al.* 2003). According to the results of Fraimout *et al.* (2017) obtained from neutral

molecular markers and Olazcuaga *et al.* (2020) using whole-genome data, all invasive populations of *D. suzukii*

experienced significant loss of genetic diversity due to various intensities of bottlenecks upon introduction.

Lesser levels of genetic variance and lower volumes in the French **G** matrices are thus consistent with the

effect of drift. Similarly, the higher volume of the US-WAT population could be explained by the genetic

admixture of two highly differentiated lineages from Asia and Hawaii (Fraimout *et al.* 2017). However,

populations' invasion history was not reflected in all **G** matrices, as other genetically admixed populations

(FR-PAR, US-DAY; Fraimout *et al.* 2017) did not show higher volumes and neither did ancestral populations

where genetic bottlenecks most likely did not occur. Contrasting with the random evolution of **G** in all

directions of the multivariate space expected under genetic drift (Arnold *et al.*, 2001; Phillips *et al.*, 2001), our

results show that most aspects of the eigenstructure of **G** (size and shape) have remained stable throughout the

invasion process.

Our analyses may lack statistical power to detect more subtle differences between **G** matrices. Indeed, while

our crossing design consists of an acceptable number of genetic lines per population, the multidimensional

nature of the *Drosophila* wing implies a large number of parameters to be estimated (n x (n-1) /2 + n; 351 in

our case), larger than the number of replicates (*i.e.* isofemale lines). Nonetheless, replicating our analyses using

fewer numbers of PCs - and thus less parameters to be estimated - led to the same conclusions (see

Supplementary material). Finally, our design did not allow us to partition additive from non-additive genetic

components of wing shape variation. Dominance genetic variance has been shown to explain up to 35% of

phenotypic variation for wing traits in *D. serrata* (Sztepanacz & Blows 2015) and it is therefore possible that

some of our estimates of genetic variance could be inflated by such non-additive components. Furthermore,

the use of a limited number of isofemale lines to capture the possibly large standing genetic variation found in

D. suzukii's wild populations could further exacerbate the effects of non-additivity on our estimates of the **G**

matrix. Specifically, establishing isofemale lines could result in inbreeding depression, where mildly

deleterious mutations otherwise sheltered in heterozygotes are expressed in the laboratory under sib-mating

(Agrawal & Whitlock 2012). This in turns could inflate the contribution of dominance genetic variance

(Charlesworth & Willis 2009). While we did not estimate the contribution of dominance variance, the use of

gravid females mated in the wild may have reduced the effect of inbreeding depression upon establishing

isofemale lines. Whether between-population differences exist in the importance of dominance variance for

wing shape is unknown, and future work including appropriate breeding designs (*e.g.,* double-cousins) should

shed light on the importance of non-additive genetic components in wing shape variation of *D. suzukii*.

Phenotypic divergence is not constrained in D. suzukii's invasive populations

We tested the hypotheses of ancestral constraints on wing shape evolution by estimating the similarity between

the ancestral **G** matrix and the divergence matrix **D**, and found that the leading vectors of both matrices were

tightly associated. Such similarity can be viewed as a mark of genetic constraints affecting divergence,

compatible with both neutral and adaptive scenarios. Genetic drift is expected to induce divergence along the

axes with the most genetic variation, and proportionally to the amount of variation in the ancestral population

(Lande 1979; Lofsvold 1998; Marroig & Cheverud, 2004). But this pattern of concordance between **G** and **D**

is also expected under selection, the main directions of genetic variation biasing the response to selection (i.e.

the lines of genetic least resistance; Schluter, 1996). However, the estimated ancestral **G** matrix for *D. suzukii*

wing shape had a relatively spherical shape, suggesting the existence of genetic variation in all directions of

the phenotypic space, and therefore a lack of genetic constraint. This is evident from the results of our

phenotypic divergence vector analysis showing that, despite the relatively small differentiation among

populations, the evolution of wing shape in France and USA has followed very different directions (Fig. 2),

suggesting no constraint. Finally, the response to selection predicted from our theoretical selection gradient

applied to the estimated ancestral **G** produced a wing shape different from that observed in any of the extant

invasive populations, further supporting an absence of genetic constraint on wing shape evolution in this

species. These results are interesting in the light of recent research investigating the link between trait

evolvability and divergence (Tsuboi *et al.* 2024, Pélabon *et al.* 2025). Recent studies have shown that population divergence across species and traits tends to scale positively with standing additive genetic variance (*i.e.,* evolvability; Opedal *et al.* 2023, Holstad *et al.* 2024), suggesting that phenotypic evolution can be channeled by ancestral genetic constraints through lines of least resistance. Here, we did not find evidence for genetic lines of least resistance, but rather, unconstrained phenotypic divergence from ancestral standing genetic variation.

Adaptive or neutral divergence?

The results from our multivariate Q_{ST}-F_{ST} support an adaptive scenario for wing shape evolution, as divergence was found to be significantly greater than expected under neutral evolution. While failure to account for demographic history and population structure can bias the outcome of Q_{ST}-F_{ST} comparisons (de Villemereuil *et al.* 2022, do O *et al.* 2025), it does not seem to be the case here. The demographic history of *D. suzukii* is known, and there is no evidence for gene flow or migration among populations (even within-country; Fraimout *et al.* 2017) that could reduce values of F_{ST}, biasing upward the result for our test of selection ($\rho st > F_{ST}$). As for a possible inflation of ρst, this could happen if all populations had undergone drastic bottlenecks and reductions in **G** while **D** remained large. We do not have evidence that this happened in our study populations, as the volume of most invasive populations' **G** is not significantly smaller than that of the native population. The divergence in wing shape among invasive populations of *D. suzukii* might thus be adaptive. The statistical power of this test is conditional on the sampling design and particularly the number of populations sampled (Martin *et al.* 2008). While the test for departure of neutrality should be robust to a small number of populations (*i.e.,* < 10 as in the current study) results of the proportionality test should be interpreted with caution. Similar analyses using long-diverged groups (*i.e.,* Teosinte and Maize), report values of ρ_{ST} several order of magnitude above the neutral expectation (Yang *et al.* 2019), suggesting that the signal of selection we observe in *D. suzukii* is rather low. The adaptive value of wing shape variation in *Drosophila* is a long-standing conundrum, and in the absence of fitness data, remains speculative (Gilchrist *et al.* 2000). Given the role of wing in flight, we leveraged available flight data (Fraimout *et al* 2018) to test whether selection on flight speed could be involved in wing shape divergence among our populations. Despite partial correspondence, the response to this selection applied to the ancestral **G** matrix, was very different from the observed evolutionary divergence,

providing no support for selection on flight speed as a driver of this divergence.

Finally, it is worth stressing that we focused our analyses on the study of male wings only. Wing shape is under

strong sexual selection in Drosophila, as males produce a variety of visual and acoustic signalling using their

wings during courtship (Debelle *et al.* 2014). By focusing on male wings only, it is possible that we are missing

an important component of evolutionary constraint, as previous work has shown that genetic covariance across

sexes can bias the evolutionary response of wing shape to selection (McGuigan & Blows 2007, Sztepanacz &

Houle 2019). Nonetheless, recent work on *D. buzzati* (Iglesias *et al.* 2023) suggests that adaptive divergence

in wing shape can be attributable to selection on courtship traits in males among populations. Additionally,

Branden & De Lisle (2026) have shown that the effect of stabilizing selection is particularly pronounced in

male Drosophila, where departure from the average wing shape is associated with lower fitness in males. Our

results could be compatible with those studies, reflecting a combination of stabilizing selection on *D. suzukii*

males' wing shape and divergent selection among populations. Without access to fitness assays and its

relationship to wing shape variance in our study, this hypothesis cannot however be formally tested with the

data at hand. The nature of the selective factors affecting wing shape evolution thus remains to be identified

in *D. suzukii*, and further work would be valuable to shed light on the relationship between wing shape and

fitness in males, and on the effect of stabilizing selection on the wing shape **G** matrix.

Conclusion

Consensual predictions on the effect of selection and drift on the **G** matrix remain scarce in the literature and

most empirical evidence are based on experimental evolution studies (*e.g.* Doroszuk *et al.*, 2008; Careau *et al.*,

2015) or comparisons among long-diverged species (e.g. McGlothlin *et al.*, 2018). In this context our study

provides new insights on the evolution (or lack thereof) of **G** over a relatively short evolutionary time scale in

natural conditions. We showed that native and invasive populations of *D. suzukii* display quantitative genetic

differentiation in wing shape across continents and at an early stage of their worldwide invasion. Our results

suggest an effect of selection acting on a poorly constraining ancestral G matrix. They provide a reference

point for future studies quantifying phenotypic evolution in this species.

Acknowledgments

AF was supported by the Agence Nationale de la Recherche, through the LabEx ANR-10-LABX-0003-BCDiv, of the program "Investissements d'avenir" (ANR-11-IDEX-0004-02). VD was funded by ANR SWING (ANR-16-CE02-0015). The authors are grateful to S. Fellous, A. Xuéreb, K. Tamura, M. Toda, P. Shearer, T. Schlenke and A. Kopp for help with flies sampling. We thank M. Guillaume for helping with the stock flies maintenance and F. Peronnet for providing flies rearing medium.

Author contribution

AF, VD and CT designed the study; AF, SC and NN analyzed the data; AF and VD obtained the samples; AF led the writing of the first draft and all authors contributed critically to the manuscript.

Data availability

Data and code necessary to replicate the analyses presented in the manuscript have been deposited on the Zenodo platform (zenodo.org).

References

Adams, D. C. (2011). Quantitative genetics and evolution of head shape in Plethodon salamanders. *Evolutionary Biology*, *38*(3), 278-286.

Adrion, J. R., Hahn, M. W., & Cooper, B. S. (2015). Revisiting classic clines in Drosophila melanogaster in the age of genomics. *Trends in Genetics, 31*(8), 434-444.

Agrawal, A. F., & Stinchcombe, J. R. (2009). How much do genetic covariances alter the rate of adaptation? *Proc Biol Sci, 276*(1659), 1183-1191.

Agrawal, A. F., & Whitlock, M. C. (2012). Mutation load: the fitness of individuals in populations where deleterious alleles are abundant. *Annual Review of Ecology, Evolution, and Systematics*, *43*(1), 115-135.

Aguirre, J. D., Hine, E., McGuigan, K., & Blows, M. W. (2014). Comparing G: multivariate analysis of genetic variation in multiple populations. *Heredity*, *112*(1), 21-29.

Arnold, S. J., Burger, R., Hohenlohe, P. A., Ajie, B. C., & Jones, A. G. (2008). Understanding the evolution and stability of the G-matrix. *Evolution, 62*(10), 2451-2461.

Arnold, S. J., Pfrender, M. E., & Jones, A. G. (2001). The adaptive landscape as a conceptual bridge between micro-and macroevolution *Microevolution Rate, Pattern, Process* (pp. 9-32): Springer.

Asplen, M. K., Anfora, G., Biondi, A., Choi, D.-S., Chu, D., Daane, K. M., Gibert, P., Gutierrez, A. P., Hoelmer, K. A., Hutchison, W. D., *et al.* (2015). Invasion biology of spotted wing Drosophila (Drosophila suzukii): a global perspective and future priorities. *Journal of Pest Science, 88*(3), 469-494.

Azevedo, R. B., James, A. C., McCabe, J., & Partridge, L. (1998). Latitudinal variation of wing: thorax size ratio and wing-aspect ratio in Drosophila melanogaster. *Evolution*, *52*(5), 1353-

1362.

Bégin, M., & Roff, D. A. (2003). The constancy of the G matrix through species divergence and the effects of quantitative genetic constraints on phenotypic evolution: a case study in crickets. *Evolution*, *57*(5), 1107-1120.

Benjamini, Y. and Hochberg, Y. (1995). Controlling the false discovery rate: A practical and powerful approach to multiple testing. *J. Roy. Statist. Soc. Ser. B*, *57*, 289–300

Berger, D., Postma, E., Blanckenhorn, W. U., & Walters, R. J. (2013). Quantitative genetic divergence and standing genetic (co)variance in thermal reaction norms along latitude. *Evolution, 67*(8), 2385-2399.

Björklund, M. (2004). Constancy of the G Matrix in Ecological Time. *Evolution, 58*(6).

Björklund, M., & Gustafsson, L. (2015). The stability of the G-matrix: The role of spatial heterogeneity. *Evolution*, *69*(7), 1953-1958.

Bjorklund, M., Husby, A., & Gustafsson, L. (2013). Rapid and unpredictable changes of the G-matrix in a natural bird population over 25 years. *Journal of Evolutionary Biology, 26*(1), 1-13.

Blows, M. W., & Hoffmann, A. A. (2005). A reassessment of genetic limits to evolutionary change. *Ecology, 86*(6), 1371-1384.

Bock, D. G., Caseys, C., Cousens, R. D., Hahn, M. A., Heredia, S. M., Hubner, S., Turner, K. G.Whitney, K. D., & Rieseberg, L. H. (2015). What we still don't know about invasion genetics. *Molecular Ecology, 24*(9), 2277-2297.

Bookstein, F. L. (1996). Combining the tools of geometric morphometrics. In *Advances in morphometrics* (pp. 131-151). Boston, MA: Springer US.

Bränden, A., & De Lisle, S. P. (2026). Apparent stabilizing selection explains micro and macroevolution of Drosophila wings. *bioRxiv*, 2026-01.

Calabria, G., Máca, J., Bächli, G., Serra, L., & Pascual, M. (2012). First records of the potential pest species Drosophila suzukii (Diptera: Drosophilidae) in Europe. *Journal of Applied Entomology, 136*(1-2), 139-147.

Camus, L., Rode, N. O., Serga, S., Loiseau, A., Chen, X., Iampietro, C., ... & Gautier, M. (2025). Adaptive Challenges of Past and Future Invasion of Drosophila suzukii: Insights From Novel Genomic Resources and Statistical Methods Combining Individual and Pool Sequencing Data. *Molecular Ecology*, e70192.

Cano, J. M., Laurila, A., Palo, J., & Merilä, J. (2004). Population differentiation in G matrix structure due to natural selection in Rana temporaria. *Evolution*, *58*(9), 2013-2020.

Careau, V., Wolak, M. E., Carter, P. A., & Garland, T. (2015). Evolution of the additive genetic variance–covariance matrix under continuous directional selection on a complex behavioural phenotype. *Proceedings of the Royal Society B: Biological Sciences*, *282*(1819).

Chantepie, S., & Chevin, L. M. (2020). How does the strength of selection influence genetic correlations?. *Evolution letters*, *4*(6), 468-478.

Chantepie, S., Charmantier, A., Delahaie, B., Adriaensen, F., Matthysen, E., Visser, M. E., ... & Teplitsky, C. (2024). Divergence in evolutionary potential of life history traits among wild populations is predicted by differences in climatic conditions. *Evolution Letters*, *8*(1), 29-42.

Chapuis, E., Martin, G., & Goudet, J. (2008). Effects of selection and drift on G matrix evolution in a heterogeneous environment: a multivariate Q st–F st test with the freshwater snail Galba truncatula. *Genetics*, *180*(4), 2151-2161.

Chenoweth, S. F., Rundle, H. D., & Blows, M. W. (2010). The contribution of selection and genetic constraints to phenotypic divergence. *The American Naturalist*, *175*(2), 186-196.

Colautti, R. I., & Lau, J. A. (2016). Contemporary evolution during invasion: evidence for differentiation, natural selection, and local adaptation. *Invasion Genetics: The Baker and Stebbins Legacy*, 101-121.

Conner, J. K., Franks, R., & Stewart, C. (2003). Expression of additive genetic variances and covariances for wild radish floral traits: comparison between field and greenhouse

environments. *Evolution, 57*(3), 487-495.

David, J. R., Gibert, P., Legout, H., Pétavy, G., Capy, P., & Moreteau, B. (2005). Isofemale lines in Drosophila: an empirical approach to quantitative trait analysis in natural populations. *Heredity*, *94*(1), 3-12.

Debelle, A., Ritchie, M. G., & Snook, R. R. (2014). Evolution of divergent female mating preference in response to experimental sexual selection. *Evolution*, *68*(9), 2524-2533.

Delahaie, B., Charmantier, A., Chantepie, S., Garant, D., Porlier, M., & Teplitsky, C. (2017). Conserved G-matrices of morphological and life-history traits among continental and island blue tit populations. *Heredity*, *119*(2), 76-87.

de Villemereuil, P., Gaggiotti, O. E., & Goudet, J. (2022). Common garden experiments to study local adaptation need to account for population structure. *Journal of Ecology*, *110*(5), 1005-1009.

Dillon, M. E., & Frazier, M. R. (2006). Drosophila melanogaster locomotion in cold thin air. *Journal of Experimental Biology*, *209*(2), 364-371.

Dlugosch, K. M., Anderson, S. R., Braasch, J., Cang, F. A., & Gillette, H. D. (2015). The devil is in the details: genetic variation in introduced populations and its contributions to invasion. *Molecular Ecology, 24*(9), 2095-2111.

do O, I., Gaggiotti, O. E., de Villemereuil, P., & Goudet, J. (2025). A method for identifying local adaptation in structured populations. *PLoS Genetics*, *21*(9), e1011871.

Doroszuk, A., Wojewodzic, M. W., Gort, G., & Kammenga, J. E. (2008). Rapid divergence of genetic variance-covariance matrix within a natural population. *The American Naturalist, 171*(3), 291-304.

Dryden, I. L., & Mardia, K. V. (1993). Multivariate shape analysis. *Sankhyā: The Indian Journal of Statistics, Series A*, 460-480.

Dugand, R. J., Aguirre, J. D., Hine, E., Blows, M. W., & McGuigan, K. (2021). The contribution of mutation and selection to multivariate quantitative genetic variance in an outbred population of Drosophila serrata. *Proceedings of the National Academy of Sciences*, *118*(31).

Eroukhmanoff., & Svensson, E. I. (2011). Evolution and stability of the G-matrix during the colonization of a novel environment. *Journal of Evolutionary Biology, 24*(6), 1363-1373.

Estoup, A., Ravigné, V., Hufbauer, R., Vitalis, R., Gautier, M., & Facon, B. (2016). Is There a Genetic Paradox of Biological Invasion? *Annual Review of Ecology, Evolution, and Systematics, 47*(1), 51-72.

Facon, B., Genton, B. J., Shykoff, J., Jarne, P., Estoup, A., & David, P. (2006). A general eco-evolutionary framework for understanding bioinvasions. *Trends in Ecology & Evolution, 21*(3), 130-135.

Fischer, E. K., Ghalambor, C. K., & Hoke, K. L. (2016). Plasticity and evolution in correlated suites of traits. *Journal of Evolutionary Biology, 29*(5), 991-1002.

Fraimout, A., Debat, V., Fellous, S., Hufbauer, R. A., Foucaud, J., Pudlo, P., ... & Estoup, A. (2017). Deciphering the routes of invasion of Drosophila suzukii by means of ABC random forest. *Molecular biology and evolution*, *34*(4), 980-996.

Fraimout, A., Jacquemart, P., Villarroel, B., Aponte, D. J., Decamps, T., Herrel, A., ... & Debat, V. (2018). Phenotypic plasticity of Drosophila suzukii wing to developmental temperature: implications for flight. *Journal of Experimental Biology*, *221*(13).

Fraimout, A., Loiseau, A., Price, D. K., Xuereb, A., Jean-Francois, M., Vitalis, R., Fellous, S., Debat, V., & Estoup, A. (2015). New set of microsatellite markers for the spotted-wing Drosophila suzukii (Diptera: Drosophilidae): A promising molecular tool for inferring the invasion history of this major insect pest. *European Journal of Entomology, 112*(4), 855-859.

Fraimout, A., Päiviö, E., & Merilä, J. (2022). Relaxed risk of predation drives parallel evolution of stickleback behavior. *Evolution*, *76*(11), 2712-2723.

Frazier, M. R., Harrison, J. F., Kirkton, S. D., & Roberts, S. P. (2008). Cold rearing improves cold-flight performance in Drosophila via changes in wing morphology. *Journal of Experimental*

Biology, *211*(13), 2116-2122.

Gilchrist, G. W., Huey, R. B., Balanyà, J., Pascual, M., & Serra, L. (2004). A Time Series of Evolution
in Action: A Latitudinal Cline in Wing Size in South American Drosophila Subobscura.
Evolution, 58(4).

Gilchrist, G. W., Huey, R. B., & Serra, L. (2001). Rapid evolution of wing size clines in Drosophila
subobscura. *Genetica*, *112*(1), 273-286.

Golani, D., Azzurro, E., Corsini-Foka, M., Falautano, M., Andaloro, F., & Bernardi, G. (2007).
Genetic bottlenecks and successful biological invasions: the case of a recent Lessepsian
migrant. *Biology letters*, *3*(5), 541-545.

Goudet, J., Jombart, T., & Goudet, M. J. (2015). Package 'hierfstat'. Estimation and Tests of
Hierarchical F-Statistics.

Guillaume, F., & Whitlock, M. C. (2007). Effects of migration on the genetic covariance matrix.
Evolution, *61*(10), 2398-2409.

Haas, H. L., & Tolley, K. A. (1998). Geographic variation of wing morphology in three Eurasian
populations of the fruit fly, Drosophila lummei. *Journal of Zoology*, *245*(2), 197-203.

Hadfield, J. D. (2010). MCMC methods for multi-response generalized linear mixed models: the
MCMCglmm R package. *Journal of Statistical Software, 33*(2), 1-22.

Hansen, T. F., & Houle, D. (2004). Evolvability, stabilizing selection, and the problem. *Phenotypic*
integration: Studying the ecology and evolution of complex phenotypes, *130*.

Hauser, M. (2011). A historic account of the invasion of Drosophila suzukii (Matsumura) (Diptera:
Drosophilidae) in the continental United States, with remarks on their identification. *Pest*
Manag Sci, 67(11), 1352-1357.

Hayes, K. R., & Barry, S. C. (2008). Are there any consistent predictors of invasion success?
Biological Invasions, 10(4), 483-506.

Henry, G. A., & Stinchcombe, J. R. (2023). G-matrix stability in clinally diverging populations of
an annual weed. *Evolution*, *77*(1), 49-62.

Hervé, M. (2016). RVAideMemoire: diverse basic statistical and graphical functions. R package v
0.9-55.

Hine, E., Chenoweth, S. F., Rundle, H. D., & Blows, M. W. (2009). Characterizing the evolution of
genetic variance using genetic covariance tensors. *Philosophical Transactions of the Royal*
Society B: Biological Sciences, *364*(1523), 1567-1578.

Hoffmann, A. A., & Parsons, P. A. (1988). The analysis of quantitative variation in natural
populations with isofemale strains. *Génétique sélection évolution*, *20*(1), 87-98.

Hoffmann, A. A., & Shirriffs, J. (2002). Geographic variation for wing shape in Drosophila serrata.
Evolution, *56*(5), 1068-1073.

Houle, D., & Meyer, K. (2015). Estimating sampling error of evolutionary statistics based on genetic
covariance matrices using maximum likelihood. *Journal of Evolutionary Biology*, *28*(8),
1542-1549.

Huey, R. B., Gilchrist, G. W., Carlson, M. L., Berrigan, D., & Serra, L. (2000). Rapid evolution of a
geographic cline in size in an introduced fly. *Science*, *287*(5451), 308-309.

Jolliffe, I. T. (2002). Graphical representation of data using principal components. In *Principal*
component analysis (pp. 64-91). New York, NY: Springer New York.

Jones, A. G., Arnold, S. J., & Bürger, R. (2003). Stability of the G-matrix in a population experiencing
pleiotropic mutation, stabilizing selection, and genetic drift. *Evolution*, *57*(8), 1747-1760.

Jones, A. G., Arnold, S. J., & Burger, R. (2004). Evolution and stability of the G-matrix on a
landscape with a moving optimum. *Evolution*, *58*(8), 1639-1654.

Jones, A. G., Bürger, R., & Arnold, S. J. (2014). Epistasis and natural selection shape the mutational
architecture of complex traits. *Nature communications, 5*(1), 1-10.

Keller, S. R., & Taylor, D. R. (2008). History, chance and adaptation during biological invasion:
separating stochastic phenotypic evolution from response to selection. *Ecology letters*, *11*(8),

852-866.

Kilkenny, F. F., & Galloway, L. F. (2013). Adaptive divergence at the margin of an invaded range.
Evolution, 67(3), 722-731.

Klingenberg, C. P., & Leamy, L. J. (2001). Quantitative genetics of geometric shape in the mouse
mandible. *Evolution*, *55*(11), 2342-2352.

Klingenberg, C. P., & McIntyre, G. S. (1998). Geometric morphometrics of developmental
instability: analyzing patterns of fluctuating asymmetry with Procrustes methods. *Evolution*,
52(5), 1363-1375.

Klingenberg, C. P. (2003). Quantitative genetics of geometric shape: heritability and the pitfalls of
the univariate approach. *Evolution*, *57*(1), 191-195.

Klingenberg, C. P. (2010). Evolution and development of shape: integrating quantitative approaches.
Nature Reviews Genetics, *11*(9), 623-635.

Koch, J. B., Dupuis, J. R., Jardeleza, M.-K., Ouedraogo, N., Geib, S. M., Follett, P. A., & Price, D.
K. (2020). Population genomic and phenotype diversity of invasive Drosophila suzukii in
Hawai 'i. *Biological Invasions*, 1-18.

Kruuk, L. E. (2004). Estimating genetic parameters in natural populations using the 'animal model'.
Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences,
359(1446), 873-890.

Krzanowski, W. J. (1979). Between-groups comparison of principal components. *Journal of the*
american statistical association, *74*(367), 703-707.

Lande, R. (1976). Natural selection and random genetic drift in phenotypic evolution. *Evolution*, 314-
334.

Lande, R. (1979). Quantitative genetic analysis of multivariate evolution, applied to brain: body size
allometry. *Evolution*, 402-416.

Lande, R., & Arnold, S. J. (1983). The measurement of selection on correlated characters. *Evolution*,
1210-1226.

Lee, C. E. (2002). Evolutionary genetics of invasive species. *Trends in Ecology & Evolution, 17*(8),
386-391.

Leinonen, T., Cano, J. M., & Merila, J. (2011). Genetics of body shape and armour variation in
threespine sticklebacks. *Journal of Evolutionary Biology, 24*(1), 206-218.

Lofsvold D. (1988). Quantitative genetics of morphological differentiation in Peromyscus. II.
Analysis of selection and drift. *Evolution, 42(1)*:54-67.

Lynch, M., & Walsh, B. (1998). *Genetics and Analysis of Quantitative Traits*: Sinauer.

Mardia, K. V., Kent, J. T., & Bibby, J. M. (1979). Multivariate analysis.

Marroig, G., & Cheverud, J. M. (2004). Did natural selection or genetic drift produce the cranial
diversification of neotropical monkeys?. *The American Naturalist*, *163*(3), 417-428.

Martin, G., Chapuis, E., & Goudet, J. (2008). Multivariate Q st–F st comparisons: a neutrality test for
the evolution of the G matrix in structured populations. *Genetics*, *180*(4), 2135-2149.

Martínez-Abadías, N., Mitteroecker, P., Parsons, T. E., Esparza, M., Sjøvold, T., Rolian, C., ... &
Hallgrímsson, B. (2012). The developmental basis of quantitative craniofacial variation in
humans and mice. *Evolutionary Biology*, *39*(4), 554-567.

Matsumura, S. (1931). *6000 illustrated insects of Japan-Empire*.

McGuigan, K., & Blows, M. W. (2007). The phenotypic and genetic covariance structure of
drosphilid wings. *Evolution*, *61*(4), 902-911.

McGlothlin, J. W., Kobiela, M. E., Wright, H. V., Mahler, D. L., Kolbe, J. J., Losos, J. B., & Brodie,
E. D., 3rd. (2018). Adaptive radiation along a deeply conserved genetic line of least resistance
in Anolis lizards. *Evolution Letters, 2*(4), 310-322.

McGuigan, K. J. M. (2006). Studying phenotypic evolution using multivariate quantitative genetics.
Molecular Ecology, 15(4), 883-896.

Merilä, J., Bjorklund, M., & Gustafsson, L. (1994). Evolution of Morphological Differences with

Moderate Genetic Correlations among Traits as Exemplified by 2 Flycatcher Species (Ficedula, Muscicapidae). *Biological Journal of the Linnean Society, 52*(1), 19-30.

Merilä, J. & Björklund, M. (2004). Phenotypic integration as a constraint and adaptation. *Phenotypic integration: studying the ecology and evolution of complex phenotypes*, 107-129.

Mezey, J. G., & Houle, D. (2003). Comparing G matrices: Are common principal components informative? *Genetics, 165*(1), 411-425.

Mezey, J. G., & Houle, D. (2005). The dimensionality of genetic variation for wing shape in Drosophila melanogaster. *Evolution*, *59*(5), 1027-1038.

Mitteroecker, P., & Bookstein, F. (2011). Linear Discrimination, Ordination, and the Visualization of Selection Gradients in Modern Morphometrics. *Evolutionary Biology, 38*(1), 100-114.

Monteiro, L. R., Duarte, L. C., & dos Reis, S. F. (2003). Environmental correlates of geographical variation in skull and mandible shape of the punar rat Thrichomys apereoides (Rodentia: Echimyidae). *Journal of Zoology, 261*(1), 47-57.

Morrissey, M. B., Hangartner, S., & Monro, K. (2019). A note on simulating null distributions for G matrix comparisons. *Evolution*, *73*(12), 2512-2517.

Myers, E. M., Janzen, F. J., Adams, D. C., & Tucker, J. K. (2006). Quantitative genetics of plastron shape in slider turtles (Trachemys scripta). *Evolution*, *60*(3), 563-572.

Olazcuaga, L., Loiseau, A., Parrinello, H., Paris, M., Fraimout, A., Guedot, C., ... & Gautier, M. (2020). A whole-genome scan for association with invasion success in the fruit fly Drosophila suzukii using contrasts of allele frequencies corrected for population structure. *Molecular biology and evolution*, 37(8), 2369-2385.

Pélabon, C., Hansen, T. F., Carter, A. J., & Houle, D. (2010). Evolution of variation and variability under fluctuating, stabilizing, and disruptive selection. *Evolution*, *64*(7), 1912-1925.

Pélabon, C., Agudelo-Cantero, G. A., Araya Ajoy, Y. G., Bolstad, G. H., Cheng, C., Galis, F., ... & Wright, J. (2025). Evolvability: progress and key questions. *BioScience*, *75*(12), 1042-1057.

Phillips, P. C., Whitlock, M. C., & Fowler, K. (2001). Inbreeding changes the shape of the genetic covariance matrix in Drosophila melanogaster. *Genetics*, *158*(3), 1137-1145.

Pitchers, W., Pool, J. E., & Dworkin, I. (2013). Altitudinal clinal variation in wing size and shape in African Drosophila melanogaster: one cline or many?. *Evolution*, *67*(2), 438-452.

Polly, P. D. (2008). Developmental dynamics and G-matrices: Can morphometric spaces be used to model phenotypic evolution?. *Evolutionary biology*, vol. 35, no 2, p. 83-96.

Prentis, P. J., Wilson, J. R., Dormontt, E. E., Richardson, D. M., & Lowe, A. J. (2008). Adaptive evolution in invasive species. *Trends in Plant Science, 13*(6), 288-294.

Puentes, A., Granath, G., & Agren, J. (2016). Similarity in G matrix structure among natural populations of Arabidopsis lyrata. *Evolution, 70*(10), 2370-2386.

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. https://www.R-project.org/

Rius, M., & Darling, J. A. (2014). How important is intraspecific genetic admixture to the success of colonising populations? *Trends in Ecology & Evolution, 29*(4), 233-242.

Roff, D. (2000). The evolution of the G matrix: selection or drift? *Heredity*, *84*(2), 135-142.

Rogers, A. R., & Harpending, H. C. (1983). Population structure and quantitative characters. *Genetics*, *105*(4), 985.

Rogers, A. R., & Harpending, H. C. (1983). Population structure and quantitative characters. *Genetics*, *105*(4), 985.

Schluter, D. (1996). Adaptive radiation along genetic lines of least resistance. *Evolution*, *50*(5), 1766-1774.

Shaw, F. H., Shaw, R. G., Wilkinson, G. S., & Turelli, M. (1995). Changes in genetic variances and covariances: G whiz!. *Evolution*, 1260-1267.

Sikkink, K. L., Reynolds, R. M., Cresko, W. A., & Phillips, P. C. (2015). Environmentally induced changes in correlated responses to selection reveal variable pleiotropy across a complex

genetic network. *Evolution, 69*(5), 1128-1142.

Siren, J., Ovaskainen, O., & Merila, J. (2017). Structure and stability of genetic variance-covariance
matrices: A Bayesian sparse factor analysis of transcriptional variation in the three-spined
stickleback. *Molecular Ecology, 26*(19), 5099-5113.

Spitze, K. (1993). Population structure in Daphnia obtusa: quantitative genetic and allozymic
variation. *Genetics*, *135*(2), 367-374.

Steppan, S. J., Phillips, P. C., & Houle, D. (2002). Comparative quantitative genetics: evolution of
the G matrix. *Trends in ecology & evolution*, *17*(7), 320-327.

Sztepanacz, J. L., & Blows, M. W. (2015). Dominance genetic variance for traits under directional
selection in Drosophila serrata. *Genetics*, *200*(1), 371-384.

Sztepanacz, J. L., & Houle, D. (2021). Allometry constrains the evolution of sexual dimorphism in
Drosophila across 33 million years of divergence. *Evolution*, *75*(5), 1117-1131.

Stuart, Y. E., Veen, T., Weber, J. N., Hanson, D., Ravinet, M., Lohman, B. K., ... & Bolnick, D. I.
(2017). Contrasting effects of environment and genetics generate a continuum of parallel
evolution. *Nature ecology & evolution*, *1*(6), 0158.

Svensson, E. I., Arnold, S. J., Bürger, R., Csilléry, K., Draghi, J., Henshaw, J. M., Jones, A. G., De
Lisle, S., Marques, D. A., & McGuigan, K. (2021). Correlational selection in the age of
genomics. *Nature Ecology and Evolution*, 1-12.

Szűcs, M., Vahsen, M. L., Melbourne, B. A., Hoover, C., Weiss-Lehman, C., & Hufbauer, R. A.
(2017). Rapid adaptive evolution in novel environments acts as an architect of population
range expansion. *Proceedings of the National Academy of Sciences*, *114*(51), 13501-13506.

Tsuboi, M., Gaboriau, T., & Latrille, T. (2024). An introduction to the special issue: inferring
macroevolutionary patterns and processes from microevolutionary mechanisms. *Journal of
evolutionary biology*, *37*(12), 1395-1401.

Turelli, M. (1988). Phenotypic evolution, constant covariances, and the maintenance of additive
variance. *Evolution*, 1342-1347.

Uesugi, A., Connallon, T., Kessler, A., & Monro, K. (2017). Relaxation of herbivore-mediated
selection drives the evolution of genetic covariances between plant competitive and defense
traits. *Evolution, 71*(6), 1700-1709.

Walsh, B., & Blows, M. W. (2009). Abundant Genetic Variation plus Strong Selection = Multivariate
Genetic Constraints: A Geometric View of Adaptation. *Annual Review of Ecology Evolution
and Systematics, 40*, 41-59.

Walsh, B., & Lynch, M. (2018). *Evolution and selection of quantitative traits*. Oxford University
Press.

Walter, G. M., Aguirre, J. D., Blows, M. W., & Ortiz-Barrientos, D. (2018). Evolution of genetic
variance during adaptive radiation. *The American Naturalist*, *191*(4), E108-E128.

Walter, G. M. (2023). Experimental evidence that phenotypic evolution but not plasticity occurs along
genetic lines of least resistance in homogeneous environments. *The American Naturalist*,
201(4), E70-E89.

Wares, J. (2005). Mechanisms that drive evolutionary change. Insights from species introduction and
invasions. *Species invasions: insights into ecology, evolution, and biogeography*, 229-257.

Whitlock, M. C. (2008). Evolutionary inference from QST. *Molecular ecology*, *17*(8), 1885-1896.

Wood, C. W., & Brodie, E. D., 3rd. (2015). Environmental effects on the structure of the G-matrix.
Evolution, 69(11), 2927-2940.

Yang, C. J., Samayoa, L. F., Bradbury, P. J., Olukolu, B. A., Xue, W., York, A. M., ... & Doebley, J.
F. (2019). The genetic architecture of teosinte catalyzed and constrained maize domestication.
Proceedings of the National Academy of Sciences, *116*(12), 5643-5652.

FIGURES AND TABLES

Figure 1. Sampling locations of the six *D. suzukii* populations. Adult *D. suzukii* were sampled from the wild in 3 countries (dark grey shading) from the native range in Japan (green circles) and the invasive range in France (blue circles) and the USA (red circles).

Figure 2. Phenotypic divergence of the invasive populations from their common ancestor. Graphical representation of the vectors of phenotypic divergence of each invasive population (solid line) from their hypothetical common ancestor (Japan, grey filled circle). The vectors are projected on the multivariate divergence space where d1 and d2 represent the first and second main axes of the multivariate divergent covariance matrix. The magnitude (length of the arrow) and direction (angles between arrows) of the vectors reflect the difference in mean wing shape phenotype among groups. Dashed lines represent the vector of phenotypic divergence of the average wing shape by country (USA or France) from the Japanese hypothetical ancestor. Colors correspond to Japanese (green), French (blue) and American (red) wing shape. Wing shape outlines represent the mean shape of each group and show the wing shape evolution from the Japanese ancestor. Outlines were obtained via a thin plate spine warping of a reference outline using MorphoJ (Klingenberg, 2011), based on the position of landmarks and are used for visualization purposes only.

Figure 3. Graphical representation of the population-specific G-matrices. For each population-specific **G** matrix, the projection of the mean **G** matrix (red circles) onto the first two axes of the eigen decomposition (PC1 & PC2) are shown (top 6 panels). For each population, a sample of 200 ellipses corresponding to 200 random posteriors drawn from the MCMC outputs (grey ellipses) illustrate the uncertainty around our estimated **G** matrices. Results from the eigen decomposition of each matrix is shown by plotting the percentage of variance explained by the first 11 PCs (blue histograms; mean total variance explained across PC and populations = 91.22%) for both observed matrices, and null matrices (black filled points). This shows that the main axes of variation (PC1 and PC2) explain more variance than expected at random. It is important to note that no conclusions should be drawn beyond PC3, as differences in eigenvectors between null and observed **G** matrix prevent interpretation.

Figure 4. Results of the genetic covariance tensor analysis. Eigenvalues of the 5 observed eigentensors of
S (black) are shown against the random expectation from the estimated null **G** matrices(grey). Overlapping of
the 95% confidence intervals indicates that the six observed **G** matrices are not more different than a random
set of **G** matrices.

Figure 5. Percentage of genetic variance along divergence vectors. A) Conceptual figure for our test of selection based on the percentage of genetic variance along divergence vectors. In a bivariate trait space, population means (black filled circle) of three populations have diverged from a common ancestor (ANC). The within-population variance (Var.pop) corresponds to the amount of genetic variance in the direction of the phenotypic change between the ancestor (ANC) and the focal population. The ancestral variance (Var.ANC) corresponds to the amount of genetic variance in the ancestral **G** matrix, along that same divergence axis. According to our hypothesis, reduction of the within-population variance (shortening of the Var.pop arrow) would be suggestive of the effect of selection. B) For each population, the observed amount of genetic variation in the direction of divergence was computed (Var. pop.) and compared to the amount of variation in the ancestral **G** matrix in that direction (Var. ANC). The percentage of genetic variance along this axis is plotted (filled grey dot) with the 95% highest posterior density interval (black horizontal lines) next to the population code on the y axis. We hypothesize that selection would result in a reduction in genetic variance in the directions of divergence. This is clearly not observed, as none of the directions shows depleted genetic variance.

Figure 6. Response to a hypothetical selection compared to observed shape changes. The observed wing shape change of the average French (blue arrows) and USA (red arrows) population are shown, representing the directions of change for each landmark, from the ancestral wing shape. The shape change associated with the selection differential defined by the shape difference between slow and fast fliers from Fraimout et al (2018) ($s_{velocity}$) is shown with black arrows on the left panels. The shape change corresponding to the response to selection (Δz) based on the MCMC estimates of the ancestral **G** and **P** matrices is shown with black arrows on right panels. The results show similarity in the shift of the basal landmarks between the divergence of French populations and the theoretical response to selection (alignment between blue and black arrows of the leftmost landmarks; upper right panel) but no congruence in any other comparison, suggesting limited evidence for selection on flight performance in these populations.

Population	JP-TOK	JP-SAP	US-DAY	US-WAT	FR-PAR
JP-SAP	1.03 [0.83:1.3]				
US-DAY	0.88 [0.69:1.08]	0.87 [0.66:1.03]			
US-WAT	1.17 [0.95:1.45]	1.14 [0.9:1.38]	**1.32 [1.09:1.65]**		
FR-PAR	0.93 [0.73:1.13]	0.89 [0.7:1.07]	1.05 [0.83:1.29]]	**0.79 [0.63:0.94]**	
FR-MON	0.93 [0.73:1.11]	0.88 [0.7:1.06]	1.04 [0.83:1.26]	**0.76 [0.62:0.92]**	0.95 [0.79:1.2]

Table 1. Pairwise comparison of G matrices' volumes. For each pair of **G** matrix the posterior mode of the ratio between volumes is shown along with the Highest Posterior Density (HPD) interval. Bold values correspond to significant differences between two matrices volumes whether the ratio is significantly > 1 (row matrix is larger) or < 1 (row matrix is smaller).