Musicians' brains at rest: Multilayer network analysis of MEG data

Kanad N Mandke[a,b], Prejaas Tewarie[c], Peyman Adjamian[d], Martin Schürmann[a,*, §], Jil Meier[e,f,g,*]

[a] *School of Psychology, University of Nottingham, Nottingham, United Kingdom*

[b] *Centre for Neuroscience in Education, Department of Psychology, University of Cambridge*

[b] *Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham, United Kingdom*

[d] *MRC, Institute of Hearing Research, University of Nottingham, Nottingham, United Kingdom*

[e] *Department of Neurology, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht, The Netherlands*

[f] *Current address: Berlin Institute of Health, Charité – Universitätsmedizin Berlin, Berlin, Germany*

[g] *Current address: Department of Neurology with Experimental Neurology, Brain Simulation Section, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany*

* *J.M. and M.S. contributed equally to this work*

[§] *Corresponding author: Dr Martin Schürmann (martin.schuermann@nottingham.ac.uk), School of Psychology, University of Nottingham, University Park, Nottingham NG7 2RD, UK, Phone: +44 115 84 679*

E-Mail-addresses of all other authors:

        Kanad Manke: knm27@cam.ac.uk

        Prejaas Tewarie: Prejaas.Tewarie@nottingham.ac.uk

Peyman Adjamian: peyman71@gmail.com

Jil Meier: jil-mona.meier@bih-charite.de

Short running title: "Multilayer networks of musician's brains"

Data Availability Statement

The data that support the findings of this study are openly available in Open MEG Archive

(OMEGA) at http://doi.org/10.23686/0015896.

Funding Statement

We thank the University of Nottingham for Vice Chancellor's Scholarship awarded to KM.

Conflicts of Interest Statement

The authors declare no conflict of interest or competing interests.

Keywords: Functional connectivity, modularity, multilayer networks, musicians, resting

state

Abstract

The ability to proficiently play a musical instrument requires a fine-grained synchronisation

between several sensorimotor and cognitive brain regions. Previous studies have

demonstrated that the brain undergoes functional changes with musical training,

identifiable also in resting-state data. These studies analysed fMRI or electrophysiological

frequency-specific brain networks in isolation. While the analysis of such "mono-layer"

networks has proven useful, it fails to capture the complexities of multiple interacting

networks. To this end, we applied a multilayer network framework for analysing publicly

available data (Open MEG Archive) obtained with magnetoencephalography (MEG). We

investigated resting-state differences between participants with musical training (n=31) and

those without (n=31). While single-layer analysis did not demonstrate any group

differences, multilayer analysis revealed that musicians show a modular organisation that

spans visuomotor and frontotemporal areas, known to be involved in musical performance

execution, which is significantly different from non-musicians. Differences between the two

groups are primarily observed in the theta (6.5-8Hz), alpha1 (8.5-10Hz) and beta1 (12.5-

16Hz) frequency bands.

We demonstrate that the multilayer method provides additional information that single-

layer analysis cannot. Overall, the multilayer network method provides a unique

opportunity to explore the pan-spectral nature of oscillatory networks, with studies of brain

plasticity as a potential future application.

Word count: 200

Introduction

Musical proficiency is a commonly used model to study brain plasticity (Herholz & Zatorre, 2012; Jäncke, 2009b; Zatorre & McGill, 2005). This research is largely motivated by the fact that achieving high levels of proficiency in any given musical instrument requires years of focused training. Numerous studies have reported structural brain changes due to musical training (Amunts et al., 1997; Elbert et al., 1995; Schlaug et al., 1995). For example, an increased size of the anterior half of the corpus callosum (Schlaug et al., 1995), and differences in the cortical representation of the hand motor area (Amunts et al., 1997; Elbert et al., 1995). More specifically, the brain also undergoes anatomical plasticity that pertains to the demands posed by the particular type of musical training, i.e., pianists showed a left hemisphere dominance, whereas violinists showed right hemisphere advantage (Bangert & Schlaug, 2006). Robust changes in grey matter specific to auditory, motor and visuo-spatial regions have also been reported (Gaser & Schlaug, 2003). This anatomical specialisation is potentially the result of years of musical training that requires fine-grained communication between several cortical areas. For example, musicians showed an increased coupling between auditory and pre-motor areas in functional magnetic resonance imaging (fMRI) (Grahn & Rowe, 2009), related to musicians' skilled actions during a performance.

It is worth noting that the training-related plastic changes are not unique to musicians but can be seen in groups of subjects with some degree of behavioural specialisation. For instance, training-related white matter changes have been observed in healthy adults learning to juggle (Scholz et al., 2010), professional golfers (Jäncke, 2009a), and London taxi drivers (Maguire et al., 2000).

To better characterise relationships between anatomical regions from task-free data,

resting-state functional connectivity approaches have proven useful. It must be noted that

either a seed-based or whole-brain functional connectivity analysis can inform about the

strengths of these connections (Friston, 2011). Going a step further, studying the pattern of

these connections can give us more information about the functional network as a whole.

Using a seed-based connectivity approach to resting-state fMRI in musicians, demonstrated

an increased functional connectivity in auditory, visual and motor areas. Similarly, (Tanaka &

Kirino, 2017) demonstrated an increased functional connectivity in the supplementary

motor area (SMA) network in musicians imagining musical performances based on fMRI

data. Using a whole-brain analysis of fMRI, (Alluri et al., 2017) reported that musicians

recruit action-based brain networks, while non-musicians recruit perception-based

networks in a naturalistic listening paradigm. In a seed-based fMRI study by (Alluri et al.,

2015a), musicians demonstrated an increased connectivity with SMA and with ventromedial

and ventrolateral cerebral and cerebellar affective regions while listening to music, whereas

non-musicians displayed a higher connectivity only with subcortical regions. In a resting-

state fMRI study, (Luo et al., 2014) demonstrated a greater local functional connectivity in

musicians compared to non-musicians in the bilateral dorsal anterior cingulate cortex,

anterior insula, and anterior temporoparietal junction. In a similar study, increased insular

connectivity in musicians was reported (Zamorano et al., 2017). As reviewed above,

evidence of functional specialisation in musicians comes from structural data (white or grey

matter changes) or haemodynamic functional data. Little attention has been paid so far to

the analysis of electrophysiological magnetoencephalography (MEG) data recorded from

musicians' brains during resting state, capturing spontaneous, non-task-related brain

activity, which could offer insights into the brain activity of musicians and the

communication between brain areas on shorter temporal time scales. Activity recorded by MEG predominantly consists of neuronal oscillations, which occur over a wide range of temporal scales (1-200Hz). Numerous studies have shown that these oscillations are involved in mediating both long-range and local communication in the brain (e.g., (Fries, 2005)).

Analysis of resting-state networks is not limited only to functional magnetic resonance imaging (fMRI) but has also been a significant part of the human electrophysiological research based on MEG (Brookes et al., 2011; Hillebrand et al., 2012) and electroencephalography (EEG) (Smit et al., 2008; Stam et al., 2005). However, to the best of our knowledge, only one study (Klein et al., 2016) has explored resting-state differences between musicians and non-musicians using an electrophysiological approach. The authors analysed high-density EEG resting-state data obtained from musicians with graph-theoretical approaches. They reported that musicians showed an increased intra- and inter-hemispheric functional connectivity between auditory, sensorimotor, and prefrontal cortex in theta (6.5-8Hz), alpha1 (8.5-10Hz), and alpha2 (10.5-12Hz) frequency ranges. A limitation of this study is that separate frequency specific networks were studied in isolation (also referred to as analysis of single layers), which ignores the pan-spectral picture of ongoing brain connectivity.

To overcome this limitation, the present study used a multilayer network framework, combining data obtained from multiple frequency bands in one single network description (Brookes et al., 2016a; Mandke et al., 2018; Tewarie, Hillebrand, et al., 2016). Generally, a multilayer network is a network of networks, which is made up of individual networks as layers that are interconnected via some relationship (a more formal introduction to multilayer networks follows in the methods section). Multilayer network approaches have

been introduced in the field of neuroscience where different layers of the network

correspond to different M/EEG frequency bands (frequency specific networks) or networks

obtained from different neuroimaging modalities (Bassett et al., 2010; Brookes et al., 2016b;

Buldú & Porter, 2018; De Domenico, 2017; Mandke et al., 2018; Tewarie, Hillebrand, et al.,

2016; Vaiana & Muldoon, 2020; Yu et al., 2017). Using frequency-based multilayer

networks, (De Domenico et al., 2016) showed that multilayer networks can better classify

between schizophrenic patients and healthy controls than single-layer or aggregated single-

layer networks. Other disease-oriented applications of multilayer networks are MEG-based

frequency-specific multilayer networks (Yu et al., 2017) and a multimodal diffusion weighted

imaging (DWI) -fMRI-MEG multilayer network (Guillon et al., 2017) to investigate

Alzheimer's disease, where multilayer network metrics of patients in the latter study

predicted individual cognitive and memory impairment.

Most complex networks – including multilayer networks – exhibit a certain level of

modularity, i.e. densely connected groups of nodes (brain regions) form clusters (modules).

Numerous neuroimaging studies have reliably shown that human brain architecture is highly

modular (Power et al., 2011; Yeo et al., 2011). Recently, (Puxeddu et al., 2021) tested

different community detection algorithms on multilayer EEG networks. Modules have been

suggested to sub-serve information processing (Sporns & Betzel, 2016). The modular

organisation of the brain begins early in life and matures during the adolescent years (Baum

et al., 2017). This development of modular organisation allows for cortical specialisation and

that reorganisation occurs with learning (Bassett et al., 2010). Accordingly, as a result of

training, brain networks in musicians may reorganise to facilitate communication between

sensory, motor and cognitive areas of the brain.

In the present methods-focused study, we investigated resting-state MEG recordings from

musicians and non-musicians obtained from the Open MEG Archive (Niso et al., 2016). The

primary aim of this study was to identify neuronal signatures of behavioural specialisation

or plasticity-induced changes following musical training in MEG resting-state networks using

multilayer network analysis. Furthermore, we also investigated changes in functional

connectivity in motor regions that have previously shown structural and functional changes

(in task-based data) following musical training (Alluri et al., 2015b; Tanaka & Kirino, 2017).

In the present study, we estimated functional connectivity using amplitude envelope

correlations between anatomically pre-defined regions. In a previous publication (Mandke

et al., 2018), we demonstrated that group comparisons using multilayer networks can be

biased by differences in link weights (e.g., by differences in average connectivity). This was

ameliorated by normalising the link weights using the singular value decomposition (SVD).

We used a seed-based connectivity analysis, an established single-layer analysis method

(Network-Based Statistics toolbox, (Zalesky et al., 2010)), and a multilayer network

approach, with layers representing MEG frequency bands. The absence of a statistically

significant difference between groups in single-layer analysis and a presence of a difference

in multilayer analysis would highlight the added value of the latter approach. Here, we

focused on identifying the modularity of resting-state networks within and between

individual MEG frequency bands ("pan-spectral" modularity). We hypothesised that the

years of musical training would have changed the modular structure of the musicians in

comparison with non-musicians, where we expected sensory, motor, and cognitive areas of

the brain to be stronger interconnected in musicians than in non-musicians, leading to less

modularity in musicians than in non-musicians. Finally, we conclude with a discussion of

some   of   the   methodological   issues   in   the   analysis   of   multilayer   networks.

Methods

Participants and MEG data pre-processing

All MEG data analysed in this study were obtained from a public access data repository,

Open MEG Archive, OMEGA (Niso et al., 2016). At the time the analysis was performed, the

OMEGA repository contained MEG data from 150 healthy controls. The repository also

includes self-report questionnaire data. Out of the questionnaire data, we used self-

reported musical experience to classify the healthy participants into two groups. We used

the 5-min resting-state MEG recordings acquired on a 275-channel CTF-MEG system (MISL,

Coquitlam, Canada) as detailed in (Niso et al., 2016). The first group included data from

participants who had no history of neuropsychiatric disorders and 5+ years of experience

playing some musical instrument (Supplementary Table 1). A total of 31 participants met

this inclusion criteria. Out of the 31 participants, 23 answered "Yes" to the questionnaire

item, "Do you consider yourself to be a musician?" and 8 answered "No". For simplicity, we

will refer to the whole group of 31 participants as musicians to reflect their 5+ years of

experience. This group was a heterogeneous sample, as it included a wide range of musical

expertise playing instruments (e.g. string instruments, percussions etc.). The second

(control) group included data from neurologically healthy OMEGA participants. We

identified 31 (age- and gender-matched) participants for our control group (non-musicians).

Further data from the total 62 participants (31 in each group, all right-handed by self-report)

are given in Table 1.

168   ***Table 1.*** *Summary of statistics for both groups.*

	Musicians	Non-musicians
Gender	M: 20; F: 11	M: 20; F: 11
Age	Mean: 28.25 years; SD: 7.25	Mean: 27.48 years; SD: 5.95
Years of experience	Mean: 13.25 years; SD: 5.04	no experience

M: male; F: female; SD: standard deviation.

The raw data included task-free, 5-mins eyes open resting-state MEG scans, sampled at

2400Hz (Niso et al., 2016). As part of our pre-processing, the data were down-sampled at

600Hz. A third order synthetic gradiometer configuration of the CTF-MEG system was

applied; a 150Hz low pass anti-aliasing filter was used. OMEGA repository also provides

participant's digitised head shape (Polhemus). Using in-house software, we performed

surface matching of the digitised head shape to an equivalent head shape extracted from

individual anatomical scans to allow for co-registration of MEG sensor geometry and brain

anatomy.

Source localisation using an atlas based beamformer

An atlas based beamforming approach (Hillebrand et al., 2012) – previously also used in

(Brookes et al., 2016b; Tewarie, Bright, et al., 2016; Tewarie, Hillebrand, et al., 2016) – was

applied for source localisation. The cortex was parcellated using the automated anatomical

labelling (AAL) (Tzourio-Mazoyer et al., 2002). Out of the 116 AAL regions, we selected 78

cortical regions that provide a full cortical coverage and ignored the remaining 38

subcortical and cerebellar regions in line with previous multilayer network studies (Brookes

et al., 2016b; Tewarie, Bright, et al., 2016; Tewarie, Hillebrand, et al., 2016). A beamformer

approach (Robinson & Vrba, 1999) was employed to generate a single time course of

electrophysiological activity within each of these regions. To achieve this, the centre of mass

was first derived for each region. Given the spatial resolution of MEG (Barnes et al., 2004),

voxels were defined on a regular 4 mm grid covering the entire region, and the

beamformer-estimated time-course of electrical activity was derived for each voxel. To

generate a single time course representing the whole region, individual voxel signals were

weighted according to their distance from the centre of mass using a Gaussian weighting

function. This procedure ensures that the regional time course is biased towards the centre

of the region, with a full width half maximum of approximately 17mm. To calculate

individual voxel time courses, a scalar beamformer was used (Robinson & Vrba, 1999).

Covariance was computed within a 0.5-150 Hz frequency window and a time window

spanning the entire experiment (300s) (Brookes et al., 2008). Regularisation was applied to

the data covariance matrix using the Tikhonov method with a regularisation parameter

equal to 5% of the maximum eigenvalue of the unregularised covariance matrix. The

forward model was based upon a dipole approximation (Sarvas, 1987) and a multiple local

sphere head model (Huang et al., 1999). Dipole orientation was determined using a non-

linear search for optimum signal to noise ratio (SNR). Beamformer time courses were sign

flipped where necessary to account for the arbitrary polarity introduced by the beamformer

source orientation estimation. This process resulted in 78 electrophysiological time courses,

each representative of a separate AAL region. This approach was applied to each subject

individually.

Functional connectivity calculation

The time courses were frequency filtered in theta (6.5-8HzHz), alpha1 (8.5-10Hz), alpha2

(10.5-12Hz), beta1 (12.5-16Hz), beta2 (16.5-20Hz), this frequency selection was motivated

by the work on training related plasticity (Klein et al., 2016; Langer et al., 2013). After

frequency filtering, to correct for signal leakage, frequency filtered time courses were

subjected to a multivariate orthogonalisation (Colclough et al., 2015). Using the Hilbert

transform, an amplitude envelope was computed by calculating the absolute value of the

analytical signal. The envelopes of all the frequency bands were down-sampled to 4Hz

(Brookes et al., 2011; Luckhoo et al., 2012). Finally, a functional connectivity matrix was

reconstructed for individual frequency bands by computing Pearson's correlation coefficient

between down-sampled envelope pairs, with each correlation coefficient forming a single

element in the weighted adjacency matrix (O'Neill et al., 2015; Tewarie, Hillebrand, et al.,

2016; van Wijk, 2014). This produced a square (78 nodes x 78 nodes) off-diagonal weighted

adjacency matrix for each frequency band under consideration.

Seed-based connectivity

Furthermore, to explore seed-based connectivity, the following regions were used – left and

right SMA (nodes 12 and 51 in AAL atlas) and left and right precentral gyrus (nodes 14 and

53 in AAL atlas) (review: (Jäncke, 2009b). A linearly constrained minimum variance

beamformer was used to project data to these regions. Following this, amplitude envelope

correlations were used to estimate functional connectivity between these regions across

participants.

Single layer analysis using Network-based statistics

To test if the effects of musical training can be detected in raw adjacency matrices, we

performed single layer analysis using the Network-Based Statistics (NBS toolbox, (Zalesky et

al., 2010). For technical details on NBS, we would like to redirect the reader to the original

article (Zalesky et al., 2010). A short summary of NBS is as follows: In a typical cross-

sectional design, NBS identifies significantly different pairwise associations between groups,

where such associations are links or connections between two nodes. A given symmetric $N \times$

N connectivity matrix has maximum $N(N-1)/2$ individual connections. Using M independent

permutations, a t-statistic is estimated for all individual connections. Then links that exceed

a certain threshold with their respective t statistic are retained, which forms the supra-

threshold network. The size of the biggest connected component found in this resulting

network is used as a test statistic. The p-value of an observed component is calculated by

finding the number of permutations for which the maximal component size exceeds the

original test statistic, normalised by M. To identify group differences using NBS, we used

M=5000 permutations with independent samples t-test ($p<0.05$) and tested a range of t-

statistic thresholds t=0.5:0.1:5.0.

Multilayer network construction

A multilayer network is a complex network structure, which is a network of networks, where

each layer is formed by an individual network. The links in such a network describe

relationships between combinations of all possible nodes and layers. In this study,

multilayer networks were constructed by integrating information from theta (6.5-8Hz),

alpha1 (8.5-10Hz), alpha2 (10.5-12Hz), beta1 (12.5-16Hz), beta2 (16.5-20Hz) frequency

bands, where each network or layer shared the same number of nodes (78 cortical regions).

The links in each layer were the amplitude envelope correlation (AEC) weights within that

frequency band. Figure 1 shows a schematic of multilayer network construction.

Mathematical representation of a multilayer network is its block adjacency matrix (De

Domenico et al., 2013; Van Mieghem, 2016). An f-layered multilayer network can be written

in the form of a block adjacency matrix (Sahneh et al., 2015) as follows:

265
$$A = \begin{bmatrix} A_1 & H_{12} & \cdots & H_{1f} \\ H_{21} & A_2 & & \vdots \\ \vdots & & \ddots & \\ H_{f1} & \cdots & & A_f \end{bmatrix}, \qquad (1)$$

where A_α corresponds to a symmetric, square adjacency matrix of a layer α, $1 \leq \alpha \leq f$. In

our case A_α corresponds to individual frequency specific networks. The matrix H_{kl}

corresponds to the coupling matrix between the layers k and l, where $1 \leq k, l \leq f, f = 5$.

The frequency specific networks, A_α, have the same dimensions for all layers (*N x N, N=78*),

implying that every layer of the network has the same number of nodes or brain regions.

To be in line with previous studies (Mandke et al., 2018; Yu et al., 2017), we chose one-to-

one between layer relationship $H_{kl} = cI$, where c is a constant and I the identity matrix

(more details on the inter-layer coupling in Supplementary material 1). Thus, we used a case

where coupling matrices are special diagonal matrices ($H_{kl} = cI$). In other words, we allow

only for links between the same nodes across all layers (i.e. we ignore cross-frequency

coupling between non-identical areas in different layers). We computed all results over a

range of values for the interlayer coupling, c=0:0.01:1.

Figure 1. *A schematic of the analysis pipeline used in the construction of multilayer networks. Here, we use a*

special case of the multilayer networks: multilayer networks with one-to-one between-layer coupling. rs:

Resting state; AEC: Amplitude envelope correlations; Theta: 6.5-8Hz; Alpha1: 8.5-10Hz; Alpha2: 10.5-12Hz;

Beta1: 12.5-16Hz; Beta2: 16.5-20Hz.

Graph analysis of multilayer networks

A multilayer network was constructed for every participant in the two groups. The absolute

value was taken of all link weights. However, comparing raw block-adjacency matrices

between groups can be biased due to differences in link weights, i.e. such a comparison can

inflate false positives or negatives (Mandke et al., 2018). These biases in the context of

multilayer networks arise due to differences in average connectivity, which can influence

the calculated network metrics (van den Heuvel et al., 2017). To correct for such biases we

used singular value decomposition (SVD) based normalisation approach (Mandke et al.,

2018). Prior to subjecting the block adjacency matrix to graph analysis, all singular values

were normalised by the largest singular value of the weighted adjacency matrix to correct

for difference in mean connectivity between groups. For a given block adjacency matrix A,

we apply a singular value decomposition $A = U \Lambda V^T$, where U and V contain the left and

right singular vectors and Λ the singular values of A. To correct for differences in average

connectivity, we rescaled Λ by the largest singular value λ_1, $\tilde{A} = U \left(\frac{10}{\lambda_1} \Lambda \right) V^T$ (the

justification of the choice of eigenvalue can be found in the Supplementary Material 2). The

rescaled matrix would thus become $\tilde{A}$. The multiplication by 10 is used to ensure that the

range of values in $\tilde{A}$ is not too small and varies between 0 and 1.

To identify community structure specific to musicians we used a community detection

algorithm. Community detection algorithms (e.g. (Blondel et al., 2008) have widely been

used to study resting-state fMRI networks and structural data (Crossley et al., 2013; Garcia

et al., 2018; Sporns & Betzel, 2016). Generally, communities in a network are obtained by

optimising a quality function, e.g. modularity (Newman, 2006), which provides an index for

within-community and between-community connections. The multilayer modularity as

expressed in (Bassett et al., 2013) is as follows:

$$Q = \frac{1}{2u_{tot}} \sum_{ijlk} \left[\left(\tilde{A}_{l,ij} - \gamma_l \frac{_ld_i \, _ld_j}{2 \sum_i \, _ld_i} \delta_{lk} \right) + \delta_{ij} H_{lk,j} \right] \delta_{g_{il},g_{jk}} \qquad (2)$$

where u_{tot} is the total link weight in the network, γ_l is the resolution parameter (set to 1,

constant over all subjects), g_{il} is the community assignment of node i in layer l, and δ_{ij} is

the Kronecker delta function (meaning that δ_{ij}=1 if i=j and δ_{ij}=0 if i≠j). Using this

community detection algorithm, we calculated individual nodal module assignments

expressed as indices. Nodes with the same index can be said to be part of the same

community. To ensure that the community organization was stable and not based on initial

settings, we repeated the algorithm 100 times at every instance.

We first performed this community detection algorithm on the multilayer matrix of each

individual subject. Using these index assignments, we calculated an agreement (consensus)

matrix (*N x N*) for every layer per group, where each element in the matrix corresponds to

the proportional number of times two given nodes were part of the same community over

all individual subjects (Baum et al., 2017). Furthermore, we obtained a collapsed agreement

matrix by averaging over all individual layer agreement matrices. This step was performed

to achieve a group-level representation of the entire multilayer network.

To compare the multilayer community structure between groups (musicians vs non-

musicians), we applied the community detection algorithm 100 times (Equation 2) to each

collapsed agreement matrix, generating 100 partitions of the 78 nodes into various

communities (Baum et al., 2017). We then repeated the process of constructing an

agreement matrix from these 100 partitions and reapplying the community detection

algorithm until achieving a stable community assignment across all 100 Louvain algorithm

repetitions. This process yielded a single community assignment for each interlayer coupling

value c. Next, we calculated the frequency with which each node was assigned to each

module across all c values.

For some of the analyses, we used the Brain Connectivity Toolbox

(https://sites.google.com/site/bctnet/, (Rubinov & Sporns, 2010)). All our used code is

publicly available at https://github.com/jilmeier/multilayer-musicians/.

Results

Functional connectivity and Network-Based Statistics

Seed-based connectivity analysis for the 4 nodes under consideration (bilateral SMA and

pre- central gyri) did not reveal any statistically significant ($p<0.05$) differences between the

two groups.

After performing SVD normalisation, for visual representation of the data, functional

connectivity estimates between all nodes for all frequency bands (theta, alpha1, alpha2,

beta1, beta2) obtained using AEC were averaged across participants (Figure 2: musicians

and Figure 3: non-musicians). Only the top 40% (i.e., the strongest) of connections were

subsequently plotted in brain space to visualise the underlying networks (see Figures 2 and

3). After computing individual connectivity matrices, we performed single layer analysis

using NBS. However, this analysis failed to show any significant effect with stringent

statistical thresholding ($p<0.05$, FDR-corrected) across frequency bands for a large range of

t-statistic thresholds. A null result using NBS is possible because of several reasons: 1) Single

layer analysis ignores the pan-spectral picture in the data, which warrants a more

integrative approach. 2) High differences in link weights can smear underlying experimental

effects and this demands some corrections as highlighted in (Mandke et al., 2018; van Wijk

et al., 2010).

Figure 2. *Averaged AEC in adjacency matrix (78 brain area labels on each axis: 1-39 left hemisphere, 40-78*

right hemisphere) across musicians (A' to E') and its corresponding source space (top 40% (i.e., the strongest) of

connections) representation (3D plots, A to E) in the 5 frequency bands under consideration. All adjacency

matrices show typical patterns of high-AEC within left hemisphere (upper left quadrant), within right

hemisphere (lower right quadrant) and between hemispheres (upper right quadrant). In the 3D plots, red lines

indicate presence of connections, with thicker lines corresponding to stronger connections. The grey circles in

the 3D plots indicate summed magnitude of link weights (node strength, weighted degree) between the node

and the rest of the brain. ROIs: Regions of interest.

Figure 3. *Averaged AEC plotted as adjacency matrices (78 brain area labels on each axis: 1-39 left hemisphere,*

40-78 right hemisphere) across non-musicians (A' to E') and their corresponding source space (top 40% (i.e., the

strongest) of connections) representation (3D plots, A to E) in 5 frequency bands under consideration. In the 3D

plots, red lines indicate presence of connections, with thicker lines corresponding to stronger connections. The

grey circles in the 3D plots indicate summed magnitude of link weights (node strength, weighted degree)

between the node and rest of the brain. ROIs: Regions of interest.

Graph analysis of the multilayer networks: Community structure

The block adjacency matrices of the two groups were subjected to community detection

algorithm after SVD normalisation. Modularity (Q) was calculated for individual subjects'

SVD normalised multilayer networks (Figure 4) and was significantly different between

groups over a large range of interlayer coupling values (*p*<0.001 for each value of

c=0:0.01:1, Wilcoxon rank-sum test). The Wilcoxon rank-sum test was conducted to

compare modularity between musicians and non-musicians. The analysis yielded statistically

significant results for all comparisons, with z-values ranging from −1.70 to - 6.76, p<0.001.

The mean effect size for the tests, calculated as rank-biserial correlation (r), was r=−0.553,

indicating a strong negative effect. Cohen's d equivalent for the mean effect size was −1.33,

further confirming a very large effect size. The negative sign indicates that the first group in

the comparisons (i.e., musicians) always had lower values. A predictive classifier based on

multilayer network feature of modularity confirmed that we can predict whether an

individual is a musician or not with above chance level (above 54%), which increased up to

100% for increasing values of c (Supplementary Figure S1). If we consider the average

modularity over all c values per individual as a classifying feature, the accuracy for

prediction was 96.78% (93.44% in a 4-fold cross-validation, more details in Supplementary

Material 3).

Figure 4: Modularity difference between musicians and non-musicians. (A) *Modularity (Q) in the two groups*

of musicians (purple) and non-musicians (green), depicted as mean values with error bars in the length of their

standard deviation over all 100 runs of the community detection algorithm and for each c in the range of

0:0.01:1. This difference was found to be statistically significant for all values of c (p<0.001 Wilcoxon rank-sum

test). **(B-C)** *Example boxplots of modularity (Q) for the two groups for c=0.02 (B) and c=0.98 (C). The boxes*

show 95% SEM and SD plotted as whiskers.

The group of musicians demonstrated different modular organisation than the non-

musicians (average adjusted Rand index over all c values: 0.6876). The group representation

of this modular organisation for both groups is shown in Figure 5. The multilayer network

community structure observed in musicians had two large communities, while non-

musicians had three communities. Musicians showed a modular structure that encompasses

both visual and motor networks, shown in yellow (Figure 5, left panel), and frontal and

temporal regions, shown in purple. In non-musicians, the modularity analysis revealed three

modules, the frontal (colour: purple), midline-temporal (colour: green) and posterior

(colour: yellow) network components. The non-musicians showed distinctly different

community structure from the group of musicians. In non-musicians, the visual and motor

networks are segregated from one another (as shown in yellow and green, respectively, see

Figure 5 right panel), whereas musicians display integrated visuo-motor networks (as shown

in yellow in left panel in Figure 5). In addition, the musicians also exhibit markedly different

patterns of connectivity spanning the fronto-temporal regions, in particular in the right

hemisphere, with integration of frontal and superior temporal regions into a single

community only in musicians (Figure 5, left panel).

Musicians   Non-musicians

Figure 5: Modular structure in musicians and non-musicians. *Within each group, regions shown in identical*

colour have been assigned to the same community (module). ***(Left panel)*** *Musicians have two modules (one*

purple, one yellow) over all values of interlayer coupling c. In both panels, the regions are coloured according to

the module to which they were most frequently assigned across all interlayer coupling values (c=0:0.1:1),

exceeding what would be expected by chance. ***(Right panel)*** *non-musicians have three modules over all*

interlayer coupling values, one frontal module (purple), one motor module (green) and one occipital module

(yellow).

We repeated the analysis without applying SVD normalisation and were able to confirm the

results obtained with SVD normalisation. Specifically, the same number of modules were

identified in the groups, and the significantly different modularity across all interlayer values

remained consistent. The module compositions differed only in a few regions, preserving

the overall interpretation of the three modules (Supplementary Figure S2).

Regarding the relation of modularity with other demographic factors, we found a significant

monotonic relationship between age and modularity (Spearman's rank correlation: r=-0.046

and $p<0.001$, over all interlayer coupling c and all runs of the community detection

algorithm), meaning that aging has a modularity-decreasing effect. We also calculated the

correlation between the age at which musicians started taking lessons and modularity and

found a significant monotonic relation (Spearman rank correlation: r=-0.0327 and p<0.001).

To further understand the link between musical training and neural connectivity findings,

we correlated years of musical training with modularity and have a significant monotonic

relationship between those two factors (Spearman's rank correlation: r=-0.0319, p<0.001).

Since we found a significant relation between age and modularity, we corrected this result

for age via partial correlation calculation (Spearman's rank correlation: r=-0.038, p<0.001).

Discussion

We studied whether an impact of musical training would be detectable in musicians' brain

function, measured with MEG during resting state. To this end, a multilayer network

framework to construct multi-frequency networks in a group of musicians and non-

musicians was used. We studied the properties of these networks using graph-theoretical

approaches. As per the hypothesis, a modular organisation in musicians spanning the

visuomotor regions and the frontal and auditory parts of the brain was found. This modular

organisation points to a stronger inter-modular connectivity, potentially underlying the

integrated organisation observed in musicians. It must be noted here that neither single

layer analysis using NBS nor seed-based connectivity analysis revealed significant group

differences. Therefore, our results highlight the added value of the multilayer analysis.

The present methods-focused study adds to the body of knowledge about brain

reorganisation in highly-trained musicians, which was so far limited to structural MRI studies

for example (Schlaug et al., 1995), (Amunts et al., 1997), (Bangert & Schlaug, 2006), and

task-based for example, (Alluri et al., 2017; Grahn & Rowe, 2009) as well as resting-state

fMRI studies for example (Luo et al., 2012, 2014; Zamorano et al., 2017). The question

whether musical training would also be reflected in electrophysiological functional networks

outside of task conditions – during resting state – has only recently been addressed, to our

knowledge in a single study so far: (Klein et al., 2016). They reported an increased intra- and

inter-hemispheric functional connectivity in resting-state EEG data of musicians in theta

(6.5-8Hz) and alpha1 (8.5-10Hz) frequency bands. However, their investigation was limited

to studying frequency specific networks in isolation. This single layer approach does not

detect pan-spectral functional networks.

The findings from our multilayer analysis underscore an important aspects about network

organisation in musicians: At a global pan-spectral level, musicians have less segregated

clusters (lower modularity), thereby allowing for an integrated network organisation over

the whole multi-frequency network. For example, (classical) musicians often need to read

sheet music, encode the visual information (alpha band network) and translate it into

signals to be executed by the motor cortex (beta band network). The between network

coordination is not limited only to visuo-motor areas, but has also been shown to span

auditory, motor and other cognitive areas, which likely facilitates higher overall network

integration (Jäncke, 2012; Proverbio et al., 2014). For instance, the coordination between

auditory and motor areas is said to be mediated via the arcuate fasciculus (Catani et al.,

2005; Catani & Mesulam, 2008). It is also plausible that this anatomical architecture

underlies the auditory-motor coupling reported in musicians (Grahn & Rowe, 2009; Jäncke,

2012).

The present study compared musicians (all with years of training in a visual-auditory-motor

skill) and non-musicians, thereby complementing earlier studies of changes in modular brain

organisation in the course of learning over weeks. In a resting-state fMRI study of

participants who learned to perform a visuo-motor task, (Bassett et al., 2015) found

segregation of visual and motor modules, increasing with the number of trials. Similarly,

during improvement of task performance, segregation of modules was observed by (Wang

et al., 2024) and found to be correlated with learning rate. In the same study, segregation

did not correlate with habit strength. Due to the relationship between habits and motor

skills (Du et al., 2022), the result reported by (Wang et al., 2024) is a possible link between

findings from learning-based studies and our result of integration rather than segregation in

musicians after years of training.

The critical difference observed in our analysis regarding between layer organization lies

primarily in the theta-beta1 and alpha1-beta1 layers. The present results observed in lower

frequency bands may reflect long-range communication (Ward, 2003). Both theta and alpha

band activity are involved in a wide range of cognitive functions (review: (Başar et al., 2001;

Ward, 2003) – such as memory and attention, which are highly active areas while playing an

instrument (Schlaug, 2015).

The relationship between behavioural specialisation and changes in the underlying neuronal

activity is not very straightforward (Zatorre et al., 2012). This problem is more pronounced

when investigating spontaneous neuronal oscillations (using M/EEG), as both increases and

decreases in activity can reflect a different brain state (Baillet, 2017). There are several

studies (Schneider et al., 2002; Shahin et al., 2003; Strait & Kraus, 2011, 2014) that have

used evoked potentials to investigate musical specialisation. However, such evidence does

not exist for resting-state data. Our results also show a significant correlation between

multilayer network measures with years of training as a behavioural variable. Modularity in

this case might not be sensitive enough to detect correlation within the musician group and

can also detect between-group differences (absence vs. presence of experience), which has

been confirmed in a predictive classifier (Supplementary Material 3). A possible solution to

circumvent this issue would be to conduct a longitudinal electrophysiological study to

establish a clearer link between behaviour and increase or decrease in the underlying

neuronal activity (using a similar approach as in the structural imaging study by (Draganski

et al., 2004)). In addition, the number of analysed group members in the musician group

(n=31) in our study is also small and could be the reason for not identifying a clear

relationship between behaviour and neural dynamics.

Brain dynamics occur across multiple spatial and temporal scales. Various neuro-imaging

techniques measure different aspects of the same underlying physiological processes, but

these data sets are often analysed in isolation. However, the multilayer network framework

has the potential to bridge this gap by taking a more integrative approach (De Domenico,

2017; Mandke et al., 2018). For example, the framework can be used to integrate data from

MEG in different frequency ranges (as in the current study), from DTI and MEG or fMRI and

MEG in the same group of participants. As a future research direction, one can also

integrate the structural network information as an additional network layer in the multilayer

framework, as e.g. investigated in (Battiston et al., 2017, 2018; Lim et al., 2019) for one

fMRI-based and an additional structural network layer for healthy controls. Recently, a

framework for including - next to structural and functional layers - also a third

morphological grey matter network layer has been proposed (Casas-Roma et al., 2022).

Therefore, the multilayer network approach allows us to study in an individual how brain

processes at different scales, from functional to structural, undergo plasticity related

changes. Our study is one of the first demonstrations that an integrative approach is useful

to study training related plasticity.

The present study is not without limitations. Firstly, the data used here were part of the

Open MEG Archive (Niso et al., 2016), which included a number of participants (musicians, 8

out of 31 participants), who played an instrument for 5+ years but did not identify

themselves as musicians. In addition, in our group of musicians, the participants did not play

the same instrument (Supplementary Table 1) nor did they have similar years of experience

(Mean: 13.25 years; SD: 5.04) playing an instrument. A future study could be designed to

replicate these findings, with a more careful selection of participants to identify a clear

relationship between resting-state network organisation and years of experience. The

neurobiological interpretation of our results should be approached with caution since a

demonstration of their reproducibility in a second independent large dataset has been left

as future work. Secondly, the MEG recordings in our study were only 5 min in duration. The

recording duration, an important aspect in capturing brain states in individual subjects

(Liuzzi et al., 2017), should be carefully considered when designing future resting-state EEG

or MEG studies. The OMEGA questionnaire data does not include more specific information

on musical training or preference of musical style. As a result, we were unable to investigate

these aspects. With more elaborate questionnaire data, one could analyse which precise

factors of musical training cause the multilayer network changes. Another limitation is the

use of self-report questionnaires in the OMEGA study, which are known to inherit multiple

biases (Schwarz, 1999). While this method is common in the field, future studies should

consider using more objective measures to quantify musical expertise such as the

short/mini-Profile of Music Perception Skills (Zentner & Strauss, 2017).

The majority of the previous studies that have reported group differences in motor cortices

of musicians in both structural and task-based functional data (see for example: (Palomar-

García et al., 2017; Schlaug, 2015; Schlaug et al., 1995; Stewart, 2008), have recruited

participants that play the same instrument professionally and have very similar levels of

expertise. A critical limitation of the present study is the heterogeneity among participants

in terms of years of experience (13.25 years ± 5.04) and a wide range of musical expertise

(string instruments, percussions etc). This heterogeneity coupled with resting-state MEG

recordings that were only 5-mins long probably make identification of seed-based

differences difficult. Hence, this methods-focused study focused on identifying global

changes in the network configuration. However, it is plausible in a future study to introduce

a task that engages both auditory and motor systems, which would likely make the

differences more apparent.

Recently, in the context of speech, (Assaneo & Poeppel, 2018) provided evidence for the

presence of an intrinsic rhythm – which mediates auditory-motor communication. It is

conceivable that such a fine-tuned mechanism might also exist in musicians, mediating the

communication between different areas and potentially leading to the formation of sub-

networks or communities. This analysis needs to be probed further by studying the

relationship between task-based data and task-free data in the same group of musicians.

To conclude, we used an open access dataset (OMEGA, (Niso et al., 2016)) to investigate the

effects of musical training related plasticity in resting-state MEG data. We studied the

frequency specific networks using NBS and by applying a multilayer network framework to

construct multi-frequency networks in musicians and non-musicians, which is a novel

framework in this context. The single layer analysis did not reveal a statistically significant

difference between the two groups. However, effects of musical training were detected in

the modularity of the network, which spanned visuo-motor and fronto-temporal areas that

are involved in execution of a musical performance. The results also indicate that musicians

show an integrated network organisation in comparison to non-musicians. Presently, the

study of musical training related plasticity lacks an integrative approach and therefore

neglects interactions between processes of plasticity occurring at different frequency bands

and/or different modalities within individuals. To overcome this limitation, we suggest the

use of the multilayer network framework.

Author contribution statement

KM: Conceptualisation, Methodology, Investigation, Software, Data curation, Formal analysis, Visualisation, Writing: original draft, review, editing.

PT: Conceptualisation, Methodology, Investigation, Software, Validation, Writing: review & editing.

PA: Methodology, Supervision.

MS: Conceptualisation, Resources, Methodology, Investigation, Project administration, Supervision, Writing: original draft, review, editing.

JM: Conceptualisation, Methodology, Investigation, Formal analysis, Verification, Supervision, Project administration, Software, Writing: original draft, review, editing.

References

Alluri, V., Brattico, E., Toiviainen, P., Burunat, I., Bogert, B., Numminen, J., & Kliuchko, M.
(2015a). Musical expertise modulates functional connectivity of limbic regions during
continuous music listening. *Psychomusicology: Music, Mind, and Brain*, *25*(4), 443–
454. https://doi.org/10.1037/pmu0000124
Alluri, V., Brattico, E., Toiviainen, P., Burunat, I., Bogert, B., Numminen, J., & Kliuchko, M.
(2015b). Musical expertise modulates functional connectivity of limbic regions
during continuous music listening. *Psychomusicology: Music, Mind, and Brain*, *25*(4),
443–454. https://doi.org/10.1037/pmu0000124
Alluri, V., Toiviainen, P., Burunat, I., Kliuchko, M., Vuust, P., & Brattico, E. (2017).
Connectivity patterns during music listening: Evidence for action-based processing in
musicians. *Human Brain Mapping*, *38*(6), 2955–2970.
https://doi.org/10.1002/hbm.23565
Amunts, K., Schlaug, G., Jäncke, L., Steinmetz, H., Schleicher, A., Dabringhaus, A., & Zilles, K.
(1997). Motor cortex and hand motor skills: Structural compliance in the human
brain. *Human Brain Mapping*, *5*(3), 206–215. https://doi.org/10.1002/(SICI)1097-
0193(1997)5:3<206::AID-HBM5>3.0.CO;2-7
Assaneo, M. F., & Poeppel, D. (2018). The coupling between auditory and motor cortices is
rate-restricted: Evidence for an intrinsic speech-motor rhythm. *Science Advances*, 1–
10. https://doi.org/10.1126/sciadv.aao3842
Baillet, S. (2017). Magnetoencephalography for brain electrophysiology and imaging. *Nature*
Neuroscience, *20*(3), 327–339. https://doi.org/10.1038/nn.4504
Bangert, M., & Schlaug, G. (2006). Specialization of the specialized in features of external
human brain morphology. *European Journal of Neuroscience*, *24*(6), 1832–1834.
https://doi.org/10.1111/j.1460-9568.2006.05031.x
Barnes, G. R., Hillebrand, A., Fawcett, I. P., & Singh, K. D. (2004). Realistic spatial sampling
for MEG beamformer images. *Human Brain Mapping*, *23*(2), 120–127.
https://doi.org/10.1002/hbm.20047
Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and
theta oscillations govern cognitive processes. *International Journal of*
Psychophysiology, *39*(2–3), 241–248. https://doi.org/10.1016/s0167-8760(00)00145-
8
Bassett, D. S., Porter, M. A., Wymbs, N. F., Grafton, S. T., Carlson, J. M., & Mucha, P. J.
(2013). Robust detection of dynamic community structure in networks. *Chaos*, *23*(1),
013142–16. https://doi.org/10.1063/1.4790830
Bassett, D. S., Wymbs, N. F., Porter, M. A., Mucha, P. J., Carlson, J. M., & Grafton, S. T.
(2010). Dynamic reconfiguration of human brain networks during learning.
Proceedings of the National Academy of Sciences, *108*(18), 7641–7646.
https://doi.org/10.1073/pnas.1018985108
Bassett, D. S., Yang, M., Wymbs, N. F., & Grafton, S. T. (2015). Learning-induced autonomy
of sensorimotor systems. *Nature Neuroscience*, *18*(5), 744–751.
https://doi.org/10.1038/nn.3993
Battiston, F., Guillon, J., Chavez, M., Latora, V., & De Vico Fallani, F. (2018). Multiplex core–
periphery organization of the human connectome. *Journal of The Royal Society*
Interface, *15*(146), 20180514. https://doi.org/10.1098/rsif.2018.0514

Battiston, F., Nicosia, V., Chavez, M., & Latora, V. (2017). Multilayer motif analysis of brain
networks. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *27*(4), 047404.
https://doi.org/10.1063/1.4979282
Baum, G. L., Ciric, R., Roalf, D. R., Betzel, R. F., Moore, T. M., Shinohara, R. T., Kahn, A. E.,
Quarmley, M., Cook, P. A., Elliot, M. A., Ruparel, K., Gur, R. E., Gur, R. C., Bassett, D.
S., & Satterthwaite, T. D. (2017). Modular segregation of structural brain networks
supports the development of executive function in youth. *Current Biology*, *27*(11),
1561–1572. https://doi.org/10.1016/j.cub.2017.04.051
Blondel, V. D., Guillaume, J.-L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of
communities in large networks. *Journal of Statistical Mechanics: Theory and*
Experiment, *2008*(10), 10008–10012. https://doi.org/10.1088/1742-
5468/2008/10/P10008
Brookes, M. J., Tewarie, P. K., Hunt, B. A. E., Robson, S. E., Gascoyne, L. E., Liddle, E. B.,
Liddle, P. F., & Morris, P. G. (2016a). A multi-layer network approach to MEG
connectivity analysis. *NeuroImage*, *132*, 425–438.
https://doi.org/10.1016/j.neuroimage.2016.02.045
Brookes, M. J., Tewarie, P. K., Hunt, B. A. E., Robson, S. E., Gascoyne, L. E., Liddle, E. B.,
Liddle, P. F., & Morris, P. G. (2016b). A multi-layer network approach to MEG
connectivity analysis. *NeuroImage*, *132*, 425–438.
https://doi.org/10.1016/j.neuroimage.2016.02.045
Brookes, M. J., Vrba, J., Robinson, S. E., Stevenson, C. M., Peters, A. M., Barnes, G. R.,
Hillebrand, A., & Morris, P. G. (2008). Optimising experimental design for MEG
beamformer imaging. *NeuroImage*, *39*(4), 1788–1802.
https://doi.org/10.1016/j.neuroimage.2007.09.050
Brookes, M. J., Woolrich, M. W., Luckhoo, H., Price, D., Hale, J. R., Stephenson, M. C.,
Barnes, G. R., Smith, S. M., & Morris, P. G. (2011). Investigating the
electrophysiological basis of resting state networks using magnetoencephalography.
Proceedings of the National Academy of Sciences, *108*(40), 16783–16788.
https://doi.org/10.1073/pnas.1112685108
Buldú, J. M., & Porter, M. A. (2018). Frequency-based brain networks: From a multiplex
framework to a full multilayer description. *Network Neuroscience*, *2*(4), 418–441.
https://doi.org/10.1162/netn_a_00033
Casas-Roma, J., Martinez-Heras, E., Solé-Ribalta, A., Solana, E., Lopez-Soley, E., Vivó, F., Diaz-
Hurtado, M., Alba-Arbalat, S., Sepulveda, M., Blanco, Y., Saiz, A., Borge-Holthoefer,
J., Llufriu, S., & Prados, F. (2022). Applying multilayer analysis to morphological,
structural, and functional brain networks to identify relevant dysfunction patterns.
Network Neuroscience, *6*(3), 916–933. https://doi.org/10.1162/netn_a_00258
Catani, M., Jones, D. K., & Ffytche, D. H. (2005). Perisylvian language networks of the human
brain. *Annals of Neurology*, *57*(1), 8–16. https://doi.org/10.1002/ana.20319
Catani, M., & Mesulam, M. (2008). The arcuate fasciculus and the disconnection theme in
language and aphasia: History and current state. *Cortex*, *44*(8), 953–961.
https://doi.org/10.1016/j.cortex.2008.04.002
Colclough, G. L., Brookes, M. J., Smith, S. M., & Woolrich, M. W. (2015). A symmetric
multivariate leakage correction for MEG connectomes. *NeuroImage*, *117*, 439–448.
https://doi.org/10.1016/j.neuroimage.2015.03.071
Crossley, N. A., Mechelli, A., Vértes, P. E., Winton-Brown, T. T., Patel, A. X., Ginestet, C. E.,
McGuire, P., & Bullmore, E. T. (2013). Cognitive relevance of the community

structure of the human brain functional coactivation network. *Proceedings of the*
National Academy of Sciences, *110*(38), 15502–15502.
https://doi.org/10.1073/pnas.1314559110

De Domenico, M. (2017). Multilayer modeling and analysis of human brain networks.
GigaScience, *6*, 1–8. https://doi.org/10.1093/gigascience/gix004

De Domenico, M., Sasai, S., & Arenas, A. (2016). Mapping Multiplex Hubs in Human
Functional Brain Networks. *Frontiers in Neuroscience*, *10*.
https://www.frontiersin.org/articles/10.3389/fnins.2016.00326

De Domenico, M., Solé-Ribalta, A., Cozzo, E., Kivelä, M., Moreno, Y., Porter, M. A., Gómez,
S., & Arenas, A. (2013). Mathematical formulation of multi-layer networks. *Physical*
Review X, *3*(4), 1–15. https://doi.org/10.1103/PhysRevX.3.041022

Draganski, B., Gaser, C., Busch, V., Schuierer, G., Bogdahn, U., & May, A. (2004). Changes in
grey matter induced by training. *Nature*, *427*(6972), Article 6972.
https://doi.org/10.1038/427311a

Du, Y., Krakauer, J. W., & Haith, A. M. (2022). The relationship between habits and motor
skills in humans. *Trends in Cognitive Sciences*, *26*(5), 371–387.
https://doi.org/10.1016/j.tics.2022.02.002

Elbert, T., Pantev, C., Wienbruch, C., Rockstroh, B., & Taub, E. (1995). Increased Cortical
Representation of the Fingers of the Left Hand in String Players. *Science*, *270*(5234),
305–307. https://doi.org/10.1126/science.270.5234.305

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through
neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480.
https://doi.org/10.1016/j.tics.2005.08.011

Friston, K. J. (2011). Functional and Effective Connectivity: A Review. *Brain Connectivity*,
1(1), 13–36. https://doi.org/10.1089/brain.2011.0008

Garcia, J. O., Ashourvan, A., Muldoon, S., Vettel, J. M., & Bassett, D. S. (2018). Applications
of Community Detection Techniques to Brain Graphs: Algorithmic Considerations
and Implications for Neural Function. *Proceedings of the IEEE*, *106*(5), 846–867.
https://doi.org/10.1109/JPROC.2017.2786710

Gaser, C., & Schlaug, G. (2003). Brain structures differ between musicians and non-
musicians. *The Journal of Neuroscience*, *23*(27), 9240–9245.
https://doi.org/10.1523/JNEUROSCI.23-27-09240.2003

Grahn, J. A., & Rowe, J. B. (2009). Feeling the Beat: Premotor and Striatal Interactions in
Musicians and Nonmusicians during Beat Perception. *Journal of Neuroscience*,
29(23), 7540–7548. https://doi.org/10.1523/JNEUROSCI.2018-08.2009

Guillon, J., Attal, Y., Colliot, O., La Corte, V., Dubois, B., Schwartz, D., Chavez, M., & De Vico
Fallani, F. (2017). Loss of brain inter-frequency hubs in Alzheimer's disease. *Scientific*
Reports, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-07846-w

Herholz, S. C., & Zatorre, R. J. (2012). Musical Training as a Framework for Brain Plasticity:
Behavior, Function, and Structure. *Neuron*, *76*(3), 486–502.
https://doi.org/10.1016/j.neuron.2012.10.011

Hillebrand, A., Barnes, G. R., Bosboom, J. L., Berendse, H. W., & Stam, C. J. (2012).
Frequency-dependent functional connectivity within resting-state networks: An
atlas-based MEG beamformer solution. *NeuroImage*, *59*(4), 3909–3921.
https://doi.org/10.1016/j.neuroimage.2011.11.005

Huang, M.-X., Mosher, J. C., & Leahy, R. M. (1999). A sensor-weighted overlapping-sphere
742   head model and exhaustive head model comparison for MEG. *Physics in Medicine*
743   *and Biology*, *44*(2), 423–440. https://doi.org/10.1088/0031-9155/44/2/010

744   Jäncke, L. (2009a). The Architecture of the Golfer's Brain. *PLoS ONE*, *101*(11), 1435–1439.
745   https://doi.org/10.1371/journal.pone.0004785

746   Jäncke, L. (2009b). The plastic human brain. *Restorative Neurology and Neuroscience*, *27*(5),
747   521–538. https://doi.org/10.3233/RNN-2009-0519

748   Jäncke, L. (2012). The dynamic audio-motor system in pianists. *Annals of the New York*
749   *Academy of Sciences*, *1252*(1), 246–252. https://doi.org/10.1111/j.1749-
750   6632.2011.06416.x

751   Klein, C., Liem, F., Hänggi, J., Elmer, S., & Jäncke, L. (2016). The "silent" imprint of musical
752   training. *Human Brain Mapping*, *37*(2), 536–546.
753   https://doi.org/10.1002/hbm.23045

754   Langer, N., von Bastian, C. C., Wirz, H., Oberauer, K., & Jäncke, L. (2013). The effects of
755   working memory training on functional brain network efficiency. *Cortex*, *49*(9),
756   2424–2438. https://doi.org/10.1016/j.cortex.2013.01.008

757   Lim, S., Radicchi, F., van den Heuvel, M. P., & Sporns, O. (2019). Discordant attributes of
758   structural and functional brain connectivity in a two-layer multiplex network.
759   *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-39243-w

760   Liuzzi, L., Gascoyne, L. E., Tewarie, P. K., Barratt, E. L., Boto, E., & Brookes, M. J. (2017).
761   Optimising experimental design for MEG resting state functional connectivity
762   measurement. *NeuroImage*, *155*(November 2016), 565–576.
763   https://doi.org/10.1016/j.neuroimage.2016.11.064

764   Luckhoo, H., Hale, J. R., Stokes, M. G., Nobre, A. C., Morris, P. G., Brookes, M. J., & Woolrich,
765   M. W. (2012). Inferring task-related networks using independent component analysis
766   in magnetoencephalography. *NeuroImage*, *62*(1), 530–541.
767   https://doi.org/10.1016/j.neuroimage.2012.04.046

768   Luo, C., Guo, Z.-W., Lai, Y., Liao, W., Liu, Q., Kendrick, K. M., Yao, D., & Li, H. (2012). Musical
769   training induces functional plasticity in perceptual and motor networks: Insights
770   from resting-state fMRI. *PLoS ONE*, *7*(5), 1–10.
771   https://doi.org/10.1371/journal.pone.0036568

772   Luo, C., Tu, S., Peng, Y., Gao, S., Li, J., Dong, L., Li, G., Lai, Y., Li, H., & Yao, D. (2014). Long-
773   term effects of musical training and functional plasticity in salience system. *Neural*
774   *Plasticity*, 1–14. https://doi.org/10.1155/2014/180138

775   Maguire, E. A., Gadian, D. G., Johnsrude, I. S., Good, C. D., Ashburner, J., Frackowiak, R. S. J.,
776   & Frith, C. D. (2000). Navigation-related structural change in the hippocampi of taxi
777   drivers. *Proceedings of the National Academy of Sciences*, *97*(8), 4398–4403.
778   https://doi.org/10.1073/pnas.070039597

779   Mandke, K., Meier, J. M., Brookes, M. J., O'Dea, R. D., Van Mieghem, P., Stam, C. J.,
780   Hillebrand, A., & Tewarie, P. K. (2018). Comparing multilayer brain networks
781   between groups: Introducing graph metrics and recommendations. *NeuroImage*,
782   *166*, 371–384. https://doi.org/10.1016/j.neuroimage.2017.11.016

783   Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of*
784   *the National Academy of Sciences*, *103*(23), 8577–8582.
785   https://doi.org/10.1073/pnas.0601602103

Niso, G., Rogers, C., Moreau, J. T., Chen, L. Y., Madjar, C., Das, S., Bock, E., Tadel, F., Evans, A.
C., Jolicoeur, P., & Baillet, S. (2016). OMEGA: The Open MEG Archive. *NeuroImage*,
124, 1182–1187. https://doi.org/10.1016/j.neuroimage.2015.04.028
O'Neill, G. C., Barratt, E. L., Hunt, B. A. E., Tewarie, P. K., & Brookes, M. J. (2015). Measuring
electrophysiological connectivity by power envelope correlation: A technical review
on MEG methods. *Physics in Medicine and Biology*, *60*(21), R271–R295.
https://doi.org/10.1088/0031-9155/60/21/R271
Palomar-García, M. Á., Zatorre, R. J., Ventura-Campos, N., Bueicheků, E., & Ávila, C. (2017).
Modulation of Functional Connectivity in Auditory-Motor Networks in Musicians
Compared with Nonmusicians. *Cerebral Cortex*, *27*(5), 2768–2778.
https://doi.org/10.1093/cercor/bhw120
Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C.,
Laumann, T. O., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2011). Functional
Network Organization of the Human Brain. *Neuron*, *72*(4), 665–678.
https://doi.org/10.1016/j.neuron.2011.09.006
Proverbio, A. M., Calbi, M., Manfredi, M., & Zani, A. (2014). Audio-visuomotor processing in
the Musician's brain: An ERP study on professional violinists and clarinetists.
Scientific Reports, *4*, 5866. https://doi.org/10.1038/srep05866
Puxeddu, M. G., Petti, M., & Astolfi, L. (2021). A Comprehensive Analysis of Multilayer
Community Detection Algorithms for Application to EEG-Based Brain Networks.
Frontiers in Systems Neuroscience, *15*.
https://www.frontiersin.org/articles/10.3389/fnsys.2021.624183
Robinson, S. E., & Vrba, J. (1999). Functional neuroimaging by synthetic aperture
magnetometry. *Recent Advances in Biomagnetism*.
Rubinov, M., & Sporns, O. (2010). Complex network measures of brain connectivity: Uses
and interpretations. *NeuroImage*, *52*(3), 1059–1069.
https://doi.org/10.1016/j.neuroimage.2009.10.003
Sahneh, F. D., Scoglio, C., & Van Mieghem, P. (2015). Exact coupling threshold for structural
transition reveals diversified behaviors in interconnected networks. *Physical Review
E - Statistical, Nonlinear, and Soft Matter Physics*, *92*(4), 1–5.
https://doi.org/10.1103/PhysRevE.92.040801
Sarvas, J. (1987). Basic mathematical and electromagnetic concepts of the biomagnetic
inverse problem. *Physics in Medicine and Biology*, *32*(1), 11–22.
https://doi.org/10.1088/0031-9155/32/1/004
Schlaug, G. (2015). Musicians and music making as a model for the study of brain plasticity.
Progress in Brain Research, *217*, 37–55. https://doi.org/10.1016/bs.pbr.2014.11.020
Schlaug, G., Jäncke, L., Huang, Y., Staiger, J. F., & Steinmetz, H. (1995). Increased corpus
callosum size in musicians. *Neuropsychologia*, *33*(8), 1047–1055.
https://doi.org/10.1016/0028-3932(95)00045-5
Schneider, P., Scherg, M., Dosch, H. G., Specht, H. J., Gutschalk, A., & Rupp, A. (2002).
Morphology of Heschl's gyrus reflects enhanced activation in the auditory cortex of
musicians. *Nature Neuroscience*, *5*(7), 688–694. https://doi.org/10.1038/nn871
Scholz, J., Klein, M. C., Behrens, T. E. J., & Johansen-Berg, H. (2010). Training induces
changes in white matter architecture. *Nature Neuroscience*, *12*(11), 1370–1371.
https://doi.org/10.1038/nn.2412
Schwarz, N. (1999). Self-reports: How the questions shape the answers. *American
Psychologist*, *54*(2), 93–105. https://doi.org/10.1037/0003-066X.54.2.93

Shahin, A., Bosnyak, D. J., Trainor, L. J., & Roberts, L. E. (2003). Enhancement of Neuroplastic
P2 and N1c Auditory Evoked Potentials in Musicians. *Journal of Neuroscience*, *23*(13),
5545–5552. https://doi.org/10.1523/JNEUROSCI.23-13-05545.2003
Smit, D. J. A., Stam, C. J., Posthuma, D., Boomsma, D. I., & De Geus, E. J. C. (2008).
Heritability of "small-world" networks in the brain: A graph theoretical analysis of
resting-state EEG functional connectivity. *Human Brain Mapping*, *29*(12), 1368–1378.
https://doi.org/10.1002/hbm.20468
Sporns, O., & Betzel, R. F. (2016). Modular brain networks. *Annual Review of Psychology*,
67(1), 613–640. https://doi.org/10.1146/annurev-psych-122414-033634
Stam, C. J., Montez, T., Jones, B. F., Rombouts, S. A. R. B., Van Der Made, Y., Pijnenburg, Y. A.
L., & Scheltens, P. (2005). Disturbed fluctuations of resting state EEG synchronization
in Alzheimer's disease. *Clinical Neurophysiology*, *116*(3), 708–715.
https://doi.org/10.1016/j.clinph.2004.09.022
Stewart, L. (2008). Do musicians have different brains? *Clinical Medicine*, *8*(3), 304–308.
https://doi.org/10.7861/clinmedicine.8-3-304
Strait, D. L., & Kraus, N. (2011). Can you hear me now? Musical training shapes functional
brain networks for selective auditory attention and hearing speech in noise. *Frontiers*
in Psychology, *2*, 1–10. https://doi.org/10.3389/fpsyg.2011.00113
Strait, D. L., & Kraus, N. (2014). Biological impact of auditory expertise across the life span:
Musicians as a model of auditory learning. *Hearing Research*, *308*, 109–121.
https://doi.org/10.1016/j.heares.2013.08.004
Tanaka, S., & Kirino, E. (2017). Reorganization of the thalamocortical network in musicians.
Brain Research, *1664*, 48–54. https://doi.org/10.1016/j.brainres.2017.03.027
Tewarie, P. K., Bright, M. G., Hillebrand, A., Robson, S. E., Gascoyne, L. E., Morris, P. G.,
Meier, J. M., Van Mieghem, P., & Brookes, M. J. (2016). Predicting haemodynamic
networks using electrophysiology: The role of non-linear and cross-frequency
interactions. *NeuroImage*, *130*, 273–292.
https://doi.org/10.1016/j.neuroimage.2016.01.053
Tewarie, P. K., Hillebrand, A., van Dijk, B. W., Stam, C. J., O'Neill, G. C., Van Mieghem, P.,
Meier, J. M., Woolrich, M. W., Morris, P. G., & Brookes, M. J. (2016). Integrating
cross-frequency and within band functional networks in resting-state MEG: A multi-
layer network approach. *NeuroImage*, *142*, 324–336.
https://doi.org/10.1016/j.neuroimage.2016.07.057
Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N.,
Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in
SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject
brain. *NeuroImage*, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978
Vaiana, M., & Muldoon, S. F. (2020). Multilayer Brain Networks. *Journal of Nonlinear*
Science, *30*(5), 2147–2169. https://doi.org/10.1007/s00332-017-9436-8
van den Heuvel, M., de Lange, S. C., Zalesky, A., Seguin, C., Yeo, T. B. T., & Schmidt, R.
(2017). Proportional thresholding in resting-state fMRI functional connectivity
networks and consequences for patient-control connectome studies: Issues and
recommendations. *NeuroImage*, *152*, 437–449.
https://doi.org/10.1016/j.neuroimage.2017.02.005
Van Mieghem, P. (2016). Interconnectivity structure of a general interdependent network.
Physical Review E, *93*(4), 1–11. https://doi.org/10.1103/PhysRevE.93.042305

van Wijk, B. C. M. (2014). Thalamo-cortical cross-frequency coupling detected with MEG.
Frontiers in Human Neuroscience, *15*(6), 884–890. https://doi.org/10.1038/nn.3101

van Wijk, B. C. M., Stam, C. J., & Daffertshofer, A. (2010). Comparing brain networks of
different size and connectivity density using graph theory. *PLoS ONE*, *5*(10).
https://doi.org/10.1371/journal.pone.0013701

Wang, X., Zwosta, K., Hennig, J., Böhm, I., Ehrlich, S., Wolfensteller, U., & Ruge, H. (2024).
The dynamics of functional brain network segregation in feedback-driven learning.
Communications Biology, *7*(1), 531. https://doi.org/10.1038/s42003-024-06210-9

Ward, L. M. (2003). Synchronous neural oscillations and cognitive processes. *Trends in
Cognitive Sciences*, *7*(12), 553–559. https://doi.org/10.1016/j.tics.2003.10.012

Yeo, T. B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M.,
Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fisch, B., Liu, H., & Buckner, R.
L. (2011). The organization of the human cerebral cortex estimated by intrinsic
functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165.
https://doi.org/10.1152/jn.00338.2011

Yu, M., Engels, M. M. A., Hillebrand, A., Van Straaten, E. C. W., Gouw, A. A., Teunissen, C.,
Van Der Flier, W. M., Scheltens, P., & Stam, C. J. (2017). Selective impairment of
hippocampus and posterior hub areas in Alzheimer's disease: An MEG-based
multiplex network study. *Brain*, *140*(5), 1466–1485.
https://doi.org/10.1093/brain/awx050

Zalesky, A., Fornito, A., & Bullmore, E. T. (2010). Network-based statistic: Identifying
differences in brain networks. *NeuroImage*, *53*(4), 1197–1207.
https://doi.org/10.1016/j.neuroimage.2010.06.041

Zamorano, A. M., Cifre, I., Montoya, P., Riquelme, I., & Kleber, B. (2017). Insula-based
networks in professional musicians: Evidence for increased functional connectivity
during resting state fMRI. *Human Brain Mapping*, *38*(10), 4834–4849.
https://doi.org/10.1002/hbm.23682

Zatorre, R. J., Fields, R. D., & Johansen-Berg, H. (2012). Plasticity in gray and white:
Neuroimaging changes in brain structure during learning. *Nature Neuroscience*,
15(4), 528–536. https://doi.org/10.1038/nn.3045

Zatorre, R. J., & McGill, J. (2005). Music, the food of neuroscience? *Nature*, *434*(7031), 312–
315. https://doi.org/10.1038/434312a

Zentner, M., & Strauss, H. (2017). Assessing musical ability quickly and objectively:
Development and validation of the Short-PROMS and the Mini-PROMS. *Annals of the
New York Academy of Sciences*, *1400*(1), 33–45. https://doi.org/10.1111/nyas.13410