Musicians' brains at rest: Multilayer network analysis of MEG data

Kanad Mandke[a,b], Prejaas Tewarie[c], Peyman Adjamian[d], Martin Schürmann[a,*], Jil Meier[e,f,g,*]

[a] School of Psychology, University of Nottingham, Nottingham, United Kingdom

[b] Centre for Neuroscience in Education, Department of Psychology, University of Cambridge

[b] Sir Peter Mansfield Imaging Centre, School of Physics and Astronomy, University of Nottingham, Nottingham, United Kingdom

[d] MRC, Institute of Hearing Research, University of Nottingham, Nottingham, United Kingdom

[e] Department of Neurology, UMC Utrecht Brain Center, University Medical Center Utrecht, Utrecht, The Netherlands

[f] Current address: Berlin Institute of Health, Charité – Universitätsmedizin Berlin, Berlin, Germany

[g] Current address: Department of Neurology with Experimental Neurology, Brain Simulation Section, Charité – Universitätsmedizin Berlin, corporate member of Freie Universität Berlin and Humboldt-Universität zu Berlin, Berlin, Germany

Page count: 35

Word count: 6305 (excluding references)

Figures: 5

* J.M. and M.S. contributed equally to this work

Abstract

The ability to proficiently play a musical instrument requires a fine-grained synchronisation between several sensorimotor and cognitive brain regions. Previous studies have demonstrated that the brain undergoes functional changes with intensive musical training, which has also been identified in resting-state data. These studies have investigated changes in brain networks using fMRI or analysing electrophysiological frequency-specific networks in

isolation (i.e., neural oscillatory networks). While the analysis of such "mono-layer" networks has proven useful, it fails to capture the complexities of multiple interacting networks. To this end, we applied a multilayer network framework for analysing publicly available data (Open MEG Archive) obtained with magnetoencephalography (MEG). We investigated resting-state differences between participants with musical training (n=31) and those without (n=31). While single-layer Network-Based Statistics analysis did not demonstrate any group differences, our multilayer analysis revealed that musicians show a modular organisation that spans visuomotor and frontotemporal areas, which are known to be involved in the execution of a musical performance. This twofold modular structure was found to be significantly different from non-musicians. The differences between the two groups are primarily seen in the theta (6.5-8Hz), alpha1 (8.5-10Hz) and beta1 (12.5-16Hz) frequency bands.

No statistically significant relationships were found between self-reported measures of musical training and network properties, which could be attributed to the heterogeneity of the dataset. However, we demonstrate that the novel analysis method provides additional information that single-layer analysis methods cannot. Overall, the multilayer network method provides a unique opportunity to explore the pan-spectral nature of oscillatory networks, with studies of brain plasticity in musicians as a potential future application of the novel method.

Word count: 263

Keywords: Functional connectivity, resting state, multilayer networks, complex networks, graph theory, musicians, modularity

Introduction

Musical proficiency is a commonly used model to study brain plasticity (Herholz & Zatorre, 2012; Jäncke, 2009b; Zatorre & McGill, 2005). This research is largely motivated by the fact that achieving high levels of proficiency in any given musical instrument requires years of focused training. Numerous studies have reported structural brain changes due to musical training (Amunts et al., 1997; Elbert et al., 1995; Schlaug et al., 1995). For example, an increased size of the anterior half of the corpus callosum (Schlaug et al., 1995), and differences in the cortical representation of the hand motor area (Amunts et al., 1997; Elbert et al., 1995). More specifically, the brain also undergoes anatomical plasticity that pertains to the demands posed by the particular type of musical training, i.e., pianists showed a left hemisphere dominance, whereas violinists showed right hemisphere advantage (Bangert & Schlaug, 2006). Robust changes in grey matter specific to auditory, motor and visuo-spatial regions have also been reported (Gaser & Schlaug, 2003). This anatomical specialisation is potentially the result of years of musical training that requires fine-grained communication between several cortical areas. For example, musicians showed an increased coupling between auditory and pre-motor areas in functional magnetic resonance imaging (fMRI) (Grahn & Rowe, 2009), related to musicians' skilled actions during a performance.

It is worth noting that the training-related plastic changes are not unique to musicians but can be seen in groups of subjects with some degree of behavioural specialisation. For instance, training-related white matter changes have been observed in healthy adults learning to juggle (Scholz et al., 2010), professional golfers (Jäncke, 2009a), and London taxi drivers (Maguire et al., 2000).

To better characterise relationships between anatomical regions from task-free data, resting-state functional connectivity approaches have proven useful. It must be noted that either a seed-based or whole-brain functional connectivity analysis can inform about the strengths of these connections (Friston, 2011). Going a step further, studying the pattern of these connections can give us more information about the functional network as a whole. Using a seed-based connectivity approach to resting-state fMRI in musicians, demonstrated an increased functional connectivity in auditory, visual and motor areas. Similarly, (Tanaka & Kirino, 2017) demonstrated an increased functional connectivity in the supplementary motor area (SMA) network in musicians imagining musical performances based on fMRI data. Using

a whole-brain analysis of fMRI, (Alluri et al., 2017) reported that musicians recruit action-based brain networks, while non-musicians recruit perception-based networks in a naturalistic listening paradigm. In a seed-based fMRI study by (Alluri et al., 2015a), musicians demonstrated an increased connectivity with SMA and with ventromedial and ventrolateral cerebral and cerebellar affective regions while listening to music, whereas non-musicians displayed a higher connectivity only with subcortical regions. In a resting-state fMRI study, (Luo et al., 2014) demonstrated a greater local functional connectivity in musicians compared to non-musicians in the bilateral dorsal anterior cingulate cortex, anterior insula, and anterior temporoparietal junction. In a similar study, increased insular connectivity in musicians was reported (Zamorano et al., 2017). As reviewed above, evidence of functional specialisation in musicians comes from structural data (white or grey matter changes) or haemodynamic functional data. Little attention has been paid so far to the analysis of electrophysiological magnetoencephalography (MEG) data recorded from musicians' brains during resting state, capturing spontaneous, non-task-related brain activity, which could offer insights into the brain activity of musicians and the communication between brain areas on shorter temporal time scales. Activity recorded by MEG predominantly consists of neuronal oscillations, which occur over a wide range of temporal scales (1-200Hz). Numerous studies have shown that these oscillations are involved in mediating both long-range and local communication in the brain (e.g., (Fries, 2005)).

Analysis of resting-state networks is not limited only to functional magnetic resonance imaging (fMRI) but has also been a significant part of the human electrophysiological research based on MEG (Brookes et al., 2011; Hillebrand et al., 2012) and electroencephalography (EEG) (Smit et al., 2008; Stam et al., 2005). However, to the best of our knowledge, only one recent study (Klein et al., 2016) has explored resting-state differences between musicians and non-musicians using an electrophysiological approach. The authors analysed high-density EEG resting-state data obtained from musicians with graph-theoretical approaches. They reported that musicians showed an increased intra- and inter-hemispheric functional connectivity between auditory, sensorimotor, and prefrontal cortex in theta (6.5-8Hz), alpha1 (8.5-10Hz), and alpha2 (10.5-12Hz) frequency ranges. A limitation of this study is that separate frequency specific networks were studied in isolation (also referred to as analysis of single layers), which ignores the pan-spectral picture of ongoing brain connectivity.

To overcome this limitation, the present study used a novel multilayer network framework, combining data obtained from multiple frequency bands in one single network description (Brookes et al., 2016a; Mandke et al., 2018; Tewarie, Hillebrand, van Dijk, et al., 2016). Generally, a multilayer network is a network of networks, which is made up of individual networks as layers that are interconnected via some relationship (a more formal introduction to multilayer networks follows in the methods section). Multilayer network approaches have recently been introduced in the field of neuroscience where different layers of the network correspond to different M/EEG frequency bands (frequency specific networks) or networks obtained from different neuroimaging modalities (Bassett et al., 2010; Brookes et al., 2016b; Buldú & Porter, 2018; De Domenico, 2017; Mandke et al., 2018; Tewarie, Hillebrand, van Dijk, et al., 2016; Vaiana & Muldoon, 2020; Yu et al., 2017). Using frequency-based multilayer networks, (De Domenico et al., 2016) showed that multilayer networks can better classify between schizophrenic patients and healthy controls than single-layer or aggregated single-layer networks. Other disease-oriented applications of multilayer networks are MEG-based frequency-specific multilayer networks (Yu et al., 2017) and a multimodal diffusion weighted imaging (DWI) -fMRI-MEG multilayer network (Guillon et al., 2017) to investigate Alzheimer's disease, where multilayer network metrics of patients in the latter study predicted individual cognitive and memory impairment.

Most complex networks – including multilayer networks – exhibit a certain level of modularity, i.e. densely connected groups of nodes (brain regions) form clusters (modules). Numerous neuroimaging studies have reliably shown that human brain architecture is highly modular (Power et al., 2011; Yeo et al., 2011). Recently, (Puxeddu et al., 2021) tested different community detection algorithms on multilayer EEG networks. Modules have been suggested to sub-serve information processing (Sporns & Betzel, 2016). The modular organisation of the brain begins early in life and matures during the adolescent years (Baum et al., 2017). This development of modular organisation allows for cortical specialisation and that reorganisation occurs with learning (Bassett et al., 2010). Accordingly, as a result of training brain networks in musicians, may reorganise to facilitate communication between sensory, motor and cognitive areas of the brain.

In the present methods-focused study, we investigated resting-state MEG recordings from musicians and non-musicians obtained from the Open MEG Archive (Niso et al., 2016). The primary aim of this study was to identify neuronal signatures of behavioural specialisation or

plasticity-induced changes following musical training in MEG resting-state networks using multilayer network analysis. Furthermore, we also investigated changes in functional connectivity in motor regions that have previously shown structural and functional changes (in task-based data) following musical training (Alluri et al., 2015b; Tanaka & Kirino, 2017). In the present study, we estimated functional connectivity using amplitude envelope correlations between anatomically pre-defined regions. In a previous publication (Mandke et al., 2018), we demonstrated that group comparisons using multilayer networks can be biased by differences in link weights (e.g., by differences in average connectivity). This was ameliorated by normalising the link weights using the singular value decomposition (SVD).

We used a seed-based connectivity analysis, an established single-layer analysis method (Network-Based Statistics toolbox, (Zalesky et al., 2010)), and a multilayer network approach, with layers representing MEG frequency bands. The absence of a statistically significant difference between groups in single-layer analysis and a presence of a difference in multilayer analysis would highlight the added value of the latter approach. Here, we focused on identifying the modularity of resting-state networks within and between individual MEG frequency bands ("pan-spectral" modularity). We hypothesised that the years of musical training would have changed the modular structure of the musicians in comparison with non-musicians, where we expected sensory, motor, and cognitive areas of the brain to be stronger interconnected in musicians than in non-musicians, leading to less modularity in musicians than in non-musicians. Finally, we conclude with a discussion of some of the methodological issues in the analysis of multilayer networks.

Methods

Participants and MEG data pre-processing

All MEG data analysed in this study were obtained from a public access data repository, Open MEG Archive, OMEGA (Niso et al., 2016). At the time the analysis was performed, the OMEGA repository contained MEG data from 150 healthy controls. The repository also includes self-report questionnaire data. Out of the questionnaire data, we used self-reported musical experience to classify the healthy participants into two groups. We used the 5-min resting-state MEG recordings acquired on a 275-channel CTF-MEG system (MISL, Coquitlam, Canada) as detailed in (Niso et al., 2016).

The first group included data from participants who had no history of neuropsychiatric disorders and 5+ years of experience playing some musical instrument. Out of this group, we excluded individuals whose structural MRI data were not suitable for the procedure of co-registration with MEG. We further excluded subjects with unusual sensor-level power spectra (Gross et al., 2013). The 31 subjects that passed our quality checks formed our experimental group (musicians). Out of the 31 musicians, 23 were professional musicians and 8 can be considered amateur musicians. This group was a heterogeneous sample, as it included a wide range of musical expertise playing instruments (e.g. string instruments, percussions etc.).

The second (control) group included data from neurologically healthy OMEGA participants who did not play any musical instrument. Using the same exclusion criteria as for the musicians' group, we identified 31 (age- and gender-matched) participants for our control group (non-musicians). Further data from the altogether 62 participants (31 in each group, all right-handed by self-report) are given in Table 1.

Table 1. Summary of statistics for both groups.

	Musicians	Non-musicians
Gender	M: 20; F: 11	M: 20; F: 11
Age	Mean: 28.25 years; SD: 7.25	Mean: 27.48 years; SD: 5.95
Years of experience	Mean: 13.25 years; SD: 5.04	no experience

M: male; F: female; SD: standard deviation.

The raw data included task-free, 5-mins eyes open resting-state MEG scans, sampled at 2400Hz (Niso et al., 2016). As part of our pre-processing, the data were down-sampled at 600Hz. A third order synthetic gradiometer configuration of the CTF-MEG system was applied; a 150Hz low pass anti-aliasing filter was used. OMEGA repository also provides participant's digitised head shape (Polhemus). Using in-house software, we performed surface matching of the digitised head shape to an equivalent head shape extracted from individual anatomical scans to allow for co-registration of MEG sensor geometry and brain anatomy.

Source localisation using an atlas based beamformer

An atlas based beamforming approach (Hillebrand et al., 2012) – previously also used in (Brookes et al., 2016b; Tewarie, Bright, et al., 2016; Tewarie, Hillebrand, van Dijk, et al., 2016) – was applied for source localisation. The cortex was parcellated using the automated anatomical labelling (AAL) (Tzourio-Mazoyer et al., 2002). Out of the 116 AAL regions, we selected 78 cortical regions that provide a full cortical coverage and ignored the remaining 38 subcortical and cerebellar regions in line with previous multilayer network studies (Brookes et al., 2016b; Tewarie, Bright, et al., 2016; Tewarie, Hillebrand, van Dijk, et al., 2016). A beamformer approach (Robinson & Vrba, 1999) was employed to generate a single time course of electrophysiological activity within each of these regions. To achieve this, the centre of mass was first derived for each region. Given the spatial resolution of MEG (Barnes et al., 2004), voxels were defined on a regular 4 mm grid covering the entire region, and the beamformer-estimated time-course of electrical activity was derived for each voxel. To generate a single time course representing the whole region, individual voxel signals were weighted according to their distance from the centre of mass using a Gaussian weighting function. This procedure ensures that the regional time course is biased towards the centre of the region, with a full width half maximum of approximately 17mm. To calculate individual voxel time courses, a scalar beamformer was used (Robinson & Vrba, 1999). Covariance was computed within a 0.5-150 Hz frequency window and a time window spanning the entire experiment (300s) (Brookes et al., 2008). Regularisation was applied to the data covariance matrix using the Tikhonov method with a regularisation parameter equal to 5% of the maximum eigenvalue of the unregularised covariance matrix. The forward model was based upon a dipole approximation (Sarvas, 1987) and a multiple local sphere head model (Huang et al., 1999). Dipole orientation was determined using a non-linear search for optimum signal

to noise ratio (SNR). Beamformer time courses were sign flipped where necessary to account for the arbitrary polarity introduced by the beamformer source orientation estimation. This process resulted in 78 electrophysiological time courses, each representative of a separate AAL region. This approach was applied to each subject individually.

Functional connectivity calculation

The time courses were frequency filtered in theta (6.5-8HzHz), alpha1 (8.5-10Hz), alpha2 (10.5-12Hz), beta1 (12.5-16Hz), beta2 (16.5-20Hz), this frequency selection was motivated by the work on training related plasticity (Klein et al., 2016; Langer et al., 2013). After frequency filtering, to correct for signal leakage, frequency filtered time courses were subjected to a multivariate orthogonalisation (Colclough et al., 2015). Using the Hilbert transform, an amplitude envelope was computed by calculating the absolute value of the analytical signal. The envelopes of all the frequency bands were down-sampled to 4Hz (Brookes et al., 2011; Luckhoo et al., 2012). Finally, a functional connectivity matrix was reconstructed for individual frequency bands by computing Pearson's correlation coefficient between down-sampled envelope pairs, with each correlation coefficient forming a single element in the weighted adjacency matrix (O'Neill et al., 2015; Tewarie, Hillebrand, van Dijk, et al., 2016; van Wijk, 2014). This produced a square (78 nodes x 78 nodes) off-diagonal weighted adjacency matrix for each frequency band under consideration.

Seed-based connectivity

Furthermore, to explore seed-based connectivity, the following regions were used – left and right SMA (nodes 12 and 51 in AAL atlas) and left and right precentral gyrus (nodes 14 and 53 in AAL atlas) (review: (Jäncke, 2009b). A linearly constrained minimum variance beamformer was used to project data to these regions. Following this, amplitude envelope correlations were used to estimate functional connectivity between these regions across participants.

Single layer analysis using Network-based statistics

To test if the effects of musical training can be detected in raw adjacency matrices, we performed single layer analysis using the Network-Based Statistics (NBS toolbox, (Zalesky et al., 2010). For technical details on NBS, we would like to redirect the reader to the original article (Zalesky et al., 2010). A short summary of NBS is as follows: In a typical cross-sectional

design, NBS identifies significantly different pairwise associations between groups, where such associations are links or connections between two nodes. A given symmetric *N x N* connectivity matrix has maximum *N(N-1)/2* individual connections. Using *M* independent permutations, a t-statistic is estimated for all individual connections. Then links that exceed a certain threshold with their respective *t* statistic are retained, which forms the supra-threshold network. The size of the biggest connected component found in this resulting network is used as a test statistic. The *p*-value of an observed component is calculated by finding the number of permutations for which the maximal component size exceeds the original test statistic, normalised by *M*. To identify group differences using NBS, we used M=5000 permutations with independent samples t-test ($p<0.05$).

Multilayer network construction

A multilayer network is a complex network structure, which is a network of networks, where each layer is formed by an individual network. The links in such a network describe relationships between combinations of all possible nodes and layers. In this study, multilayer networks were constructed by integrating information from theta (6.5-8Hz), alpha1 (8.5-10Hz), alpha2 (10.5-12Hz), beta1 (12.5-16Hz), beta2 (16.5-20Hz) frequency bands, where each network or layer shared the same number of nodes (78 cortical regions). The links in each layer were the amplitude envelope correlation (AEC) weights within that frequency band. Figure 1 shows a schematic of multilayer network construction.

Mathematical representation of a multilayer network is its block adjacency matrix (De Domenico et al., 2013; Van Mieghem, 2016). An *f*-layered multilayer network can be written in the form of a block adjacency matrix (Sahneh et al., 2015) as follows:

$$A = \begin{bmatrix} A_1 & H_{12} & \cdots & H_{1f} \\ H_{21} & A_2 & & \vdots \\ \vdots & & \ddots & \\ H_{f1} & \cdots & & A_f \end{bmatrix}, \tag{1}$$

where A_α corresponds to a symmetric, square adjacency matrix of a layer *α*, $1 \leq \alpha \leq f$. In our case A_α corresponds to individual frequency specific networks. The matrix H_{kl} corresponds to the coupling matrix between the layers *k* and *l*, where $1 \leq k, l \leq f, f = 5$. The frequency specific networks, A_α, have the same dimensions for all layers (*N x N, N=78*), implying that every layer of the network has the same number of nodes or brain regions.

To be in line with previous studies (Mandke et al., 2018; Yu et al., 2017), we chose one-to-one between layer relationship $H_{kl} = cI$, where c is a constant and *I* the identity matrix (more details on the inter-layer coupling in Supplementary material 2). Thus, we used a case where coupling matrices are special diagonal matrices ($H_{kl} = cI$) with c being the mean connectivity calculated over all the layers. In other words, we allow only for links between the same nodes across all layers (i.e. we ignore cross-frequency coupling between non-identical areas in different layers).

Figure 1. *A schematic of the analysis pipeline used in the construction of multilayer networks. Here, we use a special case of the multilayer networks: multilayer networks with one-to-one between-layer coupling. rs: Resting state; AEC: Amplitude envelope correlations; Theta: 6.5-8Hz; Alpha1: 8.5-10Hz; Alpha2: 10.5-12Hz; Beta1: 12.5-16Hz; Beta2: 16.5-20Hz.*

Graph analysis of multilayer networks

A multilayer network was constructed for every participant in the two groups. However, comparing raw block-adjacency matrices between groups can be biased due to differences in link weights, i.e. such a comparison can inflate false positives or negatives (Mandke et al., 2018). These biases in the context of multilayer networks arise due to differences in average connectivity (van den Heuvel et al., 2017). To correct for such biases we used singular value decomposition (SVD) based normalisation approach (Mandke et al., 2018). Prior to subjecting the block adjacency matrix to graph analysis, all singular values were normalised by the

largest singular value of the weighted adjacency matrix to correct for difference in mean connectivity between groups. For a given block adjacency matrix A, we apply a singular value decomposition $A = U\Lambda V^T$, where U and V contain the left and right singular vectors and Λ the singular values of A. To correct for differences in average connectivity, we rescaled Λ by the largest singular value λ_1, $\tilde{A} = U\left(\frac{10}{\lambda_1}\Lambda\right)V^T$ (the justification of the choice of eigenvalue can be found in the Supplementary Material 1). The rescaled matrix would thus become $\tilde{A}$. The multiplication by 10 is used to ensure that the range of values in $\tilde{A}$ is not too small and varies between 0 and 1.

To identify community structure specific to musicians we used a community detection algorithm, details of which are given in the next paragraph. Multilayer network formulation allows us to study various between layer relationships. Degree correlations (DC) is one of such measures that enables us to study the correspondence between hubs (defined here as high-degree nodes) in different layers. DC can be derived by calculating a Pearson correlation coefficient between degree sequences of different layers (Battiston et al., 2014; Mandke et al., 2018), where degree provides a measure for "importance" of a given node. In the case of weighted networks, node strength (i.e. the sum of the link weights surrounding a node) is used instead of degree. Here, we focus on DC as a measure of relative importance of a given node and study the modularity to go beyond individual nodes and study their higher-order organisation.

Community detection algorithms (e.g. (Blondel et al., 2008) have widely been used to study resting-state fMRI networks and structural data (Crossley et al., 2013; Garcia et al., 2018; Sporns & Betzel, 2016). Generally, communities in a network are obtained by optimising a quality function, e.g. modularity (Newman, 2006), which provides an index for within-community and between-community connections. The multilayer modularity as expressed in (Bassett et al., 2013) is as follows:

$$Q = \frac{1}{2u_{tot}}\sum_{ijlk}\left[\left(\tilde{A}_{l,ij} - \gamma_l \frac{_ld_i\,_ld_j}{2\sum_i\,_ld_i}\delta_{lk}\right) + \delta_{ij}H_{lk,j}\right]\delta_{g_{il},g_{jk}} \qquad (2)$$

where u_{tot} is the total link weight in the network, γ_l is the resolution parameter (set to 1, constant over all subjects), g_{il} is the community assignment of node i in layer l, and δ_{ij} is the Kronecker delta function (meaning that δ_{ij}=1 if $i=j$ and δ_{ij}=0 if i≠j). Using this community

detection algorithm, we calculated individual nodal module assignments expressed as indices. Nodes with the same index can be said to be part of the same community.

We first performed this community detection algorithm on the multilayer matrix of each individual subject. Using these index assignments, we calculated an agreement (consensus) matrix ($N \times N$) for every layer per group, where each element in the matrix corresponds to the proportional number of times two given nodes were part of the same community over all individual subjects (Baum et al., 2017). Furthermore, we obtained a collapsed agreement matrix by averaging over all individual layer agreement matrices. This step was performed to achieve a group-level representation of the entire multilayer network.

To compare the multilayer community structure between groups (musicians vs non-musicians), we subjected each collapsed agreement matrix to the same community detection algorithm (Equation 2), resulting in a partition of the 78 nodes into a number of communities (Baum et al., 2017). To identify whether the community assignments between the groups are similar or dissimilar we calculated the adjusted Rand index (Traud et al., 2008), which also takes into account chance correlations. For any two given partitions A and B, the adjusted Rand index achieves a maximum score of 1 if they are perfectly similar, and 0 if they have no overlap.

Results

Functional connectivity and Network-Based Statistics

Seed-based connectivity analysis for the 4 nodes under consideration (bilateral SMA and pre-central gyri) did not reveal any statistically significant ($p<0.05$) differences between the two groups.

After performing SVD normalisation, for visual representation of the data, functional connectivity estimates between all nodes for all frequency bands (theta, alpha1, alpha2, beta1, beta2) obtained using AEC were averaged across participants (Figure 2: musicians and Figure 3: non-musicians). Only the top 40% (i.e., the strongest) of connections were subsequently plotted in brain space to visualise the underlying networks (see Figures 2 and 3).

After computing individual connectivity matrices, we performed single layer analysis using NBS. However, this failed to show any significant effect with stringent statistical thresholding ($p<0.05$, FDR-corrected) across frequency bands. A null result using NBS is possible because of several reasons: 1) Single layer analysis ignores the pan-spectral picture in the data, which warrants a more integrative approach. 2) High differences in link weights can smear underlying experimental effects and this demands some corrections as highlighted in (Mandke et al., 2018; van Wijk et al., 2010).

Figure 2. *Averaged AEC in adjacency matrix (78 brain area labels on each axis: 1-39 left hemisphere, 40-78 right hemisphere) across musicians (A' to E') and its corresponding source space (top 40% (i.e., the strongest) of connections) representation (3D plots, A to E) in the 5 frequency bands under consideration. All adjacency matrices show typical patterns of high-AEC within left hemisphere (upper left quadrant), within right hemisphere (lower right quadrant) and between hemispheres (upper right quadrant). In the 3D plots, red lines indicate presence of connections, with thicker lines corresponding to stronger connections. The grey circles in the 3D plots indicate summed magnitude of link weights (node strength, weighted degree) between the node and the rest of the brain. ROIs: Regions of interest.*

Figure 3. *Averaged AEC* plotted as *adjacency* matrices (*78 brain area labels on each axis: 1-39 left hemisphere, 40-78 right hemisphere) across non-musicians (A' to E') and their corresponding source spa* ROIs *.e., the strongest) of connections) representation (3D plots, A to E) in 5 frequency bands under consideration. In the 3D plots, red lines indicate presence of connections, with thicker lines corresponding to stronger connections. The grey circles in the 3D plots indicate summed magnitude of link weights (node strength, weighted degree) between the node and rest of the brain. ROIs: Regions of interest.*

Graph analysis of the multilayer networks

   a) Community structure

The block adjacency matrices of the two groups were subjected to community detection algorithm after SVD normalisation. Modularity (Q) was calculated for individual subjects' SVD normalised multilayer networks (Figure 4a) and was significantly different between groups (*p*=0.033, Wilcoxon rank-sum test). To further understand the link between musical training and neural connectivity findings, we correlated years of musical training with modularity, but this correlation was not found to be significant (*p*=0.8). Moreover, we also correlated age at which participants started taking lessons with modularity and this was also found to be non-significant (*p*=0.1).

Figure 4. (A) Boxplots *of modularity (Q) for the two groups. This* difference *was found to be statistically significant (p=0.033 Wilcoxon rank-sum test). (B) Boxplots of degree correlations. Only the layer pairs that showed significant differences are plotted. Left panel shows significantly different DC between theta layer (layer 1) and beta1 layer (layer 4) (p=0.045, Wilcoxon rank-sum test). Right panel shows significantly different DC between layer 2 (alpha1) and layer 4 (beta1) (p=0.031, Wilcoxon rank-sum test). The box shows 95% SEM and SD plotted as whiskers.*

The group of musicians demonstrated different modular organisation than the non-musicians (adjusted Rand index: 0.568). The group representation of this modular organisation for both groups is shown in Figure 5. The multilayer network community structure observed in musicians had two large communities, while non-musicians had three communities. Musicians showed a modular structure that encompasses both visual and motor networks, shown in red (Figure 5, left panel), and frontal and temporal regions, shown in blue. In non-musicians, the modularity analysis revealed three modules, the frontal (colour: blue), midline-temporal (colour: green) and posterior (colour: red) network components. The non-musicians showed distinctly different community structure from the group of musicians. In non-musicians, the visual and motor networks are segregated from one another (as shown in red and green, respectively, see Figure 5), whereas musicians display integrated visuo-motor networks. In addition, the musicians also exhibit markedly different patterns of connectivity spanning the fronto-temporal regions, in particular in the right hemisphere, with integration of frontal and superior temporal regions into a single community only in musicians (Figure 5, left panel).

Figure 5. Schematic representation of collapsed agreement matrix computed for both groups. The collapsed agreement matrices were subjected to community detection algorithm. The community assignment was subsequently plotted in brain space to reveal group level representation. Within each group, identical colours belong to the same community.

b) Degree correlations

One aspect of between layer relationships is the correspondence between hubs in different layers, which can be quantified using DC. We calculated DC between all combinations of layers. Out of all the layer combinations only statistically significant (p<0.05, Wilcoxon rank-sum test) pairs are shown in Figure 4B. The left panel in Figure 4B shows the DC difference between layer 1 (theta) and layer 4 (beta1) and the right panel displays the DC difference between layer 2 (alpha1) and layer 4 (beta1) between groups.

Discussion

We studied whether an impact of musical training would be detectable in musicians' brain function, measured with MEG during resting state. To this end, a novel multilayer network framework to construct multi-frequency networks in a group of musicians and non-musicians was used. We studied the properties of these networks using graph-theoretical approaches. As per the hypothesis, a modular organisation in musicians spanning the visuomotor regions and the frontal and auditory parts of the brain was found. This modular organisation points to a stronger inter-modular connectivity, potentially underlying the integrated organisation observed in musicians. It must be noted here that neither single layer analysis using NBS nor seed-based connectivity analysis revealed significant group differences. Therefore, our results highlight the added value of the multilayer analysis.

The present methods-focused study adds to the body of knowledge about brain reorganisation in highly-trained musicians, which was so far limited to structural MRI studies for example (Schlaug et al., 1995), (Amunts et al., 1997), (Bangert & Schlaug, 2006), and task-based for example, (Alluri et al., 2017; Grahn & Rowe, 2009) as well as resting-state fMRI studies for example (Luo et al., 2012, 2014; Zamorano et al., 2017). The question whether musical training would also be reflected in electrophysiological functional networks outside of task conditions – during resting state – has only recently been addressed, to our knowledge in a single study so far: (Klein et al., 2016). They reported an increased intra- and inter-hemispheric functional connectivity in resting-state EEG data of musicians in theta (6.5-8Hz) and alpha1 (8.5-10Hz) frequency bands. However, their investigation was limited to studying frequency specific networks in isolation. This single layer approach does not detect pan-spectral functional networks.

In addition to studying its modularity, the multilayer network framework also allows to analyse between layer dependencies. When the brain is modelled as a multilayer network, the individual layers can act independently, in coherence of one another or at a transition point between these two extremes (Tewarie, Hillebrand, van Dijk, et al., 2016). One important aspect of this between layer dependency is the correspondence between hubs in different layers. To probe this further, we calculated the DC between all combinations of the layers and compared them between the two groups. Our analysis revealed that musicians show a significantly different degree distribution compared to non-musicians, particularly in the theta-beta1 and alpha1-beta1 coupling.

The findings from our multilayer analysis underscore two important aspects about network organisation in musicians:

1) At a global pan-spectral level, musicians have less segregated clusters (lower modularity), thereby allowing for an integrated network organisation.

2) Between layers, we find a lower DC value for musicians, which points towards different layers functioning independently within the multilayer network, i.e. the hubs in the two layers are distinctly different from one another.

Taken together, musicians possess a highly integrated organization over the whole multi-frequency network, but the different layers retain at the same time their own unique network structure. For example, (classical) musicians often need to read sheet music, encode the visual information (alpha band network) and translate it into signals to be executed by the motor cortex (beta band network). The between network coordination is not limited only to visuo-motor areas, but has also been shown to span auditory, motor and other cognitive areas, which likely facilitates higher overall network integration (Jäncke, 2012; Proverbio et al., 2014). For instance, the coordination between auditory and motor areas is said to be mediated via the arcuate fasciculus (Catani et al., 2005; Catani & Mesulam, 2008). It is also plausible that this anatomical architecture underlies the auditory-motor coupling reported in musicians (Grahn & Rowe, 2009; Jäncke, 2012).

The critical difference observed in our analysis regarding between layer organization lies primarily in the theta-beta1 and alpha1-beta1 layers. The present results observed in lower frequency bands may reflect long-range communication (Ward, 2003). Both theta and alpha band activity are involved in a wide range of cognitive functions (review: (Başar et al., 2001; Ward, 2003) – such as memory and attention, which are highly active areas while playing an instrument (Schlaug, 2015).

The relationship between behavioural specialisation and changes in the underlying neuronal activity is not very straightforward (Zatorre et al., 2012). This problem is more pronounced when investigating spontaneous neuronal oscillations (using M/EEG), as both increases and decreases in activity can reflect a different brain state (Baillet, 2017). There are several studies (Schneider et al., 2002; Shahin et al., 2003; Strait & Kraus, 2011, 2014) that have used evoked potentials to investigate musical specialisation. However, such evidence does not exist for resting-state data. Our results also show no significant correlation between multilayer

network measures with years of training as a behavioural variable. Modularity in this case might not be sensitive enough to detect correlation within the musician group but can detect between-group differences (absence vs presence of experience).

A possible solution to circumvent this issue would be to conduct a longitudinal electrophysiological study to establish a clearer link between behaviour and increase or decrease in the underlying neuronal activity (using a similar approach as in the structural imaging study by (Draganski et al., 2004)). In addition, the number of analysed group members in the musician group (n=31) in our study is also small and could be the reason for not identifying a clear relationship between behaviour and neural dynamics.

Brain dynamics occur across multiple spatial and temporal scales. Various neuro-imaging techniques measure different aspects of the same underlying physiological processes, but these data sets are often analysed in isolation. However, the multilayer network framework has the potential to bridge this gap by taking a more integrative approach (De Domenico, 2017; Mandke et al., 2018). For example, the framework can be used to integrate data from MEG in different frequency ranges (as in the current study), from DTI and MEG or fMRI and MEG in the same group of participants. As a future research direction, one can also integrate the structural network information as an additional network layer in the multilayer framework, as e.g. investigated in (Battiston et al., 2017, 2018; Lim et al., 2019) for one fMRI-based and an additional structural network layer for healthy controls. Recently, a framework for including - next to structural and functional layers - also a third morphological grey matter network layer has been proposed (Casas-Roma et al., 2022). Therefore, the multilayer network approach allows us to study in an individual how brain processes at different scales, from functional to structural, undergo plasticity related changes. Our study is one of the first demonstrations that an integrative approach is useful to study training related plasticity.

The present study is not without limitations. Firstly, the data used here were part of the Open MEG Archive (Niso et al., 2016), which included a number of participants (non-musicians, 8 out of 31 participants), who played an instrument for 5+ years but did not identify themselves as professional musicians. In addition, the participants who were professional musicians did not play the same instrument nor did they have similar years of experience (Mean: 13.25 years; SD: 5.04) playing an instrument. A future study could be designed to replicate these findings, with a more careful selection of participants to identify a clear relationship between resting-state network organisation and years of experience. Secondly, the MEG recordings in

our study were only 5 min in duration. The recording duration, an important aspect in capturing brain states in individual subjects (Liuzzi et al., 2017), should be carefully considered when designing future resting-state EEG or MEG studies. The OMEGA questionnaire data does not include more specific information on musical training or preference of musical style. As a result, we were unable to investigate these aspects. With more elaborate questionnaire data, one could analyse which precise factors of musical training cause the multilayer network changes. NBS relies on choosing an arbitrary test statistic threshold (Zalesky et al., 2010), representing a critical limitation of this method. While it is the only parameter that needs to be manually set by the user, the results depend on the appropriate selection of the test statistic.

The majority of the previous studies that have reported group differences in motor cortices of musicians in both structural and task-based functional data (see for example: (Palomar-García et al., 2017; Schlaug, 2015; Schlaug et al., 1995; Stewart, 2008), have recruited participants that play the same instrument professionally and have very similar levels of expertise. A critical limitation of the present study is the heterogeneity among participants in terms of years of experience (13.25 years ± 5.04) and a wide range of musical expertise (string instruments, percussions etc). This heterogeneity coupled with resting-state MEG recordings that were only 5-mins long probably make identification of seed-based differences difficult. Hence, this methods-focused study focused on identifying global changes in the network configuration. However, it is plausible in a future study to introduce a task that engages both auditory and motor systems, which would likely make the differences more apparent.

Recently, in the context of speech, (Assaneo & Poeppel, 2018) provided evidence for the presence of an intrinsic rhythm – which mediates auditory-motor communication. It is conceivable that such a fine-tuned mechanism might also exist in musicians, mediating the communication between different areas and potentially leading to the formation of sub-networks or communities. This analysis needs to be probed further by studying the relationship between task-based data and task-free data in the same group of musicians.

To conclude, we used an open access dataset (OMEGA, (Niso et al., 2016)) to investigate the effects of musical training related plasticity in resting-state MEG data. We studied the frequency specific networks using NBS and by applying a novel multilayer network framework

to construct multi-frequency networks in musicians and non-musicians. The single layer analysis did not reveal a statistically significant difference between the two groups. However, effects of musical training were detected in the modularity of the network, which spanned visuo-motor and fronto-temporal areas that are involved in execution of a musical performance. The results also indicate that musicians show an integrated network organisation in comparison to non-musicians. DC results showed a significantly different layer-to-layer correspondence, potentially indicating that the formation of hubs in different layers is significantly different between the two groups. Presently, the study of musical training related plasticity lacks an integrative approach and therefore neglects interactions between processes of plasticity occurring at different frequency bands and/or different modalities within individuals. To overcome this limitation, we suggest the use of the multilayer network framework.

Acknowledgements:

We thank the University of Nottingham for Vice Chancellor's Scholarship awarded to KM.

References

Alluri, V., Brattico, E., Toiviainen, P., Burunat, I., Bogert, B., Numminen, J., & Kliuchko, M. (2015a). Musical expertise modulates functional connectivity of limbic regions during continuous music listening. *Psychomusicology: Music, Mind, and Brain*, *25*(4), 443–454. https://doi.org/10.1037/pmu0000124

Alluri, V., Brattico, E., Toiviainen, P., Burunat, I., Bogert, B., Numminen, J., & Kliuchko, M. (2015b). Musical expertise modulates functional connectivity of limbic regions during continuous music listening. *Psychomusicology: Music, Mind, and Brain*, *25*(4), 443–454. https://doi.org/10.1037/pmu0000124

Alluri, V., Toiviainen, P., Burunat, I., Kliuchko, M., Vuust, P., & Brattico, E. (2017). Connectivity patterns during music listening: Evidence for action-based processing in musicians. *Human Brain Mapping*, *38*(6), 2955–2970. https://doi.org/10.1002/hbm.23565

Amunts, K., Schlaug, G., Jäncke, L., Steinmetz, H., Schleicher, A., Dabringhaus, A., & Zilles, K. (1997). Motor cortex and hand motor skills: Structural compliance in the human brain. *Human Brain Mapping*, *5*(3), 206–215. https://doi.org/10.1002/(SICI)1097-0193(1997)5:3<206::AID-HBM5>3.0.CO;2-7

Assaneo, M. F., & Poeppel, D. (2018). The coupling between auditory and motor cortices is rate-restricted: Evidence for an intrinsic speech-motor rhythm. *Science Advances*, 1–10. https://doi.org/10.1126/sciadv.aao3842

Baillet, S. (2017). Magnetoencephalography for brain electrophysiology and imaging. *Nature Neuroscience*, *20*(3), 327–339. https://doi.org/10.1038/nn.4504

Bangert, M., & Schlaug, G. (2006). Specialization of the specialized in features of external human brain morphology. *European Journal of Neuroscience*, *24*(6), 1832–1834. https://doi.org/10.1111/j.1460-9568.2006.05031.x

Barnes, G. R., Hillebrand, A., Fawcett, I. P., & Singh, K. D. (2004). Realistic spatial sampling for MEG beamformer images. *Human Brain Mapping*, *23*(2), 120–127. https://doi.org/10.1002/hbm.20047

Başar, E., Başar-Eroglu, C., Karakaş, S., & Schürmann, M. (2001). Gamma, alpha, delta, and theta oscillations govern cognitive processes. *International Journal of Psychophysiology*, *39*(2–3), 241–248. https://doi.org/10.1016/s0167-8760(00)00145-8

Bassett, D. S., Porter, M. A., Wymbs, N. F., Grafton, S. T., Carlson, J. M., & Mucha, P. J. (2013). Robust detection of dynamic community structure in networks. *Chaos*, *23*(1), 013142–16. https://doi.org/10.1063/1.4790830

Bassett, D. S., Wymbs, N. F., Porter, M. A., Mucha, P. J., Carlson, J. M., & Grafton, S. T. (2010). Dynamic reconfiguration of human brain networks during learning. *Proceedings of the National Academy of Sciences*, *108*(18), 7641–7646. https://doi.org/10.1073/pnas.1018985108

Battiston, F., Guillon, J., Chavez, M., Latora, V., & De Vico Fallani, F. (2018). Multiplex core–periphery organization of the human connectome. *Journal of The Royal Society Interface*, *15*(146), 20180514. https://doi.org/10.1098/rsif.2018.0514

Battiston, F., Nicosia, V., Chavez, M., & Latora, V. (2017). Multilayer motif analysis of brain networks. *Chaos: An Interdisciplinary Journal of Nonlinear Science*, *27*(4), 047404. https://doi.org/10.1063/1.4979282

Battiston, F., Nicosia, V., & Latora, V. (2014). Structural measures for multiplex networks. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, *89*(3), 1–14. https://doi.org/10.1103/PhysRevE.89.032804

Baum, G. L., Ciric, R., Roalf, D. R., Betzel, R. F., Moore, T. M., Shinohara, R. T., Kahn, A. E., Quarmley, M., Cook, P. A., Elliot, M. A., Ruparel, K., Gur, R. E., Gur, R. C., Bassett, D. S., & Satterthwaite, T. D. (2017). Modular segregation of structural brain networks supports the development of executive function in youth. *Current Biology*, *27*(11), 1561–1572. https://doi.org/10.1016/j.cub.2017.04.051

Blondel, V. D., Guillaume, J.-L., Lambiotte, R., & Lefebvre, E. (2008). Fast unfolding of communities in large networks. *Journal of Statistical Mechanics: Theory and Experiment*, *2008*(10), 10008–10012. https://doi.org/10.1088/1742-5468/2008/10/P10008

Brookes, M. J., Tewarie, P. K., Hunt, B. A. E., Robson, S. E., Gascoyne, L. E., Liddle, E. B., Liddle, P. F., & Morris, P. G. (2016a). A multi-layer network approach to MEG connectivity analysis. *NeuroImage*, *132*, 425–438. https://doi.org/10.1016/j.neuroimage.2016.02.045

Brookes, M. J., Tewarie, P. K., Hunt, B. A. E., Robson, S. E., Gascoyne, L. E., Liddle, E. B., Liddle, P. F., & Morris, P. G. (2016b). A multi-layer network approach to MEG connectivity analysis. *NeuroImage*, *132*, 425–438. https://doi.org/10.1016/j.neuroimage.2016.02.045

Brookes, M. J., Vrba, J., Robinson, S. E., Stevenson, C. M., Peters, A. M., Barnes, G. R., Hillebrand, A., & Morris, P. G. (2008). Optimising experimental design for MEG beamformer imaging. *NeuroImage*, *39*(4), 1788–1802. https://doi.org/10.1016/j.neuroimage.2007.09.050

Brookes, M. J., Woolrich, M. W., Luckhoo, H., Price, D., Hale, J. R., Stephenson, M. C., Barnes, G. R., Smith, S. M., & Morris, P. G. (2011). Investigating the electrophysiological basis of resting state networks using magnetoencephalography. *Proceedings of the National Academy of Sciences*, *108*(40), 16783–16788. https://doi.org/10.1073/pnas.1112685108

Buldú, J. M., & Porter, M. A. (2018). Frequency-based brain networks: From a multiplex framework to a full multilayer description. *Network Neuroscience*, *2*(4), 418–441. https://doi.org/10.1162/netn_a_00033

Casas-Roma, J., Martinez-Heras, E., Solé-Ribalta, A., Solana, E., Lopez-Soley, E., Vivó, F., Diaz-Hurtado, M., Alba-Arbalat, S., Sepulveda, M., Blanco, Y., Saiz, A., Borge-Holthoefer, J., Llufriu, S., & Prados, F. (2022). Applying multilayer analysis to morphological, structural, and functional brain networks to identify relevant dysfunction patterns. *Network Neuroscience*, *6*(3), 916–933. https://doi.org/10.1162/netn_a_00258

Catani, M., Jones, D. K., & Ffytche, D. H. (2005). Perisylvian language networks of the human brain. *Annals of Neurology*, *57*(1), 8–16. https://doi.org/10.1002/ana.20319

Catani, M., & Mesulam, M. (2008). The arcuate fasciculus and the disconnection theme in language and aphasia: History and current state. *Cortex*, *44*(8), 953–961. https://doi.org/10.1016/j.cortex.2008.04.002

Cohen, M. X., Axmacher, N., Lenartz, D., Elger, C. E., Sturm, V., & Schlaepfer, T. E. (2009). Good Vibrations: Cross-frequency Coupling in the Human Nucleus Accumbens during Reward Processing. *Journal of Cognitive Neuroscience*, *21*(5), 875–889. https://doi.org/10.1162/jocn.2009.21062

Colclough, G. L., Brookes, M. J., Smith, S. M., & Woolrich, M. W. (2015). A symmetric multivariate leakage correction for MEG connectomes. *NeuroImage*, *117*, 439–448. https://doi.org/10.1016/j.neuroimage.2015.03.071

Crossley, N. A., Mechelli, A., Vértes, P. E., Winton-Brown, T. T., Patel, A. X., Ginestet, C. E., McGuire, P., & Bullmore, E. T. (2013). Cognitive relevance of the community structure of the human brain functional coactivation network. *Proceedings of the National Academy of Sciences*, *110*(38), 15502–15502. https://doi.org/10.1073/pnas.1314559110

De Domenico, M. (2017). Multilayer modeling and analysis of human brain networks. *GigaScience*, *6*, 1–8. https://doi.org/10.1093/gigascience/gix004

De Domenico, M., Sasai, S., & Arenas, A. (2016). Mapping Multiplex Hubs in Human Functional Brain Networks. *Frontiers in Neuroscience*, *10*. https://www.frontiersin.org/articles/10.3389/fnins.2016.00326

De Domenico, M., Solé-Ribalta, A., Cozzo, E., Kivelä, M., Moreno, Y., Porter, M. A., Gómez, S., & Arenas, A. (2013). Mathematical formulation of multi-layer networks. *Physical Review X*, *3*(4), 1–15. https://doi.org/10.1103/PhysRevX.3.041022

Draganski, B., Gaser, C., Busch, V., Schuierer, G., Bogdahn, U., & May, A. (2004). Changes in grey matter induced by training. *Nature*, *427*(6972), Article 6972. https://doi.org/10.1038/427311a

Elbert, T., Pantev, C., Wienbruch, C., Rockstroh, B., & Taub, E. (1995). Increased Cortical Representation of the Fingers of the Left Hand in String Players. *Science*, *270*(5234), 305–307. https://doi.org/10.1126/science.270.5234.305

Florin, E., & Baillet, S. (2015). The brain's resting-state activity is shaped by synchronized cross-frequency coupling of neural oscillations. *NeuroImage*, *111*, 26–35. https://doi.org/10.1016/j.neuroimage.2015.01.054

Fries, P. (2005). A mechanism for cognitive dynamics: Neuronal communication through neuronal coherence. *Trends in Cognitive Sciences*, *9*(10), 474–480. https://doi.org/10.1016/j.tics.2005.08.011

Friston, K. J. (2011). Functional and Effective Connectivity: A Review. *Brain Connectivity*, *1*(1), 13–36. https://doi.org/10.1089/brain.2011.0008

Garcia, J. O., Ashourvan, A., Muldoon, S., Vettel, J. M., & Bassett, D. S. (2018). Applications of Community Detection Techniques to Brain Graphs: Algorithmic Considerations and Implications for Neural Function. *Proceedings of the IEEE*, *106*(5), 846–867. https://doi.org/10.1109/JPROC.2017.2786710

Gaser, C., & Schlaug, G. (2003). Brain structures differ between musicians and non-musicians. *The Journal of Neuroscience*, *23*(27), 9240–9245. https://doi.org/10.1523/JNEUROSCI.23-27-09240.2003

Grahn, J. A., & Rowe, J. B. (2009). Feeling the Beat: Premotor and Striatal Interactions in Musicians and Nonmusicians during Beat Perception. *Journal of Neuroscience*, *29*(23), 7540–7548. https://doi.org/10.1523/JNEUROSCI.2018-08.2009

Gross, J., Baillet, S., Barnes, G. R., Henson, R. N., Hillebrand, A., Jensen, O., Jerbi, K., Litvak, V., Maess, B., Oostenveld, R., Parkkonen, L., Taylor, J. R., van Wassenhove, V., Wibral, M., & Schoffelen, J.-M. (2013). Good practice for conducting and reporting MEG research. *NeuroImage*, *65*, 349–363. https://doi.org/10.1016/j.neuroimage.2012.10.001

Guillon, J., Attal, Y., Colliot, O., La Corte, V., Dubois, B., Schwartz, D., Chavez, M., & De Vico Fallani, F. (2017). Loss of brain inter-frequency hubs in Alzheimer's disease. *Scientific Reports*, *7*(1), Article 1. https://doi.org/10.1038/s41598-017-07846-w

Herholz, S. C., & Zatorre, R. J. (2012). Musical Training as a Framework for Brain Plasticity: Behavior, Function, and Structure. *Neuron*, *76*(3), 486–502. https://doi.org/10.1016/j.neuron.2012.10.011

Hillebrand, A., Barnes, G. R., Bosboom, J. L., Berendse, H. W., & Stam, C. J. (2012). Frequency-dependent functional connectivity within resting-state networks: An atlas-based MEG beamformer solution. *NeuroImage*, *59*(4), 3909–3921. https://doi.org/10.1016/j.neuroimage.2011.11.005

Huang, M.-X., Mosher, J. C., & Leahy, R. M. (1999). A sensor-weighted overlapping-sphere head model and exhaustive head model comparison for MEG. *Physics in Medicine and Biology*, *44*(2), 423–440. https://doi.org/10.1088/0031-9155/44/2/010

Jäncke, L. (2009a). The Architecture of the Golfer's Brain. *PLoS ONE*, *101*(11), 1435–1439. https://doi.org/10.1371/journal.pone.0004785

Jäncke, L. (2009b). The plastic human brain. *Restorative Neurology and Neuroscience*, *27*(5), 521–538. https://doi.org/10.3233/RNN-2009-0519

Jäncke, L. (2012). The dynamic audio-motor system in pianists. *Annals of the New York Academy of Sciences*, *1252*(1), 246–252. https://doi.org/10.1111/j.1749-6632.2011.06416.x

Jensen, O., & Colgin, L. L. (2007). Cross-frequency coupling between neuronal oscillations. *Trends in Cognitive Sciences*, *11*(7), 267–269. https://doi.org/10.1016/j.tics.2007.05.003

Kivelä, M., Arenas, A., Barthelemy, M., Gleeson, J. P., Moreno, Y., & Porter, M. A. (2014). Multilayer networks. *Journal of Complex Networks*, *2*(3), 203–271. https://doi.org/10.1093/comnet/cnu016

Klein, C., Liem, F., Hänggi, J., Elmer, S., & Jäncke, L. (2016). The "silent" imprint of musical training. *Human Brain Mapping*, *37*(2), 536–546. https://doi.org/10.1002/hbm.23045

Langer, N., von Bastian, C. C., Wirz, H., Oberauer, K., & Jäncke, L. (2013). The effects of working memory training on functional brain network efficiency. *Cortex*, *49*(9), 2424–2438. https://doi.org/10.1016/j.cortex.2013.01.008

Lim, S., Radicchi, F., van den Heuvel, M. P., & Sporns, O. (2019). Discordant attributes of structural and functional brain connectivity in a two-layer multiplex network. *Scientific Reports*, *9*(1), Article 1. https://doi.org/10.1038/s41598-019-39243-w

Liuzzi, L., Gascoyne, L. E., Tewarie, P. K., Barratt, E. L., Boto, E., & Brookes, M. J. (2017). Optimising experimental design for MEG resting state functional connectivity measurement. *NeuroImage*, *155*(November 2016), 565–576. https://doi.org/10.1016/j.neuroimage.2016.11.064

Luckhoo, H., Hale, J. R., Stokes, M. G., Nobre, A. C., Morris, P. G., Brookes, M. J., & Woolrich, M. W. (2012). Inferring task-related networks using independent component analysis in magnetoencephalography. *NeuroImage*, *62*(1), 530–541. https://doi.org/10.1016/j.neuroimage.2012.04.046

Luo, C., Guo, Z.-W., Lai, Y., Liao, W., Liu, Q., Kendrick, K. M., Yao, D., & Li, H. (2012). Musical training induces functional plasticity in perceptual and motor networks: Insights from resting-state fMRI. *PLoS ONE*, *7*(5), 1–10. https://doi.org/10.1371/journal.pone.0036568

Luo, C., Tu, S., Peng, Y., Gao, S., Li, J., Dong, L., Li, G., Lai, Y., Li, H., & Yao, D. (2014). Long-term effects of musical training and functional plasticity in salience system. *Neural Plasticity*, 1–14. https://doi.org/10.1155/2014/180138

Maguire, E. A., Gadian, D. G., Johnsrude, I. S., Good, C. D., Ashburner, J., Frackowiak, R. S. J., & Frith, C. D. (2000). Navigation-related structural change in the hippocampi of taxi drivers. *Proceedings of the National Academy of Sciences*, 97(8), 4398–4403. https://doi.org/10.1073/pnas.070039597

Mandke, K., Meier, J. M., Brookes, M. J., O'Dea, R. D., Van Mieghem, P., Stam, C. J., Hillebrand, A., & Tewarie, P. K. (2018). Comparing multilayer brain networks between groups: Introducing graph metrics and recommendations. *NeuroImage*, 166, 371–384. https://doi.org/10.1016/j.neuroimage.2017.11.016

Newman, M. E. J. (2006). Modularity and community structure in networks. *Proceedings of the National Academy of Sciences*, 103(23), 8577–8582. https://doi.org/10.1073/pnas.0601602103

Niso, G., Rogers, C., Moreau, J. T., Chen, L. Y., Madjar, C., Das, S., Bock, E., Tadel, F., Evans, A. C., Jolicoeur, P., & Baillet, S. (2016). OMEGA: The Open MEG Archive. *NeuroImage*, 124, 1182–1187. https://doi.org/10.1016/j.neuroimage.2015.04.028

O'Neill, G. C., Barratt, E. L., Hunt, B. A. E., Tewarie, P. K., & Brookes, M. J. (2015). Measuring electrophysiological connectivity by power envelope correlation: A technical review on MEG methods. *Physics in Medicine and Biology*, 60(21), R271–R295. https://doi.org/10.1088/0031-9155/60/21/R271

Onslow, A. C. E., Bogacz, R., & Jones, M. W. (2011). Quantifying phase–amplitude coupling in neuronal network oscillations. *Progress in Biophysics and Molecular Biology*, 105(1), 49–57. https://doi.org/10.1016/j.pbiomolbio.2010.09.007

Palomar-García, M. Á., Zatorre, R. J., Ventura-Campos, N., Bueichekú, E., & Ávila, C. (2017). Modulation of Functional Connectivity in Auditory-Motor Networks in Musicians Compared with Nonmusicians. *Cerebral Cortex*, 27(5), 2768–2778. https://doi.org/10.1093/cercor/bhw120

Power, J. D., Cohen, A. L., Nelson, S. M., Wig, G. S., Barnes, K. A., Church, J. A., Vogel, A. C., Laumann, T. O., Miezin, F. M., Schlaggar, B. L., & Petersen, S. E. (2011). Functional Network Organization of the Human Brain. *Neuron*, 72(4), 665–678. https://doi.org/10.1016/j.neuron.2011.09.006

Proverbio, A. M., Calbi, M., Manfredi, M., & Zani, A. (2014). Audio-visuomotor processing in the Musician's brain: An ERP study on professional violinists and clarinetists. *Scientific Reports*, 4, 5866. https://doi.org/10.1038/srep05866

Puxeddu, M. G., Petti, M., & Astolfi, L. (2021). A Comprehensive Analysis of Multilayer Community Detection Algorithms for Application to EEG-Based Brain Networks. *Frontiers in Systems Neuroscience*, 15. https://www.frontiersin.org/articles/10.3389/fnsys.2021.624183

Robinson, S. E., & Vrba, J. (1999). Functional neuroimaging by synthetic aperture magnetometry. *Recent Advances in Biomagnetism*.

Sahneh, F. D., Scoglio, C., & Van Mieghem, P. (2015). Exact coupling threshold for structural transition reveals diversified behaviors in interconnected networks. *Physical Review E - Statistical, Nonlinear, and Soft Matter Physics*, 92(4), 1–5. https://doi.org/10.1103/PhysRevE.92.040801

Sarvas, J. (1987). Basic mathematical and electromagnetic concepts of the biomagnetic inverse problem. *Physics in Medicine and Biology*, *32*(1), 11–22. https://doi.org/10.1088/0031-9155/32/1/004

Schlaug, G. (2015). Musicians and music making as a model for the study of brain plasticity. *Progress in Brain Research*, *217*, 37–55. https://doi.org/10.1016/bs.pbr.2014.11.020

Schlaug, G., Jäncke, L., Huang, Y., Staiger, J. F., & Steinmetz, H. (1995). Increased corpus callosum size in musicians. *Neuropsychologia*, *33*(8), 1047–1055. https://doi.org/10.1016/0028-3932(95)00045-5

Schneider, P., Scherg, M., Dosch, H. G., Specht, H. J., Gutschalk, A., & Rupp, A. (2002). Morphology of Heschl's gyrus reflects enhanced activation in the auditory cortex of musicians. *Nature Neuroscience*, *5*(7), 688–694. https://doi.org/10.1038/nn871

Scholz, J., Klein, M. C., Behrens, T. E. J., & Johansen-Berg, H. (2010). Training induces changes in white matter architecture. *Nature Neuroscience*, *12*(11), 1370–1371. https://doi.org/10.1038/nn.2412

Shahin, A., Bosnyak, D. J., Trainor, L. J., & Roberts, L. E. (2003). Enhancement of Neuroplastic P2 and N1c Auditory Evoked Potentials in Musicians. *Journal of Neuroscience*, *23*(13), 5545–5552. https://doi.org/10.1523/JNEUROSCI.23-13-05545.2003

Smit, D. J. A., Stam, C. J., Posthuma, D., Boomsma, D. I., & De Geus, E. J. C. (2008). Heritability of "small-world" networks in the brain: A graph theoretical analysis of resting-state EEG functional connectivity. *Human Brain Mapping*, *29*(12), 1368–1378. https://doi.org/10.1002/hbm.20468

Sporns, O., & Betzel, R. F. (2016). Modular brain networks. *Annual Review of Psychology*, *67*(1), 613–640. https://doi.org/10.1146/annurev-psych-122414-033634

Stam, C. J., Montez, T., Jones, B. F., Rombouts, S. A. R. B., Van Der Made, Y., Pijnenburg, Y. A. L., & Scheltens, P. (2005). Disturbed fluctuations of resting state EEG synchronization in Alzheimer's disease. *Clinical Neurophysiology*, *116*(3), 708–715. https://doi.org/10.1016/j.clinph.2004.09.022

Stam, C. J., & van Straaten, E. C. W. (2012). The organization of physiological brain networks. *Clinical Neurophysiology*, *123*(6), 1067–1087. https://doi.org/10.1016/j.clinph.2012.01.011

Stewart, L. (2008). Do musicians have different brains? *Clinical Medicine*, *8*(3), 304–308. https://doi.org/10.7861/clinmedicine.8-3-304

Strait, D. L., & Kraus, N. (2011). Can you hear me now? Musical training shapes functional brain networks for selective auditory attention and hearing speech in noise. *Frontiers in Psychology*, *2*, 1–10. https://doi.org/10.3389/fpsyg.2011.00113

Strait, D. L., & Kraus, N. (2014). Biological impact of auditory expertise across the life span: Musicians as a model of auditory learning. *Hearing Research*, *308*, 109–121. https://doi.org/10.1016/j.heares.2013.08.004

Tanaka, S., & Kirino, E. (2017). Reorganization of the thalamocortical network in musicians. *Brain Research*, *1664*, 48–54. https://doi.org/10.1016/j.brainres.2017.03.027

Tewarie, P. K., Bright, M. G., Hillebrand, A., Robson, S. E., Gascoyne, L. E., Morris, P. G., Meier, J. M., Van Mieghem, P., & Brookes, M. J. (2016). Predicting haemodynamic networks using electrophysiology: The role of non-linear and cross-frequency interactions. *NeuroImage*, *130*, 273–292. https://doi.org/10.1016/j.neuroimage.2016.01.053

Tewarie, P. K., Hillebrand, A., van Dijk, B. W., Stam, C. J., O'Neill, G. C., Van Mieghem, P., Meier, J. M., Woolrich, M. W., Morris, P. G., & Brookes, M. J. (2016). Integrating

cross-frequency and within band functional networks in resting-state MEG: A multi-layer network approach. *NeuroImage*, *142*, 324–336. https://doi.org/10.1016/j.neuroimage.2016.07.057

Tewarie, P. K., Hillebrand, A., Van Dijk, B. W., Stam, C. J., O'Neill, G. C., Van Mieghem, P., Meier, J. M., Woolrich, M. W., Morris, P. G., & Brookes, M. J. (2016). Integrating cross-frequency and within band functional networks in resting-state MEG: A multi-layer network approach. *NeuroImage*, *142*, 324–336. https://doi.org/10.1016/j.neuroimage.2016.07.057

Tewarie, P. K., Prasse, B., Meier, J., Byrne, A., De Domenico, M., Stam, C. J., Brookes, M. J., Hillebrand, A., Daffertshofer, A., & Coombes, S. (2021). Interlayer connectivity reconstruction for multilayer brain networks using phase oscillator models. *New Journal of Physics*, *23*(6), 063065.

Traud, A. L., Kelsic, E. D., Mucha, P. J., & Porter, M. A. (2008). *Comparing Community Structure to Characteristics in Online Collegiate Social Networks*. *53*(3), 526–543. https://doi.org/10.1137/080734315

Tzourio-Mazoyer, N., Landeau, B., Papathanassiou, D., Crivello, F., Etard, O., Delcroix, N., Mazoyer, B., & Joliot, M. (2002). Automated anatomical labeling of activations in SPM using a macroscopic anatomical parcellation of the MNI MRI single-subject brain. *NeuroImage*, *15*(1), 273–289. https://doi.org/10.1006/nimg.2001.0978

Vaiana, M., & Muldoon, S. F. (2020). Multilayer Brain Networks. *Journal of Nonlinear Science*, *30*(5), 2147–2169. https://doi.org/10.1007/s00332-017-9436-8

van den Heuvel, M., de Lange, S. C., Zalesky, A., Seguin, C., Yeo, T. B. T., & Schmidt, R. (2017). Proportional thresholding in resting-state fMRI functional connectivity networks and consequences for patient-control connectome studies: Issues and recommendations. *NeuroImage*, *152*, 437–449. https://doi.org/10.1016/j.neuroimage.2017.02.005

Van Mieghem, P. (2016). Interconnectivity structure of a general interdependent network. *Physical Review E*, *93*(4), 1–11. https://doi.org/10.1103/PhysRevE.93.042305

van Wijk, B. C. M. (2014). Thalamo-cortical cross-frequency coupling detected with MEG. *Frontiers in Human Neuroscience*, *15*(6), 884–890. https://doi.org/10.1038/nn.3101

van Wijk, B. C. M., Stam, C. J., & Daffertshofer, A. (2010). Comparing brain networks of different size and connectivity density using graph theory. *PLoS ONE*, *5*(10). https://doi.org/10.1371/journal.pone.0013701

Ward, L. M. (2003). Synchronous neural oscillations and cognitive processes. *Trends in Cognitive Sciences*, *7*(12), 553–559. https://doi.org/10.1016/j.tics.2003.10.012

Wulff, P., Ponomarenko, A. A., Bartos, M., Korotkova, T. M., Fuchs, E. C., Bähner, F., Both, M., Tort, A. B. L., Kopell, N. J., Wisden, W., & Monyer, H. (2009). Hippocampal theta rhythm and its coupling with gamma oscillations require fast inhibition onto parvalbumin-positive interneurons. *Proceedings of the National Academy of Sciences*, *106*(9), 3561–3566. https://doi.org/10.1073/pnas.0813176106

Yeo, T. B. T., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M., Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fisch, B., Liu, H., & Buckner, R. L. (2011). The organization of the human cerebral cortex estimated by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–1165. https://doi.org/10.1152/jn.00338.2011

Yu, M., Engels, M. M. A., Hillebrand, A., Van Straaten, E. C. W., Gouw, A. A., Teunissen, C., Van Der Flier, W. M., Scheltens, P., & Stam, C. J. (2017). Selective impairment of

hippocampus and posterior hub areas in Alzheimer's disease: An MEG-based multiplex network study. *Brain*, *140*(5), 1466–1485. https://doi.org/10.1093/brain/awx050

Zalesky, A., Fornito, A., & Bullmore, E. T. (2010). Network-based statistic: Identifying differences in brain networks. *NeuroImage*, *53*(4), 1197–1207. https://doi.org/10.1016/j.neuroimage.2010.06.041

Zamorano, A. M., Cifre, I., Montoya, P., Riquelme, I., & Kleber, B. (2017). Insula-based networks in professional musicians: Evidence for increased functional connectivity during resting state fMRI. *Human Brain Mapping*, *38*(10), 4834–4849. https://doi.org/10.1002/hbm.23682

Zatorre, R. J., Fields, R. D., & Johansen-Berg, H. (2012). Plasticity in gray and white: Neuroimaging changes in brain structure during learning. *Nature Neuroscience*, *15*(4), 528–536. https://doi.org/10.1038/nn.3045

Zatorre, R. J., & McGill, J. (2005). Music, the food of neuroscience? *Nature*, *434*(7031), 312–315. https://doi.org/10.1038/434312a

Supplementary material 1 –

The reasons for normalising the symmetric and square block adjacency matrix (as studied here) by its *largest* eigenvalue are as follows –

1) The largest eigenvalue λ_1 is called the spectral radius of the graph (λ_1, λ_2, λ_3,..., λ_n is the eigenspectrum). Previous work has shown that the epidemic threshold (critical threshold) of a network is given by the inverse of the spectral radius of the network's adjacency matrix (Kivelä et al., 2014; Van Mieghem, 2016). Moreover, (Van Mieghem, 2016) has demonstrated that it is also the threshold of phase transition of virus spread and synchronisation of coupled oscillators in networks. Therefore, by normalising the adjacency matrix by its spectral radius, the critical threshold is equalised between groups and datasets. The inverse of spectral radius is a measure of synchronisation threshold, it is related to shortest path, overall connectedness and degree diversity of a graph (for review see (Stam & van Straaten, 2012)

2) The largest eigenvalue is an upper bound for the norm of the adjacency matrix (Van Mieghem, 2016). Therefore, normalisation based on the first eigenvalue is a normalisation factor related to the norm of the adjacency matrix. Any combination of set of values in the eigen spectrum (i.e., sum over all eigenvalues or over more than one of the highest eigenvalues) does not have such a proven mathematical relationship with the matrix norm (to the best of our knowledge).

3) Lastly, the first eigenvalue also captures the main mode of variation of functional connectivity. (Tewarie, Hillebrand, Van Dijk, et al., 2016) demonstrated this by removing the first eigenvalue and calculating the average correlation over all the layers of the multilayer network. This difference led to a sharp increase in global correlations, potentially indicating that the largest eigenvalue is likely responsible for frequency specific information.

Supplementary material 2 -

Methodological considerations for multilayer networks: A note on interlayer coupling

In this work, a frequency-based multilayer network was used, where individual layers of the network were derived from individual MEG frequency bands that were identified *a priori*. The intra-layer (within layer/band) communication was quantified using AEC and the inter-layer or cross-frequency (between layer/band) communication was set to one-to-one (or parameter *c*). Presently, there is no consensus in the network science community on the appropriate way of estimating interlayer coupling parameters and it suffers from some level of arbitrariness (Tewarie et al., 2021). This problem is further compounded by the fact that interlayer coupling between frequency-specific networks can be determined by many different metrics. For instance, the commonly described cross-frequency interactions as per (Jensen & Colgin, 2007) are – (a) power to power, (b) phase to phase, (c) phase to frequency, and (d) phase to power. There is mounting experimental evidence for the presence of phase to power or phase-amplitude coupling both in human (Cohen et al., 2009; Onslow et al., 2011) and animal electrophysiology (Wulff et al., 2009). Furthermore, (Jensen & Colgin, 2007), argue that gamma oscillations (>80 Hz) tend to couple strongly with the phase of the theta cycle (4-8 Hz), which has primarily been shown in intracranial data. A similar relationship has also been shown in resting-state MEG data (Florin & Baillet, 2015), however, the highest frequency band under consideration in the present work was 16.5-20 Hz and the lowest was 6.5-8 Hz, where such a cross-frequency relationship has not been demonstrated before, making estimation of interlayer coupling a non-trivial problem. In a multilayer description, it is possible to include interlayer link weights derived from a separate measure of connectivity, such as, phase-amplitude coupling, but this will likely lead to inconsistent results and requires a careful methodological investigation. Moreover, the non-trivial problem of interlayer coupling is accentuated when considering multilayer networks where individual layers are derived from different imaging modalities (for example, layer 1: fMRI network, layer 2: MEG network).

As problematic as the interlayer coupling is, (De Domenico et al., 2016) argue that presence of such interlayer links allows us to take full advantage of tensorial algebra, allowing us to extend single-layer network metrics to multilayer networks. They also suggest that the choice of "*c*" should therefore depend on the choice of analysis. Moreover, if the value of "*c*" is kept constant between the groups under consideration, the measure of degree will scale linearly

with c. Similarly, as modularity Q depends linearly on H ($H = cI$, see Equation 2 in manuscript), it follows that Q should also scale linearly with c. Recent work on interlayer coupling using a network reconstruction algorithm demonstrated that interlayer coupling is indeed dominated by one-to-one layer coupling for alpha to beta band while strong widespread long-distance interlayer links should be considered when coupling theta and gamma band layers (Tewarie et al., 2021).

In the end, building on the work of (De Domenico et al., 2016) and to maintain consistency between findings reported by (Guillon et al., 2017; Yu et al., 2017) the inter-layer coupling parameter was set to $H_{kl} = cI$. This parameter can be thought of as a scaling parameter, which does not affect the results if it is kept constant for a given analysis. However, future work should consider exploring different interlayer couplings due to their recently discovered frequency-specific nature (Tewarie et al., 2021).

rs-MEG

T_1 MRI

Block adjacency matrix

Beta2

Beta1

Alpha2

Alpha1

Theta

Multilayer network construction

Anatomical atlas

Functional connectivity: AEC

Time series from 78 regions

SVD normalisation

Multilayer graph analysis

Musicians

A: Theta (6.5-8Hz)
B: Alpha1 (8.5-10Hz)
C: Alpha2 (10.5-12Hz)
D: Beta1 (12.5-16Hz)
E: Beta2 (16.5-20Hz)

Non-musicians

A: Theta (6.5-8Hz)
B: Alpha1 (8.5-10Hz)
C: Alpha2 (10.5-12Hz)
D: Beta1 (12.5-16Hz)
E: Beta2 (16.5-20Hz)

Collapsed agreement matrix: Musicians

Collapsed agreement matrix: Non-musicians