Novel metagenomics analysis of stony coral tissue loss disease

Jakob M. Heinz[1,2*‡], Jennifer Lu[1,2,3], Lindsay K. Huebner[4], Steven L. Salzberg[1,2,5,6*], Markus Sommer[1,2*+], Stephanie M. Rosales[7,8*+]

Affiliations:

[1]Center for Computational Biology, Johns Hopkins University; Baltimore, MD 21211, United States

[2]Department of Biomedical Engineering, Johns Hopkins School of Medicine and Whiting School of Engineering; Baltimore, MD 21218, United States

[3]Department of Pathology, Johns Hopkins School of Medicine, Baltimore, MD 21205, United States

[4]Fish and Wildlife Research Institute, Florida Fish and Wildlife Conservation Commission; St. Petersburg, FL 33701, United States

[5]Department of Computer Science, Johns Hopkins University; Baltimore, MD 21218, United States

[6]Department of Biostatistics, Johns Hopkins University; Baltimore, MD 21205, United States

[7]Cooperative Institute for Marine and Atmospheric Studies, University of Miami; Miami, FL 33149, United States

[8]Atlantic Oceanographic and Meteorological Laboratory, National Oceanographic and Atmospheric Administration, Miami, FL 33149, United States

‡ Present address: Department of Biomedical Informatics, Harvard Medical School, Boston, MA 02115, United States

+These authors contributed equally.

Corresponding Authors: jheinz@g.harvard.edu, salzberg@jhu.edu, markusjsommer@gmail.com, stephanie.rosales@noaa.gov

Running head:
Novel metagenomics of SCTLD

Keywords:
Bioinformatics, Metagenomics, SCTLD, Florida's Coral Reef, Epidemic

Abstract

Stony coral tissue loss disease (SCTLD) has devastated coral reefs off the coast of Florida and continues to spread throughout the Caribbean. Although a number of bacterial taxa have consistently been associated with SCTLD, no pathogen has been definitively implicated in the etiology of SCTLD. Previous studies have predominantly focused on the prokaryotic community through 16S rRNA sequencing of healthy and affected tissues. Here, we provide a different analytical approach by applying a bioinformatics pipeline to publicly available metagenomic sequencing samples of SCTLD lesions and healthy tissues from four stony coral species. To compensate for the lack of coral reference genomes, we used data from apparently healthy coral samples to approximate a host genome and healthy microbiome reference. These reads were then used as a reference to which we matched and removed reads from diseased lesion tissue samples, and the remaining reads associated only with disease lesions were taxonomically classified at the DNA and protein levels. For DNA classifications, we used a pathogen identification protocol originally designed to identify pathogens in human tissue samples, and for protein classifications, we used a fast protein sequence aligner. To assess the utility of our pipeline, a species-level analysis of a candidate genus, *Vibrio*, was used to demonstrate the pipeline's effectiveness. Our approach revealed both complementary and unique coral microbiome members compared to a prior metagenome analysis of the same dataset.

Article Summary

Studies of stony coral tissue loss disease (SCTLD), a devastating coral disease, have primarily used 16S rRNA sequencing approaches to identify putative pathogens. In contrast, this study applied human tissue pathogen identification protocols to SCTLD metagenomic DNA samples. Diseased samples were filtered of host sequences using a k-mer based method since host genomes were unavailable. DNA and protein-level classifications from this novel approach revealed both complementary and unique microbiome members compared to a prior metagenome analysis of the same dataset.

Introduction

Stony coral tissue loss disease (SCTLD) was discovered off the coast of Miami, FL in 2014 and has since had negative consequences on the function of coral reefs across Florida and the Caribbean (Walton *et al.* 2018; Alvarez-Filip *et al.* 2022). To date, despite many efforts, no pathogen has been definitively identified as the causative agent of SCTLD. The stony coral (order Scleractinia) microbiome is a complex system of interactions between the host, bacteria, viruses, fungi, archaea, and algal symbionts (Bourne *et al.* 2009); thus a disturbance in any number of these symbiotic relationships could be involved in SCTLD progression. Multiple studies have explored viruses that may infect stony coral symbionts, notably *Symbiodiniaceae*, but no causative relationships have been detected (Work *et al.* 2021; Veglia *et al.* 2022; Beavers *et al.* 2023; Howe-Kerr *et al.* 2023). Bacterial species are particularly under scrutiny for their potential involvement in SCTLD, due to the effectiveness of antibiotics in halting lesion progression in multiple affected coral species (Aeby *et al*. 2019, Neely *et al.* 2020; Shilling *et al.* 2021; Studivan *et al.* 2023). Consequently, SCTLD studies have predominantly focused on understanding changes in the bacterial community between apparently healthy and SCTLD-affected corals.

Studies to identify bacterial pathogens have relied primarily on small subunit 16S ribosomal RNA (rRNA) sequencing, followed by computational analysis (Callahan *et al.* 2016) typically using the Silva database (Quast *et al.* 2013) to assign and classify Amplicon Sequence Variants (ASVs) into taxa. ASVs found in diseased lesion samples are then compared to samples from apparently healthy colonies to determine which ASVs are associated with the tissue loss lesions (Meyer *et al.* 2019; Rosales *et al.* 2020; Clark *et al.* 2021). These methods have characterized many notable shifts in coral bacterial communities due to SCTLD and several bacterial taxa associated with SCTLD lesions, including *Rhizobiales*, *Clostridiales*, *Peptostreptococcales-Tissierellales*, *Rhodobacteraceae*, *Flavobacteriaceae*, and *Vibrionaceae* (Rosales *et al.* 2023). However, because of the difficulty in determining whether an associated bacterial taxon is a harmless commensal, an opportunistic secondary infection, or the primary pathogen, none of the bacterial taxa associated with SCTLD have been identified as the causative agent.

An alternative approach to understanding disease dynamics is the use of metagenomic sequencing, in which all of the DNA from a source is sequenced, including not only the host, but also viruses, bacteria, and eukaryotic species living on or within the host tissue. For example, by analyzing metagenomic sequencing data of human tissue samples taken from the site of infection, researchers have identified pathogenic agents in brain infections (Salzberg *et al.* 2016; Wilson *et al.* 2019), corneal infections (Eberhart *et al.* 2017), and other diseases (Kostic *et al.* 2012). The sensitivity of this approach relies on first, sequencing the source DNA deeply enough to capture the pathogen of interest, and second, the existence of genome assemblies closely related to the pathogen in public databases. While the number of complete genomes has grown

98   enormously over the past two decades, databases still contain few or no genomes for non-model
99   organisms, including scleractinian corals.
100
Currently, data from only one SCTLD metagenome study is publicly available. While the
authors of that study (Rosales *et al.* 2022) were able to assemble and annotate genomes for
SCTLD-associated bacterial taxa such as *Rhodobacterales*, *Rhizobiales*, and *Flavobacteriales*,
the results were focused on only five of the twenty diseased lesion tissue samples, all from the
same coral species (*Stephanocoenia intersepta*), because the majority of samples were dominated
by host sequences. In metagenomic studies, host sequences can confound results, so they are
typically removed by aligning all reads to a host reference genome (Gihawi *et al.* 2023; Lu *et al.*
2022). Currently, the GenBank database has 53 genome assemblies from scleractinian corals, of
which only seven are at the chromosome level (NCBI 2023). Of these 53 genomes, none are
from the species of corals previously investigated for SCTLD (Rosales *et al.* 2022), emphasizing
the additional challenges associated with using metagenomics in non-model organisms.
Additionally, given the complex symbiotic microbiome (i.e., algal symbiont, viruses, and
prokaryotic community) of stony corals (Bourne *et al.* 2009), the host DNA is only one of the
hurdles.
115
In this study, we applied new classification methods to understand this devastating coral disease.
We used a method to filter host reads from metagenome data by using data collected from
apparently healthy corals of the same species to approximate a species-specific healthy host coral
genome and microbiome. We then applied the Kraken software suite for pathogen identification
(Lu *et al.* 2022) using KrakenUniq (Breitwieser *et al.* 2018) to identify putative pathogens
present in diseased samples and not present in healthy ones. Using these methods, we identified a
number of taxa that have previously been associated with SCTLD, providing further support for
their involvement in SCTLD pathogenesis. Finally, from the pool of bacterial taxa we identified
as associated with SCTLD lesions, we selected a candidate genus, *Vibrio*, with which to explore
the utility of our pipeline at a finer taxonomic level.
126

Materials and methods

Data acquisition

We downloaded 58 metagenomic datasets that consisted of 150 bp paired-end reads from NCBI
Bioproject PRJNA576217 (Benson *et al.* 2017), previously generated by Rosales *et al.* (2020).
Sample SRR15960000, an apparently healthy *Diploria labyrinthiformis* sample, was removed
due to data quality problems, leaving 57 sets of paired-end samples for analysis. These were 20
diseased colony lesion (DL) samples, 20 diseased colony unaffected (DU) samples, and 17
apparently healthy colony (AH) samples from the coral species *D. labyrinthiformis*, *Dichocoenia*
stokesii, *Meandrina meandrites*, and *Stephanocoenia intersepta*. DU samples were taken from

apparently unaffected tissue from the diseased corals also sampled for DL. All samples were
collected and treated in the same manner, as previously described (Rosales *et al.* 2020, 2022). In
brief, coral tissue and mucus slurries were scraped from the coral surface with 10-ml plastic
syringes, transferred to plastic tubes on the boat, and held in a cooler on ice until being flash-
frozen in a liquid nitrogen dewar on shore; samples were held in a -80°C freezer until processed
for sequencing. Because all samples were from corals within reefs with an ongoing SCTLD
outbreak in the Florida Keys, it is possible that a primary pathogen of SCTLD could be present
in low abundance in at least one of the AH samples or that the AH microbiome was different
from that of corals in reefs where SCTLD had yet to arrive. Therefore, in this study, our findings
represent microbial communities associated with the observable surface tissue loss formation
stage of SCTLD (hereafter visual tissue loss or diseased lesion) compared to corals with no
visual signs of disease (i.e., AH).
148

The tissue samples from the four coral species were pooled by each of the three disease states,
resulting in twelve pooled read files (AH, DU, and DL for each of the four coral species). It was
assumed that a putative pathogen involved in visual tissue loss would likely show different
abundances in DL samples during different stages of lesion progression, so pooling the samples
was thought to increase the likelihood of observing a putative agent. All subsequent analyses
were based on these data, focusing primarily on the DL and AH samples. The reads from the DU
samples were explored in the preliminary analysis but were not considered in the final analysis.
Due to the proximity of the DU samples to lesion tissue, DU samples were considered likely to
represent early stages of surface tissue loss, and therefore poor choices for our methods.

Filtering reads with a healthy coral reference database

Because no sequenced genome was available for any of the four coral species, we created a
customized database to identify reads that likely originated from either the host genome or the
healthy host microbiome. To do this, we used reads from all AH samples to create a KrakenUniq
(Breitwieser *et al.* 2018) database for each coral species. We then used this database along with
KrakenUniq to classify reads from DL samples, thereby removing any read in DL samples that
matched any read in the AH samples. This filtering step produced a subset of diseased reads that
we considered unique to the DL samples, and greatly reduced the number of reads analyzed in
subsequent steps (Figure 1a).
167

The k-mer size for databases was set to 29 bp, lower than the default of 31bp, because we wanted
to filter more aggressively. For all other parameters, the default values of KrakenUniq were used.
For each coral species, the pooled DL reads were classified with KrakenUniq against the AH
reads database corresponding to that species. The original DL files were parsed to extract all
reads that were unclassified by the AH KrakenUniq database, generating the DL unique files
used for subsequent analysis (Figure 1a). With this aggressive filtration approach, we may have
lost information about changes in relative abundances between the AH and DL samples and were

therefore limited to only investigating species in high abundance, but we were left with reads that
were truly unique to the DL samples. This set would likely provide the clearest signal of
microbes associated with visual lesion formation, which was the primary goal of this study.

Genus-level classification with KrakenUniq

To first establish a baseline abundance profile before any filtering, the AH and DL reads were
classified with KrakenUniq (Breitwieser *et al.* 2018) using default parameters and the default k-
mer size of 31 bps against a microbial database (Figure 1a). The database used for classification
was built in August 2020 using all NCBI RefSeq complete bacterial, viral, and archaeal
genomes, the GRCh38 human genome, the NCBI UniVec database, and a curated set of
sequences from EuPathDB (Lu and Salzberg 2018; Amos *et al.* 2022). Then, the unique DL
reads were also classified with KrakenUniq using the same database and parameters. All reads
that were classified as *Homo* (human contamination) by KrakenUniq were removed from
subsequent analysis and results. If there are novel species associated with SCTLD, then their
genomes will not be present in public databases; however, if closely related species from the
same genera are available, then we might find DNA sequence-level matches to those genomes.
For this reason, the unique k-mers are reported at the genus level. Additionally, the relative
abundances of the genera are calculated by the unique k-mer count rather than the read count. In
general, using the unique k-mer count (i.e., sequences of length k are counted just once per
taxon, no matter how many times they occur in the raw data) rather than read count reduces the
bias introduced from using amplification-based sequencing workflows. Using the unique k-mer
count also reduces false positives that may arise from reads that contain low-complexity k-mers
(Breitwieser *et al.* 2018).
197
The report files were initially visualized and explored with Pavian (Breitwieser and Salzberg
2020). The read classifications were verified by randomly sampling 40 classified reads, aligning
them with megablast (Altschul *et al.* 1990) to the NCBI standard non-redundant (nr) nucleotide
database, and ensuring they had the same or similar classifications as with KrakenUniq. The
unique k-mers-per-read statistic served as a confidence flag. For a species that was truly present
in the sample, even with amplified metagenomic data, we expected a high number of unique k-
mers per read. A 150 bp read may contain up to 120 unique 31-mers, although repetitive k-mers
will reduce the unique count. There is also an upper bound to unique k-mers found in a genome,
which may be reached when the genome is small or when the sampling depth is high. In this
study, we considered a value of less than five unique k-mers per read as a flag that the taxon
might be a false positive. The unique k-mer-per-read count for every classified genus is reported
in Supp. Table S1.

Protein-level classification with MMseqs2

Because protein sequences are more conserved than DNA across distant species, we ran translated searches using MMSeqs2 easy-taxonomy workflow (Steinegger and Söding 2017) with the UniRef50 protein database (Suzek *et al.* 2015) to determine if this would identify more of the microbial reads than DNA sequences alone (Figure 1a). UniRef50 (Suzek *et al.* 2015) allowed for faster alignment than using the standard UniRef given that we aligned our protein sequences to clusters of similar protein sequences rather than all protein sequences. The paired reads had to be classified separately because MMseqs2 easy-taxonomy does not allow for both paired reads to be processed together. Additionally, MMSeqs2 does not report k-mer counts, only read counts, a metric that is subject to more bias from PCR amplification protocols.

To identify reads belonging to members of the algal symbiont family *Symbiodiniaceae*, which were among the most abundant reads (see Results), the MMseqs2 output was parsed to extract all protein cluster identifiers that had at least one alignment at the "f_Symbiodiniaceae" level or below for each coral species. The UniRef50 cluster identifiers were mapped to the full UniProtKB (UniProt Consortium, 2021). Their functions, if known, are reported in Supp. Table S2 as output by the UniProt ID mapping service (Huang *et al.* 2011). Supp. Table S2 was produced by inserting the number of alignments from the original MMseqs2 "tophit_report" files into the outputs of the UniProtKB ID mapping.

Contig assembly and classification

Due to the high genomic diversity of viruses (Aiewsakun *et al.* 2018), a viral agent might have been missed in our previous read analyses because it was too divergent from available DNA and protein sequences. This problem could be mitigated if the query sequences were longer, and therefore we assembled the raw reads to see if any long viral contigs were assembled. With our filtering approach, we did not expect to have the depth of coverage to assemble anything much larger than an abundant virus.

The filtered unique diseased reads from all four coral species were pooled to form a fasta file of all filtered unique diseased reads and were then assembled with MEGAHIT v1.2.9 (Li *et al.* 2016) using default parameters (Figure 1b). The contigs were classified with KrakenUniq using the same database of complete bacterial and viral genomes used above. The viral classifications from the report file were extracted to search for any viruses of interest. These steps were repeated with pooling just the filtered unique diseased reads excluding *S. intersepta,* because samples from this species represented a majority of the reads (89.8%) in our study and dominated the previous assembly (Rosales *et al*. 2022).

Candidate genus (*Vibrio*) investigation

To explore the utility of our pipeline in characterizing the SCTLD microbiome at a finer
taxonomic level, we selected a candidate bacterial genus, *Vibrio*, from the k-mers found most
abundant by KrakenUniq (see Results). We selected *Vibrio* for this exercise because this genus
was not emphasized in the previous metagenomic sequencing analysis of this same dataset
(Rosales *et al.* 2022), *V. coralliilyticus* coinfection increases the virulence of SCTLD (Ushijima
et al. 2020), and many *Vibrio* species are pathogenic to marine organisms broadly, not just to
corals (de Souza Valente and Wan 2021; Dincturk *et al.* 2023). First, the reads that were
classified to the *Vibrio* species level were investigated in further detail. To do this, the
KrakenUniq report files of the unique diseased reads for every coral species were parsed to
extract the number of unique k-mers assigned to each *Vibrio* species. The k-mer counts were
normalized by dividing them by the total number of k-mers assigned to the *Vibrio* genus within
the respective coral species. To investigate the prevalence of these *Vibrio* species across the
samples comprising coral species data, we calculated the relative proportion of different *Vibrio*
species for individual samples. The contribution of *Vibrio* species reads from each sample was
then found by parsing the KrakenUniq output to determine the sample ID number from the read
identifier. With these methods, it was not possible to determine the number of unique k-mers that
were contributed by each sample, therefore the read counts were reported. Second, to compare
our identified *Vibrio* reads with previous SCTLD *Vibrio* assemblies, the reads that were
classified at or below the *Vibrio* genus level by KrakenUniq were extracted from the original
sequence files for *S. intersepta* and *D. labyrinthiformis* only, because they contributed the
majority of the *Vibrio* genus reads (see Results). The draft genomes from an SCTLD study that
cultured *V. coralliilyticus* (Ushijima *et al.* 2020) were downloaded from NCBI Bioproject
PRJNA625269 (Benson *et al.* 2017) and a Bowtie2 index was built for each one. The extracted
Vibrio reads were aligned with Bowtie2 (Langmead and Salzberg 2012) to each of the eight draft
genomes. The Bowtie2 alignment rates to each draft genome are reported in Supp. Table S4.

Figure 1: (a) Filtering the diseased reads consisted of building a KrakenUniq database from all
healthy reads for each coral species and classifying the corresponding pooled diseased reads
against their respective database. That is every coral species was only filtered with its respective
"Healthy" Coral DB, and there was no cross-species filtration. Reads that were unclassified by
the database were considered unique to the diseased samples. This subset was classified at the
DNA level with KrakenUniq and at the protein level with MMseqs2. **(b)** The unique diseased
reads from all coral species were combined and assembled with MEGAHIT. The assembled
contigs were then classified with KrakenUniq.
280

Results

Filtering reads with a healthy coral reference database

We analyzed metagenomic reads from corals *D. labyrinthiformis, D. stokesii, M. meandrites,* and
S. intersepta from three different tissue sample types: diseased colony lesion (DL), diseased
colony unaffected (DU), and apparently healthy colony (AH). The total number of reads from
each species and sample type is shown in Table 1.
287

Table 1: Summary of total DNA sequencing reads from each coral species and tissue type. Rows labeled "Filtered" report the unique reads remaining after filtering out reads that overlapped with those found in apparently healthy samples, as described in the text. M=millions of reads.

Coral Species	Tissue Type (# of samples)	Read Counts (M)
Diploria labyrinthiformis	Apparently Healthy (4)	151.125
	Diseased Unaffected (5)	211.401
	Diseased Lesion (5)	198.816
	Filtered Diseased Lesion	0.955
Dichocoenia stokesii	Apparently Healthy (5)	179.831
	Diseased Unaffected (5)	202.168
	Diseased Lesion (5)	257.031
	Filtered Diseased Lesion	0.562
Meandrina meandrites	Apparently Healthy (3)	126.223
	Diseased Unaffected (5)	206.423
	Diseased Lesion (5)	185.556
	Filtered Diseased Lesion	4.159
Stephanocoenia intersepta	Apparently Healthy (5)	197.367
	Diseased Unaffected (5)	190.251

Diseased Lesion (5)	190.304
Filtered Diseased Lesion	50.028

Genus-level classification with KrakenUniq

Before any filtering, both the AH and the DL reads were profiled to establish baseline
abundances. The AH reads were dominated by *Synechococcus* (Supp. Fig S1). Although filtering
removed a significant majority (up to 99%) of the reads, the original DL microbial read
classifications did not have notable changes (Supp. Fig S2), indicating predominantly coral host
DNA was filtered from the DL samples, which led to substantial improvements in computational
speed in downstream analyses.

313
314
315 **Figure 2:** Unique k-mer counts from KrakenUniq genus-level classifications of unique diseased
316 reads for every coral species. The intensity of the shading represents the percent of total unique
317 k-mers assigned to the genus. Each box is annotated with the number of unique k-mers (in
318 thousands) assigned to the genus.
319
When the KrakenUniq reports for filtered DL samples were sorted by unique k-mer count within
coral species, the microbial genera *Synechococcus, Vibrio, Ruegeria, Phaeobacter,* and
Sulfitobacter were found at high proportions in all coral species, with the genera *Synechococcus*,
Vibrio, and *Ruegeria* being particularly abundant across all coral species (Figure 2).
Synechococcus was the most or second most abundant in all coral species. In *D. labyrinthiformis,*
Vibrio was the most abundant, with 4.1 times more unique k-mers than the second most
abundant genus, *Synechococcus*. In *D. stokesii*, *Synechococcus* was the most abundant, with
Vibrio being second most abundant. As in D. stokesii, *Synechococcus* was the most abundant in

M. meandrites, but in this coral *Pseudovibrio* was the second most abundant and was the only
coral in which this genus appeared in a high proportion. *Vibrio* had the sixth highest relative
abundance in *M. meandrites,* which, though lower than observed in the other coral species, still
represented a high k-mer-to-read ratio of 52.0 (Supp. Table S1). In *S. intersepta, Vibrio* was
again clearly the most abundant, having 4.3 times the amount of unique k-mers compared to the
second place *Synechococcus.* Due to their high abundances across all coral species in this
analysis, *Synechococcus*, *Vibrio,* and the *Rhodobacteraceae* family (to which *Ruegeria,*
Phaeobacter, and *Sulfitobacter* belong) appear to be associated with visual tissue loss and may
represent important agents of SCTLD. Another useful aspect of the KrakenUniq pipeline is that
by including the human reference genome in the database, human contamination can be detected
in the classification step, without the expensive additional step of aligning all reads to the human
reference genome. Human contamination, as reported by KrakenUniq, represented less than
0.2% of the filtered DL reads for all coral species. In the initial unfiltered DL and AH samples,
human contamination was reported to be between 1.5 - 3.0% of reads across samples, an
expected level for marine samples (Schmieder and Edwards 2011).
343

Protein-level classification with MMseqs2

The number of microbial reads classified for each coral species at the protein-level using
MMseqs2 (Steinegger and Söding 2017) increased approximately six-fold compared to the
DNA-based searches (Supp. Figure S3). Because we were primarily interested in whether any
new candidate taxa emerged, we did not consider the relative abundances of different taxa
classified by the protein-based search compared to the DNA-based search. Due to the decreased
specificity of a protein search, MMseqs2 classified many reads as "unclassified [family level]";
thus, the results are presented at the family rather than the genus level (Figure 3). As expected,
the read counts for each coral species were similar between the paired reads, which are classified
separately by MMSeqs2.

354
355 **Figure 3.** Classifications from MMseqs2 at the family-level for forward ("1") and reverse reads
356 ("2") from unique diseased reads from each coral species. The intensity of the shading represents
357 the proportion of total reads from the coral species that were assigned to the family. The boxes
358 are annotated with the number of reads (in thousands) assigned to each family.

359
MMseqs2 was able to classify a higher percentage of bacterial and eukaryotic reads compared to
the KrakenUniq DNA-level analysis (Supp. Figure 3), predominantly from coral algal symbionts
such as *Symbiodiniaceae*, yet we saw similar abundant taxa. *Symbiodiniaceae* was among the top
families in all coral species and particularly abundant (~48% of all classified reads) in *M.*
meandrites. Due to particular interest in the role of *Symbiodiniaceae* in SCTLD progression
(Landsberg *et al.* 2020, Beavers *et al.* 2023), the functions of the proteins in the *Symbiodiniaceae*
protein clusters identified are provided in Supp. Table S2.

367
The families *Flavobacteriaceae, Roseobacteraceae*, and *Paracoccaceae* were among the most
abundant families in all coral species. In this database, *Roseobacteraceae* and *Paracoccaceae* are
homotypic synonyms of the family *Rhodobacteraceae* in the database used for DNA-level
classifications (Göker 2022). So, we observed *Rhodobacteraceae* as before in the DNA analysis,
but *Flavobacteriaceae* emerged as another family of interest in this protein-level analysis.

Flavobacteriaceae was also found as a top family in the DNA-level classification (Supp. Figure
S4); however, no top genus was identified that belongs to this family. Additionally, the average
number of unique k-mers stemming from reads classified at or below the *Flavobacteriaceae*
family at the DNA-level was relatively low. For example, in *S. intersepta*, there were 794,258
unique k-mers from 489,794 reads, or ~1.6 k-mers per read. In *D. stokesii* there were 38,333
unique k-mers from 3,802 reads, or ~10 k-mers per read. Across all coral species, the
Flavobacteriaceae family had one of the lowest average k-mer-per-read counts of all bacterial
families identified. For example, the *Vibrionaceae* family, which has similar sized genomes to
Flavobacteriaceae (Lin *et al.* 2018; Gavriilidou *et al.* 2020), had 25 k-mers-per-read and 88 k-
mers-per-read in *S. intersepta* and *D. stokesii*, respectively.

Contig assembly and classification

In addition to characterizing the bacterial community, using metagenomic data made it possible
to explore DNA viruses found in the unique DL reads. To account for the genomic diversity of
viruses, which may not share many conserved sequences with genomes in public databases, we
assembled contigs from the unique DL reads and classified them with KrakenUniq to identify
any viral contigs that may be of interest in SCTLD etiology. This resulted in 1,014,402
assembled contigs. KrakenUniq classified 168,829 (16.6%) contigs, of which only 227 (0.02%)
were viruses. *Paracoccus* phages were the most abundant viral contigs, with *Synechococcus*
phages*, Dinoroseobacter* phages*, Vibrio* phages, and *Cyanophages* being abundant as well
(Figure 4). These phage results reflect bacteria that were found in high proportions in the results
above, providing support that the high abundances of the associated bacteria previously observed
were representative of the true metagenomic compositions of the samples. Finally, five contigs
were classified as *Chrysochromulina ericina* virus, a virus that infects the microalga
Chrysochromulina ericina (also known as *Haptolina ericina*) (Gallot-Lavallée *et al.* 2017), but
they represented only 108 unique k-mers and an assembly of 2.5 kb of a 473.6 kb genome
(Gallot-Lavallée *et al*. 2015). When aligned with BLASTN (Altschul et al. 1990) to standard
databases, only two contigs aligned best to *C. ericina* virus and the other three aligned best to
Eukarya, possibly indicating false positives and leading us to be skeptical of any significant
implications of this finding for SCTLD progression.
402

403
404 **Figure 4:** Classifications of assembled viral contigs showing the number of unique k-mers (in
405 thousands) and the total sequence length (in kb) assembled for every genus.

406
When combining the filtered diseased reads from every coral species for assembly, the reads
originating from *S. intersepta* samples represented a majority of the reads (89.8%). Therefore,
we repeated the previous steps without *S. intersepta* reads and assembled the reads from the
other three coral species. This resulted in 27,759 assembled contigs, of which 2,010 (7.2%) were
classified by KrakenUniq, and only six matched viruses (five *Pseudoalteromonas* phages and
one *Synechococcus* phage).

Candidate genus (*Vibrio*) investigation

To explore the utility of our pipeline at a finer taxonomic level, we selected a candidate genus,
Vibrio. We first investigated reads classified at the *Vibrio* species level. Vibrio reads were not
able to be classified to the species level in 6.8%, 1.0%, 8.0%, and 11% of *S. intersepta, D.*
labyrinthiformis, D.stokesii, and *M. meandrites*, respectively. The k-mers stemming from each
Vibrio species for each host coral species pooled are displayed in Figure 5a and an overview of
Vibrio read counts found in each colony sampled comprising the pooled coral species is shown
in Figure 5b. *V. europaeus* and *V. tubiashii*, represented a large portion (37% combined) of the
Vibrio k-mers in *S. intersepta* (Figure 5a). *V. mediterranei* dominated in *D. labyrinthiformis*
(96%) and *D. stokesii* (67%) but was muted in the other coral species (Figure 5a); however, one
colony per species appeared to be responsible for these high proportions: *D. labyrinthiformis*
colony 57 and *D. stokesii* colony 63 (Figure 5b). *V. sp. THAF190c* was the predominant *Vibrio*
species (17%) in *M. meandrites*. In general, other species like *V. coralliilyticus*, *V. harveyi*, *V.*
owensii, and *V. sp. THAF100* appeared consistently in all coral species, but never in high
proportions, but *V. tubiashii* and *V. owensii* contributed more than 5% of the reads in 10 samples
each (Figure 5b).

Figure 5. Species-level KrakenUniq classifications of *Vibrio* genus reads in SCTLD lesions. (**a**) The proportion of unique k-mers assigned to each *Vibrio* species grouped by samples from the same coral species. The size of the dots is relative to the proportion of unique k-mers assigned to the *Vibrio* species in each coral species. Those with over 5% are annotated with their proportion. (**b**) The species level read count proportions of the *Vibrio* for every sample. The size of the dots is relative to the proportion of read counts assigned to *Vibrio* species in each sample. The x-axis is labeled with the colony sample number, which corresponds to those assigned in the original analysis of this dataset (Rosales *et al.* 2020). Classifications that represent at least 5% of the *Vibrio* species in the sample are annotated with the number of reads.

Secondly, eight draft genomes from *V. coralliilyticus* strains isolated from a previous SCTLD study (Ushijima *et al.* 2020) allowed us to continue exploring the use of our pipeline and determine whether we identified the same *V. coralliilyticus* strains within our study. Reads classified by KrakenUniq as *Vibrio* in *S. intersepta* and *D. labyrinthiformis* were extracted and aligned to the Ushijima *et al.* (2020) draft genomes. Between 3.76% to 4.02% of the reads mapped to the *V. coralliilyticus* strains, while 7.54% of reads mapped to the McD22-P3 *Vibrio* strain, which was the control strain, and not a *V. coralliilyticus* strain (Supp. Table S4). As would

449 be expected, the proportions of reads that are aligned are similar to the proportions of reads that
450 were classified as *V. coralliilyticus* by KrakenUniq (Figure 5a).
451

Discussion

In this study, we used previously published sequencing data from coral affected by SCTLD and
developed a novel metagenomic analysis pipeline to explore the microbial communities present
in those data. The data consisted of samples from four coral species collected from Florida's
coral reefs during a SCTLD outbreak. To investigate the microbial taxonomy of these samples,
the previous study used small subunit rRNA gene assemblies and metagenome-assembled
genomes (MAGs). Our investigation differed by using the Kraken software suite and focusing on
unique k-mer count data to understand abundances. By predominantly working with k-mers
instead of MAGs, we maximized the utility of the read data, as our approach allowed us to
capture reads that may have been discarded during MAG assembly and binning. Samples with
high intrapopulation diversity can provide challenges in MAG assembly and binning (Ramos-
Barbero et al. 2019; Meziti et al. 2021), which may have hindered the generation of a MAG bin
in the previous analysis. The previous work also did not filter out host sequences because
reference genomes do not exist for the sampled coral species. However, here we applied a novel
technique to filter these reads by using data derived from apparently healthy (AH) samples as a
surrogate for reference genomes. This allowed us to examine unique sequences from DL samples
by approximating a species-specific host coral genome and reduce computational load. In
addition, in this study, we investigated the SCTLD DNA virome, which has not been previously
reported.
471

The families *Rhodobacteraceae* and *Flavobacteriaceae* were found to be associated with
SCTLD in our protein analysis, consistent with other SCTLD studies (reviewed in Papke *et al.*
2024), including previous examinations of the same sample set (Rosales *et al.* 2020, 2022).
Rhodobacteraceae is one of the most common bacterial families associated with coral diseases
(Gignoux-Wolfsohn *et al.* 2017) in diverse geographic locations, but no member has been
identified as a causative coral disease agent (Mouchka *et al.* 2010), and members of this group
are broadly found across ocean habitats and fulfill diverse ecological functions (Brinkhoff *et al.*
2008). *Flavobacteriaceae* has been enriched in White Band Disease in the Scleractinia staghorn
coral *Acropora cervicornis* (Gignoux-Wolfsohn and Vollmer 2015), but has never been
identified as a causative agent in coral tissue loss. In SCTLD, *Flavobacteriaceae* has previously
been found enriched in unaffected (non-lesion) tissue from diseased colonies, potentially
indicating colony stress and initial dysbiosis due to disease (Rosales *et al.* 2023). In contrast to
the protein analysis, which used read count data, our DNA-level analysis using k-mers did not
detect a singular genus belonging to *Rhodobacteraceae* or *Flavobacteriaceae* in high proportions
among the unique diseased reads, providing support for the idea that members of these two
families are likely a diverse set of bacteria species that associate with SCTLD opportunistically.
488

In our DNA-level analysis, *Vibrionaceae* and *Synechococcaceae* were among the most abundant
families within the unique DL reads and these high abundances showcase how the k-mer based
abundance approach can provide a different perspective of highly abundant taxa in particular
samples. These families were not observed as highly abundant in the MMSeqs2 protein analysis;
however, we were primarily interested in whether new candidates emerged from the MMSeqs2
analysis, not the relative proportions of the candidates. Interestingly, the genera *Synechococcus*
and *Vibrio* were not detected in the previous analysis of these data (Rosales *et al.* 2022). In the
16S rRNA metabarcoding study of these same samples, *Synechococcus* and *Vibrio* were
differentially abundant in *M. meandrites* samples (Rosales *et al.* 2020). These results did not
show a clear trend as *Synechococcus* was present across multiple AH samples and *Vibrio* was not
prevalent across DL coral samples. In this study, *Synechococcus* was the most or second most
proportionally abundant genus across all four coral species. *Synechococcus* belong to phylum
Cyanobacteria, which are photosynthetic picoplankton (Kim *et al.* 2018), and typically not
involved in pathogenesis. Even so, *Synechococcus* have been enriched in other SCTLD studies
that compared healthy colonies and healthy tissue on diseased colonies, and it was hypothesized
that their increase in abundance is a response to disease stress (Rosales *et al.* 2023). The high
proportion of *Synechococcus* in this study supports the suggestion that *Synechococcus* may have
some role in microbial community interactions during SCTLD.
507
Vibrio also have been associated with SCTLD (reviewed in Papke *et al.* 2024), including in these
samples (Rosales *et al.* 2020), but in contrast to *Synechococcus*, *Vibrio* have been associated
with other coral tissue loss diseases, such as white pox disease (Kemp *et al.* 2018) and yellow
band disease (Cervino *et al.* 2008), and diseases in other marine organisms, such as sponges
(Dincturk *et al.* 2023) and crustaceans (de Souza Valente and Wan 2021). In three coral species
analyzed here, *Vibrio* was one of the two most abundant genera, suggesting it may be associated
with SCTLD tissue loss. In the fourth species, *M. meandrites, Vibrio* was only the sixth most
abundant genus. However, *M. meandrites* had the fewest AH reads to create the database used
for filtering the DL reads (Table 1), which may reduce the effectiveness of our novel pipeline
that relies on reads from control groups. This was also a surprising result, since as mentioned,
Vibrio were abundant in the 16S rRNA *M. meandrites* analysis of these samples (Rosales *et al.*,
2020).
520
When using our pipeline to explore a species-level analysis within the candidate genus *Vibrio*,
we found read matches to *V. mediterranei/shilonii* (Tarazona et al. 2014), *V. coralliilyticus*, *V.*
harveyi, and *V. owensii* – all known coral pathogens associated with bleaching and tissue loss
(Kushmaro *et al.* 1997; Ben-Haim *et al.* 2003; Luna *et al.* 2007; Ushijima *et al.* 2012; Munn
2015). *V. mediterranei/shilonii*, which comprised a majority of the classified *Vibrio* species in *D.*
labyrinthiformis and *D. stokesii,* was previously found to be responsible for the annual bleaching
of the scleractinian coral *Oculina patagonica* off the Israeli coast from 1993 (Kushmaro *et al.*
1996, 1997) to 2003 (Reshef *et al.* 2006). Additionally, when *V. mediterranei*/*shilonii* is

experimentally introduced together with *V. coralliilyticus* to healthy corals, they appear to have
synergistic pathogenic effects (Rubio-Portillo *et al.* 2014). Given these associations, *V.*
mediterranei/shilonii, may be of particular interest in future SCTLD studies as potentially
increasing the virulence of SCTLD, as does *V. coralliilyticus* (Ushijima *et al.* 2020), which also
appeared in our samples. However, the very low similarity between the cultured *V. coralliilyticus*
sequences in Ushijima *et al.* (2020) and our *Vibrio* sequences leads us to believe that multiple
Vibrio species may be involved in SCTLD lesion development. It is important to note that our
picture of the SCTLD microbiome is restricted by the genomes in the databases used. The *Vibrio*
genus has been found to have a large degree of genetic plasticity between species (Gu et al.
2009), so while the classifications to different *Vibrio* species may truly represent the presence of
an array of *Vibrio* species, it may instead be the result of various reads from a novel *Vibrio*
species matching different *Vibrio* species based on closest genomic similarity. Therefore, while
matches to different *Vibrio* species and their potential role in SCTLD may offer some insights, a
more robust interpretation is to consider the implications of disease association by *Vibrio* at the
genus level.
544
The etiology of SCTLD has been hypothesized to be viral (Work *et al.* 2021), and gene
expression data show that there is an increase in coral viral immune response in corals with
SCTLD (Beavers *et al.* 2023). Researchers have explored the potential role of RNA viruses in
SCTLD, but no RNA viruses have been found exclusively in corals with SCTLD (Veglia *et al.*
2022) and these viruses are likely ubiquitous in corals without any potential relationship to
SCTLD (Howe-Kerr *et al.* 2023). However, the involvement of DNA viruses has not previously
been explored. Our data show the majority of DNA viruses in diseased samples represent phages.
Not surprisingly, phage sequences correspond with some of the most abundant bacteria identified
in this study, such as *Rhodobacteraceae, Vibrionaceae,* and *Synechococcaceae*. The *Paracoccus*
phage, which infects *Rhodobacteraceae,* and the *Vibrio* phage would be interesting to further
explore as potential avenues for disease mitigation. In addition to phages, sequences were found
with similarities to the *Chrysochromulina ericina* virus. However, with only two contigs and
little coverage of its genome, we do not believe this virus plays a role in SCTLD. Thus, we did
not find any DNA viruses with a definitive association with SCTLD. Future studies may
consider viral enrichment protocols prior to sequencing to help better characterize the SCTLD
DNA virome.
561
In addition to differences in the bacterial and viral communities, members of *Symbiodiniaceae*
were found to be abundant in DL samples, particularly in *M. meandrites*. SCTLD disrupts the
relationship between the host coral and its *Symbiodiniaceae* through symbiont necrosis and
peripheral nuclear chromatin condensation, among other physiological changes (Landsberg *et al.*
2020). This may result from an increase in *rab7* signaling among the *Symbiodiniaceae* to
degrade dead and dysfunctional cells through endocytic phagosomes (Beavers *et al.* 2023). The
Symbiodiniaceae DNA identified in diseased samples in this study may be a byproduct of this

necrosis and degradation of the symbiont. This was especially notable in *M. meandrites,* which
was the coral in this study most susceptible to acute tissue loss and mortality from SCTLD
(Precht *et al.* 2016); this accelerated tissue loss may lead to higher levels of dead and
dysfunctional symbionts being produced during visual lesion progression in *M. meandrites* than
in other coral species.
574
Although, this pipeline was initially developed under the one-pathogen-one-disease assumption
for humans, our results show that we can identify a consortium of putative pathogens with this
method (i.e., *Rhodobacteraceae*, *Flavobacteriaceae*, and *Vibrio*). However, this pipeline is
limited by the amount of prior information about pathogens in the species investigated; for
example, human pathogens are well represented in metagenomic classification databases,
whereas non-model organism pathogens may not be, which can lead to a higher rate of
unclassified or misclassified reads. In addition, disease states are more clearly defined in humans
than in non-model organisms, making the assumptions about sample health more reliable in
human analyses.
584

Conclusions

In all, we provide a novel pipeline to understand coral disease, and further investigate the role of
bacteria and DNA viruses in SCTLD. Our new pipeline found both incongruities and parallels
with the original analysis of these data. For instance, both studies revealed a prevalence of
Rhodobacteraceae and *Flavobacteriaceae* in lesion samples. One of the major differences
between the studies was the dominance of *Synechococcus* and *Vibrio* in this study, which was
likely driven by the use of k-mers to quantify taxonomic abundance (as the protein analysis with
read counts aligned more with previous results). In addition to providing novel insights, our
pipeline also reduced the computational load for downstream analysis, making large
metagenomic analysis more accessible to those with less computational resources. This method
is useful not only for coral scientists, but also for fields that study non-model organisms and lack
comprehensive genomic resources. In corals, the lack of such resources can impede the progress
of metagenomic analysis and the identification of potential pathogens. Specifically, host-
associated metagenomes without a reference genome can dominate and confound metagenomic
results (Rosales *et al.* 2022). As genome assemblies are unlikely to be available anytime soon for
the over 30 species of coral affected by SCTLD, this pipeline offers a useful tool for the
simultaneous examination of viruses, bacteria, and eukaryotic species living on or within the host
tissue that may be associated with infection for any coral species or non-model organism.
603

Data availability

The samples investigated in this study were downloaded from NCBI BioProject PRJNA576217.
Supplemental materials are available at FigShare. Table S1 contains the average k-mers per read
for every genus across all coral species. Table S2 contains the *Symbiodinaceae* protein

classifications. Table S3 contains the virulence factors identified in the *Vibrio* contigs. Table S4
contains the alignment rates of reads classified as *Vibrio* to eight draft assemblies of *Vibrio*
species previously isolated from SCTLD infected corals. The IDs of the unique diseased reads
remaining after filtering for each coral species and the assembled contigs are also available at
FigShare. The report files from the KrakenUniq and MMseqs2 analyses have been made
available at the following repository: https://github.com/jheinz27/coral_results/tree/main.

Acknowledgements

616 We thank Dr. Martin Steinegger and Dr. Christopher Pockrandt for their helpful advice and
617 exploratory work on pathogen identification in Mariana crows, which helped inform the methods
618 development here. We also thank Dr. Alaina Shumate for her valuable advice and guidance. The
619 samples were collected under permits #FKNMS-2018-057 and #FKNMS-2017-100.

Funding

622 This work was supported in part by NIH grants R35-GM130151 and R01-HG006677 to SLS.

Conflicts of interest

625 The authors declare that there is no conflict of interest.

Literature Cited

Aeby, G. S., B. Ushijima, J. E. Campbell, S. Jones, G. J. Williams *et al.*, 2019 Pathogenesis of a
tissue loss disease affecting multiple species of corals along the Florida reef tract. Front.
Mar. Sci. 6.: https://doi.org/10.3389/fmars.2019.00678

Aiewsakun, P., E. M. Adriaenssens, R. Lavigne, A. M. Kropinski, and P. Simmonds, 2018
Evaluation of the genomic diversity of viruses infecting bacteria, archaea and eukaryotes
using a common bioinformatic platform: steps towards a unified taxonomy. J. Gen. Virol.
99: 1331–1343. https://doi.org/10.1099/jgv.0.001110

Altschul, S. F., W. Gish, W. Miller, E. W. Myers, and D. J. Lipman, 1990 Basic local alignment
search tool. J. Mol. Biol. 215: 403–410. https://doi.org/10.1016/S0022-2836(05)80360-2

Alvarez-Filip, L., F. J. González-Barrios, E. Pérez-Cervantes, A. Molina-Hernández, and N.
Estrada-Saldívar, 2022 Stony coral tissue loss disease decimated Caribbean coral
populations and reshaped reef functionality. Commun Biol 5: 440.
https://doi.org/10.1038/s42003-022-03398-6

Amos, B., C. Aurrecoechea, M. Barba, A. Barreto, E. Y. Basenko *et al.*, 2022 VEuPathDB: the

eukaryotic pathogen, vector and host bioinformatics resource center. Nucleic Acids Res.

50: D898–D911. https://doi.org/10.1093/nar/gkab929

Beavers, K. M., E. W. Van Buren, A. M. Rossin, M. A. Emery, A. J. Veglia *et al.*, 2023 Stony

coral tissue loss disease induces transcriptional signatures of in situ degradation of

dysfunctional Symbiodiniaceae. Nat. Commun. 14: 2915. https://doi.org/10.1038/s41467-

023-38612-4

Ben-Haim, Y., F. L. Thompson, C. C. Thompson, M. C. Cnockaert, B. Hoste *et al.*, 2003 Vibrio

coralliilyticus sp. nov., a temperature-dependent pathogen of the coral Pocillopora

damicornis. Int. J. Syst. Evol. Microbiol. 53: 309–315. https://doi.org/10.1099/ijs.0.02402-

0

Benson, D. A., M. Cavanaugh, K. Clark, I. Karsch-Mizrachi, D. J. Lipman *et al.*, 2017 GenBank.

Nucleic Acids Res. 45: D37–D42. https://doi.org/10.1093/nar/gkw1070

Bourne, D. G., M. Garren, T. M. Work, E. Rosenberg, G. W. Smith *et al.*, 2009 Microbial

disease and the coral holobiont. Trends Microbiol. 17: 554–562.

https://doi.org/10.1016/j.tim.2009.09.004

Breitwieser, F. P., D. N. Baker, and S. L. Salzberg, 2018 KrakenUniq: confident and fast

metagenomics classification using unique k-mer counts. Genome Biol. 19: 198.

https://doi.org/10.1186/s13059-018-1568-0

Breitwieser, F. P., and S. L. Salzberg, 2020 Pavian: interactive analysis of metagenomics data for

microbiome studies and pathogen identification. Bioinformatics 36: 1303–1304.

https://doi.org/10.1093/bioinformatics/btz715

Brinkhoff, T., H.-A. Giebel, and M. Simon, 2008 Diversity, ecology, and genomics of the

Roseobacter clade: a short overview. Arch. Microbiol. 189: 531–539.

https://doi.org/10.1007/s00203-008-0353-y

Callahan, B. J., P. J. McMurdie, M. J. Rosen, A. W. Han, A. J. A. Johnson *et al.*, 2016 DADA2:

High-resolution sample inference from Illumina amplicon data. Nat. Methods 13: 581–583.

https://doi.org/10.1038/nmeth.3869

Cervino, J. M., F. L. Thompson, B. Gomez-Gil, E. A. Lorence, T. J. Goreau *et al.*, 2008 The

Vibrio core group induces yellow band disease in Caribbean and Indo-Pacific reef-building

corals. J. Appl. Microbiol. 105: 1658–1671. https://doi.org/10.1111/j.1365-

2672.2008.03871.x

Clark, A. S., S. D. Williams, K. Maxwell, S. M. Rosales, L. K. Huebner *et al.*, 2021

Characterization of the Microbiome of Corals with Stony Coral Tissue Loss Disease along

Florida's Coral Reef. Microorganisms 9.: https://doi.org/10.3390/microorganisms9112181

de Souza Valente, C., and A. H. L. Wan, 2021 Vibrio and major commercially important

vibriosis diseases in decapod crustaceans. J. Invertebr. Pathol. 181: 107527.

https://doi.org/10.1016/j.jip.2020.107527

Dinçtürk, E., F. Öndes, L. Leria, and M. Maldonado, 2023 Mass mortality of the keratose sponge

Sarcotragus foetidus in the Aegean Sea (Eastern Mediterranean) correlates with

proliferation of Vibrio bacteria in the tissues. Front. Microbiol. 14: 1272733.

https://doi.org/10.3389/fmicb.2023.1272733

Eberhart, C., Li, Z., Breitwieser, F. P., Lu, J., Jun, A. S. *et al.*, 2017 Diagnosing corneal

infections in formalin fixed specimens using next generation sequencing. Investigative

Ophthalmology & Visual Science.

Gallot-Lavallée, L., G. Blanc, and J.-M. Claverie, 2017 Comparative Genomics of

Chrysochromulina Ericina Virus and Other Microalga-Infecting Large DNA Viruses

Highlights Their Intricate Evolutionary Relationship with the Established Mimiviridae

Family. J. Virol. 91.: https://doi.org/10.1128/JVI.00230-17

Gallot-Lavallée, L., A. Pagarete, M. Legendre, S. Santini, R.-A. Sandaa *et al.*, 2015 The 474-

Kilobase-Pair Complete Genome Sequence of CeV-01B, a Virus Infecting Haptolina

(Chrysochromulina) ericina (Prymnesiophyceae). Genome Announc. 3.:

https://doi.org/10.1128/genomeA.01413-15

Gavriilidou, A., J. Gutleben, D. Versluis, F. Forgiarini, M. W. J. van Passel *et al.*, 2020

Comparative genomic analysis of Flavobacteriaceae: insights into carbohydrate

metabolism, gliding motility and secondary metabolite biosynthesis. BMC Genomics 21:

569. https://doi.org/10.1186/s12864-020-06971-7

Gignoux-Wolfsohn, S. A., F. M. Aronson, and S. V. Vollmer, 2017 Complex interactions

between potentially pathogenic, opportunistic, and resident bacteria emerge during

infection on a reef-building coral. FEMS Microbiol. Ecol. 93.:

https://doi.org/10.1093/femsec/fix080.

Gignoux-Wolfsohn, S. A., and S. V. Vollmer, 2015 Identification of Candidate Coral Pathogens

on White Band Disease-Infected Staghorn Coral. PLoS One 10: e0134416.

https://doi.org/10.1371/journal.pone.0134416

Gihawi, A., Y. Ge, J. Lu, D. Puiu, A. Xu *et al.*, 2023 Major data analysis errors invalidate cancer

microbiome findings. MBio 14: e0160723. e0160723. https://doi.org/10.1128/mbio.01607-

23

Göker, M., 2022 Filling the gaps: missing taxon names at the ranks of class, order and family.

Int. J. Syst. Evol. Microbiol. 72.: https://doi.org/10.1099/ijsem.0.005638

Gu, J., J. Neary, H. Cai, A. Moshfeghian, S. A. Rodriguez *et al.*, 2009 Genomic and systems

evolution in Vibrionaceae species. BMC Genomics 10 Suppl 1: S11.

https://doi.org/10.1186/1471-2164-10-S1-S11

Howe-Kerr, L. I., A. M. Knochel, M. D. Meyer, J. A. Sims, C. E. Karrick *et al.*, 2023

Filamentous virus-like particles are present in coral dinoflagellates across genera and ocean

basins. ISME J. https://doi.org/10.1038/s41396-023-01526-6

Huang, H., P. B. McGarvey, B. E. Suzek, R. Mazumder, J. Zhang *et al.*, 2011 A comprehensive

protein-centric ID mapping service for molecular data integration. Bioinformatics 27:

1190–1191. https://doi.org/10.1093/bioinformatics/btr101

Huntley, N., M. E. Brandt, C. C. Becker, C. A. Miller, S. S. Meiling *et al.*, 2022 Experimental

transmission of Stony Coral Tissue Loss Disease results in differential microbial responses

within coral mucus and tissue. ISME Communications 2: 1–11.

https://doi.org/10.1038/s43705-022-00126-3

Kim, Y., J. Jeon, M. S. Kwak, G. H. Kim, I. Koh *et al.*, 2018 Photosynthetic functions of

Synechococcus in the ocean microbiomes of diverse salinity and seasons. PLoS One 13:

e0190266. https://doi.org/10.1371/journal.pone.0190266

Kemp, K. M., J. R. Westrich, M. S. Alabady, M. L. Edwards, and E. K. Lipp, 2018 Abundance

and Multilocus Sequence Analysis of Vibrio Bacteria Associated with Diseased Elkhorn

Coral (Acropora palmata) of the Florida Keys. Appl. Environ. Microbiol. 84.:

https://doi.org/10.1128/AEM.01035-17

Kostic, A. D., D. Gevers, C. S. Pedamallu, M. Michaud, F. Duke *et al.*, 2012 Genomic analysis

identifies association of Fusobacterium with colorectal carcinoma. Genome Res. 22: 292–

298. https://doi.org/10.1101/gr.126573.111

Kushmaro, A., Y. Loya, M. Fine, and E. Rosenberg, 1996 Bacterial infection and coral

bleaching. Nature 380: 396–396. https://doi.org/10.1038/380396a0

Kushmaro, A., E. Rosenberg, M. Fine, and Y. Loya, 1997 Bleaching of the coral Oculina

patagonica by Vibrio AK-1. Mar. Ecol. Prog. Ser. 147: 159–165.

https://doi.org/10.3354/meps147159

Landsberg, J. H., Y. Kiryu, E. C. Peters, P. W. Wilson, N. Perry *et al.*, 2020 Stony coral tissue

loss disease in Florida is associated with disruption of host–zooxanthellae physiology.

Front. Mar. Sci. 7.: https://doi.org/10.3389/fmars.2020.576013

Langmead, B., and S. L. Salzberg, 2012 Fast gapped-read alignment with Bowtie 2. Nat.

Methods 9: 357–359. https://doi.org/10.1038/nmeth.1923

Li, D., R. Luo, C.-M. Liu, C.-M. Leung, H.-F. Ting *et al.*, 2016 MEGAHIT v1.0: A fast and

scalable metagenome assembler driven by advanced methodologies and community

practices. Methods 102: 3–11. https://doi.org/10.1016/j.ymeth.2016.02.020

Lin, H., M. Yu, X. Wang, and X.-H. Zhang, 2018 Comparative genomic analysis reveals the

evolution and environmental adaptation strategies of vibrios. BMC Genomics 19: 135.

https://doi.org/10.1186/s12864-018-4531-2

Liu, B., D. Zheng, S. Zhou, L. Chen, and J. Yang, 2022 VFDB 2022: a general classification

scheme for bacterial virulence factors. Nucleic Acids Res. 50: D912–D917.

https://doi.org/10.1093/nar/gkab1107

Lu, J., N. Rincon, D. E. Wood, F. P. Breitwieser, C. Pockrandt *et al.*, 2022 Metagenome analysis

using the Kraken software suite. Nat. Protoc. 17: 2815–2839.

https://doi.org/10.1038/s41596-022-00738-y

Lu, J., and S. L. Salzberg, 2018 Removing contaminants from databases of draft genomes. PLoS

Comput. Biol. 14: e1006277. https://doi.org/10.1371/journal.pcbi.1006277

Luna, G. M., F. Biavasco, and R. Danovaro, 2007 Bacteria associated with the rapid tissue

necrosis of stony corals. Environ. Microbiol. 9: 1851–1857. https://doi.org/10.1111/j.1462-

2920.2007.01287.x

Meziti, A., L. M. Rodriguez-R, J. K. Hatt, A. Peña-Gonzalez, K. Levy et al., 2021 The

Reliability of Metagenome-Assembled Genomes (MAGs) in Representing Natural

Populations: Insights from Comparing MAGs against Isolate Genomes Derived from the

Same Fecal Sample. Appl. Environ. Microbiol. 87. https://doi.org/10.1128/AEM.02593-20

Meyer, J. L., J. Castellanos-Gell, G. S. Aeby, C. C. Häse, B. Ushijima *et al.*, 2019 Microbial

Community Shifts Associated With the Ongoing Stony Coral Tissue Loss Disease

Outbreak on the Florida Reef Tract. Front. Microbiol. 10: 2244.

https://doi.org/10.3389/fmicb.2019.02244

Mouchka, M. E., I. Hewson, and C. D. Harvell, 2010 Coral-associated bacterial assemblages:

current knowledge and the potential for climate-driven impacts. Integr. Comp. Biol. 50:

662–674. https://doi.org/10.1093/icb/icq061

Munn, C. B., 2015 The Role of Vibrios in Diseases of Corals. Microbiol Spectr 3.:

https://doi.org/10.1128/microbiolspec.VE-0006-2014

National Center for Biotechnology Information (NCBI) Scleractinia [Internet]. Bethesda (MD):

National Library of Medicine (US), National Center for Biotechnology Information; [1988]

[cited 2023 Nov 26]. Available from:

https://www.ncbi.nlm.nih.gov/datasets/taxonomy/6125/

Neely, K. L., K. A. Macaulay, E. K. Hower, and M. A. Dobler, 2020 Effectiveness of topical

antibiotics in treating corals affected by Stony Coral Tissue Loss Disease. PeerJ 8: e9289.

https://doi.org/10.7717/peerj.9289

Papke, E., A. Carreiro, C. Dennison, J. M. Deutsch, L. M. Isma *et al.*, 2024 Stony coral tissue

loss disease: a review of emergence, impacts, etiology, diagnostics, and intervention. Front.

Mar. Sci. 10.: https://doi.org/10.3389/fmars.2023.1321271

Precht, W. F., B. E. Gintert, M. L. Robbart, R. Fura, and R. van Woesik, 2016 Unprecedented

Disease-Related Coral Mortality in Southeastern Florida. Sci. Rep. 6: 31374.

https://doi.org/10.1038/srep31374

Quast, C., E. Pruesse, P. Yilmaz, J. Gerken, T. Schweer *et al.*, 2013 The SILVA ribosomal RNA

gene database project: improved data processing and web-based tools. Nucleic Acids Res.

41: D590–6. https://doi.org/10.1093/nar/gks1219

Reshef, L., O. Koren, Y. Loya, I. Zilber-Rosenberg, and E. Rosenberg, 2006 The coral probiotic

hypothesis. Environ. Microbiol. 8: 2068–2073. https://doi.org/10.1111/j.1462-

2920.2006.01148.x

Ramos-Barbero, M. D., A.-B. Martin-Cuadrado, T. Viver, F. Santos, M. Martinez-Garcia et al.,

2019 Recovering microbial genomes from metagenomes in hypersaline environments: The

Good, the Bad and the Ugly. Syst. Appl. Microbiol. 42: 30–40.

https://doi.org/10.1016/j.syapm.2018.11.001

Rosales, S. M., A. S. Clark, L. K. Huebner, R. R. Ruzicka, and E. M. Muller, 2020

Rhodobacterales and Rhizobiales Are Associated With Stony Coral Tissue Loss Disease

and Its Suspected Sources of Transmission. Front. Microbiol. 11: 681.

https://doi.org/10.3389/fmicb.2020.00681

Rosales, S. M., L. K. Huebner, A. S. Clark, R. McMinds, R. R. Ruzicka *et al.*, 2022 Bacterial

Metabolic Potential and Micro-Eukaryotes Enriched in Stony Coral Tissue Loss Disease

Lesions. Frontiers in Marine Science 8.: https://doi.org/10.3389/fmars.2021.776859

Rosales, S. M., L. K. Huebner, J. S. Evans, A. Apprill, A. C. Baker *et al.*, 2023 A meta-analysis

of the stony coral tissue loss disease microbiome finds key bacteria in unaffected and lesion

tissue in diseased colonies. ISME Commun 3: 19. https://doi.org/10.1038/s43705-023-

00220-0

Rubio-Portillo, E., P. Yarza, C. Peñalver, A. A. Ramos-Esplá, and J. Antón, 2014 New insights

into Oculina patagonica coral diseases and their associated Vibrio spp. communities. ISME

J. 8: 1794–1807. https://doi.org/10.1038/ismej.2014.33

Salzberg, S. L., F. P. Breitwieser, A. Kumar, H. Hao, P. Burger *et al.*, 2016 Next-generation

sequencing in neuropathologic diagnosis of infections of the nervous system. Neurol

Neuroimmunol Neuroinflamm 3: e251. https://doi.org/10.1212/NXI.0000000000000251

Schmieder, R., and R. Edwards, 2011 Fast identification and removal of sequence contamination

from genomic and metagenomic datasets. PLoS One 6: e17288.

https://doi.org/10.1371/journal.pone.0017288

Shilling, E. N., I. R. Combs, and J. D. Voss, 2021 Assessing the effectiveness of two intervention

methods for stony coral tissue loss disease on Montastraea cavernosa. Sci. Rep. 11: 8566.

https://doi.org/10.1038/s41598-021-86926-4

Steinegger, M., and J. Söding, 2017 MMseqs2 enables sensitive protein sequence searching for

the analysis of massive data sets. Nat. Biotechnol. 35: 1026–1028.

https://doi.org/10.1038/nbt.3988

Studivan, M. S., R. J. Eckert, E. Shilling, N. Soderberg, I. C. Enochs *et al.*, 2023 Stony coral

tissue loss disease intervention with amoxicillin leads to a reversal of disease-modulated

gene expression pathways. Mol. Ecol. 32: 5394–5413. https://doi.org/10.1111/mec.17110

Suzek, B. E., Y. Wang, H. Huang, P. B. McGarvey, C. H. Wu *et al.*, 2015 UniRef clusters: a

comprehensive and scalable alternative for improving sequence similarity searches.

Bioinformatics 31: 926–932. https://doi.org/10.1093/bioinformatics/btu739

Tarazona, E., T. Lucena, D. R. Arahal, M. C. Macián, M. A. Ruvira *et al.*, 2014 Multilocus

sequence analysis of putative Vibrio mediterranei strains and description of Vibrio

thalassae sp. nov. Syst. Appl. Microbiol. 37: 320–328.

https://doi.org/10.1016/j.syapm.2014.05.005

UniProt: the universal protein knowledgebase in 2021, 2021 Nucleic Acids Res. 49: D480–

D489. https://doi.org/10.1093/nar/gkaa1100

Ushijima, B., A. Smith, G. S. Aeby, and S. M. Callahan, 2012 Vibrio owensii induces the tissue

loss disease Montipora white syndrome in the Hawaiian reef coral Montipora capitata.

PLoS One 7: e46717. https://doi.org/10.1371/journal.pone.0046717

Ushijima, B., J. L. Meyer, S. Thompson, K. Pitts, M. F. Marusich *et al.*, 2020 Disease

Diagnostics and Potential Coinfections by Vibrio coralliilyticus During an Ongoing Coral

Disease Outbreak in Florida. Front. Microbiol. 11: 569354.

https://doi.org/10.3389/fmicb.2020.569354

Veglia, A. J., K. Beavers, E. W. Van Buren, S. S. Meiling, E. M. Muller *et al.*, 2022

Alphaflexivirus Genomes in Stony Coral Tissue Loss Disease-Affected, Disease-Exposed,

and Disease-Unexposed Coral Colonies in the U.S. Virgin Islands. Microbiol Resour

Announc 11: e0119921. https://doi.org/10.1128/mra.01199-21

Walton, C. J., N. K. Hayes, and D. S. Gilliam, 2018 Impacts of a regional, multi-year, multi-

species coral disease outbreak in southeast Florida. Front. Mar. Sci. 5.:

https://doi.org/10.3389/fmars.2018.00323

Wilson, M. R., H. A. Sample, K. C. Zorn, S. Arevalo, G. Yu *et al.*, 2019 Clinical Metagenomic

Sequencing for Diagnosis of Meningitis and Encephalitis. N. Engl. J. Med. 380: 2327–

2340. https://doi.org/10.1056/NEJMoa1803396

Work, T. M., T. M. Weatherby, J. H. Landsberg, Y. Kiryu, S. M. Cook *et al.*, 2021 Viral-like

particles are associated with Endosymbiont pathology in Florida corals affected by stony

coral tissue loss disease. Front. Mar. Sci. 8.: https://doi.org/10.3389/fmars.2021.750658

Number of Reads (in thousands)

Family Classifications	D. lab_1	D. lab_2	D. stok_1	D. stok_2	M. mean_1	M. mean_2	S. inter_1	S. inter_2	Median
Flavobacteriaceae	20.6	19.1	12.1	11.6	29.2	28	1290.7	1229.9	24.3
Roseobacteraceae	7.2	6.8	8.2	7.9	20.6	19.5	1271.6	1209.2	13.85
Paracoccaceae	5.5	5.1	9.5	8.9	18.6	17.9	1039.1	994.3	13.7
Symbiodiniaceae	10.8	9.8	4.1	2.7	511.4	478.5	129.1	120.7	65.75
Harpacticidae	2.6	2.6	2.6	2.5	0.4	0.4	183.4	176.3	2.6
Vibrionaceae	5.9	5.7	0.5	0.5	3.1	3.1	248.9	243.4	4.4
Pocilloporidae	6.8	6.1	2.4	2	8.5	7.6	20.4	19.3	7.2
Saprospiraceae	2.7	2.6	1.4	1.4	6.2	5.8	100.4	97.6	4.25
Nymphonidae	1.2	1.1	1.4	1.2	3	2.8	161.6	155	2.1
Synechococcaceae	1.6	1.5	1.5	1.5	7.1	6.7	21.8	21.1	4.15
Planctomycetaceae	1.4	1.3	3	2.9	3.6	3.5	79.8	76.9	3.25
Eubacteriales Family XII	0.2	0.1	1.3	1.3	1.4	1.3	354.7	340.2	1.3
Merulinidae	7.1	6.6	1	0.7	5.4	5	2.6	2	3.8
Acroporidae	3.1	2.7	0.8	0.6	4.6	4.1	6	5.4	3.6
Chitinophagaceae	2.7	2.7	0.7	0.6	2.2	2.1	45	43.7	2.45
Pirellulaceae	0.8	0.8	2.2	2.1	1.9	1.9	48.1	46	2
Stappiaceae	0.3	0.3	0.4	0.4	18.8	18.3	62.7	59.4	9.35
Suessiaceae	0.6	0.6	0.4	0.3	28.1	26.4	10	9.5	5.05
Oceanospirillaceae	0.6	0.6	0.3	0.2	3.4	3.3	357.2	347.6	1.95
Cellvibrionaceae	0.3	0.2	0.2	0.2	0.7	0.8	177.3	169.8	0.5
Arcobacteraceae	0.2	0.2	0.1	0.1	1.4	1.4	504.8	490.1	0.8
Portunidae	0.2	0.2	0.1	0.1	13	12.7	5.2	5.3	2.7
Pseudobacteriovoracaceae	0.1	0.1	1.9	1.8	0.1	0.1	4.2	4.1	0.95
Majidae	0.1	0.1	0	0	14.3	13.9	1.6	1.7	0.85
Varunidae	0	0	0	0	10.3	10	1	1	0.5

Coral Species

Percent of Reads in Each Coral Species: >15%, 12%, 9%, 6%, 3%