Chromosome Structure II: Stem-loops and circle-loops.

Wenfan Ke[1]*, Miki Fujioka[2]*, Paul Schedl[1], and James B. Jaynes[2]

1: Department of Molecular Biology, Princeton University, Princeton NJ 08544

2: Department of Biochemistry and Molecular Biology, Thomas Jefferson University,

Philadelphia, PA 19107

* These authors contributed equally.

Abstract

The chromosomes in multicellular eukaryotes are organized into a series of topologically independent loops called TADs. In flies, TADs are formed by physical interactions between neighboring boundaries. Fly boundaries exhibit distinct partner preferences, and pairing interactions between boundaries are typically orientation dependent. Pairing can be head-to-tail or head-to-head. The former generates a stem-loop TAD, while the latter gives a circle-loop TAD. The TAD that encompasses the *Drosophila even skipped* (*eve*) gene is formed by the head-to-tail pairing of the *nhomie* and *homie* boundaries. To explore the relationship between loop topology and the physical and regulatory landscape, we flanked the *nhomie* boundary region with two attP sites. The attP sites were then used to generate four boundary replacements: λ *DNA*, *nhomie forward* (WT orientation), *nhomie reverse* (opposite of WT), and *homie forward* (same as WT *homie*). The *nhomie forward* replacement restores the WT physical and regulatory landscape: In MicroC experiments, the *eve* TAD is a volcano triangle topped by a plume, and the *eve* gene and its regulatory elements are sequestered from interactions with neighbors. The λ *DNA* replacement lacks boundary function: the endpoint of the "new" *eve* TAD on the *nhomie* side is ill-defined, and *eve* stripe enhancers activate a nearby gene, *eIF3j*. While *nhomie reverse* and *homie forward* restore the *eve* TAD, the topology is a circle-loop, and this changes the local physical and regulatory landscape. In MicroC experiments, the *eve* TAD interacts with its neighbors, and the plume at the top of the *eve* volcano triangle is replaced by a cloud of contacts with the next-door TADs. Consistent with the loss of isolation afforded by the stem-loop topology, the *eve* enhancers weakly activate genes in the neighboring TADs. Conversely, *eve* function is partially disrupted.

Introduction

Chromosomes in multicellular animals are organized into a series of topologically independent looped domains, called TADs (topologically associated domains: (Cavalheiro et al., 2021; Chetverina et al., 2017; Jerković et al., 2020; Matthews and White, 2019; Rowley and Corces, 2018). The arrangement of TADs in a given chromosomal DNA segment is generally (though not precisely) similar in different tissues and developmental stages, and this is a reflection of the mechanism underlying TAD formation—the endpoints of TADs are determined by a special class of elements called chromatin boundaries or insulators. While boundary-like elements have been identified in a wide range of animals and plants, the properties of this class of DNA elements have been most fully characterized in *Drosophila* (Cavalheiro *et al.*, 2021; Chetverina *et al.*, 2017). Fly boundaries have one or more large (100-400 bp) nucleosome-free nuclease hypersensitive sequences that are targets for multiple DNA binding chromosomal architectural proteins. While only a single chromosomal architectural protein, CTCF, has been characterized in mammals, there are almost two dozen such proteins in flies and the list is still growing (Heger et al., 2013; Heger and Wiehe, 2014; Schoborg and Labrador, 2010). In addition to subdividing the chromosome into a series of loops, fly boundary elements have insulating activity. When placed between enhancers or silencers and their target promoters, boundaries block regulatory interactions (Bell et al., 2001; Chetverina et al., 2014; Chetverina *et al.*, 2017). This activity provides a mechanism for delimiting units of independent gene activity: genes located between a pair of compatible boundaries are subject to regulatory interactions with enhancers/silencers present in the same chromosomal interval, while they are insulated from the effects of enhancers/silencers located beyond either boundary in adjacent regulatory neighborhoods. It is currently thought that their ability to organize the chromosome into

topologically independent loops is important for their insulating activity (Cai and Shen, 2001;

Gohl et al., 2011; Muravyova et al., 2001).

Studies dating back to the '90's have suggested that fly boundaries subdivide the

chromosome into loops by physically pairing with each other (Chetverina *et al.*, 2014;

Chetverina *et al.*, 2017). In these first experiments, regulatory interactions were observed for

transgenes inserted at distant sites in chromosome that were carrying either the *gypsy* transposon

boundary *su(Hw)* or the bithorax complex (BX-C) boundary *Mcp* (Muller et al., 1999; Sigrist and

Pirrotta, 1997; Vazquez et al., 1993). Further support for the idea that boundaries function by

pairing has come from chromatin immunoprecipitation, chromosome conformation capture

(CCC), MicroC, and direct imaging experiments (Chen et al., 2018; Li et al., 2011; Vazquez et

al., 2006). More recent studies have revealed physical interactions in the CNS, such as those

found for *su(Hw)* and *Mcp,* which "reach over" multiple intervening TADs, consistent with them

playing an important role in cell type-specific gene regulation (Mohana et al., 2023) by bringing

distant enhancers and promoters together.

The parameters governing pairing interactions have been defined using insulator bypass,

transvection and boundary competition assays. These studies have shown that fly boundaries are

able to pair not only with heterologous boundaries but also with themselves. Moreover, the

pairing interactions typically exhibit a number of characteristic features: promiscuity coupled

with clear partner preferences, and orientation dependence.

Partner preferences depend upon the chromosomal architectural proteins that interact

with each boundary. For example, in the boundary bypass assay, a set of enhancers are placed

upstream of two reporters (Cai and Shen, 2001; Kyrchanova et al., 2008a; Muravyova *et al.*,

2001). When multimerized CTCF sites are placed between the enhancers and the closest

reporters, both reporters are insulated from the enhancers. When a second set of multimerized

CTCF sites are place downstream of the closest reporter, bypass is observed. In this case the

closest reporter, which is bracketed by the multimerized CTCF sites, is still insulated from the

enhancers; however, the downstream reporter is activated (Muravyova *et al.*, 2001).

Heterologous combinations give a different result: when multimerized dCTCF sites are placed

upstream of the closest reporter and multimerized Zw5 sites are placed downstream, no bypass is

observed. Endogenous fly boundaries also show partner preferences in bypass assays and in

boundary competition experiments (Gohl *et al.*, 2011; Kyrchanova et al., 2011; Kyrchanova et

al., 2008b). On the other hand, while boundaries have partner preferences, they are also

promiscuous in their ability to establish functional interactions with other boundaries. For

example, the *Fab-8* insulator can pair with *scs'* from the *Drosophila* heat shock locus (Gohl *et*

al., 2011).

In addition, to partner preferences, pairing interactions between endogenous fly

boundaries are, with a few exceptions, orientation dependent. Self-pairing interactions are head-

to-head. This seems to be a common feature of fly boundaries and has been observed for *scs*,

scs', *iA2*, *wari*, *Mcp*, *Fab-8*, *AB-I, homie* and *nhomie* (Fujioka et al., 2016; Kyrchanova *et al.*,

2008a). In contrast, pairing interactions between heterologous boundaries can be head-to-head

or head to tail. The two boundaries bracketing the *even-skipped* (*eve*) locus, *homie* and *nhomie*,

pair with each other head-to-tail, while boundaries in the *Abdominal-B* (*Abd-B*) region of BX-C

pair with their neighbors head-to-head. The topology of the loops (TADs) generated by head-to-

tail and head-to-head pairing between neighboring boundaries in *cis* are different. As illustrated

in Fig. 1, head-to-tail pairing generates stem-loops, while head-to-head pairing generates circle-

loops. The loops could be connected to each other by unanchored loops (Fig. 1A and C), or they

could be linked directly to each other if boundaries can pair simultaneously with both neighbors

(Fig 1B and D). An alternating pattern of TADs connected by DNA segments that crosslink to

each other with reduced frequency (c.f., λ DNA below) is not observed in MicroC experiments.

Instead, most TADs appeared to be directly connected to both of their neighbors without an

intervening unanchored loop (Batut et al., 2022; Bing et al., 2023; Levo et al., 2022): see also

below). This would suggest that TAD boundaries are typically linked to both neighbors, either

simultaneously or as alternating pair-wise interactions.

Key to understanding the 3D organization of chromosomes in multicellular eukaryotes

will be the identification of TADs that are stem-loops and TADs that are circle-loops. In the

studies reported here, we have used MicroC to analyze the contact maps generated by stem-loops

and circle-loops. Stem-loops and circle-loops TADs are expected to interact differently with

their neighbors and this should be reflected in the patterns of crosslinking events between

neighboring TADs. As illustrated for linked stem-loops in Fig. 1B, TAD2, is isolated from its

next-door neighbors, TAD1 and TAD3. In this configuration, crosslinking events between

sequences in TAD2 and sequences in TAD1 and TAD3 will be suppressed. On the other hand,

TAD1 and TAD3 are in comparatively close proximity, and crosslinking between sequences in

these two TADs will be enhanced. A different pattern of neighborly interactions is expected for

circle-loop TADs. In this case, the TAD in the middle, TAD2, is expected to interact with both

of its neighbors (Fig. 1D). To test these predictions, we have first compared the MicroC contact

profiles for stem-loop and circle-loop TADs. For stem-loops we selected the *eve* TAD, while for

circle-loops we chose the four TADs that comprise the *Abd-B* parasegment-specific regulatory

domains. We show that these stem-loop and circle-loop TADs have distinctive crosslinking

signatures. To confirm these MicroC signatures we converted the topology of the *eve* TAD from

a stem-loop to a circle-loop. In addition to changing the MicroC signature of the *eve* TAD, the

change in topology is accompanied by changes in the regulatory interactions between *eve* and its

neighbors.

Results

Stem-loops versus circle-loops

The distinctive loop topologies of stem-loops and circle-loops are expected to be reflected in

the contact maps that are generated in MicroC experiments. To determine if this is the case, we

compared the MicroC contact maps for the *eve* TAD and the TADs that correspond to the four

Abd-B parasegment specific regulatory domains, *iab-5*, *iab-6*, *iab-7* and *iab-8*. The *eve* TAD is

generated by pairing interactions between the *nhomie* boundary upstream of the *eve* transcription

unit and the *homie* boundary downstream. Since *nhomie* and *homie* pair with each other head-to-

tail, the *eve* TAD has a stem-loop topology. Unlike the *eve* boundaries, the boundaries that

delimit the *Abd-B* regulatory domains are thought to pair with their neighbors head-to-head

(Chetverina *et al.*, 2017; Kyrchanova *et al.*, 2008a; Kyrchanova *et al.*, 2011; Kyrchanova *et al.*,

2008b)). This means that the parasegment-specific regulatory domain TADs, *iab-5*, *iab-6*, *iab-7*

and *iab-8*, are expected to have a circle-loop topology.

As shown in Fig. 2, the *eve* TAD and the four *Abd-B* TADs have distinctive MicroC contact

patterns. The *eve* TAD is a volcano triangle with a plume. The endpoints of the volcano triangle

are delimited by *nhomie* on the left and *homie* on the right, and within the *eve* locus (the

volcano), there are additional enhanced interactions. While the volcano triangle is generated by

contacts between sequences within the *eve* stem-loop, contacts between sequences in *eve* and in

the neighboring TAD on the left, TL (which contains multiple sub-TADs and six genes:

CG15863, CG1418, *Pal1, CG12133, eIF3j* and *CG12134*), are much reduced (L-ev in Fig 2A).

There is a similar suppression of contacts between sequences in the *eve* TAD and sequences in

the large neighboring TAD on the right, TM (which contains *TER94* and *Pka-R2*) (ev-M in Fig

2A). On the other hand, as expected from the regulatory interactions observed for stem-loops in

boundary bypass experiments (Kyrchanova *et al.*, 2008a), physical contacts between sequences

in TL and TM are enhanced compared to those between *eve* and TL (L-ev) or TM (ev-M).

Because of the preferential interactions between TADs to either side of the *eve* stem-loop, the

eve volcano triangle is topped by a plume (L-M in Fig. 2A). TM also interacts with the two of

TADs farther to the left of *eve*, TK (K-M in Fig. 2A) and TJ (J-M in Fig. 2A) (see also Fig.

Supplemental 1).

Like the *eve* boundaries, the TADs in the *Abd-B* region of BX-C region are connected to

their neighbors by the boundaries at their base. As predicted from genetic studies on BX-C

boundaries, each TAD corresponds to one of the four parasegment-specific regulatory domains.

While the MicroC contact maps for the four *Abd-B* TADs resemble the contact patterns in the

eve TAD, these *Abd-B* TADs differ from *eve* in that there are no plumes at the top of their

volcano triangle (Fig. 2B). Instead, the *Abd-B* TADs are overlaid by a series of rectangular

interlocking low-density contact (LDC) domains, or clouds. As illustrated in Fig 2B, the *iab-6*

regulatory domain is surrounded by clouds generated by crosslinking with its next door

neighbors *iab-5* (5-6) and *iab-7*(6-7), followed by crosslinking with neighbors that are a TAD

away from *iab-6*, *iab-4* (4-6) and *iab-8* (6-8). The other regulatory domains also form a unique

set of interlocking LDCs/clouds with their immediate neighbors, their next-next door neighbors,

and their next-next-next door neighbors.

TAD formation in a nhomie deletion

To further investigate the pairing properties and functioning of the *eve* boundaries, we used

CRISPR-Cas9 to add two attP sites flanking the *nhomie* region, replacing the region with a mini-

white gene. Using mini-*white* as an exchange maker, ΦC31 recombinase-mediated cassette

exchange (RMCE) was used to restore the sequence of the region, with *nhomie* modifications.

As a control for possible effects of the sequences introduced in generating the modification, we

reinserted a 597 bp *nhomie* DNA fragment in the same orientation as the endogenous *nhomie*

boundary (*nhomie forward*). To maintain roughly the same distance between *eve* and the

neighboring TAD in the *nhomie* deletion, we introduced a 606 bp DNA fragment from phage

λ (*λ DNA*). Fig. 3 (and Fig. Supplemental 1) shows the MicroC contact profiles for the *nhomie*

forward and *λ DNA* replacements. Except that the sequencing depth of the *nhomie forward*

replacement is not as great as the WT shown for the *eve* locus in Fig. 2, the profile is quite

similar. Like WT, there are sub-TADs within the *eve* TAD. One of these appears to link *homie*

and the neighboring PRE to the *eve* promoter-proximal PRE (Fujioka et al., 2008), and is marked

by an interaction dot (asterisk in Fig. 3A). Another links *nhomie* to the *eve* promoter region

(blue arrow in Fig. 3A). The *eve* TAD is topped by a plume which is generated by interactions

between sequences in neighboring TADs TL with TM (L-M). On the other hand, interactions

between *eve* and its neighbors are suppressed. Like *eve*, there are sub-TADs in the neighboring

TAD, TL. The TL sub-TAD closest to *eve* (TL4) corresponds to the *CG12134* transcription unit

(green arrowhead marks the boundary: Fig. 3), while the neighboring sub-TAD (TL3)

encompasses the *eIF3j* transcription unit (blue arrowhead: Fig. 3).

The MicroC profile of the *λ DNA* replacement (Fig. 3D) is quite different from that of

either *nhomie forward* or WT, which are similar (Figs. 2A and 3A). While *homie* still defines

the distal (relative to the centromere) end of the *eve* locus, the *λ DNA* replacement does not

function as a TAD boundary, and the leftward endpoint of the *eve* TAD is no longer well-

defined. One new "endpoint" for the *eve* locus maps to sequences between *CG12134* and *eIF3j*

(green arrowhead), which in wild type correspond to the left boundary of the TL sub-TAD TL4.

The other endpoint maps to sequences between *eIF3j* and *CG12133* (blue arrowhead), which in

wild type define the left boundary of the TL sub-TAD TL3. These interactions are not as stable

as those between *nhomie* and *homie*, as the density of interaction dots is lower. Furthermore,

they appear to flip back and forth between alternative endpoints (as indicated by the green and

blue double arrows in Fig. 3F) based on the MicroC contact profile, which is consistent with a

mixture of (at least) two conformations. The *eve* TAD also interacts with sequences in the two

other TL sub-TADs, TL1 and TL2. In addition, the *eve* promoter appears to interact with

sequences located upstream of *CG12134* (purple arrowhead in Fig. 3D and double arrow in Fig.

3F), while this interaction is not observed in the *nhomie forward* replacement. While the TL

TAD (from the TK:TL boundary to *nhomie*) is also disrupted by the *nhomie* deletion (it has a

much less distinct "volcano apex", and its right-most sub-TAD TL4 is now fused with the *eve*

TAD), the left-most TL sub-TADs (TL1, TL2, and TL3) are still present, indicating that their

formation does not depend on *nhomie*. As shown in the virtual 4C at *homie* viewpoint in Fig. 3B

and Fig. 3E, the *homie* boundary interacts with the *nhomie forward* replacement, but does not

contact the *λ DNA* replacement.

***eve* enhancers activate *eIF3j* expression in the *nhomie* deletion**

In transgene assays, boundary elements block regulatory interactions when interposed been

enhancers (or silencers) and reporter genes (Chetverina *et al.*, 2014; Chetverina *et al.*, 2017;

Kellum and Schedl, 1992). To determine whether this is also true in their endogenous context,

we compared the expression in syncytial blastoderm embryos of the two genes that flank *nhomie*

at the *eve* locus, *eIF3j* and *CG12134*. As the *nhomie* deletion eliminates *homie*'s pairing partner

and disrupts the *eve* TAD, we also examined the expression of *eve* and of the gene just beyond

the *homie* boundary, *TER94*, which has strong maternal expression through stage 11 (Fig.

Supplemental 3B, WT). Consistent with the seemingly normal MicroC profile on the *homie* side

of the *eve* TAD, we did not detect evidence of *eve*-like *TER94* expression (Figs. 4 and 5). Thus,

the formation and functioning of the TM TAD does not appear to be impacted by the loss of

nhomie and the fact that left end of the *eve* TAD is no longer properly anchored.

In the case of the gene closest to the *nhomie* deletion, *CG12134*, we were unable to

consistently detect transcription driven by the *eve* enhancers in either *nhomie forward* or *λ DNA*

embryos. In some *λ DNA* embryos, there were hints of stripes at the blastoderm stage (see Fig.

Supplemental 2); however, these "stripes" were not observed in most embryos. Since *CG12134*

(which forms the TL4 sub-TAD in wild type) is closest to the *eve* enhancers and interacts most

strongly, it is possible that the promoter is not compatible with the *eve* enhancers. A different

result was obtained for *eIF3j* in *λ DNA* embryos. As shown in the HCR-FISH experiment in Fig.

4 and quantitated in Fig. 5, we observed a series of *eIF3j* stripes over a dark background in pre-

cellular blastoderm embryos. As this background hybridization is evident in earlier stages, much

of it is likely to be of maternal origin. In contrast to the *λ DNA* replacement, these stripes are not

visible in the *nhomie forward* (control) replacement (Figs 4 and 5). While it is possible to detect

all seven stripes in *λ DNA* blastoderm stage embryos, their levels of expression are not equal.

The highest levels correspond to *eve* stripes 1, 2, 3 and 7, while *eve* stripes 4, 5 and 6 are

expressed at much lower levels. Since the stripe enhancers for 1, 2, 3 and 7 are located between

the *eve* promoter and *nhomie*, they are closer to the *eIF3j* promoter than the enhancers for stripes

4, 5 and 6, which are located downstream of the *eve* transcription unit. In addition to possible

effects of distance, the subdomain linking the *eve* PRE and *homie* to the promoter is still

observed in the *λ DNA* replacement, and this could partially sequester the enhancers located

downstream of the *eve* promoter.

We also used digoxigenin *in situ* hybridization to analyze *eIF3j* expression. With this

procedure we were able to detect a very low level of *eve*-activated *eIF3j* stripe expression in WT

and *nhomie forward* embryos at stage 5 and stages 7-8 (Fig. Supplemental 3). In stage 5

embryos when *eve* expression is driven by specific stripe enhancers, *eIF3j* expression appears to

be similar in all seven stripes (Fig. Supplemental 3). In contrast, as was observed in the HCR-

FISH experiments (Fig. 4), there is a clear bias for enhancers located upstream of *eve* in the λ

DNA replacement at this point in development. In stage 7-8 embryos, the seven-stripe enhancer

drives *eve* expression. It is located close to *nhomie* and, not surprisingly, high levels of *eIF3j*

expression are observed in all seven stripes in the λ *DNA* replacement (Fig. Supplemental 3). At

later embryonic stages, *eve* expression is driven by tissue-specific neurogenic, mesodermal, and

anal plate enhancers. However, *eIF3j* is expressed at high levels in a complex pattern in older

embryos, and we were unable to unambiguously detect expression driven by the *eve* enhancers

over this background mRNA. This is also not surprising, given that all of the enhancers for these

aspects of *eve* expression are located downstream of the *eve* promoter, like the enhancers driving

stage 5 stripes 4, 5, and 6.

While the *nhomie* deletion did not have any obvious impact on the level or pattern of *eve*

expression in blastoderm stage embryos (see Fig. 4), it seemed possible that *eve* activity was not

entirely normal. To test this possibility, we mated males homozygous for either λ *DNA* or

nhomie forward to females heterozygous for a chromosomal deficiency that includes the *eve*

gene. We then quantitated the number of missing denticle bands in embryonic cuticle preps. As

shown in Fig. 5, the frequency of larvae with "severe" defects (2 or more missing ventral

denticle bands) in *nhomie forward* is similar to that in a WT *yw* control. In contrast, in the λ

DNA replacement, the frequency of larval cuticles with 2 or more missing denticle bands is

increased nearly two-fold. Taken together, the increase in severity of the cuticle defects is

significant at the $P > 0.01$ level (one-tailed t-test). The A6 denticle belt is missing most

frequently followed by A2, A4, and then A8. These even-numbered denticle bands are those that

are lost in *eve* deficiency mutants (from which the name *even skipped* comes), suggesting that

eve stripe expression at blastoderm stages is compromised in the embryos that produce these

defective cuticles.

The eve TAD is converted from a stem-loop to a circle-loop by inverting nhomie

The orientation of boundary:boundary pairing interactions determines the topology of

each chromatin loop (Bing *et al.*, 2023; Fujioka *et al.*, 2016). Since *nhomie* and *homie* pair with

each other head-to-tail, the endogenous *eve* TAD is a stem-loop. This orientation dependence

means that one can convert the *eve* TAD from a stem-loop to a circle-loop by inverting the

nhomie boundary. If our expectations are correct, the MicroC contact pattern will also be

transformed from a volcano triangle with a plume to one in which sequences in the *eve* TAD is

surrounded by a cloud of crosslinked sequences from both neighboring TADs (TL and TM), like

that observed in the *Abd-B* region of BX-C.

We tested this prediction by inserting the *nhomie* boundary in the reverse orientation

(*nhomie reverse*). Fig. 6A shows that the *eve* TAD is reconstituted by *nhomie reverse* (compare

with Fig. 3). The sub-TAD evident in the *nhomie forward* replacement linking *nhomie* to the *eve*

promoter is also re-established (blue arrow). In addition, consistent with our expectation, the

plume topping the *eve* TAD is gone and is replaced by a much more sparsely populated LDC

domain (purple double arrow and above). The more prominent LDC TAD-TAD interactions (the

clouds) are between sequences in the *eve* TAD and the neighboring TADs. On the right, *eve*

forms an LDC interaction domain with TM (ev-TM). On the left, *eve* interacts strongly with

sequences in TL4, and progressively less strongly with sequences in the sub-TADs TL3, TL2,

and then TL1. In addition to restoring the *eve* TAD, the TL TAD is also re-established,

indicating that the *nhomie* boundary is important in defining both endpoints of the TL TAD. On

the other hand. with the exception of the *CG12133* sub-TAD, *nhomie* does not play a role in

generating the three other sub-TADs in the TL TAD. The other interesting feature is a 45º band

of interaction (to the left and below the purple double-arrow) between sequences in TL4 and

sequences in *eve* that appear to be located near the left edge of *homie*, and likely correspond to

the distal *eve* PRE (Fujioka *et al.*, 2008). A similar though weaker 45º band is also seen in

nhomie forward (Fig. 3).

Insulation is reduced in nhomie reverse

Consistent with the models for stem-loops and circle-loops in Fig. 1, the neighborly

interactions evident in the MicroC contact patterns for *nhomie forward* and *nhomie reverse* are

quite distinct. The *nhomie forward* TAD is isolated from its neighbors, and crosslinking between

eve and the neighboring TADs is suppressed (Fig. 1B and 2A). This is not true for the circle-

loop TAD generated by *nhomie reverse*: in this configuration, the *eve* TAD is not sequestered

from neighboring TADs, but instead interacts much more frequently with sequences in next-door

TADs than in the stem-loop configuration. Since the *eve* TAD is no longer isolated from its

neighbors, this could increase the frequency of "productive" interactions between *eve* regulatory

elements and genes in nearby TADs, and vice versa.

To test whether the circle-loop topology has an impact on the regulatory landscape, we

hybridized *nhomie reverse* embryos with HCR-FISH probes for *eIF3j*, *TER94* and *eve*. In early

blastoderm stage *nhomie forward* embryos, there is little evidence of *eIF3j* or *TER94* expression

driven by *eve* stripe enhancers, and the HCR-FISH hybridization pattern is uniform (see Fig. 4).

In contrast, it is possible to discern individual stripes of both *eiF3j* and *TER94* mRNA over the

background signal in a subset of *nhomie reverse* blastoderm stage embryos in HCR-FISH (Fig.

7). Since these genes are assembled into their own topologically independent looped domains

rather than being in the same domain as *eve*, the level of *eiF3j* and also *TER94* stripe expression

is lower than that observed in the *nhomie* deletion (λ *DNA*). The *nhomie reverse* circle-loop also

differs from the *nhomie* deletion (λ *DNA*) in that there is not such an obvious preference for

which *eve* enhancers activate expression. In addition, *eve*-dependent *eiF3J* and *TER94* stripes are

detected in only about half of the blastoderm stage embryos (Fig. 5A). It is possible that the

frequency of productive inter-TAD contacts differs from one embryo to the next; however, a

more likely reason is that the high background of *eIF3j* and *TER94* transcripts obscures the low

level of *eve* enhancer-driven expression at this stage. Once the seven-stripe enhancer is

activated, *eve*-dependent *TER94* expression in *nhomie reverse* is elevated, and all seven stripes

are observed (Fig. Supplemental 3B). This fits with the MicroC contact profile. As shown in

Fig. 6, the *TER94* gene is preferentially crosslinked to *eve* sequences located between *nhomie*

reverse and the *eve* promoter as compared to sequences spanning the *eve* gene and the

downstream enhancers. By contrast, there is much less seven-stripe enhancer-driven expression

of *eIF3j* (Fig. Supplemental 3A) than *TER94*. Consistent with this observation, crosslinking

between *eIF3j* and sequences in *eve* close to *nhomie reverse* and the seven-stripe enhancer

appear to occur less frequently than crosslinking to sequences on the other side of the *eve* TAD

(though this crosslinking bias is not as pronounced as that observed between *TER94* and the *eve*

sequences near the seven-stripe enhancer relative to downstream *eve* sequences.)

While *eve* stripe expression is not discernably different from wild type, the circle-loop

topology still impacts *eve* function. As shown in Fig. 5B, the fraction of *nhomie reverse*

embryos with 2 or more missing denticle bands is nearly twice that in *yw* or the *nhomie forward*

replacement. Taken together, the increase in severity of the cuticle defects is significant at the P

> 0.01 level (one-tailed t-test). As was observed for the *λ DNA* replacement, the A6 denticle

band is missing most frequently, followed by A2, A4 and then A8.

homie forward converts the eve stem-loop to a circle-loop

While the findings in the previous section show that loop topology impacts how

sequences in TADs interact with each other and with their neighbors, one might argue that the

effects we observed are a reflection of some novel properties of the *nhomie* boundary when it is

inverted. To test this possibility, we took advantage of the fact that in addition to pairing with

nhomie, the *homie* boundary pairs with itself (Fujioka *et al.*, 2016). However, unlike

nhomie:homie pairing which is head-to-tail, *homie:homie* pairing is head-to-head. This means

that it is possible to convert the *eve* TAD into a circle-loop by inserting *homie* into the *nhomie*

deletion in the forward orientation.

As shown in Fig. 6C, the MicroC contact profile of *homie forward* is similar to that

observed for *nhomie reverse*. The plume toping the *eve* TAD in wild type (Fig. 2) or in *nhomie*

forward (Fig. 4) is absent. Likewise, instead of being isolated from its neighbors, the *eve* TAD

contacts TL and TM. Also like *nhomie reverse*, *homie forward* forms a subdomain within the

eve TAD linking it to sequences in the *eve* promoter. There is also a 45° band of crosslinking

between sequences in TL4 and sequences on the distal side of *eve* that appears to correspond to

the *eve* 3' PRE.

The MicroC pattern is not the only similarity between *homie forward* and *nhomie*

reverse. The functional properties of the *homie forward eve* TAD are also similar. Fig. 7 shows

that the *eve* enhancers weakly activate both *eIF3j* and *TER94*. As was the case for *nhomie*

reverse, expression levels at the blastoderm stage are low and are observed in only about half of

the embryos (Fig. 5A). After the blastoderm stage, when the seven-stripe enhancer drives *eve*

expression, an even higher level of *TER94* expression is observed (Fig. Supplemental 3B). In

addition, the functioning of the *eve* gene when it is in the circle-loop configuration is not as

efficient, and the frequency of *homie forward* embryos with 2 or more missing denticle bands is

twice that of *nhomie forward* (Fig. 5B). Taken together, the increase in severity of the cuticle

defects is significant at the $P > 0.01$ level (one-tailed t-test).

Discussion

Two different though overlapping classes of chromosomal architectural elements have

been identified in flies. One class is the PREs found in many developmental loci. PREs were

first discovered because they induce pairing sensitive silencing of reporter genes (Americo et al.,

2002; Kassis et al., 1991). More recent studies have shown that the ability of these elements to

physically pair with each other may be their most important function (Batut *et al.*, 2022; Levo *et

al.*, 2022). The other class of architectural elements are chromatin boundaries (also called

insulators). PRE pairing in *cis* typically takes place within the context of a larger chromosomal

domain, or TAD. In contrast, boundary elements are responsible for defining the endpoints of

these looped domains (Arzate-Mejía et al., 2020; Batut *et al.*, 2022; Bing *et al.*, 2023; Chetverina

et al., 2017; Ibragimov et al., 2023; Stadler et al., 2017). While not much is known about the

parameters governing PRE pairing, the pairing interactions of fly boundaries have been studied

in some detail. The key features include an ability to engage in promiscuous pairing interactions,

distinct partner preferences, and orientation dependence. Of the endogenous (non-*gypsy*)

boundaries whose functional properties have been studied in detail, only one*, Fab-7*, appears to

be able to pair in both orientations. However, *Fab-7* may be unusual in that its boundary activity

depends upon factors that have been implicated in the functioning of PREs (Kyrchanova et al.,

2018). For all of the other boundaries studied so far, pairing interactions are orientation

dependent. When fly boundaries pair with themselves, the interactions are head-to-head

(Kyrchanova *et al.*, 2008a). This make sense, as the available evidence suggests that self-pairing

interactions in *trans* may be largely responsible for the pairing of homologs in precise register

(Erokhin et al., 2021; Fujioka *et al.*, 2016). In this case, head-to-tail self-pairing would

uncoupled the loops on the two homologs.

Unlike self-pairing in *trans*, pairing interactions between heterologous boundaries in *cis*

can be head-to-head or head-to-tail. The topological consequences are quite distinct. The former

generates a circle-loop, while the later forms a stem-loop (Chetverina *et al.*, 2017).   In the

studies reported here, we have investigated how these two different topologies impact the local

chromatin organization. We have also determined whether circle-loops and stem-loops alter the

ability of boundary elements to define units of independent gene activity and insulate against

regulatory interactions between neighboring TADs.

nhomie deletion disrupts the eve TAD

As would be predicted from many different studies (c.f., (Cavalheiro *et al.*, 2021;

Chetverina *et al.*, 2017), deletion of the *nhomie* boundary and replacement with a control λDNA

disrupts the *eve* TAD and alters the regulatory landscape. The MicroC profile shows that

disruption of the *eve* TAD and the neighboring TADs is one sided. Within the *eve* TAD, the

subdomain linking *homie* and the nearby PRE to the *eve* promoter is unaffected. Likewise the

large TAD, TM, which encompasss both *TER94* and *pka-R2*, and is defined at one end by

homie and at the other by an uncharacterized boundary element upstream of the *pka-R2*

promoter, is intact (Fig. Supplemental 1B). In contrast, on the *nhomie* side of the *eve* TAD, the

sub-TAD linking *nhomie* to the *eve* promoter is absent, and is replaced by a less well-defined

sub-TAD linking the *eve* promoter to an element near the *CG12134* promoter. However, the

end-point of the *eve* TAD is no longer distinct, and *eve* sequences are now crosslinked to the

eIF3j sub-TAD and to sequences in the more distant TL sub-TADs, TL3, TL2 and TL1.

Consistent with the alterations in the physical organization of the *eve* and neighboring TADs, the

TER94 gene is still insulated from the *eve* enhancers. While sequences in the *eve* TAD

physically interact with the gene closest to *nhomie*, *CG12134*, only the next gene over, *eIF3j*, is

clearly activated by the *eve* enhancers. Since crosslinking between sequences in the *CG12134*

and the *eve* TAD is more frequent than crosslinking between the *eIF3j* sub-TAD TL3 and the *eve*

TAD, it seems likely that *CG12134* is refractory to activation by the *eve* enhancers. This could

be due to an incompatiblity between the *CG12134* promoter and the *eve* enhancers.

Alternatively, the promoter may not be active at this stage. While *eIF3j* is activated by the *eve*

enhancers in stage 5 embryos in the *nhomie* deletion, the stripe enhancers located upstream of the

eve gene drive a higher level of expression than those located downstream. Two factors in

addition to the effects of distance could potentially account for this finding. Since the *eve*

promoter is located between the downstream enhancers and the *eIF3j* gene, activation of *eIF3j*

by the downstream enhancers could be suppressed by promoter competition. Alternatively, or in

addition, the sub-TAD formed between the 3' PRE/*homie* and the *eve* promoter/proximal PRE

(Fujioka *et al.*, 2008) could tend to sequester the downstream *eve* stripe enhancers away from

interactions with *eIF3j*.

Topology impacts local 3D genome organization and the potential for regulatory interactions

In boundary bypass experiments using endogenous fly boundaries, the ability of the

upstream enhancers to activate the downstream reporter depended on the topology of the loop

generated by the paired boundaries (Kyrchanova *et al.*, 2008a). Activation is observed for stem-

loops, as this configuration brings the upstream enhancers into close proximity with the

downstream reporter. In contrast, the enhancers and downstream reporter are not brought into

contact when the topology is a circle-loop. As would be predicted from these bypass

experiments, the stem-loop formed by the head-to-tail pairing of *nhomie* and *homie* physically

isolates the *eve* TAD from its neighbors, and this is reflected in the low density of contacts

between sequences in *eve* and the neighboring TADs (Fig. 2A). Conversely, the TADs that flank

eve are brought together and contacts between them generate the plume that is observed above

the *eve* volcano triangle.

The physical isolation afforded by the stem-loop topology is lost when the *eve* TAD is

converted to a circle-loop, either by inverting the *nhomie* boundary or by replacing *nhomie* with

the *homie* boundary inserted in the forward direction (Fig. 6). In the former case, head-to-tail

nhomie:*homie* pairing generates a circle-loop. In the later case, head-to-head pairing of *homie*

(inserted in the forward orientation) with endogenous *homie* generates a circle-loop. The

alteration in the local 3D organization induced by the conversion of *eve* to a circle-loop is

evident from the changes in the MicroC contact pattern. Instead of being isolated from

neighboring TADs, the *eve* TAD interacts not only with its immediate neighbors, but also with

more distant TADs. As a result, the plume of enhanced contacts linking TM to TL, TK, and TJ

is absent and is replaced by contacts between these TADs and the *eve* TAD.

As might be expected from the MicroC contact patterns, the conversion to a circle-loop

topology is accompanyied by alterations in regulatory interactions between *eve* and the genes in

the neighboring TADs (Fig. 7 and Fig. Supplemental 3). Unlike the *nhomie forward*

replacement, the *eve* stripe enhancers in both of the circle-loop replacements are able to weakly

activate expression of two neighboring genes, *eIF3j* and *TER94*. This pattern of activation

mirrors the enhancement in contacts between the *eve* TAD and the neighboring TL and TM

TADs evident in MicroC experiments. Thus, though *eIF3j* and *TER94* are clearly shielded from

the *eve* enhancers when the *eve* TAD is a circle-loop, a greater degree of isolation from the

action of the *eve* enhancers is afforded when the *eve* TAD is a stem-loop.

In addition to reducing insulation from regulatory interactions with genes in neighboring

TADs, the circle-loop topology impacts the functioning of the *eve* gene (Fig. 5B). For both

nhomie reverse and *homie forward*, the frequency of multiple denticle band defects compared to

the *nhomie forward* control is enhanced in a sensitized genetic background. While we did not

detect any obvious reductions in the *eve* stripes in blastoderm stage embryos, the circle-loop

provides less insulation than the stem-loop, and it is possible that the neighboring promoters

suppress *eve* expression by competing for the *eve* enhancers. Another (non-mutually exclusive)

possibility comes from the studies of Yokoshi et al. (Yokoshi et al., 2020), who used live

imaging to examine the effects of flanking a reporter with the *nhomie* and *homie* boundaries. In

their experiments, reporter expression was enhanced over two-fold when the reporter was

flanked by *nhomie* and *homie*; however, the enhancement was greater when the paired

boundaries formed a stem-loop than when they formed a circle-loop.

Stem-loops versus circle-loops

The results we have reported here raise several questions. One of these is, what is the

relative frequency of stem-loops versus circle-loops in the fly genome? The MicroC contact

patterns for the stem-loop and circle-loop versions of the *eve* TAD are quite distinct. The former

is a volcano triangle with a plume while the latter is a volcano triangle flanked by clouds. A

survey of the MicroC contact patterns elsewhere in the non-repetitive regions of the fly genome

indicates that volcanoes with plumes are rare (~30). For example, there are two volcano triangles

with plumes in the *Antennapedia* complex and they encompass the *deformed* and *fushi-tarazu*

genes (Levo *et al.*, 2022). However, since most of the "euchromatic" regions of the fly genome

are assembled into TADs whose MicroC profiles resemble that observed for *eve* circle-loops and

the *Abd-B* region of BX-C, it possible that much of the fly genome is assembled into circle-

loops, not stem-loops.

While this suggestion is consistent with the available data, it is based on contact patterns

between neighboring TADs, and important caveats remain. For one, the contact patterns

between neighboring TADs can deviate in one way or another from that seen in the *Abd-B*

region. For example, there are TADs in which interactions with one set of neighbors appears to

be suppressed as expected for stem-loops, but the classic plume is absent, as interactions are not

suppressed with the other neighbors (c.f., Fig. Supplemental 4A). In other cases, there are a

series of complicated TAD-TAD interactions topped by a rectangular plume (Fig. Supplemental

4B: purple arrow). For this reason, it will not be possible to draw firm conclusions about the

frequency of stem-loops versus circle-loops genome-wide until the relative orientation of the

paired boundaries themselves can be determined directly. On the other hand, it is clear from our

studies that both classes of TADs must exist in flies. If, as seems likely, a significant fraction of

the TADs genome-wide are circle-loops, this would effectively exclude cohesin-based loop

extrusion as a general mechanism for TAD formation in flies. In addition, though stem-loops

could be generated by a cohesin-dependent mechanism, it is unlikely that this mechanism is

operational in flies, as we have shown here and in Bing et al. (Bing *et al.*, 2023) that stem-loops

in flies are formed by orientation-dependent boundary:boundary pairing.

Another important question is whether our findings have any relevance to the formation

and topology of TADs in mammals. In the most common version of the loop extrusion model,

the mammalian genome is assembled into an alternating pattern of stem-loops and unanchored

loops (c.f., (Davidson and Peters, 2021; Higashi and Uhlmann, 2022; Perea-Resa et al., 2021).

In this case, one might expect to observe volcano triangles topped by plumes alternating with

DNA segments that have a considerablly lower densisty of internal contacts. However, this

crosslinking pattern is not observed in published MicroC data sets (Hsieh et al., 2020;

Krietenstein et al., 2020). Instead of an alternating pattern of high-density TAD triangles

separated by regions of low density contacts, the TAD triangles are generally linked to both

neighbors, just as in flies. Moreover, also like in Drosophila, there are very few stem-loop

volcano TADs topped by plumes. Instead, the crosslinking pattern between neighboring TADs

appears similar to that observed for circle-loops in flies. Of course, one problem with these

MicroC studies is that the resolution may not be sufficient to detect volcanos with plumes or the

other features predicted by the loop extrusion model. However, there are no obvious volcanoes

with plumes in the much higher resolution RCMC studies of Goel et al. (Goel et al., 2023).

Instead the MicroC profiles most closely resemble those seen in the *Abd-B* region of BX-C (c.f.

the *Ppm1g* locus in Fig. 4 of Goel et al., (Goel *et al.*, 2023)). Moreover, compromising cohesin

activity has minimal impact on the TADs in this region of the mouse genome, as evidenced from

the MicroC pattern before and after knockdown. Based on these observations, one can

reasonably question whether cohesin-mediated loop extrusion is deployed in mammals as the

mechanism for not only generating TADs but also determining TAD boundaries. Clearly

validation of the loop extrusion/CTCF road-block model as currently formulated will require a

direct demonstration that mammalian TADs are exclusively either stem-loops or unanchored

loops, and that the endpoints are always (or almost always) determined by CTCF roadblocks.

Acknowledgements: We would like to thank the Gordon Grey for running the fly food

facility at Princeton, members of the Lewis Sigler Genomics Core facility for their invaluable

assistance with DNA sequencing, and Qing Liu for excellent technical assistance. We would

also like to thank members of MOL431 for creative input. Special thanks to Olga Kyrchanova,

Daria Chetverina, Maksim Erokhin, Pavel Georigev, Tsutomu Aoki, Girish Deshpande, Airat

Ibragimov, Sergey Ryabichko, Yuri Pritykin, Alex Ostrin, Xinyang Bing, Xiao Li and Mike

Levine for stimulating discussions and sharing unpublished data.

 Materials and methods

 Key resources table

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Chemicals, peptides, and recombinant proteins		
n-Heptane	Fisher Chemical	O3008-4
Paraformaldehyde 20% solution, EM Grade	Electron Microscopy Sciences	15713-S
Formaldehyde, 16%, methanol free, Ultra Pure	Polysciences Inc	18814-10
PBS - Phosphate-Buffered Saline (10X) pH 7.4, RNase-free	Thermal Fisher	AM9624
Tween 20	Sigma	P1379
Triton X-100	Bio-Rad	161-0407
Tris base	Sigma	11814273001
Methanol	Fisher Chemical	203403
SSC, 20X	Thermal Fisher	15557044
Formamide	Thermal Fisher	17899
Dextran sulfate	Sigma	D8906
Salmon Sperm DNA	Thermal Fisher	AM9680
Ribonucleoside Vanadyl Complex	NEB	S1402S
Nuclease-free BSA	Sigma	126609
Triethylammonium Acetate	Sigma	625718
dGTP (100 MM)	VWR	76510-208
dTTP (100 MM)	VWR	76510-224
Lonza NuSieve 3:1 Agarose	Thermal Fisher	BMA50090
T4 DNA ligase	NEB	M0202L
Biotin-11-dCTP	Jen Bioscience	NU-809-BIOX

Biotin-14-dATP	Jen Bioscience	NU-835-BIO14
Qubit dsDNA HS Assay Kit	Life Technologies Corporation	Q32851
Atto 633 NHS ester	Sigma	01464
Phase Lock Gel™, QuantaBio - 2302830, Phase Lock Gel Heavy	VMR	10847-802
NEBNext Ultra II DNA Library Prep Kit for Illumina	NEB	E7645S
Ampure Xp 5ml Kit	Thermal Fisher	NC9959336
Hifi Hotstart Ready Mix	Thermal Fisher	501965217
Dynabeads MyOne Streptavidin C1	Life Technologies Corporation	65001
cOmplete™, EDTA-free Protease Inhibitor Cocktail	Sigma	11873580001
N,N-Dimethylformamide	Sigma	227056
Potassium acetate solution	Sigma	95843
DSG (disuccinimidyl glutarate)	Thermal Fisher	PI20593
T4 Polynucleotide Kinase - 500 units	NEB	M0201S
DNA Polymerase I, Large (Klenow) Fragment - 1000 units	NEB	M0210L
End-it DNA End Repair Kit	Thermal Fisher	NC0105678
Proteinase K recomb. 100 mg	Sigma	3115879001
Nuclease MicroCoccal (s7)	Thermal Fisher	NC9391488
EGS (ethylene glycol bis(succinimidyl succinate))	Thermal Fisher	PI21565
Atto 565 NHS ester	Sigma	72464
HCR™ RNA-FISH Custom Probe Set: eve	Molecular Instruments	
HCR™ RNA-FISH Custom Probe Set: ter94	Molecular Instruments	
HCR™ RNA-FISH Custom Probe Set: CG12134	Molecular Instruments	
HCR™ RNA-FISH Custom Probe Set: eIF3j	Molecular Instruments	

HCR™ Amplifier B1, 488	Molecular Instruments	
HCR™ Amplifier B2, 564	Molecular Instruments	
HCR™ Amplifier B3, 647	Molecular Instruments	
HCR™ Buffers	Molecular Instruments	
NEBNext Multiplex Oligos for Illumina	NEB	E7335S
Software and algorithms		
Fiji (ImageJ)	Schindelin et al., 2012	fiji.sc
NIS element	Nikon	microscope.healthcare.nikon.com/products/software/nis-elements
Graphpad Prism 8	GraphPad Software	graphpad.com
HiGlass	Kerpedjiev et al., 2018	higlass.io
bwa	Li H. and Durbin R., 2009	bio-bwa.sourceforge.net/
samtools	Github/open source	samtools.github.io
pairsamtools	Github/open source	github.com/mirnylab/pairsamtools
pairix	Github/open source	github.com/4dn-dcic/pairix
cooler	Abdennur, N., and Mirny, L., 2019	github.com/mirnylab/cooler
Miniconda	Anaconda	docs.conda.io/en/latest/miniconda.html
Snakemake	Github/open source	snakemake.github.io

Creation of *nhomie* deletion flies

To modify *nhomie* at the *eve* locus, we used recombinase-mediated cassette exchange

(Bateman et al., 2006). First, we inserted two nearby attP sites, using CRISPR. The donor

plasmid for this was constructed as follows. First, a mini-*white* (*mw*) gene with Glass binding

sites (Fujioka et al., 1999) was inserted into BlueScript. From the standard *mw* gene, the Wari

insulator (Chetverina et al., 2008) was deleted. Then, two 102bp attP sequences (Venken et al.,

2011) were inserted, one just 5' of the Glass binding sites and the other at the 3' end of the

modified *mw*, creating the plasmid P-attPx2-*mw*. 5' and 3' homologous arms were added to both

ends. Two gRNA sequences were cloned into plasmid pCFD4 ((Port et al., 2014), Addgene).

The donor and gRNA plasmids were injected into a Cas9 line (y[1] M{vas-Cas9.S}ZH-2A

w[1118]), Bloomington Drosophila Stock Center). This chromosomal modification resulted in

one attP site being inserted in the intron of *CG12134*, and the other being inserted between the

eve 7-stripe enhancer and the 3+7 stripe enhancer regions. This also deleted 2.2kb of

endogenous sequence, including *nhomie* and the *eve* 7-stripe enhancer.

After identifying a successful insertion (NattPmw), *mw* was replaced by each of the

following using RMCE: 1) the previously deleted 2.2kb, restoring *nhomie* and the *eve* 7-stripe

enhancer, to create "wild-type *nhomie*" (*nhomie forward*), 2) the same 2.2kb sequence, but with

600bp of phage λ DNA in place of 600bp *nhomie* (λ DNA), 3) the same 2.2kb sequence, but

with 600bp *nhomie* inverted (*nhomie reverse*), and 4) the same 2.2kb sequence, but with 600bp

nhomie replaced by a copy of ~600bp *homie* in its native orientation in the chromosome (*homie*

forward). Each of these changes was confirmed by sequencing of genomic DNA from the

transgenic fly lines.

Analysis of embryonic cuticle patterns and *in situ* hybridization

To identify defects in developing embryos, embryos were collected for 2.5 hr., and

allowed to develop for an additional 20-21 hr. at 25°C. Embryos were dechorionated and

mounted in a 1:1 mixture of Hoyer's medium and lactic acid. Mounted embryos were left at

room temperature until they cleared (12-14 days), and the patterns of ventral abdominal denticles

were examined and tallied as follows. Loss of at least one-fifth of a denticle band (in A1-A8)

was counted as "missing". Fused denticle bands, which rarely occurred, was also counted as a

"missing" band. Minor defects such as in individual denticle rows were not counted.

Digoxigenin (DIG) *in situ* hybridization was performed using DIG-labeled anti-sense

RNA against *CG12134*, *eIFj3*, and *TER94*. RNA expression was visualized using alkaline

phosphatase-conjugated anti-DIG antibody (Roche), using CBIP and NBT as substrates (Roche).

Each set of experiments was carried out in parallel to minimize experimental variation.

Representative expression patterns are shown in each figure.

HCR-FISH

The sequences of target genes were obtained from Flybase (flybase.org)(Gramates et al., 2022).

To design probes, the target gene sequences were submitted to the Molecular Instruments probe

design platform (www.molecularinstruments.com/hcr-rnafish) (Choi et al., 2016), with

parameters set to a 35 probe set size for *Drosophila melanogaster*. A similar method was

designed based on published smFISH methods (Little and Gregor, 2018; Trcek et al., 2017). 100-

200 flies were placed in a cage with an apple juice plate at the bottom of the cage. For early

stages, the embryos were collected for 7 hours, while for later-stage embryos, collection were

overnight. Embryos from each plate were washed into collection mesh and dechorionated in

bleach for 2min, then fixed in 5mL of 4% paraformaldehyde in 1X PBS and 5mL of heptane for

15min with horizontal shaking. The paraformaldehyde was then removed and replaced with

5mL methanol. The embryos were then devitellinized by vortexing for 30s, and washed in 1mL

of methanol twice. Methanol was then removed and replaced by PTw (1X PBS with 0.1%

Tween-20) through serial dilution as 7:3, 1:1, and 3:7 methanol:PTw. The embryos were washed

twice in 1mL of PTw and pre-hybridize in 200μL of probe hybridization buffer for 30 min at 37°

C. 0.4pmol of each probe sets were added to the embryos in the probe hybridization buffer and

the embryos were incubated at 37ºC for 12-14h. The embryos were then washed 3X with probe

wash buffer at 37°C for 30min and 2X with 5X SSCT(5X SSC+0.1% tween) at room

temperature for 5min. Then the embryos were pre-amplify with 300μL amplification buffer for

10 min at 25°C. Meanwhile, 6 pmol of hairpin h1 and h2 were snap cooled separately (95ºC for

1:30, cool to RT with 0.1ºC drop per second), and then mixed in 100μL of amplification buffer

at room temperature. After that, the pre-amplification solution was removed from the embryos,

and 100μL of hairpin h1/h2 mix were added to the embryos. Next, the embryos were incubated

for 12-14h at room temperature in dark. To remove the excess hairpins, the embryos were then

washed in SSCT as follows: 2X for 5min, 2X for 30min and 5X for 5min. Then, the embryos

were washed with 1mL PTw for 2min and stained with DAPI/Hoechst at 1μg/mL for 15min at

room temperature in the dark. The embryos were then washed with PTw 3X for 5min. Finally,

the embryos were mounted on microscope slides with Vectashield and a #1.5 coverslip for

imaging.

Imaging, image analysis and statistics

Embryos from hcrFISH were imaged by using a Nikon A1 confocal microscope system,

Plan Apo 20X/0.75 DIC objective. Z-stack images were taken at interval of 2μm, 4X average,

1024X1024 resolution, and the appropriate laser power and gain were set for 405, 488, 561, and

640 channels to avoid overexposure. Images were processed by ImageJ and the maximum

projection was applied to each of the stack images. To determine the presence of stripes of early

embryos, multi-channel images were first split into single channels and the stripe signal was

highlighted and detected by the MaxEntropy thresholding method. GraphPad Prism was used for

data visualization and statistical analysis. Two-way ANOVA with Tukey's multiple

comparisons test for each pair of groups was used to determine the statistical significance for the

percentage of embryos carrying stripes in *eIF3j* and *TER94* channels in each group.

MicroC library construction for the *nhomie* replacements

Embryos were collected on yeasted apple juice plates in population cages for 4 hours,

incubated for 12 hours at 25°C, then subjected to fixation as follows. Embryos were

dechorionated for 2min in 3% sodium hypochlorite, rinsed with deionized water, and transferred

to glass vials containing 5mL PBST (0.1% Triton-X100 in PBS), 7.5mL n-heptane, and 1.5mL

fresh 16% formaldehyde. Crosslinking was carried out at room temperature for exactly 15min

on an orbital shaker at 250rpm, followed by addition of 3.7 mL 2M Tris-HCl pH7.5 and shaking

for 5min to quench the reaction. Embryos were washed twice with 15mL PBST and subjected to

secondary crosslinking. Secondary crosslinking was done in 10mL of freshly prepared 3mM

final DSG and ESG in PBST for 45min at room temperature with passive mixing. The reaction

was quenched by addition of 3.7mL of 2M Tris-HCl pH7.5 for 5min, washed twice with PBST,

snap-frozen, and stored at -80°C until library construction.

Micro-C libraries were prepared as previously described (Batut *et al.*, 2022) with the following

modifications: 50μL of 12-16h embryos were used for each biological replicate. 60U of MNase

was used for each reaction to digest chromatin to a mononucleosome:dinucleosome ratio of 4.

Libraries were barcoded, pooled and subjected to paired-end sequencing on an Illumina Novaseq

S1 100nt Flowcell (read length 50 bases per mate, 6-base index read).

Micro-C data processing

MicroC data for *D. melanogaster* were aligned to custom genomes edited from the

Berkeley Drosophila Genome Project (BDGP) Release 6 reference assembly (dos Santos et al.,

2015) with BWA-MEM (Li and Durbin, 2009) using parameters **-S -P -5 -M**. The resultant

BAM files were parsed, sorted, de-duplicated, filtered, and split with Pairtools

(https://github.com/mirnylab/pairtools). We removed pairs where only half of the pair could be

mapped, or where the MAPQ score was less than three. The resultant files were indexed with

Pairix (https://github.com/4dn-dcic/pairix). The files from replicates were merged with Pairtools

before generating 100bp contact matrices using Cooler (Abdennur and Mirny, 2020). Finally,

balancing and Mcool file generation was performed with Cooler's Zoomify tool.

Virtual 4C profiles were extracted from individual replicates using FAN-C (Kruse et al.,

2020) at 400bp resolution. The values were summed across replicates and smoothed across three

bins (1.2kb). The *homie* Viewpoint was set to the 549nt *homie* sequence that was defined in

previous studies.

     Figure Legends

Figure 1. Diagram of the possible loop topologies generated by head-to-head and head-to-
tail pairing. A) Head-to-tail boundary (arrows) pairing generates a series of stem-loops linked
together by unanchored loop. In this case, the main axis of the chromosome would correspond to
the unanchored loops connecting different stem-loops. B) If boundaries pair with both neighbors
(head-to-tail) the stem-loops would be linked to each other by the paired boundaries. In this case
the main axis of the chromosome will correspond to the paired boundaries. C) Head-to-head
boundary pairing generates a series of circle-loops linked together by an unanchored loop. The
unanchored loop will be the main axis of the chromosome. D) If boundaries pair with both
neighbors (head-to-head), the chromatin fiber will be organized into a series of circle-loops
connected to each other at their base and these paired boundaries will define the chromosomal
axis. In both B) and D), the pairing interactions between the blue and red boundaries need not be
in register with the pairing of the red boundary to the next-door green boundary. In this case, the
main axis of the chromosome may bend and twist, and this could impact the relative orientation
of the stem-loops/circle-loops. More complex structures would be generated by mixtures of
stem-loops and circle-loops.
662
Figure 2. Stem-loops and circle-loops. "*Once in a while you get shown the light in the*
strangest of places if you look at it right." * MicroC contact profile for *Drosophila* wild-type
(*yw*) stage 14 embryos. A) *eve* and neighboring TADs (TI, TJ, TK TL, TM and TN). The *eve*
TAD is a volcano with a plume that is anchored by *nhomie* (*nh*) and *homie* (*h*). The plume is
generated by crosslinking of sequences in neighboring TADs. At the bottom of the plume, TL
sequences are linked to sequences in TM close to *eve* including *ter94*. At the next level,
sequences in TK are linked to TM. In addition, sequences in LM are linked to sequences in TM
located beyond the *TER94* gene. At the next level, sequences in TJ are linked to sequences in
TM. Note that interactions between sequences in TL and TJ and sequences in TM close to the
eve TAD are somewhat less frequent than those farther away from the *eve* TAD. Sequences in
the neighboring TAD also interact with each other as indicated. For example, sequences in TK
and TJ interact with each other (J-K) and also with interact with sequences in TI (I-K and I-J).
B) The Bithorax *Abd-B* gene and the parasegment (PS) specific regulatory domains, <u>*iab-3, iab-4,*</u>
<u>*iab-5, iab-6, iab-7, iab-8*</u>. *iab-4* regulates the *abd-A* gene in PS9. *iab-5* regulates *Abd-B* in
PS10, *iab-6* regulates *Abd-B* in PS11, *iab-7* regulates *Abd-B* in PS12 and *iab-8* regulates *Abd-B*
in PS13. The domains are separated from each other by the boundary elements *Fab-4*, *Mcp*,
Fab-6, *Fab-7*, and *Fab-8* as indicated. The *AB-I* boundary is located upstream of the *Abd-B*
promoter. Each regulatory domain corresponds to a TAD. Though insulated from each other,
each TAD interacts with its immediate neighbor. For example, *iab-5* interacts with its immediate
neighbor *iab-4* and *iab-6* to give *4-5* and *5-6* respectively. It also interacts with the next-next
door neighbor *iab-7* (*5-7*) and even its next-next-next door neighbor *iab-8* (*5-8*). (x: Scarlet
Begonias GD: 1974).
685
Figure 3. TAD organization of the *nhomie forward* and *λDNA* replacements. A) MicroC
contact profile of stage 14 *nhomie forward* embryos. In this replacement *nhomie* is inserted in
the same orientation as the endogenous *nhomie*. N(replicates)=2. Resolution = 200bp. *L-M*:
Interactions between TADs TL and TM flanking the *eve* locus. Asterisk: sub-TAD linking the
eve promoter to the *eve* PRE and *homie*. Dark blue arrow: sub-TAD linking the *eve* promoter to
nhomie. Light blue arrowhead: *nhomie*. Red arrowhead: *homie*. Green arrowhead: sub-TAD
boundary formed by the *CG12134* promoter region. Dark blue arrowhead: sub-TAD boundary

formed by *eIF3j* promoter region. Diagram: Map of *eve* locus and surrounding genes. B)
Virtual 4C with viewpoint from *homie* (black arrow) towards *nhomie forward*. C) Diagram of
the *eve* stem-loop TAD. D) MicroC contact profile of stage 14 *λ DNA* embryos. In this
replacement, λ DNA is inserted in place *nhomie*. N(replicates)=2. Resolution = 200bp. Asterisks:
sub-TAD linking the *eve* promoter to the *eve* PRE and *homie*. Purple arrowhead: sub-TAD
linking *CG12134* promoter region to *eve* promoter. The *eIF3j* sub-TAD (demarked by the blue
and green arrowheads) is still present. E) Virtual 4C with viewpoint from *homie* reading toward
eIF3j. F) Diagram of the "unanchored" *eve* TAD. Arrows show novel interactions.
701
Figure 4. *nhomie* deletion (*lambda DNA* replacement) exposes *eIF3j* to the *eve* enhancers.
nh forward: *nhomie* replacement with *nhomie* in the forward (WT) orientation. *λ DNA: nhomie*
replacement with λ DNA. At the syncytial blastoderm stage, a series of stripe-specific enhancers
upstream (stripes 1, 2, 3, 7) and downstream (stripes 1, 4, 5, 6) of the *eve* gene drive *eve*
expression. During cellularization of the blastoderm and gastrulation, a single enhancer located
upstream of *eve* drives expression of all seven stripes. DAPI: DNA stained with DAPI (blue).
eIF3j: Embryo hybridized with probe complementary to *eIF3j* mRNA. *eve*: Embryo hybridized
with probe complementary to *eve* mRNA. *TER94*: Embryo hybridized with probe
complementary to *TER94*. Yellow carets: *eve*-enhancer-driven *eIF3j* stripes. Control non-
specific probes for each channel indicates autofluorescence background for each channel in the
top panel.
713
Figure 5. Manipulating the *nhomie* boundary impacts the regulatory landscape. N=# of
independent replicate, n= # of embryos. Two-way ANOVA with Tukey's multiple comparisons
test for each pair of groups was used to determine the statistical significance. *$p \leq 0.05$,
**$p \leq 0.01$, ***$p \leq 0.001$ and ****$p \leq 0.0001$. A) Quantitation of numbers of embryos showing
stripe patterns in HCR-FISH for *eIF3j* and *TER94,* as shown in Fig. 4 and Fig 7. N=3. n=45 for
each group. B) Quantitation of the number of missing ventral denticle bands in larvae from a
cross of *BSC/CyO,hb-lacZ* deficiency females to males of the indicated genotypes (N=6): wild-
type control (*yw*) n = 767. The *nhomie forward* replacement n = 1099. The *λDNA* replacement n
= 1175. The *nhomie reverse* replacement: n = 1083. The *homie forward* replacement: n =
1137.
724
725
Figure 6. TAD organization of the *nhomie reverse* and *homie forward* replacements. A)
MicroC contact profile of stage 14 *nhomie reverse* embryos. In this replacement *nhomie* is
inserted in the reverse orientation compared to WT *nhomie*. N(replicates)=2. Resolution =
200bp. B) Virtual 4C with viewpoint from *homie* (black arrow) towards *nhomie reverse*. C)
MicroC contact profile of stage 14 *homie forward* embryos. In this replacement *homie* is
inserted in the forward orientation (the same as the endogenous *homie*). N(replicates)=2.
Resolution = 200bp. D) Virtual 4C with viewpoint from *homie* (black arrow) towards *homie*
forward. A) and C) Note that interactions between the TADs flanking the *eve* locus (purple
double arrow) are suppressed compared to *nhomie forward* (see Fig. 3), while interactions
between *eve* and TL, TM are enhanced (L-ev and ev-M). Asterisks: sub-TAD linking the *eve*
promoter to the *eve* PRE and *homie*. Dark blue arrow: sub-TAD linking the *eve* promoter to
nhomie reverse. Light blue bar: *nhomie reverse*. Red bar: *homie*. Green arrowhead: sub-TAD
boundary formed by the *CG12134* promoter region. Dark blue arrowhead: sub-TAD boundary
formed by the *eIF3j* promoter region. Diagram: Map of *eve* locus and surrounding genes.
740

741
Figure 7. *eve* enhancers activate neighboring genes when the *eve* TAD is a circle-loop.
hcrFISH hybridization to mRNA expressed by *eIF3j*, *eve* and *TER94* at the blastoderm stage
(embryonic stage 5). *nh forward*: *nhomie* replacement with *nhomie* in forward (normal)
orientation (same as in Fig. 3). *nh reverse: nhomie* replacement with *nhomie* in reverse
orientation. *h forward: nhomie* replacement with *homie* in forward orientation. Yellow
arrowhead: position of stripes. DAPI (blue): DNA stained with DAPI. *eIF3j* (green): Embryo
hybridized with probe complementary to *eIF3j* mRNA. *eve* (orange): Embryo hybridized with
probe complementary to *eve* mRNA. *TER94* (red): Embryo hybridized with probe
complementary to *TER94*. Control non-specific probes for each channel indicates
autofluorescence background for each channel in the top panel.
758 <u>**Figures Supplemental**</u>
759
760 **Fig. Supplemental 1. MicroC contact profiles for *nhomie forward*, *λDNA*, *nhomie reverse*,**
761 **and *homie forward* in larger scale.** N(replicates)=2. Resolution=200. A) MicroC contact maps
762 for the *nhomie forward* replacement. B) MicroC contact maps for the *λDNA* replacement. C)
763 MicroC profile of the *nhomie reverse* replacement. D) MicroC profile of the *homie forward*
764 replacement.
765
Fig. Supplemental 2. Expression of *CG12134* in WT (*yw*) and the four *nhomie*
replacements.
Digoxigenin *in situ* hybridization was used to detect expression of *CG12134* during development
in the indicated genetic backgrounds. Approximate developmental stages of the embryos in each
genetic background are shown on the right. As controls, embryos of similar stages were
hybridized with an *eve* probe (top to bottom: stages 5, 7, 10, 11, and 13).
772
Fig. Supplemental 3. Expression of *eIF3j* (*Adam*) and *TER94* in WT (*yw*) and the four
***nhomie* replacements.** Digoxigenin *in situ* hybridization was used to detect expression of A)
eIF3j and B) *TER94* during development in the indicated genetic backgrounds. Approximate
developmental stages of the embryos in each genetic background are shown on right. As
controls embryos of a similar stages were hybridized with an *eve* probe. Note that unlike the
HCR-FISH results shown in Fig. 4, we can detect a low level of *eIF4j* stripe expression in WT
(*yw*) and *nohomie forward* St 5 embryos. The difference is likely due to the fact that the signal
amplification in the digoxigenin *in situ* hybridization procedure is not linear.
781
782
Fig. Supplemental 4. MicroC patterns of DNA segments on left and right arm of
chromosome 2. A) MicroC contact profiles on left arm of chromosome 2 around 3,000,000 bp.
Black arrow indicates that interactions between neighboring TADs are suppressed as might be
expected for a stem-loop configuration. The red arrow in immediately above the black arrow
points to contacts between next-next-(next) door neighbors that are enhanced. However, the
plume that is formed is one sided as indicated by the green arrow. B) MicroC contact profiles

for TADs on the right arm of chromosome 2. The TADs indicated by the double headed blue
arrow at the bottom have a complex pattern of neighborly interactions. The asterisk and purple
arrow indicate a potential volcano plume; however, unlike the *eve* volcano triangle and plume,
this plume appears to be generated by crosslinking between sequences to either side of the
collection of TADs indicated by the doubleheader blue arrow. The TAD-to-TAD interaction
patterns are further complicated by a band of enhanced crosslinking (red arrow).
795
796

References

Abdennur, N., and Mirny, L.A. (2020). Cooler: scalable storage for Hi-C data and other genomically labeled arrays. Bioinformatics *36*, 311-316. 10.1093/bioinformatics/btz540.

Americo, J., Whiteley, M., Brown, J.L., Fujioka, M., Jaynes, J.B., and Kassis, J.A. (2002). A complex array of DNA-binding proteins required for pairing-sensitive silencing by a polycomb group response element from the Drosophila engrailed gene. Genetics *160*, 1561-1571.

Arzate-Mejía, R.G., Josué Cerecedo-Castillo, A., Guerrero, G., Furlan-Magaril, M., and Recillas-Targa, F. (2020). In situ dissection of domain boundaries affect genome topology and gene transcription in Drosophila. Nat Commun *11*, 894. 10.1038/s41467-020-14651-z.

Bateman, J.R., Lee, A.M., and Wu, C.T. (2006). Site-specific transformation of Drosophila via phiC31 integrase-mediated cassette exchange. Genetics *173*, 769-777.

Batut, P.J., Bing, X.Y., Sisco, Z., Raimundo, J., Levo, M., and Levine, M.S. (2022). Genome organization controls transcriptional dynamics during development. Science *375*, 566-570. 10.1126/science.abi7178.

Bell, A.C., West, A.G., and Felsenfeld, G. (2001). Insulators and boundaries: versatile regulatory elements in the eukaryotic genome. Science *291*, 447-450. 10.1126/science.291.5503.447.

Bing, X., Ke, W., Fujioka, M., Kurbidaeva, A., Levitt, S., Levine, M., Schedl, P., and Jaynes, J.B. (2023). Chromosome Structure I: Loop extrusion or boundary:boundary pairing?

Cai, H.N., and Shen, P. (2001). Effects of cis arrangement of chromatin insulators on enhancer-blocking activity. Science *291*, 493-495.

Cavalheiro, G.R., Pollex, T., and Furlong, E.E. (2021). To loop or not to loop: what is the role of TADs in enhancer function and gene regulation? Curr. Opin. Genet. Dev. *67*, 119-129. 10.1016/j.gde.2020.12.015.

Chen, H., Levo, M., Barinov, L., Fujioka, M., Jaynes, J.B., and Gregor, T. (2018). Dynamic interplay between enhancer–promoter topology and gene activity. Nat. Genet. *50*, 1296.

Chetverina, D., Aoki, T., Erokhin, M., Georgiev, P., and Schedl, P. (2014). Making connections: insulators organize eukaryotic chromosomes into independent cis-regulatory networks. Bioessays *36*, 163-172.

Chetverina, D., Fujioka, M., Erokhin, M., Georgiev, P., Jaynes, J.B., and Schedl, P. (2017). Boundaries of loop domains (insulators): Determinants of chromosome form and function in multicellular eukaryotes. Bioessays *39*, 1600233.

Chetverina, D., Savitskaya, E., Maksimenko, O., Melnikova, L., Zaytseva, O., Parshikov, A., Galkin, A.V., and Georgiev, P. (2008). Red flag on the white reporter: a versatile insulator abuts the white gene in Drosophila and is omnipresent in mini-white constructs. Nucleic Acids Res. *36*, 929-937.

Choi, H.M., Calvert, C.R., Husain, N., Huss, D., Barsi, J.C., Deverman, B.E., Hunter, R.C., Kato, M., Lee, S.M., Abelin, A.C., et al. (2016). Mapping a multiplexed zoo of mRNA expression. Development *143*, 3632-3637. 10.1242/dev.140137.

Davidson, I.F., and Peters, J.M. (2021). Genome folding through loop extrusion by SMC complexes. Nat Rev Mol Cell Biol *22*, 445-464. 10.1038/s41580-021-00349-7.

dos Santos, G., Schroeder, A.J., Goodman, J.L., Strelets, V.B., Crosby, M.A., Thurmond, J., Emmert, D.B., Gelbart, W.M., and FlyBase, C. (2015). FlyBase: introduction of the Drosophila melanogaster Release 6 reference genome assembly and large-scale migration of genome annotations. Nucleic Acids Res. *43*, D690-697.

Erokhin, M., Gorbenko, F., Lomaev, D., Mazina, M.Y., Mikhailova, A., Garaev, A.K., Parshikov, A., Vorobyeva, N.E., Georgiev, P., Schedl, P., and Chetverina, D. (2021). Boundaries

potentiate polycomb response element-mediated silencing. BMC Biol. *19*, 113. 10.1186/s12915-
021-01047-8.
Fujioka, M., Emi-Sarker, Y., Yusibova, G.L., Goto, T., and Jaynes, J.B. (1999). Analysis of an
even-skipped rescue transgene reveals both composite and discrete neuronal and early
blastoderm enhancers, and multi-stripe positioning by gap gene repressor gradients.
Development *126*, 2527-2538.
Fujioka, M., Mistry, H., Schedl, P., and Jaynes, J.B. (2016). Determinants of Chromosome
Architecture: Insulator Pairing in *cis* and in *trans*. PLoS Genet. *12*, e1005889.
Fujioka, M., Yusibova, G.L., Zhou, J., and Jaynes, J.B. (2008). The DNA-binding Polycomb-
group protein Pleiohomeotic maintains both active and repressed transcriptional states through a
single site. Development *135*, 4131-4139.
Goel, V.Y., Huseyin, M.K., and Hansen, A.S. (2023). Region Capture Micro-C reveals
coalescence of enhancers and promoters into nested microcompartments. Nat. Genet. *55*, 1048-
1056. 10.1038/s41588-023-01391-1.
Gohl, D., Aoki, T., Blanton, J., Shanower, G., Kappes, G., and Schedl, P. (2011). Mechanism of
chromosomal boundary action: roadblock, sink, or loop? Genetics *187*, 731-748.
Gramates, L.S., Agapite, J., Attrill, H., Calvi, B.R., Crosby, M.A., Dos Santos, G., Goodman,
J.L., Goutte-Gattat, D., Jenkins, V.K., Kaufman, T., et al. (2022). FlyBase: a guided tour of
highlighted features. Genetics *220*. 10.1093/genetics/iyac035.
Heger, P., George, R., and Wiehe, T. (2013). Successive gain of insulator proteins in arthropod
evolution. Evolution *67*, 2945-2956.
Heger, P., and Wiehe, T. (2014). New tools in the box: an evolutionary synopsis of chromatin
insulators. Trends Genet. *30*, 161-171.
Higashi, T.L., and Uhlmann, F. (2022). SMC complexes: Lifting the lid on loop extrusion. Curr.
Opin. Cell Biol. *74*, 13-22. 10.1016/j.ceb.2021.12.003.
Hsieh, T.S., Cattoglio, C., Slobodyanyuk, E., Hansen, A.S., Rando, O.J., Tjian, R., and Darzacq,
X. (2020). Resolving the 3D Landscape of Transcription-Linked Mammalian Chromatin
Folding. Mol. Cell *78*, 539-553.e538. 10.1016/j.molcel.2020.03.002.
Ibragimov, A., Bing, X.Y., Shidlovskii, Y.V., Levine, M., Georgiev, P., and Schedl, P. (2023).
lncRNA read-through regulates the BX-C insulator Fub-1. Elife *12*. 10.7554/eLife.84711.
Jerković, I., Szabo, Q., Bantignies, F., and Cavalli, G. (2020). Higher-Order Chromosomal
Structures Mediate Genome Function. J. Mol. Biol. *432*, 676-681. 10.1016/j.jmb.2019.10.014.
Kassis, J.A., VanSickle, E.P., and Sensabaugh, S.M. (1991). A fragment of engrailed regulatory
DNA can mediate transvection of the white gene in Drosophila. Genetics *128*, 751-761.
Kellum, R., and Schedl, P. (1992). A group of scs elements function as domain boundaries in an
enhancer-blocking assay. Mol. Cell. Biol. *12*, 2424-2431.
Krietenstein, N., Abraham, S., Venev, S.V., Abdennur, N., Gibcus, J., Hsieh, T.S., Parsi, K.M.,
Yang, L., Maehr, R., Mirny, L.A., et al. (2020). Ultrastructural Details of Mammalian
Chromosome Architecture. Mol. Cell *78*, 554-565.e557. 10.1016/j.molcel.2020.03.003.
Kruse, K., Hug, C.B., and Vaquerizas, J.M. (2020). FAN-C: a feature-rich framework for the
analysis and visualisation of chromosome conformation capture data. Genome Biol *21*, 303.
10.1186/s13059-020-02215-9.
Kyrchanova, O., Chetverina, D., Maksimenko, O., Kullyev, A., and Georgiev, P. (2008a).
Orientation-dependent interaction between Drosophila insulators is a property of this class of
regulatory elements. Nucleic Acids Res. *36*, 7019-7028.
Kyrchanova, O., Ivlieva, T., Toshchakov, S., Parshikov, A., Maksimenko, O., and Georgiev, P.
(2011). Selective interactions of boundaries with upstream region of Abd-B promoter in

Drosophila bithorax complex and role of dCTCF in this process. Nucleic Acids Res. *39*, 3042-3052.

Kyrchanova, O., Kurbidaeva, A., Sabirov, M., Postika, N., Wolle, D., Aoki, T., Maksimenko, O., Mogila, V., Schedl, P., and Georgiev, P. (2018). The bithorax complex iab-7 Polycomb response element has a novel role in the functioning of the Fab-7 chromatin boundary. PLoS Genet. *14*, e1007442.

Kyrchanova, O., Toshchakov, S., Podstreshnaya, Y., Parshikov, A., and Georgiev, P. (2008b). Functional interaction between the Fab-7 and Fab-8 boundaries and the upstream promoter region in the Drosophila Abd-B gene. Mol. Cell. Biol. *28*, 4188-4195.

Levo, M., Raimundo, J., Bing, X.Y., Sisco, Z., Batut, P.J., Ryabichko, S., Gregor, T., and Levine, M.S. (2022). Transcriptional coupling of distant regulatory genes in living embryos. Nature *605*, 754-760. 10.1038/s41586-022-04680-7.

Li, H., and Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler transform. Bioinformatics *25*, 1754-1760. 10.1093/bioinformatics/btp324.

Li, H.-B., Muller, M., Bahechar, I.A., Kyrchanova, O., Ohno, K., Georgiev, P., and Pirrotta, V. (2011). Insulators, not Polycomb response elements, are required for long-range interactions between Polycomb targets in Drosophila melanogaster. Mol. Cell. Biol. *31*, 616-625.

Little, S.C., and Gregor, T. (2018). Single mRNA Molecule Detection in Drosophila. Methods Mol Biol *1649*, 127-142. 10.1007/978-1-4939-7213-5_8.

Matthews, N.E., and White, R. (2019). Chromatin Architecture in the Fly: Living without CTCF/Cohesin Loop Extrusion?

Mohana, G., Dorier, J., Li, X., Mouginot, M., Smith, R.C., Malek, H., Leleu, M., Rodriguez, D., Khadka, J., Rosa, P., et al. (2023). Chromosome-level organization of the regulatory genome in the Drosophila nervous system. Cell *186*, 3826-3844.e3826. 10.1016/j.cell.2023.07.008.

Muller, M., Hagstrom, K., Gyurkovics, H., Pirrotta, V., and Schedl, P. (1999). The mcp element from the Drosophila melanogaster bithorax complex mediates long-distance regulatory interactions. Genetics *153*, 1333-1356.

Muravyova, E., Golovnin, A., Gracheva, E., Parshikov, A., Belenkaya, T., Pirrotta, V., and Georgiev, P. (2001). Loss of insulator activity by paired Su(Hw) chromatin insulators. Science *291*, 495-498.

Perea-Resa, C., Wattendorf, L., Marzouk, S., and Blower, M.D. (2021). Cohesin: behind dynamic genome topology and gene expression reprogramming. Trends Cell Biol. *31*, 760-773. 10.1016/j.tcb.2021.03.005.

Port, F., Chen, H.-M., Lee, T., and Bullock, S.L. (2014). Optimized CRISPR/Cas tools for efficient germline and somatic genome engineering in Drosophila. Proceedings of the National Academy of Sciences *111*, E2967-E2976.

Rowley, M.J., and Corces, V.G. (2018). Organizational principles of 3D genome architecture. Nature Reviews Genetics *19*, 789-800.

Schoborg, T.A., and Labrador, M. (2010). The phylogenetic distribution of non-CTCF insulator proteins is limited to insects and reveals that BEAF-32 is Drosophila lineage specific. J. Mol. Evol. *70*, 74-84. 10.1007/s00239-009-9310-x.

Sigrist, C.J., and Pirrotta, V. (1997). Chromatin insulator elements block the silencing of a target gene by the Drosophila polycomb response element (PRE) but allow trans interactions between PREs on different chromosomes. Genetics *147*, 209-221.

Stadler, M.R., Haines, J.E., and Eisen, M.B. (2017). Convergence of topological domain boundaries, insulators, and polytene interbands revealed by high-resolution mapping of chromatin contacts in the early Drosophila melanogaster embryo. eLife *6*, 17.

Trcek, T., Lionnet, T., Shroff, H., and Lehmann, R. (2017). mRNA quantification using single-
molecule FISH in Drosophila embryos. Nat Protoc *12*, 1326-1348. 10.1038/nprot.2017.030.
Vazquez, J., Farkas, G., Gaszner, M., Udvardy, A., Muller, M., Hagstrom, K., Gyurkovics, H.,
Sipos, L., Gausz, J., Galloni, M., and et al. (1993). Genetic and molecular analysis of chromatin
domains. Cold Spring Harb Symp Quant Biol *58*, 45-54. 10.1101/sqb.1993.058.01.008.
Vazquez, J., Muller, M., Pirrotta, V., and Sedat, J.W. (2006). The Mcp element mediates stable
long-range chromosome-chromosome interactions in Drosophila. Molecular Biology of the Cell
17, 2158-2165.
Venken, K.J.T., Schulze, K.L., Haelterman, N.A., Pan, H., He, Y., Evans-Holm, M., Carlson,
J.W., Levis, R.W., Spradling, A.C., Hoskins, R.A., and Bellen, H.J. (2011). MiMIC: a highly
versatile transposon insertion resource for engineering Drosophila melanogaster genes. Nat.
Methods *8*, 737-743.
Yokoshi, M., Segawa, K., and Fukaya, T. (2020). Visualizing the Role of Boundary Elements in
Enhancer-Promoter Communication. Mol. Cell *78*, 224-235.e225. 10.1016/j.molcel.2020.02.007.

982 **FIGURES**

983
984
985

986
987 **Figure 1. Diagram of the possible loop topologies generated by head-to-head and head-to-**
988 **tail pairing.**

989
990
991

A

B

992
993 **Figure 2. Stem-loops and circle-loops.**
994

Figure 3. TAD organization of the *nhomie forward* and *λDNA* replacements.

Figure 4. *nhomie* deletion (*lambda DNA* replacement) exposes *eIF3j* to the *eve* enhancers.

1019
1020 **Figure 5. Manipulating the *nhomie* boundary impacts the regulatory landscape.**

Figure 6. TAD organization of the *nhomie reverse* and *homie forward* replacements.

Figure 7. *eve* enhancers activate neighboring genes when the *eve* TAD is a circle-loop.