ABSTRACT

Patients present a wide range of clinical severities in response SARS-CoV-2 infection, but the underlying molecular and cellular reasons why clinical outcomes vary so greatly within the population remains unknown. Here, we report that negative clinical outcomes in severely ill patients were associated with divergent RNA transcriptome profiles in peripheral immune cells compared with mild cases during the first weeks after disease onset. Protein-protein interaction analysis indicated that early-responding cytotoxic NK cells were associated with an effective clearance of the virus and a less severe outcome. This innate immune response was associated with the activation of select cytokine-cytokine receptor pathways and robust Th1/Th2 cell differentiation profiles. In contrast, severely ill patients exhibited a dysregulation between innate and adaptive responses affiliated with divergent Th1/Th2 profiles and negative outcomes. This knowledge forms the basis of clinical triage that may be used to preemptively detect high-risk patients before life-threatening outcomes ensue.

LONGITUDINAL TRANSCRIPTIONAL CHANGES REVEAL GENES FROM THE NATURAL KILLER CELL-MEDIATED CYTOTOXICITY PATHWAY AS CRITICAL PLAYERS UNDERLYING COVID-19 PROGRESSION

Matias A. Medina[1], Francisco Fuentes-Villalobos[2], Claudio Quevedo[3], Felipe Aguilera[3], Raul Riquelme[1,4], Maria Luisa Rioseco[1,4], Sebastian Barria[4], Yazmin Pinos[5], Mario Calvo[6], Ian Burbulis[1], Camila Kossack[1], Raymond A Alvarez[7], COVID 19 South Chile Group, Jose Luis Garrido[1*], Maria Ines Barria[1*]

[1]Facultad de Medicina y Ciencia, Universidad San Sebastián, Puerto Montt, Chile

[2]Departamento de Microbiología, Facultad de Ciencias Biológicas, Universidad de Concepción, Concepción, Chile

[3]Departamento de Bioquímica y Biología Molecular, Facultad de Ciencias Biológicas, Universidad de Concepción, Concepción, Chile.

[4]Hospital Dr. Eduardo Schütz Schroeder, Puerto Montt, Chile.

[5]Hospital Base San José, Osorno, Chile.

[6]Instituto de Medicina, Facultad de Medicina, Universidad Austral, Valdivia, Chile.

[7]Division of Infectious Diseases, Department of Medicine, Immunology Institute, Icahn School of Medicine at Mount Sinai, New York, NY 10029, USA.

*Corresponding authors: Jose Luis Garrido, jlgarri@gmail.com; Maria Ines Barria, maria.barriac@uss.cl

Highlights

- Mild COVID-19 patients displayed an early transcriptional commitment with NK cell function, whereas severe patients do so with neutrophil function.

- The identified co-expressed genes give insights into a coordinated transcriptional program of NK cell cytotoxic activity being associated with mild patients.

- Key checkpoints of NK cell cytotoxicity that were enriched in mild patients include: *KLRD1*, *CD247*, and *IFNG*.

- The early innate immune response related to NK cells connects with the Th1/Th2 adaptive immune responses, supporting their relevance in COVID-19 progression.

Introduction

The severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) promotes several dysfunctions in human immune responses while triggering a broad spectrum of clinical presentations that range from asymptomatic infection to a mild, moderate, or sometimes lethal severe symptomatology (Ge et al., 2020; The, 2020; Wu & McGoogan, 2020). Convalescent patients report prolonged COVID-19 symptoms beyond the time course for typical cold and flu events, which highlights the possibility of long-term tissue damage generated by this virus (Ladds et al., 2020; Nalbandian et al., 2021; Ryan et al., 2022; Subramanian et al., 2022). Similarly as other respiratory viruses, it is known that SARS-CoV-2 triggers an immune response involving the recruitment, activation, and differentiation of innate and adaptive immune cells (Newton, Cardani, & Braciale, 2016; Shen et al., 2023; Wauters et al., 2021). For mildly-ill patients, these coordinated immunological efforts resolve infection but for unknown reasons the virus evades these responses in severely-ill patients to produce life-threatening COVID-19 (Park, 2020; Rashid et al., 2022; Sun, Xie, Bu, Zhong, & Zeng, 2022; Thorne et al., 2022). The genetic background and physiological health of individual patients certainly plays a major role in the clinical presentation of COVID-19 but the exact mechanisms of how the virus evades innate and adaptive responses is not known, or why there is such great variability in severity of clinical presentation among patients. This information is critical for developing new diagnostics that detect patients who will eventually progress to severe COVID-19 before respiratory failure ensues, and furthermore, provide host and virus targets to engineer effective treatments (X. Li et al., 2021; Samadizadeh et al., 2021).

Biomarkers linked to COVID-19 severity hold promise for detecting patients that will eventually develop severe COVID-19 (Janssen et al., 2021; The, 2020). In this context, blood-derived cues were associated with severe COVID-19, including an imbalance in immune cell populations that included neutrophil abundance, lymphopenia, myeloid dysfunction, and T cell activation/exhaustion (Ahern et al., 2022; Chen & John Wherry, 2020; Mann et al., 2020; Wauters et al., 2021). The differential expression of select chemokines and their receptors (Khalil, Elemam, &

Maghazachi, 2021) with associated cytokine storm drove monocyte and megakaryocyte dysfunction in severely ill patients (Ren et al., 2021). Comprehensive knowledge of host immune responses against SARS-CoV-2 is still limited but these divergent cell profiles implicated cell-to-cell signaling events occurring between the innate and adaptive cell compartments as critical for the progression of severe COVID-19 (Daamen et al., 2021; Rabaan et al., 2022; C. Wang et al., 2022).

One approach to identifying changes in immune-responses affiliated with severe COVID-19 is to monitor autocrine, paracrine and endocrine signaling in individual patients over time. Temporal events associated with each type of signaling is obviously difficult to disentangle from measuring the activities of circulating peripheral cells alone because there are distinct events happening in localized microenvironments, e.g., the spleen and lymph nodes. A complementary tactic to access information about these events is to monitor gene expression for the synthesis of chemokines and cell-associated receptors as a proxy of biochemical events happening in distinct immune effector cells. Based on current knowledge, we hypothesized that critical events occurring at the earliest stages of infection necessary for effective viral clearance are either perturbed or disrupted so as to promote cytokine storm and other pathologies associated with severe outcomes. We predicted these pathological immune events may be observable by measuring changes gene expression reflecting activities in distinct effector cells during the first weeks of infection (Ahern et al., 2022; Bernardes et al., 2020; Notarbartolo et al., 2021; Xiong et al., 2020; Zheng et al., 2020). However, these types of experiments require careful design because the type and quantity of all immune responses are dynamic during infections and comparing poorly-matched PBMCs may confound identification of bonafide immune dysregulation evident between patients (Bernardes et al., 2020; Notarbartolo et al., 2021; Zheng et al., 2020).

Here, we designed a longitudinal comparison between mild and severe patients, choosing the appropriate samples according to the clinical progression and the unbiased gene expression profile to study how changes in gene expression in distinct immune effector cells changed during the earliest time points since peak of

symptoms and during progression of clinical disease. We repeatedly measured
whole-transcriptome profiles of peripheral blood mononuclear cells (PBMCs) from
the same cohort of mildly- and severely-ill patients to identify molecular pathways
that were enriched during the clinical trajectory of COVID-19 over time. Briefly, to
gain more insights into our findings and get more comprehensive knowledge of the
phenomenon, we used a pairwise comparison of gene expression, gene set
enrichment, and weight-correlated gene network analyses. By doing so, we
identified pathways of genes involved with the NK cell cytotoxicity enriched in mild
patients when compared to severe. Besides focusing on a particular molecular
pathway, we investigated the interactions to better comprehend the underlying
phenomena of a successful immune response, contributing to an integrated point of
view throughout the transcriptomic analyses of functional pathways to mitigate
potential biases attributed to focusing the study on a single pathway. In this regard,
we revealed that the NK signaling pathway was intricately related to other
transcriptional circuits, such as those governing Th1/Th2 cell differentiation and
cytokine-cytokine receptor signaling pathways. These interactions highlight the
importance of these pathways as bridges between the innate and adaptive immune
responses throughout the disease, implying that the innate NK signaling pathway
(cell cytotoxic activity) is beneficial, and possibly a critical activity required to
effectively eradicate coronavirus. We also observed that an adaptive immune
response including early cell-mediated immunity was significant in lowering disease
severity. The link between the primary innate NK cell activity and the transcriptional
priming of adaptive Th1 and Th2 cell responses appears to be more robust in mild
patients than in severe. This work provides clear guidance to develop better medical
practices and prevention tactics against SARS-CoV-2 and other related infectious
respiratory virus (Haitao et al., 2020; Ponti, Maccaferri, Ruini, Tomasi, & Ozben,
2020; L. Zhang & Guo, 2020).

Results

Clinical features and temporal gene expression patterns in SARS-CoV-2 infected patients

A total of 22 peripheral blood samples were obtained from eight COVID-19 patients. These samplings following a longitudinal schedule complemented with 2 samples from healthy donors. All patients were recruited after an average period of 5 days after symptoms onset (Figure 1A). Some samples were taken from patients at the Hospital of Osorno and the Hospital of Puerto Montt, which are cities located in the region Los Lagos. The remaining samples were collected from patients at the Hospital Base and Clínica Alemana in Valdivia, a city located in the Region Los Ríos. All infections occurred between November 2020 and May 2021 (Table 1).

A crucial issue with longitudinal studies is defining an appropriate sampling schedule that provides a reasonable comparison between patients during the time course of naturally occurred infections. To align the comparability and consistency of data measured between patients, we designed a protocol consisting of three donations per patient to monitor events occurring during both acute infection and the recovery phase. We collected peripheral blood samples on days 0, 7, and 28 (D0, D7, and D28) during the peaks of symptoms (Figure 1B).

Clinical features of COVID-19 patients with mild and severe symptoms were determined by medical personnel at the hospitals mentioned above and used to describe their disease trajectories over time using the WHO ordinal scale (Ahern et al., 2022) (Figure 1B). In contrast to mild patients, all four severe patients experienced symptoms such as fever, cough, headache, chills, diarrhea, myalgia, and dyspnea (Table 1). These patients received mechanical ventilation on sampling D0. In general, symptoms from severe and mild patients diminished gradually up to D28 after recruitment, with the exception of one mild and another severe patient who still experienced mild symptoms.

Mild and severely ill patients displayed different transcriptional programs at the beginning of disease onset. To determine the gene expression profiles of each

patient over the time of disease progression, we developed an RNA-seq approach

that takes advantage of the longitudinal sampling scheme. By using all expressed

genes, we performed a Principal Component Analysis to unbiasedly compare the

transcriptional signatures of each patient and two healthy donors. All peripheral

blood samples from severe patients on D0 (represented by red circles) were widely

dispersed over the left of Principal Component 1 (PC1) (Figure 1C), in contrast with

mild patients on D0 (represented by blue circles), suggesting that both groups of

patients displayed different transcriptional programs at the beginning of the disease.

The transcriptional profiles of severely ill patients changed during the recovery phase

to be consistent with that observed in mildly ill patients. Gradually, along with disease

progression and medical treatments, samples from severely ill patients shifted to the

right of PC1 (D7 and D28). Interestingly, on D28, when the majority of patients had

recovered, samples from severely ill patients were pooled together with those mild

patients who had already recovered. These observations indicated that despite the

transcriptional profiles being closer to that of mild patients at D28 as compared to

D0, severely ill patients still exhibited higher variability between themselves and

controls (Figure 1C). In contrast, every mild COVID-19 patient was separated from

the severe group on D0 and D7. Notably, only one mild COVID-19 patient (M1)

clustered with severe patients at D0. This donor showed a broader set of symptoms

over time when compared with the rest of mild patients (Figure 1B). This evidence

indicated a dissimilar transcriptional response in that specific patient at the onset of

disease. Over time, and after medical treatments, the transcriptional program of this

patient shifted to be consistent with the other mild patients (Figure 1C).

The timing of COVID-19 related gene expression differed between mild and severely

ill patients. We focused on the temporal variation of gene expression to identify

differentially expressed genes associated with COVID-19 progression. We found

statistically-significant differences in the timing of differential gene expression

between mild and severely ill individuals (Figure 1D and figure supplement 1). We

observed that severe patients displayed a transcriptional response completely

different from that of mild patients at the sequential time points of D0 and D7 (Figure

1D). Previous longitudinal studies identified molecular markers associated with

severe COVID-19 (Bernardes et al., 2020; Notarbartolo et al., 2021; Zheng et al., 2020), We detected these same molecular markers in our severely ill cohort (Figure 1D). The expression profiles of those genes varied significantly between mild and severe patients. For instance, the expression of MMP9 metalloproteinase (Zheng et al., 2020), S100A8/A9 alarmins (Bernardes et al., 2020), PADI2 (Notarbartolo et al., 2021) and IL18Rap peptidyl-arginine deiminases (Masood et al., 2021; Schultze & Aschenbrenner, 2021) were higher in severe patients on D0 than mild or control patients (Figure 1D). In addition, we found that *IFNG CCL2,* and CXCL10 cytokines, which were previously described as molecular markers in severely ill patients (Sette & Crotty, 2021; Vabret et al., 2020), displayed low expression in our severe COVID-19 patients in comparison with mildly ill patients during the progression of disease (Figure 1D).

The immune response of mild and severe patients is activated differentially during the acute phase of the COVID-19 infection

Most of the variations observed in the gene expression profiles of mild and severely ill patients occurred during the acute phase of disease. We performed pairwise gene expression comparisons between mild and severe patients and found differentially expressed genes (DEGs) mainly on D0 and D7. On D0, we found a total of 812 DEGs including 298 up-regulated and 514 down-regulated genes (Figure 1 - figure supplement 2). On D7, the number of DEGs was similar to D0, with 319 genes showing higher expression and 563 genes with lower expression. We found no differential gene expression between mild and severe patients at D28, supporting the interpretation that most imbalances in the gene expression profiles in the PBMCs of severely ill patients leveled out by D28 (Figure 1 - supplement 2).

Functional pathways involved with humoral immunity were enriched in severely ill patients during the acute phase compared to pathways involved with cell-mediated immunity in mild patients. The above results provided only a course overview of the transcriptional responses during COVID-19 progression. We expanded our focus to detect molecular mechanisms and pathways involved in the immune responses of

all patients by linking functional pathways to deferentially expressed genes (DEGs)

detected between severely ill, mildly ill and control patients. We used a 2-fold change

in gene expression level as a threshold to identify DEGs between mild and severe

patients on D0 and D7. We found upregulated expression for genes involved in

biological processes that included the T receptor signaling pathway, positive

regulation of T cell activation, and regulation of leukocyte cell adhesion in mild

COVID-19 patients at D0 (Figure 2A). We observed genes involved with

immunoglobulin production, antigen receptor-mediated signaling pathway, and

adaptive immune response were up regulated at D7 (Figure 2B). In contrast, we

observed enrichment of gene expression in pathways involved with complement

activation, humoral immune response mediated by circulated immunoglobulin, and

defense response to bacterium on D0 in severe COVID-19 patients. Furthermore,

DEGs in functional pathways mediating hydrogen peroxide metabolic processes,

cellular oxidant detoxification and reactive oxygen species were enriched on D7 of

infection in this group (Figures 2A and 2B). Biological pathways consistent with a

robust lymphocyte cellular immune response were enriched on D0 in mild patients.

This functional profile is distinctly different to the antibody / complement-dependent

humoral immune responses observed in severely ill individuals at the same time

point (Figure 2A). Nonetheless, differential expression of genes associated with

immunoglobulin function were mainly enriched in mild patients at D7 (Figure 2B),

while severe patients showed enrichment for genes related to inflammation, reactive

oxygen species (ROS), and responses against bacteria at that time of infection

(Figure 2B).

In addition to enriched biological processes, we also focused on KEGG pathway

enrichment among DEGs at D0 and D7 after COVID-19 infection. On D0, mild and

severe patients showed considerable differences in terms of the innate response,

with the Natural Killer-mediated cytotoxicity pathway enriched in mild-infected

patients, while neutrophil extracellular trap formation was enriched in severe ones

(Figure 2C). Furthermore, DEGs associated with the antigen processing and

presentation pathways are enriched in mild COVID-19 patients, in contrast with the

enrichment of complement and coagulation cascade pathways in severely ill patients

(Figure 2C). On the other hand, on D7 of the COVID-19 infections, Natural Killer cell

mediated cytotoxicity is one of the main enriched pathways in the mild-infected

group, whereas IL-17 signaling is the most significant pathway in the severe-infected

group (Figure 2D). This finding is remarkable because besides COVID19 IL-17 is

affiliated with other clinical pathophysiologies in which a dysregulation between

innate and adaptive immune responses such as myocarditis and lupus (S. Y. Lee et

al., 2019; Rangachari et al., 2006; Sadeghi et al., 2021).

Taken together, we show that there are distinct transcriptional responses along the

COVID-19 progression, which suggest that immune responses to SARS-CoV-2

infection occur differently in individuals; thus, there might exist a differential

imprinting associated with the severity of the COVID-19 infection.

Higher expression of NK cell hub-genes is a core event of acute phase that

distinguishes mild from severe symptoms in COVID-19 individuals.

Given that our findings pointed out changes in the immune response after SARS-

CoV-2 infection of the patients cataloged as mild and severely ill, we decided to

uncover molecular pathways that might be responsible of the differences observed

between patient groups during COVID-19 progression. To do so, we first identified

genes that were differentially expressed between severity groups, and second, we

chose only those that also showed changes in their trajectories across sampling

times. In doing so, we found 828 genes that exhibited temporal differences in

expression level during disease progression. Then using the Enrichr platform, we

discovered additional biological processes and KEGG pathways that were

differentially enriched during the COVID-19 progression in mild and severe patients

(Figure 3). For instance, mild-infected patients exhibited expression of genes

involved in kinase activity, enzyme-linked receptor activity, and apoptotic process

not only at D0 (acute phase) but also at D7 (middle phase) (Figures 3A and 3C). In

contrast, severely ill patients exhibited high level expression of genes involved in

neutrophil activity. This observation was the most notorious outcome elicited by

SARS-CoV-2 during acute COVID-19 in this group (Figures 3B, 3D, and 3F). We

observed that Natural Killer cell cytotoxicity was the most enriched pathway among the temporal and differential expressed genes in mildly ill patients during the acute phase (Figures 3B and 3C). Among these enriched genes, we found abundant membrane receptor genes that included *KLRC1*, *KLRC3*, *KLRD1*, *KIR3DL2*, *NCR3*, as well as other intra- and extra-cellular effectors that included *SH2D1A*, *PRF1*, *GZMB*, *FASLG*, *ZAP70*, *IFNG*, *CD247*, and *LAT*. Furthermore, the *ZAP70*, *CD4*, *IFNG*, *IL2RB*, *STAT4*, *CD247*, *DLL1*, *LAT*, and IL12RB2 genes were enriched in COVID-19 mild patients during the acute phase (Figure 3 - table supplement 1 and 2). This data indicated that the Th1/Th2 cell differentiation pathway was robust and active during this phase and likely played an important role in the effective adaptation to dynamic events during the progression of the infection that protected mildly ill patients from experiencing severe symptoms. Interestingly, metabolic pathways involved with hematopoietic cell lineages were enriched in severely ill patients at this matched moment in time with the mildly ill patients (Figure 3B and 3D). Collectively, these observations indicate that coordination between humoral and cell mediated immunity were more tightly regulated in mildly ill patients than in severely ill patients. To confirm the importance of the differentially enriched pathways between mild and severe COVID-19 patients, we focused on analyzing the context of gene-gene interactions (Figure 4A and Figure 4 - figure supplement 1) and changes in their quantitative expression levels overtime graphed as a heatmap (Figure 4B). The genes displayed in this KEGG pathway graph represent the up-regulated genes (red boxes) in mild patients and their interactions involved in NK cell-mediated cytotoxicity (Figure 4A). Interestingly, all these genes showed overtime trajectories with high levels on days 0 and 7 in mild patients. These gene expression levels became roughly equivalent by D28 in both the mild and severe groups (Figure 4B). Complementing these observations, we constructed a protein-protein interaction (PPI) network using only upregulated genes during the early phase (days 0 and 7), followed by a clustering process that detected proteins with more significant interactions among the selected genes (Figure 4C). Notably, we detected KLRD1, CD247, and IFNG as central nodes of protein-protein interaction networks. This finding makes sense because these proteins exhibit numerous interactions with

other proteins involved in activating or inhibiting NK cell cytotoxicity (e.g., KLRC1, KLRC3, and KIR3DL2), as well as Th1/Th2 cell differentiation (CD4) and cytokine-cytokine receptor interaction (IL5RA, IL2RB). In figure 4D, we show the comparative trajectories of these node-genes between both groups of severity. Interestingly, we found a convergence of KLRD1 and CD247 genes on D28, while IFNG remained differentially expressed between patient groups.

Once we identified the trajectories of NK cell hub-genes participating in COVID-19 progression, we asked whether there were any DEGs (adj. $p \leq 0.05$ and $\log_2$-fold change ≥ 2.0) obtained from a pairs-wise comparison of mildly and severely ill patients at days 0 and 7 that would have been left out from the longitudinal analysis. Given that the number of DEGs at each time point is higher when compared to the list of genes exhibiting differential trajectories, we performed a GO and pathways analysis with the new set list of genes (Figure 4 - figure supplement 2). The main result showed that Natural Killer cell mediated cytotoxicity was predominant on D7. This finding reinforced the interpretation that there is a dysregulation of innate immunity, as previously suggested in severe patients (Paludan & Mogensen, 2022), with an over-representation of neutrophil activation. The results shown in Figure 4 - figure supplement 3 and 4 summarize the pathway and PPI network analysis for these genes on D0 and D7, respectively, and show the predominant enrichment of NK genes. Taken together, these data are consistent with an active and regulated innate NK cytotoxic immune response mounted during the acute phase of infection in mild COVID-19 patients. This observation contrasts with the humoral- and neutrophil-biased response observed in severely ill individuals.

Previous comparisons were done with the assumption that both cohorts were at the peak of their symptoms on D0. However, taking into account the delta at the days of symptoms onset, we also analyzed the pairwise comparison for D7 in mild patients, but this time comparing it to D0 in severe patients. The Figure 4 - figure supplement 5 highlights GO terms related to Natural Killer cell response and enriched pathways for NK cell cytotoxicity in mild patients. This result supports the idea that the transcriptional program differs between mild and severe individuals with the outstanding contribution of NK cells in mild patients. Otherwise, severe patients

352 exhibit an inflammatory response reflected by the complement and coagulation
353 pathways, which is according to our previous analysis.

355 Gene co-expression identifies NK hub genes linked to the innate and adaptive
356 immune response of mild COVID-19 patients

We identified genes that were coordinately expressed during COVID-19. We
developed a weighted gene correlation network (WGCN) to simultaneously analyze
all peripheral blood samples collected from patients during the longitudinal protocol
and those from heathy donors to identify genes with coordinated expression. By
using a differential co-expression approach, we identified ten modules of co-
expressed genes (Figure 5A). We then used these networks to correlate each
module with available clinical information of the patients by calculating the module
significance (MS) for each module-trait correlation. Not surprisingly, we found that
most module eigen genes grouped according to the degree of COVID-19 infection
(i.e., mild or severe patients) (Figure 5B). Among co-expressed gene modules, we
focused on three modules that contained the largest number of genes. These
modules correspond to blue (704 genes), brown (508 genes) and turquoise (712
genes). The blue and brown modules, which are correlated positively with mild
patients (Figure 5B), were enriched with genes related to T-cell activation and
platelet function, respectively (Figure 5C). In contrast, the turquoise module, which
was correlated positively with severe COVID-19 patients (Figure 5B), was enriched
with genes related to neutrophil activation and inflammatory responses (Figure 5C).
As shown thus far, the Th1/Th2 cell differentiation pathway was relevant in the
immune response of mild COVID-19 patients, and because the blue module is
enriched in lymphocyte-based immune response genes, we performed a gene-gene
network analysis to determine how the genes from this module might have worked
in the context of an adaptive immune response. In this analysis, we found genes
belonging to the NK cell-mediated cytotoxicity pathway grouped together with the
cytokine-cytokine receptor interaction and Th1/Th2 cell differentiation pathways
(Figure 5D). Furthermore, these genes were previously identified as differentially

expressed in the NK cytolytic pathway, like *KLRD1*, *KLRC3,* and *KLRC1* receptors,

as well as *FASLG*, *SH2DB1A/B,* and *LAT*. All of this evidence is consistent with the

interpretation that highly interconnected genes from blue module had a functional

significance in limiting the progression COVID-19 in mild patients. In the other

modules, the brown network (Figure 5E) depicts enriched pathways for platelet

activation, extracellular matrix-receptor interaction, hematopoietic cell lineage, gap

junction, and complement and coagulation cascades which complemented with GO

terms is an important focus of interest in COVID19 (S.B et al., 2024). Furthermore,

the enriched pathways for the turquoise module include leishmaniasis, malaria,

osteoclast differentiation, and nitrogen metabolism (Figure 5F), some of which are

implicated in a neutrophil response as indicated by the GO terms (Babatunde &

Adenuga, 2022; Passelli, Billion, & Tacchini-Cottier, 2021).

The remaining seven modules were analyzed for Gene Ontology (Figure 5 - figure

supplement 1) which reveals different aspects related to the immunological response

to viral infection. In addition, we searched for enriched pathways and displayed the

results in the form of a network graph (Figure 5 - figure supplement 2) that can be

further investigated in future studies.

Discussion

We systematically analyzed transcriptomic features of PBMCs from COVID-19

patients with mild and severe symptoms at three sequential time-points (D0, D7, and

D28) during the peak of the symptoms. Our longitudinal analysis revealed key

temporal features of immune responses that distinguished mild from severe patients

during acute disease. We observed a prominent role of Natural Killer (NK) cell

mediated cytotoxicity function pathways during COVID-19 progression. These

pathways include genes such as *KLRC1*, *KLRC3*, *KLRD1*, *KIR3DL2*, and *NCR3*

receptors, as well as other effectors like *SH2D1A*, *PRF1*, *GZMB*, *FASLG*, *ZAP70*,

IFNG, *CD247*, and *LAT*. Most, if not all, of these genes are implicated in regulatory

processes of cytotoxicity and attraction of NK cells as part of the viral infection control

mechanism (Björkström, Strunz, & Ljunggren, 2022). Antiviral NK cell cytotoxicity

depends on a steady state for survival, basal turn-over and their function

maintenance, which are monitored by several checkpoints (Björkström et al., 2022; Masselli et al., 2020; Vivier, Tomasello, Baratin, Walzer, & Ugolini, 2008). We found a dynamic transcriptomic profile of a NK cell gene-hub characterized by higher gene expression levels in individuals with mild disease compared with those with severe symptoms across 0 and 7 days. However, expression levels of these NK cell gene-hubs became more similar between mild and severe patients on D28. In contrast to this orchestrated transcriptional response of dominant NK cells activities, we found an up-regulated gene signature consistent with dominant neutrophil activities in our severe cohort even after recovery. This finding was previously observed with the concomitant increase of IgG production and complement activation at the earliest phase of disease (J. Wang et al., 2020; B. Zhang et al., 2020; Zuo et al., 2020), (Figure 2).

In our NK cell gene hub, we recognized activating (*KLRC3*, *NCR3*), and inhibitory (*KLRC1*, *KIR3DL2*) genes of cytotoxicity, as well as regulatory and effector proteins (*KLRD1, GZMB, and PRF1*). All these genes could be participating in the balancing of a well-coordinated NK cell activity profile. In this sense, the *KRLD1* gene, which encodes the CD94 protein, stands out as an important node interconnecting proteins networks. This node regulates activating (NKG2E from *KLRC3 gene*) and inhibitory functions (NKG2A from *KLRC1* gene), and thus modulates NK cell cytotoxicity (Borrego, Masilamani, Marusina, Tang, & Coligan, 2006). Supporting this role, a previous study demonstrated the importance of CD94:NKG2E heterodimeric receptor in response to the lethal mousepox virus (Fang et al., 2011). This node may be relevant for an efficient response against SARS-CoV-2 infection given the high conservation of receptors and ligands between the human and mouse pathways (Borrego et al., 2006). Wauters et al. found that mild COVID-19 patients displayed an interaction of CD94:NKG2E/HLA-F between their T cells and neutrophils in bronchoalveolar lavage samples (Wauters et al., 2021). On the other hand, NKG2A is an important inhibitory receptor that interacts with CD94 and together regulate NK cell functions (Borrego, Masilamani, Kabat, Sanni, & Coligan, 2005; N. Lee et al., 1998). Our analysis showed a higher expression of NKG2A in mild than severe patients during the acute phase. Regarding the same receptor, previous research

showed that NKG2A was more highly expressed in lymphocytes and NK cells during infection compared with healthy controls (Zheng et al., 2020). In parallel, Zheng et al. showed a decrease of expression of NKG2A in recovering patients along with an increase of NK cell number. Collectively, this evidence supports the conclusion that CD94, and its partners, play important roles in regulating both activating and inhibitory checkpoints related to NK cell cytolytic functions. Furthermore, it substantiates the relevance of innate NK cell immune responses in combating SARS-CoV-2, and likely other coronaviruses. This pathway might also be a prominent player in controlling other infectious respiratory virus infections to promote a mild presentation of disease.

Other genes located in the NK cell hub included membrane proteins such as *SH2D1A*, *LAT*, *CD247*, *FASLG*, the enzyme *ZAP70*, and the cytokine *IFNG*. Remarkably, genes encoding *IFNG* and *CD247* were also identified as important nodes within the protein-protein interaction network during the acute phase. Considering the interactions of these nodes with proteins involved in cytokine-cytokine receptor interactions and Th1/Th2 cell differentiation pathways, it is possible that they coordinately regulated these immune responses with NK cell cytolytic functions. In this context, cytokine-cytokine receptor interaction and Th1/Th2 cell differentiation were well-represented pathways in mild patients during the acute phase highlighting that both innate and adaptive immune responses were active and effective in these patients. Particularly, CD247 (CD3ζ) protein is part of the T-cell antigen receptor (TCR) complex, whose low expression levels have been related to chronic inflammation and decreased T cell activity (Y. Li et al., 2021). In the same line, IFNG protein is a critical player between innate and adaptive immunity after viral infection (Kang, Brown, & Hwang, 2018). Giving support to this connection between innate and adaptive immune responses, it would be expected that adaptive CD8+ T cell cytolytic functions would also be enriched in mild patients due to its important role controlling viral infections (Prager & Watzl, 2019; Uzhachenko & Shanker, 2019). Interestingly, GO/pathway-based analyses did not detect these functions as a differential player in clinical COVID-19 progression, despite the fact that some genes are shared with NK cytotoxic gene hub (Uzhachenko & Shanker, 2019). This

observation suggested that Th1/Th2 cell differentiation may be more essential for a

successful adaptive response against SARS-CoV-2 than CD8+ T cell cytolytic

function in mild patients, at least during the early phase of COVID-19. If this

interpretation is credible, then cell-mediated cytolytic activities should rely on the

well-regulated activity of innate NK cell subset as a primary immune response.

Taking into account all these data, it is reasonable to interpret that an early fate-

compromise towards NK cell activity instead of a Neutrophil effector activity may

have had an important effect on subsequent processes regulating adaptive

immunity. This model favors a robust integration of innate and adaptive immune

response during an effective control of COVID-19.

Until now, we have discussed relevant genes involved in immune pathways enriched

in mild or severe COVID-19 progression. However, we also decided to look for genes

exhibiting coordinated gene expression patterns across all our samples. We found

that one module (blue module), which has a strong positive correlation with mild

patients, included genes involved with metabolic pathways regulating T cell

activation, kinase activity, and antigen presentation. In contrast, the turquoise

module, which exhibited a strong positive correlation with severely ill individuals,

contained genes associated with neutrophil-related biological processes. This

finding was indicative of an opposed early fate of innate immune responses between

mild and severe COVID-19 cases. Neutrophil long-term differential enrichment seen

across severe cases could be related to other repercussions of SARS-CoV-2

infection, like neutrophil-induced platelet aggregation (Jevtic & Nazy, 2022).

Consistent with this interpretation, dysfunction of platelets has been associated with

abnormal clot formation in severe COVID-19 cases (Litvinov et al., 2021). In this

sense, our results show that the brown module, which is negatively correlated with

severe patients, displays biological processes linked to platelet degranulation activity

and negative regulation of aggregation. Hence, these results are consistent with a

platelet dysfunction pathology linked to severely ill patients.

We performed a pathway enrichment analysis to understand how the positive

correlation of genes in the blue co-expression module related to immune response

functions in mildly ill patients. Not surprisingly, the main pathways enriched included Th1/Th2 cell differentiation, cytokine-cytokine receptor interaction, antigen processing, and NK cell-mediated cytotoxicity pathways. Albeit the co-expression analysis included all samples, regardless of the severity of the disease or the longitudinal sampling. This effort revealed transcriptional programs of immune response that were consistent with the profiles detected in mild and severe patients reported in our previous investigations. This evidence supports the idea that the transcriptional regulation of cell mediated immunity in mildly ill patients is more robust than that observed in patients with severe clinical progression.

In order to identify differences in transcriptional programs associated with mild or severe outcomes, we carefully compared changes in gene expression during the acute phase. This analysis detected a broader list of genes than those found using the longitudinal analysis alone. This accomplishment was resulted from only considering the differences in gene expression between mild and severe groups, independently of quantitative changes in gene expression over-time. These results consistently showed a NK cell hub of genes being differentially expressed in mild patients. Importantly, we found novel DEGs including *KLRK1*, *KIR2DS4*, and *KLRC2*. The gene *KLRK1* codes for NKG2D protein, an activating receptor with critical importance due its interaction with the major histocompatibility complex-class-I (Zingoni et al., 2018). We found that NKG2D is comparatively over-expressed between days 0 and 7 in the mild group. In line with this finding, Varchetta et al. found an increase of circulating NKG2D(-) NK cells using cell cytometry during the acute phase, which was linked to exhaustion in severe COVID-19 patients. Importantly, their sampling times ranged from hours to days after onset symptoms (Varchetta et al., 2021). Additionally, the regulatory role of NKG2D in COVID-19 is also supported by Lee et al. where their results show that the viral non-structural protein 1 (Nsp1) of SARS-CoV-2 mediates its immune escape by downregulating NKG2D-Ligands, therefore decreasing NKG2D-dependent NK cytotoxic responsiveness and conferring resistance to infected cells (M. J. Lee et al., 2022).

Complementary to this scenario of NK cell activating receptors being important in

mitigating symptom severity as previously reported (Gardiner, 2008), we found that

the *KIR2DS4* gene, which belongs to the KIR receptors gene-family, was correlated

with mild progression of COVID-19. This activating receptor was more highly

expressed in mild patients than in severe patients at both 0 and 7 days (4-fold and

6.8-fold, respectively). In this regard, Bernal et al. found that a low expression of

KIR2DS4 was part of a distinctive immunophenotype in peripheral NK cells that was

increased in severe COVID-19 individuals (Bernal et al., 2021). On the contrary,

Casado et al. found an enriched KIR2DS4(+) subset of CD56brightCD16neg

peripheral NK cells in hospitalized individuals compared to mild patients, indicating

a positive correlation with severity (Casado et al., 2022). However, as their mild

patient cohort were recruited at a mean of 60 days after diagnosis, a direct

comparison with our data is not precise. Additionally, the absence of severe patients,

as well as a lack of longitudinal sampling, were some of the inconsistencies between

the study designs of Casado et. al. and ours, which may account for these

inconsistent observations.

Another NK cell activating receptor (the NKG2C protein) encoded by the *KLRC2*

gene was previously implicated as a COVID-19 marker (Fielding et al., 2022;

Maucourant et al., 2020; Vietzen et al., 2021). Although we did not find a significant

difference in the gene expression of *KLRC2* between mild and severe groups (being

excluded by our threshold criteria), we found it to be comparatively lower among

severe patients compared to mild patients in both 0 and 7 days (-1.6 fold and -2.7

fold, respectively). We observed a similar trend in the control patients between 0 and

7 days in the acute phase (-1.6 fold and -1.7 fold, respectively). Surprisingly, when

we compared the quantitative expression of the NKG2C gene in the severe group to

the control group on D28, we discovered an inverted pattern with respect to the acute

phase in which NKG2C over-passed the levels of controls (2.1-fold). In this context,

Maucourant et al, in a scRNA-Seq study performed with bronchio-alveolar lavage

(BAL) from severe COVID-19 patients, showed increased NKG2C levels linked to

the adaptive response of NK cells (Maucourant et al., 2020). These data additionally

reinforce the interpretation that NKG2C is required to mount an effective NK cell response against SARS-CoV-2 infection. Vietzen et al. found that a deletion in the NKG2C gene resulted in a significant correlation with severe COVID-19 (Vietzen et al., 2021). This evidence supports the idea that the innate and adaptive immune responses are being differentially modulated in severe COVID-19 than mild patients. Another important observation detected in our comparative analysis included enrichment of the IL-17 pathway in severe patients on D7. Our analysis identified a group of 10 genes (*FOSL1, CXCL6, CEBPB, LCN2, TNFAIP3, CXCL1, CXCL2, MMP9, S100A9,* and *S100A*) associated with this time point. IL-17 has diverse biological functions that promote protective immunity against many pathogens but also driving inflammatory pathology during autoimmunity. Interleukin-17-driven inflammation is normally controlled by regulatory T cells expressing the anti-inflammatory cytokines IL-10, TGFbeta, and IL-35 (Pacha, Sallman, & Evans, 2020). One explanation may be that an imbalance in T cells and cytokine secretions mediated by IL-17 promote an inflammatory phenotype in patients with severe symptoms. Notably, Th17 cells were elevated on D7 in patients with mild symptoms. These Th17 cells can display plasticity in cytokine production *in vivo* and can switch from predominantly producing IL-17 to predominantly producing IFNγ, thereby resembling Th1 cells (Y. K. Lee, Mukasa, Hatton, & Weaver, 2009). Sequential activation of STAT1 by IFNG and STAT4 by IL-12 drives optimal expression of T-bet (TBX21), a central transcription factor for Th1 programming (Y. K. Lee et al., 2009). Otherwise, the activation of STAT6 by IL-4 upregulates GATA3, which is central to Th2 programming (Y. K. Lee et al., 2009). All the genes related to Th1 activity (*IFNG, STAT4, TBX21,* and *IL-12*) were up-regulated in our cohort of patients exhibiting mild symptoms, which is consistent with this potential regulatory circuit.

Even though the primary criterion for comparing mild with severe was the peak of symptoms in earlier analyses, we also investigate according to the time of the symptoms onset. As a result, we also performed a complement analysis comparing mild-D7 to severe-D0 to identify the enriched pathways that represent this comparison. Notably, the most important GO-terms and enriched pathways in mild patients correspond to Natural Killer cell cytotoxicity, represented by the same set of

genes reported in previous comparisons. This lends significant support to the
primary findings of our study, which show that a transcriptional program is
differentially regulated in mild patients and is focused on innate immune response
associated with Natural Killer cell cytotoxicity.

In conclusion, the longitudinal trajectories of gene expression, the differential
GO/Pathways, and protein-protein interactions analyses, together with the co-
expressed gene-gene correlation network is consistent with the existence of a
regulatory transcriptional program linked to an early activation of NK cell cytotoxicity
in mild COVID-19 patients. This work establishes the notion that innate immune
responses are crucial for the progression of COVID-19 severity, and reinforces the
importance of the NK cell cytotoxicity pathway in distinguishing between mild and
severe COVID-19 progression. Taken together, these differential responses are
complemented by cytokine activities and Th1/Th2 cell differentiation programs
indicating a well-regulated crosstalk between innate and adaptive immune
responses in the mild COVID-19 progression.

Methods

Patient cohort and Peripheral Blood Mononuclear Cells (PBMCs) sampling

We recruited a total of 8 patients diagnosed to be suffering from COVID-19 who were
separated into two groups, one composed of 4 mild outpatients and another,
conformed of 4 severe hospitalized individuals. Peripheral venous blood samples
were obtained by using the venipuncture technique in Vacutainer K2 EDTA tubes
(BD, USA) from each patient three times, including two clinical stages (acute phase
and convalescence). PBMC were isolated from each fresh heparinized peripheral
blood sample, through density gradient centrifugation on Ficoll-Paque Plus (GE
Healthcare Life Sciences, USA) by centrifuging at 1600 rpm for 30 minutes (using
minimum acceleration and no deceleration configurations). PBMC-containing
fraction was then washed two times with 2 mM EDTA in PBS and stored in RNAlater
solution (Sigma, USA) at -20° C until RNA extraction. The detailed clinical features
of all patients and the detailed sampling time are shown in Figure 1 and Table 1. All

samples were processed in a qualified BSL-2 laboratory and, according to protocols

and approval from Institutional Review Boards, CEC-SSLR Ord N°226 and Ord

N°399. Written informed consent was received before the participation of each

patient.

RNA extraction, library preparation, and PBMC transcriptome sequencing

Total RNA extraction was performed from PBMC by Diagenode (Belgium). RNA

samples were quantified using QubitTM RNA BR Assay Kit (Thermo Fisher

Scientific, USA) and secondly checked for integrity using HS RNA Kit (Agilent, USA)

on a Fragment analyzer system (Agilent, USA). The library preparation was

performed using NEBnext ultraII Directional Kit and sequencing of the samples was

performed on an Illumina NovaSeq 6000 instrument producing 150bp paired-end

reads running Control Software 1.7.0.

Identification of differentially expressed genes along COVID-19 progression

and between disease severities.

Sequencing-quality check was performed using FastQC (Andrews, 2010), and low-

quality reads were trimmed using Trim_Galore! (Krueger) with --clip_R1 3 and --

clip_R2 3 options. High-quality reads were aligned to the human reference genome

version GRCh38 with STAR v2.6.1a_08-27 (S. Zhang et al., 2022). Transcript counts

were generated using featureCounts v1.6.3 (Bastard et al., 2020) with default

settings. Differential gene expression analysis was performed in two ways using

edgeR package v3.36.0 (Delorey et al., 2021). Temporally and differentially

expressed genes and differentially expressed genes between mild and severe

COVID-19 patients at each sampling timepoint were identified using the generalized

log-linear model (GLM) option in edgeR. Genes were considered as differentially

expressed either with temporal expression differences or disease severity condition

using a false discovery rate (FDR; Benjamini–Hochberg), an adjusted p-value of

<0.05, and an absolute log2 fold change of 2. Transcript counts (normalized using

TMM approach) were used to generate heatmaps for visualization of differentially

expressed genes using the pheatmap R package. Expression was scaled by row z-

scores for visualization, taking into account the mild and severe patients to obtain the mean and standard deviation of each group.

Gene set enrichment analysis (GSEA)

In order to perform GSEA (Schultze & Aschenbrenner, 2021), after DEGs analysis only values of log2 fold change equal to or greater than 1 were considered. Then the GSEA was performed through ClusterProfiler package (v.3.16) (Yu, Wang, Han, & He, 2012) in R, we evaluated the significantly enriched biological process (GO) using gseGO function and pathways from Kyoto encyclopedia of gene and genomes (KEGG) using gseKEGG function.

Enrichment analysis of differentially expressed genes along COVID-19 progression and between mild and severe patients

Gene ontology (GO) and pathways enrichment analyses were performed with both upregulated and downregulated genes using Enrichr platform (Xiong et al., 2020). Significant GO terms (Biological processes and Molecular functions) and pathways (KEGG and Reactome) were calculated from the adjusted p-value (q-value) using the Benjamini-Hochberg method for correction for multiple hypothesis. Considering a difference of one unit in z-score between the two severity groups, for mild patients we analyzed 365, 359, and 101 genes for days 0, 7, and 28 respectively, while for severe patients we analyzed 369, 414, and 136 genes for day 0, 7, and 28 respectively.

Weighted correlation network analysis (WGCNA)

Based on the assumption that differentially expressed genes may explain transcriptional differences observed between mild and severe COVID-19 patients. Read counts from differentially expressed genes among all samples were selected as a reference set for construction of a weighted gene co-expression networks and modules detection. Co-expressed gene modules were constructed using WGCNA R package v1.71 (Langfelder & Horvath, 2008) under a signed networks approach because it provides a better understanding of molecular regulatory mechanisms at

the systemic level, facilitating better separation of modules in terms of biological

performances. To do so, we removed outliers using the adjacency function and a

standardized connectivity score of < -2.0; then we used the pickSoftThreshold

function to identify the soft thresholding power ß value, which was subsequently

transformed into a Topological Overlap Matrix (TOM). Next, an average linkage

hierarchical clustering analysis was performed based on the TOM dissimilarity (1-

TOM) and modules were detected through a dynamic hybrid tree cutting algorithm.

After the modules were identified, the module eigengene (ME) was summarized by

the first principal component of the module expression levels. Module-trait

relationships (MTRs) were estimated using Pearson correlation between MEs and

disease severity. To evaluate the correlation strength, we calculated the module

significance (MS) that is defined as the average absolute gene significance (GS) of

all genes involved in a module.

PPI network

All genes with differential trajectories over time (#827 genes) were included to

construct a protein-protein interaction network (PPI) by STRING V11.5 (Szklarczyk

et al., 2020). After clusterization, we focused our analysis on the principal cluster (1

of 3) with the highest connectivity between genes and highlighted the genes involved

in the most significant pathways according to KEGG.

Acknowledgements

711 We especially thank all blood donors who participated in this study. COVID-19 South Chile Group: Renato Ocampo, Christian Esveile, Leonila Ferreira, Johanna Cabrera, Geraro Solís, Marcelo Pozas, Equipo de enfermería Servicio de Medicina Interna Hospital Base Osorno, María Paz Contreras, Equipo de enfermería de la Universidad de Los Lagos Osorno, Catherine Fernández, Camila Rojas, Paulina Lagos, Rocío Mejías, Melissa Canales, Patricio Suazo, Pamela Ángel, Romina Inostroza, Pamela Silva, Felipe Collado, Vanina Cuevas, Rodrigo Oñate, Daniel Salamanca, Javier Briones, Vanessa Villagrán, Diana Bocaz, Andrés Umanzor.

This work was supported by the National Agency of Research and Development (ANID) projects COVID0422 and FONDEF ID20I10192. We also thank USS-FIN-23-PDOC-02.

References

Ahern, D. J., Ai, Z., Ainsworth, M., Allan, C., Allcock, A., Angus, B., . . . Zurke, Y.-X. (2022). A blood atlas of COVID-19 defines hallmarks of disease severity and specificity. *Cell, 185*(5), 916-938.e958. doi:https://doi.org/10.1016/j.cell.2022.01.012

Andrews, S. (2010). FastQC: A quality control tool for high throughput sequence data. Retrieved from https://www.bioinformatics.babraham.ac.uk/projects/fastqc/

Babatunde, K. A., & Adenuga, O. F. (2022). Neutrophils in malaria: A double-edged sword role. *Frontiers in immunology, 13*, 922377. doi:10.3389/fimmu.2022.922377

Bastard, P., Rosen, L. B., Zhang, Q., Michailidis, E., Hoffmann, H.-H., Zhang, Y., . . . Casanova, J.-L. (2020). Autoantibodies against type I IFNs in patients with life-threatening COVID-19. *370*(6515), eabd4585. doi:10.1126/science.abd4585 %J Science

Bernal, E., Gimeno, L., Alcaraz, M. J., Quadeer, A. A., Moreno, M., Martínez-Sánchez, M. V., . . . group, o. b. o. M. C. S. (2021). Activating Killer-cell Immunoglobulin-like Receptors are associated with the severity of COVID-19. *The Journal of Infectious Diseases*. doi:10.1093/infdis/jiab228

Bernardes, J. P., Mishra, N., Tran, F., Bahmer, T., Best, L., Blase, J. I., . . . Ziebuhr, J. (2020). Longitudinal Multi-omics Analyses Identify Responses of Megakaryocytes, Erythroid Cells, and Plasmablasts as Hallmarks of Severe COVID-19. *Immunity, 53*(6), 1296-1314.e1299. doi:https://doi.org/10.1016/j.immuni.2020.11.017

Björkström, N. K., Strunz, B., & Ljunggren, H.-G. (2022). Natural killer cells in antiviral immunity. *Nature Reviews Immunology, 22*(2), 112-123. doi:10.1038/s41577-021-00558-3

Borrego, F., Masilamani, M., Kabat, J., Sanni, T. B., & Coligan, J. E. (2005). The cell biology of the human natural killer cell CD94/NKG2A inhibitory receptor. *Molecular Immunology, 42*(4), 485-488. doi:https://doi.org/10.1016/j.molimm.2004.07.031

Borrego, F., Masilamani, M., Marusina, A. I., Tang, X., & Coligan, J. E. (2006). The CD94/NKG2 family of receptors. *Immunologic Research, 35*(3), 263-277. doi:10.1385/IR:35:3:263

Casado, J. L., Moraga, E., Vizcarra, P., Velasco, H., Martín-Hondarza, A., Haemmerle, J., . . . Vallejo, A. (2022). Expansion of CD56dimCD16neg NK Cell Subset and Increased Inhibitory KIRs in Hospitalized COVID-19 Patients. *Viruses, 14*(1), 46. Retrieved from https://www.mdpi.com/1999-4915/14/1/46

Chen, Z., & John Wherry, E. (2020). T cell responses in patients with COVID-19. *Nature Reviews Immunology, 20*(9), 529-536. doi:10.1038/s41577-020-0402-6

Daamen, A. R., Bachali, P., Owen, K. A., Kingsmore, K. M., Hubbard, E. L., Labonte, A. C., . . . Lipsky, P. E. (2021). Comprehensive transcriptomic analysis of COVID-19 blood, lung, and airway. *Scientific Reports, 11*(1), 7052. doi:10.1038/s41598-021-86002-x

Delorey, T. M., Ziegler, C. G. K., Heimberg, G., Normand, R., Yang, Y., Segerstolpe, Å., . . . Regev, A. (2021). COVID-19 tissue atlases reveal SARS-CoV-2 pathology and cellular targets. *Nature, 595*(7865), 107-113. doi:10.1038/s41586-021-03570-8

Fang, M., Orr, Mark T., Spee, P., Egebjerg, T., Lanier, Lewis L., & Sigal, Luis J. (2011). CD94 Is Essential for NK Cell-Mediated Resistance to a Lethal Viral Disease. *Immunity, 34*(4), 579-589. doi:https://doi.org/10.1016/j.immuni.2011.02.015

Fielding, C. A., Sabberwal, P., Williamson, J. C., Greenwood, E. J. D., Crozier, T. W. M., Zelek, W., . . . Stanton, R. J. (2022). SARS-CoV-2 host-shutoff impacts innate NK cell functions, but antibody-dependent NK activity is strongly activated through non-spike antibodies. *eLife, 11*, e74489. doi:10.7554/eLife.74489

Gardiner, C. M. (2008). Killer cell immunoglobulin-like receptors on NK cells: the how, where and
why. *International Journal of Immunogenetics, 35*(1), 1-8.
doi:https://doi.org/10.1111/j.1744-313X.2007.00739.x
Ge, H., Wang, X., Yuan, X., Xiao, G., Wang, C., Deng, T., . . . Xiao, X. (2020). The epidemiology and
clinical information about COVID-19. *European Journal of Clinical Microbiology &*
Infectious Diseases, 39(6), 1011-1019. doi:10.1007/s10096-020-03874-z
Haitao, T., Vermunt, J. V., Abeykoon, J., Ghamrawi, R., Gunaratne, M., Jayachandran, M., . . .
Garovic, V. D. (2020). COVID-19 and Sex Differences: Mechanisms and Biomarkers. *Mayo*
Clinic Proceedings, 95(10), 2189-2203. doi:https://doi.org/10.1016/j.mayocp.2020.07.024
Janssen, N. A. F., Grondman, I., de Nooijer, A. H., Boahen, C. K., Koeken, V., Matzaraki, V., . . . van
de Veerdonk, F. L. (2021). Dysregulated Innate and Adaptive Immune Responses
Discriminate Disease Severity in COVID-19. *J Infect Dis, 223*(8), 1322-1333.
doi:10.1093/infdis/jiab065
Jevtic, S. D., & Nazy, I. (2022). The COVID Complex: A Review of Platelet Activation and Immune
Complexes in COVID-19. *Frontiers in immunology, 13*, 807934.
doi:10.3389/fimmu.2022.807934
Kang, S., Brown, H. M., & Hwang, S. (2018). Direct Antiviral Mechanisms of Interferon-Gamma.
Immune Netw, 18(5). Retrieved from https://doi.org/10.4110/in.2018.18.e33
Khalil, B. A., Elemam, N. M., & Maghazachi, A. A. (2021). Chemokines and chemokine receptors
during COVID-19 infection. *Computational and Structural Biotechnology Journal, 19*, 976-
988. doi:https://doi.org/10.1016/j.csbj.2021.01.034
Krueger, F. *Trim Galore!* Retrieved from
https://www.bioinformatics.babraham.ac.uk/projects/trim_galore/
Ladds, E., Rushforth, A., Wieringa, S., Taylor, S., Rayner, C., Husain, L., & Greenhalgh, T. (2020).
Persistent symptoms after Covid-19: qualitative study of 114 "long Covid" patients and
draft quality principles for services. *BMC Health Serv Res, 20*(1), 1144.
doi:10.1186/s12913-020-06001-y
Langfelder, P., & Horvath, S. (2008). WGCNA: an R package for weighted correlation network
analysis. *BMC Bioinformatics, 9*(1), 559. doi:10.1186/1471-2105-9-559
Lee, M. J., Leong, M. W., Rustagi, A., Beck, A., Zeng, L., Holmes, S., . . . Blish, C. A. (2022). SARS-
CoV-2 escapes direct NK cell killing through Nsp1-mediated downregulation of ligands for
NKG2D. *Cell Reports, 41*(13). doi:10.1016/j.celrep.2022.111892
Lee, N., Llano, M., Carretero, M., Ishitani, A., Navarro, F., López-Botet, M., & Geraghty, D. E.
(1998). HLA-E is a major ligand for the natural killer inhibitory receptor CD94/NKG2A.
Proceedings of the National Academy of Sciences, 95(9), 5199-5204.
doi:doi:10.1073/pnas.95.9.5199
Lee, S. Y., Lee, S. H., Seo, H. B., Ryu, J. G., Jung, K., Choi, J. W., . . . Cho, M. L. (2019). Inhibition of IL-
17 ameliorates systemic lupus erythematosus in Roquin(san/san) mice through regulating
the balance of TFH cells, GC B cells, Treg and Breg. *Sci Rep, 9*(1), 5227.
doi:10.1038/s41598-019-41534-1
Lee, Y. K., Mukasa, R., Hatton, R. D., & Weaver, C. T. (2009). Developmental plasticity of Th17 and
Treg cells. *Current Opinion in Immunology, 21*(3), 274-280.
doi:https://doi.org/10.1016/j.coi.2009.05.021
Li, X., Zhong, X., Wang, Y., Zeng, X., Luo, T., & Liu, Q. (2021). Clinical determinants of the severity of
COVID-19: A systematic review and meta-analysis. *PLOS ONE, 16*(5), e0250602.
doi:10.1371/journal.pone.0250602

Li, Y., Chen, S., Li, X., Wang, X., Li, H., Ning, S., & Chen, H. (2021). CD247, a Potential T Cell–Derived
Disease Severity and Prognostic Biomarker in Patients With Idiopathic Pulmonary Fibrosis.
Frontiers in immunology, 12. doi:10.3389/fimmu.2021.762594
Litvinov, R. I., Evtugina, N. G., Peshkova, A. D., Safiullina, S. I., Andrianova, I. A., Khabirova, A. I., . . .
Weisel, J. W. (2021). Altered platelet and coagulation function in moderate-to-severe
COVID-19. *Scientific Reports, 11*(1), 16290. doi:10.1038/s41598-021-95397-6
Mann, E. R., Menon, M., Knight, S. B., Konkel, J. E., Jagger, C., Shaw, T. N., . . . Hussell, T. (2020).
Longitudinal immune profiling reveals key myeloid signatures associated with COVID-19.
Science Immunology, 5(51), eabd6197. doi:doi:10.1126/sciimmunol.abd6197
Masood, K. I., Yameen, M., Ashraf, J., Shahid, S., Mahmood, S. F., Nasir, A., . . . Hasan, Z. (2021).
Upregulated type I interferon responses in asymptomatic COVID-19 infection are
associated with improved clinical outcome. *Scientific Reports, 11*(1). doi:10.1038/s41598-
021-02489-4
Masselli, E., Vaccarezza, M., Carubbi, C., Pozzi, G., Presta, V., Mirandola, P., & Vitale, M. (2020). NK
cells: A double edge sword against SARS-CoV-2. *Advances in Biological Regulation, 77*,
100737. doi:https://doi.org/10.1016/j.jbior.2020.100737
Maucourant, C., Filipovic, I., Ponzetta, A., Aleman, S., Cornillet, M., Hertwig, L., . . . Karolinska, C.-S.
G. (2020). Natural killer cell immunotypes related to COVID-19 disease severity. *Sci
Immunol, 5*(50), eabd6832. doi:10.1126/sciimmunol.abd6832
Nalbandian, A., Sehgal, K., Gupta, A., Madhavan, M. V., McGroder, C., Stevens, J. S., . . . Wan, E. Y.
(2021). Post-acute COVID-19 syndrome. *Nat Med, 27*(4), 601-615. doi:10.1038/s41591-
021-01283-z
Newton, A. H., Cardani, A., & Braciale, T. J. (2016). The host immune response in respiratory virus
infection: balancing virus clearance and immunopathology. *Seminars in Immunopathology,
38*(4), 471-482. doi:10.1007/s00281-016-0558-0
Notarbartolo, S., Ranzani, V., Bandera, A., Gruarin, P., Bevilacqua, V., Putignano, A. R., . . .
Abrignani, S. (2021). Integrated longitudinal immunophenotypic, transcriptional and
repertoire analyses delineate immune responses in COVID-19 patients. *Science
Immunology, 6*(62). doi:10.1126/sciimmunol.abg5021
Pacha, O., Sallman, M. A., & Evans, S. E. (2020). COVID-19: a case for inhibiting IL-17? *Nature
Reviews Immunology, 20*(6), 345-346. doi:10.1038/s41577-020-0328-z
Paludan, S. R., & Mogensen, T. H. (2022). Innate immunological pathways in COVID-19
pathogenesis. *Science Immunology, 7*(67), eabm5505.
doi:doi:10.1126/sciimmunol.abm5505
Park, M. D. (2020). Immune evasion via SARS-CoV-2 ORF8 protein? *Nature Reviews Immunology,
20*(7), 408-408. doi:10.1038/s41577-020-0360-z
Passelli, K., Billion, O., & Tacchini-Cottier, F. (2021). The Impact of Neutrophil Recruitment to the
Skin on the Pathology Induced by Leishmania Infection. *Frontiers in immunology, 12*.
doi:10.3389/fimmu.2021.649348
Ponti, G., Maccaferri, M., Ruini, C., Tomasi, A., & Ozben, T. (2020). Biomarkers associated with
COVID-19 disease progression. *Critical Reviews in Clinical Laboratory Sciences, 57*(6), 389-
399. doi:10.1080/10408363.2020.1770685
Prager, I., & Watzl, C. (2019). Mechanisms of natural killer cell-mediated cellular cytotoxicity.
Journal of Leukocyte Biology, 105(6), 1319-1329. doi:https://doi.org/10.1002/JLB.MR0718-
269R
Rabaan, A. A., Smajlovic, S., Tombuloglu, H., Cordic, S., Hajdarevic, A., Kudic, N., . . . Al-Suhaimi, E.
(2022). SARS-CoV-2 infection and multi-organ system damage: a review. *Bosn J Basic Med
Sci*. doi:10.17305/bjbms.2022.7762

Rangachari , M., Mauermann , N., Marty , R. R., Dirnhofer , S., Kurrer , M. O., Komnenovic , V., . . .
Eriksson , U. (2006). T-bet negatively regulates autoimmune myocarditis by suppressing
local production of interleukin 17. *Journal of Experimental Medicine, 203*(8), 2009-2019.
doi:10.1084/jem.20052222
Rashid, F., Xie, Z., Suleman, M., Shah, A., Khan, S., & Luo, S. (2022). Roles and functions of SARS-
CoV-2 proteins in host immune evasion. *Frontiers in immunology, 13*.
doi:10.3389/fimmu.2022.940756
Ren, X., Wen, W., Fan, X., Hou, W., Su, B., Cai, P., . . . Zhang, Z. (2021). COVID-19 immune features
revealed by a large-scale single-cell transcriptome atlas. *Cell, 184*(7), 1895-1913.e1819.
doi:https://doi.org/10.1016/j.cell.2021.01.053
Ryan, F. J., Hope, C. M., Masavuli, M. G., Lynn, M. A., Mekonnen, Z. A., Yeow, A. E. L., . . . Lynn, D. J.
(2022). Long-term perturbation of the peripheral immune system months after SARS-CoV-
2 infection. *BMC Medicine, 20*(1), 26. doi:10.1186/s12916-021-02228-6
S.B, M. J., Chacko, B., Selvarajan, S., Peter, J. V., Geevar, T., Dave, R. G., . . . Srivastava, A. (2024).
Biomarkers of coagulation, endothelial, platelet function, and fibrinolysis in patients with
COVID-19: a prospective study. *Scientific Reports, 14*(1), 2011. doi:10.1038/s41598-024-
51908-9
Sadeghi, A., Tahmasebi, S., Mahmood, A., Kuznetsova, M., Valizadeh, H., Taghizadieh, A., . . .
Ahmadi, M. (2021). Th17 and Treg cells function in SARS-CoV2 patients compared with
healthy controls. *Journal of Cellular Physiology, 236*(4), 2829-2839.
doi:https://doi.org/10.1002/jcp.30047
Samadizadeh, S., Masoudi, M., Rastegar, M., Salimi, V., Shahbaz, M. B., & Tahamtan, A. (2021).
COVID-19: Why does disease severity vary among individuals? *Respir Med, 180*, 106356.
doi:10.1016/j.rmed.2021.106356
Schultze, J. L., & Aschenbrenner, A. C. (2021). COVID-19 and the human innate immune system.
Cell, 184(7), 1671-1692. doi:10.1016/j.cell.2021.02.029
Sette, A., & Crotty, S. (2021). Adaptive immunity to SARS-CoV-2 and COVID-19. *Cell*.
doi:https://doi.org/10.1016/j.cell.2021.01.007
Shen, J., Fan, J., Zhao, Y., Jiang, D., Niu, Z., Zhang, Z., & Cao, G. (2023). Innate and adaptive
immunity to SARS-CoV-2 and predisposing factors. *Frontiers in immunology, 14*.
doi:10.3389/fimmu.2023.1159326
Subramanian, A., Nirantharakumar, K., Hughes, S., Myles, P., Williams, T., Gokhale, K. M., . . .
Haroon, S. (2022). Symptoms and risk factors for long COVID in non-hospitalized adults.
Nat Med, 28(8), 1706-1714. doi:10.1038/s41591-022-01909-w
Sun, C., Xie, C., Bu, G.-L., Zhong, L.-Y., & Zeng, M.-S. (2022). Molecular characteristics, immune
evasion, and impact of SARS-CoV-2 variants. *Signal Transduction and Targeted Therapy,
7*(1), 202. doi:10.1038/s41392-022-01039-2
Szklarczyk, D., Gable, A. L., Nastou, K. C., Lyon, D., Kirsch, R., Pyysalo, S., . . . von Mering, C. (2020).
The STRING database in 2021: customizable protein–protein networks, and functional
characterization of user-uploaded gene/measurement sets. *Nucleic Acids Research,
49*(D1), D605-D612. doi:10.1093/nar/gkaa1074
The, C.-H. G. I. (2020). The COVID-19 Host Genetics Initiative, a global initiative to elucidate the
role of host genetic factors in susceptibility and severity of the SARS-CoV-2 virus
pandemic. *European Journal of Human Genetics, 28*(6), 715-718. doi:10.1038/s41431-020-
0636-6
Thorne, L. G., Bouhaddou, M., Reuschl, A.-K., Zuliani-Alvarez, L., Polacco, B., Pelin, A., . . . Krogan,
N. J. (2022). Evolution of enhanced innate immune evasion by SARS-CoV-2. *Nature,
602*(7897), 487-495. doi:10.1038/s41586-021-04352-y

Uzhachenko, R. V., & Shanker, A. (2019). CD8+ T Lymphocyte and NK Cell Network: Circuitry in the
Cytotoxic Domain of Immunity. *Frontiers in immunology, 10*.
doi:10.3389/fimmu.2019.01906
Vabret, N., Britton, G. J., Gruber, C., Hegde, S., Kim, J., Kuksin, M., . . . Laserson, U. (2020).
Immunology of COVID-19: Current State of the Science. *Immunity, 52*(6), 910-941.
doi:https://doi.org/10.1016/j.immuni.2020.05.002
Varchetta, S., Mele, D., Oliviero, B., Mantovani, S., Ludovisi, S., Cerino, A., . . . Mondelli, M. U.
(2021). Unique immunological profile in patients with COVID-19. *Cellular & Molecular
Immunology, 18*(3), 604-612. doi:10.1038/s41423-020-00557-9
Vietzen, H., Zoufaly, A., Traugott, M., Aberle, J., Aberle, S. W., & Puchhammer-Stöckl, E. (2021).
Deletion of the NKG2C receptor encoding KLRC2 gene and HLA-E variants are risk factors
for severe COVID-19. *Genetics in Medicine, 23*(5), 963-967.
doi:https://doi.org/10.1038/s41436-020-01077-7
Vivier, E., Tomasello, E., Baratin, M., Walzer, T., & Ugolini, S. (2008). Functions of natural killer
cells. *Nature Immunology, 9*(5), 503-510. doi:10.1038/ni1582
Wang, C., Huyan, T., Zhou, X., Zhang, X., Duan, S., Gao, S., . . . Li, Q. (2022). Development of Single-
Cell Transcriptomics and Its Application in COVID-19. *Viruses, 14*(10).
doi:10.3390/v14102271
Wang, J., Li, Q., Yin, Y., Zhang, Y., Cao, Y., Lin, X., . . . Qiu, Y. (2020). Excessive Neutrophils and
Neutrophil Extracellular Traps in COVID-19. *Frontiers in immunology, 11*.
doi:10.3389/fimmu.2020.02063
Wauters, E., Van Mol, P., Garg, A. D., Jansen, S., Van Herck, Y., Vanderbeke, L., . . . collaborators, C.
(2021). Discriminating mild from critical COVID-19 by innate and adaptive immune single-
cell profiling of bronchoalveolar lavages. *Cell Research, 31*(3), 272-290.
doi:10.1038/s41422-020-00455-9
Wu, Z., & McGoogan, J. M. (2020). Characteristics of and Important Lessons From the Coronavirus
Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72 314 Cases From
the Chinese Center for Disease Control and Prevention. *JAMA, 323*(13), 1239-1242.
doi:10.1001/jama.2020.2648
Xiong, Y., Liu, Y., Cao, L., Wang, D., Guo, M., Jiang, A., . . . Chen, Y. (2020). Transcriptomic
characteristics of bronchoalveolar lavage fluid and peripheral blood mononuclear cells in
COVID-19 patients. *Emerging microbes & infections, 9*(1), 761-770.
doi:10.1080/22221751.2020.1747363
Yu, G., Wang, L.-G., Han, Y., & He, Q.-Y. (2012). clusterProfiler: an R Package for Comparing
Biological Themes Among Gene Clusters. *OMICS: A Journal of Integrative Biology, 16*(5),
284-287. doi:10.1089/omi.2011.0118
Zhang, B., Zhou, X., Zhu, C., Song, Y., Feng, F., Qiu, Y., . . . Wang, J. (2020). Immune Phenotyping
Based on the Neutrophil-to-Lymphocyte Ratio and IgG Level Predicts Disease Severity and
Outcome for Patients With COVID-19. *Frontiers in Molecular Biosciences, 7*.
doi:10.3389/fmolb.2020.00157
Zhang, L., & Guo, H. (2020). Biomarkers of COVID-19 and technologies to combat SARS-CoV-2.
Advances in Biomarker Sciences and Technology, 2, 1-23.
doi:https://doi.org/10.1016/j.abst.2020.08.001
Zhang, S., Cooper-Knock, J., Weimer, A. K., Shi, M., Kozhaya, L., Unutmaz, D., . . . Snyder, M. P.
(2022). Multiomic analysis reveals cell-type-specific molecular determinants of COVID-19
severity. *Cell Systems*. doi:https://doi.org/10.1016/j.cels.2022.05.007

Zheng, H.-Y., Xu, M., Yang, C.-X., Tian, R.-R., Zhang, M., Li, J.-J., . . . Zheng, Y.-T. (2020). Longitudinal
957   transcriptome analyses show robust T cell immunity during recovery from COVID-19.
958   *Signal Transduction and Targeted Therapy, 5*(1), 294. doi:10.1038/s41392-020-00457-4
959 Zingoni, A., Molfetta, R., Fionda, C., Soriani, A., Paolini, R., Cippitelli, M., . . . Santoni, A. (2018).
960   NKG2D and Its Ligands: "One for All, All for One". *Frontiers in immunology, 9*.
961   doi:10.3389/fimmu.2018.00476
962 Zuo, Y., Yalavarthi, S., Shi, H., Gockman, K., Zuo, M., Madison, J. A., . . . Knight, J. S. (2020).
963   Neutrophil extracellular traps in COVID-19. *JCI Insight, 5*(11).
964   doi:10.1172/jci.insight.138999

Figure 1. Clinical profile and gene expression patterns for mild and severe COVID-19 patients during 28 days. **A.** Longitudinal sampling schedule for severe and mild COVID-19 donors of peripheral blood (22 samples in total from 8 donors). Three sampling times (black dots) are displayed with respect to the recruitment day (D0). In addition, panel A shows the diagnosis day with positive PCR (green star), symptoms onset day (orange star), and hospitalization day (red cross). **B.** The COVID-19 progression according to the WHO ordinal scale describes the temporal disease severity for each donor. Severe patients (S1-S4) are displayed in lines with colors scaling from green to red, while mild patients (M1-M4) are shown in blue line colors. The X-axis represents the days relative to the recruitment day. The vertical dot lines indicate the three sampling times used in this study (days 0, 7, and 28). **C.** Principal Component Analysis plot based on gene expression profiles for mild (blue) and severe (red) COVID-19 patients and grouped by their sampling time (0, 7, and 28 days after recruitment). In addition, peripheral blood samples from two healthy

donors are shown in green. PC = Principal Component. **D.** Heatmap of the 100 most

significant differentially expressed genes related to the COVID-19 disease

progression. At the bottom, each column corresponds to the sampling points (0, 7,

and 28 days since recruitment) of mild and severe patients. Genes are displayed as

horizontal rows and are clustered by the similarity of expression profiles, represented

by the dendrogram to the left of the heatmap. Red indicates higher expression, while

blue means lower expression represented by the z-score of normalized read counts.

Some COVID-19 severity-associated genes previously reported are indicated to the

right of the heatmap in red.

Figure 2. Gene Set Enrichment Analysis (GSEA) shows biological processes and

enriched pathways associated to innate and adaptive immune response after SARS-

CoV-2 infection. (**A** – **B**) Bubble plots show the biological processes (BP) of enriched

genes for DEGs between mild and severe for D0 and D7 after recruitment,

respectively. (**C** – **D**). Bubble plots show the KEGG enriched pathways of enriched

genes for DEGs between mild and severe for D0 and D7, respectively. The *Count*

(black circles) represents the number of genes included on each set. *Generatio* is

the ratio between numbers of genes found in the set and total genes of set. The

scale-color bar indicates the p-adjustment of each BP or KEGG pathway.

1000

Figure 3. Biological processes and KEGG pathway for genes with differential expression levels over time in mild versus severe patients. Bar plots show Gene ontology analysis for biological processes in light-blue/dark-blue and KEGG pathway in light-green/dark-green, using upregulated genes (**A** - **C** - **E**) and downregulated genes (**B** – **D** - **F**). The size of each bar is according to its -log10 p-value, and name pathways with asterisk indicates a q-value ≤ 0.05.

1007

Figure 4. Gene function network for Natural Killer cell hub-genes with differential expression levels between mild and severe patients during acute phase of COVID-19. (**A**) KEGG pathway of Natural Killer cell mediated cytotoxicity represents the set NK cell hub-genes upregulated (red-boxes) in mild versus severe patients. The green boxes correspond to genes without differential expression. (**B**) Heatmap shows the differential expression levels of NK cell hub-genes over time (D0, D7, and D28 after recruitment) separated by mild and severe groups. The expression levels are represented by the z-score of normalized counts. Dendrogram shows the hierarchical clustering of genes. (**C**) Protein-protein interaction (PPI) network for upregulated genes during the acute phase in mild patients. The network corresponds to the principal clusters with more interaction between proteins and highlight the three most represented pathways: Cytokine-cytokine receptor interaction (blue); NK cell mediated cytotoxicity (red); and Th1 and Th2 cell differentiation (yellow). (**D**)

1021 Time course expression levels for the main protein nodes identified in PPI network
1022 during the acute phase of COVID-19. The trajectories of these genes are graphed
1023 as days after recruitment (D0, D7, and D28) for mild (red triangle) and severe (green
1024 circle) groups and their enrichment is represented by the z-score of normalized
1025 counts.

Figure 5. Gene co-expression network analysis among the longitudinal transcriptomic profiling. **(A)** Gene hierarchical clustering dendrogram of ten detected modules based on Topological Overlap Matrices (TOM) measure. The branches and color bands represent the assigned module. **(B)** Module-trait relationships (MTRs)

between detected modules and disease severity of COVID-19. MTRs are obtained

by calculating Pearson correlation between the traits and the module eigengenes.

The red and blue colors indicate strong positive or negative correlations,

respectively. Rows represent module eigengene (ME) and columns indicate the

disease severity of COVID-19, where correlations from each patient are shown from

left to right according to their sampling time (D0, D7, and D28). **(C)** GO enrichment

analysis of genes in the blue, brown, and turquoise modules. The color in the bar

graphs refers to the module eigengene (ME). Enrichment results are sorted by -

log10(p-value) (higher on top) of each biological process GO term. **(D-E-F)** Network

visualization of enriched pathways based on co-expression genes from blue, brown,

and turquoise modules, respectively.

Table 1. Clinical characteristics of the cohort

CLINICAL CHARACTERISTICS	MILD	SEVERE	ALL
Sex, n (female/male)	(2/2)	(2/2)	(4/4)
Total, n	4	4	8
Median age, years ± SD	39.0 ± 3.9	46.7 ± 8	42.9 ± 7
Days from onset of symptoms to recruitment, median ± SD	1.2 ± 1.3	10.0±1.8	5 ± 4.7
Days from COVID-19 diagnosis to recruitment, median ± SD	3.0 ± 0.7	8.5 ± 2.1	6 ± 3.4
Symptoms, n (%)			
Fever	1 (25%)	3 (75%)	4 (50%)
Chills	1 (25%)	2 (50%)	3 (37.5%)
Fever feeling	1 (25%)	2 (50%)	3 (37.5%)
Odynophagia	2 (50%)	1 (25%)	3 (37.5%)
Cough	1 (25%)	4 (100%)	5 (62.5%)
Expectoration	-	-	-
Dyspnoea	-	4 (100%)	4 (50%)
Thoraxic pain	1 (25%)	1 (25%)	2 (25%)
Diarrhea	1 (25%)	2 (50%)	3 (37.5%)
Anosmia	1 (25%)	-	1 (12.5%)
Ageusia	1 (25%)	-	1 (12.5%)
Myalgia	1 (25%)	2 (50%)	3 (37.5%)
Headache	2 (50%)	1 (25%)	3 (37.5%)
Treatment			
Hospitalization, n (%)	-	4 (100%)	4 (50%)
Intubation, n (%)	-	4 (100%)	4 (50%)
Mechanical ventilation, n (%)	-	4 (100%)	4 (50%)
Samples, n	12	10	22