1 **"Two-Birds-One-Stone" Oral Nanotherapeutic Designed to Target Intestinal**

2 **Integrins and Regulate Redox Homeostasis for Ulcerative Colitis Treatment**

4 Long Huang[1,2], Wei Hu[2], Long Qun Huang[1], Qin Xuan Zhou[2], Zheng Yang Song[2],
Heng Yu Tao[2],

Bing Xu[3], Can Yang Zhang[2, *], Yi Wang[1, *], Xin-Hui Xing[1,2,3,4, *]

1. Institute of Biochemical Engineering, Department of Chemical Engineering, Tsinghua University,

Beijing 100084, China.

2. Institute of Biopharmaceutical and Health Engineering, Shenzhen International Graduate School,

Tsinghua University, Shenzhen 518055, China.

3. Institute of Biomedical Health Technology and Engineering, Shenzhen Bay Laboratory,

Shenzhen 518118, China.

4. Center for Synthetic and Systems Biology, Tsinghua University, Beijing 100084, China.

[†]Corresponding author: Xin-Hui Xing

Tel: 86-10-62794771, E-mail: xhxing@tsinghua.edu.cn

[†]Corresponding author: Yi Wang

Tel: 86-10-62773004, E-mail: wangyi671@mail.tsinghua.edu.cn

[†]Corresponding author: Can Yang Zhang

E-mail: zhang.cy@sz.tsinghua.edu.cn

Long Huang, E-mail: huanglon22@mails.tsinghua.edu.cn

Abstract

Designing highly efficient orally administered nanotherapeutics with specific inflammatory site-targeting functions in the gastrointestinal (GI) tract for ulcerative colitis (UC) management is a significant challenge. Straightforward and adaptable modular multifunctional nanotherapeutics represent groundbreaking advancements and are crucial to promoting broad application in both academic research and clinical practice. In this study, we focused on exploring a specific targeting modular and functional oral nanotherapy, serving as "one stone", for the directed localization of inflammation and the regulation of redox homeostasis, thereby achieving effects against "two birds" for UC treatment. The designed nanotherapeutic agent OPNs@LMWH, which has a core-shell structure composed of oxidation-sensitive ε-polylysine nanoparticles (OPNs) in the core and low-molecular-weight heparin (LMWH) in the shell, exhibited specific active targeting effects and therapeutic efficacy simultaneously. We qualitatively and quantificationally confirmed that OPNs@LMWH possessed high integrin αM-mediated immune cellular uptake efficiency and preferentially accumulated in inflamed lesions. Compared with bare OPNs, OPNs@LMWH exhibited enhanced intracellular reactive oxygen species (ROS) scavenging and anti-inflammatory effects. After oral administration of OPNs@LMWH to mice with dextran sulfate sodium (DSS)-induced colitis, robust resilience was observed. OPNs@LMWH effectively ameliorated oxidative stress and inhibited the activation of inflammation-associated signalling pathways while simultaneously bolstering the protective mechanisms of the colonic epithelium. Overall, these findings underscore the compelling dual functionalities of OPNs@LMWH, which enable effective oral delivery to inflamed sites, thereby facilitating precise UC management.

Introduction

The global incidence and prevalence of ulcerative colitis (UC), a chronic remitting-relapsing inflammatory disorder of the large intestine, have been increasing rapidly worldwide, yet this disease is incurable and has a multifaceted pathophysiology[1-3]. The European Crohn's and Colitis Organization (ECCO) and American Gastroenterological Association (AGA) recommend that patients follow a step-up approach with chemical drugs or monoclonal antibodies to achieve the desired remission because there is no known clinically available or effective cure for UC[4-6]. However, the frequent and long-term use of conventional small-molecule therapeutics or biologic-based immunosuppressive drugs can lead to severe complications, such as opportunistic infections, malignancies, autoimmunity, and liver toxicity[7-9]. Therefore, designing and developing effective therapeutics and treatment strategies for UC is very important, which may be achieved based on active component discovery and precise delivery by drawing inspiration from the evidence-based understanding of inflammatory signalling and drug delivery methods[10].

In prior research, precise drug delivery has been identified as a significant challenge[11,12]. The success of precise delivery is fundamentally dependent on the judicious choice of targeting strategy. A variety of strategies have been devised to target intestinal inflammation sites, such as microbial enzyme degradation, pH adjustment, and mucosal charge reversal at inflamed locations[13-17]. However, the complex and ever-changing environment of gastrointestinal diseases significantly undermines the reliability and precision of UC treatment outcomes[18]. Given the propensity of immune cells to accumulate at disease sites, the use of ligand

molecules to target immune cells seems more promising. Specific immune cells, such as
monocytes, innate lymphoid cells, and T cells, in addition to various associated regulatory
cytokines, have been extensively highlighted in the context of inflammatory bowel disease[19-
21]. Furthermore, studies have pinpointed the altered trafficking of immune cell circuits as key
drivers of mucosal inflammation and tissue destruction in UC[22], likening it to a severe "traffic
jam" at lesion sites. Consequently, integrins such as α4β7 and αMβ2, which have been
implicated in immune cell-related adhesion and signalling, could serve as excellent targets[23].
Recently, more studies have tended to focus on interference with integrin function[24], however,
few studies have explored functional integrin-targeting molecules suitable for the complex
gastrointestinal environment of UC. As a result, there is scant research on oral therapeutic
systems that can specifically target inflammatory cell infiltration via integrins.

When contemplating this issue, we thought of heparin. Heparin is an anionic polysaccharide
comprising highly sulfated disaccharide units. Low-molecular-weight heparin (LMWH),
prepared by the depolymerization of unfractionated heparin (UFH), has been recognized as
an attractive anticoagulant and has been successfully used in clinical settings for several
decades[25,26]. Intriguingly, due to the complex sequence of disaccharide units, where
anticoagulant fragments account for only 30 % of the polysaccharide[27], LMWH has been
demonstrated to have potential beyond its anticoagulant properties and has been reported
to be effective in the treatment of many other diseases, including infectious diseases, lung
cancer and Alzheimer's disease[27-29]. More importantly, heparin reportedly serves as an
adhesive ligand for integrins α4 and αM[23,30,31], known as CD49d and CD11b, respectively.
Inspired by these findings, we hypothesize that LMWH via oral administration could be a
promising inflammatory site-targeting ligand for developing precise oral therapy for UC.

Since conventional immunosuppressive drugs for UC management fail to fully address
treatment needs, it is crucial to develop groundbreaking strategies with multiple therapeutic
effects for rebalancing intestinal disorders in UC patients. Reactive oxygen species (ROS), as
integral factors affecting intestinal homeostasis, are signalling molecules extensively involved
in numerous cellular processes[32]. However, the disproportionate generation of ROS during
the UC pathogenesis process, which can lead to an increase by a factor of 10 to 100[33], inflicts
severe tissue damage on the intestinal epithelium and triggers the overproduction of
inflammatory mediators. While small-molecule drugs have been found to have unavoidable
limitations[34,35], there is mounting evidence that biocompatible polymer-based materials,
typically those based on glucose-related polysaccharides, show significant potential in
suppressing oxidative stress and inflammatory mediators, thereby regulating the redox
balance in the colon[9,36-40]. Unfortunately, the interactions of multiple hydroxy groups on the
glucose residues of polysaccharides introduce structural uncertainty. To design optimal ROS
scavenging materials with a defined structure and well-controlled function, alternative
biocompatible polymers need to be explored. For instance, ε-polylysine (EPL), a cationic and
biodegradable polypeptide produced by microbial fermentation, has garnered increased
interest in food chemistry and oral delivery systems[41]. The presence of a single amino group
on each repeating unit of EPL enables an available defined structure.

Here, we strongly emphasize the importance of creating precise nanotherapies that
integrate multiple functions. Therefore, we propose a modular orally administered
nanotherapeutic consisting of oxidation-sensitive ε-polylysine (OsEPL) nanoparticles (OPNs)

as the core, positively charged chitosan (CS) as the middle layer and negatively charged
LMWH as the outermost layer to achieve "two-birds-one-stone" treatment of UC.
Considering the adaptability and prospective engineering applications of these materials, we
utilized nanoprecipitation and layer-by-layer self-assembly methodologies in the fabrication
process. We hypothesize that orally administrated nanoparticles can not only target integrin-
associated immune cell recruitment and trafficking through the specific interaction between
LMWH and integrin αM but also regulate intestinal redox homeostasis through the ROS
scavenging function of OsEPL. Overall, we expect that our OPNs@LMWH may be translated
to topical UC therapy with high patient compliance *via* oral administration with high
therapeutic efficacy and few side effects. Moreover, the strategy of "two-birds-one-stone"
might be straightforward and effective for dealing with complex clinical problems.
123
Results
Preparing integrin-targeting OPNs@LMWH by nanoprecipitation and a layer-by-layer
process.
The oxidation-sensitive material (OsEPL) was designed and synthesized by introducing a
hydrophobic phenylboronic ester group that is sensitive to ROS into the side chain amino
group, as shown in Supplementary Scheme 1 and Supplementary Fig. 1. The structures of the
intermediate and OsEPL were confirmed by ^{1}H nuclear magnetic resonance (^{1}H-NMR)
(Supplementary Figs. 2-4), indicating successful synthesis. Additionally, we constructed a
series of OsEPLs with different feeding ratios of phenylboronic acid pinacol ester to amino
moiety. As the molar ratio of the hydrophobic side chain increased, the OsEPL gradually
formed an amphiphilic polymer in the aqueous phase. In addition, with a max feeding ratio,
OsEPL underwent rapid disintegration when incubated with peroxide (Supplementary Fig. 5).
Then, we obtained OPNs and OPNs@LMWH by nanoprecipitation and layer-by-layer
methods, as illustrated in Fig. 1a. To obtain nanoparticles coated with LMWH, the OPNs were
dispersed in CS buffer, which helped to form nanoparticles with a positive charge (OPNs@CS),
followed by incubation with an aqueous solution of LMWH to prepare OPNs@LMWH. The
OPNs@LMWH presented a spherical core-shell structure with an average particle size of
approximately 600 nm and a low polydispersity index (PDI; less than 0.2 on average)
according to scanning electron microscopy (SEM) (Fig. 1b), transmission electron microscopy
(TEM) (Fig. 1b, c) and dynamic light scattering (DLS) analyses (Fig. 1d). We also observed
charge reversal between the OPNs, OPNs@CS and OPNs@LMWH (Fig. 1e), which indicated
successful coating. Considering the goal of using this nanotherapeutic for oral administration,
its stability within the gastrointestinal tract is a matter of concern. The stability of
OPNs@LMWH was investigated by incubation with both dd-H$_2$O (4 °C) and simulated
gastric/intestinal fluids (SGF and SIF) (Supplementary Fig. 7a). Our results indicated that the
OPNs@LMWH had excellent stability both during storage (Supplementary Fig. 7b) and after
oral administration (Supplementary Fig. 7c-e).

Fig. 1 | **Preparation and characterization of OPNs@LMWH. a,** Schematic illustration of the preparation of OPNs@LMWH. **b,** Representative SEM and TEM images of OPNs and OPNs@LMWH. Scale bar, 200 nm. **c,** The elements C, B, and S in OPNs@LMWH. Scale bar, 200 nm. **d, e,** Particle size (**d**) and zeta potential (**e**) measurements of different nanoparticles determined by DLS. The data are presented as the means ± SDs. **f, g,** Schematic of the oral administration of OPNs@LMWH targeting immune cell recruitment and regulating intestinal redox homeostasis. OPNs@LMWH, when administered orally, can accumulate at the site of colitis inflammation (**f**). OPNs@LMWH efficiently regulate redox homeostasis by targeting the inflammation-related integrin (**g**).

Internalization by M1 macrophages and accumulation in inflamed colon tissue infiltrated by inflammatory cells.

After the preparation procedures were established, we next compared the ability of OPNs@LMWH and OPNs to be taken up by proinflammatory immune cells. To mimic the interactions between nanoparticles and proinflammatory immune cells at the lesion site,

RAW264.7 cells were activated by lipopolysaccharides (LPS) and IFN-γ to form M1
macrophages, which were subsequently incubated with fluorescein amine (FA)-labelled OPNs
(FA-OPNs) or FA-labelled OPNs@LMWH (FA-OPNs@LMWH). The samples were
subsequently imaged via confocal laser scanning microscopy (CLSM), as shown in Fig. 2a and
Supplementary Fig. 8. Compared with the OPN sample (untreated), the OPNs@LMWH sample
showed a much greater green fluorescence intensity, indicating significantly enhanced cellular
uptake efficiency. Then, we pretreated RAW264.7 cells with an antibody against the known
receptor integrin αM to interfere with the potential interaction between LMWH and integrin
αM. Consistent with our expectations, reduced attachment of OPNs@LMWH was observed
(Fig. 2a, b). For further illustration, we confirmed that free fluorescein was not taken up by
cells, while both OPNs and OPNs@LMWH were taken up after 8 h of incubation, which
suggested that the enhanced cellular uptake efficiency relied on the LMWH coating layer
(Supplementary Fig. 9).
Administering drinking water containing dextran sulfate sodium (DSS) is a well-recognized
method to construct a murine model that simulates human UC. We generated DSS-induced
colitis model mice to study the in vivo distribution of orally administered OPNs@LMWH. Since
we successfully confirmed that OPNs@LMWH targeted integrin αM in vitro, we subsequently
confirmed that integrin αM was enriched in colitis model mice because of immune cell
recruitment to the colonic submucosal layer (Supplementary Fig. 10). As expected, we
successfully observed high accumulation of OPNs@LMWH at the lesion site in vivo. After oral
administration of indocyanine green (ICG)-labelled OPNs@LMWH (ICG-OPNs@LMWH),
ICG-OPNs@LMWH presented a prolonged retention time in vivo in colitis model mice, and
the fluorescence signal peaked at 12 h (Fig. 2c, d). Compared with OPNs@LMWH, bare OPNs
were eliminated more rapidly by both healthy and colitis model mice (Fig. 2c, d). In the colitis
model mice administered with OPNs@LMWH, we also observed a significantly greater
particle accumulation in the colon, which was approximately 2-fold greater than that in colitis
model mice given indocyanine green-labelled OPNs (ICG-OPNs) and 12-fold greater than
that in healthy mice given ICG-OPNs or ICG-OPNs@LMWH (Fig. 2e, f). Notably, both OPNs
and OPNs@LMWH were eliminated from the colon in a short time in healthy mice, while
OPNs@LMWH were remained located in the inflamed colon extensively (Fig. 2e),
demonstrating the excellent inflammation-targeting capability of OPNs@LMWH. These
findings suggested that OPNs@LMWH possessed an inflammation-targeting effect in vivo.
To further observe the distribution of the particles in detail, frozen sections of colon tissue
were obtained 24 h after administering FA-OPNs or FA-OPNs@LMWH. We found that there
was almost no residual fluorescence signal from the particles in normal mouse colon tissue
and believed that the structurally intact intestinal barrier prevented the entry of the particles
and successfully excreted them (Fig. 2g, Supplementary Fig. 11). A small amount of OPNs
remained in the inflamed colon tissue. This could be attributed to nonspecific factors such as
disruption of the intestinal barrier and abnormal colon function. More significantly, a
substantial amount of OPNs@LMWH not only accumulated at the site of inflammation but
also penetrated deeper into the submucosal layer heavily infiltrated by integrin αM+
inflammatory cells (Fig. 2g, Supplementary Fig. 11), which suggested the excellent integrin
αM-targeting effect of LMWH.

Fig. 2 | OPNs@LMWH were internalized by M1 macrophages and accumulated in
inflamed colon tissue infiltrated by inflammatory cells. a, Activated M1-type RAW264.7
cells were incubated with FA-OPNs or FA-OPNs@LMWH after pretreatment with or without
the anti-integrin αM antibody, and visualized via CLSM. Scale bar, 50 μm. **b,** Quantitative
analysis of the FA fluorescence intensity in images taken by CLSM. **c, d,** Representative healthy
and colitis model mice were orally administered ICG-OPNs or ICG-OPNs@LMWH, imaged
by an in vivo imaging system (IVIS) and (**c**) the ICG fluorescence signal around the abdomen
was quantified (**d**). **e, f,** After 24 h, the colons were obtained and imaged with an IVIS (**e**), after
which the ICG fluorescence signal was quantified (**f**). **g,** Healthy or colitis model mice were
orally administered FA-OPNs or FA-OPNs@LMWH, and colon tissues were collected after 24
h and stained with DAPI, CM-Dil and integrin αM for visualization via CLSM. Scale bar, 50 μm.
Representative images or quantitative analysis of n=6 biologically independent samples (**a, b**)
and n=3 animals from two independent experiments (**c-g**) are shown. The data are presented

as the means ± SDs. * p < 0.05, ** p < 0.01, *** p < 0.001, **** p < 0.0001 and ns (not
significant) p > 0.05, analysed by two-way ANOVA (**b, d**) or ordinary one-way ANOVA (**f**) for
multiple comparisons.
227
Suppression of ROS and proinflammatory cytokines by OPNs@LMWH
Since the precise targeting properties of OPNs@LMWH have been illustrated, we next
evaluated the efficacy of OPNs@LMWH for immune regulation, including ROS scavenging
and anti-inflammatory effects. First, a 24 h cytotoxicity test was performed to evaluate cell
viability in vitro. The results indicated that OPNs@LMWH had the desired biocompatibility
and negligible cytotoxicity at a concentration of 100 μg mL^{-1} with both macrophages and
intestinal epithelial cells (Supplementary Fig. 12a-d). To investigate the ROS scavenging ability
of OPNs@LMWH, RAW264.7 cells were stimulated with phorbol 12-myristate 13-acetate
(PMA) after pretreatment with different nanoparticles. We observed significant
concentration-dependent intracellular ROS scavenging behaviour by both the OPNs and
OPNs@LMWH, while OPNs@LMWH exhibited greater scavenging capability due to their
enhanced cellular uptake efficiency (Fig. 3a-d, Supplementary Fig. 13a-c). In a confirmatory
experiment, we also found that LMWH had little anti-ROS activity (Supplementary Fig. 14a-
d) and that the mass proportion of LMWH in OPNs@LMWH did not exceed 1 %
(Supplementary Fig. 15), suggesting that the enhanced cellular uptake efficiency rather than
the potential oxidative sensitivity of LMWH resulted in enhanced intracellular ROS scavenging
behaviour. Then we evaluated the protective effects of the nanoparticles on peroxide-induced
apoptosis in NCM460 cells, which represents the extracellular ROS scavenging behaviour and
protection of the intestinal epithelium. We observed comparable results for OPNs and
OPNs@LMWH (Fig. 3e, f), suggesting comparable extracellular ROS scavenging behaviour.
Surprisingly, OPNs can also suppress the production of IL-6 and TNF-α, which are two
essential proinflammatory cytokines that are involved in most autoimmune diseases. This
effect was improved by the LMWH coating (Fig. 3g, h). Considering the mutually reinforcing
relationship between oxidative stress and inflammation, we attribute this improvement in
anti-inflammatory activity to increased cellular uptake and the consequent increase in
intracellular ROS scavenging efficiency, where LMWH plays an important role. These results
demonstrated that OPNs@LMWH are more effective at comprehensively reversing oxidative
stress and inflammation.

256
Fig. 3 | Suppression of ROS and proinflammatory cytokines by OPNs@LMWH. a-d,
RAW264.7 cells were stimulated with PMA and treated with OPNs or OPNs@LMWH (10 µg
mL^{-1}). The level of intracellular ROS was detected by DCFH-DA, visualized by fluorescence
microscopy (scale bar, 50 µm) (**a**) and determined by flow cytometry (**b**). The fluorescence
intensity (**c**) and percentage of ROS-positive cells were calculated (**d**). **e, f,** NCM460 cells were
incubated with H$_2$O$_2$ (500 µM) and treated with OPNs or OPNs@LMWH (10 µg mL^{-1}).
Peroxide-induced apoptosis was detected by staining with Annexin-V and propidium iodide
(PI) (**e**). The percentages of total apoptotic cells in the different treatment groups were
calculated (**f**). **g, h,** RAW264.7 cells were stimulated with LPS to produce proinflammatory
cytokines and incubated with OPNs or OPNs@LMWH (10 µg mL^{-1}). The concentrations of IL-
6 (**g**) and TNF-α (**h**) were determined via ELISA. Representative images, gating strategy and
quantitative analysis of n=4 biologically independent samples (**a-f**) and n=6 biologically
independent samples (**g, h**) are shown. The data are presented as the means ± SDs. $\star$ $p <$
0.05, $\star\star$ $p < 0.01$, $\star\star\star$ $p < 0.001$, $\star\star\star\star$ $p < 0.0001$ and ns $p > 0.05$, analysed by ordinary one-
way ANOVA (**c, d, f-h**) for multiple comparisons.

Therapeutic efficacy of OPNs@LMWH administered orally for UC treatment.

We next investigated the therapeutic efficacy of OPNs@LMWH in alleviating the
manifestations of colitis in mice. We first examined the in vivo safety of OPNs@LMWH to
confirm that a high daily dosage of OPNs@LMWH administered orally for two weeks did not
cause significant damage to the mice. Our results indicated that there was no evidence-based
damage in five main organs after constant daily oral administration at various concentrations,
which preliminary suggested the ideal biosafety of the formulation (Supplementary Figs. 16-
18). We subsequently investigated the therapeutic efficacy of OPNs and OPNs@LMWH in
colitis model mice. Fig. 4a schematically shows the development of the DSS-induced colitis
mouse model and the oral treatment strategies with OPNs or OPNs@LMWH. Compared with
water, both OPNs and OPNs@LMWH significantly protected mice against colitis, including
body weight loss (Fig. 4b), colon length shortening (Fig. 4c, e) and enlargement of the spleen
(Fig. 4d, f). Importantly, compared with that of bare OPNs, a significant improvement was
observed with OPNs@LMWH. According to the histopathological examination, treatment
with OPNs@LMWH also protected the colonic epithelium from pathological damage, and the
structures of the crypts were significantly more complete and more stable after OPNs@LMWH
treatment (Fig. 4g, h). These results indicated that OPNs@LMWH could effectively alleviate
disease status and that the LMWH coating could significantly improve the efficacy, which was
consistent with the results of in vitro experiments and our speculation.

Fig. 4 | Evaluation of the therapeutic efficacy of OPNs@LMWH in mice with colitis. a,
Schematic illustration of the development of the DSS-induced colitis mouse model. Mice were
administered 50 mg kg^{-1} OPNs or OPNs@LMWH daily or in drinking water. **b,** Bodyweight
changes in each group measured daily. **c-f,** Mice were sacrificed on day 7, and the colon (**c**)
and spleen (**d**) were collected. The colon length (**e**) and spleen index (**f**) were quantified. **g,**
Representative images of colons stained with haematoxylin and eosin. Scale bar, 100 μm. **h,**
The histological score was calculated to quantify tissue damage. Representative images or
quantitative analyses of n= 10 animals (**b-f**) and n=5 biologically independent samples (**g, h**)
are shown. The data are presented as the means ± SDs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$,
**** $p < 0.0001$ and ns $p > 0.05$, analysed by two-way ANOVA (**b**) or ordinary one-way
ANOVA (**e, f, h**) for multiple comparisons.

OPNs@LMWH restore intestinal homeostasis and protect the colonic epithelium.

Our results described above revealed the potential of OPNs@LMWH for restoring intestinal
redox homeostasis. Inspired by these data, we examined the mechanisms of OPNs@LMWH

in UC management. Our enzyme-linked immunosorbent assay (ELISA) results confirmed that
there were lower levels of proinflammatory cytokines in both the peripheral blood and lamina
propria of colitis model mice treated with OPNs@LMWH than in those treated with pure water
or OPNs (Fig. 5a-d). These results revealed the enhanced immune regulation ability of
OPNs@LMWH compared with OPNs. According to the myeloperoxidase (MPO)
immunohistochemistry (IHC) results, we observed the least oxidative stress in the colons of
the colitis model mice in the OPNs@LMWH treatment group compared with the other groups.
Remarkably, the decrease in the proportion of MPO-positive cells in the OPNs@LMWH
treatment group was primarily attributed to the significant reduction in the submucosal layer,
which is rich in recruited proinflammatory immune cells (Fig. 5e). These results indicated that
OPNs@LMWH precisely influenced the production of excess proinflammatory cytokines in
immune cells by scavenging ROS, which is consistent with our in vitro results. Moreover, the
IHC results showed decreased activation of the JAK-STAT signalling pathway in the lamina
propria after OPNs@LMWH treatment (Fig. 5f). As an essential signal transduction pathway,
JAK-STAT is activated mainly by proinflammatory cytokines in colitis lesions, hence,
decreasing activation of this pathway indicates a reduction in overall inflammatory activity.
An important sign of the restoration of intestinal homeostasis is the normalization of crypt
maintenance and self-renewal. The stability and rapid renewal of the intestinal mucosal layer
depend on the orderly arrangement of crypt cells and the performance of each functional cell,
including maintaining the epithelial barrier and regulating cell proliferation and apoptosis[42].
Interestingly, by immunofluorescence staining, we found that the epithelial cells were restored
and arranged more regularly in colitis model mice treated with OPNs@LMWH, which also
indicated that intestinal mucosal permeability was restored (Fig. 5g, h). We utilized cleaved
caspase 3 (CC3), a marker of apoptosis, to investigate the occurrence of apoptosis in colitis
model mice. Our findings revealed substantial overlap between cells exhibiting high
expression of CC3 and E-cadherin in untreated colitis model mice (Fig. 5h, i, k). Concurrently,
the proliferation (marked by PCNA) of transit amplifying (TA) cells and intestinal stem cells
located at the base of the crypts was virtually halted (Fig. 5j). This finding suggested that the
mucosal layer in colitis model mice had lost its barrier function and self-repair capabilities.
However, for colitis model mice treated with OPNs or OPNs@LMWH, a significant reduction
in the proportion of apoptotic epithelial cells was observed, demonstrating that the
proliferation of crypt cells was restored (Fig. 5i-k). More importantly, most of the PCNA-
positive cells were located at the bottom of the crypts after treatment with OPNs@LMWH.
These results indicated orderly proliferation and differentiation were ensured (Fig. 5j). Taken
together, these findings suggested that OPNs@LMWH can significantly ameliorate UC in vivo
by restoring intestinal homeostasis and protecting the colonic epithelium.

344
Fig. 5 | OPNs@LMWH restore intestinal homeostasis and protect the colonic epithelium.
a-d, Colitis model mice were treated with OPNs or OPNs@LMWH. The concentrations of IL-
6 in the serum (**a**) and IL-6 (**b**), TNF-α (**c**) and IL-1β (**d**) in colon tissue were determined via
ELISAs. **e, f,** Representative images of immunohistochemical staining for MPO (**e**) and p-
STAT3 (**f**). Scale bar, 100 μm. **g-k,** Representative images of immunofluorescence staining for
proliferation and apoptosis showing DAPI (**g**), E-cadherin (**h**), PCNA (**i**), CC3 (**j**), and merged
images (**k**) obtained via CLSM. Scale bar, 100 μm. Quantitative analysis or representative
images from n=5 biologically independent samples (**a-d**) and n= 3 biologically independent
samples (**e-k**) are shown. The data are presented as the means ± SDs. * $p < 0.05$, ** $p < 0.01$,
*** $p < 0.001$, **** $p < 0.0001$ and ns $p > 0.05$, analysed by ordinary one-way ANOVA (**a-d**)

for multiple comparisons.

Discussion

The intricacies of UC present significant challenges for existing treatment methods. Currently
available clinical drugs have notable limitations, and the off-target effects of oral medications
are concerning[8,43]. To address these issues, numerous studies on precision drug delivery
systems for UC have been conducted, yielding promising results[11,12]. Despite the varying
approaches and outcomes described in previous research, there is consensus that designing
new multifunctional targeted therapies for the immune response holds great potential[18].

In our work, we considered the transport and infiltration phenomena of inflammatory cells
at UC lesion sites and proposed a "two-birds-one-stone" precise nanotherapy based on
integrin αM targeting and ROS scavenging. By adopting a modular integration approach,
OsEPL was combined with LMWH through nanoprecipitation and layer-by-layer self-
assembly to form core-shell nanoparticles. Given the imbalance in redox homeostasis at
inflammatory sites, we opted for a drug-free approach using biocompatible ROS scavenging
materials derived from modified EPL as a regulator of intestinal oxidative stress.

Despite the discovery of the heparin–integrin interaction mechanism in previous research,
our knowledge regarding its potential in drug delivery remains sparse. Our findings robustly
validated the enhanced immune cellular uptake of heparin-coated nanoparticles and
highlighted the crucial mediating role of LMWH-integrin αM interactions. OPNs@LMWH also
demonstrated a significantly extended retention time in the colon of colitis model mice.
Remarkably, 24 h after oral administration, the concentration of OPNs@LMWH was
approximately double that of the bare OPNs. Conversely, normal mice rapidly eliminated
OPNs@LMWH, underscoring its superior inflammation-targeting properties. A closer
examination of colon tissue sections revealed that OPNs@LMWH penetrated more deeply
into submucosal tissues with inflammatory cell infiltration than bare OPNs did. As anticipated,
OPNs@LMWH successfully targeted intestinal inflammatory cell migration, accumulation, and
infiltration in colitis model mice.

To understand how OPNs@LMWH improve colitis, we conducted comprehensive in vivo
and in vitro experiments. In our in vitro studies, we noted a reduction in the levels of
intracellular ROS and in the secretion of proinflammatory cytokines following the application
of OPNs@LMWH. Interestingly, in experiments involving cells damaged by exogenous
peroxides, the performance of OPNs@LMWH was comparable to that of OPNs. These
findings suggest that while the cellular uptake efficiency of OPNs@LMWH is better than that
of OPNs, their inherent antioxidant functionality remains effective. In subsequent in vivo trials,
OPNs@LMWH demonstrated superior effectiveness in alleviating the symptoms of colitis. Our
results revealed that following treatment with OPNs@LMWH, there were significant
reductions in the levels of proinflammatory cytokines in peripheral blood and colonic lamina
propria in mice with colitis. Additionally, we noted a marked decrease in oxidative stress within
immune cells and reduced activation of the JAK-STAT signalling pathway. The process of
apoptosis in the intestinal epithelial cells of the mice was effectively suppressed, and their
orderly proliferation at the base of the crypts was successfully reinstated.

Our findings provide robust validation for our "two-birds-one-stone" strategy that
combines LMWH-based integrin targeting with the regulation of intestinal redox homeostasis,

thereby highlighting the application of OPNs@LMWH in the treatment of UC. More
promisingly, our research underscores the feasibility and flexibility of employing LMWH via
oral administration as an integrin-targeted treatment for UC. We postulate that the LMWH
surface coating could be universally implemented across a variety of nanoparticle designs
targeting integrins αM. This straightforward and adaptable modular design could stimulate
further research into integrin-targeted treatment strategies for autoimmune diseases.

Experimental section
Materials:
ε-Polylysine (EPL), 4-hydroxyphenyl boronic acid pinacol ester, dichloromethane (DCM),
carbonyl diimidazole (CDI), sodium chloride (NaCl), sodium sulfate (Na$_2$SO$_4$), 4-
dimethylaminopyridine (DMAP), dimethyl sulfoxide (DMSO), lecithin, ammonium acetate
(CH$_3$COONH$_4$) and hydrogen peroxide solution (H$_2$O$_2$) were purchased from Aladdin
(Shanghai, CHN). Heparin was purchased from Changshan Biotechnology Co., Ltd. (Hebei,
CHN). Chitosan (CS, >75 % deacetylated), indocyanine green (ICG), fluorescein amine (FA),
azure A, lipopolysaccharide (LPS), phorbol 12-myristate 13-acetate (PMA), sucrose and
haematoxylin solution were purchased from Sigma–Aldrich (St. Louis, MO, USA). Cell
Counting Kit-8 (CCK-8), a BCA protein assay kit, protease and phosphatase inhibitor cocktail
for mammalian cell and tissue extracts (P1050) and DCFH-DA were purchased from Beyotime
Biotechnology (Shanghai, CHN). Foetal bovine serum (FBS) was purchased from Gemini (USA).
Dulbecco's modified Eagle's medium (DMEM), Roswell Park Memorial Institute (RPMI) 1640
medium, and Opti-MEM I reduced serum medium were purchased from Gibco (USA). Dextran
sulfate sodium (DSS, colitis grade) was purchased from MP Biomedicals (USA). The anti-E-
cadherin antibody (14-3249-82), CM-Dil (C7001), and the ELISA kits for IL-6 (88-7064), TNF-
α (88-7324), and IL-1β (88-7013) were purchased from Invitrogen (USA). The Annexin-V/PI
kit was purchased from BioLegend, Inc. (USA). The antibodies against phospho-STAT3
(Tyr705, 9145), cleaved caspase-3 (Asp175, 9579) and PCNA (2586) were purchased from Cell
Signaling Technology (USA). The antibodies against myeloperoxidase (ab208670) and integrin
αM (ab133357) were purchased from Abcam (Cambridge, UK). DAPI was purchased from
Thermo Scientific (USA). Secondary antibodies conjugated with Alexa Fluor 488 (A-21208),
Alexa Fluor 594 (A-21207), and Alexa Fluor 647 (A-31573 and A-31571) were purchased from
Invitrogen (USA). Secondary antibodies conjugated with horseradish peroxidase (HRP), 3,3'-
diaminobenzidine (DAB), peroxidase blocking solution and serum-free protein blocking
solution were purchased from Dako (DNK). The O.C.T. compound was purchased from Sakura
(USA).

Synthesis of OsEPL: Oxidation-sensitive ε-polylysine (OsEPL) was synthesized inspired by
Broaders et al.[44]. Briefly, 4-hydroxyphenyl boronic acid pinacol ester (10 mmol) was dissolved
in DCM (10 mL). Then, CDI (20 mmol) was added, and the mixture was stirred at RT for 1 h.
The solution was diluted by adding another 10 mL of DCM and washed with dd-H$_2$O (20 mL)
three times. The obtained organic phase was subsequently washed with saturated brine (20
mL) three times, dried over anhydrous Na$_2$SO$_4$ for 30 min and concentrated in a rotary
evaporator to obtain imidazoyl carbamate (Compound 1, 8.5 mM, 85 %). To generate OsEPL,
Compound 1 (820 mg), EPL (320 mg) and DMAP (305 mg) were dissolved in anhydrous DMSO

(10 mL). The mixture was stirred at RT for 24 h. Then, OsEPL was precipitated in dd-H₂O (40

mL) and isolated by centrifugation (8000 × g, 10 min). The precipitate was washed with dd-

H₂O (20 mL) three times and lyophilized (72 h) for further use.

Preparation of OPNs@LMWH: LMWH (Mw=8.5 kDa) was prepared by a controlled

heparinase depolymerization method according to previous methods.[26,45] The molecular

weights of unfractionated heparin (UFH) and LMWH were determined by high-performance

liquid chromatography (HPLC, 1260 Infinity II, Agilent, USA) with a GPC column (TSK Gel

G2000SWXL, 7.8 mm × 300 mm, Tosoh Co., JPN), and the results are shown in Supplementary

Fig. 6. OPNs@LMWH were prepared by a nanoprecipitation/layer-by-layer method. Briefly,

lecithin (20 mg) was dispersed in 1 mL of anhydrous ethanol and subsequently added

dropwise to 8 mL of CH₃COONH₄ buffer (0.1 M, pH = 5.00) with magnetic stirring (600 rpm)

to form a dispersion. OsEPL (20 mg) was dissolved in DMSO (1 mL) and subsequently added

dropwise to the dispersion. The nanoparticles were collected by centrifugation (30000 × g,

10 min). Bare OPNs were harvested after washing three times with dd-H₂O. CS (20 mg) was

dissolved in 10 mL of CH₃COONH₄ buffer (0.1 M, pH = 5.00). Then, the bare OPNs were

dispersed in the CS solution and incubated at RT for 30 min with magnetic stirring (600 rpm).

The CS-coated OPNs were collected after washing three times with dd-H₂O, dispersed in 10

mL of LMWH aqueous solution (2 mg mL^{-1}) and incubated at RT for 30 min with magnetic

stirring (600 rpm) again. Finally, the OPNs@LMWH were washed three additional times and

collected by centrifugation (30000 × g, 10 min) for further use. For in vivo imaging, ICG (1 mg

mL^{-1}) or FA (1 mg mL^{-1}) was added to the primary DMSO solution containing OsEPL, and two

kinds of fluorescently labelled OPNs or OPNs@LMWH were prepared via a method same to

that described above. The fluorescence intensity of the fluorescently labelled nanoparticles

was normalized before use.

Characterization of OPNs@LMWH: The surface morphologies of the OPNs and

OPNs@LMWH were determined via scanning electron microscopy (SEM) (Merlin, Zeiss, DEU),

transmission electron microscopy (TEM) (Tecnai G2 20, FEI, USA) and aberration-corrected

transmission electron microscopy (AC-TEM) (Titan 80-300, FEI, USA). The sizes, polydispersity

index (PDI) and surface charges of the OPNs and OPNs@LMWH were analysed via dynamic

light scattering (DLS) (Zetasizer Nano ZS90, Malvern, UK). For stability assays, simulated

gastric fluid (SGF) and simulated intestinal fluid (SIF) were prepared according to the methods

of the United States Pharmacopeia. For stability assay, the nanoparticles were dispersed and

incubated in deionized water (48 h) at 4 °C or in SGF (2 h) or SIF (44 h) at 37 °C. Degradation

was evaluated by measuring the absorbance at 500 nm with a microplate reader (Infinite

M200 Pro, Tecan Trading AG, Switzerland). The mass proportion of LMWH in OPNs@LMWH

was calculated by measuring the concentration of free LMWH in the supernatant before and

after coating. The concentration of LMWH was detected by incubation with Azure A.

Cell culture: RAW264.7 (mouse) macrophages and NCM460 (human) colonic enterocytes

were kindly provided by Prof. Zhijie Chang (School of Medicine, Tsinghua University).

RAW264.7 cells were cultured in DMEM supplemented with 10 % FBS, and NCM460 cells were

cultured in RPMI 1640 medium supplemented with 10 % FBS. For in vitro assays, the medium

was replaced with Opti-MEM I reduced serum medium after the cells were seeded overnight
to avoid potential impacts from FBS.
489
In vitro cellular uptake assays: RAW264.7 cells were seeded in 12-well plates (Corning 3513,
USA) at a density of 3×10^5 cells per well and stimulated with LPS (100 ng mL^{-1}) and IFN-γ
(20 ng mL^{-1}) for 12 h to form M1 macrophages. Then, FA-labelled nanoparticles (20 μg mL^{-1})
were added. After 8 h of incubation, the cells were collected and washed with PBS three times.
The fluorescence intensity of the cells was immediately assessed via flow cytometry
(LSRFortessa, BD, USA). To confirm the improved cellular uptake efficiency, free FA with the
same fluorescence intensity as the nanoparticles was used as a control group. For
immunofluorescence (IF) imaging, cells were seeded on round coverslips and pretreated with
anti-integrin αM antibodies for 1 h before 8 h of incubation with the nanoparticles. Cells on
coverslips were fixed in 4 % paraformaldehyde for 15 min, stained with CM-Dil and DAPI and
washed three times. Confocal laser scanning microscopy (CLSM; FV3000RS, Olympus, JPN)
was used to obtain images.
502
In vitro ROS scavenging behaviour and anti-inflammatory effects: Intracellular ROS levels
were evaluated via the DCFH-DA method. Briefly, RAW264.7 cells were seeded in 12-well
plates (Corning 3513, USA) at a density of 3×10^5 cells per well. After incubation for 12 h, the
medium was replaced with Opti-MEM containing nanoparticles at a predetermined
concentration, and the mixture was incubated for another 12 h. Then, PMA (100 ng mL^{-1}) and
DCFH-DA were added. After incubation for 1 h, the cells were collected and washed with PBS
three times. The fluorescence intensity of the cells was immediately assessed by flow
cytometry. To confirm the ROS scavenging function of the nanoparticles, we evaluated the
reduction in peroxide-induced apoptosis via the Annexin-V/PI method. Briefly, NCM460 cells
were seeded in 12-well plates (Corning 3513, USA) at a density of 3×10^5 cells per well. After
incubation for 12 h, the medium was replaced with Opti-MEM containing nanoparticles (10
μg mL^{-1}), and the cells were incubated for 1 h. Then, H$_2$O$_2$ (500 μM) was added, and the plate
was incubated for 12 h. The cells were collected, stained with an Annexin-V/PI apoptosis kit
and assessed immediately by flow cytometry. For the in vitro anti-inflammatory assay,
RAW264.7 cells were seeded in 96-well plates (Corning 3599, USA) at a density of 3×10^4
cells per well. After incubation for 12 h, the medium was replaced with Opti-MEM containing
LPS (100 ng mL^{-1}). After 30 min of stimulation with LPS, nanoparticles at a predetermined
concentration (10 μg mL^{-1}) were added, and the cells were incubated for an additional 24 h.
The supernatant was collected to determine the concentration of proinflammatory cytokines
by ELISA kits.
523
Animals and colitis model: All animal studies complied with the regulations of the Laboratory
Animal Resources Center, Tsinghua University (23-XXH2). Six-week-old male C57BL/6J mice
(Specific Pathogen Free, SPF) were obtained from Charles River (Beijing, CHN) and maintained
on a 12-h light–dark cycle at 25 °C in SPF barrier environment. Mice supplied with food and
water were cohoused for 7 days before random assignment to the experimental groups. A
colitis model was established by providing drinking water containing 3 % DSS, and this
drinking water was renewed every other day.

In vivo therapeutic efficacy: Mice were orally administered drinking water or nanoparticle dispersions at a dose of 50 mg kg^{-1} every day for 7 days (beginning on day 0 and ending on day 6). On day 7, the mice were sacrificed, and the spleens, colons and serum were excised. To obtain histological sections, colons were fixed in 4 % paraformaldehyde for 24 h, dehydrated and embedded in paraffin. The samples were cut into 5 μm thick sections by using a microtome (RM2235, Leica, DEU) and stored at -20 °C until further use. Haematoxylin and eosin staining was performed for histopathological examination. The histological scores were determined according to a previously reported method[46]. Briefly, infiltration of inflammatory cells (0: none, 1: mild, 2: moderate, 3: severe), depth of ulceration involvement (0: none, 1: mucosa, 2: submucosa, 3: muscular layer), area of ulceration involvement (0: none, 1: 0 %–25 %, 2: 26 %–50 %, 3: 51 %–75 %, 4: 76 %–100 %), and crypt destruction (0: none, 1: lower than one third, 2: lower than two thirds, 3: more than two thirds, 4: all crypts and epithelium) were assessed. To determine the concentrations of cytokines in colon tissues, colon segments were homogenized (tissue homogenizer, 30 Hz, 2 min) in PBS containing 2 % protease and phosphatase inhibitor cocktail. The supernatants were collected, and cytokine levels were assessed via ELISA kits. All samples were normalized to equivalent protein concentrations by using a BCA protein assay kit.

Immunohistochemistry and immunofluorescence imaging of histological sections: Immunohistochemistry (IHC) and IF were used to investigate the underlying mechanisms. Briefly, histological sections were obtained by the same method as described for "In vivo therapeutic efficacy", followed by deparaffinization and hydration of tissue sections through xylene and gradient ethanol treatment, respectively. Antigen retrieval was performed at 100 °C for 15 min in Tris-EDTA buffer (pH = 9.00). For IHC, a peroxidase blocking solution was applied to eliminate nonspecific staining, while for IF, a serum-free protein blocking solution was applied. The tissue slides were incubated overnight at 4 °C with primary antibodies. All the slides were washed with PBS (containing 0.05 % Tween-20) and incubated with secondary antibodies for 30 min at RT, where antibodies conjugated with HRP were used for IHC and antibodies conjugated with fluorophores were used for IF. For IHC, DAB was applied to develop the colour, followed by haematoxylin staining. For IF, nuclear staining was performed by using DAPI. Then, all the slides were mounted with mounting medium (Leica), protected from light and stored at 4 °C.

In vivo biodistribution: Both healthy mice and mice with colitis were orally administered fluorescently labelled OPNs/OPNs@LMWH. Briefly, after fasting overnight, mice orally administered ICG-OPNs/OPNs@LMWH were imaged by an IVIS (Lumina III, USA) at 6 h, 12 h and 24 h after gavage and then sacrificed for colon imaging. To investigate the specific distribution of the labelled OPNs/OPNs@LMWH in colon tissue, mice orally administered FA-OPNs/OPNs@LMWH were sacrificed 24 h after gavage. The colons were obtained and stored in PBS containing 30 % sucrose for 24 h. Then, the solution was removed, and all the samples were embedded in O.C.T. compound. Histological sections were obtained by using a freezing microtome (CM3050S, Leica, DEU). Slides were stained with DAPI, CM-Dil and integrin-αM, mounted, protected from light and stored at 4 °C.

Safety assays: In vitro safety assays utilized cells. Briefly, cells were cultured in 96-well plates with different concentrations of particles for 24 h. Then, CCK-8 reagent was added, and the plates were incubated for another 30 min. The absorbance of each well at 450 nm was measured, and this value was used to calculate cell viability. In vivo safety assays were based on mouse data. Briefly, mice were orally administered different doses of nanoparticles daily for 14 days, and their body weights were recorded every day. Then, all the mice were sacrificed on day 14. Blood samples were collected and used for blood haematology examinations. The spleens, hearts, lungs, livers and kidneys of the mice were collected to obtain sections that were stained with haematoxylin and eosin.

Statistical analysis: The data are expressed as the mean ± standard deviation (SD). Statistical analysis was performed by student's t tests for unpaired data and by one-way ANOVA or two-way ANOVA for multiple comparisons. A two-tailed p value of < 0.05 was considered to indicate statistical significance, denoted as * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ and ns (not significant) for $p > 0.05$. GraphPad Prism 8.3 software was used for all statistical analyses.

References

1. Kobayashi, T. *et al.* Ulcerative colitis. *Nat. Rev. Dis. Primers.* **6**, 74 (2020).

2. Ungaro, R., Mehandru, S., Allen, P. B., Peyrin-Biroulet, L. & Colombel, J. F. Ulcerative colitis. *Lancet* **389**, 1756-1770 (2017).

3. Kaplan, G. G. & Ng, S. C. Understanding and preventing the global increase of inflammatory bowel disease. *Gastroenterology* **152**, 313-321 .e312 (2017).

4. Raine, T. *et al.* ECCO Guidelines on Therapeutics in Ulcerative Colitis: Medical Treatment. *J. Crohns Colitis* **16**, 2-17 (2022).

5. Ko, C. W. *et al.* AGA Clinical Practice Guidelines on the Management of Mild-to-Moderate Ulcerative Colitis. *Gastroenterology* **156**, 748-764 (2019).

6. Feuerstein, J. D. *et al.* AGA Clinical Practice Guidelines on the Management of Moderate to Severe Ulcerative Colitis. *Gastroenterology* **158**, 1450-1461 (2020).

7. Bernstein, C. N. *et al.* World gastroenterology organization practice guidelines for the diagnosis and management of IBD in 2010. *Inflamm. Bowel. Dis.* **16**, 112-124 (2010).

8. Matsuoka, K. *et al.* Evidence-based clinical practice guidelines for inflammatory bowel disease. *J. Gastroenterol.* **53**, 305-353 (2018).

9. Shi, C. *et al.* A nanoparticulate dual scavenger for targeted therapy of inflammatory bowel disease. *Sci. Adv.* **8**, eabj2372 (2022).

10. Tabas, I. & Glass, C. K. Anti-Inflammatory therapy in chronic disease: challenges and opportunities. *Science* **339**, 166-172 (2013).

11. Awad, A. *et al.* Clinical translation of advanced colonic drug delivery technologies. *Adv. Drug Deliv. Rev.* **181**, 114076 (2022).

12. Duran-Lobato, M., Niu, Z. & Alonso, M. J. Oral delivery of biologics for precision medicine. *Adv. Mater.* **32**, 1901935 (2020).

13. Bertoni, S. *et al.* pH and reactive oxygen species-sequential responsive nano-in-micro composite for targeted therapy of inflammatory bowel disease. *Adv. Funct. Mater.* **28**, 1806175 (2018).

14. Ding, Y. F. *et al.* Oral colon-targeted konjac glucomannan hydrogel constructed through noncovalent cross-linking by cucurbit 8 uril for ulcerative colitis therapy. *ACS Appl. Bio Mater.* **3**, 10-19 (2020).

15. Neurath, M. F. & Travis, S. P. L. Mucosal healing in inflammatory bowel diseases: a systematic review. *Gut* **61**, 1619-1635 (2012).

16. Nugent, S. G., Kumar, D., Rampton, D. S. & Evans, D. F. Intestinal luminal pH in inflammatory bowel disease: possible determinants and implications for therapy with aminosalicylates and other drugs. *Gut* **48**, 571-577 (2001).

17. Zhang, S. F. *et al.* An inflammation-targeting hydrogel for local drug delivery in inflammatory bowel disease. *Sci. Transl. Med.* **7**, 300ra128-300ra128 (2015).

18. Kotla, N. G. *et al.* Bioresponsive drug delivery systems in intestinal inflammation: State-of-the-art and future perspectives. *Adv. Drug Deliv. Rev.* **146**, 248-266, doi:10.1016/j.addr.2018.06.021 (2019).

19. Neurath, M. F. Cytokines in inflammatory bowel disease. *Nat. Rev. Immunol.* **14**, 329-342 (2014).

20. Biton, M. *et al.* T helper cell cytokines modulate intestinal stem cell renewal and differentiation. *Cell* **175**, 1307-1320 e1322 (2018).

21   Schwartz, D. M. *et al.* JAK inhibition as a therapeutic strategy for immune and
inflammatory diseases. *Nat. Rev. Drug Discov.* **16**, 843-862 (2017).
22   Neurath, M. F. Targeting immune cell circuits and trafficking in inflammatory bowel
disease. *Nat. Immunol.* **20**, 970-979 (2019).
23   Ley, K., Rivera-Nieves, J., Sandborn, W. J. & Shattil, S. Integrin-based therapeutics:
biological basis, clinical use and new drugs. *Nat. Rev. Drug Discov.* **15**, 173-183 (2016).
24   Mitroulis, I. *et al.* Leukocyte integrins: role in leukocyte recruitment and as therapeutic
targets in inflammatory disease. *Pharmacol. Therapeut.* **147**, 123-135 (2015).
25   Mulloy, B., Hogwood, J., Gray, E., Lever, R. & Page, C. P. Pharmacology of heparin and
related drugs. *Pharmacol. Rev.* **68**, 76-141 (2016).
26   Ji, Y. *et al.* A heparin derivatives library constructed by chemical modification and
enzymatic depolymerization for exploitation of non-anticoagulant functions. *Carbohydr.*
Polym. **249**, 116824 (2020).
27   Beurskens, D. M. H. *et al.* The anticoagulant and nonanticoagulant properties of heparin.
Thromb. Haemost. **120**, 1371-1383 (2020).
28   Wang, P. *et al.* Heparin: an old drug for new clinical applications. *Carbohydr. Polym.* **295**,
119818 (2022).
29   Yan, Y. *et al.* Mucoadhesive nanoparticles-based oral drug delivery systems enhance
ameliorative effects of low molecular weight heparin on experimental colitis. *Carbohydr.*
Polym. **246**, 116660 (2020).
30   Tyler, C. J. *et al.* Antibody secreting cells are critically dependent on integrin
α4β7/MAdCAM-1 for intestinal recruitment and control of the microbiota during chronic
colitis. *Mucosal Immunol.* **15**, 109-119 (2022).
31   Diamond, M. S., Alon, R., Parkos, C. A., Quinn, M. T. & Springer, T. A. Heparin is an adhesive
ligand for the leukocyte Mac-1(CD11b, CD18). *J. Cell Biol.* **130**, 1473-1482 (1995).
32   Bhattacharyya, A., Chattopadhyay, R., Mitra, S. & Crowe, S. E. Oxidative stress: an essential
factor in the pathogenesis of gastrointestinal mucosal diseases. *Physiol. Rev.* **94**, 329-354
(2014).
33   Wilson, D. S. *et al.* Orally delivered thioketal nanoparticles loaded with TNF-α-siRNA
target inflammation and inhibit gene expression in the intestines. *Nat. Mater.* **9**, 923-928
(2010).
34   Stallmach, A., Hagel, S. & Bruns, T. Adverse effects of biologics used for treating IBD. *Best*
Pract. Res. Clin. Gastroenterol. **24**, 167-182 (2010).
35   Lee, Y. *et al.* Hyaluronic acid-bilirubin nanomedicine for targeted modulation of
dysregulated intestinal barrier, microbiome and immune responses in colitis. *Nat. Mater.*
19, 118-126 (2020).
36   Lee, S., Stubelius, A., Hamelmann, N., Tran, V. & Almutairi, A. Inflammation-responsive
drug-conjugated dextran nanoparticles enhance anti-inflammatory drug efficacy. *ACS*
Appl. Mater. Interfaces **10**, 40378-40387 (2018).
37   Stubelius, A., Lee, S. & Almutairi, A. The chemistry of boronic acids in nanomaterials for
drug delivery. *Acc. Chem. Res.* **52**, 3108-3119 (2019).
38   Nathan, C. & Cunningham-Bussel, A. Beyond oxidative stress: an immunologist's guide to
reactive oxygen species. *Nat. Rev. Immunol.* **13**, 349-361 (2013).
39   Sies, H. *et al.* Defining roles of specific reactive oxygen species (ROS) in cell biology and

physiology. *Nat. Rev. Mol. Cell Biol.* **23**, 499-515 (2022).

40 Gao, C. *et al.* Treatment of atherosclerosis by macrophage-biomimetic nanoparticles via
targeted pharmacotherapy and sequestration of proinflammatory cytokines. *Nat.*
Commun. **11**, 2622 (2020).

41 Du, X. *et al.* Polylysine and cysteine functionalized chitosan nanoparticle as an efficient
platform for oral delivery of paclitaxel. *Carbohydr. Polym.* **229**, 115484 (2020).

42 Yen, T. H. & Wright, N. A. The gastrointestinal tract stem cell niche. *Stem Cell Rev. Rep.* **2**,
203-212 (2006).

43 Hindryckx, P., Jairath, V. & D'Haens, G. Acute severe ulcerative colitis: from
pathophysiology to clinical management. *Nat. Rev. Gastroenterol. Hepatol.* **13**, 654-664,
doi:10.1038/nrgastro.2016.116 (2016).

44 Broaders, K. E., Grandhe, S. & Frechet, J. M. A biocompatible oxidation-triggered carrier
polymer with potential in therapeutics. *J. Am. Chem. Soc.* **133**, 756-758 (2011).

45 Wu, J., Zhang, C., Mei, X., Li, Y. & Xing, X. H. Controllable production of low molecular
weight heparins by combinations of heparinase I/II/III. *Carbohydr. Polym.* **101**, 484-492
(2014).

46 Xu, J. *et al.* Probiotic-inspired nanomedicine restores intestinal homeostasis in colitis by
regulating redox balance, immune responses, and the gut microbiome. *Adv. Mater.* **35**,
e2207890 (2023).

Data availability
All the data supporting the results of this study are available within the paper and its
Supplementary Information. Additional processed data are available from the corresponding
authors upon request.

Acknowledgements
The authors thank Prof. Zhibing Zhang, FREng, University of Birmingham, for his invaluable
advice in the preparation of manuscript. The authors thank Prof. Zhijie Chang, Tsinghua
University, for generously providing the cell lines. We thank the National Center for Electron
Microscopy in Beijing, School of Materials Science, Tsinghua University, for assistance with
electron microscopy imaging; the Laboratory Animal Resources Center, Tsinghua University,
for technical support; and the Center of Biomedical Analysis, Tsinghua University, for
assistance with imaging technology and flow cytometry. This work was supported by grants
from the Key Research and Development Program of the Ministry of Science and Technology
(2023YFA0914300), the Shenzhen Medical Research Fund (SMRF, B2302009) and the National
Natural Science Foundation of China (22161142014).

Author contributions
XH. X. Y. W. and CY. Z. oversaw the research. L. H., B. X., Y. W., CY. Z. and XH. X. designed the
research strategy. L. H. synthesized the materials and fabricated the nanotherapy. L. H., LQ.
H., W. H., QX. Z., ZY. S., and HY. T. performed the experiments. L. H. and Y. W. analysed the
experimental data. L. H., Y. W., CY. Z. and XH. X. prepared the manuscript. All the authors
approved the manuscript.

Competing interests

XH. X., L. H. and Y. W. are inventors on related patents filed by Tsinghua University describing the materials and nanotherapy reported in this study. The other authors declare no competing interests.

730   **Supplementary Information**

733   **"Two-Birds-One-Stone" Oral Nanotherapeutic Designed to**

734   **Target Intestinal Integrins and Regulate Redox Homeostasis for**

735   **Ulcerative Colitis Treatment**

Contents

765
766   **Supplementary Scheme 1 | Synthesis of OsEPL.**

767
768

769
770   **Supplementary Fig. 1 | Schematic illustration of the preparation and degradation of**
771   **OsEPL.**
772

773
774 **Supplementary Fig. 2 | ^{1}H-NMR spectrum of imidazoyl carbamate (Compound 1) in**
775 **DMSO-d₆.**
776
777

778
779 **Supplementary Fig. 3 | ^{1}H-NMR spectrum of EPL in DMSO-d₆.**
780

Supplementary Fig. 4 | ^{1}H-NMR spectrum of OsEPL in DMSO-d_6.

Supplementary Fig. 5 | Physicochemical properties of OsEPL. a, A series of OsEPLs with
different feeding ratios of phenylboronic acid pinacol ester and amino moieties presented
varying amphiphilicities. **B,** OsEPL was incubated with 100 mM H_2O_2 for one hour.

Supplementary Fig. 6 | Molecular weight distributions of UFH and LMWH determined by
GPC.

Supplementary Fig. 7 | Stability of OPNs and OPNs@LMWH. a, OPNs and OPNs@LMWH
were incubated under storage conditions (deionized water, 4 °C) or subjected to a simulated
digestion process (SGF, 2 h, 37 °C; SIF, 46 h, 37 °C). **b,** The ratio of hydrolysed nanoparticles
after storage. The data shown are from quantitative analyses of n=3 biologically independent
samples. **c-e,** Representative SEM images (scale bar, 200 nm) (**c**), particle sizes (**d**) and zeta
potentials (**e**) of OPNs@LMWH subjected to a simulated digestion process. The data are
presented as the means ± SDs.

DAPI/Naparticles/CM-Dil

804
805 **Supplementary Fig. 8 | OPNs@LMWH internalization by M1 macrophages.** M1-type
806 RAW264.7 cells were incubated with FA-OPNs or FA-OPNs@LMWH after pretreatment with
807 or without the anti-integrin αM antibody and visualized via CLSM. Scale bar, 50 μm.
808 Representative images of n=6 biologically independent samples are shown.
809
810

811
Supplementary Fig. 9 | Quantitative analysis of the FA fluorescence intensity of the
internalized nanoparticles. FA fluorescence intensity measurements by flow cytometry
confirmed that free FA did not enter the cells. Quantitative analysis of 4 biologically
independent samples is shown. The data are presented as the means ± SDs. $*$ p < 0.05, $**$ p

< 0.01, *** p < 0.001, **** p < 0.0001 and ns p > 0.05, analysed by two-way ANOVA multiple
comparisons test.
818
819

820
821 **Supplementary Fig. 10 | The high level of CD11b in the colonic submucosal layer of colitis**
822 **model mice.** Representative images of immunohistochemical staining for CD11b in healthy
823 (**a**) or colitis (**b**) mice. Scale bar, 100 μm. Representative images from n= 3 biologically
824 independent samples are shown.
825
826

Supplementary Fig. 11 | OPNs@LMWH accumulate in inflamed colon tissue infiltrated

by inflammatory cells. Healthy and colitis model mice were orally administered FA-OPNs or

FA-OPNs@LMWH, and colon tissues were collected after 24 h and stained with DAPI, CM-

Dil and integrin-αM and visualized via CLSM. Scale bar, 50 μm. Representative images of n=3

animals are shown.

Supplementary Fig. 12 | Cytotoxicity determined by CCK-8 assays. a, b, Viability of

RAW264.7 cells (**a**) and NCM460 cells (**b**) after incubation with OPNs for 24 h. **c, d,** Viability

of RAW264.7 cells (**c**) and NCM460 cells (**d**) after incubation with OPNs@LMWH for 24 h.

Quantitative analysis of n=6 biologically independent samples (**a-d**) is shown. The data are

presented as the means ± SDs.

842
Supplementary Fig. 13 | Concentration-dependent intracellular ROS scavenging
behaviour. a, RAW264.7 cells were stimulated with PMA and treated with different
concentrations of OPNs or OPNs@LMWH. **b, c,** Fluorescence intensities (**b**) and percentages
of ROS-positive cells (**c**) were determined. The representative gating strategies or quantitative
analyses of 4 biologically independent samples (**a-c**) are shown. The data are presented as
the means ± SDs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ and ns $p > 0.05$,
analysed by two-way ANOVA (**b, c**).
850
851

Supplementary Fig. 14 | LMWH has little anti-ROS activity. a, b, RAW264.7 cells were
stimulated with PMA and treated with LMWH (20 μg mL^{-1}). The level of intracellular ROS was
detected by DCFH-DA, determined by flow cytometry (**a**) and visualized by fluorescence
microscopy (scale bar, 50 μm) (**b**). **c, d,** Fluorescence intensities (**c**) and percentages of ROS-
positive cells (**d**) were determined. Representative gating strategies, images and quantitative
analyses of n= 3 or 4 biologically independent samples (**a-d**) are shown. The data are
presented as the means ± SDs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ and ns
$p > 0.05$, analysed by an unpaired t test (**c, d**).

863
864 **Supplementary Fig. 15 | Concentration of free LMWH determined by Azure A.** The
amount of free LMWH in the supernatant before and after coating was detected, which
indicated that LMWH was present in a low mass proportion in OPNs@LMWH. The
quantitative analysis of 4 biologically independent samples is shown.

Supplementary Fig. 16 | Safety profile of OPNs@LMWH (body weight). Mice were orally administered different doses of nanoparticles for 14 days. Daily body weight changes were recorded. Quantitative analysis of n=6 animals is shown. The data are presented as the means ± SDs. * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, **** $p < 0.0001$ and ns $p > 0.05$, analysed by two-way ANOVA (**b, c**).

Supplementary Fig. 17 | Safety profile of OPNs@LMWH (major organs). Mice were
sacrificed on day 14. Major organ (heart, liver, kidney, lung, spleen, and colon) sections were
stained with haematoxylin and eosin. Scale bar, 500 μm. Representative images from n= 3
animals are shown.

Supplementary Fig. 18 | Safety profile of OPNs@LMWH (blood haematology). Mice were
sacrificed on day 14. Blood samples were collected and analysed by using blood haematology;
the results included red blood cell (RBC) (**a**), haemoglobin (HGB) (**b**), haematocrit (HCT) (**c**),
platelet (PLT) (**d**), white blood cell (WBC) (**e**), neutrophil (NEUT) (**f**), lymphocyte (LYMPH) (**g**),
monocyte (MONO) (**h**), eosinophil (EO) (**i**) and basophil (BASO) (**j**) counts. The quantitative
analysis of n= 4-6 animals is shown. The data are presented as the means ± SDs. ∗ $p < 0.05$,
∗∗ $p < 0.01$, ∗∗∗ $p < 0.001$, ∗∗∗∗ $p < 0.0001$ and ns $p > 0.05$, analysed by ordinary one-way
ANOVA multiple comparisons test.

894