TITLE

Interspecies secreted surfactants induce emergent motility in *Pseudomonas aeruginosa*

AUTHORS

Delayna L. Warrell[a]*, Tiffany M. Zarrella[a,b,c#]* , Christopher Machalek[a], Anupama Khare[a#]

[a]Laboratory of Molecular Biology, Center for Cancer Research, National Cancer Institute, National

Institutes of Health, Bethesda, MD, USA

[b]Postdoctoral Research Associate Training Program, National Institute of General Medical

Sciences, National Institutes of Health, Bethesda, MD, USA

[c]Current address: Department of Biology, Georgetown University, Washington, DC, USA

Running head: Surfactants induce motility in *P. aeruginosa*

[#]Address correspondence to Tiffany M. Zarrella, tiffany.zarrella@georgetown.edu, or Anupama

Khare, anupama.khare@nih.gov

*Delayna L. Warrell and Tiffany M. Zarrella contributed equally to this work. Author order was

determined alphabetically.

Key words: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, surfactants, motility,

polymicrobial interactions, phenol-soluble modulins

The authors declare no conflict of interest.

ABSTRACT

In most natural environments, bacteria live in polymicrobial communities where secreted

molecules from neighboring species alter bacterial behaviors including motility, but such

interactions are understudied. *Pseudomonas aeruginosa* is a motile opportunistic pathogen that

exists in diverse multispecies environments such as the soil and is frequently found in human

wound and respiratory tract co-infections with other bacteria including *Staphylococcus aureus*.

Here we show that *P. aeruginosa* can co-opt secreted molecules from other species for

flagellar-based emergent motility. We found that in the presence of exoproducts from *S. aureus*

and other bacteria, *P. aeruginosa* switched from swarming to an alternative form of motility

where it slid and spread on semi-solid surfaces, and it showed similar behavior on hard agar

where it was otherwise unable to move. Surfactants from these species were required for the *P.*

aeruginosa emergent motility, and it was also induced by the addition of numerous exogenous

biological and synthetic surfactants. Mutant analysis indicated that this motility was similar to a

previously described mucin-based *P. aeruginosa* motility, 'surfing', albeit with divergent

regulation. Thus, our study establishes a major *P. aeruginosa* surfing-like emergent motility

where secreted surfactants from the host as well as neighboring bacterial and interkingdom

species serve as public goods, allowing *P. aeruginosa* to access different environmental niches.

INTRODUCTION

Bacteria are often found in dynamic, complex communities where interspecies secreted factors influence bacterial behaviors, such as antibiotic resistance [1-3], biofilm formation [4-7], and motility [8-10]. However, many of the polymicrobial interactions leading to the modification of these pathogenic traits have not been elucidated. The study of pairwise interactions has the potential to reveal behavioral changes that occur within communities as well as the underlying molecular mechanisms [11, 12].

Motility allows bacterial cells to disperse and migrate to new niches for nutrients, seek out prey, and adapt to new environments; thus, modulation of motility behaviors due to ecological interactions is likely to be advantageous for bacterial fitness. Several examples of changes in bacterial motility in mixed cultures compared to monocultures have been previously reported, including the inhibition of motility via secreted volatiles [9], and emergent motility in co-culture conditions where the monocultures did not move. These include social spreading between two non-motile species [13], exploration by non-motile bacteria upon sensing fungal volatiles [8], and non-motile bacteria inducing motility in a neighboring species, and then hitchhiking on the moving population [14].

Pseudomonas aeruginosa is a social, motile microbe that typically inhabits polymicrobial environments in soil, wounds, and persistent infections of the respiratory tracts of people with cystic fibrosis (CF) [15-17], in which the presence of *P. aeruginosa* is associated with worsened lung function and disease outcomes [18]. *P. aeruginosa* is commonly found in concert with *Staphylococcus aureus* in wound infections and nosocomial pneumonia, and these two species are the most frequently isolated bacterial pathogens from the airways of people with CF [19]. Chronic co-infections of *P. aeruginosa* and *S. aureus* are associated with increased disease severity [20-24]. Since these two organisms are often found together, we posited that interactions with *S. aureus* may alter *P. aeruginosa* pathogenic behaviors, including motility.

P. aeruginosa displays several well-established forms of motility: swimming, twitching, and

swarming **(Supp. Fig. 1A)**. Motility is intricately linked with virulence regulation, antibiotic

resistance, and biofilm formation [25-27], and flagellar competency is associated with

dissemination and pathogenesis [28-31]. During swimming motility *P. aeruginosa* uses a single

polar flagellum to propel through liquid [32], while for twitching, type IV pili extension and retraction

are required for movement on hard surfaces [33]. Swarming is a social behavior with a

characteristic tendril formation on semi-solid surfaces that requires flagella and the secretion of

rhamnolipids as a surfactant [34, 35]. Another type of motility that has been described is 'surfing',

where *P. aeruginosa* spreads over semi-solid media containing the glycoprotein mucin, that is

abundant in the mucus accumulated in the CF airways and gastrointestinal tract [36, 37]. This

motility is thought to depend on the viscous nature and lubricant properties of mucin and requires

P. aeruginosa flagellar function while pili and rhamnolipids are expendable [36, 38].

P. aeruginosa motility is known to be affected by interspecies interactions. Fungal volatiles

farnesol and ethanol at high concentrations inhibit *P. aeruginosa* motility, while lower

concentrations of ethanol induce swarming when carbon sources are limited, possibly signaling

the availability of nutrients [39-41]. Further, it has been observed that *P. aeruginosa* engages in

directional pili-based 'exploratory motility' in response to the *S. aureus* secreted cytotoxins

phenol-soluble modulins (PSMs) [42]. *P. aeruginosa* explorers sense PSMs via the Pil-Chp

chemosensory system to twitch on surfaces toward *S. aureus* [43]. However, how other *P.*

aeruginosa motility behaviors are affected by *S. aureus* is not known.

In this study, we investigate whether additional *P. aeruginosa* motility behaviors are

altered by *S. aureus* exoproducts. We observe that *P. aeruginosa* displays emergent motility in

the presence of *S. aureus* secreted molecules, that enables it to slide and spread on semi-solid,

and even hard agar surfaces where it is otherwise nonmotile. This motility depends on flagella,

but does not require type IV pili, rhamnolipids, the three quorum sensing systems, or several other

regulators that affect motility including the Gac/Rsm system. Further, transcriptomics of cells

undergoing emergent motility did not reveal responses in motility genes, suggesting that the

transition to emergent motility may not be due to transcriptional regulation. Multiple CF clinical

isolates of *P. aeruginosa* and *S. aureus* also exhibit and induce emergent motility, respectively,

indicating that these behaviors are widespread within these species. Further, secreted products

from other bacterial species that produce biosurfactants, such as *Bacillus subtilis* and *P.*

aeruginosa itself, but not from species that do not make this class of molecules, also promote

emergent motility. Mutants that are deficient for the respective biosurfactants in these species are

unable to induce emergent sliding and spreading, while diverse interkingdom biological and

synthetic surfactants can, demonstrating that *P. aeruginosa* displays this motility in the presence

of molecules with surfactant properties. Finally, a comparison with mucin-mediated surfing

suggests that emergent sliding and spreading is a surfing-like motility, although movement on

mucin requires the LasR quorum sensing regulator, while movement on surfactants is LasR-

independent.

RESULTS

***P. aeruginosa* displays emergent motility in the presence of *S. aureus* secreted products.**

To determine the effect of *S. aureus* secreted products on *P. aeruginosa* motility, we

performed *P. aeruginosa* twitching, swimming, and swarming assays in the presence of 25% (v/v)

S. aureus cell-free supernatant, or media salts as a control. As reported previously [42], we

observed that *P. aeruginosa* twitched farther with the addition of *S. aureus* supernatant **(Supp.**

Fig. 1B). In the swimming assay, while *P. aeruginosa* swam uniformly creating a circular swim

area in the control agar, the swim patterns were irregular and oval shaped in the presence of *S.*

aureus supernatant **(Supp. Fig. 1C)**.

In a standard swarm assay on semi-solid agar plates (containing 0.5% agar) *P. aeruginosa*

swarmed with characteristic tendril formation on the control plate **(Fig. 1A)**. However, on plates

with 25% *S. aureus* supernatant, *P. aeruginosa* exhibited a motility unlike swarming where the

bacteria first slid on the agar and spread out across the plate, and then showed some tendril

formation **(Fig. 1A)**. *P. aeruginosa* exhibited similar motility on agar containing 12.5% *S. aureus*

supernatant and an intermediate motility on 5% supernatant, while swarming resumed on plates

with 1% supernatant **(Fig. 1A)**. Next, we tested if this phenotype occurs on hard agar plates

(containing 1.5% agar) in which *P. aeruginosa* is traditionally unable to move. While *P. aeruginosa*

did not move on the hard agar control or on the plates containing 5% and 1% *S. aureus*

supernatant **(Fig. 1B),** it exhibited emergent motility on the plates containing 25% and 12.5% *S.*

aureus supernatant **(Fig. 1B)**. Thus, *S. aureus* secreted products enable a distinct form of *P.*

aeruginosa motility, even in conditions where *P. aeruginosa* is normally nonmotile.

***P. aeruginosa* emergent motility requires flagella, but not pili or rhamnolipids.**

The established forms of *P. aeruginosa* motility include swimming, which requires flagella

[32], twitching, requiring pili [33], and swarming in which pili, flagella, and the surfactant

rhamnolipids all contribute **(Supp. Fig. 1A)** [34, 35]. Additionally, motility in *P. aeruginosa* is also

affected by quorum sensing systems and other regulatory pathways [34, 44, 45]. We therefore

tested the requirement of flagella, pili, rhamnolipids, and known motility regulators for emergent

sliding and spreading exhibited by *P. aeruginosa* in the presence of *S. aureus* secreted products.

First, we tested several mutants that lack the following structural components of the *P.*

aeruginosa flagella: the proximal rod protein FlgB, the hook protein FlgE, or the flagellin subunit

FliC [46], and are therefore defective in swimming **(Supp. Fig. 2A)**. As expected, on the semi-

solid control agar the WT was able to swarm while the flagellar mutants did not, and no motility

was observed for all strains on the control hard agar **(Fig. 2A and Supp. Fig. 2B)**. On the semi-

solid and hard agar plates with *S. aureus* supernatant, the *flgB::*tn, *flgE::*tn, and Δ*fliC* mutants only

slid on the plates without spreading out, unlike the motility exhibited by the WT strain **(Fig. 2A**

and Supp. Fig. 2B). To examine the motility in a species that does not encode flagella, we tested

S. aureus itself, and found that it was immobile on the control agar and slid on the plates with *S.*

aureus supernatant like the *P. aeruginosa* flagellar mutants **(Fig. 2B and Supp. Fig. 2C)**. To

determine if these flagella-deficient strains could move in the presence of motile *P. aeruginosa*,

these strains were fluorescently labeled, mixed, and added to hard agar with *S. aureus*

supernatant. As expected, two distinctly labeled *P. aeruginosa* WT strains co-cultured on hard

agar exhibited emergent motility equally **(Fig. 2C)**. However, the Δ*fliC* mutant and *S. aureus* still

only slid on the plate even in co-culture with *P. aeruginosa* WT cells that showed surface

spreading **(Fig. 2C)**, suggesting that this motility is an active process, and the presence of a motile

population cannot compensate for loss of competent flagella. Hence, *P. aeruginosa* emergent

motility in response to *S. aureus* secreted products requires active flagellar function.

We next determined if pili, which are required for twitching and contribute to swarming

[34], are necessary for emergent sliding and spreading. We tested multiple twitching-deficient pili

mutants **(Supp. Fig. 3A)** - two mutants deficient in pili retraction, *pilT*::tn and *pilU*::tn [47], two

mutants lacking the extended pili fiber, *pilB*::tn, via deficiency in pili extension [48], and Δ*pilA* due

to the lack of the pilin subunit [33], and two mutants deficient for pili-mediated chemotaxis, Δ*pilJ*

and Δ*pilG* [49]. While these mutants showed differences in swarming on the control plates, all of

these strains, except *pilB*::tn, displayed emergent motility on plates with *S. aureus* supernatant

(Fig. 2D, Supp. Fig. 3B and C, Supp. Fig. 4A). The pili-extension mutant *pilB*::tn slid on the plate

without spreading out, similar to the movement displayed by the flagellar mutants, **(Fig. 2A)**, and

there may be cross-regulation between pili and flagella [50, 51]. We therefore tested the pili

mutants for swimming motility as a measure of their flagellar functionality. The *pilU*::tn and *pilB*::tn

mutants had diminished swim rings, while the other mutants showed a normal swim phenotype

(Supp. Fig. 3D). Since swimming requires flagella, it is likely that the flagellar dysfunction in the

pilB::tn mutant also leads to a lack of emergent motility. In the *pilU*::tn strain, it is possible that the

flagella is defective for swimming but still functional for emergent motility. Taken together, *P.*

aeruginosa emergent motility in the presence of *S. aureus* secreted products requires flagella but

not pili.

While swarming, *P. aeruginosa* secretes rhamnolipids, biosurfactants that lubricate the

area and allow the cells to move over the surface [34, 35]. We found that while a mutant deficient

in rhamnolipid production, Δ*rhlA*, was unable to swarm as previously reported [34, 35], it still slid

and spread on both semi-solid and hard agar plates containing *S. aureus* cell-free supernatant

(Fig. 2E and Supp. Fig. 4B), suggesting that rhamnolipids are not required for emergent motility.

Finally, to determine how this motility is regulated we analyzed the role of diverse *P.*

aeruginosa signaling pathways in this motility. *P. aeruginosa* utilizes three quorum sensing

systems: Las, Rhl, and PQS [52]. To examine the role of quorum sensing in this motility we tested

mutant strains lacking each individually (Δ*rhlR, ΔlasR, ΔpqsA*)*,* or in combination (Δ*rhlR ΔlasR*

Δ*pqsA*). The Δ*rhlR, ΔlasR,* and Δ*rhlR ΔlasR ΔpqsA* mutants did not swarm on the control plates,

while the Δ*pqsA* mutant did **(Supp. Fig. 5A)**. However, all mutants displayed emergent motility

on the plates with *S. aureus* supernatant to varying degrees, with the Δ*pqsA* strain being hyper-

motile compared to the other strains and covering the entire semi-solid agar plate **(Supp. Fig. 5A**

and B). We also tested mutants from additional regulatory pathways that control motility: Δ*gacA,*

response-regulator of the Gac/Rsm pathway involved in virulence and motility; Δ*rsmA,* a

posttranscriptional regulator of the Gac/Rsm pathway; Δ*retS*, a response regulator which can

block GacA function [53]; Δ*wspR,* the response regulator of the Wsp chemosensory pathway [54,

55]; and Δ*toxR,* a c-di-GMP binding protein involved in motility and biofilm regulation [56]. The

Δ*retS* and Δ*rsmA* mutants did not swarm, while the Δ*gacA, ΔwspR,* and Δ*toxR* mutants exhibited

swarming similar to WT *P. aeruginosa* **(Supp. Fig. 6)***.* Despite these differences in swarming, all

these regulatory mutants exhibited emergent sliding and spreading **(Supp. Fig. 6)**. Therefore,

these regulatory pathways are not required for *P. aeruginosa* emergent motility, although quorum

sensing may affect motility patterns*.*

Cells undergoing emergent motility do not transcriptionally regulate motility pathways.

To identify which pathways are differentially regulated during emergent sliding and spreading, we performed RNA sequencing (RNA-seq) on *P. aeruginosa* cells scraped from the leading edge on semi-solid or hard agar plates containing the media salts control or *S. aureus* supernatant at 17 hours **(Supp. Fig. 7A)**. We found approximately 100 to 350 genes that were upregulated and downregulated on the supernatant compared to the media salts control in both cases **(Supp. Tables 1 and 2)**. We reasoned that the differentially regulated genes would include genes specific to this motility as well as genes that were regulated upon exposure to unrelated *S. aureus* secreted molecules. We therefore compared the genes to those we previously identified as being differentially regulated in early log-phase *P. aeruginosa* planktonic cultures 2 hours after the addition of *S. aureus* supernatant compared to the addition of media salts as a control [57], and found significant overlaps **(Fig. 3A)**. Some of the highest upregulated genes in both motility conditions were genes involved in tricarboxylic acids (TCA) uptake and acetoin catabolism, which are upregulated in planktonic conditions as well, likely due to *S. aureus* secreted citrate and acetoin, respectively [57] **(Fig. 3B, Supp. Fig. 7B and C, Supp. Tables 2, 3, and 4)**. Next, we performed Gene Ontology (GO) enrichment analysis of the *P. aeruginosa* genes differentially regulated in the individual motility conditions, both motility conditions (semi-solid and hard agar plates), as well as those regulated exclusively in both motility conditions, but not in the planktonic phase [58-60] **(Fig. 3C, Supp. Fig. 7D and E, and Supp. Tables 3, 4, and 5)**. Genes involved in aerobic respiration and response to heat were enriched in the genes downregulated in both motility conditions **(Supp. Fig. 7E and Supp. Table 5)**, but these genes were also downregulated in the planktonic phase and are likely unrelated to the motility **(Supp. Fig. 7E and Supp. Tables 4 and 5)**.

The *P. aeruginosa* genes differentially regulated exclusively in both motility conditions, but not in the planktonic phase, were not enriched for any motility pathways **(Fig. 3C and Supp.**

Table 5), suggesting that transcriptional regulation may not play an important role during

emergent motility, consistent with our results that major motility regulators are not required for this

motility **(Supp. Fig. 5 and 6)**. Instead, genes downregulated exclusively in the motility conditions

showed enrichment of the Type 3 Secretion System (T3SS) and heme biosynthesis pathways,

while the upregulated genes showed no statistically enriched pathways **(Fig. 3C and D, and**

Supp. Table 5). Additionally, on the hard agar there was an upregulation in transcripts from the

phage φCTX **(Fig. 3B, Supp. Fig. 7B, and Supp. Table 2)**, indicating prophage induction in this

condition. This likely explains the presence of plaque-like clearings on the hard agar (e.g. in **Fig.**

1B), which was especially apparent for the LasR mutant **(Supp. Fig. 5)**. Multiple cellular stresses

can lead to both induction of prophage and downregulation of T3SS [61-64], indicating that *P.*

aeruginosa senses physiological stress while undergoing emergent motility. Further, heme

biosynthesis and associated denitrification genes (e.g. *nirS*, *nirQ*, *norC, norB)* are induced in

anaerobic conditions [65]. The relative downregulation of these genes during emergent motility

may be indicative that the control cells (swarming cells in the semi-solid agar and non-motile cells

in the hard agar) are switching to anaerobic metabolism, while cells undergoing emergent motility

continue with aerobic respiration. Thus, *P. aeruginosa* may experience physiological stress and

metabolic changes while undergoing emergent motility in the presence of *S. aureus* exoproducts.

Interspecies surfactants facilitate emergent motility in *P. aeruginosa*.

To determine the generality of this emergent motility, we performed the motility assay with

clinical isolates of both species. We tested four clinical isolates of *P. aeruginosa* collected from

different people with CF (CF017, CF033, CF057, and CF095) **(Supp. Table 6)** on both semi-solid

and hard agar and observed that the isolates CF017 and CF057 swarmed and exhibited emergent

motility, while isolates CF033 and CF095 did not **(Supp. Fig. 8A and B)**. To determine if the

motility deficiencies were due to the dysfunction of motility appendages, we tested these isolates

for swimming and twitching. Isolate CF017 swam similar to *P. aeruginosa* PA14, while isolates

CF033, CF057, CF095 all had swimming deficiencies **(Supp. Fig. 9A)**. Additionally, the twitch

zone of CF033 was similar to PA14, but CF017 and CF057 had smaller twitch zones and CF095

did not twitch **(Supp. Fig. 9B)**. Thus, the isolates that do not exhibit the emergent motility may

have flagellar and/or pili deficiencies.

These *P. aeruginosa* strains were co-isolated with four *S. aureus* strains, CF019, CF032,

CF058, and CF100, respectively **(Supp. Table 6)**. Cell-free supernatants from each of these *S.*

aureus clinical isolates could induce emergent motility in *P. aeruginosa* PA14 **(Supp. Fig. 10A**

and B). These results demonstrate that while there is some variation, clinical strains of *P.*

aeruginosa and *S. aureus* can exhibit and induce emergent surface spreading, respectively.

Next, we investigated whether other species induce this emergent motility, and found

that *P. aeruginosa* exhibited this motility on plates with supernatant from *S. aureus, P. aeruginosa,*

and *Bacillus subtilis* ZK3814 **(Fig. 4A)**, but swarmed on plates with supernatant from *B. subtilis*

PY79, *Klebsiella pneumoniae, Vibrio cholerae, Escherichia coli, Burkholderia cenocepacia*

BAA245, and *B. cenocepacia* 25608 **(Fig. 4A and B)**.

A common class of molecules known to be secreted by each of the inducing strains, but

none of the non-inducing strains, and previously implicated in motility, is biosurfactants [66]. *S.*

aureus produces phenol soluble modulins (PSMs) [67-69], *P. aeruginosa* produces rhamnolipids

[70], and *B. subtilis* ZK3814 produces fengycin and surfactin [71]. We therefore tested the

supernatant of mutants deficient in each of these biosurfactants for their ability to induce motility

in *P. aeruginosa*. The supernatants of the *S. aureus* JE2 Δ*psmaδ*, which is deficient in the

production of PSMs α1-4 and delta toxin, as well as a Δ*psmaβδ* mutant (that lacks all PSMs) in

the *S. aureus* LAC strain, did not induce emergent motility in *P. aeruginosa* **(Fig. 4C and Supp.**

Fig. 11A, B, and C). Further, supernatant from *agrA*::tn, *agrB*::tn, or *agrC*::tn, mutants of the *S.*

aureus agr quorum sensing system which is required for the production of PSMs, also did not

lead to *P. aeruginosa* emergent motility **(Supp. Fig. 11A and D)**. Similarly, we found that *P.*

aeruginosa did not exhibit this motility on plates containing supernatant from the *P. aeruginosa*

Δ*rhlA* mutant lacking the biosurfactant rhamnolipids **(Fig. 4D and Supp. Fig. 12A)**. Finally, for *B.*

subtilis, supernatant from Δ*srfA*, a mutant deficient for surfactin production, was unable to induce

P. aeruginosa emergent motility, unlike the parental WT *B. subtilis* ZK3814 as well as Δ*fenA*, a

mutant deficient for fengycin production **(Fig. 4E and Supp. Fig. 12B)**. Collectively, these data

show that the production of biosurfactants such as *S. aureus* PSMs, *P. aeruginosa* rhamnolipids,

and *B. subtilis* surfactin, is required for the interspecies induction of sliding and spreading motility

in *P. aeruginosa*.

Detergents and surfactants are sufficient to induce emergent motility in *P. aeruginosa*.

Given that bacterial-secreted surfactants were necessary to induce emergent motility in

P. aeruginosa, we investigated if the addition of biological and synthetic surfactants was sufficient

to induce this motility. We tested commercially obtained biosurfactants from the motility-inducing

species, including rhamnolipids from *P. aeruginosa* and surfactin from *B. subtilis* ZK3814*;* mucin,

which has been previously implicated in *P. aeruginosa* surfing motility [38]; and the synthetic

surfactants sodium dodecyl sulfate (SDS), cetrimonium bromide (CTAB), and Triton X-100. *P.*

aeruginosa exhibited emergent motility, to varying degrees, on plates with *S. aureus* supernatant,

the biosurfactants surfactin, rhamnolipids, and mucin, and the synthetic surfactants SDS and

Triton X-100, but motility was inhibited on both semi-solid and hard agar plates with the addition

of CTAB **(Fig. 5A)**. Further, we tested biosurfactants from interkingdom species, such as

mannosylerythritol lipid A (MEL-A) and sophorolipid from yeasts, plant-produced saponin, and

human cathelicidin LL-37 and pulmonary surfactant dipalmitoylphosphatidylcholine (DPPC). All

these interkingdom surfactants, except DPPC, induced the sliding and spreading motility,

especially on semi-solid agar **(Fig. 5B)**. Thus, diverse biological and synthetic surfactants are

sufficient to induce *P. aeruginosa* emergent motility.

P. aeruginosa surfactant-induced surface spreading is a surfing-like motility.

Previously, it was reported that _P. aeruginosa_ exhibits 'surfing', a surface-associated, flagellar-based motility which occurs on semi-solid agar containing mucin [36, 38, 72, 73]. To compare the genetic requirements in _P. aeruginosa_ of mucin-based surfing behavior to microbial surfactant-induced emergent motility, we tested motility appendage, quorum sensing, and other regulatory mutants on semi-solid plates containing _S. aureus_ supernatant or mucin. The appendage mutants showed similar phenotypes for motility on _S. aureus_ supernatant and mucin, where _flgB_::tn, Δ_fliC,_ and _pilB_::tn showed no motility on either, while _pilT_::tn and Δ_pilA_ moved on both, though _pilT_::tn moved only moderately on mucin **(Supp. Fig. 13A)**.

Previously, Rhl and Las as well as components of Gac/Rsm were reported to contribute to mucin-based surfing [36, 72, 73]. We observed that while the Δ_pqsA_ mutant moved on both the _S. aureus_ and mucin plates, the Δ_rhlR_ and Δ_rhlR_ Δ_lasR_ Δ_pqsA_ mutants moved moderately on _S. aureus_ supernatant and had no movement on the plates with mucin **(Supp. Fig. 13B)**. Interestingly, the Δ_lasR_ mutant moved on the _S. aureus_ supernatant but not on mucin, demonstrating a divergence of quorum sensing requirements for mucin-based surfing [36, 38] and surfactant-dependent motility. The Δ_retS_, Δ_gacA_, Δ_rsmA_, Δ_wspR_, and Δ_toxR_ regulatory mutants all moved on both _S. aureus_ supernatant and mucin, but Δ_retS_ and Δ_rsmA_ had less movement on both plate types **(Supp. Fig. 14)**.

To further delineate the role of LasR in motility on a variety of surfactants, we tested the Δ_lasR_ mutant on semi-solid plates with supernatants from _S. aureus_, _P. aeruginosa_, and _B. subtilis_ ZK3814, as well as with the addition of mucin and the surfactant Triton X-100. While the Δ_lasR_ mutant did not move on mucin, it exhibited the motility on all of the other supernatant or surfactant-containing plates **(Fig. 6A)**. Overall, the surfactant-dependent emergent motility we observed and mucin-dependent surfing motility have similar appendage and regulatory

requirements, indicating that surfactant-based emergent sliding and spreading is a surfing-like

motility where the exact quorum-sensing regulation may differ between movement on mucin

compared to other surfactants.

DISCUSSION

Bacterial locomotion is an advantageous trait that enables migration to more favorable

environments and access to fresh nutrients. Within microbial communities, interspecies secreted

factors can affect bacterial motility, by activation [8], inhibition [9], or endowing nonmotile strains

the capability to move [13, 14]. Many of these observations were reported by examining pairwise

interactions, which can reveal novel behavioral phenomena as well as the underlying mechanisms

[11]. Here, we interrogated how *P. aeruginosa* motility is affected by *S. aureus* secreted molecules

and uncovered a surfing-like motility that depends on the presence of exogenous interspecies or

synthetic surfactants **(Fig. 6B)**. The surfactant-based motility we observed requires flagellar

function, but not type IV pili, rhamnolipids, quorum sensing, or other known motility regulatory

systems like Gac/Rsm. It permits emergent motility when *P. aeruginosa* is typically nonmotile and

is induced by surfactants from a variety of species, including co-infecting pathogens, putative

environmental neighbors, as well as infection hosts, demonstrating that *P. aeruginosa* can take

advantage of specific cues from nearby species to explore new niches.

We observed that in the presence of interspecies surfactants, *P. aeruginosa* cells first slid

on the agar surface similar to nonmotile bacteria, like *S. aureus* or non-flagellated *P. aeruginosa*

strains **(Fig. 2A and B)**. The cells then spread over the surface, an active process that required

flagellar function. This was followed by tendril formation on semi-solid agar, or likely dispersal of

individual cells on hard agar that resulted in colonies at the mobile leading edge. The motility was

induced by the presence of diverse types of biological and synthetic molecules of a variety of

chemical compositions, suggesting that their shared properties of reducing surface tension and

acting as wetting agents led to the motility. The reduced surface tension likely allows for increased

movement, resulting in motility even on hard agar, where *P. aeruginosa* is otherwise nonmotile,

and enabling surface motility without rhamnolipid production, unlike what is seen for swarming.

Intrinsic rhamnolipid production occurs and is required during swarming, while the presence of

exogenous rhamnolipids leads to emergent motility **(Fig. 5A)**. Given that *P. aeruginosa*

rhamnolipid production is quorum-sensing dependent and requires high cell density, it is likely

that in the presence of foreign surfactants, *P. aeruginosa* cells immediately sense the exogenous

surfactants resulting in emergent motility before they produce rhamnolipids.

Motility with similar characteristics has been previously reported in *P. aeruginosa*. When

other gelling agents, the polysaccharides gellan gum and carrageenan, are used in place of agar

in semi-solid media, *P. aeruginosa* exhibits flagellar-dependent, rhamnolipid-independent surface

motility due to the reduction in surface tension [74], like the surfactant-mediated motility we

observed. Further, surfing motility in *P. aeruginosa* has previously been described to occur on

semi-solid agar plates (0.5 - 1.0% agar) containing the glycopeptide mucin and require the use of

flagella [36, 38]. While these motility types were thought to be distinct [36, 38, 74], our data

suggests that these are likely variations of the same *P. aeruginosa* surface motility that occurs

upon reduction of surface tension and requires active flagella, but not pili or rhamnolipids, where

the main distinction is that only mucin-mediated surfing requires LasR quorum sensing **(Fig. 6A)**

[36, 38, 74]. We propose that the term 'surfing' encompasses all these variations, and our study

thus establishes surfing as a major form of *P. aeruginosa* motility that occurs on both semi-solid

and hard surfaces in the presence of a variety of biotic and abiotic surfactants but is distinct from

other established motility types like swimming, twitching, or swarming.

The signal transduction pathway from sensing environmental changes brought on by

surfactants leading to flagellar motility remains elusive. We examined the requirement of several

regulators that control motility [75], such as components of Pil-Chp, quorum sensing, Gac/Rsm,

and Wsp surface sensing, but none were needed for surfactant-based surfing. Nonetheless, there

were noticeable differences in the appearance of quorum sensing mutant surfers. On semi-solid

agar, the final step of tendril formation was absent in Rhl mutants that are expected to lack

rhamnolipid production, indicating that the tendrils here may be similar to those seen in swarming,

and require intrinsic production of rhamnolipids. In addition, mutants devoid of the PQS pathway,

which plays a role in swarming tendril repulsion [76], are hyper-motile on semi-solid agar,

suggesting that the PQS pathway may play a role in limiting surfactant-based surfing. Given the

non-essentiality of these pathways for surfing, it is possible that the *P. aeruginosa* polar flagellum

directly senses the surfactant by the alleviation of impediments on flagellar rotation [77].

Moreover, many of the signaling cascades that lead to changes in flagellar motion involve the

motile-to-sessile second messenger cyclic di-GMP [54], which is a putative target for future

exploration of the mechanisms underlying this interspecies-induced motility.

P. aeruginosa and *S. aureus* often co-infect wounds and the airways of people with CF

where they are associated with faster lung function decline, an increase in intravenous antibiotic

treatments, and an increase in lower airway inflammation [20, 21]. We observed that the presence

of *S. aureus* PSMs, which are amphipathic alpha-helical peptides with surfactant properties,

induces surfing motility in *P. aeruginosa*. Previously, these *S. aureus* secreted toxins have been

implicated in enabling motility in both species. In *S. aureus*, PSMs allow nonmotile *S. aureus* to

slide on surfaces [68]. Additionally, *P. aeruginosa* senses PSMs via the Pil-Chp chemosensory

system to induce pili-dependent directional 'exploratory motility' [42, 43], which is a motility distinct

from the one we describe. *P. aeruginosa* explorers require pili and components of the Pil-Chp

system, such as the chemoreceptor PilJ [43], which are not necessary for flagellar-dependent

surfactant-based surfing. Sensing of surface hardness and viscosity plays a role in stimulating

motility behaviors [78, 79], with twitching and swarming occurring on hard and semi-solid

surfaces, respectively. However, we observed that even on hard agar surfactant-based surfing

still solely required flagella and not pili, in contrast to the directionality of exploratory motility

detected on hard agar [42, 43]. PSMs produced by neighboring *S. aureus* cells are also thought

to alter the flow of *P. aeruginosa* rhamnolipids, leading to repulsion of *P. aeruginosa* swarm

tendrils [80]. Thus, there is an overarching role for PSMs in inducing and altering motility in *P.*

aeruginosa, yet the mode of locomotive response likely differs based on additional environmental

stimuli and/or how these molecules are sensed by *P. aeruginosa*.

In addition to *S. aureus*-produced PSMs [68], many microbes secrete diverse classes of

biosurfactants that enable locomotion [81], such as the lipopeptide surfactin by *Bacillus subtilis*

[82], the glycolipids rhamnolipids by *Pseudomonas aeruginosa* and *Burkholderia cenocepacia*

[83, 84], and the aminolipid serrawettin by *Serratia* spp. [85]. As secreted factors, interspecies

surfactants can serve as public goods and enable other species to co-opt these wetting agents

for motility, and we observed that *P. aeruginosa* can utilize diverse intra- and interkingdom

surfactants for surfing. In phyllosphere-dwelling bacterial species, secretion of the surfactant

viscosin B by *Pseudomonas* sp. FF1 leads to co-swarming with *Pantoea eucalypti* 299R [86]. This

has also been observed in conditions where *B. cenocepacia* is individually non-motile (and does

not produce surfactant) but uses *P. aeruginosa* rhamnolipids for motility [87, 88]. Further, flagella-

deficient *P. aeruginosa* that produces rhamnolipids is incapable of swarming but can co-swarm

with *B. cenocepacia* that is using the secreted surfactant [88], providing benefits for each species.

Despite this display of hitchhiking on motile *B. cenocepacia*, in our study we did not observe

cooperation between surfing *P. aeruginosa* and flagellar mutants **(Fig. 2C)**, demonstrating that

intraspecies hitchhiking is unlikely to contribute to this form of *P. aeruginosa* motility. However,

the species providing the surfactants may benefit from the interaction if able to hitchhike on *P.*

aeruginosa, although we did not see hitchhiking of *S. aureus* on the surfing *P. aeruginosa*

population **(Fig. 2C)**.

Biosurfactants not only reduce surface tension, but their properties like charge and toxicity

can contribute to their roles as antimicrobials, inhibitors of biofilm formation, and emulsifiers [89].

For example, different classes of *P. aeruginosa* secreted surfactants mono- and di-rhamnolipids

and their precursor 3-(3-hydroxyalkanoyloxy) alkanoic acids (HAAs) have opposing roles in

coordinating tendril formation and motility during swarming [90]. In our experiments, we saw that

in contrast to non-ionic Triton X-100 and anionic SDS, addition of the cationic detergent CTAB

inhibited motility **(Fig. 5A)** [91], suggesting that charge could impact swarming and surfing. Taken

together, we posit that biophysical characteristics of surfactants allow emergent motility, but their

exact chemical attributes may alter the specifics of surfing.

Along with physical factors, nutrients such as carbon and nitrogen sources as well as iron

availability also affect *P. aeruginosa* swarming [34, 41, 92, 93]. Transcriptional analysis of surfers

compared to swarming or nonmotile cells, respectively, did not show changes in motility related

genes. In conjunction with the observation that known major transcriptional regulators of motility

are not required for surfing **(Supp. Fig. 5 and 6)**, this suggests that the transition to surfing may

not be regulated at the transcriptional level and may possibly be due to sensing of altered surface

properties directly by the flagella [77].

There was an enrichment in heme biosynthesis and T3SS pathways in the downregulated

genes in surfing cells. Further, on hard agar, transcripts of the phage φCTX were upregulated,

and visible plaques were present especially in Las-deficient cells undergoing surfing on both semi-

solid and hard agar **(Supp. Fig. 5A and B)**. The T3SS is known to be regulated by a complex

network involving both cyclic AMP and cyclic di-GMP signaling [63], where it is thought to be

induced by low Ca^{2+} and host cell contact, and downregulated in a number of stress conditions,

including DNA damage [61, 62]. DNA damage and other stresses also trigger prophage induction

[64], suggesting that *P. aeruginosa* undergoing emergent motility may be sensing cellular stress.

Heme biosynthesis genes and neighboring genes involved in nitrite reduction are induced by the

anaerobic transcriptional regulator Dnr in conditions of low oxygen, as part of the nitrate

respiration pathway [65]. Downregulation of these genes during surfing compared to swarming or

non-motile cells suggests that swarming and non-motile cells are likely transitioning to anaerobic

metabolism while surfers may still be engaging in oxidative respiration.

Some of the transcriptional responses seen in surfing cells are likely due to sensing

unrelated *S. aureus* exoproducts. The upregulation of genes involved in TCA uptake and acetoin

catabolism when surfing was also observed in our previous study in planktonic *P. aeruginosa*

cells, where we determined the upregulation is a specific response to *S. aureus* secreted citrate

and acetoin, respectively [57]. Thus, these staphylococcal intermediate metabolites are sensed

by *P. aeruginosa* irrespective of the mode of growth and motility. Citrate and acetoin are likely

not sufficient to induce surfing since the Agr-deficient *S. aureus* supernatant has similar levels of

these nutrients as the WT [57], and still did not enable motility **(Supp. Fig. 11A and D)**, but it is

unclear how these or other nutrients may affect *P. aeruginosa* surfing. Additionally, viscosity,

surface hardness, and osmolarity have been previously associated with altering *P. aeruginosa*

motility [78, 91]. We observed surfing on both semi-solid and hard agar, a condition in which *P.*

aeruginosa is nonmotile, however the effect of other physical parameters on surfing are yet to be

characterized. While a diverse class of surfactants all transition *P. aeruginosa* to surfing surface

motility, it is likely that the combination of physical and chemical cues from the host, neighboring

microbial species, and the environment influence this motility and determine the resulting niche

expansion and fitness changes in natural habitats.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Bacterial strains used in this study are listed in **Supp. Table 6**. *P. aeruginosa* UCBPP-PA14 [94],

S. aureus JE2 [95], their derivatives, and other species were grown in a modified M63 medium

[57] containing 1× M63 salts (13.6 g·L^{-1} KH$_2$PO$_4$, 2 g·L^{-1} (NH$_4$)$_2$SO$_4$, 0.8 µM ferric citrate, 1 mM

MgSO$_4$; pH adjusted to 7.0 with KOH) supplemented with 0.3% glucose, 1× ACGU solution

(Teknova), 1× supplement EZ (Teknova), 0.1 ng·L^{-1} biotin, and 2 ng·L^{-1} nicotinamide, at 37°C,

with shaking at 300 rpm. *V. cholerae* was grown in M63 with 2% NaCl added. *S. aureus* and *P.*

aeruginosa clinical isolates from the Cystic Fibrosis Foundation Isolate Core were selected at

random from four different patients. *P. aeruginosa* and *S. aureus* transposon mutants were

sourced from the PA14NR Set [96] and the Nebraska Transposon Mutant Library (NTML) [95],

respectively, and confirmed by PCR using primers flanking the transposon insertion listed in

Supp. Table 7. For cloning and mutant construction, strains were cultured in Luria Bertani (Miller)

broth or on agar plates with 15 g·L^{-1} agar supplemented with 50 µg·mL^{-1} gentamicin and/or 25

µg·mL^{-1} irgasan as needed for selection.

Preparation of cell-free supernatant.

Overnight cultures of bacterial strains were diluted to OD$_{600}$ of 0.05 in fresh media and grown for

24 h in flasks before harvesting at 4000 rpm for 20 minutes. The supernatants were filter-sterilized

with a Steriflip with a 0.2 µm polyethersulfone filter (MilliporeSigma). Supernatants were stored at

-30°C until use.

Agar plate motility assays.

LB agar plates (for twitch plates) or M63 agar plates (for twitch, swim, swarm, and hard agar

plates) containing the specified percentage of agar (twitch: 1.5%; swim: 0.3%; swarm: 0.5%; hard:

1.5%) and 25% (v/v) media salts, media salts with the indicated additives, or the indicated species'

supernatant were prepared the day of the assay. For the twitch assays, M63 plates were used for

experiments shown in **Supp. Fig. 1B**, and LB plates for all the other experiments. Freshly poured

plates were allowed to rest at room temperature for 4 h before inoculation with the specified

bacterial species. Plates were inoculated as follows: twitch, *P. aeruginosa* colony stabbed through

the agar to the bottom of the plate; swim, *P. aeruginosa* overnight culture stabbed halfway through

the agar; swarm and hard agar, 2 µL of *P. aeruginosa* overnight culture or mixed culture (equal

volumes of both strains being mixed) was spotted on the center of the plate. Plates were then

incubated for 24 h at 37°C in a single layer of petri dishes in a Mini Low Temperature Incubator

(Fisherbrand) or for the twitch plates, for 48 h at 30°C. When noted, Type II mucin from porcine

stomach (Sigma Aldrich), Triton X-100 (Fisher), sodium dodecyl sulfate (SDS) (Fisher

BioReagents), hexadecyltrimethylammonium bromide (CTAB) (Sigma Aldrich), surfactin from *B.*

subtilis (Sigma Aldrich), rhamnolipids from *P. aeruginosa* (Sigma Aldrich), Mannosylerythritol

Lipid A (MEL-A) from *Pseudozyma* yeasts (Cayman), lactonic sophorolipid (Sigma Aldrich), LL-

37 (Peptide Sciences), 1,2-Dipalmitoyl-sn-glycero-3-phosphocholine (DPPC) (Sigma Aldrich),

and saponin from *Quillaja* bark (Sigma Aldrich) were added at the indicated concentrations to

25% media salts. After incubation, all plates were imaged either using a Chemidoc (Bio-Rad) on

the white tray using the Coomassie blue setting or the black tray using the Cy2 (for GFP) or Cy3

(for dsRed or mKO) settings with optimal exposure, or for the twitch plates, using an Epson

Perfection V700 Photo scanner with bottom illumination. All experiments were repeated three

times.

Construction of *P. aeruginosa* mutants.

Bacterial strains and plasmids are listed in **Supp. Table 6**. For mutant construction, homologous

downstream and upstream arms of genes of interest were amplified using the primers listed in

Supp. Table 7. The amplified fragments were cloned into pDONRPEX18Gm *attP* sites using the

Gateway BP Clonase II Enzyme mix (ThermoFisher). Plasmids were transformed into *E. coli* S17-

1 λ-pir and confirmed by sequencing prior to conjugation. Conjugants were streaked onto LB

plates (without NaCl) + 10% sucrose, and then tested for gentamicin resistance. Gentamicin-

sensitive strains were tested for the deletion by PCR and sequencing.

RNA extraction and library preparation.

P. aeruginosa was scraped after 17 h from 1.5% or 0.5% agar M63 plates containing 25% (v/v)

salts or *S. aureus* cell-free supernatant, resuspended in 2 mL salts mixed with 4 mL of RNAprotect

Bacteria Reagent (Qiagen), and then incubated for 5 min at room temperature for stabilization

before centrifugation at 4000 rpm for 10 min. Supernatants were completely removed before

storage of the pellets at -80°C. RNA was extracted using the Total RNA Purification Plus Kit

(Norgen) according to the manufacturer's instructions for Gram-negative bacteria. The extracted

RNA was subjected to an additional genomic DNA removal by DNase I treatment in solution using

the TURBO DNA-free Kit (Invitrogen) and checked by PCR for the absence of contaminating

DNA. Integrity of the RNA preparation was confirmed by running on an agarose gel and observing

intact rRNA bands. Next, rRNA was removed using the *P. aeruginosa* riboPOOLs rRNA Depletion

Kit (siTOOLs Biotech) followed by library preparation with the NEBNext Ultra II Directional RNA

Library Prep Kit for Illumina (New England Biolabs). The sequencing was performed at the Center

for Cancer Research (CCR) Genomics Core Facility. Two biological replicates were collected.

RNA-seq analysis.

The sequencing files were processed with Cutadapt [97] and Trimmomatic [98]. Alignment to the

P. aeruginosa UCBPP-PA14 genome (NCBI) and pairwise comparisons were made using

Rockhopper (Wellesley College) [99, 100]. Gene expression was set at a minimum baseline value

of 4 and p values of 0 were changed to the lowest values recorded within a dataset. Ribosomal

RNAs and predicted RNAs were removed from each dataset. Upregulated and downregulated

genes were based on transcripts that had $p < 0.05$ and $\log_2$ fold change ≥ 1 or ≤ -1. The location

of prophages in the genome of *P. aeruginosa* UCBPP-PA14 NC_008463.1 was determined using

PHASTER [101, 102]. Venn diagrams were generated using matplotlib_venn package with venn3

using Python [103].

Gene ontology (GO) enrichment analysis.

For *P. aeruginosa* PA14 pathway analysis, the open reading frame designations for the

corresponding PAO1 orthologs of *P. aeruginosa* UCBPP-PA14 genes were obtained using the

Pseudomonas Genome Database (www.pseudomonas.com/rbbh/pairs) [104]. The list of

designations from the RNA-seq analysis were analyzed at the Gene Ontology Resource

(www.geneontology.org) by the PANTHER Overrepresentation Test (released 20221013 and

20231017, as noted) (Annotation Version and Release Date: GO Ontology database DOI:

10.5281/zenodo.7942786 Released 2023-05-10) for enriched biological processes by Fisher's

exact test with Bonferroni correction. All genes identified in the GO enrichment categories 'heme

biosynthesis' and 'T3SS' as well as genes in the same or related neighboring operons, or known

to be co-regulated, that were downregulated in both semi-solid and hard agar conditions, are

shown in the heatmap in **Fig. 3D**.

Accession number(s). The RNA-seq data have been deposited at NCBI Gene Expression

Omnibus (GEO) (https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE249024.

ACKNOWLEDGMENTS

We would like to acknowledge the Center for Cancer Research (CCR) Genomics Core for RNA-

sequencing and whole-genome sequencing, the Cystic Fibrosis Foundation Isolate Core for

providing clinical isolate strains, and the Adhya, Otto, and Ramamurthi labs at NIH, the Brinsmade

lab at Georgetown University, and the Limoli lab at the University of Iowa for kindly providing

bacterial strains. This work utilized the computational resources of the NIH High Performance

Computing Biowulf Cluster (http://hpc.nih.gov). The diagram in Figure 6 was created with

Biorender (http://biorender.com). We thank Susan Gottesman, Gisela Storz, Anthony Martini, and

Kalinga Pavan Thushara Silva for comments on the manuscript, and members of the Khare,

Gottesman, Storz, and Ramamurthi labs for helpful discussion and feedback. This work was

supported by the Intramural Research Program of the NIH, National Cancer Institute, Center for

Cancer Research. TMZ was supported by a Postdoctoral Research Associate Training (PRAT)

Fellowship award 1FI2GM137843-01 from the National Institute of General Medical Sciences.

COMPETING INTERESTS

No competing interests declared.

FIGURES

Figure 1. *S. aureus* secreted products induce *P. aeruginosa* emergent motility in a dose-

dependent manner. *P. aeruginosa* was inoculated on **(A)** semi-solid or **(B)** hard agar plates

containing the indicated percentages of media salts control or *S. aureus* supernatant, and plates

were imaged after 24 hours incubation. Representative images of three independent replicates

are shown. Additional replicates are shown in **Extended Figure 1**.

Figure 2. Emergent motility requires flagella, but not pili or rhamnolipids. The indicated

strains of **(A,D,E)** *P. aeruginosa* or **(B)** *S. aureus* were inoculated on **(A,B,D,E)** semi-solid agar

plates containing 25% media salts control or *S. aureus* supernatant as indicated. **(C)** From left

to right, hard agar plates were inoculated with *P. aeruginosa* WT-mKO mixed with WT-GFP, *P.*

aeruginosa WT-mKO mixed with Δ*fliC*-GFP, and *P. aeruginosa* WT-GFP mixed with *S. aureus*

WT-dsRed. **(A-E)** Motility was imaged after 24 hours incubation. Representative images of three

independent replicates are shown. Additional replicates are shown in **Extended Figure 2**.

Figure 3. *P. aeruginosa* differentially regulates genes involved in T3SS, and heme

biosynthesis, but not motility, during emergent motility on *S. aureus* secreted products.

Transcript levels were compared between *P. aeruginosa* cells exposed to *S. aureus* supernatant

and those exposed to media salts control on semi-solid and hard agar after 17 hours. **(A)** Venn

diagrams of upregulated and downregulated genes on hard and semi-solid agar, as well as in

planktonic *P. aeruginosa* cells after *S. aureus* supernatant exposure compared to media salts

control [57]. **(B)** Scatter plot of $\log_2$(fold-change) transcript levels on hard agar compared to the

$\log_2$(fold-change) of transcript levels on semi-solid agar. Differentially regulated genes in

pathways enriched in the common downregulated genes, as well as in TCA uptake, acetoin

catabolism, and φCTX are shown. **(C)** GO enrichment of *P. aeruginosa* genes differentially

expressed on both semi-solid and hard agar plates, but not in planktonic phase [58-60].

Nonredundant categories are shown. **(D)** Heatmap showing the $\log_2$(fold-change) of genes

associated with enriched pathways for heme biosynthesis and T3SS on semi-solid and hard

agar.

Figure 4. Interspecies secreted surfactants induce emergent motility in *P. aeruginosa*.

(A-E) *P. aeruginosa* was inoculated on semi-solid agar plates containing 25% media salts

control or supernatant from the indicated species: *S. aureus* (JE2), *P. aeruginosa* (PA14), *B.*

subtilis, K. pneumoniae, V. cholerae, E. coli (MG1655)*,* and *B. cenocepacia*. Supernatant was

from **(A, B)** WT strains and **(C-E)** mutants for the biosynthesis of surfactants. Images were

taken after 24 hours incubation. Representative images of three independent replicates are

shown. Additional replicates are shown in **Extended Figure 4**.

Figure 5. The addition of diverse biotic and synthetic surfactants is sufficient to induce

emergent motility in *P. aeruginosa*. (A,B) *P. aeruginosa* was inoculated on **(top)** semi-solid or

(bottom) hard agar plates containing **(A)** (from left to right) 25% media salts control or *S.*

aureus supernatant, or 25% media salts with the addition of 50 µg·mL⁻¹ rhamnolipids, 5 µg·mL⁻¹

surfactin, 0.4% mucin, 0.1% SDS, 0.1% CTAB, or 0.1% Triton X-100; or **(B)** 25% media salts

control or *S. aureus* supernatant; or 25% media salts with the addition of 25 µg·mL⁻¹ MEL-A, 25

µg·mL⁻¹ sophorolipid, 25 µg·mL⁻¹ saponin, 100 µg·mL⁻¹ LL-37, or 100 µg·mL⁻¹ DPPC. **(A,B)**

Images were taken after 24 hours incubation. Representative images of three independent

replicates are shown. Additional replicates are shown in **Extended Figure 5**.

Figure 6. The Las system is required for surfing motility on mucin, but not on

surfactants. (A) *P. aeruginosa* WT and Δ*lasR* were inoculated on semi-solid agar plates

containing (from left to right) 25% media salts control or supernatant from the indicated species:

S. aureus (JE2), *P. aeruginosa* (PA14), or *B. subtilis* (ZK3814); or 25% media salts with the

addition of 0.1% Triton X-100 or 0.4% mucin. Images were taken after 24 hours incubation.

Representative images of three independent replicates are shown. Additional replicates are in

Extended Figure 6. **(B)** Working model of *P. aeruginosa* motility in response to exogenous

surfactants. The glycopeptide mucin, PSMs secreted from *S. aureus*, and other biotic

surfactants secreted by human cells, plants, yeast, and bacteria, as well as synthetic

surfactants, are sensed by *P. aeruginosa.* In response, *P. aeruginosa* exhibits flagella-

640   dependent emergent motility on surfaces. In contrast to the motility in the presence of

641   surfactants, mucin-mediated surfing motility requires the LasR quorum sensing system in *P.*

642   *aeruginosa.*

REFERENCES

1.   Jean-Pierre F, Hampton TH, Schultz D, Hogan DA, Groleau MC, Deziel E, et al. Community composition shapes microbial-specific phenotypes in a cystic fibrosis polymicrobial model system. Elife. 2023;12:e81604. Epub 20230120. doi: 10.7554/eLife.81604. PubMed PMID: 36661299; PubMed Central PMCID: PMCPMC9897730.

2.   Groenhagen U, Baumgartner R, Bailly A, Gardiner A, Eberl L, Schulz S, et al. Production of bioactive volatiles by different *Burkholderia ambifaria* strains. J Chem Ecol. 2013;39(7):892-906. Epub 2013/07/09. doi: 10.1007/s10886-013-0315-y. PubMed PMID: 23832658.

3.   Trizna EY, Yarullina MN, Baidamshina DR, Mironova AV, Akhatova FS, Rozhina EV, et al. Bidirectional alterations in antibiotics susceptibility in *Staphylococcus aureus‐Pseudomonas aeruginosa* dual-species biofilm. Sci Rep. 2020;10(1):14849. Epub 2020/09/11. doi: 10.1038/s41598-020-71834-w. PubMed PMID: 32908166; PubMed Central PMCID: PMCPMC7481796.

4.   Shank EA, Kolter R. New developments in microbial interspecies signaling. Curr Opin Microbiol. 2009;12(2):205-14. Epub 2009/03/03. doi: 10.1016/j.mib.2009.01.003. PubMed PMID: 19251475; PubMed Central PMCID: PMCPMC2709175.

5.   Shank EA, Klepac-Ceraj V, Collado-Torres L, Powers GE, Losick R, Kolter R. Interspecies interactions that result in *Bacillus subtilis* forming biofilms are mediated mainly by members of its own genus. Proc Natl Acad Sci U S A. 2011;108(48):E1236-43. Epub 2011/11/15. doi: 10.1073/pnas.1103630108. PubMed PMID: 22074846; PubMed Central PMCID: PMCPMC3228442.

6.   Zhou Y, Smith D, Leong BJ, Brannstrom K, Almqvist F, Chapman MR. Promiscuous cross-seeding between bacterial amyloids promotes interspecies biofilms. J Biol Chem. 2012;287(42):35092-103. Epub 20120813. doi: 10.1074/jbc.M112.383737. PubMed PMID: 22891247; PubMed Central PMCID: PMCPMC3471717.

7.   Scoffield JA, Duan D, Zhu F, Wu H. A commensal streptococcus hijacks a *Pseudomonas aeruginosa* exopolysaccharide to promote biofilm formation. PLoS Pathog. 2017;13(4):e1006300. Epub 2017/04/28. doi: 10.1371/journal.ppat.1006300. PubMed PMID: 28448633; PubMed Central PMCID: PMCPMC5407764.

8.   Jones SE, Ho L, Rees CA, Hill JE, Nodwell JR, Elliot MA. *Streptomyces* exploration is triggered by fungal interactions and volatile signals. Elife. 2017;6:4268–73. Epub 2017/01/04. doi: 10.7554/eLife.21738. PubMed PMID: 28044982; PubMed Central PMCID: PMCPMC5207766.

9.   Kim KS, Lee S, Ryu CM. Interspecific bacterial sensing through airborne signals modulates locomotion and drug resistance. Nat Commun. 2013;4:1809. Epub 2013/05/09. doi: 10.1038/ncomms2789. PubMed PMID: 23651997.

10.   Lloyd DG, Whitworth DE. The Myxobacterium *Myxococcus xanthus* can sense and respond to the quorum signals secreted by potential prey organisms. Front Microbiol.

2017;8:439. Epub 2017/03/30. doi: 10.3389/fmicb.2017.00439. PubMed PMID: 28352265;
PubMed Central PMCID: PMCPMC5348527.

11.   Khare A. Experimental systems biology approaches reveal interaction mechanisms in
model multispecies communities. Trends Microbiol. 2021. Epub 2021/04/19. doi:
10.1016/j.tim.2021.03.012. PubMed PMID: 33865676.

12.   Pierce EC, Dutton RJ. Putting microbial interactions back into community contexts. Curr
Opin Microbiol. 2022;65:56-63. Epub 2021/11/06. doi: 10.1016/j.mib.2021.10.008. PubMed
PMID: 34739927.

13.   McCully LM, Bitzer AS, Seaton SC, Smith LM, Silby MW. Interspecies social spreading:
interaction between two sessile soil bacteria leads to emergence of surface motility. Msphere.
2019;4(1):e00696–18. Epub 2019/02/01. doi: 10.1128/mSphere.00696-18. PubMed PMID:
30700513; PubMed Central PMCID: PMCPMC6354810.

14.   Hagai E, Dvora R, Havkin-Blank T, Zelinger E, Porat Z, Schulz S, et al. Surface-motility
induction, attraction and hitchhiking between bacterial species promote dispersal on solid
surfaces. ISME J. 2014;8(5):1147-51. Epub 2013/12/07. doi: 10.1038/ismej.2013.218. PubMed
PMID: 24304675; PubMed Central PMCID: PMCPMC3996695.

15.   Buch PJ, Chai Y, Goluch ED. Treating polymicrobial infections in chronic diabetic
wounds. Clin Microbiol Rev. 2019;32(2):e00091. Epub 2019/01/18. doi: 10.1128/CMR.00091-
18. PubMed PMID: 30651226; PubMed Central PMCID: PMCPMC6431129.

16.   Bisht K, Baishya J, Wakeman CA. *Pseudomonas aeruginosa* polymicrobial interactions
during lung infection. Curr Opin Microbiol. 2020;53:1-8. Epub 2020/02/18. doi:
10.1016/j.mib.2020.01.014. PubMed PMID: 32062024; PubMed Central PMCID:
PMCPMC7244363.

17.   Crone S, Vives-Florez M, Kvich L, Saunders AM, Malone M, Nicolaisen MH, et al. The
environmental occurrence of *Pseudomonas aeruginosa*. APMIS. 2020;128(3):220-31. Epub
2019/11/12. doi: 10.1111/apm.13010. PubMed PMID: 31709616.

18.   Malhotra S, Hayes D, Jr., Wozniak DJ. Cystic fibrosis and *Pseudomonas aeruginosa*:
the host-microbe interface. Clin Microbiol Rev. 2019;32(3):e00138-18. Epub 2019/05/31. doi:
10.1128/CMR.00138-18. PubMed PMID: 31142499; PubMed Central PMCID:
PMCPMC6589863.

19.   Foundation CF. Cystic Fibrosis Foundation patient registry 2018 annual data report.
Cystic Fibrosis Foundation, 2018.

20.   Sagel SD, Gibson RL, Emerson J, McNamara S, Burns JL, Wagener JS, et al. Impact of
Pseudomonas and *Staphylococcus* infection on inflammation and clinical status in young
children with cystic fibrosis. J Pediatr. 2009;154(2):183-8. Epub 2008/09/30. doi:
10.1016/j.jpeds.2008.08.001. PubMed PMID: 18822427; PubMed Central PMCID:
PMCPMC2654617.

21.   Maliniak ML, Stecenko AA, McCarty NA. A longitudinal analysis of chronic MRSA and
Pseudomonas aeruginosa co-infection in cystic fibrosis: a single-center study. J Cyst Fibros.
2016;15(3):350-6. doi: 10.1016/j.jcf.2015.10.014. PubMed PMID: 26610860.

22.   Limoli DH, Yang J, Khansaheb MK, Helfman B, Peng L, Stecenko AA, et al.
Staphylococcus aureus and *Pseudomonas aeruginosa* co-infection is associated with cystic
fibrosis-related diabetes and poor clinical outcomes. Eur J Clin Microbiol Infect Dis.
2016;35(6):947-53. Epub 2016/03/20. doi: 10.1007/s10096-016-2621-0. PubMed PMID:
26993289.

23.   Hubert D, Reglier-Poupet H, Sermet-Gaudelus I, Ferroni A, Le Bourgeois M, Burgel PR,
et al. Association between *Staphylococcus aureus* alone or combined with *Pseudomonas*
aeruginosa and the clinical condition of patients with cystic fibrosis. J Cyst Fibros.
2013;12(5):497-503. Epub 2013/01/08. doi: 10.1016/j.jcf.2012.12.003. PubMed PMID:
23291443.

24.   Hotterbeekx A, Kumar-Singh S, Goossens H, Malhotra-Kumar S. *In vivo* and *in vitro*
interactions between *Pseudomonas aeruginosa* and *Staphylococcus* spp. Front Cell Infect
Microbiol. 2017;7:106. doi: 10.3389/fcimb.2017.00106. PubMed PMID: 28421166; PubMed
Central PMCID: PMCPMC5376567.

25.   Overhage J, Bains M, Brazas MD, Hancock RE. Swarming of *Pseudomonas aeruginosa*
is a complex adaptation leading to increased production of virulence factors and antibiotic
resistance. J Bacteriol. 2008;190(8):2671-9. Epub 2008/02/05. doi: 10.1128/JB.01659-07.
PubMed PMID: 18245294; PubMed Central PMCID: PMCPMC2293252.

26.   Tremblay J, Deziel E. Gene expression in *Pseudomonas aeruginosa* swarming motility.
BMC Genomics. 2010;11:587. Epub 2010/10/22. doi: 10.1186/1471-2164-11-587. PubMed
PMID: 20961425; PubMed Central PMCID: PMCPMC3091734.

27.   Kazmierczak BI, Schniederberend M, Jain R. Cross-regulation of *Pseudomonas* motility
systems: the intimate relationship between flagella, pili and virulence. Curr Opin Microbiol.
2015;28:78-82. Epub 2015/10/20. doi: 10.1016/j.mib.2015.07.017. PubMed PMID: 26476804;
PubMed Central PMCID: PMCPMC4688086.

28.   Drake D, Montie TC. Flagella, motility and invasive virulence of *Pseudomonas*
aeruginosa. J Gen Microbiol. 1988;134(1):43-52. Epub 1988/01/01. doi: 10.1099/00221287-134-
1-43. PubMed PMID: 3141566.

29.   Feldman M, Bryan R, Rajan S, Scheffler L, Brunnert S, Tang H, et al. Role of flagella in
pathogenesis of *Pseudomonas aeruginosa* pulmonary infection. Infect Immun. 1998;66(1):43-
51. Epub 1998/01/10. doi: 10.1128/IAI.66.1.43-51.1998. PubMed PMID: 9423837; PubMed
Central PMCID: PMCPMC107856.

30.   Arora SK, Neely AN, Blair B, Lory S, Ramphal R. Role of motility and flagellin
glycosylation in the pathogenesis of *Pseudomonas aeruginosa* burn wound infections. Infect
Immun. 2005;73(7):4395-8. Epub 2005/06/24. doi: 10.1128/IAI.73.7.4395-4398.2005. PubMed
PMID: 15972536; PubMed Central PMCID: PMCPMC1168557.

31.   Coleman SR, Pletzer D, Hancock REW. Contribution of swarming motility to
dissemination in a *Pseudomonas aeruginosa* murine skin abscess infection model. J Infect Dis.
2021;224(4):726-33. Epub 2020/12/23. doi: 10.1093/infdis/jiaa778. PubMed PMID: 33349847;
PubMed Central PMCID: PMCPMC8366438.

32.   Ha DG, Kuchma SL, O'Toole GA. Plate-based assay for swimming motility in
Pseudomonas aeruginosa. Methods in molecular biology. 2014;1149:59-65. Epub 2014/05/14.
doi: 10.1007/978-1-4939-0473-0_7. PubMed PMID: 24818897; PubMed Central PMCID:
PMCPMC9007281.

33.   Burrows LL. *Pseudomonas aeruginosa* twitching motility: type IV pili in action. Annu Rev
Microbiol. 2012;66:493-520. Epub 2012/07/04. doi: 10.1146/annurev-micro-092611-150055.
PubMed PMID: 22746331.

34.   Köhler T, Curty LK, Barja F, van Delden C, Pechère JC. Swarming of *Pseudomonas
aeruginosa* is dependent on cell-to-cell signaling and requires flagella and pili. J Bacteriol.
2000;182(21):5990-6. Epub 2000/10/13. doi: 10.1128/JB.182.21.5990-5996.2000. PubMed
PMID: 11029417; PubMed Central PMCID: PMCPMC94731.

35.   Caiazza NC, Shanks RM, O'Toole GA. Rhamnolipids modulate swarming motility
patterns of *Pseudomonas aeruginosa*. J Bacteriol. 2005;187(21):7351-61. Epub 2005/10/21.
doi: 10.1128/JB.187.21.7351-7361.2005. PubMed PMID: 16237018; PubMed Central PMCID:
PMCPMC1273001.

36.   Yeung AT, Parayno A, Hancock RE. Mucin promotes rapid surface motility in
Pseudomonas aeruginosa. mBio. 2012;3(3):e00073-12. Epub 2012/05/03. doi:
10.1128/mBio.00073-12. PubMed PMID: 22550036; PubMed Central PMCID:
PMCPMC3569861.

37.   Morrison CB, Markovetz MR, Ehre C. Mucus, mucins, and cystic fibrosis. Pediatr
Pulmonol. 2019;54 Suppl 3(Suppl 3):S84-S96. Epub 2019/11/13. doi: 10.1002/ppul.24530.
PubMed PMID: 31715083; PubMed Central PMCID: PMCPMC6853602.

38.   Sun E, Liu S, Hancock REW. Surfing motility: a conserved yet diverse adaptation among
motile bacteria. J Bacteriol. 2018;200(23):e00394. Epub 2018/09/19. doi: 10.1128/JB.00394-18.
PubMed PMID: 30224438; PubMed Central PMCID: PMCPMC6222206.

39.   McAlester G, O'Gara F, Morrissey JP. Signal-mediated interactions between
Pseudomonas aeruginosa and *Candida albicans*. J Med Microbiol. 2008;57(Pt 5):563-9. Epub
2008/04/26. doi: 10.1099/jmm.0.47705-0. PubMed PMID: 18436588.

40.   Lewis KA, Baker AE, Chen AI, Harty CE, Kuchma SL, O'Toole GA, et al. Ethanol
decreases *Pseudomonas aeruginosa* flagellar motility through the regulation of flagellar stators.
J Bacteriol. 2019;201(18):e00285-19. Epub 2019/05/22. doi: 10.1128/JB.00285-19. PubMed
PMID: 31109994; PubMed Central PMCID: PMCPMC6707923.

41.   Badal D, Jayarani AV, Kollaran MA, Prakash D, P M, Singh V. Foraging signals promote
swarming in starving *Pseudomonas aeruginosa*. mBio. 2021;12(5):e0203321. Epub 2021/10/06.
doi: 10.1128/mBio.02033-21. PubMed PMID: 34607460; PubMed Central PMCID:
PMCPMC8546858.

42.   Limoli DH, Warren EA, Yarrington KD, Donegan NP, Cheung AL, O'Toole GA.
Interspecies interactions induce exploratory motility in *Pseudomonas aeruginosa*. Elife.
2019;8:e47365. doi: 10.7554/eLife.47365. PubMed PMID: 31713513; PubMed Central PMCID:
PMCPMC6910820.

43.   Yarrington KD, Shendruk, T. N., Limoli, D. H. Twitching cells use a chemoreceptor to
detect bacterial competitors. *bioRxiv*. 2022. doi: 10.1101/2022.11.28.518211.

44.   Guo Q, Kong W, Jin S, Chen L, Xu Y, Duan K. PqsR-dependent and PqsR-independent
regulation of motility and biofilm formation by PQS in *Pseudomonas aeruginosa* PAO1. J Basic
Microbiol. 2014;54(7):633-43. Epub 2013/09/03. doi: 10.1002/jobm.201300091. PubMed PMID:
23996096.

45.   Jimenez PN, Koch G, Thompson JA, Xavier KB, Cool RH, Quax WJ. The multiple
signaling systems regulating virulence in *Pseudomonas aeruginosa*. Microbiol Mol Biol Rev.
2012;76(1):46-65. Epub 2012/03/07. doi: 10.1128/MMBR.05007-11. PubMed PMID: 22390972;
PubMed Central PMCID: PMCPMC3294424.

46.   Jyot J, Ramphal, R. Flagella and pili of *Pseudomonas aeruginosa*. In: Rehm BHA, editor.
Pseudomonas: Model Organism, Pathogen, Cell Factory: Wiley Online Books; 2008. p. 85-108.

47.   Whitchurch CB, Mattick JS. Characterization of a gene, *pilU*, required for twitching
motility but not phage sensitivity in *Pseudomonas aeruginosa*. Mol Microbiol. 1994;13(6):1079-
91. Epub 1994/09/01. doi: 10.1111/j.1365-2958.1994.tb00499.x. PubMed PMID: 7854122.

48.   Bertrand JJ, West JT, Engel JN. Genetic analysis of the regulation of type IV pilus
function by the Chp chemosensory system of *Pseudomonas aeruginosa*. J Bacteriol.
2010;192(4):994-1010. Epub 2009/12/17. doi: 10.1128/JB.01390-09. PubMed PMID: 20008072;
PubMed Central PMCID: PMCPMC2812951.

49.   Kearns DB, Robinson J, Shimkets LJ. *Pseudomonas aeruginosa* exhibits directed
twitching motility up phosphatidylethanolamine gradients. J Bacteriol. 2001;183(2):763-7. Epub
2001/01/03. doi: 10.1128/JB.183.2.763-767.2001. PubMed PMID: 11133973; PubMed Central
PMCID: PMCPMC94935.

50.   Kilmury SL, Burrows LL. Type IV pilins regulate their own expression via direct
intramembrane interactions with the sensor kinase PilS. Proc Natl Acad Sci U S A.
2016;113(21):6017-22. Epub 2016/05/11. doi: 10.1073/pnas.1512947113. PubMed PMID:
27162347; PubMed Central PMCID: PMCPMC4889343.

51.   Kilmury SLN, Burrows LL. The *Pseudomonas aeruginosa* PilSR two-component system
regulates both twitching and swimming motilities. mBio. 2018;9(4):e01310-18. Epub 2018/07/26.
doi: 10.1128/mBio.01310-18. PubMed PMID: 30042200; PubMed Central PMCID:
PMCPMC6058289.

52.   Schuster M, Greenberg EP. A network of networks: quorum-sensing gene regulation in
Pseudomonas aeruginosa. Int J Med Microbiol. 2006;296(2-3):73-81. Epub 2006/02/16. doi:
10.1016/j.ijmm.2006.01.036. PubMed PMID: 16476569.

53.   Shao X, Yao C, Ding Y, Hu H, Qian G, He M, et al. The transcriptional regulators of
virulence for *Pseudomonas aeruginosa*: Therapeutic opportunity and preventive potential of its
clinical infections. Genes Dis. 2023;10(5):2049-63. Epub 2023/07/26. doi:
10.1016/j.gendis.2022.09.009. PubMed PMID: 37492705; PubMed Central PMCID:
PMCPMC10363592.

54.   Romling U, Galperin MY, Gomelsky M. Cyclic di-GMP: the first 25 years of a universal
bacterial second messenger. Microbiol Mol Biol Rev. 2013;77(1):1-52. Epub 2013/03/09. doi:
10.1128/MMBR.00043-12. PubMed PMID: 23471616; PubMed Central PMCID:
PMCPMC3591986.

55.   Guvener ZT, Harwood CS. Subcellular location characteristics of the *Pseudomonas
aeruginosa* GGDEF protein, WspR, indicate that it produces cyclic-di-GMP in response to
growth on surfaces. Mol Microbiol. 2007;66(6):1459-73. Epub 2007/11/22. doi: 10.1111/j.1365-
2958.2007.06008.x. PubMed PMID: 18028314; PubMed Central PMCID: PMCPMC4105145.

56.   Dubern JF, Romero M, Mai-Prochnow A, Messina M, Trampari E, Gijzel HN, et al. ToxR
is a c-di-GMP binding protein that modulates surface-associated behaviour in *Pseudomonas
aeruginosa*. NPJ Biofilms Microbiomes. 2022;8(1):64. Epub 2022/08/19. doi: 10.1038/s41522-
022-00325-9. PubMed PMID: 35982053; PubMed Central PMCID: PMCPMC9388670.

57.   Zarrella TM, Khare A. Systematic identification of molecular mediators of interspecies
sensing in a community of two frequently coinfecting bacterial pathogens. PLoS Biol.
2022;20(6):e3001679. Epub 2022/06/22. doi: 10.1371/journal.pbio.3001679. PubMed PMID:
35727825; PubMed Central PMCID: PMCPMC9249247.

58.   Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. PANTHER version 14: more
genomes, a new PANTHER GO-slim and improvements in enrichment analysis tools. Nucleic
Acids Res. 2019;47(D1):D419-D26. doi: 10.1093/nar/gky1038. PubMed PMID: 30407594;
PubMed Central PMCID: PMCPMC6323939.

59.   Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, et al. Gene Ontology:
tool for the unification of biology. The Gene Ontology Consortium. Nat Genet. 2000;25(1):25-9.
doi: 10.1038/75556. PubMed PMID: 10802651; PubMed Central PMCID: PMCPMC3037419.

60.   Gene Ontology C. The Gene Ontology resource: enriching a GOld mine. Nucleic Acids
Res. 2021;49(D1):D325-D34. doi: 10.1093/nar/gkaa1113. PubMed PMID: 33290552; PubMed
Central PMCID: PMCPMC7779012.

61.   Wu W, Jin S. PtrB of *Pseudomonas aeruginosa* suppresses the Type III Secretion
System under the stress of DNA damage. J Bacteriol. 2005;187(17):6058-68. Epub 2005/08/20.
doi: 10.1128/JB.187.17.6058-6068.2005. PubMed PMID: 16109947; PubMed Central PMCID:
PMCPMC1196158.

62.   Yahr TL, Wolfgang MC. Transcriptional regulation of the *Pseudomonas aeruginosa* Type
III Secretion System. Mol Microbiol. 2006;62(3):631-40. Epub 2006/09/26. doi: 10.1111/j.1365-
2958.2006.05412.x. PubMed PMID: 16995895.

63.   Williams McMackin EA, Djapgne L, Corley JM, Yahr TL. Fitting pieces into the puzzle of
Pseudomonas aeruginosa type III secretion system gene expression. J Bacteriol.
2019;201(13):e00209-19. Epub 2019/04/24. doi: 10.1128/JB.00209-19. PubMed PMID:
31010903; PubMed Central PMCID: PMCPMC6560140.

64.   Silpe JE, Duddy OP, Bassler BL. Induction mechanisms and strategies underlying
interprophage competition during polylysogeny. PLoS Pathog. 2023;19(5):e1011363. Epub
2023/05/18. doi: 10.1371/journal.ppat.1011363. PubMed PMID: 37200239; PubMed Central
PMCID: PMCPMC10194989.

65.   Trunk K, Benkert B, Quack N, Munch R, Scheer M, Garbe J, et al. Anaerobic adaptation
in *Pseudomonas aeruginosa*: definition of the Anr and Dnr regulons. Environ Microbiol.
2010;12(6):1719-33. Epub 2010/06/18. doi: 10.1111/j.1462-2920.2010.02252.x. PubMed PMID:
20553552.

66.   Sharma J, Sundar D, Srivastava P. Biosurfactants: potential agents for controlling
cellular communication, motility, and antagonism. Front Mol Biosci. 2021;8:727070. Epub
2021/10/29. doi: 10.3389/fmolb.2021.727070. PubMed PMID: 34708073; PubMed Central
PMCID: PMCPMC8542798.

67.   Tsompanidou E, Sibbald MJ, Chlebowicz MA, Dreisbach A, Back JW, van Dijl JM, et al.
Requirement of the *agr* locus for colony spreading of *Staphylococcus aureus*. J Bacteriol.
2011;193(5):1267-72. Epub 2010/12/21. doi: 10.1128/JB.01276-10. PubMed PMID: 21169484;
PubMed Central PMCID: PMCPMC3067592.

68.   Tsompanidou E, Denham EL, Becher D, de Jong A, Buist G, van Oosten M, et al.
Distinct roles of phenol-soluble modulins in spreading of *Staphylococcus aureus* on wet
surfaces. Appl Environ Microbiol. 2013;79(3):886-95. Epub 2012/11/28. doi:
10.1128/AEM.03157-12. PubMed PMID: 23183971; PubMed Central PMCID:
PMCPMC3568538.

69.   Otto M. Phenol-soluble modulins. Int J Med Microbiol. 2014;304(2):164-9. Epub
2014/01/23. doi: 10.1016/j.ijmm.2013.11.019. PubMed PMID: 24447915; PubMed Central
PMCID: PMCPMC4014003.

70.   Abdel-Mawgoud AM, Lepine F, Deziel E. Rhamnolipids: diversity of structures, microbial
origins and roles. Appl Microbiol Biotechnol. 2010;86(5):1323-36. Epub 2010/03/26. doi:
10.1007/s00253-010-2498-2. PubMed PMID: 20336292; PubMed Central PMCID:
PMCPMC2854365.

71.   Piewngam P, Zheng Y, Nguyen TH, Dickey SW, Joo HS, Villaruz AE, et al. Pathogen
elimination by probiotic *Bacillus* via signalling interference. Nature. 2018;562(7728):532-7. Epub
2018/10/12. doi: 10.1038/s41586-018-0616-y. PubMed PMID: 30305736; PubMed Central
PMCID: PMCPMC6202238.

72.   Sun E, Gill EE, Falsafi R, Yeung A, Liu S, Hancock REW. Broad-spectrum adaptive
antibiotic resistance associated with *Pseudomonas aeruginosa* mucin-dependent surfing
motility. Antimicrob Agents Chemother. 2018;62(9):e00848-18. Epub 2018/07/04. doi:
10.1128/AAC.00848-18. PubMed PMID: 29967020; PubMed Central PMCID:
PMCPMC6125533.

73.   Pletzer D, Sun E, Ritchie C, Wilkinson L, Liu LT, Trimble MJ, et al. Surfing motility is a
complex adaptation dependent on the stringent stress response in *Pseudomonas aeruginosa*
LESB58. PLoS Pathog. 2020;16(3):e1008444. Epub 2020/03/26. doi:
10.1371/journal.ppat.1008444. PubMed PMID: 32208458; PubMed Central PMCID:
PMCPMC7122816.

74.   Morin CD, Deziel E. Use of alternative gelling agents reveals the role of rhamnolipids in
Pseudomonas aeruginosa surface motility. Biomolecules. 2021;11(10):1468. Epub 2021/10/24.
doi: 10.3390/biom11101468. PubMed PMID: 34680106; PubMed Central PMCID:
PMCPMC8533327.

924 75. Khan F, Pham DTN, Oloketuyi SF, Kim YM. Regulation and controlling the motility
925 properties of *Pseudomonas aeruginosa*. Appl Microbiol Biotechnol. 2020;104(1):33-49. Epub
926 2019/11/27. doi: 10.1007/s00253-019-10201-w. PubMed PMID: 31768614.

927 76. Bru JL, Rawson B, Trinh C, Whiteson K, Høyland-Kroghsbo NM, Siryaporn A. PQS
928 produced by the *Pseudomonas aeruginosa* stress response repels swarms away from
929 bacteriophage and antibiotics. J Bacteriol. 2019;201(23):10.1128/jb.00383-19. Epub
930 2019/08/28. doi: 10.1128/JB.00383-19. PubMed PMID: 31451543; PubMed Central PMCID:
931 PMCPMC6832071.

932 77. Belas R. Biofilms, flagella, and mechanosensing of surfaces by bacteria. Trends
933 Microbiol. 2014;22(9):517-27. Epub 2014/06/05. doi: 10.1016/j.tim.2014.05.002. PubMed PMID:
934 24894628.

935 78. Kamatkar NG, Shrout JD. Surface hardness impairment of quorum sensing and
936 swarming for *Pseudomonas aeruginosa*. PLoS One. 2011;6(6):e20888. Epub 2011/06/21. doi:
937 10.1371/journal.pone.0020888. PubMed PMID: 21687741; PubMed Central PMCID:
938 PMCPMC3110244.

939 79. Fauvart M, Phillips P, Bachaspatimayum D, Verstraeten N, Fransaer J, Michiels J, et al.
940 Surface tension gradient control of bacterial swarming in colonies of. Soft Matter. 2012;8(1):70-
941 6. doi: 10.1039/c1sm06002c. PubMed PMID: WOS:000297561400011.

942 80. Bru JL, Kasallis SJ, Chang R, Zhuo Q, Nguyen J, Pham P, et al. The great divide:
943 rhamnolipids mediate separation between *P. aeruginosa* and *S. aureus*. Front Cell Infect
944 Microbiol. 2023;13:1245874. Epub 2023/10/02. doi: 10.3389/fcimb.2023.1245874. PubMed
945 PMID: 37780859; PubMed Central PMCID: PMCPMC10540625.

946 81. Kearns DB. A field guide to bacterial swarming motility. Nat Rev Microbiol.
947 2010;8(9):634-44. Epub 2010/08/10. doi: 10.1038/nrmicro2405. PubMed PMID: 20694026;
948 PubMed Central PMCID: PMCPMC3135019.

949 82. Kearns DB, Losick R. Swarming motility in undomesticated *Bacillus subtilis*. Mol
950 Microbiol. 2003;49(3):581-90. Epub 2003/07/17. doi: 10.1046/j.1365-2958.2003.03584.x.
951 PubMed PMID: 12864845.

952 83. Huber B, Riedel K, Hentzer M, Heydorn A, Gotschlich A, Givskov M, et al. The *cep*
953 quorum-sensing system of *Burkholderia cepacia* H111 controls biofilm formation and swarming
954 motility. Microbiology (Reading). 2001;147(Pt 9):2517-28. Epub 2001/09/06. doi:
955 10.1099/00221287-147-9-2517. PubMed PMID: 11535791.

956 84. Soberón-Chávez G, Lépine F, Déziel E. Production of rhamnolipids by *Pseudomonas
957 aeruginosa*. Appl Microbiol Biot. 2005;68(6):718-25. doi: 10.1007/s00253-005-0150-3. PubMed
958 PMID: WOS:000232856700002.

959 85. Lindum PW, Anthoni U, Christophersen C, Eberl L, Molin S, Givskov M. N-Acyl-L-
960 homoserine lactone autoinducers control production of an extracellular lipopeptide biosurfactant
961 required for swarming motility of *Serratia liquefaciens* MG1. J Bacteriol. 1998;180(23):6384-8.
962 Epub 1998/11/26. doi: 10.1128/JB.180.23.6384-6388.1998. PubMed PMID: 9829950; PubMed
963 Central PMCID: PMCPMC107727.

86.   Schreiber L, Schlechter RO, Kunzler M, Remus-Emsermann MNP. Hitching a ride in the
phyllosphere: Surfactant production of *Pseudomonas* spp. causes co-swarming of *Pantoea*
eucalypti 299R. bioRxiv. 2023:2023.11. 07.566084.

87.   Bernier SP, Hum C, Li X, O'Toole GA, Magarvey NA, Surette MG. *Pseudomonas*
aeruginosa-derived rhamnolipids and other detergents modulate colony morphotype and motility
in the *Burkholderia cepacia* complex. J Bacteriol. 2017;199(13):e00171-17. Epub 2017/04/26.
doi: 10.1128/JB.00171-17. PubMed PMID: 28439038; PubMed Central PMCID:
PMCPMC5472820.

88.   Morin C, Landry M, Groleau MC, Deziel E. Surface motility favors codependent
interaction between *Pseudomonas aeruginosa* and *Burkholderia cenocepacia*. Msphere. 2022.
doi: 10.1128/msphere.00153-22. PubMed PMID: WOS:000821730500001.

89.   Pardhi DS, Panchal RR, Raval VH, Joshi RG, Poczai P, Almalki WH, et al. Microbial
surfactants: a journey from fundamentals to recent advances. Front Microbiol. 2022;13:982603.
Epub 2022/08/23. doi: 10.3389/fmicb.2022.982603. PubMed PMID: 35992692; PubMed Central
PMCID: PMCPMC9386247.

90.   Tremblay J, Richardson AP, Lepine F, Deziel E. Self-produced extracellular stimuli
modulate the *Pseudomonas aeruginosa* swarming motility behaviour. Environ Microbiol.
2007;9(10):2622-30. Epub 2007/09/07. doi: 10.1111/j.1462-2920.2007.01396.x. PubMed PMID:
17803784.

91.   Yang A, Tang WS, Si T, Tang JX. Influence of physical effects on the swarming motility
of *Pseudomonas aeruginosa*. Biophys J. 2017;112(7):1462-71. Epub 2017/04/14. doi:
10.1016/j.bpj.2017.02.019. PubMed PMID: 28402888; PubMed Central PMCID:
PMCPMC5389960.

92.   Inoue T, Shingaki R, Fukui K. Inhibition of swarming motility of *Pseudomonas*
aeruginosa by branched-chain fatty acids. FEMS Microbiol Lett. 2008;281(1):81-6. Epub
2008/03/06. doi: 10.1111/j.1574-6968.2008.01089.x. PubMed PMID: 18318842.

93.   Mattingly AE, Weaver AA, Dimkovikj A, Shrout JD. Assessing travel conditions:
environmental and host influences on bacterial surface motility. J Bacteriol.
2018;200(11):e00014-18. Epub 2018/03/21. doi: 10.1128/JB.00014-18. PubMed PMID:
29555698; PubMed Central PMCID: PMCPMC5952383.

94.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. Common
virulence factors for bacterial pathogenicity in plants and animals. Science.
1995;268(5219):1899-902. doi: 10.1126/science.7604262. PubMed PMID: 7604262.

95.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, et al. A genetic
resource for rapid and comprehensive phenotype screening of nonessential *Staphylococcus*
aureus genes. MBio. 2013;4(1):e00537-12. doi: 10.1128/mBio.00537-12. PubMed PMID:
23404398; PubMed Central PMCID: PMCPMC3573662.

96.   Liberati NT, Urbach JM, Miyata S, Lee DG, Drenkard E, Wu G, et al. An ordered,
nonredundant library of Pseudomonas aeruginosa strain PA14 transposon insertion mutants.
Proc Natl Acad Sci U S A. 2006;103(8):2833-8. Epub 2006/02/16. doi:

10.1073/pnas.0511100103. PubMed PMID: 16477005; PubMed Central PMCID:
PMCPMC1413827.

97.   Martin M. Cutadapt removes adapter sequences from high-throughput sequencing
reads. EMBnet J. 2011;17:10-2. doi: 10.14806/ej.17.1.200.

98.   Bolger AM, Lohse M, Usadel B. Trimmomatic: a flexible trimmer for Illumina sequence
data. Bioinformatics. 2014;30(15):2114-20. doi: 10.1093/bioinformatics/btu170. PubMed PMID:
24695404; PubMed Central PMCID: PMCPMC4103590.

99.   McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA, et al.
Computational analysis of bacterial RNA-seq data. Nucleic Acids Res. 2013;41(14):e140. Epub
2013/05/30. doi: 10.1093/nar/gkt444. PubMed PMID: 23716638; PubMed Central PMCID:
PMCPMC3737546.

100.   Tjaden B. *De novo* assembly of bacterial transcriptomes from RNA-seq data. Genome
Biol. 2015;16:1. Epub 2015/01/15. doi: 10.1186/s13059-014-0572-2. PubMed PMID: 25583448;
PubMed Central PMCID: PMCPMC4316799.

101.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. PHAST: a fast phage search tool.
Nucleic Acids Res. 2011;39(Web Server issue):W347-52. Epub 2011/06/16. doi:
10.1093/nar/gkr485. PubMed PMID: 21672955; PubMed Central PMCID: PMCPMC3125810.

102.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, et al. PHASTER: a better, faster
version of the PHAST phage search tool. Nucleic Acids Res. 2016;44(W1):W16-21. Epub
2016/05/05. doi: 10.1093/nar/gkw387. PubMed PMID: 27141966; PubMed Central PMCID:
PMCPMC4987931.

103.   matplotlib_venn [Internet]. 2020 [cited 2020]. Available from:
https://github.com/konstantint/matplotlib-venn.

104.   Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. Enhanced annotations
and features for comparing thousands of *Pseudomonas genomes* in the *Pseudomonas* genome
database. Nucleic Acids Res. 2016;44(D1):D646-53. doi: 10.1093/nar/gkv1227. PubMed PMID:
26578582; PubMed Central PMCID: PMCPMC4702867.

1031