TITLE

Interspecies surfactants serve as public goods enabling surface motility in *Pseudomonas*

aeruginosa

AUTHORS

Delayna L. Warrell[a]*, Tiffany M. Zarrella[a,b,c#]* , Christopher Machalek[a], Anupama Khare[a#]

[a]Laboratory of Molecular Biology, Center for Cancer Research, National Cancer Institute, National

Institutes of Health, Bethesda, MD, USA

[b]Postdoctoral Research Associate Training Program, National Institute of General Medical

Sciences, National Institutes of Health, Bethesda, MD, USA

[c]Current address: Department of Biology, Georgetown University, Washington, DC, USA

Running head: Surfactants enable motility in *P. aeruginosa*

[#]Address correspondence to Tiffany M. Zarrella, tiffany.zarrella@georgetown.edu, or Anupama

Khare, anupama.khare@nih.gov

*Delayna L. Warrell and Tiffany M. Zarrella contributed equally to this work. Author order was

determined alphabetically.

Key words: *Pseudomonas aeruginosa*, *Staphylococcus aureus*, surfactants, motility,

polymicrobial interactions, phenol-soluble modulins

The authors declare no conflict of interest.

ABSTRACT

In most natural environments, bacteria live in polymicrobial communities where secreted

molecules from neighboring species alter bacterial behaviors including motility, but such

interactions are understudied. *Pseudomonas aeruginosa* is a motile opportunistic pathogen that

exists in diverse multispecies environments such as the soil and is frequently found in human

wound and respiratory tract co-infections with other bacteria including *Staphylococcus aureus*.

Here we show that *P. aeruginosa* can co-opt secreted surfactants from other species for flagellar-

based surface motility. We found that exogenous surfactants from *S. aureus*, other bacteria, and

interkingdom species enabled *P. aeruginosa* to switch from swarming to an alternative surface

spreading motility on semi-solid surfaces and allowed for the emergence of surface motility on

hard agar where *P. aeruginosa* was otherwise unable to move. This motility was distinct from the

response of other motile bacteria in the presence of exogenous surfactants. Mutant analysis

indicated that this *P. aeruginosa* motility was similar to a previously described mucin-based

motility, 'surfing', albeit with divergent regulation. Thus, our study demonstrates that secreted

surfactants from the host as well as neighboring bacterial and interkingdom species act as public

goods facilitating *P. aeruginosa* flagella-mediated surfing-like surface motility, thereby allowing it

to access different environmental niches.

INTRODUCTION

Bacteria are often found in dynamic, complex communities where interspecies secreted factors influence bacterial behaviors, such as antibiotic resistance (1-3), biofilm formation (4-7), and motility (8-10). However, many of the polymicrobial interactions leading to the modification of these pathogenic traits have not been elucidated. The study of pairwise interactions has the potential to reveal behavioral changes that occur within communities as well as the underlying molecular mechanisms (11, 12).

Motility allows bacterial cells to disperse and migrate to new niches for nutrients, seek out prey, and adapt to new environments, and modulation of motility behaviors due to ecological interactions is thus likely to be advantageous for bacterial fitness. Several examples of changes in bacterial motility in mixed cultures compared to monocultures have been previously reported, including the inhibition of motility via secreted volatiles (9), and emergence of motility in co-culture conditions where the monocultures did not move. Such emergent motility includes social spreading between two non-motile species (13), exploration by non-motile bacteria upon sensing fungal volatiles (8), and non-motile bacteria inducing motility in a neighboring species, and then hitchhiking on the moving population (14).

Pseudomonas aeruginosa is a social, motile microbe that typically inhabits polymicrobial environments in soil, wounds, and persistent infections of the respiratory tracts of people with cystic fibrosis (CF) (15-17). Motility in *P. aeruginosa* is intricately linked with virulence regulation, antibiotic resistance, and biofilm formation (18-20), and flagellar competency is associated with dissemination and pathogenesis (21-24). *P. aeruginosa* displays several well-established forms of motility: swimming, twitching, and swarming. During swimming motility *P. aeruginosa* uses a single polar flagellum to propel through liquid (25), while for twitching, type IV pili extension and retraction are required for movement on hard surfaces (26). Swarming is a social behavior with a characteristic tendril formation on semi-solid surfaces that requires flagella and the secretion of rhamnolipids as a surfactant, while pili also contribute (27, 28). Another type of motility that has

been described is 'surfing', where *P. aeruginosa* spreads over semi-solid media containing the

glycoprotein mucin, that is abundant in the mucus accumulated in the CF airways and

gastrointestinal tract (29, 30). This motility is thought to depend on the viscous nature and

lubricant properties of mucin and requires *P. aeruginosa* flagellar function while pili and

rhamnolipids are expendable (29, 31). *P. aeruginosa* also exhibits sliding motility which requires

only the production of rhamnolipids, but no motility appendages (32). Therefore, physical

parameters are an important cue for *P. aeruginosa*, where the viscosity of the environment

dictates the type of motility, with liquid and low viscosity environments facilitating swimming, and

hard surfaces inducing twitching.

P. aeruginosa motility is affected by interspecies interactions. Fungal volatiles farnesol

and ethanol at high concentrations inhibit *P. aeruginosa* motility, while lower concentrations of

ethanol induce swarming when carbon sources are limited, possibly signaling the availability of

nutrients (33-35). Further, *P. aeruginosa* is commonly found in concert with *Staphylococcus*

aureus in wound infections and nosocomial pneumonia, and these two species are the most

frequently isolated bacterial pathogens from the airways of people with CF (36). It has been

observed that *P. aeruginosa* engages in directional pili-based 'exploratory motility' via the Pil-Chp

chemosensory system in response to the *S. aureus* secreted cytotoxins phenol-soluble modulins

(PSMs) (37, 38).

In this study, we investigate whether additional *P. aeruginosa* motility behaviors are

altered by *S. aureus* exoproducts. We show that the presence of surfactants from *S. aureus*, the

soil bacterium *Bacillus subtilis*, and interkingdom species, such as fungi, plants, and humans, as

well as synthetic surfactants enables a flagellar-based surface spreading motility in *P. aeruginosa*.

Bacterial surface motility has been classified into four types: swarming, twitching, sliding, and

gliding (39), and we demonstrate that surface spreading facilitated by exogenous surfactants is

an additional distinct type of surface motility. Flagellar-based surface spreading in the presence

of exogenous surfactants is not a general bacterial response, but specific to *P. aeruginosa*. This

motility occurs on semi-solid as well as hard agar surfaces, where *P. aeruginosa* is otherwise

non-motile, thereby demonstrating an emergent phenotype by co-opting already existing

interspecies surfactants from the environment. Finally, a comparison with mucin-mediated surfing

suggests that this surfactant-facilitated surface spreading is a surfing-like motility, although only

movement on mucin requires a quorum sensing regulator.

RESULTS

***P. aeruginosa* displays emergent surface spreading in the presence of *S. aureus* secreted**

products.

To determine the effect of *S. aureus* secreted products on *P. aeruginosa* motility, we

performed *P. aeruginosa* twitching, swimming, and swarming assays in the presence of 25% (v/v)

S. aureus cell-free supernatant, or media salts as a control. As reported previously (37), we

observed that *P. aeruginosa* twitched farther with the addition of *S. aureus* supernatant **(Supp.**

Fig. 1A). In the swimming assay, while *P. aeruginosa* swam uniformly creating a circular swim

area in the control agar, the swim patterns were irregular and oval-shaped in the presence of *S.*

aureus supernatant **(Supp. Fig. 1B)**.

In a standard swarm assay on semi-solid agar plates (containing 0.5% agar) *P. aeruginosa*

swarmed with characteristic tendril formation on the control plate **(Fig. 1A)**. However, on plates

with 25% *S. aureus* supernatant, *P. aeruginosa* exhibited a motility unlike swarming where the

bacteria spread out across the plate, and then showed some swarming-like tendril formation **(Fig.**

1A). *P. aeruginosa* exhibited similar motility on agar containing 12.5% *S. aureus* supernatant and

an intermediate motility on 5% supernatant, while swarming resumed on plates with 1%

supernatant **(Fig. 1A)**. Next, we tested if this phenotype occurs on hard agar plates (containing

1.5% agar) on which *P. aeruginosa* is traditionally unable to move. While *P. aeruginosa* did not

move on the hard agar control or on the plates containing 5% and 1% *S. aureus* supernatant **(Fig.**

1B), it exhibited emergent surface spreading on the plates containing 25% and 12.5% *S. aureus*

supernatant **(Fig. 1B)**. There was no tendril formation on hard agar, likely because the tendrils

are due to swarming after surface spreading. Thus, *S. aureus* secreted products enable a distinct

form of *P. aeruginosa* motility, even in conditions where *P. aeruginosa* is normally nonmotile.

To determine the generality of this motility, we performed the surface spreading assay

with clinical isolates of both species. We tested four clinical isolates of *P. aeruginosa* collected

from different people with CF (CF017, CF033, CF057, and CF095) **(Supp. Table 1)** on both semi-

solid and hard agar and observed that the isolates CF017 and CF057 swarmed and exhibited

surface spreading to varying degrees, while isolates CF033 and CF095 did not **(Supp. Fig. 2A**

and B). To determine if these isolates also had other motility deficiencies, we tested these isolates

for swimming and twitching. Isolate CF017 swam similar to *P. aeruginosa* PA14, while isolates

CF033, CF057, CF095 all had swimming deficiencies **(Supp. Fig. 3A)**. Additionally, the twitch

zone of CF033 was similar to PA14, but CF017 and CF057 had smaller twitch zones and CF095

did not twitch **(Supp. Fig. 3B)**. Therefore, the isolates that do not exhibit the emergent motility

may also have flagellar and/or pili deficiencies. These *P. aeruginosa* strains were co-isolated with

four *S. aureus* strains, CF019, CF032, CF058, and CF100, respectively **(Supp. Table 1)**. Cell-

free supernatants from each of these *S. aureus* clinical isolates could facilitate this motility in *P.*

aeruginosa PA14 **(Supp. Fig. 4A and B)***.* These results demonstrate that while there is some

variation, clinical strains of *P. aeruginosa* and *S. aureus* can exhibit and enable surface spreading,

respectively*.*

Interspecies surfactants facilitate surface spreading in *P. aeruginosa.*

To identify the molecules that enable surface spreading in *P. aeruginosa*, we investigated

whether other species promote this motility, and found that *P. aeruginosa* also exhibited similar

motility on plates with supernatant from *P. aeruginosa* and *Bacillus subtilis* ZK3814 **(Fig. 2A)** but

swarmed on plates with supernatant from *B. subtilis* PY79, *Klebsiella pneumoniae, Vibrio*

cholerae, Escherichia coli, Burkholderia cenocepacia BAA245, and *B. cenocepacia* 25608 **(Fig.**

2A and B).

A common class of molecules previously implicated in motility, and known to be secreted

by each of the inducing strains, but none of the non-inducing strains, is biosurfactants (40). *S.*

aureus produces phenol soluble modulins (PSMs) (41-43), *P. aeruginosa* produces rhamnolipids

(44), and *B. subtilis* ZK3814 produces fengycin and surfactin (45). We therefore tested the

supernatant of mutants deficient in each of these biosurfactants for their ability to facilitate motility

in *P. aeruginosa*. The supernatants of the *S. aureus* JE2 and LAC strains that lack production of

PSMs did not enable motility in *P. aeruginosa* **(Fig. 2C and Supp. Fig. 5A, B, and C)**. Further,

supernatant from *agrA*::tn, *agrB*::tn, or *agrC*::tn, mutants of the *S. aureus agr* quorum sensing

system which is required for the production of PSMs, also did not lead to *P. aeruginosa* surface

spreading **(Supp. Fig. 5A and D)**. Similarly, we found that *P. aeruginosa* did not exhibit this

motility on plates containing supernatant from the *P. aeruginosa* ∆*rhlA* mutant deficient for

rhamnolipids **(Fig. 2D and Supp. Fig. 6A)**. Finally, supernatant from a *B. subtilis* mutant without

surfactin production (∆*srfA*) was unable to promote *P. aeruginosa* surface spreading, unlike the

parental WT *B. subtilis* ZK3814 as well as ∆*fenA*, a mutant deficient for fengycin production **(Fig.**

2E and Supp. Fig. 6B). Collectively, these data show that the production of biosurfactants is

required for the interspecies facilitated surface spreading in *P. aeruginosa*.

Detergents and surfactants are sufficient to facilitate surface spreading in *P. aeruginosa*.

Given that bacterial-secreted surfactants were necessary to facilitate surface spreading in

P. aeruginosa, we investigated if the addition of biological and synthetic surfactants was sufficient

to enable this motility. We tested commercially available biosurfactants from the motility-inducing

species, including rhamnolipids from *P. aeruginosa* and surfactin from *B. subtilis;* mucin, which

has been previously implicated in *P. aeruginosa* surfing motility (31); and the synthetic surfactants

sodium dodecyl sulfate (SDS), cetrimonium bromide (CTAB), and Triton X-100. *P. aeruginosa*

exhibited spreading motility on plates containing all these surfactants to varying degrees, except

CTAB, which inhibited motility **(Fig. 3A)**. Further, we tested biosurfactants from interkingdom

species, such as mannosylerythritol lipid A (MEL-A) and sophorolipid from yeasts, plant-produced

saponin, and human cathelicidin LL-37 and pulmonary surfactant dipalmitoylphosphatidylcholine

(DPPC). All these interkingdom surfactants, except DPPC, caused the spreading motility,

especially on semi-solid agar **(Fig. 3B)**. Thus, diverse biological and synthetic surfactants are

sufficient to enable *P. aeruginosa* surface spreading motility.

***P. aeruginosa* surface spreading requires flagella, but not pili or rhamnolipids.**

P. aeruginosa swarming motility requires endogenous production of the biosurfactant

rhamnolipids, therefore we tested whether surface spreading was similar to swarming, by

comparing their respective genetic requirements. Pili, flagella, and the surfactant rhamnolipids all

contribute to *P. aeruginosa* swarming (27, 28). Additionally, swarming is also affected by quorum

sensing systems and other regulatory pathways (27, 46, 47).

First, we tested several mutants that lack the following structural components of the *P.*

aeruginosa flagella: the proximal rod protein FlgB, the hook protein FlgE, or the flagellin subunit

FliC (48), and are therefore defective in swimming **(Supp. Fig. 7A)** and swarming **(Fig. 4A)**. On

the semi-solid and hard agar plates with *S. aureus* supernatant, the *flgB::*tn, *flgE::*tn, and Δ*fliC*

mutants only slid on the plates without spreading out, unlike the motility exhibited by the WT strain

(Fig. 4A and Supp. Fig. 7B), indicating that flagella are required for surface spreading, and that

sliding is likely passive, and occurs only in the presence of surfactants possibly due to a slight

difference in the leveling of the agar. To examine the motility in a species that does not encode

flagella, we tested *S. aureus* itself, and found that it was immobile on the control agar and slid on

the plates with *S. aureus* supernatant like the *P. aeruginosa* flagellar mutants **(Fig. 4B and Supp.**

Fig. 7C). To determine if these flagella-deficient strains could move in the presence of motile *P.*

aeruginosa, these strains were fluorescently labeled, mixed, and added to hard agar with *S.*

aureus supernatant. As expected, two distinctly labeled *P. aeruginosa* WT strains co-cultured on

hard agar exhibited surface spreading equally **(Fig. 4C)**. However, the Δ*fliC* mutant and *S. aureus*

still only slid on the plate even in co-culture with *P. aeruginosa* WT cells that showed surface

spreading **(Fig. 4C)**, suggesting that this motility is an active process, unlike passive sliding, and

the presence of a motile population cannot compensate for loss of competent flagella. Hence, *P.*

aeruginosa surface spreading in the presence of interspecies surfactants requires active flagellar

function.

We next determined if pili, which are required for twitching and contribute to swarming

(27), are necessary for this motility. We tested multiple twitching-deficient pili mutants **(Supp. Fig.**

8A and B) - two mutants deficient in pili retraction, *pilT*::tn and *pilU*::tn (49), two mutants lacking

the extended pili fiber, Δ*pilB*, via deficiency in pili extension (50), and Δ*pilA* due to the lack of the

pilin subunit (26), and two mutants deficient for pili-mediated chemotaxis, Δ*pilJ* and Δ*pilG* (51).

While these mutants showed differences in swarming, all of these strains, displayed surface

spreading on plates with *S. aureus* supernatant **(Fig. 4D and E, Supp. Fig. 8C and D, Supp. Fig.**

9A). In particular, the *pilT*::tn mutant was unable to swarm, but still showed surface spreading,

albeit without tendril formation, suggesting that the surface spreading is distinct from swarming,

although the tendril formation likely indicates swarming.

Previously, it was demonstrated that *P. aeruginosa* shows passive 'sliding motility' that

does not require any appendages but is impeded by pili (32). In that study, flagellar mutants did

not slide, while pili mutants and a mutant for both pili and flagella showed sliding motility (32). We

therefore tested a Δ*fliC* Δ*pilA* double mutant and found that this mutant only slid on the plates with

S. aureus supernatant and did not demonstrate surface spreading **(Fig. 4E)**. Taken together, *P.*

aeruginosa surface spreading in the presence of *S. aureus* secreted products is an active motility

that requires flagella but not pili.

While swarming, *P. aeruginosa* secretes rhamnolipids, biosurfactants that lubricate the

area and allow the cells to move over the surface (27, 28). We found that while a mutant deficient

in endogenous rhamnolipid production, Δ*rhlA*, was unable to swarm as previously reported (27,

28), it still spread on both semi-solid and hard agar plates containing *S. aureus* cell-free

supernatant **(Fig. 4F and Supp. Fig. 9B)**, suggesting that endogenously produced rhamnolipids

are not required for surface spreading. This mutant did not exhibit tendril formation after the

surface spreading on semi-solid agar, reiterating that the tendrils likely indicate swarming.

Finally, quorum sensing and other motility regulators are known to be important for

swarming (27, 46, 47). *P. aeruginosa* utilizes three quorum sensing systems: Las, Rhl, and PQS

(52). To examine the role of quorum sensing in surface spreading we tested mutant strains lacking

each individually (Δ*rhlR,* Δ*lasR,* Δ*pqsA*)*,* or in combination (Δ*rhlR* Δ*lasR* Δ*pqsA*). The Δ*rhlR,* Δ*lasR,*

and Δ*rhlR* Δ*lasR* Δ*pqsA* mutants did not swarm on the control plates, while the Δ*pqsA* mutant did

(Supp. Fig. 10A). However, all mutants displayed surface spreading on the plates with *S. aureus*

supernatant to varying degrees, with the Δ*pqsA* strain being hyper-motile compared to the other

strains and covering the entire semi-solid agar plate **(Supp. Fig. 10A and B)**. We also tested

mutants from additional regulatory pathways that control motility, including the Gac/Rsm and Wsp

pathways (53-56). The Δ*retS* and Δ*rsmA* mutants did not swarm, while the Δ*gacA,* Δ*wspR,* and

Δ*toxR* mutants exhibited swarming **(Supp. Fig. 11)***.* However, all these regulatory mutants

exhibited surface spreading **(Supp. Fig. 11)**, indicating that these regulatory pathways are not

required for *P. aeruginosa* surface spreading, although quorum sensing may affect motility

patterns*.* Thus, this interspecies surfactant-facilitated motility is distinct from swarming since it

does not require processes that are necessary for swarming, such as pili retraction or quorum

sensing and other regulatory pathways.

Other bacterial species show distinct motility phenotypes in the presence of exogenous surfactants

Given that *P. aeruginosa* requires flagella for surface spreading, and the non-motile *S. aureus* only slides in the presence of exogenous surfactants, we tested other bacterial species like *B. cenocepacia*, *K. pneumoniae*, *V. cholerae*, and *E. coli* on semi-solid plates containing *S. aureus* supernatant, Triton X-100, or saponin **(Supp. Fig. 12)**. All these species, except *K. pneumoniae*, are known to be flagellated (57, 58), and *E. coli* and *B. cenocepacia* demonstrated swimming motility **(Supp Fig. 13A and B)**. While *V. cholerae* showed some surface spreading in the presence of *S. aureus* supernatant, none of the other species showed robust movement beyond sliding. Thus, surface spreading as exhibited by *P. aeruginosa* is not a ubiquitous bacterial motility, even for flagellated swimming-proficient species in the presence of exogenous surfactants.

We also tested *B. subtilis* strains ZK3814 that demonstrated swimming motility (**Supp Fig. 13A**) and produces surfactin and PY79 that does not, on semi-solid agar containing either media salts as a control, or *S. aureus* supernatant, rhamnolipids, or saponin **(Supp. Fig. 14)**. Interestingly, *B. subtilis* ZK3814 swarmed in all these conditions, and did not switch to surface spreading, unlike what is seen in *P. aeruginosa*. Further, *B. subtilis* PY79 was nonmotile on the control plate, but showed similar motility to *B. subtilis* ZK3814 in the presence of exogenous surfactants, possibly preceded by sliding and surface spreading. Therefore, the presence of exogenous surfactants likely restores swarming motility in surfactant-deficient *B. subtilis*, unlike the *P. aeruginosa* rhamnolipid-deficient mutant, which displayed only surface spreading **(Fig. 4F)**.

Cells undergoing surface spreading do not differentially transcriptionally regulate motility pathways.

Given the lack of requirement of all major motility regulators for surface spreading, and the diversity of surfactant molecules that facilitate it, this motility could potentially be regulated by

alternative transcriptional regulators or exclusively by the physical cue of reduced surface tension.

Thus, to test for transcriptional regulation during surface spreading, we performed RNA

sequencing (RNA-seq) on *P. aeruginosa* cells scraped from the leading edge on semi-solid or

hard agar plates containing the media salts control or *S. aureus* supernatant at 17 hours **(Supp.**

Fig. 15A). We found approximately 100 to 350 genes that were upregulated and downregulated

on the supernatant compared to the media salts control in both cases **(Supp. Tables 2 and 3)**.

We reasoned that the differentially regulated genes would include genes specific to this motility

as well as genes that were regulated upon exposure to unrelated *S. aureus* secreted molecules.

To distinguish between generalized responses to *S. aureus* products and motility specific

transcripts, we performed a secondary comparison of both motility conditions to genes we

previously identified as differentially regulated in early log-phase *P. aeruginosa* planktonic

cultures 2 hours after the addition of *S. aureus* supernatant compared to media salts as a control

(59), and found significant overlaps **(Fig. 5A)**. Some of the highest upregulated genes in both

motility conditions as well as in planktonic conditions were genes involved in tricarboxylic acids

(TCA) uptake and acetoin catabolism, likely due to *S. aureus* secreted citrate and acetoin,

respectively (59) **(Fig. 5B, Supp. Fig. 15B and C, Supp. Tables 3, 4, and 5)**. Next, we performed

Gene Ontology (GO) enrichment analysis of the *P. aeruginosa* genes differentially regulated in

the individual motility conditions, both motility conditions (semi-solid and hard agar plates), as well

as those regulated exclusively in both motility conditions, but not in the planktonic phase (60-62)

(Fig. 5C, Supp. Fig. 15D and E, and Supp. Tables 4, 5, and 6). Genes involved in aerobic

respiration and response to heat were enriched in the genes downregulated in both motility

conditions **(Supp. Fig. 15E and Supp. Table 6)**, but these genes were also downregulated in the

planktonic phase and are likely unrelated to the motility **(Supp. Fig. 15E and Supp. Tables 5 and**

6).

The *P. aeruginosa* genes differentially regulated exclusively in both motility conditions, but

not in the planktonic phase, were not enriched for any motility pathways **(Fig. 5C and Supp.**

Table 6), suggesting that transcriptional regulation may not play an important role during surface

spreading, consistent with our results that major motility regulators are not required for this motility

(Supp. Fig. 10 and 11). Instead, genes downregulated exclusively in the motility conditions

showed enrichment of the Type 3 Secretion System (T3SS) and heme biosynthesis pathways,

while the upregulated genes showed no statistically enriched pathways **(Fig. 5C and D, and**

Supp. Table 6). Additionally, on the hard agar there was an upregulation in transcripts from the

phage φCTX that encode the R- and F-type pyocins **(Fig. 5B, Supp. Fig. 15B, and Supp. Table**

3) (63), indicating pyocin induction in this condition. This likely explains the presence of plaque-

like clearings on the hard agar (e.g. in **Fig. 1B**), which was especially apparent for the LasR

mutant **(Supp. Fig. 10)**. Multiple cellular stresses can lead to both induction of prophage and

downregulation of T3SS (64-67), indicating that *P. aeruginosa* senses physiological stress while

undergoing surface spreading. Further, heme biosynthesis and associated denitrification genes

(e.g. *nirS*, *nirQ*, *norC, norB)* are induced in anaerobic conditions (68). The relative downregulation

of these genes during surface spreading may be indicative that the control cells (swarming cells

in the semi-solid agar and non-motile cells in the hard agar) are switching to anaerobic

metabolism, while cells undergoing surface spreading continue with aerobic respiration. These

results indicate that *P. aeruginosa* surface spreading motility is unlikely to be regulated at the

transcriptional level, although cells may experience physiological stress and metabolic changes

while undergoing surface spreading in the presence of *S. aureus* exoproducts.

P. aeruginosa **surfactant-mediated surface spreading is a surfing-like motility.**

We next tested whether interspecies surfactant-facilitated surface spreading is similar to

mucin-induced surfing (29, 31, 69, 70), by comparing the genetic requirements for surface

spreading on semi-solid plates containing *S. aureus* supernatant or mucin. Flagellar and pili

mutants showed similar phenotypes for motility on each, where *flgB*::tn, and Δ*fliC* showed no

motility, while *pilT*::tn and Δ*pilA* moved on both **(Supp. Fig. 16A)**.

Previously, Rhl and Las as well as components of Gac/Rsm were reported to contribute

to mucin-based surfing (29, 69, 70). We observed that while the Δ*pqsA* mutant moved on both

the *S. aureus* and mucin plates, the Δ*rhlR* and Δ*rhlR* Δ*lasR* Δ*pqsA* mutants moved moderately on

S. aureus supernatant and had no movement on the plates with mucin **(Supp. Fig. 16B)**.

Interestingly, the Δ*lasR* mutant moved on the *S. aureus* supernatant but not on mucin,

demonstrating a divergence of quorum sensing requirements for mucin-based surfing (29, 31)

and surfactant-enabled motility. The Δ*retS*, Δ*gacA*, Δ*rsmA*, Δ*wspR*, and Δ*toxR* regulatory mutants

all moved on both *S. aureus* supernatant and mucin to varying degrees **(Supp. Fig. 17)**. To further

delineate the role of LasR in motility on a variety of surfactants, we tested the Δ*lasR* mutant on

semi-solid plates with supernatants from *S. aureus*, *P. aeruginosa*, and *B. subtilis* ZK3814, as

well as with the addition of mucin and the surfactant Triton X-100. While the Δ*lasR* mutant did not

move on mucin, it exhibited surface spreading on all of the other surfactant-containing plates **(Fig.**

6A). Overall, the surfactant-dependent surface spreading we observed, and mucin-dependent

surfing motility have similar appendage and regulatory requirements, indicating that surfactant-

based surface spreading is a surfing-like motility.

DISCUSSION

Bacterial locomotion is an advantageous trait that enables migration to more favorable

environments and access to fresh nutrients. Within microbial communities, interspecies secreted

factors can affect bacterial motility, by activation (8), inhibition (9), or endowing nonmotile strains

the capability to move (13, 14). Many of these observations were reported by examining pairwise

interactions, which can reveal novel behavioral phenomena as well as the underlying mechanisms

(11). Here, we interrogated how *P. aeruginosa* motility is affected by *S. aureus* secreted

molecules and uncovered a surfing-like surface spreading motility that depends on the presence

of exogenous interspecies or synthetic surfactants **(Fig. 6B)**, likely due to the low surface tension

environment provided by the physical properties of these surfactants. The surfactant-based

motility we observed requires flagellar function, but not type IV pili or endogenous production of

rhamnolipids, suggesting it is distinct from previously described motility types in *P. aeruginosa*

(Fig. 6C). It permits the emergence of motility when *P. aeruginosa* is typically nonmotile and is

facilitated by surfactants from a variety of species, including co-infecting pathogens, putative

environmental neighbors, as well as infection hosts, demonstrating that *P. aeruginosa* can take

advantage of pre-existing public goods from nearby species to explore new niches.

We observed that in the presence of exogenous surfactants, *P. aeruginosa* cells spread

over the surface, an active process that required flagellar function. This was followed by swarming

tendril formation on semi-solid agar, or likely dispersal of individual cells on hard agar that resulted

in colonies at the mobile leading edge. The motility was facilitated by the presence of diverse

types of biological and synthetic molecules of a variety of chemical compositions, suggesting that

their shared properties of reducing surface tension and acting as wetting agents led to the motility.

Previously it had been shown that when other gelling agents, the polysaccharides gellan gum and

carrageenan, are used in place of agar in semi-solid media, *P. aeruginosa* exhibits a similar

flagellar-dependent, rhamnolipid-independent surface spreading motility, possibly because the

alternate gelling agents also acted as wetting agents (71). It is possible that the *P. aeruginosa*

polar flagellum directly senses surfactant properties by the alleviation of impediments on flagellar

rotation (72), and transitions cells to surface spreading.

The reduced surface tension due to exogenous surfactants likely allows for increased

movement, resulting in the emergence of motility even on hard agar, where *P. aeruginosa* is

otherwise nonmotile, and enabling surface motility without rhamnolipid production, unlike what is

seen for swarming. Intrinsic rhamnolipid production occurs and is required during swarming, while

the presence of exogenous rhamnolipids leads to the emergent motility phenotype **(Fig. 3A)**.

Given that *P. aeruginosa* rhamnolipid production is quorum-sensing dependent and requires high

cell density, it is likely that in the presence of exogenous surfactants, *P. aeruginosa* cells rapidly

sense the change in environmental conditions resulting in surface spreading motility before they

produce rhamnolipids.

While reduced surface tension likely facilitates *P. aeruginosa* surface spreading via

flagellar function, this is not a general phenotype for bacteria. We observe that *E. coli* and *B.*

cenocepacia strains with functional flagella do not show surface spreading, suggesting that

additional parameters likely affect this motility. It is possible that cell surface or other physical

characteristics of *E. coli* and *B. cenocepacia* prevent surface spreading, unlike what is seen in *P.*

aeruginosa. Further, whereas *P. aeruginosa* switches from swarming to surface spreading in the

presence of exogenous surfactants, *B. subtilis* swarming is unaltered. Additionally, a *B. subtilis*

strain deficient for surfactin production (PY79) resumes swarming in the presence of exogenous

surfactants, unlike what is seen for the *P. aeruginosa* rhamnolipid mutant Δ*rhlA* **(Fig. 4F)**, which

demonstrates only surface spreading. Thus, the *P. aeruginosa* surface spreading in the presence

of exogenous surfactants is a motility response distinct from that displayed by other motile

bacteria.

We systematically examined the requirement of several regulators that control motility

(73), such as components of Pil-Chp, quorum sensing, Gac/Rsm, and Wsp surface sensing, but

none were needed for surfactant-based surface spreading, except for mucin-based surfing

requiring the LasR quorum sensing regulator **(Fig. 6A)** (29, 69). Nonetheless, there were

noticeable differences in the appearance of surface spreading quorum sensing mutants. On semi-

solid agar, the final step of tendril formation was absent in Rhl mutants that are expected to lack

rhamnolipid production, indicating that the tendrils here are swarming, and require intrinsic

production of rhamnolipids. In addition, mutants devoid of the PQS pathway, which plays a role

in swarming tendril repulsion (74), are hyper-motile on semi-solid agar, suggesting that the PQS

pathway may play a role in limiting surfactant-based surface spreading.

P. aeruginosa and *S. aureus* often co-infect wounds and the airways of people with CF.

We observed that the presence of *S. aureus* PSMs, which are amphipathic alpha-helical peptides

with surfactant properties, facilitates surface spreading motility in *P. aeruginosa*. Previously, these

S. aureus secreted toxins have been implicated in enabling motility in both species. In *S. aureus*,

PSMs allow nonmotile *S. aureus* to slide on surfaces (42). Additionally, *P. aeruginosa* senses

PSMs via the Pil-Chp chemosensory system to induce pili-dependent directional 'exploratory

motility' (37, 75), which is a motility distinct from the one we describe. *P. aeruginosa* explorers

require pili and components of the Pil-Chp system, such as the chemoreceptor PilJ (75), which

are not necessary for flagellar-dependent surfactant-based surface motility. Sensing of surface

hardness and viscosity plays a role in stimulating motility behaviors (76, 77), with twitching and

swarming occurring on hard and semi-solid surfaces, respectively **(Fig. 6C)**. However, we

observed that even on hard agar surface spreading still solely required flagella and not pili, in

contrast to the directionality of exploratory motility detected on hard agar (37, 75). PSMs produced

by neighboring *S. aureus* cells are also thought to alter the flow of *P. aeruginosa* rhamnolipids,

leading to repulsion of *P. aeruginosa* swarm tendrils (78). Thus, there is an overarching role for

PSMs in inducing and altering motility in *P. aeruginosa*, and the mode of locomotive response

likely differs based on additional environmental stimuli and/or how these molecules are sensed

by *P. aeruginosa*.

In addition to *S. aureus*-produced PSMs (42), many microbes secrete diverse classes of

biosurfactants that enable locomotion (79), such as the lipopeptide surfactin by *B. subtilis* (80),

the glycolipids rhamnolipids by *P. aeruginosa* and *B. cenocepacia* (81, 82), and the aminolipid

serrawettin by *Serratia* spp. (83). As secreted factors, interspecies surfactants can serve as public

goods and enable other species to co-opt these wetting agents for motility, and we observed that

P. aeruginosa can utilize diverse intra- and interkingdom surfactants for this surfing-like motility.

It has been observed that in conditions where *B. cenocepacia* is individually non-motile (and does

not produce surfactant), it can use *P. aeruginosa* rhamnolipids for motility (84, 85). Further,

flagella-deficient *P. aeruginosa* that produces rhamnolipids is incapable of swarming but can co-

swarm with *B. cenocepacia* that is using the secreted surfactant (85), providing benefits for each

species. Despite this display of hitchhiking on motile *B. cenocepacia*, in our study we did not

observe cooperation or hitchhiking between surface spreading *P. aeruginosa* and non-spreading

flagellar mutants or *S. aureus* **(Fig. 4C)**.

Biosurfactants properties like charge and toxicity can contribute to their roles as signaling

molecules, inhibitors of biofilm formation, and emulsifiers (86). In our experiments, we saw that in

contrast to non-ionic Triton X-100 and anionic SDS, addition of the cationic detergent CTAB

inhibited motility **(Fig. 3A)** (87), suggesting that charge could impact swarming and surface

spreading. Taken together, we posit that biophysical characteristics of surfactants allow surface

spreading, but their exact chemical attributes may alter the specifics of the surface motility.

Viscosity, surface hardness, and osmolarity have been previously associated with altering

P. aeruginosa motility (76, 87). Further, nutrients such as carbon and nitrogen sources as well as

iron availability also affect *P. aeruginosa* swarming (27, 35, 39, 88). We observed surface

spreading on both semi-solid and hard agar, a condition in which *P. aeruginosa* is nonmotile,

however the effect of other physical parameters on this motility are yet to be characterized. While

a diverse class of surfactants all transition *P. aeruginosa* to surfing surface motility, it is likely that

the combination of physical and chemical cues from the host, neighboring microbial species, and

the environment influence this motility and determine the resulting niche expansion and fitness

changes in natural habitats.

MATERIALS AND METHODS

Bacterial strains and growth conditions.

Bacterial strains used in this study are listed in **Supp. Table 1**. *P. aeruginosa* UCBPP-PA14 (89),

S. aureus JE2 (90), their derivatives, and other species were grown in a modified M63 medium

(59) containing $1\times$ M63 salts (13.6 g·L^{-1} KH$_2$PO$_4$, 2 g·L^{-1} (NH$_4$)$_2$SO$_4$, 0.8 μM ferric citrate, 1 mM

MgSO$_4$; pH adjusted to 7.0 with KOH) supplemented with 0.3% glucose, $1\times$ ACGU solution

(Teknova), $1\times$ supplement EZ (Teknova), 0.1 ng·L^{-1} biotin, and 2 ng·L^{-1} nicotinamide, at 37°C,

with shaking at 300 rpm. *V. cholerae* was grown in M63 with 2% NaCl added. *S. aureus* and *P.*

aeruginosa clinical isolates from the Cystic Fibrosis Foundation Isolate Core were selected at

random from four different patients. *P. aeruginosa* and *S. aureus* transposon mutants were

sourced from the PA14NR Set (91) and the Nebraska Transposon Mutant Library (NTML) (90),

respectively, and confirmed by PCR using primers flanking the transposon insertion listed in

Supp. Table 7. For cloning and mutant construction, strains were cultured in Luria Bertani (Miller)

broth or on agar plates with 15 $g \cdot L^{-1}$ agar supplemented with 50 $\mu g \cdot mL^{-1}$ gentamicin and/or 25

$\mu g \cdot mL^{-1}$ irgasan as needed for selection.

Preparation of cell-free supernatant.

462 Overnight cultures of bacterial strains were diluted to OD_{600} of 0.05 in fresh media and grown for

463 24 h in flasks before harvesting at 4000 rpm for 20 minutes. The supernatants were filter-sterilized

464 with a Steriflip with a 0.2 µm polyethersulfone filter (MilliporeSigma). Supernatants were stored at

465 -30°C until use.

Agar plate motility assays.

LB agar plates (for *P. aeruginosa* twitch, and multiple species swim and swarm plates) or M63

agar plates (for *P. aeruginosa* twitch, swim, swarm, and hard agar plates) containing the specified

percentage of agar (twitch: 1.5%; swim: 0.3%; swarm: 0.5%; hard: 1.5%) and 25% (v/v) media

salts, media salts with the indicated additives, or the indicated species' supernatant which were

mixed into the agar medium, were used for the motility assays. For the twitch assays, M63 plates

were used for experiments shown in **Supp. Fig. 1A**, and LB plates for all the other experiments.

For the swim assays, LB plates were used for experiments shown in **Supp. Fig. 13**, and M63

plates for all the other experiments. For the swarm assays, LB plates were used for experiments

shown in **Supp. Fig. 12 and 14**, and M63 plates for all the other experiments. Freshly poured 25

mL plates were allowed to rest at room temperature for 4 h before inoculation with the specified

bacterial species. Twitch plates were prepared prior to the day of the assay. Plates were

inoculated as follows: twitch, *P. aeruginosa* colony stabbed through the agar to the bottom of the

plate; swim, *P. aeruginosa* or indicated species overnight culture stabbed halfway through the

agar; swarm and hard agar, 2 µL of *P. aeruginosa* overnight culture, indicated species, or mixed

culture (equal volumes of both strains being mixed) was spotted on the center of the plate. Plates

were then incubated for 24 h at 37°C in a single layer of petri dishes in a Mini Low Temperature

Incubator (Fisherbrand) for the swarm and surface spreading plates, or for the swim plates, for

24 h or 48 h (as noted) at room temperature, or for the twitch plates, for 48 h at 30°C. When noted,

Type II mucin from porcine stomach (Sigma Aldrich), Triton X-100 (Fisher), sodium dodecyl

sulfate (SDS) (Fisher BioReagents), hexadecyltrimethylammonium bromide (CTAB) (Sigma

Aldrich), surfactin from *B. subtilis* (Sigma Aldrich), rhamnolipids from *P. aeruginosa* (Sigma

Aldrich), Mannosylerythritol Lipid A (MEL-A) from *Pseudozyma* yeasts (Cayman), lactonic

sophorolipid (Sigma Aldrich), LL-37 (Peptide Sciences), 1,2-Dipalmitoyl-sn-glycero-3-

phosphocholine (DPPC) (Sigma Aldrich), and saponin from *Quillaja* bark (Sigma Aldrich) were

added at the indicated concentrations to 25% media salts. After incubation, all plates were imaged

either using a Chemidoc (Bio-Rad) on the white tray using the Coomassie blue setting or the black

tray using the Cy2 (for GFP) or Cy3 (for dsRed or mKO) settings with optimal exposure, or for the

twitch plates, using an Epson Perfection V700 Photo scanner with bottom illumination. All

experiments were repeated three times.

Construction of *P. aeruginosa* mutants.

Bacterial strains and plasmids are listed in **Supp. Table 1**. For mutant construction, homologous

downstream and upstream arms of genes of interest were amplified using the primers listed in

Supp. Table 7. The amplified fragments were cloned into pDONRPEX18Gm *attP* sites using the

Gateway BP Clonase II Enzyme mix (ThermoFisher). Plasmids were transformed into *E. coli* S17-

1 λ-pir and confirmed by sequencing prior to conjugation. Conjugants were streaked onto LB

plates (without NaCl) + 10% sucrose, and then tested for gentamicin resistance. Gentamicin-

sensitive strains were tested for the deletion by PCR and sequencing.

RNA extraction and library preparation.

P. aeruginosa was scraped after 17 h from 1.5% or 0.5% agar M63 plates containing 25% (v/v)

salts or *S. aureus* cell-free supernatant, resuspended in 2 mL salts mixed with 4 mL of RNAprotect

Bacteria Reagent (Qiagen), and then incubated for 5 min at room temperature for stabilization

before centrifugation at 4000 rpm for 10 min. Supernatants were completely removed before

storage of the pellets at -80°C. RNA was extracted using the Total RNA Purification Plus Kit

(Norgen) according to the manufacturer's instructions for Gram-negative bacteria. The extracted

RNA was subjected to an additional genomic DNA removal by DNase I treatment in solution using

the TURBO DNA-free Kit (Invitrogen) and checked by PCR for the absence of contaminating

DNA. Integrity of the RNA preparation was confirmed by running on an agarose gel and observing

intact rRNA bands. Next, rRNA was removed using the *P. aeruginosa* riboPOOLs rRNA Depletion

Kit (siTOOLs Biotech) followed by library preparation with the NEBNext Ultra II Directional RNA

Library Prep Kit for Illumina (New England Biolabs). The sequencing was performed at the Center

for Cancer Research (CCR) Genomics Core Facility. Two biological replicates were collected.

RNA-seq analysis.

The sequencing files were processed with Cutadapt (92) and Trimmomatic (93). Alignment to the

P. aeruginosa UCBPP-PA14 genome (NCBI) and pairwise comparisons were made using

Rockhopper (Wellesley College) (94, 95). Gene expression was set at a minimum baseline value

of 4 and p values of 0 were changed to the lowest values recorded within a dataset. Ribosomal

RNAs and predicted RNAs were removed from each dataset. Upregulated and downregulated

genes were based on transcripts that had $p < 0.05$ and $\log_2$ fold change ≥ 1 or ≤ -1. The location

of prophages in the genome of *P. aeruginosa* UCBPP-PA14 NC_008463.1 was determined using

PHASTER (96, 97). Venn diagrams were generated using matplotlib_venn package with venn3

using Python (98).

Gene ontology (GO) enrichment analysis.

For *P. aeruginosa* PA14 pathway analysis, the open reading frame designations for the

corresponding PAO1 orthologs of *P. aeruginosa* UCBPP-PA14 genes were obtained using the

Pseudomonas Genome Database (www.pseudomonas.com/rbbh/pairs) (99). The list of

designations from the RNA-seq analysis were analyzed at the Gene Ontology Resource

(www.geneontology.org) by the PANTHER Overrepresentation Test (released 20221013 and

20231017, as noted) (Annotation Version and Release Date: GO Ontology database DOI:

10.5281/zenodo.7942786 Released 2023-05-10) for enriched biological processes by Fisher's

exact test with Bonferroni correction. All genes identified in the GO enrichment categories 'heme

biosynthesis' and 'T3SS' as well as genes in the same or related neighboring operons, or known

to be co-regulated, that were downregulated in both semi-solid and hard agar conditions, are

shown in the heatmap in **Fig. 5D**.

Accession number(s). The RNA-seq data have been deposited at NCBI Gene Expression

Omnibus (GEO) (https://www.ncbi.nlm.nih.gov/geo/) under accession number GSE249024.

ACKNOWLEDGMENTS

We would like to acknowledge the Center for Cancer Research (CCR) Genomics Core for RNA-

sequencing and whole-genome sequencing, the Cystic Fibrosis Foundation Isolate Core for

providing clinical isolate strains, and the Adhya, Otto, and Ramamurthi labs at the NIH, the

Brinsmade lab at Georgetown University, the Limoli lab at the University of Iowa, and the Goldberg

lab at the Emory University School of Medicine for kindly providing bacterial strains. This work

utilized the computational resources of the NIH High Performance Computing Biowulf Cluster

(http://hpc.nih.gov). The diagrams in Figure 6 was created with Biorender (http://biorender.com).

We thank Susan Gottesman, Gisela Storz, Anthony Martini, and Kalinga Pavan Thushara Silva

for comments on the manuscript, and members of the Khare, Gottesman, Storz, and Ramamurthi

labs for helpful discussion and feedback. This work was supported by the Intramural Research

Program of the NIH, National Cancer Institute, Center for Cancer Research. TMZ was supported

by a Postdoctoral Research Associate Training (PRAT) Fellowship award 1FI2GM137843-01

from the National Institute of General Medical Sciences.

COMPETING INTERESTS

No competing interests declared.

FIGURES

Figure 1. *S. aureus* secreted products enable *P. aeruginosa* surface spreading in a dose-

dependent manner. *P. aeruginosa* was inoculated on **(A)** semi-solid or **(B)** hard agar plates

containing the indicated percentages of media salts as a control or *S. aureus* supernatant, and

plates were imaged after 24 hours incubation. **(A,B)** Representative images of three independent

replicates are shown. Additional replicates are shown in **Extended Figure 1**.

Figure 2. Interspecies secreted surfactants facilitate surface spreading in *P. aeruginosa*.

(A-E) *P. aeruginosa* was inoculated on semi-solid agar plates containing 25% media salts as a control or supernatant from the indicated species: *S. aureus* (JE2), *P. aeruginosa* (PA14), *B. subtilis, K. pneumoniae, V. cholerae, E. coli* (MG1655)*,* and *B. cenocepacia*. Supernatant was from **(A, B)** WT strains and **(C-E)** mutants for the biosynthesis of surfactants. Images were taken after 24 hours incubation. Representative images of three independent replicates are shown. Additional replicates are shown in **Extended Figure 2**.

Figure 3. The addition of diverse biotic and synthetic surfactants is sufficient to enable surface spreading in *P. aeruginosa*. (A,B) *P. aeruginosa* was inoculated on (top) semi-solid or (bottom) hard agar plates containing **(A)** (from left to right) 25% media salts as a control or *S. aureus* supernatant, or 25% media salts with the addition of 50 μg·mL^{-1} rhamnolipids, 5 μg·mL^{-1} surfactin, 0.4% mucin, 0.1% SDS, 0.1% CTAB, or 0.1% Triton X-100; or **(B)** 25% media salts control or *S. aureus* supernatant; or 25% media salts with the addition of 25 μg·mL^{-1} MEL-A, 25 μg·mL^{-1} sophorolipid, 25 μg·mL^{-1} saponin, 100 μg·mL^{-1} LL-37, or 100 μg·mL^{-1} DPPC. **(A,B)** Images were taken after 24 hours incubation. Representative images of three independent replicates are shown. Additional replicates are shown in **Extended Figure 3**.

Figure 4. Surface spreading requires flagella, but not pili or rhamnolipids. The indicated

strains of **(A,D,E,F)** *P. aeruginosa* or **(B)** *S. aureus* were inoculated on **(A,B,D,E,F)** semi-solid

agar plates containing 25% media salts as a control or *S. aureus* supernatant as indicated. **(C)**

From left to right, *P. aeruginosa* WT-mKO mixed with WT-GFP, *P. aeruginosa* WT-mKO mixed

with Δ*fliC*-GFP, and *P. aeruginosa* WT-GFP mixed with *S. aureus* WT-dsRed were inoculated on

599   hard agar plates containing 25% *S. aureus* supernatant. **(A-F)** Motility was imaged after 24 hours

600   incubation. Representative images of three independent replicates are shown. Additional

601   replicates are shown in **Extended Figure 4**.

Figure 5. *P. aeruginosa* differentially regulates genes involved in T3SS, and heme biosynthesis, but not motility, during surface spreading on *S. aureus* secreted products. Transcript levels were compared between *P. aeruginosa* cells exposed to *S. aureus* supernatant and those exposed to media salts as a control on semi-solid and hard agar after 17 hours. **(A)** Venn diagrams of upregulated and downregulated genes on hard and semi-solid agar, as well as in planktonic *P. aeruginosa* cells after *S. aureus* supernatant exposure compared to media salts

control (59). **(B)** Scatter plot of log_2(fold-change) transcript levels on hard agar compared to the

log_2(fold-change) of transcript levels on semi-solid agar. Differentially regulated genes in

pathways enriched in the common downregulated genes, as well as in TCA uptake, acetoin

catabolism, and φCTX are shown. **(C)** GO enrichment of *P. aeruginosa* genes differentially

expressed in response to *S. aureus* supernatant on both semi-solid and hard agar plates, but not

in planktonic phase (60-62). Nonredundant categories are shown. **(D)** Heatmap showing the

log_2(fold-change) of genes associated with enriched pathways for heme biosynthesis and T3SS

on semi-solid and hard agar.

Figure 6. The Las system is required for surfing motility on mucin, but not on surfactants.

(A) *P. aeruginosa* WT and Δ*lasR* were inoculated on semi-solid agar plates containing (from left

to right) 25% media salts as a control or supernatant from the indicated species: *S. aureus* (JE2),

P. aeruginosa (PA14), or *B. subtilis* (ZK3814); or 25% media salts with the addition of 0.1% Triton

X-100 or 0.4% mucin. Images were taken after 24 hours incubation. Representative images of

three independent replicates are shown. Additional replicates are in **Extended Figure 6**. **(B)**

Working model of *P. aeruginosa* motility in response to exogenous surfactants. The glycopeptide

mucin, PSMs secreted from *S. aureus*, and other biotic surfactants secreted by human cells,

plants, yeast, and bacteria, as well as synthetic surfactants, are co-opted by *P. aeruginosa.* In the

presence of exogenous surfactants, *P. aeruginosa* exhibits flagella-dependent spreading motility

on surfaces. In contrast to surfactant-facilitated surface spreading, mucin-mediated surfing

motility requires the LasR quorum sensing system in *P. aeruginosa*. **(C)** (Left to right) *P.*

aeruginosa swimming motility occurs in liquid conditions and requires flagella (25), twitching

motility occurs on hard surfaces and requires type IV pili (26), swarming motility occurs on semi-

solid surfaces and requires flagella, type IV pili, and rhamnolipids (27, 28), sliding motility occurs

on semi-solid surfaces and endogenous or exogenous surfactants (32), and surface

spreading/surfing motility occurs on semi-solid and hard surfaces and requires flagella and

exogenous surfactants (29, 31, 69, 70).

REFERENCES

1. Jean-Pierre F, Hampton TH, Schultz D, Hogan DA, Groleau MC, Deziel E, O'Toole GA. 2023. Community composition shapes microbial-specific phenotypes in a cystic fibrosis polymicrobial model system. Elife 12:e81604.

2. Groenhagen U, Baumgartner R, Bailly A, Gardiner A, Eberl L, Schulz S, Weisskopf L. 2013. Production of bioactive volatiles by different *Burkholderia ambifaria* strains. J Chem Ecol 39:892-906.

3. Trizna EY, Yarullina MN, Baidamshina DR, Mironova AV, Akhatova FS, Rozhina EV, Fakhrullin RF, Khabibrakhmanova AM, Kurbangalieva AR, Bogachev MI, Kayumov AR. 2020. Bidirectional alterations in antibiotics susceptibility in *Staphylococcus aureus-Pseudomonas aeruginosa* dual-species biofilm. Sci Rep 10:14849.

4. Shank EA, Kolter R. 2009. New developments in microbial interspecies signaling. Curr Opin Microbiol 12:205-14.

5. Shank EA, Klepac-Ceraj V, Collado-Torres L, Powers GE, Losick R, Kolter R. 2011. Interspecies interactions that result in *Bacillus subtilis* forming biofilms are mediated mainly by members of its own genus. Proc Natl Acad Sci U S A 108:E1236-43.

6. Zhou Y, Smith D, Leong BJ, Brannstrom K, Almqvist F, Chapman MR. 2012. Promiscuous cross-seeding between bacterial amyloids promotes interspecies biofilms. J Biol Chem 287:35092-35103.

7. Scoffield JA, Duan D, Zhu F, Wu H. 2017. A commensal streptococcus hijacks a *Pseudomonas aeruginosa* exopolysaccharide to promote biofilm formation. PLoS Pathog 13:e1006300.

8. Jones SE, Ho L, Rees CA, Hill JE, Nodwell JR, Elliot MA. 2017. *Streptomyces* exploration is triggered by fungal interactions and volatile signals. Elife 6:4268–73.

9. Kim KS, Lee S, Ryu CM. 2013. Interspecific bacterial sensing through airborne signals modulates locomotion and drug resistance. Nat Commun 4:1809.

10. Lloyd DG, Whitworth DE. 2017. The Myxobacterium *Myxococcus xanthus* can sense and respond to the quorum signals secreted by potential prey organisms. Front Microbiol 8:439.

11. Khare A. 2021. Experimental systems biology approaches reveal interaction mechanisms in model multispecies communities. Trends Microbiol doi:10.1016/j.tim.2021.03.012.

12. Pierce EC, Dutton RJ. 2022. Putting microbial interactions back into community contexts. Curr Opin Microbiol 65:56-63.

13. McCully LM, Bitzer AS, Seaton SC, Smith LM, Silby MW. 2019. Interspecies social spreading: interaction between two sessile soil bacteria leads to emergence of surface motility. mSphere 4:e00696–18.

14. Hagai E, Dvora R, Havkin-Blank T, Zelinger E, Porat Z, Schulz S, Helman Y. 2014.
Surface-motility induction, attraction and hitchhiking between bacterial species promote
dispersal on solid surfaces. ISME J 8:1147-51.

15. Buch PJ, Chai Y, Goluch ED. 2019. Treating polymicrobial infections in chronic diabetic
wounds. Clin Microbiol Rev 32:e00091.

16. Bisht K, Baishya J, Wakeman CA. 2020. *Pseudomonas aeruginosa* polymicrobial
interactions during lung infection. Curr Opin Microbiol 53:1-8.

17. Crone S, Vives-Florez M, Kvich L, Saunders AM, Malone M, Nicolaisen MH, Martinez-
Garcia E, Rojas-Acosta C, Catalina Gomez-Puerto M, Calum H, Whiteley M, Kolter R,
Bjarnsholt T. 2020. The environmental occurrence of *Pseudomonas aeruginosa*. APMIS
128:220-231.

18. Overhage J, Bains M, Brazas MD, Hancock RE. 2008. Swarming of *Pseudomonas
aeruginosa* is a complex adaptation leading to increased production of virulence factors
and antibiotic resistance. J Bacteriol 190:2671-9.

19. Tremblay J, Deziel E. 2010. Gene expression in *Pseudomonas aeruginosa* swarming
motility. BMC Genomics 11:587.

20. Kazmierczak BI, Schniederberend M, Jain R. 2015. Cross-regulation of *Pseudomonas*
motility systems: the intimate relationship between flagella, pili and virulence. Curr Opin
Microbiol 28:78-82.

21. Drake D, Montie TC. 1988. Flagella, motility and invasive virulence of *Pseudomonas
aeruginosa*. J Gen Microbiol 134:43-52.

22. Feldman M, Bryan R, Rajan S, Scheffler L, Brunnert S, Tang H, Prince A. 1998. Role of
flagella in pathogenesis of *Pseudomonas aeruginosa* pulmonary infection. Infect Immun
66:43-51.

23. Arora SK, Neely AN, Blair B, Lory S, Ramphal R. 2005. Role of motility and flagellin
glycosylation in the pathogenesis of *Pseudomonas aeruginosa* burn wound infections.
Infect Immun 73:4395-8.

24. Coleman SR, Pletzer D, Hancock REW. 2021. Contribution of swarming motility to
dissemination in a *Pseudomonas aeruginosa* murine skin abscess infection model. J
Infect Dis 224:726-733.

25. Ha DG, Kuchma SL, O'Toole GA. 2014. Plate-based assay for swimming motility in
Pseudomonas aeruginosa. Methods Mol Biol 1149:59-65.

26. Burrows LL. 2012. *Pseudomonas aeruginosa* twitching motility: type IV pili in action.
Annu Rev Microbiol 66:493-520.

27. Köhler T, Curty LK, Barja F, van Delden C, Pechère JC. 2000. Swarming of
Pseudomonas aeruginosa is dependent on cell-to-cell signaling and requires flagella and
pili. J Bacteriol 182:5990-6.

28.   Caiazza NC, Shanks RM, O'Toole GA. 2005. Rhamnolipids modulate swarming motility
712   patterns of *Pseudomonas aeruginosa*. J Bacteriol 187:7351-61.

713   29.   Yeung AT, Parayno A, Hancock RE. 2012. Mucin promotes rapid surface motility in
714   *Pseudomonas aeruginosa*. mBio 3:e00073-12.

715   30.   Morrison CB, Markovetz MR, Ehre C. 2019. Mucus, mucins, and cystic fibrosis. Pediatr
716   Pulmonol 54 Suppl 3:S84-S96.

717   31.   Sun E, Liu S, Hancock REW. 2018. Surfing motility: a conserved yet diverse adaptation
718   among motile bacteria. J Bacteriol 200:e00394.

719   32.   Murray TS, Kazmierczak BI. 2008. *Pseudomonas aeruginosa* exhibits sliding motility in
720   the absence of type IV pili and flagella. J Bacteriol 190:2700-8.

721   33.   McAlester G, O'Gara F, Morrissey JP. 2008. Signal-mediated interactions between
722   *Pseudomonas aeruginosa* and *Candida albicans*. J Med Microbiol 57:563-569.

723   34.   Lewis KA, Baker AE, Chen AI, Harty CE, Kuchma SL, O'Toole GA, Hogan DA. 2019.
724   Ethanol decreases *Pseudomonas aeruginosa* flagellar motility through the regulation of
725   flagellar stators. J Bacteriol 201:e00285-19.

726   35.   Badal D, Jayarani AV, Kollaran MA, Prakash D, P M, Singh V. 2021. Foraging signals
727   promote swarming in starving *Pseudomonas aeruginosa*. mBio 12:e0203321.

728   36.   Cystic Fibrosis Foundation. 2018. Cystic Fibrosis Foundation patient registry 2018
729   annual data report. Cystic Fibrosis Foundation, Bethesda, MD.,

730   37.   Limoli DH, Warren EA, Yarrington KD, Donegan NP, Cheung AL, O'Toole GA. 2019.
731   Interspecies interactions induce exploratory motility in *Pseudomonas aeruginosa*. Elife
732   8:e47365.

733   38.   Yarrington KD, Shendruk TN, Limoli DH. 2024. The type IV pilus chemoreceptor PilJ
734   controls chemotaxis of one bacterial species towards another. PLoS Biol 22:e3002488.

735   39.   Mattingly AE, Weaver AA, Dimkovikj A, Shrout JD. 2018. Assessing travel conditions:
736   environmental and host influences on bacterial surface motility. J Bacteriol 200:e00014-
737   18.

738   40.   Sharma J, Sundar D, Srivastava P. 2021. Biosurfactants: potential agents for controlling
739   cellular communication, motility, and antagonism. Front Mol Biosci 8:727070.

740   41.   Tsompanidou E, Sibbald MJ, Chlebowicz MA, Dreisbach A, Back JW, van Dijl JM, Buist
741   G, Denham EL. 2011. Requirement of the *agr* locus for colony spreading of
742   *Staphylococcus aureus*. J Bacteriol 193:1267-72.

743   42.   Tsompanidou E, Denham EL, Becher D, de Jong A, Buist G, van Oosten M, Manson
744   WL, Back JW, van Dijl JM, Dreisbach A. 2013. Distinct roles of phenol-soluble modulins
745   in spreading of *Staphylococcus aureus* on wet surfaces. Appl Environ Microbiol 79:886-
746   95.

747   43.   Otto M. 2014. Phenol-soluble modulins. Int J Med Microbiol 304:164-9.

748   44.   Abdel-Mawgoud AM, Lepine F, Deziel E. 2010. Rhamnolipids: diversity of structures,
749   microbial origins and roles. Appl Microbiol Biotechnol 86:1323-36.

750   45.   Piewngam P, Zheng Y, Nguyen TH, Dickey SW, Joo HS, Villaruz AE, Glose KA, Fisher
751   EL, Hunt RL, Li B, Chiou J, Pharkjaksu S, Khongthong S, Cheung GYC, Kiratisin P, Otto
752   M. 2018. Pathogen elimination by probiotic *Bacillus* via signalling interference. Nature
753   562:532-537.

754   46.   Jimenez PN, Koch G, Thompson JA, Xavier KB, Cool RH, Quax WJ. 2012. The multiple
755   signaling systems regulating virulence in *Pseudomonas aeruginosa*. Microbiol Mol Biol
756   Rev 76:46-65.

757   47.   Guo Q, Kong W, Jin S, Chen L, Xu Y, Duan K. 2014. PqsR-dependent and PqsR-
758   independent regulation of motility and biofilm formation by PQS in *Pseudomonas
759   aeruginosa* PAO1. J Basic Microbiol 54:633-43.

760   48.   Jyot J, Ramphal, R. 2008. Flagella and pili of *Pseudomonas aeruginosa*, p 85-108. *In*
761   Rehm BHA (ed), *Pseudomonas*: Model Organism, Pathogen, Cell Factory. Wiley Online
762   Books.

763   49.   Whitchurch CB, Mattick JS. 1994. Characterization of a gene, *pilU*, required for twitching
764   motility but not phage sensitivity in *Pseudomonas aeruginosa*. Mol Microbiol 13:1079-91.

765   50.   Bertrand JJ, West JT, Engel JN. 2010. Genetic analysis of the regulation of type IV pilus
766   function by the Chp chemosensory system of *Pseudomonas aeruginosa*. J Bacteriol
767   192:994-1010.

768   51.   Kearns DB, Robinson J, Shimkets LJ. 2001. *Pseudomonas aeruginosa* exhibits directed
769   twitching motility up phosphatidylethanolamine gradients. J Bacteriol 183:763-7.

770   52.   Schuster M, Greenberg EP. 2006. A network of networks: quorum-sensing gene
771   regulation in *Pseudomonas aeruginosa*. Int J Med Microbiol 296:73-81.

772   53.   Shao X, Yao C, Ding Y, Hu H, Qian G, He M, Deng X. 2023. The transcriptional
773   regulators of virulence for *Pseudomonas aeruginosa*: Therapeutic opportunity and
774   preventive potential of its clinical infections. Genes Dis 10:2049-2063.

775   54.   Romling U, Galperin MY, Gomelsky M. 2013. Cyclic di-GMP: the first 25 years of a
776   universal bacterial second messenger. Microbiol Mol Biol Rev 77:1-52.

777   55.   Guvener ZT, Harwood CS. 2007. Subcellular location characteristics of the
778   *Pseudomonas aeruginosa* GGDEF protein, WspR, indicate that it produces cyclic-di-
779   GMP in response to growth on surfaces. Mol Microbiol 66:1459-73.

780   56.   Dubern JF, Romero M, Mai-Prochnow A, Messina M, Trampari E, Gijzel HN, Chan KG,
781   Carabelli AM, Barraud N, Lazenby J, Chen Y, Robertson S, Malone JG, Williams P,
782   Heeb S, Camara M. 2022. ToxR is a c-di-GMP binding protein that modulates surface-
783   associated behaviour in *Pseudomonas aeruginosa*. NPJ Biofilms Microbiomes 8:64.

57. Berg HC. 2003. The rotary motor of bacterial flagella. Annu Rev Biochem 72:19-54.

58. Chaban B, Hughes HV, Beeby M. 2015. The flagellum in bacterial pathogens: For
motility and a whole lot more. Semin Cell Dev Biol 46:91-103.

59. Zarrella TM, Khare A. 2022. Systematic identification of molecular mediators of
interspecies sensing in a community of two frequently coinfecting bacterial pathogens.
PLoS Biol 20:e3001679.

60. Mi H, Muruganujan A, Ebert D, Huang X, Thomas PD. 2019. PANTHER version 14:
more genomes, a new PANTHER GO-slim and improvements in enrichment analysis
tools. Nucleic Acids Res 47:D419-D426.

61. Ashburner M, Ball CA, Blake JA, Botstein D, Butler H, Cherry JM, Davis AP, Dolinski K,
Dwight SS, Eppig JT, Harris MA, Hill DP, Issel-Tarver L, Kasarskis A, Lewis S, Matese
JC, Richardson JE, Ringwald M, Rubin GM, Sherlock G. 2000. Gene Ontology: tool for
the unification of biology. The Gene Ontology Consortium. Nat Genet 25:25-9.

62. Gene Ontology C. 2021. The Gene Ontology resource: enriching a GOld mine. Nucleic
Acids Res 49:D325-D334.

63. Saha S, Ojobor CD, Li ASC, Mackinnon E, North OI, Bondy-Denomy J, Lam JS,
Ensminger AW, Maxwell KL, Davidson AR. 2023. F-type pyocins are diverse
noncontractile phage tail-like weapons for killing *Pseudomonas aeruginosa*. J Bacteriol
205:e0002923.

64. Wu W, Jin S. 2005. PtrB of *Pseudomonas aeruginosa* suppresses the Type III Secretion
System under the stress of DNA damage. J Bacteriol 187:6058-68.

65. Yahr TL, Wolfgang MC. 2006. Transcriptional regulation of the *Pseudomonas
aeruginosa* Type III Secretion System. Mol Microbiol 62:631-40.

66. Williams McMackin EA, Djapgne L, Corley JM, Yahr TL. 2019. Fitting pieces into the
puzzle of *Pseudomonas aeruginosa* type III secretion system gene expression. J
Bacteriol 201:e00209-19.

67. Silpe JE, Duddy OP, Bassler BL. 2023. Induction mechanisms and strategies underlying
interprophage competition during polylysogeny. PLoS Pathog 19:e1011363.

68. Trunk K, Benkert B, Quack N, Munch R, Scheer M, Garbe J, Jansch L, Trost M,
Wehland J, Buer J, Jahn M, Schobert M, Jahn D. 2010. Anaerobic adaptation in
Pseudomonas aeruginosa: definition of the Anr and Dnr regulons. Environ Microbiol
12:1719-33.

69. Sun E, Gill EE, Falsafi R, Yeung A, Liu S, Hancock REW. 2018. Broad-spectrum
adaptive antibiotic resistance associated with *Pseudomonas aeruginosa* mucin-
dependent surfing motility. Antimicrob Agents Chemother 62:e00848-18.

70. Pletzer D, Sun E, Ritchie C, Wilkinson L, Liu LT, Trimble MJ, Wolfmeier H, Blimkie TM,
Hancock REW. 2020. Surfing motility is a complex adaptation dependent on the

stringent stress response in *Pseudomonas aeruginosa* LESB58. PLoS Pathog
16:e1008444.

71.   Morin CD, Deziel E. 2021. Use of alternative gelling agents reveals the role of
rhamnolipids in *Pseudomonas aeruginosa* surface motility. Biomolecules 11:1468.

72.   Belas R. 2014. Biofilms, flagella, and mechanosensing of surfaces by bacteria. Trends
Microbiol 22:517-27.

73.   Khan F, Pham DTN, Oloketuyi SF, Kim YM. 2020. Regulation and controlling the motility
properties of *Pseudomonas aeruginosa*. Appl Microbiol Biotechnol 104:33-49.

74.   Bru JL, Rawson B, Trinh C, Whiteson K, Høyland-Kroghsbo NM, Siryaporn A. 2019.
PQS produced by the *Pseudomonas aeruginosa* stress response repels swarms away
from bacteriophage and antibiotics. J Bacteriol 201:10.1128/jb.00383-19.

75.   Yarrington KD, Shendruk, T. N., Limoli, D. H. 2022. Twitching cells use a chemoreceptor
to detect bacterial competitors. *bioRxiv* doi:10.1101/2022.11.28.518211.

76.   Kamatkar NG, Shrout JD. 2011. Surface hardness impairment of quorum sensing and
swarming for *Pseudomonas aeruginosa*. PLoS One 6:e20888.

77.   Fauvart M, Phillips P, Bachaspatimayum D, Verstraeten N, Fransaer J, Michiels J,
Vermant J. 2012. Surface tension gradient control of bacterial swarming in colonies of.
Soft Matter 8:70-76.

78.   Bru JL, Kasallis SJ, Chang R, Zhuo Q, Nguyen J, Pham P, Warren E, Whiteson K,
Hoyland-Kroghsbo NM, Limoli DH, Siryaporn A. 2023. The great divide: rhamnolipids
mediate separation between *P. aeruginosa* and *S. aureus*. Front Cell Infect Microbiol
13:1245874.

79.   Kearns DB. 2010. A field guide to bacterial swarming motility. Nat Rev Microbiol 8:634-
44.

80.   Kearns DB, Losick R. 2003. Swarming motility in undomesticated *Bacillus subtilis*. Mol
Microbiol 49:581-90.

81.   Huber B, Riedel K, Hentzer M, Heydorn A, Gotschlich A, Givskov M, Molin S, Eberl L.
2001. The *cep* quorum-sensing system of *Burkholderia cepacia* H111 controls biofilm
formation and swarming motility. Microbiology (Reading) 147:2517-2528.

82.   Soberón-Chávez G, Lépine F, Déziel E. 2005. Production of rhamnolipids by
Pseudomonas aeruginosa. Applied Microbiology and Biotechnology 68:718-725.

83.   Lindum PW, Anthoni U, Christophersen C, Eberl L, Molin S, Givskov M. 1998. N-Acyl-L-
homoserine lactone autoinducers control production of an extracellular lipopeptide
biosurfactant required for swarming motility of *Serratia liquefaciens* MG1. J Bacteriol
180:6384-8.

84.   Bernier SP, Hum C, Li X, O'Toole GA, Magarvey NA, Surette MG. 2017. *Pseudomonas*
857   *aeruginosa*-derived rhamnolipids and other detergents modulate colony morphotype and
858   motility in the *Burkholderia cepacia* complex. J Bacteriol 199:e00171-17.

859   85.   Morin C, Landry M, Groleau MC, Deziel E. 2022. Surface motility favors codependent
860   interaction between *Pseudomonas aeruginosa* and *Burkholderia cenocepacia*. Msphere
861   doi:10.1128/msphere.00153-22.

862   86.   Pardhi DS, Panchal RR, Raval VH, Joshi RG, Poczai P, Almalki WH, Rajput KN. 2022.
863   Microbial surfactants: a journey from fundamentals to recent advances. Front Microbiol
864   13:982603.

865   87.   Yang A, Tang WS, Si T, Tang JX. 2017. Influence of physical effects on the swarming
866   motility of *Pseudomonas aeruginosa*. Biophys J 112:1462-1471.

867   88.   Inoue T, Shingaki R, Fukui K. 2008. Inhibition of swarming motility of *Pseudomonas*
868   *aeruginosa* by branched-chain fatty acids. FEMS Microbiol Lett 281:81-6.

869   89.   Rahme LG, Stevens EJ, Wolfort SF, Shao J, Tompkins RG, Ausubel FM. 1995.
870   Common virulence factors for bacterial pathogenicity in plants and animals. Science
871   268:1899-902.

872   90.   Fey PD, Endres JL, Yajjala VK, Widhelm TJ, Boissy RJ, Bose JL, Bayles KW. 2013. A
873   genetic resource for rapid and comprehensive phenotype screening of nonessential
874   *Staphylococcus aureus* genes. MBio 4:e00537-12.

875   91.   Liberati NT, Urbach JM, Miyata S, Lee DG, Drenkard E, Wu G, Villanueva J, Wei T,
876   Ausubel FM. 2006. An ordered, nonredundant library of *Pseudomonas aeruginosa* strain
877   PA14 transposon insertion mutants. Proc Natl Acad Sci U S A 103:2833-8.

878   92.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing
879   reads. EMBnet J 17:10-12.

880   93.   Bolger AM, Lohse M, Usadel B. 2014. Trimmomatic: a flexible trimmer for Illumina
881   sequence data. Bioinformatics 30:2114-20.

882   94.   McClure R, Balasubramanian D, Sun Y, Bobrovskyy M, Sumby P, Genco CA,
883   Vanderpool CK, Tjaden B. 2013. Computational analysis of bacterial RNA-seq data.
884   Nucleic Acids Res 41:e140.

885   95.   Tjaden B. 2015. *De novo* assembly of bacterial transcriptomes from RNA-seq data.
886   Genome Biol 16:1.

887   96.   Zhou Y, Liang Y, Lynch KH, Dennis JJ, Wishart DS. 2011. PHAST: a fast phage search
888   tool. Nucleic Acids Res 39:W347-52.

889   97.   Arndt D, Grant JR, Marcu A, Sajed T, Pon A, Liang Y, Wishart DS. 2016. PHASTER: a
890   better, faster version of the PHAST phage search tool. Nucleic Acids Res 44:W16-21.

891   98.   Tretyakov K. 2020. matplotlib_venn, GitHub repository.

892   99.   Winsor GL, Griffiths EJ, Lo R, Dhillon BK, Shay JA, Brinkman FS. 2016. Enhanced
893   annotations and features for comparing thousands of *Pseudomonas genomes* in the
894   *Pseudomonas* genome database. Nucleic Acids Res 44:D646-53.