RBM7 deficiency promotes breast cancer metastasis by coordinating MFGE8

splicing switch and NF-kB pathway

Fang Huang[1†], Zhenwei Dai[1†], Jinmiao Yu[1†], Kainan Wang[†2], Chaoqun Chen[1], Dan

Chen[3], Jinrui Zhang[4], Jinyao Zhao[4], Mei Li[4], Jinyao Zhao[4], Wenjing Zhang[4], Xiaojie

Li[5*], Yangfan Qi[4,6*], Yang Wang[1*]

[1]Sino-US Research Center for Cancer Translational Medicine of the Second Affiliated

Hospital of Dalian Medical University & Institute of Cancer Stem Cell, Dalian

Medical University, Dalian 116023, China

[2]Department of Oncology & Sino-US Research Center for Cancer Translational

Medicine, the Second Affiliated Hospital, Dalian Medical University, Dalian 116023,

China

[3]Department of Pathology, the First Affiliated Hospital of Dalian Medical University,

Dalian Medical University, Dalian 116011, China

[4]Institute of Cancer Stem Cell, Dalian Medical University, Dalian 116044, China

[5]Department of Prosthodontics, College of Stomatology, Dalian Medical University,
Dalian, 116044, China.

[6]Soochow University Cancer Institute, Suzhou, China

[†] These authors contributed equally to this work.

[*] Correspondence:

Yang Wang; Sino-US Research Center for Cancer Translational Medicine of the

Second Affiliated Hospital of Dalian Medical University & Institute of Cancer Stem

Cell, Dalian Medical University, Dalian 116023, China; Tel: +86-411-86110531;

Email: yangwang@dmu.edu.cn

Yangfan Qi; Institute of Cancer Stem Cell, Dalian Medical University, Dalian, China

116044; Tel: +86-411-86110531; Email: yangfanqi@dmu.edu.cn

Running title: RBM7 presents anti-metastasis activity by regulating splicing

We declare no conflict of interest on the studies described in this manuscript.

Abstract

Aberrant alternative splicing is well-known to be closely associated with

tumorigenesis of various cancers. However, the intricate mechanisms underlying

breast cancer metastasis driven by deregulated splicing events remain largely

unexplored. Here, we unveiled that RBM7 is decreased in lymph node and distant

organ metastases of breast cancer as compared to primary lesions and low expression

of RBM7 is correlated with the reduced disease-free survival of breast cancer patients.

Breast cancer cells with RBM7 depletion exhibited an increased potential for lung

metastasis compared to scramble control cells. The absence of RBM7 stimulated

breast cancer cell migration, invasion, and angiogenesis. Mechanistically, RBM7

controlled the splicing switch of MFGE8, favoring the production of the predominant

isoform of MFGE8, MFGE8-L. This resulted in the attenuation of STAT1

phosphorylation and alterations in cell adhesion molecules. MFGE8-L exerted an

inhibitory effect on the migratory and invasive capability of breast cancer cells, while

the truncated isoform MFGE8-S, which lack the second F5/8 type C domain had the

opposite effect. In addition, RBM7 negatively regulates the NF-κB cascade and an

NF-κB inhibitor could obstruct the increase in HUVEC tube formation caused by

RBM7 silencing. Clinically, we noticed a positive correlation between RBM7

expression and MFGE8 exon7 inclusion in breast cancer tissues, providing new

mechanistic insights for molecular-targeted therapy in combating breast cancer.

Keywords: RBM7; alternative splicing; breast cancer; metastasis

Introduction

Alternative splicing (AS) emerges as a substantial contributor to the production of a myriad of transcripts from limited genes, which enriched protein variation and phenotypic diversity in multicellular organisms. Pre-mRNA splicing reactions require a step-wise assembly of spliceosome, a megadalton complex that recognize consensus regulatory sequences, including 5' and 3' splice sites marking exon-intron boundary, and the branch point site (BPS), to accomplish exon definition (Shi, 2017) (Wilkinson, Charenton et al., 2020). While a splice site is suboptimal, AS emerges under control of inherent sequence acting in cis, namely exonic or intronic splicing enhancer or silence, and the cognate trans-acting splicing factors, which can strengthen or weaken the recognition of splice sites by spliceosome.

It is well documented that AS dysregulation leads to multiple human diseases, including cancer (Montes, Sanford et al., 2019) (Singh & Cooper, 2012). Tumorigenicity has been linked with cancer-specific splicing switch arising from either cis-elements mutations or deregulated splicing regulatory factors. In some cases, splicing factors, particularly spliceosomal components SF3B1 and U2AF1, are frequently mutated across diverse cancer types (Baralle, 2005) (Harbour, Roberson et al., 2013) (Smith, Choudhary et al., 2019) (Cheruiyot, Li et al., 2021). For instance, recurrent SF3B1 mutations led to retention of intronic region in BRD9 pre-mRNA, conferring loss of core subunit function to the ncBAF chromatin remodeling complex and metastatic advantage to melanoma cells (Inoue, Chew et al., 2019). Another layer of regulation involves the abnormal expression of splicing factors derived from

altered regulatory pathway. As an archetypal example, a set of members in SR protein family such as SRSF1 and SRSF3 are upregulated in breast tumors, lung tumors, glioma and exhibit oncogenic ability via altering AS products (for example, Bcl-x, MYO1B, NUMB and p53) (Anczuków, Akerman et al., 2015) (Zhou, Wang et al., 2019) (Tang, Horikawa et al., 2012) (Zhou, Gong et al., 2020). Otherwise, many splicing factors such as RBM (RNA binding motif) proteins can either functions as oncoproteins or tumor suppressors in cancer-type dependent manner (Li, Guo et al., 2021), which may be attributed to tissue-specific AS patterns.

Breast cancer is the most common diagnosed malignancy and remain the second leading cause of cancer mortality (over 41,000 deaths annually in the USA) among females worldwide (Siegel, Miller et al., 2019). Metastasis to vital organs accounts for approximately 90% cancer-related death. The 5-year overall survival rate of patients with metastatic breast cancer is drastically decreased to about 25%, as compared to the primary cancer patients' prognosis with a survival rate greater than 80% (Rabbani & Mazar, 2007) (Valastyan & Weinberg, 2011). Despite advances in early diagnosis and comprehensive interventions for breast cancer treatment, limited effective strategy against metastasis has been an outstanding challenge for improving prognostication of patients. This highlights the urgent need for deepened understanding of mechanisms underlying development of metastasis. Tumor metastasis is multi-step process involving a sequential cascade of events, such as epithelial-mesenchymal transition (EMT), local invasion from primary tumor site, lymphogenic or hematogenic dissemination, cancer cells seeding and metastatic

lesions formation (Liang, Zhang et al., 2020). AS dysregulation has been implicated in

multistage process of metastasis. For example, the switch of CD44 or FGFR2 splicing

isoforms can lead to EMT phenotype (Chen, Zhao et al., 2018) (Warzecha, Sato et al.,

2009). VEGF-A or XBP1 splicing transition is tightly linked to angiogenesis in cancer

(Bates, 2008) (Zeng, Xiao et al., 2013). On account of the phenotypic plasticity in

metastatic cells conferred by alternative splicing, further investigation in potential role

of critical splicing regulatory factors and the associated splicing events in breast

cancer metastasis is warranted.

RNA-binding proteins (RBPs) are a large class of conserved proteins that

participate in cancer development through precisely control RNA metabolism,

including RNA splicing. RBM7, a member of RBP family, is commonly known to

exert RNA surveillance activity as a component of the human nuclear exosome target

(NEXT) complex (Meola, Domanski et al., 2016) . Aberrant expression of RBM7 is

not only associated with lung fibrosis onset and motor neuronopathy, but also impacts

cancer cell proliferation (Fukushima, Satoh et al., 2020) (Xi, Zhang et al., 2020).

Previous findings have established a direct connection of RBM7 with spliceosome

factors SF3B2 and SF3B4 (Falk, Finogenova et al., 2016), yet whether RBM7 is

involved in the regulation of AS remain elusive. In this study, we report that RBM7 is

frequently reduced in a subset of breast cancer metastases from lymph nodes lesions

and distant organ as compared to primary tumors and its low expression predicts

dismal survival outcomes of patients. Breast cancer cells with loss of RBM7 gain

enhanced aggressive capability and metastatic potential. Mechanistically, RBM7

ablation enhanced the splicing switch of MFGE8 pre-mRNA from canonical isoform

MFGE8-L to a truncated isoform MFGE8-S. Ectopic expression of MFGE8-L caused

marked suppression of migratory and invasive ability of breast cancer cells, whereas

MFGE8-S showed opposite function/ disabled such function. At another layer, RBM7

discordantly regulated the phosphorylation of p65. In concert, NF-κB inhibitor

abolished pro-angiogenesis effect of RBM7 ablation in breast cancer. Above all, our

findings uncovered a previously unidentified role of RBM7 in breast cancer

metastasis via coupling MFGE8 splicing switch and NF-κB signaling activation,

suggesting a new therapeutic target and prognostic predictor for breast cancer.

Results

RBM7 is reduced in metastatic lesions of breast cancer and positively correlated

with patients' prognosis

To determine the clinical relevance of RBM7 in breast cancer, we assessed the

correlation between RBM7 and important clinical outcomes. Analysis of TCGA, GEO

and the METABRIC breast cancer datasets showed that RBM7 expression positively

correlates with overall survival of breast cancer patients (Figure 1A and

Supplementary Figure 1A-1B). RBM7 low expression predicts the reduced

disease-free survival in patients with breast cancer (Figure 1B and Supplementary

Figure 1C). Congruently, we found RBM7 is decreased in breast cancer compared to

normal tissues (Figure 1C and Supplementary Figure 1D). Further dissection in a

cohort of breast cancer patients from the cBioPortal unveiled cases with lymph node

metastasis have lower expression of RBM7 as compared to cases without metastasis

(Figure 1D). We next performed immunohistochemistry (IHC) detection of RBM7 in

breast cancer tissue microarrays. The results showed that patients bearing clinical

stage III tumors have significantly decreased RBM7 expression than patients bearing I

and II clinical stage tumors (Figure 1E). In tissue chip containing triple negative

breast cancer, the more aggressive and mesenchymal breast cancer subtype, RBM7

exhibits reduced protein levels in tumors as compared to paracancer tissues (Figure

1F). In addition, we collect specimen consisted of tumor tissues from 21 breast

primary cancer, in 9 cases, from metastatic loci and took advantage of IHC analysis to

determine the RBM7 expression. Most (25% strong positive;67% moderate positive)

of primary tumor samples displayed a positive IHC staining for RBM7, whereas most

of lung/liver metastatic lesions showed weak or negative RBM7 expression (Figure

1G). We also obtained paired primary breast cancer and lymph node metastatic loci

and found, when compared to the corresponding primary tumor, the expression of

RBM7 is significantly decreased in lymph node metastases (Figure 1H-1I). These

results associated RBM7 with breast cancer metastasis, prompting us to further

investigate the functional role of RBM7 in metastatic processes of breast cancer.

RBM7 silencing provokes metastatic potential of breast cancer

In light with the above findings, we generated RBM7-silenced breast cancer cells

using lentiviral expression of short hairpin RNA (shRNA) followed by profiling

high-throughput transcriptome by RNA sequencing. Strikingly, Gene Ontology and

pathway analysis displayed a wide range of differentially expressed genes were

enriched in extracellular matrix organization, cell migration and angiogenesis (Figure

2A-2B), which have been characterized as crucial processes in tumor metastasis. We

then conducted qRT-PCR analysis of theses metastasis-associated genes in

RBM7-depleted breast cancer cells and verified that they were up-regulated upon

RBM7 knockdown shown as heatmap representation (Figure 2C). The lung metastasis

is one of the most prevalent distant metastases that breast cancer is prone to and

associated with poor prognosis of patients (Medeiros & Allan, 2019). Thus, we

performed tail-vein transfer model to evaluate the potential impact of RBM7 on

pulmonary metastasis of breast cancer cells. 6×10^5 4T1 cells, the mouse breast cancer

cell line simulating stage IV human breast tumor, with stable RBM7 silence or control

vector were implanted into BALB/c mice via tail vein. RBM7-silenced 4T1 cells were

able to develop more spontaneous lung metastases than control cells as judged by

pulmonary nodules number and haematoxylin-eosin (HE) staining of lung tissues

(Figure 2D-2F and Supplementary Figure 2A). Cancer metastasis is comprised of a

complex cascade of events, such as migration, invasion and angiogenesis, thus we

next to determine which process is mediated by RBM7-loss in facilitating breast

cancer metastasis. We utilized transwell chamber with or without matrix coat and

would healing assays to examine the potential function of RBM7 in migration and

invasion capabilities and found the depletion of RBM7 led to a visible increase in

mobility and invasiveness of both the triple-negative breast cancer (TNBC) cell line

MDA-MB-231, estrogen receptor positive breast cancer cell line MCF7 and 4T1 cells

(Figure 2G-2H and Supplementary Figure 2B-2E). Conversely, RBM7 overexpression

notably repressed MDA-MB-231 and MCF7 cells to migrate or invade across

transwell inserts (Figure 2I-2J). Similar results were obtained in two additional TNBC

cell lines BT549 and HCC1937 (Supplementary Figure 2F-2G), which proved that

RBM7 possessed the capability to suppress the migration and invasion of breast

cancer cells. To examine the impact of RBM7 on angiogenesis in breast cancer, we

cultured human umbilical vein endothelia cells (HUVEC) with condition medium

obtained from culture supernatants of RBM7-depleted or scramble shRNA transduced

cells for tube formation analysis. Compared to the scramble control, HUVEC

incubated with culture medium from RBM7-knockdown breast cancer cells exhibited

accelerated tube formation length and branch junction (Figure 2K), suggesting the

inhibitory effect of RBM7 on breast cancer angiogenesis. Taken together, the above

data demonstrated RBM7 as tumor suppressor by regulating breast cancer migration,

invasion and angiogenesis.

RBM7 exhibits splicing regulatory function in breast cancer

In previous studies, RBM7 is commonly characterized as an RNA-binding

constituent of trimeric NEXT complex, which directs the RNA exosome to

non-coding transcriptome for degradation (Lubas, Andersen et al., 2015). However,

the mechanistical role of RBM7 in cancer remain largely unknown. Intriguingly, we

observed that, attributed to the ablation of RBM7, almost 551 splicing events occurred

significant decrease in PSI (percent spliced in), whereas 289 splicing events endured

increased PSI (Figure 3A). The changed events were divided among various types of

AS (alternative splicing), including SE (skipping exon), IR (intron retention), A5E

(alternative 5' splice site exon), A3E (alternative 3' splice site exon), and MXE

(mutually exclusive exons), of note, SE events were the most prominently regulated

(Figure 3B). A majority of AS events in particular of SE and A5E category were in

decrease of PSI upon RBM7 depletion (Figure 3C-3D), indicating that the presence of

RBM7 may promote the usage of alternative splice sites of SEs or A5Es.

Subsequently, we extracted the candidate binding motifs which reside within or near

cassette exons or A5E and found UUUCUU motif, simiar to RBM7 binding site

(polyU) were enriched in RBM7 positively regulated AS events to a greater extent

218 that control exons unimpacted by RBM7 (Figure 3E), raising the possibility that

219 RBM7 may serve as a splicing activator through directly tethering cis-element in

220 pre-mRNA. This is coherent with the nuclear localization of RBM7 in breast cancer

221 cells (Supplementary Figure 3A). In order to explore the splicing regulation activity

222 of RBM7 in distinct pre-mRNA context, we validated a series of

223 RBM7-significantly-changed splicing events across different AS types according to

224 the RNA-seq results. The RT-PCR analysis demonstrated that the PSI of selected gene

225 related to cancer progression was either increased or decreased endogenously upon

226 RBM7 depletion (Figure 3F and Supplementary Figure 3B). Considering that the AS

227 is a sophisticated multi-step reaction that is regulated by both pre-mRNA-interacting

228 splicing factor and other assistant factors in a context-dependent manner (Wright,

229 Smith et al., 2022), we doubted the divergent effect of RBM7 on distinct splicing

230 targets was due to a more complicated mechanism. Gene ontology (GO) enrichment

231 analysis identified a majority of genes undergoing AS alterations were involved in

232 cytoskeleton assembly and organization (Figure 3G). While globally analyzing the

233 network enrichment of RBM7-regulated downstream splicing events, we observed its

234 targets are functionally connected to well-linked network containing gene subgroups

235 (Figure 3H), which are associated with biological processes in tumor cell metastasis,

236 again suggesting the cancer-related function of RBM7.

238 **RBM7 depletion switches splicing outcomes of MFGE8 by decreasing MFGE8-L**

239 **isoform**

To ascertain the functional targets that explain the inhibitory role of RBM7 in

breast cancer metastasis, we analyzed RNA immunoprecipitation sequencing data for

RBM7 and whole-transcriptome sequencing data for breast cancer cells expressing

two shRNAs targeting RBM7. Venn diagram of 8 hub genes obtained by overlapping

three gene sets pointed forward a hotspot gene, namely MFGE8 (Figure 4A).

RNA-seq data showed that the depletion of RBM7 resulted in a significant alteration

in ratio of MFGE8 two isoforms, of which the canonical isoform (MFGE8-L) has

been demonstrated to be correlated with tumor formation and aggressiveness of breast

cancer (Yang, Hayashida et al., 2011). Importantly, the exon 7 located in F5/8 type C

2 domain, which is obligated to bind to phosphatidylserine, was lacked in the short

isoform of MFGE8 (MFGE8-S) (Figure 4B). Further RT-PCR identification verified

RBM7 ablation indeed potentiated the production of MFGE8-S in various TNBC cell

lines (Figure 4C). Similar results were obtained from MCF7 cells, suggesting that

RBM7 regulated the AS of MFGE8 independently of the ER status. In separate

experiments, we transfected breast cancer cells with an exogenous MFGE8 splicing

reporter containing exon 6-8 and 600bp of flanking intron sequences to determine the

splicing regulatory function of RBM7. Accordingly, the depletion of RBM7 still

induced the splicing shift of the minigene reporter by elevating MFGE8-S variant

(Figure 4D). To test whether such splicing regulatory effect was directly or indirectly,

we analyzed the sequence of alternative exon 7 and the motifs nearby its 5' or 3'

splice sites, and found two poly(U) motifs are positioned in proximal to the pseudo 3'

splice site. Subsequent RT-PCR assay for the precipitation in RIP assays confirmed

RBM7 could bind to the upstream sequence containing UUUCUU motifs adjacent to

3' splice site of MFGE8 cassette exon, but not another region nearby it (Figure 4E).

To pinpoint the location for the potential cis-element for AS regulation by RBM7, we

designed antisense oligonucleotides (ASOs), which have been chemically modified to

avoid endogenous nuclease degradation without recruiting RNase H, to block the

UUUCUU residues in proximity to intron6/exon7 junction. As shown in Figure 4F,

when compared to control ASO, treatment with ASOs targeting the potential binding

site (UUUCUU) of RBM7, contributed to the exclusion of exon7 of MFGE8. In

separate experiments, the depletion of RBM7 induced the splicing shift of the

minigene reporter by elevating MFGE8-S variant. While the binding motif was

depleted or mutated, RBM7 failed to affect the splicing outcomes of MFGE8

(Supplementary Figure SC). Due to its close proximity to 3' splice site, UUUCUU

residues bound by RBM7 might participated in spliceosome assembly on the

upstream 3' splice site of exon7, which may explain why disruption of the motif led

to almost complete exon7 skipping. Oue data suggest that RBM7 regulates the exon

skipping of MFGE8 by directly binding to UUUCUU located at six nucleotides

upstream of the 3' splice-site of exon7.

The short variant of MFGE8 abolishes the inhibitory function of its classical

isoform on breast cancer cell migration and invasion

MFGE8 encodes a secreted protein, Milk fat globule-EGF factor 8 (MFG-E8),

which participates in a variety of cellular processes, including mammary gland

branching morphogenesis, angiogenesis and mucosal healing, etc (Raymond, Ensslin

et al., 2009). We applied MFGE8-L stably overexpressing- or depleted-breast cancer

cells to transwell assays and observed the cellular migratory and invasive capabilities

was visibly suppressed by MFGE8-L overexpression (Supplementary Figure 4A-4B).

On the contrary, the knockdown of MFGE8-L provoked cell migration and invasion

(Supplementary Figure 4C). Strikingly, while MDA-MB-231 or HCC1937 expressing

MFGE8-L showed a remarkable decrease in migratory and invasive ability compared

to control vector, MFGE8-S not only totally abolished the inhibitory function of

MFGE8-L but elicited modest pro-migration/ invasion actions on breast cancer cells

(Figure 5A and Supplementary Figure 4D-4E). We also performed Fluorescent

Gelatin Degradation Assays for investigating invadopodia formation and found

MFGE8-L up-regulation suppressed breast cancer cells invasion through a layer of

extracellular matrix, whereas breast cancer cells with ectopic expression of MFGE8-S

acquired enhanced ability to degrade matrix and invasion (Figure 5B).

Immunofluorescence assays revealed the MFGE8-L forms foci in cytoplasm,

particularly enriched near cell-cell contact sites, whereas MFGE8-S is more evenly

distributed in the nucleus and cytoplasm (Figure 5C). As shown in diagram of domain

configuration for two variants, in comparison with MFGE8-L isoform containing

N-terminal EGF-like domain and C-terminal twice-repeated F5/8 type C domains, the

protein encoded by MFGE8-S loses the second F5/8 type C domain, implying a

shortage of the function corresponding to membrane-binding (Supplementary Figure

4F). The different structural bases of two isoforms may also be linked to their distinct

cellular function in tumor invasion (Supplementary Figure 4G). Furthermore, we

compared transcriptome profiles between MFGE8-L and MFGE8-S ectopically

expressing breast cells and found a set of functional genes were down-regulated by

MFGE8-L, whereas this negative regulation was abrogated in response to MFGE8-S

(Supplementary Figure 4H). The differentially expressed genes between two splice

variants were predominantly enriched in cell adhesion molecules and JAK-STAT

cascade as revealed by KEGG pathway and Gene ontology analysis (Figure 5D and

Figure 5E). Subsequent experimental validation demonstrated that the activation of

STAT1 was hindered by MFGE8-L, but not affected by MFGE8-S (Figure 5F).

Considering that STAT1 signaling activation plays an important role in tumor cell

metastasis via cell adhesion molecules induction (Marimuthu, Lakshmanan et al.,

2022) (Chen, Hao et al., 2019), it is highly plausible that MFGE8-L, but not

MFGE8-S, impeded breast cancer cell motility and invasive capability in a

STAT1-dependent manner.

RBM7 depletion-stimulated aggressiveness of breast cancer is dependent on

MFGE8 splicing switch and NF-kB pathway activation

Next, we sought to assess if MFGE8 splicing switch is responsible for

acquisition of aggressive feature in RBM7-silenced breast cancer cells. Flag-tagged

MFGE8-L or an empty vector was co-expressed in MDA-MB-231 cells together with

RBM7 shRNA (Supplementary Figure 5A). Following seeding into transwell chamber,

cells transfected with RBM7 shRNA exerted a higher migratory and invasive

capability than scramble control, consistent with above data. Importantly, the

restoration of MFGE8-L expression diminished the acquired aggressive properties of

MDA-MB-231 cells resulted from RBM7 depletion (Figure 6A). Similar results were

obtained using RBM7-depleted MCF7 cells with or without MFGE8-L complement

(Figure 6A). To corroborate the role of MFGE8 splicing shift in RBM7- regulated

breast cancer cell invasion, we transfected triple-negative breast cancer cells with

ASOs directed against RBM7-binding motif and examined the potential impact on

cell aggressiveness. The results showed an obvious increase in exon7-skipped variant

as compared to the scramble negative control ASOs (Supplementary Figure 5B).

Meanwhile, the migrative and invasive ability of breast cancer cells treated with

ASOs was significantly boosted (Figure 6B), further suggesting that

RBM7-knockdown stimulated aggressiveness of breast cancer at least partially relies

on the control of MFGE8 AS. However, a series of shRBM7-upregulated secreted

factors that have been reported to drive tumor angiogenesis (Gopinathan, Milagre et

al., 2015) (Fousek, Horn et al., 2021) (Feng, Ke et al., 2019), were not impacted by

MFGE8-L/S (Figure 6C and Supplementary Figure 5C). Condition medium derived

from breast cancer cells overexpressing MFGE8-L/S showed negligible effect on

HUVEC tube formation (Supplementary Figure 5D, implying that there exists

additional mechanism underlying breast cancer metastasis attributed to RBM7

reduction.

Given that consecutive gene set enrichment analysis (GSEA) of highest-ranking

gene sets based on transcriptomic profile of mRNA expression showed a signature for

NF-κB pathway that was augmented upon depletion of RBM7 (Figure 6D), we suspected that the transcriptional activation of downstream genes in NF-κB signaling known to facilitate tumor angiogenesis might be involved in shRBM7-exerted pro-metastatic function. As expected, the phosphorylation of p65, a core transcription factor well-known to switch on NF-κB pathway, was notably enhanced by RBM7 knockdown. Meanwhile, IκBα phosphorylation that can blockade p65 activation was curbed upon RBM7 knockdown (Figure 6E). Conversely, RBM7 overexpression impeded the activation of p65 (Supplementary Figure 5E). Yet MFGE8-L/S had no effect on phosphorylation of p65 or IκBα (Supplementary Figure 5F). We found RBM7 depletion remarkably promoted the expression of IL-1β as judged by qPCR and ELISA assays (Supplementary Figure 5G-5I). IL-1β, commonly known as a pro-inflammatory cytokine, could bind to IL-1R and initiate a multistage enzymatic reaction that triggers the activation of NF-κB pathway (Weber, Kracht et al., 2010) (Guo, Jin et al., 2024). Thus, it is plausible the upregulation of IL-1β might be a causal factor in RBM7-depletion-induced activation of NF-kB signaling. We further carried out Matrigel tube formation assays and found NF-κB inhibitor PDTC drastically reversed the promotion of HUVEC angiogenesis by conditional medium harvested from RBM7-depleted cells (Figure 6F-6G), implying that RBM7-p65 axis may participate in HUVEC angiogenesis through reprogramming the secretome of breast cancer cells. Similar effect was observed with respect to cell motility and invasion (Supplementary Figure 5J). Taken together, these data above indicated that both MFGE8 splice switch and NF-κB pathway activation are required for mediating

the pro-metastasis activity of RBM7 reduction in breast cancer.

The PSI of MFGE8 exon 7 is decreased in breast cancer patients and positively

correlated with RBM7 expression

To study the clinical relevance of MFGE8 isoform switch, we analyzed the PSI

changes of cassette exon7 in TCGA breast cancer cohort and found exon7 skipping

occurs more frequently in tumor than in normal tissues (Figure 7A). Consistently, in

comparison with normal counterparts, the splicing of MFGE8 is shifted towards a

reduced proportion of MFGE8-L across multiple cancer types, including LUAD,

HNSC, COAD, etc (Supplementary Figure 6A). RT-PCR assay was performed in

surgical tumor specimens in breast cancer patients and showed increased proportion

of exon7 exclusion in most tumor samples when compared to normal tissues (case#1:

86:94; case#2: 84:86; case#3: 79:85; case#4: 63:75; case#5: 69:93; case#6: 71:80)

(Figure 7B). On the other hand, the lymph node metastases contain a higher

proportion of MFGE8 variant with skipped exon7 in comparison with paired primary

tumor tissues (case#1: 80:95; case#2: 86:97; case#3: 84:90; case#4: 70:78; case#5:

83:89) (Figure 7C). This is coherent with decreased RBM7 expression levels found in

breast cancer with lymph node metastasis. In addition, patients with high expression

of MFGE8 exon7 had a visibly better overall survival than patients with low

expression (Figure 7D). We performed Pearson correlation test and revealed a

noticeably positive correlation between the expression of RBM7 and exon7 inclusion

of MFGE8 in breast cancer (Figure 7E), further supporting MFGE8 isoform switch

regulated by RBM7.

Discussion

Previous reports have recognized the RNA-binding protein RBM7 as a

component of the NEXT complex that could facilitate RNA exosome targeting to a

subset of non-coding RNAs and aberrant transcripts (Meola et al., 2016) (Pascal

Preker, 2008), including short-lived exosome substrates and pre-mRNAs, for

surveillance and turnover . In addition to this specific characteristic, RBM7 was

shown to interact with subunits of 7SK snRNP complex and thus activate P-TEF

downstream genes in response to DNA damage (Bugai, Quaresma et al., 2019).

Herein, we discover a novel role of RBM7 in alternative splicing control through

directly binding to its cognate cis-element in pre-mRNA. Using whole-transcriptome

sequencing, we found the depletion of RBM7 contributed to exon exclusion in

approximately 70% SE/A5E events, implying RBM7 mainly functions as a splicing

activator in the context of SE/A5E in breast cancer. However, the increase of PSI also

occurred due to RBM7 knockdown in a smaller faction of splicing events, such as

ZMYND8. We speculated this converse impact may be associated with the

intrinsically sophisticated splicing processes driven by distinct core splicing factors,

cognate cis-acting element and co-factors in a context-dependent manner (e.g.,

pre-mRNA structure, base modification, antagonistic or cooperative effect of multiple

splicing factors). Similar to many splicing factors, like SR proteins and hnRNPs,

RBM7 has an RRM-domain required for target RNA recognition in the N-terminus.

We performed co-IP experiment and found that RBM7 could interact with RNA

splicing factor SF3B2, a component of spliceosomal U2 SnRNP complex

(Supplementary Figure 6B). Consistent with the AS regulation of MFGE8 by RBM7,

the depletion of SF3B2 also promoted exon7 skipping (Supplementary Figure 6C-6D),

implying a cooperative effect of the two proteins in regulating MFGE8 splicing. This

is in concert with a previous study that RRM domain of RBM7 could bind a

proline-rich segment within SF3B2 (Falk, Finogenova et al., 2016). The interaction

mode with strong similarity to RBM7^{RRM}–ZCCHC8^{Proline} interaction in the NEXT

complex indicated mutually exclusive binding of SF3B2 and ZCCHC8 to RBM7.

Thus, RBM7 appears to play dual, but not conflicting, roles during RNA processes

depending on its interaction with the spliceosome or exosome.

Currently, the pathobiological function of RBM7 in breast cancer is

underexplored except one study that claimed an oncogenic role of RBM7 since its

knockdown inhibited breast cancer cell proliferation (Xi et al., 2020). Nevertheless, in

our study, RBM7 knockdown in breast cancer displayed visibly pro-metastasis

function with activated NF-κB signaling. More importantly, according to TCGA

database (total cases > 1000 used for analysis), RBM7 shows decreased expression in

breast cancer, particularly in the metastasis loci and has positive correlation with

patients' survival outcomes. It is therefore highly doubtful that RBM7 could serve as

an oncogene in breast cancer as previously reported. Moreover, we can't exclude

additional biological function of RBM7 in breast cancer. RBM7-targeted silencing for

breast cancer would likely be detrimental for neighbouring heathy tissue and immune

environment as a result of NF-κB-dependent pro-inflammatory mediators and cytokines secretion.   Other splicing events switch triggered by RBM7 might also be linked to tumor development. For example, we found RBM7 regulated alternative splicing of AKT2, which is involved in glucose metabolism (Hay, 2011). Further investigation is warranted to elaborate more splicing mechanisms of RBM7 in breast cancer.

In conclusion, we discovered that down-regulated RBM7 and its cognate elevated exon7-exclusion MFGE8 were tightly related to enhanced breast cancer metastatic potential and poor prognosis of patients. MFGE8-L, the canonical variant, possessed the capability to trigger STAT1 phosphorylation and inhibit the motility and invasion of breast cancer cells, whereas splice switch to MFGE8-S was found loss-of-function for this tumor-suppressive role. At another layer, RBM7 reduction led to NF-κB pathway activation, resulting in enhancement of breast tumor aggressiveness. Overall, our findngs implicated that activated NF-κB pathway worked in concert with MFGE8 splicing transition to spur metastasis in RBM7-depleted breast cancer. Additionally, although the detailed clinical significance underlying their cooperation need further in-depth investigation, we propose that RBM7 and its regulated splicing variants might serve as potential therapeutic biomarker and prognostic indicator in breast cancer patients.

Reference

Anczuków O, Akerman M, Cléry A, Wu J, Shen C, Shirole Nitin H, Raimer A, Sun S, Jensen Mads A, Hua Y, Allain Frédéric HT, Krainer Adrian R (2015) SRSF1-Regulated Alternative Splicing in Breast Cancer. *Molecular Cell* 60: 105-117

Baralle D (2005) Splicing in action: assessing disease causing sequence changes. *Journal of Medical Genetics* 42: 737-748

Bates SJHaDO (2008) VEGF-A splicing: the key to anti-angiogenic therapeutics? *NATURE REVIEWS* 7

Bugai A, Quaresma AJC, Friedel CC, Lenasi T, Düster R, Sibley CR, Fujinaga K, Kukanja P, Hennig T, Blasius M, Geyer M, Ule J, Dölken L, Barborič M (2019) P-TEFb Activation by RBM7 Shapes a Pro-survival Transcriptional Response to Genotoxic Stress. *Molecular Cell* 74: 254-267.e10

Chen C, Zhao S, Karnad A, Freeman JW (2018) The biology and role of CD44 in cancer progression: therapeutic implications. *Journal of Hematology & Oncology* 11

Chen Y, Hao X, Sun R, Wei H, Tian Z (2019) Natural Killer Cell–Derived Interferon-Gamma Promotes Hepatocellular Carcinoma Through the Epithelial Cell Adhesion Molecule–Epithelial-to-Mesenchymal Transition Axis in Hepatitis B Virus Transgenic Mice. *Hepatology* 69: 1735-1750

Cheruiyot A, Li S, Nonavinkere Srivatsan S, Ahmed T, Chen Y, Lemacon DS, Li Y, Yang Z, Wadugu BA, Warner WA, Pruett-Miller SM, Obeng EA, Link DC, He D, Xiao F, Wang X, Bailis JM, Walter MJ, You Z (2021) Nonsense-Mediated RNA Decay Is a Unique Vulnerability of Cancer Cells Harboring SF3B1 or U2AF1 Mutations. *Cancer Research* 81: 4499-4513

Falk S, Finogenova K, Melko M, Benda C, Lykke-Andersen S, Jensen TH, Conti E (2016) Structure of the RBM7–ZCCHC8 core of the NEXT complex reveals connections to splicing factors. *Nature*

Communications 7

Feng P-C, Ke X-F, Kuang H-L, Pan L-L, Ye Q, Wu J-B (2019) RETRACTED ARTICLE: BMP2

secretion from hepatocellular carcinoma cell HepG2 enhances angiogenesis and tumor growth in

endothelial cells via activation of the MAPK/p38 signaling pathway. *Stem Cell Research & Therapy* 10

Fousek K, Horn LA, Palena C (2021) Interleukin-8: A chemokine at the intersection of cancer plasticity,

angiogenesis, and immune suppression. *Pharmacology & Therapeutics* 219

Fukushima K, Satoh T, Sugihara F, Sato Y, Okamoto T, Mitsui Y, Yoshio S, Li S, Nojima S, Motooka D,

Nakamura S, Kida H, Standley DM, Morii E, Kanto T, Yanagita M, Matsuura Y, Nagasawa T,

Kumanogoh A, Akira S (2020) Dysregulated Expression of the Nuclear Exosome Targeting Complex

Component Rbm7 in Nonhematopoietic Cells Licenses the Development of Fibrosis. *Immunity* 52:

542-556.e13

Gopinathan G, Milagre C, Pearce OMT, Reynolds LE, Hodivala-Dilke K, Leinster DA, Zhong H,

Hollingsworth RE, Thompson R, Whiteford JR, Balkwill F (2015) Interleukin-6 Stimulates Defective

Angiogenesis. *Cancer Research* 75: 3098-3107

Harbour JW, Roberson EDO, Anbunathan H, Onken MD, Worley LA, Bowcock AM (2013) Recurrent

mutations at codon 625 of the splicing factor SF3B1 in uveal melanoma. *Nature Genetics* 45: 133-135

Hay N (2011) Akt isoforms and glucose homeostasis – the leptin connection. *Trends in Endocrinology*

& Metabolism 22: 66-73

Inoue D, Chew G-L, Liu B, Michel BC, Pangallo J, D'Avino AR, Hitchman T, North K, Lee SC-W,

Bitner L, Block A, Moore AR, Yoshimi A, Escobar-Hoyos L, Cho H, Penson A, Lu SX, Taylor J, Chen

Y, Kadoch C et al. (2019) Spliceosomal disruption of the non-canonical BAF complex in cancer.

Nature 574: 432-436

Li Z, Guo Q, Zhang J, Fu Z, Wang Y, Wang T, Tang J (2021) The RNA-Binding Motif Protein Family

in Cancer: Friend or Foe? *Frontiers in Oncology* 11

Liang Y, Zhang H, Song X, Yang Q (2020) Metastatic heterogeneity of breast cancer: Molecular

mechanism and potential therapeutic targets. *Seminars in Cancer Biology* 60: 14-27

Lubas M, Andersen Peter R, Schein A, Dziembowski A, Kudla G, Jensen Torben H (2015) The Human

Nuclear Exosome Targeting Complex Is Loaded onto Newly Synthesized RNA to Direct Early

Ribonucleolysis. *Cell Reports* 10: 178-192

Marimuthu S, Lakshmanan I, Muniyan S, Gautam SK, Nimmakayala RK, Rauth S, Atri P, Shah A,

Bhyravbhatla N, Mallya K, Grandgenett PM, Hollingsworth MA, Datta K, Jain M, Ponnusamy MP,

Batra SK (2022) MUC16 Promotes Liver Metastasis of Pancreatic Ductal Adenocarcinoma by

Upregulating NRP2-Associated Cell Adhesion. *Molecular Cancer Research* 20: 1208-1221

Medeiros B, Allan AL (2019) Molecular Mechanisms of Breast Cancer Metastasis to the Lung: Clinical

and Experimental Perspectives. *International Journal of Molecular Sciences* 20

Meola N, Domanski M, Karadoulama E, Chen Y, Gentil C, Pultz D, Vitting-Seerup K, Lykke-Andersen

S, Andersen Jens S, Sandelin A, Jensen Torben H (2016) Identification of a Nuclear Exosome Decay

Pathway for Processed Transcripts. *Molecular Cell* 64: 520-533

Montes M, Sanford BL, Comiskey DF, Chandler DS (2019) RNA Splicing and Disease: Animal

Models to Therapies. *Trends in Genetics* 35: 68-87

Pascal Preker JN, Susanne Kammler, Søren Lykke-Andersen, Marianne S. Christensen, Christophe K.

Mapendano, Mikkel H. Schierup, Torben Heick Jensen (2008) RNA Exosome Depletion Reveals

Transcription Upstream of Active Human Promoters. *SCIENCE* 322: 4

Rabbani SA, Mazar AP (2007) Evaluating distant metastases in breast cancer: from biology to

outcomes. *Cancer and Metastasis Reviews* 26: 663-674

Raymond A, Ensslin MA, Shur BD (2009) SED1/MFG-E8: A Bi-Motif protein that orchestrates diverse

cellular interactions. *Journal of Cellular Biochemistry* 106: 957-966

Shi Y (2017) Mechanistic insights into precursor messenger RNA splicing by the spliceosome. *Nature*

Reviews Molecular Cell Biology 18: 655-670

Siegel RL, Miller KD, Jemal A (2019) Cancer statistics, 2019. *CA: A Cancer Journal for Clinicians* 69:

7-34

Singh RK, Cooper TA (2012) Pre-mRNA splicing in disease and therapeutics. *Trends in Molecular*

Medicine 18: 472-482

Smith MA, Choudhary GS, Pellagatti A, Choi K, Bolanos LC, Bhagat TD, Gordon-Mitchell S, Von

Ahrens D, Pradhan K, Steeples V, Kim S, Steidl U, Walter M, Fraser IDC, Kulkarni A, Salomonis N,

Komurov K, Boultwood J, Verma A, Starczynowski DT (2019) U2AF1 mutations induce oncogenic

IRAK4 isoforms and activate innate immune pathways in myeloid malignancies. *Nature Cell Biology*

21: 640-650

Tang Y, Horikawa I, Ajiro M, Robles AI, Fujita K, Mondal AM, Stauffer JK, Zheng ZM, Harris CC

(2012) Downregulation of splicing factor SRSF3 induces p53β, an alternatively spliced isoform of p53

that promotes cellular senescence. *Oncogene* 32: 2792-2798

Valastyan S, Weinberg Robert A (2011) Tumor Metastasis: Molecular Insights and Evolving Paradigms.

Cell 147: 275-292

Warzecha CC, Sato TK, Nabet B, Hogenesch JB, Carstens RP (2009) ESRP1 and ESRP2 Are Epithelial

Cell-Type-Specific Regulators of FGFR2 Splicing. *Molecular Cell* 33: 591-601

Wilkinson ME, Charenton C, Nagai K (2020) RNA Splicing by the Spliceosome. *Annual Review of*

Biochemistry 89: 359-388

Wright CJ, Smith CWJ, Jiggins CD (2022) Alternative splicing as a source of phenotypic diversity.

Nature Reviews Genetics 23: 697-710

Xi P-W, Zhang X, Zhu L, Dai X-Y, Cheng L, Hu Y, Shi L, Wei J-F, Ding Q (2020) Oncogenic action of

the exosome cofactor RBM7 by stabilization of CDK1 mRNA in breast cancer. *npj Breast Cancer* 6

Yang C, Hayashida T, Forster N, Li C, Shen D, Maheswaran S, Chen L, Anderson KS, Ellisen LW,

Sgroi D, Schmidt EV (2011) The Integrin αvβ3-5 Ligand MFG-E8 Is a p63/p73 Target Gene in

Triple-Negative Breast Cancers but Exhibits Suppressive Functions in ER+ and erbB2+ Breast Cancers.

Cancer Research 71: 937-945

Zeng L, Xiao Q, Chen M, Margariti A, Martin D, Ivetic A, Xu H, Mason J, Wang W, Cockerill G, Mori

K, Li JY-s, Chien S, Hu Y, Xu Q (2013) Vascular Endothelial Cell Growth–Activated XBP1 Splicing in

Endothelial Cells Is Crucial for Angiogenesis. *Circulation* 127: 1712-1722

Zhou X, Wang R, Li X, Yu L, Hua D, Sun C, Shi C, Luo W, Rao C, Jiang Z, Feng Y, Wang Q, Yu S

(2019) Splicing factor SRSF1 promotes gliomagenesis via oncogenic splice-switching of MYO1B.

Journal of Clinical Investigation 129: 676-693

Zhou Z, Gong Q, Lin Z, Wang Y, Li M, Wang L, Ding H, Li P (2020) Emerging Roles of SRSF3 as a

Therapeutic Target for Cancer. *Frontiers in Oncology* 10

Axel Weber PW, Michael Kracht* (2010) Interleukin-1 (IL-1) Pathway. SCIENCESIGNALING 3:105

Guo Q, Jin Y, Chen X, Ye X, Shen X, Lin M, Zeng C, Zhou T and Zhang J (2024) NF-κB in biology

and targeted therapy: new insights and translational implications. Signal Transduction and Targeted

Therapy 9:53

Materials and Methods

Cell culture

Human breast cancer cell lines (MDA-MB-231, MCF7, HCC1937 and BT-549) and human embryonic kidney cell HEK293T in this article were obtained from the American Type Culture Collection (ATCC). MDA-MB-231 cells were cultured without carbon dioxide at 37°C in Leibovitz's L-15 medium (Gibco) contained 10% FBS (SINSAGE TECH). MCF7 and HEK293T cells were cultured with DMEM mediun (Gibco) supplemented with 10% FBS. HCC1937 cells were cultured with RPMI-1640 media (Gibco) supplemented with 10% FBS, while BT-549 was maintained with RPMI-1640 media (Gibco) supplemented with 10% FBS and 0.023 U/ml insulin. HUVECs were cultured with Endothelial Cell Medium (ScienCell) supplemented with ECGS (ScienCell), P/S solution (ScienCell) and 10% FBS (ScienCell). Except for MDA-MB-231, all the cells above were cultured in a 37 °C humidified incubator.

Stable breast cancer cell lines construction

To generate RBM7 overexpression vector, its coding regions (NM_016090.4) were cloned into pCDH-CMV-MCS-EF1-Puro with N-terminal Flag tag or HA-tag.To construct MFGE8-L/S vectors, the coding regions of MFGE8-L (NM_005928.4) and MFGE8-S (NM_001114614.3) were cloned into pCDH-CMV-MCS-EF1-Puro with N-terminal Flag tag. For RBM7 knockdown, target-specific shRNA (RBM7-sh1: 5'-CGAAAGGACTATGGATAACAT-3';RBM7-sh2:5'-CTACCTAGCAGATAGACA TTA-3') was cloned into pLKO.1 vector. And HEK293T cells were transfected these over-expression (or low-expression) lentivirus plasmisd together with lentivirus packaging plasmids pPAX2 and pMD2 according to the manufacturer's instructions. Viruses were collected 48 hours after transfection and then to infect breast cancer cell line with 8 ug/ml polybrene (beyotime biotechnology) for 24 hours. Positive cells were selected with 3 ug/ml puromycin.

Western Blot assay

Cells were lysed with RIPA buffer (10 mM Tris-HCl pH 7.5, 150 mM NaCl, 0.1% SDS, 1% Triton X-100, and 1% sodium deoxycholate) supplemented with protein tyrosine phosphatase inhibitor Na_3VO_4 (1mM), serine protease inhibitor PMSF (1mM) and protease inhibitor Cocktail. Immunoblotting assay was performed using standard methods. The following antibodies were used: anti-RBM7 (Sigma-Aldrich, HPA013993), anti-Flag (Sigma-Aldrich, F1804), anti-GAPDH (Proteintech, 10494-1-AP), anti-α-Tubulin (Proteintech, 11224-1-AP), anti-HA tag (Abcam, ab9110), anti-Phospho-NF-κB p65 (Cell Signaling Technology, #3033), anti-NF-κB p65 (Cell Signaling Technology, #8242), anti-IκBα (Cell Signaling Technology, #4814), anti-Phospho-IκBα (Ser32) (Cell Signaling Technology, #2859), anti-Phospho-stat1 (Tyr701) (Cell Signaling Technology, #8062), and anti-MFGE8 (R&D systems, MAB2767). The protein bands were detected with ECL enhanced chemiluminescence regent kits (NCM, Biotech), visualized using the MiniChemi™ Cheminescent image system (Beijing Sage Creation, China).

RNA isolation, RNA sequencing, and PCR

Total RNA from cells and tumor tissues was extracted using Trizol (Invitrogen) according to the manufacturer's instructions. Total mRNA was cleaned up using RNeasy Lipid Tissue Mini Kit (Qiagen, USA). And RNA-seq was carried out on platform of Novogene Illumina sequencing. cDNAs were obtained using *Evo M-MLV* Plus 1st Strand cDNA Synthesis Kit (Accurate Biotechnology, AG11615) with random primers. qRT-PCR was carried out using MonAmp™ Universal ChemoHS Specificity Plus qPCR Mix. GAPDH mRNA was examined as control for normalization. Gene expression relative to GAPDH was calculated using the formula $2^{-\Delta\Delta CT}$. For identified alternative splicing events, primers of the upstream and downstream of variable exons or introns were designed and PCR assay was performed. PCR products were subject to DNA agarose gel electrophoresis. AS events change of percent-spliced-in (PSI) values were the ratio of retained splices to total transcripts. The amount of each splicing isoform was measured with ImageJ. All the primers are showed in Table1.

Migration and invasion assays

For transwell migration assay, 1×10^5 MDA-MB-231 or other breast cancer cells (MCF7, BT-549 and HCC1937) were plated in the upper chamber of 6.5 mm insert (Costar, USA) with 8.0 μm polycarbonate membrane with 100 μL serum-free medium. For invasion assay, 1×10^5 breast cancer cells were seeded in the top chamber of each Matrigel-coated insert (BD Biosciences, USA). After 24 hours or 48 hours of incubation at 37 °C, migrated or invaded cells were fixed with 4% PFA and stained in 0.1% crystal violet. The cell number was counted by ImageJ software in 3 views from 3 independent experiments.

Wound healing

Knockdown RBM7 in MDA-MB-231 and 4T1 cells or control cells were implanted into 6-well culture dishes. When the cells grew about 90% confluence, lines of the same width were drawn on the bottom of the culture dish using autoclaved sterile tips. Images were captured at 0 and 36 hours after the wounding. Data shown were representative of three independent repeats.

Gelatin Degradation Assay

To evaluate the ability of breast cancer cells to form invadopodia and degrade matrix, QCM Gelatin Invadopodia Assay (ECM670-1, Merck Millipore) was performed according to the manufacturers' protocol. Briefly, 1×10^4 MFGE8-L/S-overexpressing or control MDA-MB-231 cells were suspended in L-15 medium with 10% FBS and seeded into the pre-treated 96-well plate. After 48 hours, the cells were washed twice with PBS. Fix cells with 3.7% formaldehyde for 30 minutes. Then these cells were stained respectively with TRITC-phalloidin and DAPI for 1 hour. The ability of cells to degrade the matrix was imaged using immunofluorescence microscopy (LEICA DMi8, Germany) and the area of invadopodia was measured using the Image J software.

Tube formation of HUVECs assay

Capillary tube formation assay was used to observe angiogenesis of HUVECs. HUVECs were seeded into 48-well plate coated with growth factor reduced Matrigel (BD Biosciences). HUVECs cells were treated with conditioned medium for 4-5

hours and then formation of capillary-like tubes was observed and stained with calcein for 30 minutes. Images of blood vessels were taken using fluorescence microscopy. And the tube length was analyzed by ImageJ software.

Immunofluorescence assay

After seeded on coverslips for 36 hours, the MDA-MB-231 cells were immersion fixed with 4% PFA for 20 minutes at room temperature, washed 3 times in PBS. Then the cells were treated with 0.2 % Triton-X100 in PBS for 10 minutes and blocked with 3% BSA for 10 minutes. Next the cells were immersed in primary antibody at 4℃ overnight, washed in PBS 3 times for 5 minutes each, and incubated with secondary antibody labelled with Alexa fluorophore 488/592 for 1 hour at RT. After washing 3 times in PBS, the cells were dyed with DAPI. In addition, antibodies targeting RBM7 (Sigma, HPA013993), and Flag (Sigma, 3165) were used in the assay. The pictures are captured with fluorescent microscope (Leica, USA).

RNA immunoprecipitation assay

RNA immunoprecipitation was performed as previously described. In brief, HEK293T cells were transfected Flag-RBM7 or control plasmid. After 48 hours of transfection, Flag-RBM7 or control cells were collected and cross-linked by 1% formaldehyde, homogenized in lysis buffer (50 mM Tris-HCl Ph 7.5, 0.4 M NaCl, 1 mM EDTA, 1mM DTT, 0.5% TritonX-100, 10% Glycerol supplemented with RNase inhibitor and broad-spectrum protease inhibitors Cocktail), and broke with an ultrasonic crusher. And solubilized cell lysate was obtained and precleared by ProteinG-agarose beads together with nonspecific tRNA to remove nonspecific binding. Then binding RNAs with RBM7 protein were precipitated with Anti-Flag M2 Affinity beads. Co-precipitated RNAs were subject to RT-PCR. Input controls (total RNAs) and Flag-controls were assayed simultaneously to prove that the detected signals were the result of RNAs specifically binding to RBM7. The gene-specific primers were designed MFGE8-RIP1-F:5'-tatttatgccccctcaccgc-3'; MFGE8-RIP1-R:5'-atggtggctgcctgtaat-3';MFGE8-RIP2-F:5'-ctgaggctcacccaagtag-3'; MFGE8-RIP2-R:5'-GTAGGATGCCACAAACTG-3'). GAPDH mRNA was as negative control (GAPDH-F 5'-GGGGAAGGTGAAGGTCG-G-3'; GAPDH-R 5'-TTGAGGTCAATGAAGGGG-3').

Antisense oligonucleotide (ASO)-mediated splicing blockade

MFGE8-ASOs were synthesized with a full-length phosphorothioate backbone and 2'-O-methyl modified ribose molecules (GenScriptBiotech Corp). ASO was transfected at final concentration of 500 nM using Lipofectamine3000 (Thermo Fisher Scientific) according to the manufacturer's instructions. The MCF7 cells were harvested for RNA and splicing assays 48 h after transfection.

Animal experiments

All procedures and experimental protocols involving mice were approved by the Institutional Animal Care and Use Committee (IACUC) of Dalian Medical University (approval no. AEE21015). Female BALB/C mice (6 weeks old, 4-5 mice per group as indicated in figures and/or figure legends) were injected into tail vein with murine 4T1 breast cancer cells (6×10^5 cells) to develop mice model of pulmonary metastasis. After about 3 weeks, all mice were euthanized, and lungs were harvested, fixed and

paraffin embedded. After lungs were H&E stained, the number of metastatic lesions were enumerated by pathologist using brightfield microscopy.

Clinical tissues samples collection

All human breast cancer tissues or normal tissue were obtained with written informed consent from patients or their guardians prior to participation in the study. The Institutional Review Board of the Dalian Medical University approved use of the tumor specimens in this study (approval no.PJ-KS-KY-2022-208).

Bioinformatics analysis

RBM7 expression data and survival data (including OS and DFS) were obtained from the METABRIC database using the Cbioportal web tool. Based on the expression level of RBM7, the patients were divided into four categories (i.e., Q1, Q2, Q3, Q4) by quartiles, and then log-rank was employed to compare the survival differences of patients in different categories. Cbioportal cited (Cerami E et al. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. Cancer Discov. 2012 May;2(5):401-4). Through the analysis of alternative splicing events in RBM7 knocked-down MDA-MB-231 cells compared to control, exon skipping events were screened out, and then selected the sequence of the 500nt region upstream of the alternative exon for subsequent base sequence enrichment analysis. The XSTREME tool provided by the online tool meme-suite (https://meme-suite.org/meme/) was used to perform Motif Discovery and Enrichment Analysis. E-value < 0.05 as the significance standard to output the results.

Statistical analyses

Statistical analyses and statistical graphs were conducted by GraphPad Prism 8 software. To compare the differences between the two variables, Student's T-test, one-way ANOVA with Dunnett multiple comparisons, and one-way ANOVA with Tukey's multiple comparisons according to the specific situation were applied. $*P < 0.05$, $**P < 0.01$, $***P < 0.001$, and $****P < 0.0001$ indicated the significant difference.

Data availability

The authors declare that all the data supporting the findings of this study are available within the article and its supplementary information files. RNA-seq data in this study have been deposited in Gene Expression Omnibus of NCBI with the accession code GSE248938.

Acknowledgements

This work was supported by the National Key R&D Program of China (2022YFA1104002, 2023YFE0117500 to YW); the National Natural Science Foundation of China (82225034, 81830088 to YW; 82103148 to YQ; 81872247 to WZ); the Science and Technology Innovation Foundation of Dalian (2022JJ11CG009

to YW), Liaoning Revitalization Talents Program (XLYC2202027 to YW);

Interdisciplinary Research Cooperation Project Team Funding of Dalian Medical

University (JCHZ2023008 to YQ); Joint Research Program of Natural Science

Foundation of Liaoning, China (2023-MSLH-023 to YQ); the Science and

Technology Innovation Talent Support Program of Dalian (2022RQ056 YQ); Dalian

High Level Talents Renovation Supporting Program (2019RQ097 to WZ).

Study Approval

The Institutional Animal Care and Use Committee of the Dalian Medical University

approved use of animal models in this study (approval no. AEE21015). All human

tumor tissues were obtained with written informed consent from patients or their

guardians prior to participation in the study. The Institutional Review Board of of the

First Affiliated Hospital of Dalian Medical University approved use the tumor

specimens in this study (approval no. PJ-KS-KY-2022-208).

Figure legends

Figure 1. Decreased RBM7 expression was associated with poor prognosis of breast cancer. (A-B) The correlation between RBM7 mRNA expression and overall survival (OS) or disease-free-survival (DFS) of breast cancer patients (n=1980) was analyzed based on the METABRIC dataset. The samples were divided into four equal parts, containing lower quartile Q1, median quartile Q2, upper quartile Q3 and higher quartile Q4 according to the expression of RBM7. Data are presented as means ± SD and P values were obtained by Mantel-Cox log-rank test. (C) The expression of RBM7 in primary breast carcinoma (n=1097) compared to normal tissues (n=114) was analyzed through UALCAN dataset. (D) Analysis of RBM7 expression in breast cancer lymph node metastases in comparison with breast cancer tissues with no lymph node metastasis based on TCGA dataset. BRCA samples were classified into N0 (No regional lymph node involvement) (n=515), and metastases in lymph node (N1-N3) (n=565). (E) Representative IHC images of RBM7 staining for patients at high (n=32) or low (n=58) clinical stages on tissue microarray of breast cancer specimens. Scale bars = 300 μm (top) or 30 μm (bottom). (F) Representative images of the IHC staining of RBM7 in a tissue microarray containing triple-negative breast cancer (n=119) and para-carcinoma tissues (n=20). Scale bars = 300 μm (top) and 30 μm (bottom). (G) Quantitative analysis of RBM7 expression according to IHC staining scores in primary breast cancer (n=12) and distant metastases (breast cancer lung metastases, n=4; breast cancer liver metastases, n=5). (H-I) Representative IHC images and quantitative analyses for RBM7 staining in 3 paired primary breast cancer and lymphatic metastases are shown. Scale bar = 100 μm. Data are presented as means ± SD and P values were obtained by unpaired Student t test (C-I). **$P < 0.01$, ***$P < 0.001$, ****$P < 0.0001$.

Figure 2. RBM7 negatively regulated breast cancer metastatic potential. (A) Gene Ontology analysis showed the significantly affected biological process upon RBM7 knockdown in MDA-MB-231 cells. (B) Functional association network of RBM7-regulated targets. The genes were analyzed using the STRING database, and subgroups are marked according to their function. (C) A heatmap showing the

qRT-PCR analysis of differentially expressed genes upon RBM7 knockdown in breast

cancer cells from three biological replicates. (D-F) 4T1 cells without or with RBM7

knockdown were injected into tail vein of immunodeficient BALB/c mice to establish

a lung metastasis model. Spontaneous pulmonary metastases were assessed after

about 3 weeks, the white arrow indicates lung lesions macroscopically (D). H&E

stained lung sections were quantified for the number of spontaneous metastatic

lesions from BALB/c mice (n=6, 5 and 9). Data are presented as mean ± SD and *P*

values were determined by one-way ANOVA with Dunnett's multiple comparisons

test (E). Representative images of H&E stained lung sections are shown. Top panel

scale bars indicate 7 mm; bottom panel scale bars indicate 300 μm (F) . (G-J) The

metastatic ability of breast cancer cells with RBM7 depletion (G) or ectopic

expression of RBM7 (I) were evaluated by transwell assay. Scale bars: 100 μm. Data

are mean ± SD from three random fields. *P* values were determined by one-way

ANOVA with Dunnett's multiple comparisons test (H) or unpaired Student's T test (J).

(K) Representative images of the tube formation of HUVEC cells treated with

conditional medium from RBM7-KD cells or control cells for 12h. Scale bars: 100

μm. Quantification of number of junctions formed by HUVEC was calculated by

Image J software. **P < 0.01, ***P < 0.001, ****P < 0.0001.

Figure 3. Global identification of alternative splicing events regulated by RBM7.

(A-E) RNA-seq analysis was performed on RBM7-knockdown MDA-MB-231 cells

or control cells, and the changes in splicing events were analyzed. (A) Volcano plot

illustrating up-regulated/down-regulated alternative splicing events upon RBM7

depletion. (B) Comparison of the differential splicing events in five types of AS

events affected by RBM7 depletion. (C) The relative fraction of splicing events

undergoing percent spliced in (PSI) increase or decrease induced by RBM7 depletion.

(D) Alternative splicing of MFGE8 was chosen to represent a decrease of PSI, and

numbers of exon junction reads were indicated. (E) The hit top enriched motifs

identified in the differential spliced genes regulated by RBM7. (F) RBM7-regulated

exon skipping events were identified by semiquantitative RT-PCR using

RBM7-depleted or control MDA-MB-231 cells (see also Figure 3-source data 1/

source data 2). The mean ± SD of PSI from 3 independent experiments was plotted,

with *P* values determined by one-way ANOVA with Dunnett's multiple comparisons

test. (G) Gene Ontology analysis showed the significantly enriched biological

processes of changed splicing events affected upon RBM7 knockdown in

MDA-MB-231 cells. (H) KEGG pathway enrichment analysis of functional

association network of the RBM7-controlled AS targets.

Figure 4. RBM7 knockdown promoted exon 7 skipping of MFGE8. (A)

Comparison of RBM7-binding candidates from the RNA immunoprecipitation (RIP)

sequencing (GSE144075 dataset) and two gene sets containing top 43 differentially

AS genes in RNA-seq data of breast cancer cells expressing RBM7 shRNA1/2 as

presented by venn diagram. The gene lists on the right were shown according to the

FDR *P* value of top genes with significantly changed AS in RNA-seq. (B) Upper:

schematics of human MFGE8 pre-mRNA (NM_005928.4). MFGE8-long isoform

included the exon 7 (hereafter referred to as MFGE8-L), whereas it is skipped in

MFGE8-short isoform (hereafter referred to as MFGE8-S). The arrow indicates the

direction of transcription. Lower: primers were designed on exon 6/8, and RT-PCR

was performed to identify PSI changes using RBM7-knockdown or control

MDA-MB-231 cells. The base peak diagram of sanger sequencing of RT-PCR results

showed the splicing junction sites (filled with gray). (C) Various breast cancer cells

lines with stable depletion of RBM7 or control were constructed. Alternative splicing

events of MFGE8 regulated by RBM7 were examined by semi-quantitative RT-PCR.

The knockdown efficiency of RBM7 expression was detected by Western Blotting.

(see also Figure 4-source data 1/ source data 2). The mean ± SD of PSIs from three

independent repeated experiments were plotted with *P* value calculated by one-way

ANOVA with Dunnett's multiple comparisons test. (D) Left, Schematics of MFGE8

mini-splicing reporter including exons 6-8 and two flanking introns in the 600bp

region upstream and downstream of 5'/3' splice sites; middle, RBM7 stable

knockdown or control cells were transfected with splicing reporter and collected for

RNA extraction after 48h. The splicing changes of exogenous MFGE8 were detected

by RT-PCR with specific primers on the reporter gene. The level of RBM7 protein

was detected in MCF7 cells by western blotting (see also Figure 4-source data 3/ source data 4). *P* value was calculated by one-way ANOVA with Dunnett's multiple comparisons test from three repeated experiments. (E) As shown in schematic diagram, orange box indicates cassette exon 7 and primers were designed in the putative binding regions P1 and P2. Binding of MFGE8 pre-mRNA with RBM7 was examined by RIP assay in HEK293T cells expressing Flag-RBM7. (see also Figure 4-source data 5/ source data 6) (F) Upper: the red line in diagram indicates ASOs targeting region which contain UUUCUU residues; down: MCF7 and BT-549 cells were transfected with ASOs targeting MFGE8 pre-mRNA for 48h and then applied for RT-PCR identification (see also Figure 4-source data 7/ source data 8). *$P < 0.05$, **$P < 0.01$, ***$P < 0.001$, and ****$P < 0.0001$.

Figure 5. MFGE8-L inhibited breast cancer cell migration and invasion. (A) The function of MFGE8-L/S on breast cancer cell migration and invasion were evaluated by transwell assays. (B) Gelatin degradation assay was performed to test the effect of RBM7 knockdown on invadopodia function. 10000 cells were plated onto FITC-gelatin substrates (Green) and cultured for 48 h. Representative images are shown (red, Cy3-phalloidin; blue, DAPI) and the degraded areas were quantified by Image J software. Scar bars= 50 μm. *P* values were determined by one-way ANOVA with Tukey's multiple comparisons test (n = 3). (C) The subcellular localizations of MFGE8 isoforms. MDA-MB-231 cells were transfected with Flag-tagged MFGE8-L or MFGE8-S, and visualized with immunofluorescence assay. Scale bar = 10 μm. The percent of cells with foci were quantified and plotted (about 80 cells were captured and quantified in both samples). (D-E) The KEGG pathway and Gene ontology analysis of genes up-regulated by MFGE-S compared with MFGE8-L. (F) The expression levels of p-STAT1(Tyr701) in HCC1937 cells overexpressing MFGE8-L/S was detected by western blotting. (see also Figure 5-source data 1/ source data 2). *P* values were determined by one-way ANOVA with Tukey's multiple comparisons test. **$P < 0.01$, ****$P < 0.0001$.

Figure 6. RBM7 knockdown enhanced aggressiveness of breast cancer relying on MFGE8 splicing switch to the short variant and NF-κB pathway activation. (A)

Breast cancer cells MDA-MB-231 and MCF7 were cotransfected with shRNA

targeting RBM7 or control and shRNA-resistant Flag-MFGE8-L vectors and then

subjected to transwell migration assay or invasion assay. Migrated or invaded cells

were counted from random sites of the transwell with Image J. *P* values were

determined by one-way ANOVA with Tukey's multiple comparison test. (B)

Representative transwell analysis of migrative/invasive capability of breast cancer

cells transfected with 500 nM ASO directed against RBM7-binding region in MFGE8

pre-mRNA. Cells migrating through transwell membrane with matrigel or not were

imaged and representative images are shown (upper) along with quantification (down).

P values were determined by one-way ANOVA with Tukey's multiple comparison test.

(C) The expression of metastasis-related factors in HCC1937 and BT-549 cells stably

expressing MFGE8-L, MFGE8-S or control vector was measured by real-time PCR

assay. (D) NF-κB signaling pathway was enrichment via KEGG analysis in RBM7

knockdown breast cancer cells. (E) Western blotting showed the expression of p65,

p-p65, IκBα and p-IκBα in NF-κB pathway in RBM7-depleted or control

MDA-MB-231 and BT-549 breast cancer cells (see also Figure 6-source data 1/

source data 2). (F) Western blotting was performed to test the expression of RBM7,

p65 and p-p65 in RBM7-depleted MDA-MB-231 cells treated with or without NF-κB

inhibitor PDTC 10 μm for 48h (see also Figure 6-source data 3/ source data 4). (G)

Representative images of the tube formation of HUVEC cells treated with conditional

medium from RBM7-depleted   MDA-MB-231 in the presence or absence of 10 μm

NF-κB inhibitor PDTC. Scale bars: 100 μm. Quantification of junctions of endothelial

network was conducted by ImageJ software. *P* values were determined by one-way

ANOVA with Tukey's multiple comparison test (n = 3). $*P < 0.05$, $**P < 0.01$,

$***P < 0.001$, and $****P < 0.0001$.

Figure 7. Splicing shift of MFGE8 toward exon7 exclusion was negatively

correlated with RBM7 expression in patients with breast cancer. (A) MFGE8

exon7 expression level was analyzed in breast carcinoma in comparison with normal

tissues based on the TCGA dataset. Plotted are the mean ± SD from 114 normal

tissues and 1094 tumor tissues, with **** $P<0.0001$ as determined by unpaired Student' t test. (B) The splicing alteration of MFGE8 in 6 pairs of primary breast cancer tissues and adjacent normal tissues was examined using RT-PCR (see also Figure 7-source data 1/ source data 2). The quantification of PSI vales was based on relative band intensities using Image J software. (C) The splicing alteration of MFGE8 in primary breast cancer tissues and corresponding lymph node metastases was identified by RT-PCR assays (see also Figure 7-source data 3/ source data 4). The quantification of PSI vales was determined by Image J software. (D) The correlation between the expression of MFGE8 exon7 and overall survival (OS) of breast cancer patients was analyzed based on the TCGA dataset. P value was assessed with Mantel-Cox log-rank test. (E) Correlation of RBM7 expression with MFGE8 exon7 levels was analyzed using Pearson's correlation coefficient method. (F) The mechanistic model of how RBM7 regulated metastasis of breast cancer through regulating MFGE8 splicing switch.

Figure 1-figure supplement 1. The low expression of RBM7 in breast cancer was positively correlated with poor prognosis of breast cancer patients. (A) The correlation between RBM7 expression and overall survival of breast cancer patients based on data from the Kaplan–Meier Plotter cohort. (B-C) METABRIC analysis of the correlation between RBM7 expression and overall survival (OS) or disease-free survival (DFS) of breast cancer patients based on the METABRIC dataset. P value was obtained by log-rank test (A, B, C). (D) RBM7 mRNA level was analyzed in breast invasive carcinoma based on lymph nodal metastasis status from the UALCAN dataset. P value was calculated by unpaired Student's t-test by comparing different types of lymph node metastasis of breast cancer with normal tissue (n=114). N, Lymph node metastases: N0, no metastases (n=516); N1, 1－3 metastases (n=362); N2, 4－9 metastases (n=120); N3, 10 or more metastases (n=77).

Figure 2-figure supplement 1. RBM7 depletion promoted breast cancer cell migration and invasion. (A) The mouse cell line 4T1 with RBM7 knockdown was constructed. The knockdown efficiency of RBM7 was detected by real-time quantitative PCR. (B) The migration and invasion ability of RBM7-knockdown 4T1

cells was detected by transwell chambers coated with or without Matrigel. (C) The metastatic capability of RBM7-depleted 4T1 cells was evaluated by wound healing assay. (D) Western blotting showed the expression of RBM7 in MDA-MB-231 and MCF7 cells with RBM7 depletion (see also Figure 2-figure supplementary 1-source data 1/ source data 2). (E) The metastatic capability of RBM7 on MDA-MB-231 cells was evaluated by wound healing assay. (F) Western blotting showed the expression of RBM7 in various breast cancer cell lines with ectopic expression of RBM7 (see also Figure 2-figure supplementary 1-source data 3/ source data 4). (G) The metastatic ability of BT-549 cells and HCC1937 cells upon RBM7 overexpression was evaluated by transwell assay. Data are mean ± SD from three random fields. *P* values were determined by unpaired Student's T test (G). *P* value was calculated by one-way ANOVA with Dunnett's multiple comparisons test (A, B, C, E).

Figure 3-figure supplement 1. RBM7 had a potential role in regulating alternative splicing of breast cancer cells. (A) Immunofluoresence shows endogenous RBM7 subcellular localization in breast cancer cell lines. (B) Semi-quantitative RT-PCR was performed to validate the RI (intron retention) and A5E (alternative 5' splice site exon) splicing events in MDA-MB-231 cells with RBM7 knockdown compared with vector control (see also Figure 3-figure supplementary 1-source data 1/ source data 2). *P* value was calculated by one-way ANOVA with Dunnett's multiple comparisons test. (C) MFGE8 min-splicing reporters with mutation in the RBM7 binding site or a non-specific binding were generated and shown in cartoon (left); RT-PCR assays were performed to identify the splicing outcomes of MFGE8 reporter while RBM7 was depleted in breast cancer cells (right). See also Figure 3-figure supplementary 1-source data 3/ source data 4.

Figure 5-figure supplement 1. MFGE8-L suppressed breast cancer cells migration and invasion. (A) The protein expression of MFGE8 was determined by western blot in MDA-MB-231 cells with over-expression or knockdown of MFGE8-L. SE denotes long exposure and LE denotes long exposure (see also Figure 5-figure supplementary 1-source data 1/ source data 2). (B-C) Transwell assay with Matrigel or not was performed using MDA-MB-231 cells with overexpression (B) or depletion

(C) of MFGE8. *P* value was calculated by unpaired Student's t test. (D) Western blot

showed the extrinsic expression of MFGE8-L or MFGE8-S isoform in MDAMB-231

and HCC1937 transfected with Flag-MFGE8-L, Flag-MFGE8-S or empty vectors

respectively (see also Figure 5-figure supplementary 1-source data 3/ source data 4).

The red arrow indicates the position of MFGE8-L/S protein band . (E) MDA-MB-231

cell line over-expressing MFGE8-L, MFGE8-S or vector control were analyzed by

wound healing assay. The scratch healing rates were measured in triplicate

experiments with mean$\pm$SD plotted (*P* values were calculated by one-way ANOVA

with Tukey's multiple comparisons test). (F) Schematic diagram of the domain

composition of two MFGE8 isoforms. Upper: the full-length variant with exon7

indicated by yellow square; down: the truncated variant with exon7 skipping. (G) The

model structure of two MFGE8 isoforms was implemented using SwissModel

software. The F5/8 type C2 protein domain excluded from MFGE8-S variant was

marked in red. (H) Transcriptome sequencing was performed using MDA-MB-231

cells overexpressing MFGE8-L or MFGE8-S. Venn diagram of 782 MFGE8-L

down-regulated genes compared to control and 647 MFGE8-S up-regulated genes

compared to MFGE8-L.

Figure 6-figure supplement 1. MFGE8-L/S isoforms had no effect on NF-κB

pathway in breast cancer cells. (A) MCF7 and MDA-MB-231 cells stably

expressing shRNA against RBM7 with ectopic MFGE8-L or control vector were

generated. The expression of RBM7 and MFGE8 was measured by qRT-PCR assay.

(B) RT-PCR quantification of two MFGE8 isoforms after transfecting breast cancer

cells with 500 nM ASO directed against RBM7-binding region in MFGE8 pre-mRNA.

See also Figure 6-figure supplementary 1-source data 1/ source data 2. (C) HCC1937

and BT-549 cells stably expressing MFGE8-L, MFGE8-S or control were generated.

The expression of MFGE8 was measured by qRT-PCR assay. (D) Representative

images of the tube formation of HUVEC cells treated with conditional medium from

breasst cancer cells with ectopic expression of MFGE8-L or MFGE8-S for 48h. Scale

bars: 100 μm. Quantification of number of junctions formed by HUVEC was

determined by Image J software. (E) Western blotting results showed the

phosphorylation and expression of p65 in RBM7-overexpressing breast cancer cells

(see also Figure 6-figure supplementary 1-source data 3/ source data 4). (F) Western

blotting was performed to detect the indicated proteins in breast cancer cells with

up-expression of MFGE8-L/S (see also Figure 6-figure supplementary 1-source data

5/ source data 6). (G) The mRNA levels of IL-1β in MDA-MB-231 or BT549 cells

with stable RBM7 knockdown or control vector were examined by qRT-PCR

approach. (H) Supernatants from RBM7-knockdown MDA-MB-231 or BT549 cells

were collected and protein expression of IL-1β was measured by ELISA kit. (I) The

knockdown efficiency of RBM7 in two breast cancer cell lines were determined by

qRT-PCR approach. (J) RBM7-depleted MDA-MB-231 cells were treated with or

without NF-κB inhibitor PDTC 10 μm for 48h. Then Transwell assays were

performed to assess cell migration ability. *P* values were calculated by one-way

ANOVA with Tukey's multiple comparisons.

Figure 7-figure supplement 1. MFGE8 exon7 skipping is enhanced in a variety of

cancers. (A) The PSI values for the MFGE8 exon7 were analyzed in various cancers

according to TCGA SpliceSeq dataset. *P* value was calculated by unpaired Student's

t-test. (B) Immunoprecipitation assay was performed in breast cancer cells expressing

HA-RBM7 and Flag-SF3B2 or empty vector. The Flag-tagged precipitated complexes

and lysates were analyzed through western blotting. See also Figure 7-figure

supplementary 1-source data 1/ source data 2. (C) The splicing shift of MFGE8 upon

SF3B2 knockdown in breast cancer cells was examined by RT-PCR approach. See

also Figure 7-figure supplementary 1-source data 3/ source data 4. The mean ± SD of

PSI values derived from three independent replicates is shown. (D) The SF3B2

knockdown efficiency was examined by qRT-PCR assays. *P* value was calculated by

unpaired Student's t-test.

1014

1015

1016

1017

Legends for source data

Figure 3—source data 1

Original file for the RT-PCR analysis of splicing shifts in Figure 3F (MFGE8, AKT2, ZMYN8 or MBNL1).

Figure 3—source data 2

PDF containing Figure 3F and original scans of the relevant RT-PCR analysis of splicing shifts (MFGE8, AKT2, ZMYN8 or MBNL1) with highlighted bands and sample labels.

Figure 4—source data 1

Original file for the RT-PCR analysis of MFGE8 splicing switch and Western blot identification of RBM7 knockdown in various breast cancer cell lines in Figure 4C.

Figure 4—source data 2

PDF containing Figure 4C and original scans of the relevant RT-PCR analysis for MFGE8 splicing switch and Western blot identification of RBM7 knockdown with highlighted bands and sample labels.

Figure 4—source data 3

Original file for the RT-PCR analysis of splicing switch in MFGE8 mini-splicing reporter and Western blot identification of RBM7 knockdown in Figure 4D.

Figure 4—source data 4

PDF containing Figure 4D and original scans of the relevant RT-PCR analysis of splicing switch in MFGE8 mini-splicing reporter and Western blot identification of RBM7 knockdown with highlighted bands and sample labels.

Figure 4—source data 5

Original file for the RT-PCR analysis of RBM7 protein- MFGE8 pre-mRNA binding from RIP assay in Figure 4E.

Figure 4—source data 6

PDF containing Figure 4E and original scans of the RT-PCR analysis of RBM7

protein- MFGE8 pre-mRNA binding from RIP assay with highlighted bands and sample labels.

Figure 4—source data 7

Original file for the RT-PCR analysis of MFGE8 splicing switch upon ASO treatment in Figure 4F.

Figure 4—source data 8

PDF containing Figure 4F and original scans of the relevant RT-PCR analysis for splicing switch of MFGE8 variants with highlighted bands and sample labels.

Figure 5—source data 1

Original file for the Western blot analysis in Figure 5F (anti-P-STAT1, anti-MFGE8 and anti-GAPDH).

Figure 5—source data 2

PDF containing Figure 5F and original scans of the relevant Western blot analysis (anti-P-STAT1, anti-MFGE8 and anti-GAPDH) with highlighted bands and sample labels.

Figure 6—source data 1

Original file for the Western blot analysis in Figure 6E (anti-p65, anti-P-p65, anti-IκB, anti-P-IκB, anti-RBM7 and anti-GAPDH).

Figure 6—source data 2

PDF containing Figure 6E and original scans of the relevant Western blot analysis (anti-p65, anti-P-p65, anti-IκB, anti-P-IκB, anti-RBM7 and anti-GAPDH)) with highlighted bands and sample labels.

Figure 6—source data 3

Original file for the Western blot analysis in Figure 6F (anti-p65, anti-P-p65, anti-RBM7 and anti-GAPDH)).

Figure 6—source data 4

PDF containing Figure 6F and original scans of the relevant Western blot analysis

(anti-p65, anti-P-p65, anti-RBM7 and anti-GAPDH)) with highlighted bands and

sample labels.

Figure 7—source data 1

Original file for the RT-PCR analysis of MFGE8 splicing switch in primary breast

cancer tissues and adjacent normal tissues in Figure 7B.

Figure 7—source data 2

PDF containing Figure 7B and original scans of the relevant RT-PCR analysis for

MFGE8 splicing switch with highlighted bands and sample labels in primary breast

cancer tissues and adjacent normal tissues .

Figure 7—source data 3

Original file for the RT-PCR analysis of MFGE8 splicing switch in primary breast

cancer tissues and corresponding lymph node metastases in Figure 7C.

Figure 7—source data 4

PDF containing Figure 7C and original scans of the relevant RT-PCR analysis for

splicing switch of MFGE8 variants with highlighted bands and sample labels in

primary breast cancer tissues and corresponding lymph node metastases.

1088

Figure 2-figure supplement 1—source data 1

Original file for the Western blot analysis in Figure 2-figure supplement 1D

(anti-RBM7 and anti-tubulin).

Figure 2-figure supplement 1—source data 2

PDF containing Figure 2-figure supplement 1D and original scans of the relevant

Western blot analysis (anti-RBM7 and anti-tubulin) with highlighted bands and

sample labels.

Figure 2-figure supplement 1—source data 3

Original file for the Western blot analysis in Figure 2-figure supplement 1F

(anti-RBM7 and anti-tubulin).

Figure 2-figure supplement 1—source data 4

PDF containing Figure 2-figure supplement 1F and original scans of the relevant

Western blot analysis (anti-RBM7 and anti-tubulin) with highlighted bands and

sample labels.

Figure 3-figure supplement 1—source data 1

Original file for the RT-PCR analysis of MAP7D1 or HNRNPC splicing switch in

Figure 3-figure supplement 1B.

Figure 3-figure supplement 1—source data 2

PDF containing Figure 3-figure supplement 1B and original scans of the relevant

RT-PCR analysis for splicing switch of MAP7D1 or HNRNPC variants with

highlighted bands and sample labels.

Figure 3-figure supplement 1—source data 3

Original file for RT-PCR analysis of splicing switch of MFGE8 minigene reporter in

Figure 3-figure supplement 1C.

Figure 3-figure supplement 1—source data 4

PDF containing Figure 3-figure supplement 1C and original scans of the relevant

RT-PCR analysis of MFGE8 splicing switch with highlighted bands and sample

labels.

Figure 5-figure supplement 1—source data 1

Original file for the Western blot analysis in Figure 5-figure supplement 1A

(anti-MFGE8 and anti-GAPDH).

Figure 5-figure supplement 1—source data 2

PDF containing Figure 5-figure supplement 1A and original scans of the relevant

Western blot analysis (anti-MFGE8 and anti-GAPDH) with highlighted bands and

sample labels.

Figure 5-figure supplement 1—source data 3

Original file for the Western blot analysis in Figure 5-figure supplement 1D (anti-Flag

and anti-GAPDH).

Figure 5-figure supplement 1—source data 4

PDF containing Figure 5-figure supplement 1D and original scans of the relevant
Western blot analysis (anti-Flag and anti-GAPDH) with highlighted bands and sample
labels.

Figure 6-figure supplement 1—source data 1

Original file for the RT-PCR analysis of MFGE8 splicing switch upon ASO treatment
in Figure 6-figure supplement 1B.

Figure 6-figure supplement 1—source data 2

PDF containing Figure 6-figure supplement 1B and original scans of the relevant
RT-PCR analysis for splicing switch of MFGE8 variants with highlighted bands and
sample labels.

Figure 6-figure supplement 1—source data 3

Original file for the Western blot analysis in Figure 6-figure supplement 1E (anti-p65,
anti-P-p65, anti-IκB, anti-P-IκB, anti-Flag and anti-GAPDH).

Figure 6-figure supplement 1—source data 4

PDF containing Figure 6-figure supplement 1E and original scans of the relevant
Western blot analysis (anti-p65, anti-P-p65, anti-IκB, anti-P-IκB, anti-Flag and
anti-GAPDH)) with highlighted bands and sample labels.

Figure 6-figure supplement 1—source data 5

Original file for the Western blot analysis in Figure 6-figure supplement 1F (anti-p65,
anti-P-p65, anti-RBM7 and anti-GAPDH)).

Figure 6-figure supplement 1—source data 6

PDF containing Figure 6-figure supplement 1F and original scans of the relevant
Western blot analysis (anti-p65, anti-P-p65, anti-RBM7 and anti-GAPDH)) with
highlighted bands and sample labels.

Figure 7-figure supplement 1—source data 1

Original file for the Western blot analysis of Flag-tagged precipitated complexes in

Figure 7-figure supplement 1B (anti-Flag, anti-HA and anti-GAPDH).

Figure 7-figure supplement 1—source data 2

PDF containing Figure 7-figure supplement 1B and original scans of the relevant

Western blot analysis of Flag-tagged precipitated complexes (anti-Flag, anti-HA and

anti-GAPDH) with highlighted bands and sample labels.

Figure 7-figure supplement 1—source data 3

Original file for RT-PCR analysis of splicing switch of MFGE8 upon SF3B2

knockdown in Figure 7-figure supplement 1C.

Figure 7-figure supplement 1—source data 4

PDF containing Figure 7-figure supplement 1C and original scans of the relevant

RT-PCR analysis of MFGE8 splicing switch upon SF3B2 knockdown with

highlighted bands and sample labels.

A

MFGE8_exon7 expression level

Tumor Normal

****

B

Patient: #1 #2 #3 #4 #5 #6

MFGE8

GAPDH

L
S

T N T N T N T N T N T N

400bp
200bp

100bp

PSI (%)

#1 #2 #2 #4 #5 #6

BRCA
Normal

C

Patient: #1 #2 #3 #4 #5

BC LM BC LM BC LM BC LM BC LM

MFGE8
(LE)

MFGE8
(SE)

GAPDH

L
S

L
S

400bp
200bp

400bp
200bp

100bp

PSI (%)

#1 #2 #3 #4 #5

BRCA
LM

D

Probability of Overall Survival

Days

MFGE8_exon7_high
MFGE8_exon7_low

P=0.13

E

Expression level of MFGE8 exon7

Expression level of RBM7

R=0.36
P<0.001

F

RBM7-high expression

6 RBM7 7 8

MFGE8 Pre-mRNA

Splicing

MFGE8-L 6 7 8

Translation

MFGE8-L

STAT1

Metastasis supression

RBM7

p65

Inhibition of NF-κB targets transcription

RBM7-low expression

6 RBM7 7 8

MFGE8 Pre-mRNA

Splicing

MFGE8-S 6 8

Translation

MFGE8-S

p65

MMP9 IL6 IL8 IL1β

Promotion of NF-κB targets transcription

Breast cancer Metastasis