Title: Kernels of Motor Memory Formation: Temporal Generalization in

Bimanual Adaptation

Abbreviated title: Temporal Generalization in Bimanual Adaptation

Ian S. Howard[1*], Sae Franklin[2] and David W. Franklin[2,3,4]

[1] School of Engineering, Computing and Mathematics, University of Plymouth,

Plymouth, United Kingdom.

[2] Neuromuscular Diagnostics, Department Health and Sport Sciences, TUM

School of Medicine and Health, Technical University of Munich, Germany.

[3] Munich Institute of Robotics and Machine Intelligence (MIRMI), Technical

University of Munich, Munich, Germany.

[4] Munich Data Science Institute (MDSI), Technical University of Munich,

Munich, Germany.

*Correspondence should be addressed to: ISH: ian.howard@plymouth.ac.uk

Telephone: +44 1752 586324
18

Journal Section: Behavioral/Cognitive

Number of pages: 47 (double spaced)

Number of figures: 5

Number of tables: 0

Number of words for abstract: 220

Number of words for introduction: 647

Number of words for discussion: 1492

Statement Financial interests or conflicts of interest: The authors declare that

they have no financial, personal, or professional interests that could be

construed to have influenced the paper.

Acknowledgements: Financial support was provided by the Technical

University of Munich and the SECAM at the University of Plymouth. During

the preparation of this work, ISH utilized ChatGPT-4o for proofreading. We

thank Yiming Liu for assistance with the Generalized Additive Mixed Models.

Keywords: motor learning, bimanual context, temporal generalization, dynamic

learning, human, viscous curl field

Abstract

In daily life, we coordinate both simultaneous and sequential bimanual movements to manipulate objects. Our ability to rapidly account for different object dynamics suggests there are neural mechanisms to quickly deal with them. Here we investigate how actions of one arm can serve as a contextual cue for the other arm, and facilitate adaptation. Specifically, we examine the temporal characteristics that underlie motor memory formation and recall, by testing the contextual effects of prior, simultaneous, and post contralateral arm movements. To do so, we measure their temporal generalization in three bimanual interference tasks. Importantly, the timing context of the learned action plays a pivotal role in the temporal generalization. While motor memories trained with post adaptation contextual movements generalize broadly, motor memories trained with prior contextual movements exhibit limited generalization, and motor memories trained with simultaneous contextual movements do not generalize to prior or post contextual timings. This highlights temporal tuning in sensorimotor plasticity: different training conditions yield substantially different temporal generalization characteristics. Since these generalizations extend far beyond any variability in training times, we suggest that the observed differences may stem from inherent differences in the use of prior, current and post-adaptation contextual information in the generation of natural behavior. This would imply differences in the underlying neural circuitry involved in learning and executing the corresponding coordinated bimanual movements.

Significance Statement

This study addresses a fundamental question in the field of sensorimotor neuroscience of how multiple movements are temporally linked within a single motor memory. We examine the temporal generalization of motor memory formation by varying the timing of contextual movements associated with a learned motor memory across a range of prior, current, and post adaptation movement times using a bimanual motor learning task. We observed distinct patterns of temporal generalization based on whether the contextual movements occurred prior to, simultaneously with, or after the adaptation movement. For the first time, our findings reveal that the timing of contextual movements is crucial in the formation and generalization of motor memories.

Introduction

Many everyday tasks require coordinated actions of our two arms. While some tasks, such as opening a jar, require the simultaneous control of both arms, others, such as juggling, involve an initial movement of one hand followed by a subsequent movement of the other hand. Although juggling involves the manipulation of objects that vary in size, weight, or shape, professional jugglers demonstrate impressive hand-eye coordination, producing rapid and accurate object trajectories. Such effective manipulation requires that our sensorimotor system accurately accounts for the dynamics of the objects currently being manipulated, as well as those to be manipulated in the future. Motor learning of

such object dynamics during bimanual movement is therefore a fundamental

process underpinning human skills.

Adaptation to dynamics occurs through the formation of a motor memory

(Lackner and Dizio, 1994; Shadmehr and Mussa-Ivaldi, 1994; Kawato, 1999;

Shadmehr et al., 2010). However, these motor memories are not limited to the

specific tasks learned, but they generalize to similar movements and tasks,

allowing flexible and adaptable motor control (Conditt et al., 1997;

Thoroughman and Shadmehr, 2000; Donchin et al., 2003; Orschiedt and

Franklin, 2023). As this generalization is thought to stem from learning via

neural basis functions, measurement of the extent and type of generalization

allows for the investigation of the tuning of these neural basis functions within

the sensorimotor control system. Here we use the term temporal kernel to

describe the temporal tuning of the generalization, which may arise from the

underlying neural basis functions.

To study the formation, selection and generalization of motor memories and the

effect of context, two opposing force fields are often randomly presented during

reaching movements. Without any additional cues, this random presentation

results in interference: learning one force field interferes with the memory of the

other (Shadmehr and Brashers-Krug, 1997; Karniel and Mussa-Ivaldi, 2002;

Osu et al., 2004; Krakauer and Shadmehr, 2006). However, the association of

the force field with certain contextual cues allows learning of the two

independent motor memories and appropriate switching of the memories,

termed dual adaptation (Nozaki et al., 2006; Hirashima and Nozaki, 2012;

Howard et al., 2012). This paradigm allows us to examine which contextual cue

types can select and switch between different motor memories within the

sensorimotor control system, and identify those cues that do not have such

functions (Howard et al., 2013; Forano et al., 2021).

While the tuning of neural basis functions that underly generalization is typically

examined as a function of both adaptation and contextual movement directions

(Thoroughman and Shadmehr, 2000; Donchin et al., 2003; Howard and

Franklin, 2015, 2016; Sarwary et al., 2015), and of contextual movement

kinematics (Howard et al., 2020; Orschiedt and Franklin, 2023), the temporal

generalization of these motor memories, examined by varying the timing of

contextual movements, has yet to be investigated.

Here we use an interference task to examine how different temporal

relationships between motion of the two hands affect adaptation, specifically

focusing on prior, simultaneous, and post contextual motions. In unimanual

tasks, both prior contextual lead-in movements and post adaptation contextual

follow-through movements allow dual adaptation (Howard et al., 2012, 2015;

Sheahan et al., 2016). While in bimanual tasks, simultaneous (Nozaki et al.,

2006; Howard et al., 2010) and prior (Gippert et al., 2023) contextual

movements support dual adaptation, little is known about the role of post

adaptation contextual movements. Moreover, we have no measure of the

temporal generalization of these motor memories: how learning at one timing

relation transfers to other timing relations. Here, we first investigate how prior,

simultaneous and post adaptation contextual movements of the contralateral

hand affect learning opposing force fields. For each trained timing relationship,

we then examine the recall of this learned motor memory to a range of different

temporal relations, enabling us to map out the temporal generalization of these

motor memories across different temporal relations, and extract the temporal

kernels of motor memory formation and recall.

Methods

Experimental Setup

Participants. Twenty-six human participants were randomly allocated to three

experiments. Two participants were excluded from further analysis purely

based on their adaptation to the curl force field, leaving twenty-four.

Specifically, one participant was excluded as they did not adapt at all to the

viscous curl force field (final adaptation level at the end of day 2 < 10%). The

other participant was removed as the kinematic error reduced to below 5 mm

within the first 100 trials in the force field while force compensation remained

variable and low throughout the experiment, suggesting either co-contraction

or more cognitive strategies involved in adaptation. The generalization data of

these two participants was not analyzed. This resulted in eight participants for

each of Experiment 1 (3 females, average age 24.6 ± 2.4 years), Experiment 2

(5 females, average age 28.4 ± 8.5 years), and Experiment 3 (3 females,

average age 25.6 ± 2.7 years). All participants were right-handed as determined

by the Edinburgh Handedness Questionnaire (Oldfield, 1971) and naïve to the

purposes of the experiment. Prior to the experiment, each participant provided

written informed consent. The experiments were conducted for Plymouth

participants (n=3), in accordance with ethics approval by the University of

Plymouth ethics committee and conducted for Munich participants (n=23) in

accordance with ethics approval by the Ethics Committee of the Medical faculty

of the Technical University of Munich. Each participant completed one

experiment, consisting of two three–hour experimental sessions over two

consecutive days, totaling approximately six hours for each participant.

Figure 1 about here

Apparatus. Participants performed all experiments using a bimanual vBOT

planar robotic manipulandum paired with an associated virtual reality system

(Howard et al., 2009b) (Fig. 1A). Handle positions were calculated based on

measures using optical motor encoders sampled at 1000 Hz, and torque-

controlled motors applied end-point forces. A force transducer (Nano 25; ATI),

mounted under the left handle, measured the applied forces. These force

signals were low-pass filtered at 500 Hz using 4th-order analogue Bessel filters

before digitization. To minimize body movement, participants were comfortably

seated in a sturdy chair positioned in front of the apparatus and were firmly

strapped against the backrest with a four-point seatbelt.

During the experiment, participants held the left and right robot handles with

their respective hands, with their forearms supported by air sleds. This setup

effectively restricted arm movements to the horizontal plane and provided full

support of the mass of their arms. Participants could not directly observe their

arms or hands; instead, veridical visual feedback was provided using a 2D

virtual reality system to overlay images of the left- and right-hand starting and

target locations (1.25 cm radius) and hand cursors (a 0.5 cm radius). This

arrangement ensured that the visual cursor appeared to the participant in the

same plane as their hand. Data were sampled at 1000 Hz and stored for offline

analysis.

Force Fields. Right-hand movements, termed contextual movements, were

always performed in a null field. Left-hand movements, termed adaptation

movements, were performed in one of three environments: null field, curl force

field, or channel trials. In the null field trials, no external forces were applied on

the hands by the robotic manipulandum. In the curl force field trials, a velocity-

dependent curl force field (Gandolfo et al., 1996) was implemented as:

$$\begin{bmatrix} F_x \\ F_y \end{bmatrix} = b \begin{bmatrix} 0 & -1 \\ 1 & 0 \end{bmatrix} \begin{bmatrix} \dot{x} \\ \dot{y} \end{bmatrix} \quad (1)$$

The viscous field constant b was set to ±13 Nm^{-1}s, with the sign determining

the force field's direction, which was either clockwise (CW) or counter clockwise

(CCW). Each participant experienced both force field directions, where the

direction of the force field was consistently associated with specific contextual

movements of the participant's right-hand. The relationship between the

directions of the contextual movements and the curl field direction (CW/CCW)

was counterbalanced across participants. In channel trials, a mechanical

channel, from the movement start position to the center of the final target, was

applied perpendicular to the motion direction (Scheidt et al., 2000; Milner and

Franklin, 2005) with a stiffness of 6,000 Nm^{-1} and a damping of 30 Nm^{-1}s. More

specifically, the channel was initiated at the time of the go cue and was oriented

from the participant's hand position at the time of the go cue to the center of the

final target. Channel trials were only conducted for movements towards the 0°

target.

Experiments

We conducted three bimanual interference experiments, each involving a

different participant group, to investigate the temporal generalization of

contralateral arm movements. Each experiment trained participants at a

specific timing relation between the two hands and then assessed their ability

to generalize this training across a broad range of timing relations (Fig. 1B).

The main difference between experiments was the bimanual timing relationship

for the training movements, which is the temporal gap between contextual and

adaptation movements (Fig. 1B, colored bars). Almost all other aspects of the

three experiments were identical.

In all experiments, the right-hand (contextual movement) moved from a start

location to the center point, while the left-hand (adaptation movement) moved

from the center point to a target location. All trials involved 12 cm active

movements of both hands. The bimanual timing relation of the training

movements was different for each experiment (Fig. 1B).

Experiment 1 – Post Adaptation Context. Participants first executed the left-hand adaptation movement, followed by the right-hand contextual movement. The adaptation movement starts 750 ms before the contextual movement, and is defined as a bimanual timing of +750 ms.

Experiment 2 – Simultaneous Context. Participants performed both the contextual and adaptation hand movements simultaneously, corresponding to a bimanual movement timing of 0 ms.

Experiment 3 – Prior Context. Participants first executed the right-hand contextual movement, followed by the left-hand adaptation movement. The contextual movement starts 750 ms before the adaptation movement. This timing is defined as -750 ms.

Experimental Design

The left-hand adaptation movement was associated with either a clockwise (CW) or counter clockwise (CCW) curl force field during the exposure phase of the experiment. Adaptation movements were oriented at 0°, 90°, 180° and 270°, each paired with two contextual movements rotated at either 45° CW or 45° CCW, to provide the contextual cue (see Fig. 1C). This resulted in eight unique combinations of contextual and adaptation movements. For half of the participants, context 1 was associated with the CW curl force field and context 2 was associated with the CCW curl force field while this was reversed for the

remaining participants. Although the physical locations of the handles were

shifted away from the center of the workspace to avoid hand collisions, the

visual feedback for the task was centered in the workspace, aligning the

contextual movement endpoint with the adaptation movement start. Right-hand

contextual movements always occurred in null fields. During the exposure

phase, the left-hand adaptation movements were always performed in either a

curl force field or a mechanical channel.

All trials consisted of a two-part movement produced by both hands. Each trial

began when the two cursors, representing the two hands, were placed in their

respective starting circles. Participants could not see their arms and hands, but

the hand cursors were color coded: red for the left cursor and blue for the right

cursor. The starting circle for the right-hand was grey, while that for the left-

hand was blue. Participants were instructed to keep their hands in these circles

until an auditory cue signaled the start of the movements. The order of the hand

movements was displayed at the bottom of the screen, with red and blue bars

indicating the relative timing of each hand's movements on the time scale along

the x-axis (Fig. 1E), which prepares the participant to move the hands with a

specific order and relative timing. The exact timing of the movements for both

the right- and left-hands were then cued in a trial with auditory beeps.

Upon the presentation of each auditory cue, participants moved each of their

hands (order as indicated by the display at the bottom of the screen) to the

appropriate target circle and stopped. Once the red cursor (left-hand) was

successfully placed into the final target circle, the circle's color changed from

pink to red. Similarly, when the blue cursor (right-hand) reached the center

target circle, the circle's color changed from light to dark blue. Importantly,

visual feedback of the trajectory of the contextual movement was never shown

to the participants due to its strong contextual effect (Howard et al., 2012, 2020;

Howard and Franklin, 2016; Franklin et al., 2023). Instead, participants relied

on this color change to confirm successful cursor placement in the target circle.

In each experiment, the time intervals between the start of the movement for

the left- and right-hands were set at +750 ms, 0 ms, or -750 ms for the training

trials. The trial ended once both hands remained stationary within their

respective target locations. At the end of each trial, the robotic manipulandum

passively moved the participants' hands to the next starting locations. The next

trial began once the handles had been stationary within the start locations for

300 ms. Participants were required to take short breaks approximately every

200 trials but could also rest at any time between trials by moving the cursors

out of the start targets.

At the end of each trial, participants received feedback regarding the peak

movement velocity of each hand. This speed feedback appeared immediately

after every trial as two vertical bars displayed at the top left of the screen, each

representing a hand. The desired speed range was indicated by two horizontal

lines. If a hand's peak speed was within this range, its bar turned green.

Conversely, if the movement was too fast or too slow, the bar turned red.

Participants were encouraged to maintain their peak movement speed within

the range of 38.25 to 51.75 cm/s.

If a participant's hand moved from the start position before the auditory cue, or

if they failed to complete a trial by moving their hands to the targets within the

maximum time limit (5 seconds), the trial was cancelled. Trials were also

cancelled if the time difference between initiating the movements of both arms

exceeded 200 ms for simultaneous context experiments and 950 ms for the

prior and post context experiments. Cancelled trials were repeated at the end

of each block. A trial was considered successful if both hands moved with a

peak speed within the specified range. For each successful trial, one point was

added to the participant's score, displayed at the top right of the screen.

Probe Channel Trials.

To estimate both the predictive compensation for the force field as part of the

adaptation process, and to measure the temporal generalization across a range

of timing relations between the two limbs, we introduced probe channel trials.

Throughout the experiment, on randomly selected trials, channel trials

matching the training condition were introduced to estimate the feedforward

adaptation to the curl force field. These channel trials only occurred on

adaptation movements (left-hand) moving forward to the final target, and were

associated with two possible angular contextual movements (as shown in Fig.

1D).

To examine the generalization of learning related to contextual movements made at specific temporal training relationships, we assessed the recall of adaptation across a range of timing relationships. Specifically, channel trials were conducted at 21 different bimanual timing relationships: [-1.8, -1.65, -1.5, -1.35, -1.2, -1.05, -0.9, -0.75, -0.6, -0.45, 0, 0.45, 0.6, 0.75, 0.9, 1.05, 1.2, 1.35, 1.5, 1.65, 1.8] seconds. The bimanual timing relationships used in the three experiments, as well as those used to probe generalization, are illustrated in Fig. 1B (black dots). All probe channel trials were limited to the forward reaching movements (target 0°) only.

Protocol

Participants performed two experimental sessions on two consecutive days. On day 1, participants completed 1608 trials, and on day 2, they completed 1594 trials, totaling 3202 trials (Fig. 1F). Temporal generalization probe trials were performed on day 1 in both pre-exposure and generalization phases to familiarize participants with executing movements of the two arms at different timing relations.

Day 1.

Pre-exposure phase: 216 trials. Participants began with 36 trials in the null field, divided into two blocks of 18 trials each. Each block consisted of sixteen null field trials (two for every movement type) and two channel trials (one for each context). This was followed by two blocks of mixed null field and generalization trials, totaling 180 trials. Each of these blocks comprised 48 null field trials and

42 generalization channel probe trials (1 for each context at each of the 21 different timing relationships).

Training phase: 180 trials. Participants performed ten blocks of trials in the curl force fields, where each curl force field was associated with one of the two contextual cues. Each block, consisting of 18 trials, included 16 curl field trials (two for every movement) and 2 channel trials (one for each context).

Generalization phase: 1212 trials. Participants performed six blocks in the curl force fields. Each block was composed of 160 curl field trials (20 for each of the 8 possible movements) and 42 generalization channel probe trials (one for each of the 21 different timing relations for both contexts).

Day 2.

Training phase: 180 trials. Participants performed ten blocks where each block comprised 16 curl field trials and 2 channel trials.

Generalization phase: 1414 trials. Participants performed seven blocks. Each block consisted of 160 curl field trials and 42 generalization channel probe trials (1 for each of the 21 different timing relations for both contexts). Data from this phase was used to measure the temporal generalization.

Data Analysis

The analysis was conducted offline using MATLAB R2023a. To examine learning and generalization, kinematic error and force compensation were calculated for each trial.

Kinematic Error. For each trial performed in null and viscous curl force fields,

the kinematic error was determined in the adaptation phase of the movement

by calculating the signed Maximum Perpendicular Error (MPE). The MPE is

defined as the maximum deviation of the hand's path from a straight line

between the starting location and the target. For each participant, the curl-field

direction dependent average MPE was computed over a set of 8 trials. In

addition, an average MPE was computed over both curl-field directions over a

set of sixteen trials for each participant. To combine results from Clockwise

(CW) and Counterclockwise (CCW) field trials, the sign of the MPE was

appropriately adjusted. Subsequently, the mean and standard error (SE) of the

MPE were calculated across all participants.

Force Compensation. During each channel trial, the force exerted by the

participants perpendicular to the wall of the simulated channel was measured

to estimate predictive feedforward adaptation. This measure was used to

assess both learning and generalization to different temporal relations between

the two hands. Force compensation was calculated by regressing the

measured force on the channel wall against the force required to perfectly

compensate for the force field (calculated as the movement velocity multiplied

by the force field strength b) (Smith et al., 2006). No offset term was used in

the fitting procedure. This method offers greater accuracy than relying solely on

a reduction in kinematic error, as it avoids the effect of limb impedance due to

muscle co-contraction (Franklin et al., 2003; Milner and Franklin, 2005; Franklin

and Franklin, 2021, 2023). Additionally, it provides an estimate of the

adaptation throughout the entire movement, rather than at just a single time

point. To examine adaptation, the sign of the force compensation values for the

two different contexts were maintained, so that adaptation to both field

directions could be observed independently. In additional analysis, the sign-

corrected compensation values were averaged over the two adjacent channel

trials within each block of 18 trials, representing the two different contexts. In

both cases, the mean and SE of these force compensation values were

calculated and analyzed across all participants throughout the experiment.

Bimanual Timing Relation. Channel probe trials examined the generalization of

force compensation across 21 different timing relations between the limbs,

spanning a ±1.8 second range. Although participants were cued to initiate

movements with these specific timing relations, they often moved with slightly

different timing relations. To estimate the actual bimanual timing relations

executed by participants, we calculated the start of each movement based on

specific position and speed criteria. A hand was considered to have initiated

movement when it deviated more than 2 mm from the starting target radius

(1.27 cm from the center of the start) and was traveling at a speed greater than

5 cm/s. A movement was considered to have terminated when it reached within

2 mm from the target radius (1.27 cm from the center). The bimanual temporal

relation for a given trial was calculated by subtracting the start time of the left

(adaptation) hand from the start time of the right (contextual) hand. Negative

values indicate that the contextual movement was made before the adaptation

movement.

Temporal Generalization. To quantify the temporal generalization of the motor

memory, we used force compensation data from the channel trials during the

Generalization Phase on the second day (refer to Fig. 1E) as participants had

achieved a higher level of adaptation. Temporal generalization was quantified

based on the actual timing relations between the two hands. For each

participant, compensation estimates from the channel trials were first allocated

to 29 temporal bins based on their bimanual timings. The center of each bin

ranged from -2.1 to +2.1 seconds, spaced by 150 ms intervals, matching the

cue timings (and widths) used in the generalization probe channel trials.

For each bin, we calculated the mean and the number of bin counts for each

participant. Bins with no data for a participant were removed to avoid misleading

estimates. To estimate generalization as a function of temporal relationships,

we calculated the mean and standard error of the mean (SEM) for the bin

values across all eight participants.

Statistics

Statistical analyses were performed in JASP 0.18.1 (JASP Team, 2023). A

Mauchly test was applied to check for sphericity where applicable. In the case

that the Mauchly test was significant, the degrees of freedom were adjusted

using a Greenhouse-Geisser correction. The α-level was set to .05 in all tests,

and omega squared (ω^2) was reported as the effect size. Omega squared is

an estimate of how much variance in the measurement can be accounted for

the explanatory variables.

For each experiment, we performed repeated measures ANOVAs to examine

differences in the MPE and force compensation within each experiment. We

compared the final pre-exposure MPE (mean of last block in pre-exposure),

initial exposure MPE (mean of first block in exposure) and final exposure MPE

(mean of final block on day2) using a repeated measures ANOVA with

repeated factor of phase (3 levels). If a significant main effect was found,

post-hoc comparisons were performed (Holm-Bonferroni corrected). For the

force compensation, we compared the final pre-exposure (mean of last block

in pre-exposure) and final exposure (mean of final block on day2) using a

repeated measures ANOVA with a repeated factor of phase (2 levels: pre-

exposure and final exposure).

To compare across experiments, we performed ANOVAs on the MPE and force

compensation in the final exposure block with a factor of experiment condition

(3 levels). If a significant main effect was found, post-hoc comparisons were

performed (Holm-Bonferroni corrected).

To compare the temporal generalization within each experiment and across the

three experiments we performed a repeated measures ANOVA with a repeated

factor of timing (21 levels) and between subjects factor of experimental

condition (3 levels). If a significant main effect was found, post-hoc

comparisons were performed (Holm-Bonferroni corrected for the whole family

of comparisons). Only post-hoc comparisons which were relevant for the study

were examined and reported. Specifically, we only examined differences

across timings within one experiment, or differences across experiments for

each timing. To support these findings without binning the data or averaging

trials which reduces the variability of the data, we also performed statistical

analysis on the full data set (all trials from all participants) using a Generalized

Additive Mixed Model (GAMM) in R (4.4.1). We fit the model (force

compensation as a function of time; random effect of participants) both with

conditions (3 experiments; full model) and without conditions (reduced model)

as a factor. We then compare the two models (full and reduced models) with a

likelihood ratio test to determine which model has a significantly better fit. To

perform pairwise comparisons between experimental conditions, we use a

similar procedure with each pairwise comparison of experiments. For each

comparison, we fit both the full model with two conditions and the model with

no conditions. Again, a likelihood ratio test was then used to test which model

best explains the data. If the model with conditions better explains the data, this

indicates that there is a significant effect of condition.

Results

Three groups of participants performed separate experiments, with each group

experiencing a different timing relationship in bimanual training while adapting

to two opposing force fields (Fig. 1B, C). Each group tested whether prior,

simultaneous, or post contralateral arm movements can provide a contextual

cue for the ipsilateral arm. Here the ipsilateral arm was exposed to opposing

viscous curl fields where the direction of the field was determined by the

contralateral arm motion. After this adaptation phase, the generalization of the

learned motor memory was assessed using probe channel trials (Fig. 1D) across various timing intervals between the two arms. These intervals included timings before (prior context), during (simultaneous context), and after (post context) the adaptation movement (Fig. 1B).

The generalization was examined across the three conditions and the different timings using a repeated measures ANOVA finding a significant main effect of timing (Greenhouse-Geisser corrected: $F_{3.204, 67.279}=5.863$; $p=0.001$; $\omega^2=0.131$), between subjects' effect of experiment condition ($F_{,2,21}=12.493$; $p<0.001$; $\omega^2=0.258$), and interaction between timing and experimental condition (Greenhouse-Geisser corrected: $F_{6.407, 67.279}=23.296$; $p<0.001$; $\omega^2=0.544$). Post-hoc comparisons (Holm-Bonferroni corrected) of the interaction effects were performed and reported in subsequent sections.

Experiment 1: Post Adaptation Context

Experiment 1 (post adaptation context) tested the training condition of +750 ms, in which the right-hand contextual movements were performed 750 ms after a preceding left-hand adaptation movement. That is, the movement of the right-hand acts as a planned follow-through movement that takes place after the adaptation movement of the left-hand. Before the introduction of the force field, participants made straight movements to each of the four targets with their left-hand (Fig. 2A). Once the force field was introduced, participants' movements were strongly curved away from the straight line, with the disturbance direction depending on the sign of the force field (Fig. 2B). However, by the end of the

exposure phase the movements were disturbed less laterally, and participants

made straighter movements to the targets for both force field directions

suggesting adaptation to the dynamics (Fig. 2C).

Figure 2 about here

To quantify adaptation, we examined two measures: maximum perpendicular

error (MPE) and force compensation. A repeated measures ANOVA indicated

that MPE showed significant variations between the pre-exposure, initial

exposure and final exposure phases ($F_{2,14}$=129.514; p<0.001; ω^2=0.864),

which were further tested with post-hoc comparisons. MPE (Fig. 2D) was close

to zero in the pre-exposure phase, but increased immediately upon the

introduction of the force field (p<0.001). MPE then reduced over the course of

the experiment, with final levels smaller than during initial exposure (p<0.001)

but not as low as in pre-exposure (p<0.001). Force compensation, as measured

in channel trials for the training timing relation, was close to zero in the pre-

exposure phase. However, force compensation increased slowly during the

exposure phase, reaching around 40% by the end of the second day (Fig. 2E).

A repeated measures ANOVA indicated a significant increase in force

compensation between final pre-exposure and final exposure levels

($F_{1,7}$=188.732; p<0.001; ω^2=0.887).

The temporal generalization of the learned motor memory was examined

towards the end of the exposure phase by introducing probe channel trials. In

these trials, the timing between the right contextual hand movement and the

left-hand adaptation movement was changed from +750 ms to one of 21 values

between -1.8 and +1.8 seconds (Fig. 1B). The force compensation observed in

these trials enabled us to assess the extent to which the learned motor memory

was specific to the trained condition, or whether it generalized to other timing

relations. These timing relations included both those matching the trained

relation between the hands (prior adaptation contextual movements) and

extended to those that did not (synchronous or post adaptation contextual

movements).

The temporal generalization of the motor memory after training at +750 ms

demonstrates consistent and effective generalization to other post adaptation

contextual movements (Fig. 2G), with all timing relations between +1.8 and

+0.45 s exhibiting around 40% compensation. However, there was little transfer

to adaptation from prior contextual movements (approximately 10%). Post-hoc

comparisons supported these findings with no significant differences across

any post movement timings (all p=1.0) but significant differences between all

post movements and all prior movements (all p<0.001), and no significant

differences between all post movement timings (all p=1.0). While the

simultaneous condition was different than the nearest prior condition (p=0.041),

there were no significant differences with any of the other post (all p>0.078) or

prior conditions (all p>0.188). This generalization pattern was consistent across

all but one participant, who showed flat generalization across all training

relations (Fig. 2F). We believe this flat generalization may indicate a more

explicit adaptation occurred in this participant, although we have no evidence

to test this possibility directly. Overall, we find that the post contextual bimanual

movement generalizes strongly to all other post contextual movement timings

and partially to the synchronous condition. However, there is little or no

generalization to the prior adaptation movement contextual conditions.

Experiment 2: Simultaneous Context

In the second experiment (simultaneous context), the onset of both the

adaptation and contextual movements occurred simultaneously (0 ms). In the

pre-exposure phase, participants made straight movements towards each of

the four targets with their left-hand (Fig. 3A). Again, once the force field was

introduced, participants' movements were strongly disturbed in the direction of

the applied force field (Fig. 3B). However, by the end of the exposure phase,

the participants made straight movements to the targets for both force field

directions (Fig. 3C). The kinematic error (MPE) increased immediately upon

introduction of the force field (Fig. 3D), but reduced over the course of force

field exposure. A repeated measures ANOVA found significant differences

across the phases of the experiments ($F_{2,14}$=105.044; p<0.001; ω^2=0.886), with

initial exposure larger than pre-exposure (p<0.001) and final exposure

(p<0.001), but no difference between pre-exposure and final exposure

(p=0.097). Force compensation steadily increased throughout the exposure

phase, reaching approximately 50% by the end of the second day (Fig. 3E). A

repeated measures ANOVA found a significant increase in force compensation

between final pre-exposure and final exposure levels ($F_{1,7}$=117.522; p<0.001;

ω^2=0.867). Thus, participants were able to adapt to the two opposing force

fields with simultaneous contextual movements of the other hand.

Figure 3 about here

The temporal generalization of the learned motor memory was again examined
at the end of the second day of the experiment, using probe trials at different
contextual timing relations ranging between -1.8 and +1.8 s. The temporal
generalization of the motor memory after training at 0 ms had a very strong
decline in compensation when the onset of the contextual movement was either
advanced or delayed (Fig. 3G). Post-hoc comparisons supported the significant
decreased away from the trained condition. The simultaneous timing was
significantly different than all post (all $p<0.001$) and all prior (all $p<0.001$)
timings, but there were no significant differences between any of the post and
prior timings (all $p=1.0$). This generalization pattern was consistently expressed
by all participants (Fig. 3F). Overall, we find that while strong learning occurs in
the 0 ms condition (50%), there is limited transfer to post contextual movements
(<10%), and no transfer to prior contextual movements (0%).

Experiment 3: Prior Adaptation Context

In the third experiment (prior context), right-hand contextual movements were
conducted prior to left-hand adaptation movements (-750 ms). During the pre-
exposure phase, participants made straight reaching movements (Fig. 4A),
which were strongly disturbed in the initial exposure of the force field (Fig. 4B).
However, by the end of the exposure phase the participants made fairly straight
movements to the targets (Fig. 4C) indicating that some compensation to the

applied dynamics was achieved. Again, the kinematic error (MPE) increased

immediately upon introduction of the force field (Fig. 4D), but reduced over the

course of force field exposure. A repeated measures ANOVA found significant

differences across the phases of the experiments ($F_{2,14}$=228.999; p<0.001;

ω^2=0.944), with initial exposure larger than pre-exposure (p<0.001) and final

exposure (p<0.001), although final exposure was still higher than pre-exposure

(p<0.001). Force compensation increased throughout the exposure phase,

reaching approximately 40% by the end of the second day (Fig. 4E). A repeated

measures ANOVA found a significant increase in force compensation between

final pre-exposure and final exposure levels ($F_{1,7}$=67.810; p<0.001; ω^2=0.818).

Figure 4 about here

The temporal generalization of the learned motor memory was once again

examined at the end of the second day of the experiment, using probe trials at

different contextual timing relations ranging between -1.8 and +1.8 s. The

temporal generalization of the motor memory after training at -750 ms shows a

marked decrease in compensation when the onset of the contextual movement

deviated from the trained timing relation (Fig. 4G). The trained timing (-0.75 s)

force compensation generalizes well to nearby timings (-0.45 s, -0.6 s, -0.9 s),

with no significant differences between any of the three timings (all p>0.685).

However, the force decreases away strongly from the three times with the

strongest responses (-0.6 s, -0.75 s, and -0.9 s) to the 0 ms timing (all p<0.001),

all post timings (all p<0.001), and all prior timings less than or equal to -1.6 s

(all p<0.01). The decay to longer prior adaptation contextual movement times

already decreased from the peak compensation by the 1.05 s timing (p=0.022). However, no significant differences were found between any of the post, simultaneous, or prior timings less than -1.35 s (all p>0.491). This generalization pattern was consistent across all participants (Fig. 4F). Overall, we observed a significant decrease in compensation when the onset of the contextual movement was either advanced or delayed relative to the training time, even for other prior adaptation contextual timing relations. There was no transfer to the synchronous timing relation or to prior movements.

Comparison Across Experiments

The three experiments followed identical designs, except for the timing relation between the two arms in the training condition, allowing for comparison across experiments. At the end of the force field exposure (final block of day 2), the simultaneous 0ms condition (Experiment 2, blue) yielded the straightest left-hand trajectories (Fig. 3B), whereas both the post (Experiment 1, red) and prior (Experiment 3, green) contexts remained slightly more curved (Figs. 2B, 4B). These trends align with the final levels of MPE values observed at the end of the experiments (Fig. 5A, B). The simultaneous context (blue) exhibited the most rapid decrease in MPE, whereas the prior adaptation context (green) and post contexts (red) had slower reductions in MPE. However, all three experiments showed clear reductions in MPE over the course of the experiment. An ANOVA comparing the final levels of MPE in the last block of the experiment found significant differences between the experiments ($F_{2,21}$=7.503; p=0.003; ω^2=0.028), with the simultaneous context showing a

lower final MPE than either the post (p=0.004) or prior (p=0.023) contexts, but

no significant differences between the post and prior contexts (p=0.341).

Force compensation exhibited similar patterns across all three experiments

(Fig. 5C). The simultaneous context (blue) showed the fastest increase in

adaptation, with little differences in the development of the force compensation

between the post (red) and prior (green) contexts. However, all three

experiments achieved comparable final compensation levels at the end of the

experiment (Fig. 5D), with no significant differences across the three conditions

($F_{2,21}$=1.350; p=0.281; ω^2=0.351). Overall, the results indicate that the timing

relationships in all three contextual timing relationships act as effective cues for

learning and adapting to opposing dynamics in the contralateral arm, facilitating

the adaptation to opposing force fields.

Figure 5 about here

A comparison of the temporal generalization across the three experiments

reveals the distinct effects resulting from training at the three distinct timing

relations (Fig. 5E). The synchronous context (blue) has the highest peak of

adaptation at the training condition with negligible generalization to either post

or prior timing relations. In contrast, both the post (red) and prior (green) timing

training contexts demonstrate generalization across various timing relations.

However, the prior context (green) shows selective generalization within

different prior timings (decreasing away from the training condition), and no

transfer to the synchronous or post timings. Conversely, the post context (red)

exhibits strong generalization across all post timing relations, partial transfer to the synchronous condition, and limited transfer (10%) to prior conditions. Post-hoc comparisons showed significant differences between the three experimental conditions, with the post context (red) exhibiting significant differences (all $p<0.001$) to both the simultaneous and prior contexts at every post timing (+1.8 s to +0.45 s). All three contexts were significantly different from one another at the simultaneous 0.0 s timing (all $p<0.03$). Similarly, the prior context was different from both other contexts at the trained 0.7 s timing (all $p<0.001$). The prior context (green) was also significantly different from the simultaneous context (blue) over all tested timings between 0 s and -0.9 s (all $p<0.043$). However, there were no significant differences between any of the three contexts in the timings from -1.06 to -1.8 s (all $p>0.123$). Overall, we find distinct temporal generalization across the three experiments.

To perform a similar comparison but without binning the data, we additionally used generalized Additive Mixed Models (GAMM). The GAMM analyses demonstrated that the effect of time on the force compensation significantly varies across conditions, as indicated by a likelihood ratio test comparing the full model (which allowed trends to differ by condition) to the reduced model (which assumed a common trend) ($\chi^2(4)=1009.33$, $p<0.0001$). Subsequent pairwise comparisons confirmed that the temporal force trends differ significantly between each pair of conditions (all $p<0.0001$), highlighting the distinct temporal dynamics associated with each condition.

We propose that the variations in the temporal generalization patterns across

the three experiments are indicative of different tuning functions within the

sensorimotor control system for contextual movements. These variations may

play a key role in the learning of coordinated actions across limbs. However,

an alternative explanation could be different levels of variability in the training

times, which might directly contribute to these patterns. To explore this

possibility, we quantified the variability in training timing for each experiment

(Fig. 5F). Here each histogram represents the distribution of the relative start

times between the arms during the training movements for each experiment.

The synchronous context (blue) had the narrowest spread of timings in the

training condition, with times very close to the requested 0 ms separation. In

contrast, the training conditions for the post (red) and prior movement

adaptation (green) contexts exhibited slightly greater variability in onset times.

However, this variability was significantly smaller than the width of the

generalization tuning functions observed, particularly in the case of the post

and prior contexts. This implies that the observed generalization is more likely

to reflect underlying neural circuitry rather than being merely a result of

variations in the training conditions.

To further quantify movement timing relationships, for Experiments 1 and 3 we

computed time differences histograms between the timing between the start of

the contextual movements after the end of the adaptation movements, and

between the end of the contextual movements and start of the adaptation

movements respectively (Figs. 5G, H). These values reflect the dwell time

between the subsequent movements, with mean values of these distributions

of 427 ms and 471 ms respectively.

Discussion

We examined the relationship between right arm contextual movements and

motor learning and recall in the left arm using a force field interference task. In

three experiments, participants simultaneously adapted to two opposing force

fields, where the force field directions were uniquely associated with the

movement angle between arms. Each experiment trained participants at a

distinct timing relation between the two arm movements (prior, simultaneous,

or post-context) and assessed temporal generalization. Motor memory

generalization across the experiments yielded three key findings. First, training

in the simultaneous context led to recall primarily only at the learned timing,

with limited transfer. Second, post-adaptation movement training resulted in

broad generalization to other post timings and to a lesser extent the

simultaneous timing. Lastly, prior-adaptation context training led to

generalization to similar prior timings, and no significant transfer to

simultaneous or post contexts. Overall, we find distinct patterns of temporal

generalization based on the training, emphasizing the unique characteristics

and implications of prior, simultaneous, and post-adaptation contexts in

bimanual dual adaptation.

The results confirm that bimanual motion is a strong contextual cue for learning

opposing dynamics. The relative motion of one limb predictively influences

forces in the other limb, regardless of timing. This aligns with research showing

that simultaneous movements reduce interference during dual adaptation

(Nozaki et al., 2006; Howard et al., 2010; Yokoi et al., 2011). For example,

switching between motor memories is facilitated when one hand is moving or

stationary (Nozaki et al., 2006), when the limbs move in the same or opposite

directions (Howard et al., 2010), or when the relative angle between hands

varies (Yokoi et al., 2011).

In unimanual movements, prior (Howard et al., 2012, 2020; Howard and

Franklin, 2015) and post adaptation movements (Howard et al., 2015; Sheahan

et al., 2016), facilitate learning separate motor memories. Prior motion also

influences learning in bimanual movements (Gippert et al., 2023). Our study

demonstrates that post adaptation contextual movements also drive learning of

independent motor memories in the contralateral limb, similar to unimanual

studies (Howard et al., 2012, 2015; Sheahan et al., 2016). While we found

comparable adaptation levels in both prior and post adaptation contexts,

unimanual studies showed almost double compensation for prior (lead-in)

compared to post-adaptation (follow-through) movements.

In unimanual dual adaptation, prior contextual movements are most effective

when they directly precede adaptation movements, with the contextual effect

diminishing as the dwell time (the time between the end of the contextual lead-

in movement and the start of the adaptation movement) increases (Howard et

al., 2012). In Experiment 3 the dwell time of 471 ms produced about 40% force compensation. Although this dwell time produced similar levels of adaptation in lead-in unimanual movements (Howard et al., 2012), higher adaptation occurred for shorter dwell times. In Experiment 1 the dwell time of 421 ms produced 40% adaptation, similar to that for follow-through unimanual movements (40%) with dwell times close to zero (Howard et al., 2015).

One key difference in our experimental design to previous bimanual studies (Nozaki et al., 2006; Howard et al., 2010; Yokoi et al., 2011; Gippert et al., 2023), is that our participants never received online visual feedback about the movement of their contextual (right) arm, whereas such feedback was provided in the previous research. This control is crucial because visual cursor motion alone is a strong contextual cue (Howard et al., 2012, 2020; Howard and Franklin, 2016; Franklin et al., 2023). Therefore, the inclusion of visual motion in previous bimanual studies might reflect the impact of this visual element rather than the physical motion of the contralateral limb. Although Gippert and colleagues (Gippert et al., 2023) included a visual 'bimanual' condition showing little adaptation, the visual cursor was offset by 10 cm from the adaptation movement. This offset might have weakened the contextual strength or acted as a weaker peripheral visual cue (Howard et al., 2013). Our study demonstrates that prior, simultaneous, and post adaptation movements of the contralateral limb can facilitate the separate learning of motor memories, even without visual information. This finding supports the interpretations of previous studies (Nozaki et al., 2006; Howard et al., 2010; Yokoi et al., 2011), and

highlights the robustness of limb movement as a contextual cue in motor

learning.

Numerous studies have demonstrated generalization of a motor memory

across movement directions (Thoroughman and Shadmehr, 2000; Donchin et

al., 2003; Gonzalez Castro et al., 2011), kinematics (Joiner et al., 2011; Howard

et al., 2020; Orschiedt and Franklin, 2023) and postures (Shadmehr and

Mussa-Ivaldi, 1994; Berniker et al., 2014; Leib and Franklin, 2021). Our study

is the first to show temporal generalization. Just as generalization across

directions has been used to infer the neural basis functions underlying motor

memories (Poggio and Bizzi, 2004; Shadmehr, 2004; Joiner et al., 2011;

Kadiallah et al., 2012), we propose that the pattern of the temporal

generalization reflects the temporal tuning of these neural basis functions.

The three experiments exhibit different levels of variability in the timing between

limbs during training (Fig. 5F). While the prior and post adaptation movement

contexts show similar timing variability, the simultaneous context has a

narrower range, indicating greater consistency. This is unsurprising as

simultaneous bimanual movements are generally easier to perform (Kelso,

1995; Howard et al., 2009a; Kobayashi and Nozaki, 2023). Kinematic variability

of contextual movements during adaptation reduces the learning rate (Howard

et al., 2015, 2017). Therefore, temporal variability of contextual movements

likely had a similar effect, as we observed slower final adaptation levels for the

prior and post contexts. Our study is unable to separate the effect of temporal

variability from the training condition. The prior and post conditions exhibit

broader temporal generalization than the simultaneous condition, which shows

almost no generalization to other timings. Despite similar training time

variability, strong differences in generalization between prior and post contexts

exist. The post adaptation context generalizes to simultaneous and prior

adaptation times, suggesting generalization is not confined to training

variability. Both prior and post contexts show generalization beyond training

variability. We propose that the observed differences stem from variations in

the inherent use of contextual information for generating natural behavior, and

therefore in the underlying neural circuitry involved in learning and executing

coordinated bimanual movements.

Neural activity associated with generating and controlling bimanual movement

(Swinnen, 2002) involves interactions between both cerebral hemispheres

(Gerloff and Andres, 2002). This includes both contralateral and ipsilateral

primary motor cortex, where bimanual and unimanual movements exhibit

different neural activity patterns (Donchin et al., 1998; Steinberg et al., 2002;

Cross et al., 2020). The cerebellum (Wolpert et al., 1998; Imamizu et al., 2000;

Ito, 2005), parietal, and frontal areas also play significant roles (Scott, 2012).

While we cannot elaborate on the neural regions responsible, our work

examines the temporal tuning of the neural basis functions that lead to motor

memories through generalization. Local neural basis functions are suggested

to underlie motor memory formation (Poggio and Bizzi, 2004; Shadmehr, 2004;

Joiner et al., 2011; Kadiallah et al., 2012), allowing generalization to similar

states. These basis functions often exhibit Gaussian-like tuning. However, our

findings reveal that temporal generalization can exhibit different forms

depending on the training conditions, ranging from broad temporal

generalization in the prior context to narrow tuning in the simultaneous context.

A normative perspective can provide insights into the human nervous system's

operation (Franklin et al., 2016; Makino et al., 2016; Česonis and Franklin,

2022; Leib et al., 2024). Prior movements are crucial for anticipatory muscle

activation patterns in real-world tasks (Klapp, 1979), especially in bimanual

tasks, where coordination and anticipatory adjustments between limbs ensure

synchronized actions (Swinnen and Wenderoth, 2004). As actions at one time

can require compensatory actions at different times, this might explain why both

prior and past movements show strong temporal generalization. Adjustments

to forces produced by the other limb are relevant in simultaneous movements,

such as opening a jar or pouring a glass of water. However, these interactions

are so closely linked in time that are often taken as evidence for predictive

models (Desmurget and Grafton, 2000; Wolpert and Flanagan, 2001; Flanagan

et al., 2003; Franklin and Wolpert, 2011). This might explain why trained

simultaneous interactions show limited generalization to prior or post-

adaptation contexts.

Considering an evolutionary perspective is useful (Cisek, 2022; Cisek and

Hayden, 2022). In locomotion on variable ground, movement of one limb

requires compensatory motions in the other limb, both for prior movements and

in preparation for future steps. The temporal interval between steps can vary

widely, which might explain the plastic adaptation and generalization to prior

and post-adaptation movements, while narrow generalization of simultaneous

movements reflects tight coupling in object manipulation.

This work highlights the sensorimotor control system's plasticity, showing

different training conditions produce widely different temporal generalization.

These differences might be explained by considering the influence of different

tasks on the neural circuitry over evolutionary timescales. While state-space

(Lee and Schweighofer, 2009; Forano and Franklin, 2020), and contextual

inference models (Heald et al., 2021, 2023) can explain dual-adaptation, these

models cannot yet predict different patterns of temporal generalization.

Author Contributions

The study was conceived and designed by ISH and DWF. The study was

implemented by ISH. The experiments were conducted by SF. Data analysis

was carried out by ISH. Figures were prepared by ISH and DWF. Statistics

were performed by DWF. The manuscript was written by ISH, SF and DWF.

The final manuscript was reviewed and edited by ISH, SF and DWF.

References

Berniker M, Franklin DW, Flanagan JR, Wolpert DM, Kording K (2014) Motor
learning of novel dynamics is not represented in a single global coordinate
system: evaluation of mixed coordinate representations and local learning. J
Neurophysiol 111:1165–1182.

Česonis J, Franklin DW (2022) Contextual cues are not unique for motor learning: Task-dependant switching of feedback controllers Haith AM, ed. PLoS Comput Biol 18:e1010192.

Chow J, Carlton L, Lim Y-T, Chae W-S, Shim J-H, Kuenster A, Kokubun K (2003) Comparing the pre- and post-impact ball and racquet kinematics of elite tennis players' first and second serves: a preliminary study. Journal of Sports Sciences 21:529–537.

Cisek P (2022) Evolution of behavioural control from chordates to primates. Phil Trans R Soc B 377:20200522.

Cisek P, Hayden BY (2022) Neuroscience needs evolution. Phil Trans R Soc B 377:20200518.

Conditt MA, Gandolfo F, Mussa-Ivaldi FA (1997) The motor system does not learn the dynamics of the arm by rote memorization of past experience. J Neurophysiol 78:554–560.

Cross KP, Heming EA, Cook DJ, Scott SH (2020) Maintained Representations of the Ipsilateral and Contralateral Limbs during Bimanual Control in Primary Motor Cortex. J Neurosci 40:6732–6747.

Desmurget M, Grafton S (2000) Forward modeling allows feedback control for fast reaching movements. Trends in cognitive sciences 4:423–431.

Donchin O, Francis JT, Shadmehr R (2003) Quantifying generalization from trial-by-trial behavior of adaptive systems that learn with basis functions: theory and experiments in human motor control. J Neurosci 23:9032–9045.

Donchin O, Gribova A, Steinberg O, Bergman H, Vaadia E (1998) Primary motor cortex is involved in bimanual coordination. Nature 395:274–278.

Flanagan JR, Vetter P, Johansson RS, Wolpert DM (2003) Prediction precedes control in motor learning. Curr Biol 13:146–150.

Forano M, Franklin DW (2020) Timescales of motor memory formation in dual-adaptation. PLoS Comput Biol 16:e1008373.

Forano M, Schween R, Taylor JA, Hegele M, Franklin DW (2021) Direct and indirect cues can enable dual adaptation, but through different learning processes. J Neurophysiol 126:1490–1506.

Franklin DW, Burdet E, Osu R, Kawato M, Milner TE (2003) Functional significance of stiffness in adaptation of multijoint arm movements to stable and unstable dynamics. Exp Brain Res 151:145–157.

Franklin DW, Reichenbach A, Franklin S, Diedrichsen J (2016) Temporal Evolution of Spatial Computations for Visuomotor Control. J Neurosci 36:2329–2341.

Franklin DW, Wolpert DM (2011) Computational mechanisms of sensorimotor control. Neuron 72:425–442.

Franklin S, Franklin DW (2021) Feedback Gains modulate with Motor Memory
Uncertainty. Neurons, Behavior, Data analysis, and Theory 5.

Franklin S, Franklin DW (2023) Visuomotor feedback tuning in the absence of visual
error information. Neurons, Behavior, Data analysis, and Theory.

Franklin S, Leib R, Dimitriou M, Franklin DW (2023) Congruent visual cues speed
dynamic motor adaptation. J Neurophysiol 130:319–331.

Gandolfo F, Mussa-Ivaldi FA, Bizzi E (1996) Motor learning by field approximation.
PNAS 93:3843–3846.

Gerloff C, Andres FG (2002) Bimanual coordination and interhemispheric interaction.
Acta Psychologica 110:161–186.

Gippert M, Leupold S, Heed T, Howard IS, Villringer A, Nikulin VV, Sehm B (2023)
Prior Movement of One Arm Facilitates Motor Adaptation in the Other. J
Neurosci 43:4341–4351.

Gonzalez Castro LN, Monsen CB, Smith MA (2011) The binding of learning to
action in motor adaptation. PLoS Comput Biol 7:e1002052.

Heald JB, Lengyel M, Wolpert DM (2021) Contextual inference underlies the
learning of sensorimotor repertoires. Nature 600:489–493.

Heald JB, Lengyel M, Wolpert DM (2023) Contextual inference in learning and
memory. Trends in Cognitive Sciences 27:43–64.

Hirashima M, Nozaki D (2012) Distinct Motor Plans Form and Retrieve Distinct
Motor Memories for Physically Identical Movements. Current Biology
22:432–436.

Howard IS, Ford C, Cangelosi A, Franklin DW (2017) Active lead-in variability
affects motor memory formation and slows motor learning. Sci Rep 7:7806.

Howard IS, Franklin DW (2015) Neural Tuning Functions Underlie Both
Generalization and Interference. PloS one 10:e0131268.

Howard IS, Franklin DW (2016) Adaptive tuning functions arise from visual
observation of past movement. Sci Rep 6:28416.

Howard IS, Franklin S, Franklin DW (2020) Asymmetry in kinematic generalization
between visual and passive lead-in movements are consistent with a forward
model in the sensorimotor system. PLoS ONE 15:e0228083.

Howard IS, Ingram JN, Franklin DW, Wolpert DM (2012) Gone in 0.6 seconds: the
encoding of motor memories depends on recent sensorimotor States. J
Neurosci 32:12756–12768.

Howard IS, Ingram JN, Körding KP, Wolpert DM (2009a) Statistics of natural
movements are reflected in motor errors. J Neurophysiol 102:1902–1910.

Howard IS, Ingram JN, Wolpert DM (2009b) A modular planar robotic
manipulandum with end-point torque control. J Neurosci Meth 181:199–211.

Howard IS, Ingram JN, Wolpert DM (2010) Context-dependent partitioning of motor
learning in bimanual movements. J Neurophysiol 104:2082–2091.

Howard IS, Wolpert DM, Franklin DW (2013) The effect of contextual cues on the
encoding of motor memories. J Neurophysiol 109:2632–2644.

Howard IS, Wolpert DM, Franklin DW (2015) The value of the follow-through
derives from motor learning depending on future actions. Curr Biol 25:397–
401.

Imamizu H, Miyauchi S, Tamada T, Sasaki Y, Takino R, Pütz B, Yoshioka T, Kawato
M (2000) Human cerebellar activity reflecting an acquired internal model of a
new tool. Nature 403:192–195.

Ito M (2005) Bases and implications of learning in the cerebellum — adaptive control
and internal model mechanism. Progress in Brain Research 148:95–109.

Joiner WM, Ajayi O, Sing GC, Smith MA (2011) Linear hypergeneralization of
learned dynamics across movement speeds reveals anisotropic, gain-encoding
primitives for motor adaptation. J Neurophysiol 105:45–59.

Kadiallah A, Franklin DW, Burdet E (2012) Generalization in adaptation to stable and
unstable dynamics. PloS one 7:e45075.

Karniel A, Mussa-Ivaldi FA (2002) Does the motor control system use multiple
models and context switching to cope with a variable environment? Exp Brain
Res 143:520–524.

Kawato M (1999) Internal models for motor control and trajectory planning. Current
opinion in neurobiology 9:718–727.

Kelso JAS (1995) Dynamic patterns: the self-organization of brain and behavior.
Cambridge, Mass.: MIT Press.

Klapp ST (1979) Doing two things at once: The role of temporal compatibility.
Memory & Cognition 7:375–381.

Kobayashi T, Nozaki D (2023) Motor control and adaptation in a redundant motor
task manipulating a stick with both hands. bioRxiv.

Krakauer JW, Shadmehr R (2006) Consolidation of motor memory. Trends in
neurosciences 29:58–64.

Lackner JR, Dizio P (1994) Rapid adaptation to Coriolis force perturbations of arm
trajectory. J Neurophysiol 72:299–313.

Lee J-Y, Schweighofer N (2009) Dual adaptation supports a parallel architecture of
motor memory. J Neurosci 29:10396–10404.

Leib R, Franklin DW (2021) Error Prediction Determines the Coordinate System
Used for Novel Dynamics Representation. bioRxiv.

Leib R, Howard IS, Millard M, Franklin DW (2024) Behavioral Motor Performance.
Comprehensive Physiology:5179–5224.

Makino H, Hwang EJ, Hedrick NG, Komiyama T (2016) Circuit Mechanisms of
Sensorimotor Learning. Neuron 92:705–721.

Milner TE, Franklin DW (2005) Impedance control and internal model use during the
initial stage of adaptation to novel dynamics in humans. J Physiol 567:651–
664.

Nozaki D, Kurtzer I, Scott SH (2006) Limited transfer of learning between unimanual
and bimanual skills within the same limb. Nature Neuroscience 9:1364–1366.

Oldfield RC (1971) The assessment and analysis of handedness: the Edinburgh
inventory. Neuropsychologia 9:97–113.

Orschiedt J, Franklin DW (2023) Learning context shapes bimanual control strategy
and generalization of novel dynamics. PLoS Comput Biol 19:e1011189.

Osu R, Hirai S, Yoshioka T, Kawato M (2004) Random presentation enables subjects
to adapt to two opposing forces on the hand. Nature Neuroscience 7:111–112.

Poggio T, Bizzi E (2004) Generalization in vision and motor control. Nature
431:768–774.

Sarwary AME, Stegeman DF, Selen LPJ, Medendorp WP (2015) Generalization and
transfer of contextual cues in motor learning. J Neurophysiol 114:1565–1576.

Scheidt RA, Reinkensmeyer DJ, Conditt MA, Rymer WZ, Mussa-Ivaldi FA (2000)
Persistence of motor adaptation during constrained, multi-joint, arm
movements. J Neurophysiol 84:853–862.

Scott SH (2012) The computational and neural basis of voluntary motor control and
planning. Trends in cognitive sciences 16:541–549.

Shadmehr R (2004) Generalization as a behavioral window to the neural mechanisms
of learning internal models. Human Movement Science 23:543–568.

Shadmehr R, Brashers-Krug T (1997) Functional stages in the formation of human
long-term motor memory. J Neurosci 17:409–419.

Shadmehr R, Mussa-Ivaldi FA (1994) Adaptive representation of dynamics during
learning of a motor task. J Neurosci 14:3208–3224.

Shadmehr R, Smith MA, Krakauer JW (2010) Error correction, sensory prediction,
and adaptation in motor control. Annual review of neuroscience 33:89–108.

Sheahan HR, Franklin DW, Wolpert DM (2016) Motor planning, not execution,
separates motor memories. Neuron 92:773–779.

Smith MA, Ghazizadeh A, Shadmehr R (2006) Interacting adaptive processes with
different timescales underlie short-term motor learning. PLoS Biology 4:e179.

Steinberg O, Donchin O, Gribova A, De Oliveira SC, Bergman H, Vaadia E (2002)
Neuronal populations in primary motor cortex encode bimanual arm
movements. Eur J of Neuroscience 15:1371–1380.

Swinnen SP (2002) Intermanual coordination: From behavioural principles to neural-
network interactions. Nat Rev Neurosci 3:348–359.

Swinnen SP, Wenderoth N (2004) Two hands, one brain: cognitive neuroscience of
bimanual skill. Trends in Cognitive Sciences 8:18–25.

Thoroughman KA, Shadmehr R (2000) Learning of action through adaptive
combination of motor primitives. Nature 407:742–747.

Wolpert DM, Flanagan JR (2001) Motor prediction. Curr Biol 11:R729-32.

Wolpert DM, Miall RC, Kawato M (1998) Internal models in the cerebellum. Trends
in Cognitive Sciences 2:338–347.

Yokoi A, Hirashima M, Nozaki D (2011) Gain Field Encoding of the Kinematics of
Both Arms in the Internal Model Enables Flexible Bimanual Action. J
Neurosci 31:17058–17068.

1088

1089

1090

1091

1092

1093

1094 **Figure Legends**

1095

Figure 1. Experimental Setup and Design. **A.** Experiments were conducted using a bimanual vBOT setup in a virtual reality environment. The participant sits while grasping both handles of the vBOT manipulandum and receiving visual feedback in the plane of movement. **B.** Bimanual Movement Timing Relationships. Three separate experiments were conducted with training at timing relationships of +750 ms (Post Adaptation Context; red), 0 ms (Simultaneous Context; blue), and -750 ms (Prior Context; green). The training time relation between the two hands is shown with the colored bar for each experiment. To test temporal generalization, channel trials were presented at 21 distinct timing relationships, represented by black circles, ranging from -1.8 to +1.8 seconds. Each temporal relationship was tested seven times for each of the two contexts, amounting to a total of 14 tests per relationship. **C.** Experimental protocol for training and testing. The top four rows depict the contextual relationships used in both null and training trials for the four possible adaptation movement directions. The movement of the right-hand provides context, with four distinct starting locations (at 45°, 135°, 225°, and 315°) leading to a central location on the right. The left-hand performs the adaptation movement, executed either in a null field or a viscous curl field. This hand also has four movement directions, beginning from the central position on the left and progressing to specific target locations (0°, 90°, 180°, and 270°). Notably, each adaptation movement direction is paired with two unique contextual movement directions (two columns indicating the two possible contexts). It can be seen that while the two hands were separated by 30 cm, the visual representations of the two hands' movements were aligned such that the end of the movement of the right-hand ended in the center target where the left-hand starts its movement. This involved a visual shift of 15 cm for each hand. **D.** The row (brown shaded region) displays the contextual relationships for channel trials, designed to estimate feedforward adaptation to the viscous curl field. These channel trials proceed solely along the y-axis to an upper target and are linked with two distinct angular contextual movements originating at 135° (right column) and 225° (left column). **E.** Visual feedback provided to the participants. At the bottom of the screen the temporal relation of the right-hand (blue bar) and left-hand (red bar) for the next trial was displayed. The red bar was always in the center of the screen but the blue bar was moved depending on the timing of the relation. Cursors representing the right (blue) and left (red) hands were shown, along with the start and end targets for both movements. The red cursor was moved to the red target while

the blue cursor was moved to the blue target. A score representing the number of successful

targets was presented at the top right of the screen. **F.** The experimental trial schedule was

identical across all three experiments, but each experiment used different training timing

relationships. Day 1 started with pre-exposure (Pre-exp) in the null field trials which included

testing probe channel trials. The white color indicates that all non-channel trials were performed

in a null field to familiarize participants with the manipulanda and the testing procedures. The

curl force fields (grey shaded region) were introduced in the Train session (180 trials), followed

by a Generalization phase (1212 trials) that provided practice to the bimanual movement

relations. Day 2 started with further training (Train), followed by a final Generalization session

to study generalization.

1136

Figure 2: Results for Experiment 1 where the right-hand contextual movement follows the left-hand training movement by 750 ms (+750 ms condition). **A.** Hand paths in the last block of pre-exposure in the null field condition. The mean and standard error (shaded region) of left-hand paths across participants from the central location to the target are shown for both contextual cue movement directions. Movements that will be associated with the CW curl force field are shown in orange whereas movements that will be associated with the CCW force field are

shown in purple. Within the selected block, the last 8 unique movement directions per participant were used for the calculation. **B.** The hand paths on initial exposure to the curl field. **C.** The hand paths in the final block of curl field exposure. **D.** The kinematic error, measured as maximum perpendicular error (MPE) for the left-hand, plotted over the course of the experiment. The mean (line) and standard error (shaded area) are calculated across the participants for each curl force field direction separately. The grey shaded region indicates the period where the force field is applied to the left-hand during the adaptation movements. **E.** Force compensation, expressed as a percentage of complete adaptation to the force field, over the course of the experiment for each curl force field direction separately. Positive values indicate adaptation to CW field whereas negative values indicate adaptation to the CCW field. Grey region indicates the application of the force field, and the dark grey region indicates the period over which the generalization to different timing relations between the two hands was measured. **F.** The temporal generalization of the learned motor memory to all timing relations between the two hands shown for individual participants. Each line denotes a different participant, and the red dashed vertical line indicates the trained relation. **G.** The mean and standard error of the temporal generalization of the learned motor memory across participants. Red circles denote the center of the window over which values were averaged. We see that training at +750 ms led to effective generalization to post adaptation contextual movement timing relationships, but not to prior adaptation contextual movements.

Figure 3: Results for Experiment 2 using training at a 0 ms synchronous context timing relationship **A.** Hand paths in the last block of pre-exposure in the null field condition are shown for both contextual cues. **B.** The hand paths on initial exposure to the curl field. **C.** The hand paths in the final block of curl field exposure. **D.** The kinematic error, measured as maximum perpendicular error (MPE) for the left-hand, plotted over the experiment for each curl force field direction separately. **E.** Force compensation for each curl force field direction separately,

expressed as a percentage of complete adaptation to the force field, over the experiment. Grey

region indicates the application of the force field, and the dark grey region indicates the period

over which the generalization to different timing relations between the two hands was

measured. **F.** The temporal generalization of the learned motor memory to all timing relations

between the two hands, shown for individual participants. Each line denotes a different

participant and the blue dashed vertical line indicates the trained relation. **G.** The mean and

standard error of the temporal generalization of the learned motor memory across participants.

Training at 0 ms leads to limited generalization to post or prior contextual movement timing

relationships.

1182

Figure 4: Results for Experiment 3 where the left-hand adaptation movement followed the right-hand contextual movement by 750 ms (-750 ms condition). **A.** Hand paths in the last block of pre-exposure in the null field condition. **B.** The hand paths on initial exposure to the curl field. **C.** The hand paths in the final block of curl field exposure. **D.** The kinematic error (MPE) for the left-hand for each curl force field direction separately. **E.** Force compensation for each curl force field direction separately, expressed as a percentage of complete adaptation to the force field.

F. The temporal generalization of the learned motor memory to all timing relations between the

two hands shown for individual participants. Each line denotes a different participant and the

green dashed vertical line shows the trained relation. **G.** The mean and standard error of the

temporal generalization of the learned motor memory across participants. Training at -750ms

leads to a swift decline in generalization to other prior adaptation contextual movement

temporal relationships, with no generalization to post contextual movements.

1197

Figure 5. Comparison across the three experiments where participants were trained with either

post (+750 ms; red), synchronous (0 ms; blue) or prior (-750 ms; green) timing relation. **A.**

Comparison of kinematic error (MPE) changes across the experiment. **B.** Final levels of MPE

at the end of the experiment. Colored bars indicate the mean MPE across participants in the

final exposure block, while black circles represent individual participant values. **C.** Force

compensation evolution across the three experiments. **D.** Final force compensation values

across the three experiments (mean and individual participants) are shown for the final

exposure block. **E.** Comparison of temporal generalization across bimanual timing relations

after training at one of the three timing relations. Colored vertical lines illustrate the trained

relation for each experiment. Solid lines depict the mean, and shared regions the standard error

of the mean, across participants. Dot placements on the plot indicate the central location and

the estimated mean compensation value for each bin. **F.** Histograms present relative movement

initiation offsets between the two arms for all null practice and curl field training trials across the

three experiments. Results for the +750 ms post movement context experiment are in red, the

synchronous 0 ms movement context experiment in blue, and the -750 ms prior movement

context experiment in green. **G.** Histograms of the dwell time in the prior adaptation movement

context. **H.** Histograms of the dwell time in the post adaptation movement context.