Alternative splicing plays a nonredundant role in greater amberjack

(*Seriola dumerili*) in the adaptation to ambient salinity fluctuations

Chunyu Zhao[1*], Yuqi Liu[1*], Panpan Zhang[1], Xinhui Xia[1], Yuchen Yang[1,2*]

1 State Key Laboratory of Biocontrol, School of Ecology, Sun Yat-sen University, Shenzhen

Guangdong, China

2 Guangdong Provincial Key Laboratory of Marine Resources and Coastal Engineering, Guangzhou,

Guangdong, China

* The authors contribute equally to this work.

Abstract

Alternative splicing (AS) is an important post-transcriptional mechanism for adaptation of fish to

environmental stress. Here, we performed a genome-wide investigation to explore the biological

importance of AS dynamics in greater amberjack (*Seriola dumerili*), an economical marine teleost

species, in response to hypo- (10 ppt) and hyper-salinity (40 ppt) stresses. The results revealed high

level of differential splicing in both gills and kidney upon the exposure to undesired salinity regimes.

In gills, genes involved in energy metabolism, stimulus response and epithelial cell differentiation

were differentially spliced in response to the deviation of normal water salinity, while sodium ion

transport, erythrocyte homeostasis and cellular amide metabolism were enhanced in kidney to

combat the adverse impacts of salinity changes. More importantly, the majority of the differentially

spliced genes were not differentially expressed, and AS was found to regulate different biological

processes from differential gene expression, indicative of the functionally nonredundant role of AS

in modulating salinity acclimation in greater amberjack. Together, our study highlights the important

contribution of post-transcriptional mechanisms to the adaptation of fish to ambient salinity

fluctuations, and provides a theoretical guidance to the conservation of marine fishery resources

under the increasingly extreme environmental challenges.

1 introduction

Occupying more than two-thirds of the Earth's surface, the oceans are the largest ecosystem on the

planet, hosting a huge number of organisms and playing crucial roles in maintaining the diversity

and balance of life on Earth. However, in the recent few decades, global climate changes, such as

global warming, glacier melting, and precipitation decreasing, have substantially changed the global

water cycle and altered worldwide salinity distribution among different sea areas (Durack and

Wijffels, 2010; Helm et al., 2011; Durack et al., 2012; Skliris et al., 2016; Ullah et al., 2021). One

noticeable trend is that the freshwater is getting fresher while the saltwater becomes saltier (Durack

et al., 2012). Till 2017, the mean salinity contrast between the low and high salinity regions had

increased by more than 5% (Cheng et al., 2020). Furthermore, this imbalance has been found to be

amplified by human activities (Omerspahic et al., 2022; Röthig et al., 2023). For instance, brine

discharge during seawater desalination can lead to an abnormal increase in salinity of the receiving

water body in coastal marine environment, while the enhanced urban drainage may freshen

nearshore seawater (Omerspahic et al., 2022; Röthig et al., 2023). Salinity fluctuations in seawater

would alter the dissolved oxygen level, nutrient concentration, and stratification of the oceans,

thereby posting an adverse impact on the respiration, metabolism, reproduction, and even disease

immunity of marine organisms (Lehtonen et al., 2016; Guo et al., 2020; Galkanda-Arachchige et al.,

2021; Lu et al., 2022; Lacy et al., 2023; Röthig et al., 2023). For example, the increased water

salinity was found to disrupt the normal energy allocation, and physiological and metabolic

processes of fish, such as grass carp (*Ctenopharyngodon idella*) and African catfish (*Clarias*

gariepinus), and finally inhibit their growth and survival (Smyth and Elliott, 2016; Zidan et al., 2022;

Liu et al., 2023). Moreover, salinity change-induced biodiversity loss of plankton, macroalgae, coral

reef, and seagrass may cause trophic collapse of marine ecosystem and lead to drops in fishery yield,

which threaten ecosystem functions and food supply to human (Röthig et al., 2023).

Seriola dumerili, also known as greater amberjack, is a large saltwater bony fish of the family

Carangidae. They are globally distributed in tropical and subtropical seas, including the Atlantic,

Indian, Pacific, and Mediterranean Oceans (Zupa et al., 2017; Araki et al., 2018). Greater amberjack

is an economically important fish that is widely popular for its tasty meat, and it is also a major

predator and inhabitant of artificial reefs such as coral reefs and shipwrecks, maintaining the

structure and function of these ecosystems (Pereira et al., 2023). It has been widely reported that the

growth, development, and physiology of greater amberjack were sensitive to the changes of water

salinity, temperature, and seawater pressure (Barany et al., 2021; Navarro-Guillen et al., 2022; Peng

et al., 2022; Liu et al., 2023; Ru et al., 2023). Comparative transcriptomic analyses showed that

salinity alternations substantially disrupt tissue growth and development, energy metabolism, and

immune responses in greater amberjack (Peng et al., 2022; Liu et al., 2023). Correspondingly,

muscle water content, gill and intestinal Na-K-ATPase activity, and growth-related processes were

enhanced to combat the adverse impacts (Barany et al., 2021; Liu et al., 2023). However, most of

the current researches only focused on the responsive regulations at transcriptional level. No

attentions have ever been paid to the role of post-transcriptional mechanisms in modulating the

resistance to seawater salinity fluctuations in greater amberjack.

Alternative splicing (AS) is one type of post-transcriptional regulatory mechanism in

eukaryotes, and it can produce multiple isoforms with different combinations of exons, thus giving

rise to multiple protein products from one single gene (Tian et al., 2022a). Therefore, AS can extend

the diversity of structures and functions of proteins without increasing gene numbers. In fish, AS

has been demonstrated to play critical roles in regulating the growth, development, and

environmental adaptation of fish (Wan and Su, 2015; Tan et al., 2019; Long et al., 2020; Tian et al.,

2020; Tian et al., 2022b; Xiao and Wang, 2022). The differential splicing of the genes related to

spliceosome was indicated to assist the maintenance of gill homeostasis and improve the tolerance

to hypoxia stress of spotted sea bass (*Lateolabrax maculatus*) (Ren et al., 2022). In another example,

the expression of AS variants of *trim25* gene, which encodes an E3 ubiquitinated ligase of Tripartite

Motif (TRIM) family, were up-regulated in Japanese flounder (*Paralichthys olivaceus*) in response

to poly I:C treatment, indicative of their positively regulatory role of antiviral activity (Guo et al.,

2022). Therefore, a comprehensive understanding of AS regulations associated with salt stress

tolerance in greater amberjack allows us to fully understand the genetic basis underlying their

physiological responses to fluctuating ambient salinity.

In this study, we performed a genome-wide investigation of AS dynamics in greater amberjack

juveniles in response to the exposure to undesired salinity regimes. We further compared AS

regulations to differential gene expression to explore their respective contributions to tolerance to

water salinity fluctuations. Our findings will enrich our understanding of the molecular mechanisms

underlying the adaptation to salinity stress in euryhaline fish, which provides theoretical guidance

for the conservation of fishery resources under the increasingly extreme global climate change.

2 Materials and methods

2.1 Data preprocessing and sequence alignment

In our previous study, we sequenced the transcriptomes for two osmoregulatory tissues, gills and

kidney, for greater amberjack juveniles reared under different salinity regimes: 10, 30 and 40 parts

per thousand (ppt) (National Center for Biotechnology Information (NCBI) accession number:

GSE220485) (Liu et al., 2023). Here, we re-used these RNA-seq data to investigate the role of AS

regulations in greater amberjack in response to ambient salinity changes. Following the data

preprocessing pipeline described in Liu et al. (2023), adapter contamination and low-quality reads

were filtered out using Trim Galore v. 0.6.6 (Krueger, 2012) with a quality score cut-off of 25 and

a read-length threshold of 50 bp. A genome index was constructed for the reference genome of

greater amberjack (NCBI GCA_002260705.1) by the hisat2-built utility of HISAT2 package v. 2.2.1

(Kim et al., 2019). High-quality reads were mapped to the reference genome by the hisat2 program.

For each sample, only the uniquely aligned reads were retained for downstream analysis. The

alignment output was further assembled into full-length transcripts using StringTie v2.2.1(Kovaka

et al., 2019). All sets of assembled transcripts were merged into a nonredundant set of transcripts

using StringTie's merge mode. The assembled transcripts were annotated by comparing against the

reference annotations obtained from the NCBI RefSeq database using GFFcompare v. 0.11.2 (Pertea

and Pertea, 2020).

2.2 Alternative splicing investigation and functional enrichment analysis

rMATS v. 4.0.2 (Shen et al., 2014) was employed to detect alternative splicing events of five major

types: skipped exon (SE), alternative 5' splice site (A5SS), alternative 3' splice site (A3SS),

mutually exclusive exons (MXE), and retained intron (RI) (Mehmood et al., 2020). For each event,

the inclusion level of an alternative fragment in each sample was measured by "percent spliced in

(PSI)". The changes in average PSI values were computed between the hypo- (10 ppt) or hyper-

salinity (40 ppt) treatment and the control group (30 ppt, CK) in gills and kidney, respectively, to

investigate the shift in splicing patterns in response to ambient salinity alterations. For each

comparison, the events with an FDR < 0.05 were considered to be significantly different between

the two groups, and the genes with any differential splicing event were denoted as "DSGs". Similar

to other animals, SE events were the dominant AS type in greater amberjack, thus here we primarily

focused on the functional importance of SE events. For the convenience, the genes with a larger PSI

value of SE event under a treatment condition than CK were going to produce a higher proportion

of the transcripts containing the alternative exon, and were denoted as "more exon-inclusive DSGs",

while those with a smaller PSI value were denoted as "less exon-inclusive DSGs", which would

give rise to more transcripts with the alternative exon skipped.

To access the functional importance of the DSGs, an OrgDb database was first built for greater

amberjack using the R package AnnotationForge v. 1.38.1 (Carlson and Pages, 2017), based on the

gene annotations obtained in our previous study (Liu et al., 2023). For both gills and kidney, Gene

Ontology (GO) enrichment analysis was carried out for the DSGs via the enrichGO function of

clusterProfiler package v. 4.6.2 (Wu et al., 2021). P-value < 0.05 was set as the criteria for statistical

significance. For each tissue, the more exon-inclusive DSGs were compared across different salinity

treatments, and the results were visualized by UpsetR package v. 1.4.0 (Conway et al., 2017). GO

enrichment analysis was implemented for the DSGs 1) specific to 10 ppt saline treatment, 2) specific

to 40 ppt saline treatment, and 3) identified under both treatments, respectively.

2.3 Premature termination codons (PTCs) induced by skipped exon events

One of the impacts of alternative splicing is the introduction of premature termination codons (PTCs)

to transcribed isoforms, which may cause deficiencies of protein structure and function or even lead

to nonfunctional proteins (Yeh and Hwang, 2020). PTCs account for approximately 20% of genetic

defects in human diseases and are often associated with severe phenotypes (Oren et al., 2017). Here,

we identified putative PTCs induced by SE events in greater amberjack and investigated their

impacts on the product proteins. The coding sequence (CDS) regions were first predicted for each

isoform based on the annotations of the assembled transcripts. Only those with unambiguous

annotations of CDS, translation initiation codon, and termination codon were retained for

downstream analysis. The coding sequences of each target isoform were extracted from the genome

of greater amberjack, and all the putative termination codons were screened out in the coding regions.

For the isoforms with multiple termination codons, the first appeared termination codon was

considered as the PTC. The impact of PTCs on the structure of product proteins was assessed

according to the distance between the PTC and the 5' end of transcripts, where a small distance

meant a large influence on the completeness of the protein structure.

2.4 Comparison between alternative splicing and differential gene expression

To further investigate the different roles between AS and gene expression in modulating salinity

oscillation tolerance in greater amberjack, we compared the more exon-inclusive DSGs to the up-

regulated differentially expressed genes (DEGs) under each treatment scenario. The DEGs were

detected following the same criteria described in our previous study (Liu et al., 2023). GO

enrichment analyses were then conducted for 1) the genes specifically regulated by differential AS

(DS-specific genes), and 2) genes specifically regulated by differential expression (DE-specific

genes).

3 Results

3.1 Differential splicing regulate salinity adaptation in gills of greater amberjack

In the gills of greater amberjack, compared to the CK group, RI events exhibited the largest

alterations under both hypo- and hyper-salinity treatments, followed by SE and A3SS events (Figure

1B). When exposed to 10 ppt saline, 583 SE events were detected in 461 genes, where the

transcriptional products of 182 genes displayed a higher inclusive level of the alternative exons,

while the rest 279 genes were prone to skip the alternative exons. GO enrichment analysis of these

more-exon inclusive genes revealed an enrichment of proton transmembrane transport and energy

metabolic processes, such as glucose and alcohol metabolism, and sulfur compound biosynthesis

(Figure 1C). Under 40 ppt salinity stress, 187 more exon-inclusive and 239 less exon-inclusive

genes were identified compared to CK, where the more exon-inclusive genes were significantly

overrepresented in the biological processes related to stimulus response, proton transmembrane

transport, cilium assembly and epithelial cell differentiation (Figure 1D).

These DSGs were then grouped into 8 clusters according to their changes of PSI across the

different salinity regimes (10, 30 and 40 ppt) (Figure 1E). In Cluster 1, 2 and 6, the majority of

DSGs exhibited the highest inclusive level of the alternative exons in 10 ppt saline than the other

two conditions. GO enrichment results showed that these genes were mainly involved in erythrocyte

differentiation, reactive oxygen species (ROS) metabolism, sodium ion transport, pH regulation,

regulation of developmental process, and positive regulation of Wnt signaling pathway. In contrast,

the DSGs in Cluster 3 and 4 were of the highest exon inclusion in hyper-salinity treatment and these

genes mainly participated in sodium ion transport, lipid transport, and regulation of cell division.

These results suggested that greater amberjack activated sodium ion transport in gills to cope with

the deviation of normal water salinity. A large group of DSGs displayed less exon inclusion under

either hypo- or hyper-salinity than CK (Cluster 5 and 8), including those associated with pigment

metabolic process, neurogenesis, cell polarity establishment, spindle localization, and DNA

recombination.

Figure 1. Alternative splicing reprogramming underlying salinity acclimation in gills of greater

amberjack. (A) Schematic diagram showing five primary types of alternative splicing events we

analyzed in this study. (B) Number of differential splicing events of each alternative splicing type

in gills in response to hypo- (10 ppt) and hyper-salinity (40 ppt) stresses. (C) Representative Gene

Ontology (GO) terms significantly enriched for the more exon-inclusive genes upon the treatment

of 10 ppt saline. (D) Representative GO terms significantly enriched for the more exon-inclusive

genes upon the treatment of 40 ppt saline. (E) Heatmap showing the percent spliced in (PSI) of the

alternative exon of the exon-skipping genes across different salinity regimes. Representative GO

terms enriched for the genes of each cluster are listed above the heatmap.

3.2 Alternative splicing responds to salinity changes in kidney of greater amberjack

In the kidney, 759 significant SE events in 595 DSGs were detected under salinity stress at 10 ppt

(Figure 2A). Of them, 276 DSGs displayed higher inclusion level of the alternative exons and these

genes were found to be overrepresented in the GO terms of sodium ion transmembrane transport,

organonitrogen compound catabolic process, regulation of pH, erythrocyte homeostasis, response

to mechanical stimulus, myeloid cell homeostasis, and organophosphate metabolic process (Figure

S1). While the 319 less exon-inclusive DSGs were highly representative in pigment granule

transport, epithelial cell differentiation involved in kidney development, chromosome organization,

and regulation of growth (Figure S1). Under 40 ppt salinity stress, 215 genes related to sulfur

compound metabolism, aspartate family amino acid biosynthesis, lipid metabolism, and cellular

amide metabolism process were found to include more exons than CK (Figure S2). Conversely, 287

genes were identified as less exon-inclusive DSGs and were enriched in regionalization, cell

population proliferation, pigment metabolism, and epidermis development (Figure S2).

A cross comparison showed that, the exon inclusion level of 234 genes significantly increased

under both 10 and 40 ppt saline treatments over CK, and these DSGs were mainly associated with

sulfur compound biosynthetic process, peroxisome organization, organonitrogen compound

catabolic process, and cellular lipid metabolic process (Figure 2B). Comparatively, 361 and 268

more-exon inclusive DSGs were specifically induced upon 10 and 40 ppt salinity stress, respectively

(Figure 2B). Genes specifically responding to hypo-salinity stress were overrepresented in the GO

terms of homeostatic process, regulation of growth, sodium ion transport, cation homeostasis,

epithelial cell proliferation, and erythrocyte development, while those involved in hyper-salinity

responses were enriched in epidermal cell differentiation, epidermis development, Notch signaling

pathway, and pigment metabolism (Figure 2B).

Figure 2. Differential splicing in the kidney of greater amberjack in response to salinity changes.

(A) Number of differential splicing events of each alternative splicing type in kidney in response to

hypo- (10 ppt) and hyper-salinity (40 ppt) stresses. (B) Overlap of the more exon-inclusive genes

between 10 and 40 ppt saline treatments. The number of genes detected in each category was listed

in the upset intersection diagram. Representative Gene Ontology (GO) terms significantly enriched

for the genes of each category were listed at right, with the background color corresponding to the

bars of the upset plot.

3.3 Splicing-induced premature termination codons affect completeness of product proteins

Alternative cassette exons can moreover introduce multiple terminations to mRNAs and finally alter

the coding potential of product proteins (Nikonova et al., 2020). In greater amberjack, SE events

were found to give rise to PTCs under all four salinity treatments, which largely affect the

completeness and function of the resulting proteins (Figure 3). It is worthy to note that, in gills, the

majority of the introduced PTCs were located at the first 20% length of isoforms), that is, resulting

in truncated protein products with a >80% loss in length (Figure 3). These transcripts were likely to

be targeted by UPF factors and degraded by nonsense-mediated mRNA decay (NMD) pathway

(Brogna and Wen, 2009). Comparatively, in kidney, a large proportion of PTCs occurred in the

middle of the complete isoforms, indicating relatively smaller impacts on the functions of proteins.

Figure 3. Distribution of PTCs in different locations on transcripts upon hypo- (10 ppt) and hyper-

salinity (40 ppt) treatments in gills and kidney of greater amberjack. The location intervals of

transcripts are listed along the vertical axis, and dot sizes correspond to the relative proportion of

the PTCs in each location interval.

3.4 AS regulates different biological processes from gene expression under both hypo- and hyper-

salinity

We then compared the differentially spliced genes to those of differential expression under each

treatment to investigate their different roles in greater amberjack in mitigating the adverse impacts

of salinity stress. For both gills and kidney, the majority of genes were either modulated by

differential AS or differential expression, while only few genes were under the regulations of both

mechanisms. More importantly, gene transcription and alternative splicing were found to participate

in different physiological processes to enhance the tolerance of greater amberjack to water salinity

oscillations. In the gills, when treated with 10 ppt saline, DS-specific genes were mainly involved

in sulfur compound biosynthesis and proton transport, while the DE-specific genes were highly

represented in the GO terms of growth, cartilage and osteoblast development, oxygen transport, and

epithelium migration (Figure 4A). When exposed to 40 ppt salinity, the genes related to myeloid

cell development, proton transport, and positive regulation of stimulus responses were mainly

modulated by differential splicing, while those associated with cartilage development, extracellular

matrix organization, regulation of BMP signaling pathway and response to external stimulus were

differentially expressed (Figure 4B).

In kidney, when exposed to low salinity environment, alternative splicing mainly affected the

processes related to sodium ion transport, pH regulation, and erythrocyte homeostasis and

differentiation, while transcription regulation was found to modulate the processes associated with

kidney development (Figure 4C). Upon 40 ppt salinity, DSGs were significantly enriched in the GO

terms of cellular amide metabolism, sulfur compound biosynthesis and metabolism, and post-

transcriptional regulation of gene expression, while DEGs were mainly related to stimulus response,

ammonium transport, pH regulation, and cell growth (Figure 4D).

Figure 4. Representive Gene Ontology (GO) terms enriched for the genes that were specifically

regulated by differential expression (DE-specific genes), and specifically regulated by alternative

splicing (DS-specific genes).

4 Discussion

Water salinity is a key environmental factor affecting the yield of aquaculture (Callaway et al., 2012).

In general, fish can only adapt to a limited range of water salinity changes; for instance, the optimum

range of greater amberjack is 29 ‰−36 ‰ (Kültz, 2015; Ru et al., 2023). Deviation from the

appropriate salinity range would disrupt their normal physiological and immune activities, which

threatens their growth and survival (Zhang et al., 2017; Cui et al., 2019). Indeed, a previous study

revealed a substantial up-regulation of the genes associated with organ development in both gills

and kidney of greater amberjack juveniles to combat the adverse impact of salinity oscillations (Liu

et al., 2023). In this study, we performed a genome-wide investigation of AS dynamics in greater

amberjack juveniles in response to the exposure to undesired salinity regimes. The results revealed

that both osmoregulatory organs, gills and kidney, were subjected to a high level of differential

splicing during acclimation to hypo- and hyper-salinity stress, and these changes were mainly

associated with sodium ion transport, energy metabolism, and cellular renewal and development,

(Figures 1 and 2; Figure S1 and S2), which highlights the important role of AS in supporting the

physiological reprogramming of greater amberjack to cope with short-term salinity fluctuation.

In gills, the genes involved in energy metabolism were significantly differentially spliced when

exposed to 10 ppt saline (Figure 1C). For example, one of them encodes an enzyme hexose-6-

phosphate dehydrogenase (H6PD) of endoplasmic reticulum. H6PD catalyzes the first two reactions

of the oxidative branch of the pentose phosphate pathway: the conversion from glucose-6-phosphate

and $NADP^+$ to 6-phosphogluconate and NADPH (Senesi et al., 2010; Ge et al., 2020). The produced

NADPH can reduce oxidized glutathione (GSSG) to reduced glutathione (GSH), which acts a key

role in scavenging ROS and maintaining cellular homeostasis, especially upon environmental

stresses (Senesi et al., 2010; Tomanek, 2012; Tomanek, 2015). Moreover, alcohol metabolism was

also enhanced in the gills. For most saltwater fish, a decrease in water salinity alters their oxygen

requirements in gills, thus may increase the production of lactate (Ern et al., 2014; Taugbøl et al.,

2022). To combat the adverse impact of lactate buildup, crucian carp and goldfish (*Carassius*,

Cyprinidae) can further covert lactic acid to ethanol and excrete it into ambient water via the special

coupling of pyruvate dehydrogenase complex (PDC) and alcohol dehydrogenase (ADH) in gills,

which largely improve their tolerance to low oxygen environment (Fagernes et al., 2017). However,

no studies yet have investigated the detailed alcohol metabolic processes in greater amberjack, and

future work is still required. When reared in a hyposalinity environment, the genes related to

stimulus response, for example, *RNF111*, were activated in gills (Figure 1D). The coding product of

RNF111, ring finger protein 111, was revealed to positively regulate DNA damage recognition in

nucleotide excision repair, thus alleviating stress-induced DNA damage (Poulsen et al., 2013).

Moreover, cilum assembly and epithelial cell differentiation were also differentially regulated by

AS. Previous studies have revealed that cell proliferation and differentiation of gill epithelia were

substantially activated in response to high salinity exposure, which can promote the osmoprotection

of gill cells (Chretien, 1986; Fiol et al., 2009). Moreover, the leakiness of epithelial tight junctions

was also increased to enhance the extrusion of sodium ions across the gill epithelium (Kültz, 2015).

These findings highlighted the key role of epithelial remodeling in fish gills in hypo-osmoregulation.

In kidney, sodium ion transport and pH regulation were enhanced to combat the decrease in

water salinity (Figure S1). Kidney is the main osmoregulatory organ in fish. The enhancement of

ion transport in kidney can increase salt excretion and maintain the ion balance and blood osmolality

in greater amberjack. Moreover, the splicing modes of the genes related to erythrocyte homeostasis

were also significantly changed. A low ambient salinity may lower the internal osmolality of fish,

which further alters the size and shape of red blood cells (Jensen et al., 2002). A rapid and efficient

regulation of blood cell volume can guarantee the oxygen transfer between red blood cells and

kidney. When exposed to hyper-salinity stress, cellular amide metabolic process, epidermis

development, Notch signaling pathway, and pigment metabolism were activated to cope with the

increased salinity (Figure 2B; Figure S2). The breakdown of amide/peptide bonds of proteins

moreover gives rise to free amino acids, which are important cellular osmolytes in euryhaline fish

(Aragão et al., 2010). A meta-analysis across 106 study cases showed that the cellular content of

free amino acids was positively correlated with the elevation of salinity, highlighting their essential

role in maintaining the homeostasis of osmotic pressure in fish in hyper-salinity environments

(Huang et al., 2023). It is noteworthy that energy metabolism, including organonitrogen compound

catabolism and lipid metabolism, were substantially activated by both 10 and 40 ppt saline

treatments (Figure 2B). For both hypo- and hyper-salinity acclimation, extra energy costs are

required by fish kidney for the osmoregulatory work, and the metabolism of lipids and proteins can

serve as alternative fuel sources to enhance the energy supply (Aragão et al., 2010; Nguyen et al.,

2016).

It is worthy to note that, in both gills and kidney, the most majority of differentially spliced

genes were not differentially expressed, and these two mechanisms were found to modulate different

biological processes (Figure 4). For example, in kidney, the genes involved in sodium ion transport

and pH regulation were specifically differentially spliced upon 10 ppt salinity stimulus, while those

associated with pronephros development were primarily under the regulation of differentially

expression (Figure 4C). These indicated that AS is functionally nonredundant and plays a

complementary role to gene transcription in regulating protein expression in greater amberjack in

response to ambient salinity fluctuations. It is consistent with observations in other fish species,

such as salmonid fish (*Salvelinus alpinus*) (Jacobs and Elmer, 2021)and spotted sea bass (Tian et al.,

2020). AS and gene transcription are regulated mainly by independent mechanisms. The alternative

procedures of pre-mRNA splicing are determined by the different recognitions and usages of

splicing sites of spliceosome complex (Wang et al., 2015; Liu et al., 2022). In addition, the spliced

mRNA products carrying PTCs would be targeted and degraded by the mRNA-surveillance

mechanism NMD pathway (Stamm et al., 2005; Brogna and Wen, 2009). Our study also observed

a large impact of PTCs on the structures and functions of the resulting proteins in both gills and

kidney of greater amberjack (Figure 3). These features allow AS to extend and modulate the mRNA

and protein repertoire of selectively constrained genes without altering their initial expressions,

which acts as a key mechanism in the development of phenotypic plasticity and rapid ecological

adaptation (Marden, 2008; Barbosa-Morais et al., 2012; Bradley et al., 2012; Jacobs and Elmer,

2021; Steward et al., 2022). Our current study adds further evidence for the nonnegligible role of

AS in regulating stress tolerance in teleost fishes.

5 Conclusion

In this study, we provide a new insight into the role of alternative splicing in regulating physiological

remodeling of greater amberjack to cope with the adverse impacts of fluctuated water salinity. The

results show that alternative splicing plays a relatively independent role in fine-tuning protein

expression in both gills and kidney of greater amberjacker in the early responses to salinity changes.

Differential splicing primarily results in structural and functional changes of the proteins involved

in sodium ion and proton transport, energy metabolism, and cell development and differentiation,

which may improve the adaptation of greater amberjacker to salinity stress. These findings unravel

the deeper side of the genetic basis underlying salinity acclimation in greater amberjack in response

to the fluctuating marine environment, and provide a genomic view for the conservation of marine

fishery resources.

Acknowledgement

This research was financed by grants from the National Natural Science Foundation of China

(32201420).

Data availability statements

Data were obtained from the National Center for Biotechnology Information (NCBI) with accession

number: GSE220485.

Conflict of interst statement

The authors declare that the research was conducted in the absence of any commercial or financial

relationships that could be construed as a potential conflict of interest.

References

Aragão, C., Costas, B., Vargas-Chacoff, L., Ruiz-Jarabo, I., Dinis, M. T., Mancera, J. M., and Conceição, L. E. 2010. Changes in plasma amino acid levels in a euryhaline fish exposed to different environmental salinities. Amino acids, 38: 311-317.

Araki, K., Aokic, J.-y., Kawase, J., Hamada, K., Ozaki, A., Fujimoto, H., Yamamoto, I., et al. 2018. Whole genome sequencing of greater amberjack (*Seriola dumerili*) for SNP identification on aligned scaffolds and genome structural variation analysis using parallel resequencing. International journal of genomics, 2018.

Barany, A., Gilannejad, N., Alameda-López, M., Rodríguez-Velásquez, L., Astola, A., Martínez-Rodríguez, G., Roo, J., et al. 2021. Osmoregulatory plasticity of juvenile greater amberjack (*Seriola dumerili*) to environmental salinity. Animals, 11: 2607.

Barbosa-Morais, N. L., Irimia, M., Pan, Q., Xiong, H. Y., Gueroussov, S., Lee, L. J., Slobodeniuc, V., et al. 2012. The evolutionary landscape of alternative splicing in vertebrate species. science, 338: 1587-1593.

Bradley, R. K., Merkin, J., Lambert, N. J., and Burge, C. B. 2012. Alternative splicing of RNA triplets is often regulated and accelerates proteome evolution. PLoS biology, 10: e1001229.

Brogna, S., and Wen, J. 2009. Nonsense-mediated mRNA decay (NMD) mechanisms. Nature structural & molecular biology, 16: 107-113.

Callaway, R., Shinn, A. P., Grenfell, S. E., Bron, J. E., Burnell, G., Cook, E. J., Crumlish, M., et al. 2012. Review of climate change impacts on marine aquaculture in the UK and Ireland. Aquatic Conservation: Marine and Freshwater Ecosystems, 22: 389-421.

Carlson, M., and Pages, H. 2017. AnnotationForge: code for building annotation database packages. R package version, 1.

Cheng, L., Trenberth, K. E., Gruber, N., Abraham, J. P., Fasullo, J. T., Li, G., Mann, M. E., et al. 2020. Improved estimates of changes in upper ocean salinity and the hydrological cycle. Journal of Climate, 33: 10357-10381.

Chretien, M. 1986. Cell renewal and differentiation in the gill epithelium of fresh-or salt-water-adapted euryhaline fish as revealed by (^ 3H)-thymidine radioautography. Biol. Cell., 56: 137-150.

Conway, J. R., Lex, A., and Gehlenborg, N. 2017. UpSetR: an R package for the visualization of intersecting sets and their properties. bioinformatics, 33: 2938-2940.

Cui, Q., Qiu, L., Yang, X., Shang, S., Yang, B., Chen, M., Liu, X., et al. 2019. Transcriptome profiling of the low-salinity stress responses in the gills of the juvenile *Pseudopleuronectes yokohamae*. Comparative Biochemistry and Physiology Part D: Genomics and Proteomics, 32: 100612.

Durack, P. J., and Wijffels, S. E. 2010. Fifty-year trends in global ocean salinities and their relationship to broad-scale warming. Journal of Climate, 23: 4342-4362.

Durack, P. J., Wijffels, S. E., and Matear, R. J. 2012. Ocean salinities reveal strong global water cycle intensification during 1950 to 2000. science, 336: 455-458.

Ern, R., Huong, D. T. T., Phuong, N. T., Wang, T., and Bayley, M. 2014. Oxygen delivery does not limit thermal tolerance in a tropical eurythermal crustacean. Journal of Experimental Biology, 217: 809-814.

Fagernes, C. E., Stensløkken, K.-O., Røhr, Å. K., Berenbrink, M., Ellefsen, S., and Nilsson, G. E. 2017. Extreme anoxia tolerance in crucian carp and goldfish through neofunctionalization of duplicated genes creating a new ethanol-producing pyruvate decarboxylase pathway. Scientific

reports, 7: 7884.

Fiol, D. F., Sanmarti, E., Sacchi, R., and Kültz, D. 2009. A novel tilapia prolactin receptor is functionally

distinct from its paralog. Journal of Experimental Biology, 212: 2007-2015.

Galkanda-Arachchige, H. S., Davis, R. P., Nazeer, S., Ibarra-Castro, L., and Davis, D. A. 2021. Effect of

salinity on growth, survival, and serum osmolality of red snapper, *Lutjanus campechanus*. Fish

Physiology and Biochemistry, 47: 1687-1696.

Ge, T., Yang, J., Zhou, S., Wang, Y., Li, Y., and Tong, X. 2020. The role of the pentose phosphate pathway

in diabetes and cancer. Frontiers in Endocrinology, 11: 507678.

Guo, T., Yang, Y., Meng, F., Wang, S., Xia, S., Qian, Y., Li, M., et al. 2020. Effects of low salinity on gill

and liver glycogen metabolism of great blue-spotted mudskippers (*Boleophthalmus*

pectinirostris). Comparative Biochemistry and Physiology Part C: Toxicology & Pharmacology,

230: 108709.

Guo, Y., Sun, Z., Zhang, Y., Wang, G., He, Z., Liu, Y., Ren, Y., et al. 2022. Molecular identification and

function characterization of four alternative splice variants of trim25 in Japanese flounder

(*Paralichthys olivaceus*). Fish & Shellfish Immunology, 120: 142-154.

Helm, K. P., Bindoff, N. L., and Church, J. A. 2011. Observed decreases in oxygen content of the global

ocean. Geophysical Research Letters, 38.

Huang, M., Gao, Q., Yang, X., Jiang, W., Hao, L., Yu, Y., and Tian, Y. 2023. Free amino acids in response

to salinity changes in fishes: relationships to osmoregulation. Fish Physiology and Biochemistry,

49: 1031-1042.

Jacobs, A., and Elmer, K. R. 2021. Alternative splicing and gene expression play contrasting roles in the

parallel phenotypic evolution of a salmonid fish. Molecular Ecology, 30: 4955-4969.

Jensen, F. B., Lecklin, T., Busk, M., Bury, N., Wilson, R., Wood, C., and Grosell, M. 2002. Physiological

impact of salinity increase at organism and red blood cell levels in the European flounder

(*Platichthys flesus*). Journal of experimental marine biology and ecology, 274: 159-174.

Kim, D., Paggi, J. M., Park, C., Bennett, C., and Salzberg, S. L. 2019. Graph-based genome alignment

and genotyping with HISAT2 and HISAT-genotype. Nature biotechnology, 37: 907-915.

Kovaka, S., Zimin, A. V., Pertea, G. M., Razaghi, R., Salzberg, S. L., and Pertea, M. 2019. Transcriptome

assembly from long-read RNA-seq alignments with StringTie2. Genome biology, 20: 1-13.

Krueger, F. 2012. Trim Galore: a wrapper tool around Cutadapt and FastQC to consistently apply quality

and adapter trimming to FastQ files, with some extra functionality for MspI-digested RRBS-

type (Reduced Representation Bisufite-Seq) libraries. URL http://www. bioinformatics.

babraham. ac. uk/projects/trim_galore/.(Date of access: 28/04/2016): 71.

Kültz, D. 2015. Physiological mechanisms used by fish to cope with salinity stress. The Journal of

experimental biology, 218: 1907-1914.

Lacy, E. R., Nicholas, J. S., and Colglazier, J. 2023. Changes in the composition of coelomic fluid in the

Atlantic stingray, *Hypanus sabina*, upon dilution of the environmental salinity: osmoregulatory

and pH implications. Ichthyological Research, 70: 293-300.

Lehtonen, T. K., Wong, B. B., and Kvarnemo, C. 2016. Effects of salinity on nest-building behaviour in

a marine fish. BMC ecology, 16: 1-9.

Liu, Q., Fang, L., and Wu, C. 2022. Alternative splicing and isoforms: from mechanisms to diseases.

Genes, 13: 401.

Liu, Y., Yang, Y., Qin, R., Peng, Y., Huang, Y., Zhu, C.-H., Li, G., et al. 2023. Divergent molecular

responses of greater amberjack (*Seriola dumerili*) to acute salinity stress revealed by

comparative transcriptome analysis. Frontiers in Marine Science, 10: 1185015.

Long, Y., Li, X., Liu, R., and Cui, Z. 2020. Transcriptional programs underlying cold acclimation of
common carp (*Cyprinus carpio* L.). Frontiers in Genetics, 11: 556418.

Lu, M., Su, M., Liu, N., and Zhang, J. 2022. Effects of environmental salinity on the immune response
of the coastal fish *Scatophagus argus* during bacterial infection. Fish & Shellfish Immunology,
124: 401-410.

Marden, J. 2008. Quantitative and evolutionary biology of alternative splicing: how changing the mix of
alternative transcripts affects phenotypic plasticity and reaction norms. Heredity, 100: 111-120.

Mehmood, A., Laiho, A., Venäläinen, M. S., McGlinchey, A. J., Wang, N., and Elo, L. L. 2020.
Systematic evaluation of differential splicing tools for RNA-seq studies. Briefings in
bioinformatics, 21: 2052-2065.

Navarro-Guillen, C., Yufera, M., and Perera, E. 2022. Biochemical features and modulation of digestive
enzymes by environmental temperature in the greater amberjack, *Seriola dumerili*. Frontiers in
Marine Science, 9: 960746.

Nguyen, T. V., Jung, H., Nguyen, T. M., Hurwood, D., and Mather, P. 2016. Evaluation of potential
candidate genes involved in salinity tolerance in striped catfish (*Pangasianodon hypophthalmus*)
using an RNA-Seq approach. Marine Genomics, 25: 75-88.

Nikonova, E., Kao, S.-Y., and Spletter, M. L. 2020. Contributions of alternative splicing to muscle type
development and function. *In* Seminars in Cell & Developmental Biology, pp. 65-80. Elsevier.

Omerspahic, M., Al-Jabri, H., Siddiqui, S. A., and Saadaoui, I. 2022. Characteristics of desalination brine
and its impacts on marine chemistry and health, with emphasis on the Persian/Arabian gulf: a
review. Frontiers in Marine Science, 9: 845113.

Oren, Y. S., Pranke, I. M., Kerem, B., and Sermet-Gaudelus, I. 2017. The suppression of premature
termination codons and the repair of splicing mutations in CFTR. Current opinion in
pharmacology, 34: 125-131.

Peng, Y., Shi, H., Liu, Y., Huang, Y., Zheng, R., Jiang, D., Jiang, M., et al. 2022. RNA Sequencing
analysis reveals divergent adaptive response to hypo-and hyper-salinity in greater amberjack
(*Seriola dumerili*) juveniles. Animals, 12: 327.

Pereira, I. R., Valls, M., Aoufi, S., Bienentreu, O. C., Huang, Y., and Cabanellas-Reboredo, M. 2023.
Socio-ecological overview of the greater amberjack fishery in the Balearic Islands. Fishes, 8:
184.

Pertea, G., and Pertea, M. 2020. GFF utilities: GffRead and GffCompare. F1000Research, 9.

Poulsen, S. L., Hansen, R. K., Wagner, S. A., van Cuijk, L., van Belle, G. J., Streicher, W., Wikström, M.,
et al. 2013. RNF111/Arkadia is a SUMO-targeted ubiquitin ligase that facilitates the DNA
damage response. Journal of Cell Biology, 201: 797-807.

Ren, Y., Tian, Y., Mao, X., Wen, H., Qi, X., Li, J., Li, J., et al. 2022. Acute hypoxia changes the gene
expression profiles and alternative splicing landscape in gills of spotted sea bass (*Lateolabrax*
maculatus). Frontiers in Marine Science, 9: 1024218.

Röthig, T., Trevathan-Tackett, S. M., Voolstra, C. R., Ross, C., Chaffron, S., Durack, P. J., Warmuth, L.
M., et al. 2023. Human-induced salinity changes impact marine organisms and ecosystems.
Global change biology, 29: 4731-4749.

Ru, X., Huang, Y., Shi, H., Peng, Y., Hao, R., Yang, T., Zhu, K., et al. 2023. RNA-seq of hypo-and hyper-
salinity stress-response transcriptome in the liver of greater amberjack (*Seriola dumerili*)
juveniles. Aquaculture Reports, 29: 101498.

Senesi, S., Csala, M., Marcolongo, P., Fulceri, R., Mandl, J., Banhegyi, G., and Benedetti, A. 2010.
Hexose-6-phosphate dehydrogenase in the endoplasmic reticulum.

Shen, S., Park, J. W., Lu, Z.-x., Lin, L., Henry, M. D., Wu, Y. N., Zhou, Q., et al. 2014. rMATS: robust
and flexible detection of differential alternative splicing from replicate RNA-Seq data.
Proceedings of the National Academy of Sciences, 111: E5593-E5601.

Skliris, N., Zika, J. D., Nurser, G., Josey, S. A., and Marsh, R. 2016. Global water cycle amplifying at
less than the Clausius-Clapeyron rate. Scientific reports, 6: 38752.

Smyth, K., and Elliott, M. 2016. Effects of changing salinity on the ecology of the marine environment.
Stressors in the marine environment: 161-174.

Stamm, S., Ben-Ari, S., Rafalska, I., Tang, Y., Zhang, Z., Toiber, D., Thanaraj, T., et al. 2005. Function
of alternative splicing. Gene, 344: 1-20.

Steward, R. A., de Jong, M. A., Oostra, V., and Wheat, C. W. 2022. Alternative splicing in seasonal
plasticity and the potential for adaptation to environmental change. Nature Communications,
13: 755.

Tan, S., Wang, W., Tian, C., Niu, D., Zhou, T., Jin, Y., Yang, Y., et al. 2019. Heat stress induced alternative
splicing in catfish as determined by transcriptome analysis. Comparative Biochemistry and
Physiology Part D: Genomics and Proteomics, 29: 166-172.

Taugbøl, A., Solbakken, M. H., Jakobsen, K. S., and Vøllestad, L. A. 2022. Salinity-induced
transcriptome profiles in marine and freshwater threespine stickleback after an abrupt 6-hour
exposure. Ecology and Evolution, 12: e9395.

Tian, B., Bian, Y., Bian, D.-J., Gao, Y., Zhang, X., Zhou, S.-W., Zhang, Y.-H., et al. 2022a. Knowledge
mapping of alternative splicing of cancer from 2012 to 2021: A bibliometric analysis. Frontiers
in Oncology, 12: 1068805.

Tian, Y., Gao, Q., Dong, S., Zhou, Y., Yu, H., Liu, D., and Yang, W. 2022b. Genome-wide analysis of
alternative splicing (AS) mechanism provides insights into salinity adaptation in the livers of
three euryhaline teleosts, including *Scophthalmus maximus*, *Cynoglossus semilaevis* and
Oncorhynchus mykiss. Biology, 11: 222.

Tian, Y., Wen, H., Qi, X., Zhang, X., Sun, Y., Li, J., He, F., et al. 2020. Alternative splicing (AS)
mechanism plays important roles in response to different salinity environments in spotted sea
bass. International journal of biological macromolecules, 155: 50-60.

Tomanek, L. 2012. Environmental proteomics of the mussel *Mytilus*: implications for tolerance to stress
and change in limits of biogeographic ranges in response to climate change. Oxford University
Press.

Tomanek, L. 2015. Proteomic responses to environmentally induced oxidative stress. The Journal of
experimental biology, 218: 1867-1879.

Ullah, A., Ahmed, M., Raza, S. A., and Ali, S. 2021. A threshold approach to sustainable development:
nonlinear relationship between renewable energy consumption, natural resource rent, and
ecological footprint. Journal of environmental management, 295: 113073.

Wan, Q., and Su, J. 2015. Transcriptome analysis provides insights into the regulatory function of
alternative splicing in antiviral immunity in grass carp (*Ctenopharyngodon idella*). Scientific
reports, 5: 12946.

Wang, Y., Liu, J., Huang, B., Xu, Y. M., Li, J., Huang, L. F., Lin, J., et al. 2015. Mechanism of alternative
splicing and its regulation. Biomedical reports, 3: 152-158.

Wu, T., Hu, E., Xu, S., Chen, M., Guo, P., Dai, Z., Feng, T., et al. 2021. clusterProfiler 4.0: A universal

enrichment tool for interpreting omics data. The innovation, 2.

Xiao, J., and Wang, W.-X. 2022. Patterns and Crucial Regulation of Alternative Splicing During Early

Development in Zebrafish. Journal of Molecular Biology, 434: 167821.

Yeh, J. T., and Hwang, T. C. 2020. Positional effects of premature termination codons on the biochemical

and biophysical properties of CFTR. The Journal of physiology, 598: 517-541.

Zhang, Y. T., Huang, S., Qiu, H. T., Li, Z., Mao, Y., Hong, W. S., and Chen, S. X. 2017. Optimal salinity

for rearing Chinese black sleeper (*Bostrychus sinensis*) fry. Aquaculture, 476: 37-43.

Zidan, E. M., Goma, A. A., Tohamy, H. G., Soliman, M. M., and Shukry, M. 2022. Insight study on the

impact of different salinity levels on behavioural responses, biochemical stress parameters and

growth performance of African catfish (*Clarias gariepinus*). Aquaculture Research, 53: 2750-

2759.

Zupa, R., Rodríguez, C., Mylonas, C. C., Rosenfeld, H., Fakriadis, I., Papadaki, M., Pérez, J. A., et al.

2017. Comparative study of reproductive development in wild and captive-reared greater

amberjack *Seriola dumerili* (Risso, 1810). PLoS One, 12: e0169645.

564