How much warming can mosquito vectors tolerate?

Lisa I. Couper, Desire Uwera Nalukwago, Kelsey P. Lyberger, Johannah E. Farner, Erin A. Mordecai

Abstract

Climate warming is expected to substantially impact the global landscape of mosquito-borne disease, but these impacts will vary across disease systems and regions. Understanding which diseases, and where within their distributions, these impacts are most likely to occur is critical for preparing public health interventions. While research has centered on potential warming-driven expansions in vector transmission, less is known about the potential for vectors to experience warming-driven stress or even local extirpations. In conservation biology, species risk from climate warming is often quantified through vulnerability indices such as thermal safety margins – the difference between an organism's upper thermal limit and its habitat temperature. Here, we estimated thermal safety margins for 8 mosquito species (including *Aedes aegypti* and *Anopheles gambiae*) that are the vectors of malaria, dengue, chikungunya, Zika, West Nile and other major arboviruses, across their known ranges to investigate which mosquitoes and regions are most and least vulnerable to climate warming. We find that several of the most globally important mosquito vector species have positive thermal safety margins across the majority of their ranges when realistic assumptions of mosquito behavioral thermoregulation are incorporated. On average, the lowest climate vulnerability, in terms of both the magnitude and duration of thermal safety, was just south of the equator and at northern temperate range edges, and the highest climate vulnerability was in the subtropics. Mosquitoes living in regions largely comprised of desert and xeric shrubland biomes, including the Middle East, the western Sahara, and southeastern Australia have the highest climate vulnerability across vector species.

Introduction

Climate warming is already causing shifts in the distribution and species interactions of many taxa, affecting ecosystem processes and health, and these effects are projected to accelerate with continued carbon emissions (Elsen *et al.* 2022; Grimm *et al.* 2013; Urban 2015). In particular, climate warming affects many ectotherms due to their temperature-sensitive life history and population dynamics, altering their geographic ranges and imposing thermal stress near their upper thermal limits (Deutsch *et al.* 2008; Parmesan 2006). Mosquitoes, which transmit many of the most important and high-burden infectious diseases worldwide, including malaria, dengue, Zika, and West Nile, are projected to experience changes in population dynamics, range limits, and transmission potential due to climate change (Harrigan *et al.* 2014; Kraemer *et al.* 2019; Ryan *et al.* 2019, 2021). Near cool range edges, warming is widely expected to promote mosquito population growth and disease transmission (Chen *et al.* 2010; Medlock & Leach 2015; Mordecai *et al.* 2019; Parmesan & Yohe 2003; Ryan *et al.* 2019). However, the impacts of

warming across other portions of mosquito ranges are less clear and will depend on the extent of
warming (*e.g.*, the magnitude and duration of high temperatures), as well as species
physiological limits and response strategies. In particular, ectotherm physiology theory and
empirical work predict that upper thermal limits may restrict species persistence in currently
suitable ranges under excessive warming (Deutsch *et al.* 2008; Huey & Berrigan 2001; Kearney
et al. 2009a; Kellermann *et al.* 2012; Pinsky *et al.* 2019; Pörtner *et al.* 2006; Sunday *et al.* 2014).
Understanding which mosquito species and regions will be most impacted by warming is critical
for preparing targeted vector control and disease prevention strategies.
The potential impacts of climate change across species and broad spatial scales are often
quantified through vulnerability indices such as 'thermal safety margins'—the difference
between an organism's critical thermal maximum and its habitat temperature—(sometimes used
interchangeably with 'warming tolerance'; Angilletta 2009; Clusella-Trullas *et al.* 2021; Deutsch
et al. 2008). Thermal safety margins, which reflect the amount of additional warming an
organism could experience before trait performance is inhibited, have been estimated for
hundreds of ectotherm species including lizards, snakes, fish, and insects (Deutsch *et al.* 2008;
Diamond *et al.* 2011; Sunday *et al.* 2019, 2014; Vinagre *et al.* 2019). These prior studies have
often found the lowest thermal safety margins, and thus the highest vulnerability to warming, for
species in the tropics. From a conservation standpoint, this finding has raised concern given the
high biodiversity contained in the tropics. However, whether this pattern of maximum thermal
stress in the tropics applies for species that threaten human health, such as mosquitoes that vector
pathogens, has not been rigorously evaluated.
Thermal safety margins can provide biased estimates if the estimated critical thermal
maximum and/or environmental temperatures do not reflect those relevant to the organism
(Clusella-Trullas *et al.* 2021). For example, air temperature in full sun, as is often captured by
weather station data sources, can differ drastically from an organism's body temperatures
(Kearney *et al.* 2009b; Sunday *et al.* 2014). Similarly, environmental temperature estimates with
coarse temporal resolution (*e.g.,* daily or monthly scales) may fail to capture short-term thermal
extremes, which can have major impacts on organismal fitness and population growth (Buckley
& Huey 2016a, b; Dowd *et al.* 2015; Vasseur *et al.* 2014). To address these limitations, recent
advances in climate monitoring and microclimate and biophysical modeling have enabled
increasingly accurate estimates of realized environmental temperatures on time scales relevant to
short ectotherm life cycles (Hersbach *et al.* 2020; Kearney & Porter 2017, 2020).
To date, no work has systematically estimated thermal safety margins for the mosquitoes
that transmit major human pathogens. Here, we incorporate these recent advances in
physiological and bioclimatic modeling to estimate thermal safety margins for 8 major mosquito
vector species across their ranges to identify the disease systems and locations that are most and
least vulnerable to climate warming. We specifically ask: how much buffering do mosquitoes
have from thermal extremes in their current ranges? We define thermal safety margins as the
difference between a species' critical thermal maximum, estimated in the laboratory for a range
of life history traits under constant temperatures, and the hottest hourly body temperature

experienced in the coolest microhabitat available (Clusella-Trullas *et al.* 2021; Pinsky *et al.*
2019). To ensure that our estimates provide a meaningful metric of climate vulnerability, we
explicitly: 1) incorporate the realistic potential for mosquito movement to cooler, fully-shaded
microhabitats (*i.e.*, 'behavioral thermoregulation'); 2) estimate operative mosquito body
temperature rather than air temperature, as these can differ markedly (Kearney *et al.* 2009b;
Sunday *et al.* 2014; Woods *et al.* 2015); and 3) use hourly temperature data to capture the impact
of short-term thermal extremes that may be missed with more temporally coarse temperature
estimates (*e.g.*, daily, monthly). Further, as mosquito climate vulnerability will depend not only
on the difference in critical thermal maxima and body temperatures, but also how often and for
how long critical thermal maxima are exceeded, we also estimated the time spent in thermal
danger (here, the number of consecutive hours or days when body temperatures exceed critical
thermal maxima).
93
Methods
Mosquito thermal tolerance data
We included the following 8 mosquito species in our investigation as these constituted major
disease vectors with available thermal tolerance estimates: *Aedes aegypti, Ae. albopictus,*
Anopheles gambiae, An. stephensi, Culex annulirostris, Cx. pipiens, Cx. quinquefasciatus, and
Cx. tarsalis (Table 1). Together, this includes the major vectors of 15 of the most important
human vector-borne pathogens (Table 1). As our measure of the estimated critical thermal
maximum (CT_{max}) for each species, we used the estimated thermal maxima (T_{max}) calculated
from thermal performance curves derived from laboratory experimental data on adult survival
and synthesized in Mordecai *et al.* (2019), Villena *et al.* (2022), and Cruz-Loya *et al.* (2024)
(Table 1). While for many species, thermal performance estimates are available for multiple life
history traits (*e.g.*, immature development rate, immature survival, biting rate), we used estimates
for adult survival for all species for consistency and because it is likely the life history trait that is
the most accurately and precisely estimated (*i.e.,* because the decline in trait performance above
the thermal optimum can be readily observed experimentally and estimated mathematically).
Further, it is likely the trait most strongly linked to overall fitness as mortality instantly and
irreversibly reduces fitness to zero. Lastly, the immature life stages occur in water, where
temperatures may differ substantially from that of the surrounding air, for which our
environmental temperature estimates apply. These estimates of the critical thermal maximum
(CT_{max}), as well as the critical thermal minima (CT_{min}) and optima (T_{opt}) were derived from
Bayesian thermal response models fit to empirical data on mosquito adult survival at different
constant temperatures. As a sensitivity analysis, we estimated thermal safety margins for these 8
species using the critical thermal maximum from the most thermally tolerant life history trait
(Supplemental Table S1, Supplemental Figure S1).
There is no consensus methodology for estimating organismal thermal tolerance, but
rather a wide range of methods including static and dynamic heat tolerance assays (*e.g.*, 'thermal
knockdowns') of varying durations (Bates & Morley 2020; Clusella-Trullas *et al.* 2021;

Hoffmann *et al.* 2013; Jørgensen *et al.* 2019; Lutterschmidt & Hutchison 1997; Rezende *et al.* 2014; Terblanche *et al.* 2007). Experimental methodology can impact the estimated CT_{max} (*e.g.*, Terblanche *et al.* 2007; Woods *et al.* 2018), with longer duration heat stress assays—as in the constant temperature lab exposures from which our CT_{max} estimates are derived—typically yielding lower estimated CT_{max} than short duration assays (Bates & Morley 2020; Peck *et al.* 2009; Woods *et al.* 2018). However, thermal knockdown assays measure a single trait—the ability for a mosquito to stay standing and right itself at a given heat exposure—with an unknown linkage to key demographic, life history, and transmission-related traits; and these assays often measure time duration rather than temperature as their output. Moreover, these assays have not been conducted systematically across mosquito species. We instead chose to use CT_{max} values from constant temperature experiments because these capture performance that is meaningfully linked to mosquito population dynamics and distributions, but which may be lower than values derived from short-term knockdown experiments. For a given species, these empirical estimates often come from a lab colony representing a single, potentially lab-adapted population. Thus, although mosquito thermal tolerance can vary between populations, we applied the available thermal tolerance estimate to the species across its range.

Table 1. Species-specific trait values used in biophysical models of mosquito body temperature. For each species, the thermal limit values (CT_{max}, CT_{min}, and T_{opt}) are for adult survival. The following abbreviations are used below: MDR for mosquito immature development rate, YFV for yellow fever virus, RRV for Ross river virus, WNV for West Nile fever, MVEV for Murray Valley encephalitis virus, JEV for Japanese encephalitis virus, SLEV for St. Louis encephalitis virus, and WEEV for Western equine encephalitis virus.

Species	Human pathogens vectored	CT_{min} (°C)	T_{opt} (°C)	CT_{max} (°C)	Activity pattern (diurnal, nocturnal, crepuscular)
Aedes aegypti	Dengue virus, chikungunya virus, Zika virus, YFV	10.4	23.9	37.5	Diurnal, crepuscular
Ae. albopictus	Dengue virus, chikungunya virus, Zika virus, YFV	13.5	22.5	31.4	Diurnal, crepuscular
An. gambiae	malaria, *Wucheria bancrofti* (cause of lymphatic filariasis)	8.5	22.3	38.9	Nocturnal, crepuscular
Anopheles stephensi	malaria	7.0	21.5	39.1	Nocturnal, crepuscular
Culex annulirostris	RRV, MVEV, Barmah forest virus, Kunjin virus, JEV	13.1	23.4	33.6	Nocturnal, crepuscular
Cx. pipiens	WNV	4.6	10.5	35.3	Nocturnal, crepuscular

Cx. quinquefasciatus	WNV, *Wucheria bancrofti*, SLEV, WEEV	4.8	11.9	34.6	Nocturnal, crepuscular
Cx. tarsalis	WNV, SLEV, WEEV	4.6	10.3	38.1	Nocturnal, crepuscular

Mosquito occurrence data

To determine the distribution over which to estimate thermal vulnerability, we used published and/or publicly available occurrence records (*i.e.*, a collection with an accompanying latitude and longitude) from the Global Biodiversity Information Facility (GBIF; see Supplemental Table S2 for full GBIF doi) for each species (Supplemental Figure S2). For species with hundreds of occurrence records (*i.e., Ae. aegypti*), we used a spatially stratified sampling approach to select 20-30 records from each 10° latitudinal band on which to estimate thermal safety margins. That is, within each 10° latitudinal band, we grouped records by longitude and randomly sampled from within each group to minimize overrepresentation in specific regions (*e.g.,* cities or highly visited natural areas). While our approach was not designed to comprehensively cover the entire range of a given species, in cases of clear data gaps in GBIF (*e.g., Anopheles stephensi* in Kenya), we further supplemented these occurrence records with those from published data sources through targeted literature searches. We used the associated latitude and longitude to classify occurrence records into one of 14 biomes (Olson *et al.* 2001) and as 'tropical' (0-23.5°), 'subtropical' (23.5-35.0°), and 'temperate' (35.0-66.5°).

Estimating mosquito body temperatures

For each vector species and collection location, we estimated mosquito body temperatures using NicheMapR: a suite of R programs for microclimate and biophysical modeling (Kearney & Porter 2020). In the first step, we used the microclimate model with hourly ERA5 weather data as input, to estimate microclimate conditions in both full shade and full sun (Kearney & Porter 2017). Specifically, the microclimate model input includes ERA5 data on wind speed, longwave radiation flux, solar radiation, surface pressure, 2 m temperature and 2 m dewpoint temperature, total precipitation, and cloud cover (Klinges *et al.* 2022). These data are available at a spatial resolution of 0.25° and are downscaled to an approximately 30 m resolution based on local terrain effects (*i.e.,* slope, aspect, elevation). The downscaled microclimate conditions are then used as the input in the biophysical model, which is used to estimate steady-state body temperatures. In addition to the microclimate conditions, the biophysical model utilizes information on organismal functional traits including body size, daily activity patterns (*e.g.*, diurnal, nocturnal, crepuscular), and thermal limits (Kearney & Porter 2020). We supplied species-specific parameter values for daily activity patterns and thermal limit estimates (*e.g.*, point estimates and 95% credible intervals for T_{opt}, CT_{max}, CT_{min} for adult survival; Table 1, Supplemental Table S1). In the biophysical model, T_{opt} is considered the organisms' preferred body temperature (*i.e.*, the temperature that it will attempt to maintain), while CT_{max} and CT_{min} impact the organisms' thermoregulatory behaviors (Kearney & Porter 2020). We assumed that

all mosquito species had a body size of approximately 3 mg with 85% solar absorptivity (Brust
1967). Using these parameter values, we estimated species' thermal safety margins with
behavioral thermoregulation (*i.e.*, the ability to move across fully exposed to fully shaded
microhabitats; main results). Results without behavioral thermoregulation are presented in
Supplemental Figure S3. These models have been extensively validated in a wide range of field
conditions (Kearney *et al.* 2014; Kearney & Maino 2018; Kearney & Porter 2017).
186
Incorporating impacts of drought on vector life cycles
In environments with highly seasonal precipitation, mosquito populations may reduce activity or
aestivate during drought (Adamou *et al.* 2011; Lehmann *et al.* 2010), and thus avoid exposure to
high temperature extremes experienced during that time. To incorporate potential seasonal and
aestivation responses driven by low moisture availability, we masked out any periods in which
the prior 30 or more days had soil moisture below 5%. That is, if the prior 30+ days each had soil
surface wetness <5%, then hourly thermal conditions from that point onward were excluded from
our thermal vulnerability calculations until soil moisture rose above 5% again. This soil moisture
metric roughly corresponded to precipitation events (Supplemental Figure S4). Although some
mosquito species may persist during drought by using artificial water sources (*e.g.*, *Aedes*
aegypti; Trpiš 1972), we used this drought mask for all species for consistency, as
comprehensive information on the drought responses of mosquito populations across their ranges
is not available. For all species, incorporating this drought mask had minimal impact on the
thermal safety estimates (Supplemental Tables S4-S6, Supplemental Figure S5).
201
Estimating thermal safety margins
To quantify mosquito vulnerability to climate warming, we calculated thermal safety margins:
the difference between an organism's critical thermal maximum (CT_{max}) and the warmest
temperature the organism experiences (T_e) in the coolest microhabitat available (Deutsch *et al.*
2008; also referred to as 'warming tolerance'). We interpret regions and species with low
thermal safety margins as having higher thermal vulnerability, but do not attempt to draw
specific cut-offs in what constitutes vulnerability based on this metric. By using highly resolved
microclimate data (*i.e.,* spatial resolution of approximately 30 m), and estimating T_e both with
and without behavioral thermoregulation, we believe the thermal safety margins calculated here
provide a relevant metric of thermal vulnerability. To capture both the magnitude and duration of
thermal safety, we specifically calculated three related indices: 1) thermal safety margins, 2) the
longest continuous period (in hours) that T_e exceeds CT_{max} (*i.e.*, 'thermal danger'), and 3) the
longest streak of consecutive days in thermal danger (where a day is counted if $T_e > CT_{max}$ for at
least one hour). Together, these capture both average thermal safety and seasonal and diurnal
variation in thermal safety. After estimating these indices at each collection location for each
species, we used generalized additive models (GAM) to estimate latitudinal patterns of thermal
vulnerability (Supplemental Methods). We fit all models using a smoothed function of up to
eight knots estimated using restricted maximum likelihood with the 'mgcv' R package (Wood

2017). We included only occurrence records collected below 2,500 m of elevation to focus on
latitudinal variation on temperature and avoid confounding, regionally-varying effects of
elevation (which excluded 3 total occurrence records; Supplemental Table S2).
223
Identifying maxima and minima in thermal safety
To identify the latitudes with the highest and lowest thermal safety for each species, we searched
for global and local maxima and minima in thermal safety margins, respectively, in the GAM
fits. That is, we searched for latitudes in which the first derivative of the fitted function switched
from positive to negative, and in which the estimated thermal safety margins were the highest (or
lowest) of their neighboring latitudes (*i.e.*, highest or lowest in a sequence of three). We further
confirmed the locations of maximum and minimum thermal safety margins by visually
inspecting the plotted GAM fits for each species. To estimate uncertainty, we drew 1,000
samples from the posterior-fitted GAM parameter vector, multiplied by the linear predictors (*i.e.*,
the sum of the product of the predictors and their associated regression coefficients), and
detected maxima and minima in each sampled fit, as in Pinsky *et al.* (2019) (Wood 2017;
Supplemental Figure S6).
236
237   **Results**
238
Patterns of thermal safety across species and options for avoidance
We estimated that all 8 vector species currently have portions of their distributions in thermal
safety and portions in thermal danger (*i.e.,* positive and negative thermal safety margins,
respectively) (Figure 1). Only *Ae. aegypti, An. gambiae*, and *Cx. tarsalis* were thermally safe
across the majority of their ranges, with average thermal safety margins of approximately 1°C,
(Supplemental Table S4). The remaining species—*Ae. albopictus, An. stephensi, Cx.*
annulirostris, Cx. pipiens, and *Cx. quinquefasciatus*—were in thermal danger for most of their
distributions with average thermal safety margins ranging from -0.3 to -3.6°C. In particular, *Ae.*
albopictus, An. stephensi, and Cx. annulirostris were estimated to be in thermal danger by
several degrees Celsius across nearly their entire distributions (Figure 1, Supplemental Table
S4).
250   These results above pertain to estimates under the main model specification, in which
options for avoiding thermal extremes—namely, behavioral thermoregulation and reduced
activity or aestivation during drought—were included. Thermal safety margin estimates were
highly similar when excluding aestivation during drought, but were, on average, 0.13 ± 0.19°C
lower under this specification (Supplemental Figure S5, Supplemental Table S4). However,
when the ability to behaviorally thermoregulate was removed (*i.e.*, when vectors were restricted
to fully exposed habitats), thermal safety margins were substantially lower (3.15 ± 0.34°C lower)
(Supplemental Table S4), and most species were in thermal danger across the majority of their
ranges (Supplemental Figure S3), highlighting the importance of access to thermal refugia during
high temperature extremes. However, we note that our thermal safety margin estimates use

CT$_{max}$ estimated under constant temperatures, and that short-term exposure to higher
temperatures may be tolerable, potentially with a physiological cost (as described in the
Methods: Mosquito thermal tolerance data and Discussion). As expected, thermal safety margin
estimates were substantially higher when using CT$_{max}$ estimates from the most thermally tolerant
life history trait for a given mosquito species (*e.g.*, immature survival, biting rate), rather than
adult survival for consistency across species (Supplemental Figure S1, Supplemental Table S1).
Under this specification, all vector species were thermally safe across the majority of their
distributions.
268

Figure 1. Thermal safety margins across latitude for individual mosquito vectors. Points are
estimates at individual occurrence records for a given species; solid and dotted lines show the
mean and 95% confidence intervals from the GAM fits. These are estimated with behavioral
thermoregulation and with the drought mask, including only observations from below 2,500 m
elevation (see Supplemental Figures S3 and S5 for corresponding plots under alternative
assumptions).

Latitudinal patterns of vector thermal safety margins

For vector species with distributions spanning both hemispheres, thermal safety margins typically peaked just south of the equator and at the northern temperate range edges (Figure 2, Supplemental Figures S7-S9). For these species, thermal safety margins were lowest between 23 – 35° (N or S)—the latitudinal extent of the subtropics—with the exception of *An. gambiae,* for which thermal safety was lowest in the tropics (~21°S and 18°N). For the two species with distributions restricted to the northern hemisphere (*An. stephensi, Cx. tarsalis*), thermal safety margins were either highest around the edges of their distribution (~22°N and 44°N) and lowest around 30-35°N (*Cx. tarsalis*) or peaked at their tropical range edge (~2°N) and decreased monotonically into the sub-tropics (*An. stephensi*). For all species, these patterns of thermal safety were mostly driven by variation in temperature experienced in fully shaded habitats across latitude (Supplemental Figure S10). That is, the highest hourly mosquito body temperatures were typically estimated for species occurrences in the subtropics, with relatively cooler maximum body temperatures around the equator and northern temperate range edges. These patterns held true when excluding the drought mask (Supplemental Figure S5, Supplemental Table S4).

Figure 2. Latitude of maxima (blue points) and minima (red points) in thermal safety margins across the latitudinal ranges (gray rectangles) analyzed for each species. Points and lines denote the mean and 95% confidence interval, respectively, for each species. For *An. stephensi,* no peaks or valleys were identified as the estimated thermal safety margins decrease monotonically across its range. See 'Methods: *Identifying maxima and minima in thermal safety'* for details on how locations were estimated. See Supplemental Table S3 and Supplemental Figure S6 for estimates and uncertainties.

Biogeographical patterns of vector thermal safety margins

In addition to thermal safety margins typically being lowest in the subtropics, thermal safety margins varied biogeographically (Figure 3). In particular, regions dominated by deserts and xeric shrublands, including in the Middle East, the western Sahara, and southeastern Australia.

had the lowest thermal safety margins across vector species (Figure 3; Supplemental Figure S11)

(Olson et al. 2001). Further, the longest consecutive periods of thermal danger (*i.e.*, when

estimated mosq⟨...⟩ssed

further below) t⟨...⟩

Figure 3. Vector thermal safety margins by grid cell (top) and biome (bottom). For each vector

species, thermal safety margins were scaled from 0 to 1 before being combined (by taking the

average of each grid cell) as shown here. In the top panel, grid cells are 4°x4°. In the bottom

panel, biomes refer to the 14 biomes classified by the World Wildlife Fund. The biome numbers

listed on the x-axis correspond to 1) Tropical & Subtropical Moist Broadleaf Forests, 2) Tropical

& Subtropical Dry Broadleaf Forests, 4) Temperate Broadleaf & Mixed Forests, 5) Temperate

Conifer Forests, 7) Tropical & Subtropical Grasslands, Savannas & Shrublands, 8) Temperate

Grasslands, Savannas & Shrublands, 9) Flooded Grasslands & Savannas, 12) Mediterranean

Forests, Woodlands & Scrub, 13) Deserts & Xeric Shrublands. Lines within each violin plot

denote the median relative TSM for that biome. Points are scaled estimates of TSMs for

individual occurrence records within each biome (jittered to improve visibility). The number of
occurrence records assigned to each biome is noted above each violin plot. See Supplemental
Figures S7-S9 for scaled TSMs for each individual vector species.
324
Time scales of thermal danger
The length of time spent in thermal danger (*i.e.*, when estimated mosquito body temperatures
exceed species' critical thermal maxima), varied widely between species and regions (Figure 4,
Supplemental Figure S12, Supplemental Tables S5). In particular, the longest continuous period
in thermal danger throughout the year ranged from an average of 1.9 ± 3.4 hours (*Cx. tarsalis*) to
8.0 ± 6.7 hours (*Ae. albopictus*) (mean ± s.d. across occurrence points; Supplemental Figure S12,
Supplemental Table S5). Further, the longest period in thermal danger was 8.1 ± 9.4 hours in the
subtropics compared to 3.2 ± 4.4 and 4.1 ± 4.5 hours (mean ± s.d. across occurrence records for
all species) in tropical and temperate regions, respectively, matching patterns observed in
thermal safety margins. Without the ability to behaviorally thermoregulate, the longest period in
thermal danger increased by several hours from an average of 4.3 ± 2.1 across all species and
regions to 6.1 ± 2.0 (Supplemental Table S5).
These patterns were highly similar when considering the number of consecutive days in
which thermal danger occurred for at least one hour. As above, the longest streak of days in
thermal danger was lowest for *Cx. tarsalis* (2.1 ± 6.9 days) and highest for *Ae. albopictus* (18.8 ±
27.1 days), and lower in tropical and temperate regions (8.2 ± 18.8, 3.1 ± 5.3 days, respectively)
compared to subtropical regions (11.3 ± 17.5 days) (Figure 4, Supplemental Figure S12,
Supplemental Table S6). These periods were also substantially longer when the ability to
behaviorally thermoregulate was removed, increasing by roughly two- to three-fold for all
species. Similarly, when the drought mask was removed, the streaks of thermal danger increased
by approximately 2 days (to 9.59 ± 6.07 days) across all species and had a particularly large
impact on *An. stephensi* (10.8 ± 15.8 to 17.5 ± 30.6 days) and *Cx. pipiens* (4.1 ± 11.2 to 10.1 ±
30.0 days) (Supplemental Table S6).

Figure 4. Longest consecutive periods in thermal danger by region for all vector species
combined. Y-axes show the longest streak of consecutive hours (left) or days (right) in thermal
danger (*i.e.*, when body temperatures exceed species thermal maxima). The horizontal line
within each boxplot denotes the median for that region. See Supplemental Figure S12 and Table
S5-S6 for species-specific values of these metrics, and Supplemental Figure S13 for patterns of
thermal danger by biome).

Discussion
We investigated the risk of heat stress posed by climate warming for 8 major mosquito vector
species by estimating thermal safety margins—here, the difference between a species' critical
thermal maximum and the hottest hourly body temperature it experiences across its geographic
range. We estimated that, when able to access shaded microhabitats, several of the most major
vector species, namely *Ae. aegypti, An. gambiae*, and *Cx. tarsalis*, were thermally safe across the
majority of their distributions, suggesting low risk from warming (Figure 1). However, when
limited to exposed habitats, thermal safety margins for all species were approximately 3°C
lower, placing most species in thermal danger (*i.e.,* body temperature would exceed critical
thermal maxima) across large portions of their range (Supplemental Figure S3). Our results thus
suggest that behavioral thermoregulation is likely already a pervasive strategy for buffering
mosquitoes from high temperature extremes. The reliance on behavior and access to cooler
microhabitats to avoid thermal danger has been well documented in other ectotherm species
(Kearney *et al.* 2009b; Pinsky *et al.* 2019; Sunday *et al.* 2014). For mosquitoes, behavioral
avoidance of high temperatures in laboratory settings has been observed in *Aedes, Anopheles*,
and *Culex* species (Blanford *et al.* 2009; Thomson 1938; Verhulst *et al.* 2020), and preference for
cooler, shaded oviposition sites in field settings has been documented for *Aedes* (Barrera *et al.*
2006; Vezzani & Albicócco 2009) and *Culex* (Vezzani & Albicócco 2009) species. However, the
extent and potential fitness costs of mosquito behavioral thermoregulation remain largely
unknown. Our results highlight the importance of understanding mosquito behavioral
thermoregulation and of incorporating microhabitat availability when investigating mosquito
responses to warming, as behavior could altered projected impacts of future warming (Huey &
Kingsolver 2019).
We find that buffering against climate warming for most vector species was highest
around the equator and at their northern temperate range edges, and lowest around the subtropics
(*i.e.*, 23-35° N or S) (Figures 1-3). While this finding contradicts the expectation that tropical
species are most vulnerable to climate warming from seminal studies such as Deutsch et al. 2008
and Tewksbury et al. 2008, this same pattern was found in a recent meta-analysis including over
400 ectotherm species (Pinsky *et al.* 2019). Similarly, studies of ectotherm range shifts in
response to warming, reflecting evidence of climate vulnerability, have found the fastest range
shifts at higher latitudes (Ramalho *et al.* 2023), as well as more nuanced responses including
east-west and equator-ward shifts (Lenoir & Svenning 2015; Pinsky *et al.* 2013; VanDerWal *et*
al. 2013). Our findings thus contribute to a growing body of evidence that species' risks and

responses to climate warming do not vary unidirectionally with latitude, and current risks may be
highest in subtropical, rather than tropical, regions because of their higher thermal extremes
(Johansson *et al.* 2020; Kingsolver *et al.* 2013; Pinsky *et al.* 2019; Ramalho *et al.* 2023).
Our finding that climate vulnerability is greater in the subtropics is driven by the high
short-term temperature extremes experienced there. Across species, we estimated that the highest
mosquito body temperatures experienced for one or more hours occurred around 29°N and 38°S
(Supplemental Figure S10). This is consistent with expectations from climatology that, although
average daily mean temperatures typically peak at the equator and decrease monotonically
towards the poles, average daily maximum temperatures typically peak around the subtropics,
and are relatively lower at the equator and higher-latitude temperate regions (Buckley & Huey
2016a; Hoffmann 2010; Kingsolver & Buckley 2017; Supplemental Figure S14). These short-
term thermal extremes, even if rare, are known to cause major declines in individual fitness,
scaling up to drive population and species-level impacts on demographic rates in other ectotherm
taxa (Buckley & Huey 2016a, b; Ma *et al.* 2015). Our findings thus suggest that subtropical
mosquito populations may be under the greatest pressure to shift their ranges, seasonality, and/or
adapt to warming in coming decades, with relatively higher stability in tropical and temperate
populations. However, this will depend on the relative fitness costs of exposure to high short-
term thermal extremes versus high mean temperatures, which is not well understood and
constitutes a key future research direction (Bates & Morley 2020)
While we describe patterns of mosquito climate vulnerability across broad spatial scales
(*i.e.,* between tropical, subtropical, and temperate regions) and latitude, regional-scale
biogeographical differences clearly drive variation in vulnerability (Figure 3). We found that
thermal safety margins were lowest in specific regions including the Middle East, the western
Sahara, and southeastern Australia, which are largely classified as desert and xeric shrubland
biomes (Olson *et al.* 2001), but are highly ecologically and climatologically distinct from each
other. As described for the subtropics broadly, these regions each experience the highest short-
term thermal extremes, although not necessarily the warmest annual mean temperatures
(Supplemental Figure S14). Further, while we assumed all locations contained the full range of
microhabitat conditions (*i.e.,* full sun to full shade), the availability of thermal refugia, such as
logs, tree holes, and epiphytes, varies widely by region (Scheffers *et al.* 2014), and is likely
lower in desert and xeric shrubland biomes, further compounding the higher climate
vulnerability there. Humidity, which is important for mosquito survival and may modulate
temperature sensitivity, may also limit tolerance of hot temperature extremes, especially in xeric
environments (Brown *et al.* 2023).
As climate vulnerability depends not only on the magnitude of temperature extremes, but
also on their duration and frequency, we estimated the longest continuous periods that mosquito
species spent in thermal danger. As before, we found that the longest stretches of thermal danger,
in terms of both consecutive hours and days, occurred in subtropical regions (Figure 4).
Comparing across species, we found that thermal danger, when it occurred, typically lasted no
more than 2 to 8 hours (for *Cx. tarsalis* and *Ae. aegypti* to *Ae. albopictus*, respectively,

Supplemental Table S5, Supplemental Figure S12). At these time scales, rapid hardening or heat
shock responses, which have been well documented in other ectotherm species, could
substantially increase short-term critical thermal maxima (Kellermann *et al.* 2017; King &
MacRae 2015; Ma *et al.* 2021). As heat shock responses are highly conserved across ectotherm
species (Lindquist & Craig 1988), this may be a pervasive strategy for mosquitoes mitigating
thermal danger under current and future climatic extremes. However, the precise time scales over
which heat shock responses operate, and their overall fitness costs, remain poorly understood. In
other ectotherm species, the production of heat shock proteins has been associated with
decreased critical thermal maxima on subsequent days (Bai *et al.* 2019) and overall decreases in
development and reproduction (Feder & Hofmann 1999; McMillan *et al.* 2005; Sørensen *et al.*
2003). Further, heat injury may be additive, such that recurring exposure to thermal stress may
further reduce individual fitness and population growth, particularly if there is insufficient time
(*e.g.,* <6 h) for injury repair (Jørgensen *et al.* 2021; Kashmerry & Bowler 1977). We estimated
that certain vector species – namely *Ae. albopictus*, *Ae. stephensi*, and *Cx. annulirostis* – may
experience nearly 10 or more consecutive days of thermal danger in portions of their distribution,
which would likely exceed the capacity for heat injury repair and result in reduced fitness
(Supplemental Table S6, Supplemental Figure S12). However, other species such as *Ae. aegypti*
and *Cx. tarsalis* may experience few consecutive days (*e.g.*, <3) of thermal stress, potentially
enabling recovery and mitigating impacts on fitness. Understanding the impacts of heat stress on
mosquitoes at these time scales and frequencies will improve estimates of mosquito climate
vulnerability and is an important direction for future research.
The time dimension of heat stress is also important for accurately estimating species
critical thermal maxima (Bates & Morley 2020). In this study, we used estimates of critical
thermal maxima derived from laboratory experiments, which measured adult mosquito survival
under constant temperatures, typically occurring for several hours, days, or weeks (Supplemental
Table S1; summarized in Mordecai et al. 2019 and Villena et al. 2022). Prior work has
demonstrated that the duration of heat exposure and the rate of temperature change used in heat
tolerance assays, as well as the thermal history of the organism itself, including cross-
generational effects, can impact the estimated critical thermal maxima (Heerwaarden &
Kellermann 2020; Kingsolver *et al.* 2011, 2015; Schiffer *et al.* 2013; Terblanche *et al.* 2007;
Waite & Sorte 2022). As a result, there is no single definitive critical thermal maximum for a
given taxon or individual, but instead a continuum depending on temporal dynamics of heat
exposure and the specific trait of interest (Bates & Morley 2020; Clusella-Trullas *et al.* 2021;
Hoffmann *et al.* 2013; Jørgensen *et al.* 2019; Kellermann *et al.* 2012; Lutterschmidt & Hutchison
1997). In general, critical thermal maxima estimated from longer duration heat stress assays, as
used here, are lower than those estimated from acute heat exposure assays such as thermal
knockdowns (Bates & Morley 2020; Peck *et al.* 2009; Woods *et al.* 2018). Thus, we may have
been biased towards calculating smaller thermal safety margins, and thereby overestimating the
extent of thermal danger and mosquito climate vulnerability. We used estimates of critical
thermal maxima from thermal performance curves derived from constant-temperature laboratory

experiments because they are widely available across the focal taxa, they are consistent in their
temporal dimension, and they have relevance to fitness in the field (Mordecai *et al.* 2019, 2020;
Shocket *et al.* 2018, 2020; Tesla *et al.* 2018). Further, we used the estimate of critical thermal
maximum from adult survival for consistency across species, and because this trait is likely the
most accurately and precisely estimated, and likely has largest impact on overall fitness.
However, if positive population growth requires all life history traits to be within their thermal
limits, we could be underestimating thermal danger. Other measures that could be more locally
relevant (*e.g.*, responses to short-term heat exposures following more realistic daily temperature
variation) have not been conducted systematically across taxa and would likely not be broadly
representative and comparable across species entire ranges, as we have examined here.
While we sought to provide an ecologically realistic estimate of mosquito risk to climate
warming, there are several additional limitations in our risk metric. First, mosquito populations
may suffer declines before thermal safety falls to zero, given the typically steep drop-off in
thermal performance between a species' thermal optima and maxima (Mordecai *et al.* 2019), and
potential trade-offs in mosquito life history traits. Further, although mosquito thermal tolerance
can vary between populations (*e.g.*, Couper *et al.* 2023; Reisen 1995; Ruybal *et al.* 2016), we
applied a single prior estimate of mosquito critical thermal maxima to each species across its
range, as population-specific estimates are not yet available. Additionally, seasonality in
mosquito life cycles and prolonged periods of dormancy may buffer populations from thermal
extremes (Lehmann *et al.* 2010). As mosquito seasonality and dormancy is largely driven by
precipitation and moisture conditions (Huestis & Lehmann 2014; Lehmann *et al.* 2010; Shocket
et al. 2021), we attempted to account for this by applying a drought mask, in which periods
following 30 or more days of dry surface soils were excluded from our thermal safety margin
estimates (this drought mask had minimal impact on our findings). Incorporating regionally-
specific, population-level mosquito responses would enable more accurate estimates of climate
vulnerability, but these data are not available for most species and locations. In addition to these
limitations, the ultimate impacts of climate warming on mosquito species will depend not only
on current thermal vulnerability, but on rates of warming, which are expected to be spatially
heterogeneous but generally greatest around the poles (Clem *et al.* 2020), as well as changes in
other abiotic and biotic drivers (*e.g.*, drought, humidity, urbanization, vector control measures;
Franklinos *et al.* 2019). For example, urbanization or land clearing could compound the impacts
of climate warming by eliminating thermal refugia provided by canopy cover (Alkama &
Cescatti 2016). Ongoing work on mosquito population responses to temperature in the face of
concurrent global changes will help to further refine estimates of warming-associated risk.
Despite these complexities, our findings are largely in agreement with patterns of
ectotherm range shifts in response to warming, suggesting that our metric provides meaningful
information on mosquito climate vulnerability. Our results suggest that the subtropics may be
most likely to experience shifts in the seasonality and/or intensity of mosquito-borne disease
transmission due to climate change in coming decades, potentially necessitating shifts in the
types and timing of vector control strategies for effective disease prevention. Further, the

tropics—which currently experience the highest mosquito-borne disease burden—could remain
relatively favorable for transmission under near-term climate warming, highlighting the need for
sustained vector control in this region.
512
513 **Conflict of Interest**
514 The authors declare that they have no conflicts of interest.
515
516 **Data accessibility**
517 All data and code used in this project are publicly available on GitHub:
518 https://github.com/lcouper/VectorThermalSafety/
519

References

Adamou, A., Dao, A., Timbine, S., Kassogué, Y., Yaro, A., Diallo, M., *et al.* (2011). The contribution of aestivating mosquitoes to the persistence of Anopheles gambiae in the Sahel. *Malar J*, 10, 151.

Alkama, R. & Cescatti, A. (2016). Biophysical climate impacts of recent changes in global forest cover. *Science*, 351, 600–604.

Angilletta, M.J. (2009). *Thermal Adaptation: A Theoretical and Empirical Synthesis*. OUP Oxford.

Bai, C.-M., Ma, G., Cai, W.-Z. & Ma, C.-S. (2019). Independent and combined effects of daytime heat stress and night-time recovery determine thermal performance. *Biology Open*, 8, bio038141.

Barrera, R., Amador, M. & Clark, G.G. (2006). Ecological Factors Influencing Aedes aegypti (Diptera: Culicidae) Productivity in Artificial Containers in Salinas, Puerto Rico. *JOURNAL OF MEDICAL ENTOMOLOGY*, 43.

Bates, A.E. & Morley, S.A. (2020). Interpreting empirical estimates of experimentally derived physiological and biological thermal limits in ectotherms. *Can. J. Zool.*, 98, 237–244.

Blanford, S., Read, A.F. & Thomas, M.B. (2009). Thermal behaviour of Anopheles stephensi in response to infection with malaria and fungal entomopathogens. *Malaria Journal*, 8.

Brown, J.J., Pascual, M., Wimberly, M.C., Johnson, L.R. & Murdock, C.C. (2023). Humidity – The overlooked variable in the thermal biology of mosquito-borne disease. *Ecology Letters*, 26, 1029–1049.

Brust, R.A. (1967). Weight and development time of different stadia of mosquitoes reared at various constant temperatures. *Can Entomol*, 99, 986–993.

Buckley, L.B. & Huey, R.B. (2016a). How Extreme Temperatures Impact Organisms and the Evolution of their Thermal Tolerance. *Integr. Comp. Biol.*, 56, 98–109.

Buckley, L.B. & Huey, R.B. (2016b). Temperature extremes: geographic patterns, recent changes, and implications for organismal vulnerabilities. *Global Change Biology*, 22, 3829–3842.

Chen, S.-C., Liao, C.-M., Chio, C.-P., Chou, H.-H., You, S.-H. & Cheng, Y.-H. (2010). Lagged temperature effect with mosquito transmission potential explains dengue variability in southern Taiwan: Insights from a statistical analysis. *Science of The Total Environment*, 408, 4069–4075.

Clem, K.R., Fogt, R.L., Turner, J., Lintner, B.R., Marshall, G.J., Miller, J.R., *et al.* (2020). Record warming at the South Pole during the past three decades. *Nat. Clim. Chang.*, 10, 762–770.

Clusella-Trullas, S., Garcia, R.A., Terblanche, J.S. & Hoffmann, A.A. (2021). How useful are thermal vulnerability indices? *Trends in Ecology & Evolution*, 36, 1000–1010.

Couper, L.I., Farner, J.E., Lyberger, K.P., Lee, A.S. & Mordecai, E.A. (2024). Mosquito thermal tolerance is remarkably constrained across a large climatic range. *Proceedings of the Royal Society B: Biological Sciences*, 291.

Deutsch, C.A., Tewksbury, J.J., Huey, R.B., Sheldon, K.S., Ghalambor, C.K., Haak, D.C., *et al.* (2008). Impacts of climate warming on terrestrial ectotherms across latitude. *Proceedings of the National Academy of Sciences*, 105, 6668–6672.

Diamond, S.E., Frame, A.M., Martin, R.A. & Buckley, L.B. (2011). Species' traits predict phenological responses to climate change in butterflies. *Ecology*, 92, 1005–1012.

Dowd, W.W., King, F.A. & Denny, M.W. (2015). Thermal variation, thermal extremes and the physiological performance of individuals. *Journal of Experimental Biology*, 218, 1956–1967.

Elsen, P.R., Saxon, E.C., Simmons, B.A., Ward, M., Williams, B.A., Grantham, H.S., *et al.* (2022). Accelerated shifts in terrestrial life zones under rapid climate change. *Global Change Biology*, 28, 918–935.

Feder, M.E. & Hofmann, G.E. (1999). HEAT-SHOCK PROTEINS, MOLECULAR CHAPERONES, AND THE STRESS RESPONSE: Evolutionary and Ecological Physiology. *Annu. Rev. Physiol.*, 61, 243–282.

Franklinos, L.H.V., Jones, K.E., Redding, D.W. & Abubakar, I. (2019). The effect of global change on mosquito-borne disease. *The Lancet Infectious Diseases*, 19, e302–e312.

Grimm, N.B., Chapin III, F.S., Bierwagen, B., Gonzalez, P., Groffman, P.M., Luo, Y., *et al.* (2013). The impacts of climate change on ecosystem structure and function. *Frontiers in Ecology and the Environment*, 11, 474–482.

Harrigan, R.J., Thomassen, H.A., Buermann, W. & Smith, T.B. (2014). A continental risk assessment of West Nile virus under climate change. *Global Change Biology*, 20, 2417–2425.

Heerwaarden, B. van & Kellermann, V. (2020). Does Plasticity Trade Off With Basal Heat Tolerance? *Trends in Ecology & Evolution*, 35, 874–885.

Hersbach, H., Bell, B., Berrisford, P., Hirahara, S., Horányi, A., Muñoz-Sabater, J., *et al.* (2020). The ERA5 global reanalysis. *Quarterly Journal of the Royal Meteorological Society*, 146, 1999–2049.

Hoffmann, A.A. (2010). Physiological climatic limits in Drosophila: patterns and implications. *Journal of Experimental Biology*, 213, 870–880.

Hoffmann, A.A., Chown, S.L. & Clusella-Trullas, S. (2013). Upper thermal limits in terrestrial ectotherms: how constrained are they? *Functional Ecology*, 27, 934–949.

Huestis, D.L. & Lehmann, T. (2014). Ecophysiology of Anopheles gambiae s.l.: Persistence in the Sahel. *Infection, Genetics and Evolution*, 28, 648–661.

Huey, R.B. & Berrigan, D. (2001). Temperature, Demography, and Ectotherm Fitness. *The American Naturalist*, 158, 204–210.

Huey, R.B. & Kingsolver, J.G. (2019). Climate Warming, Resource Availability, and the Metabolic Meltdown of Ectotherms. *The American Naturalist*, 194, E140–E150.

Johansson, F., Orizaola, G. & Nilsson-Örtman, V. (2020). Temperate insects with narrow seasonal activity periods can be as vulnerable to climate change as tropical insect species. *Sci Rep*, 10, 8822.

Jørgensen, L.B., Malte, H., Ørsted, M., Klahn, N.A. & Overgaard, J. (2021). A unifying model to estimate thermal tolerance limits in ectotherms across static, dynamic and fluctuating exposures to thermal stress. *Sci Rep*, 11, 12840.

Jørgensen, L.B., Malte, H. & Overgaard, J. (2019). How to assess Drosophila heat tolerance: Unifying static and dynamic tolerance assays to predict heat distribution limits. *Functional Ecology*, 33, 629–642.

Kashmerry, A.M.S. & Bowler, K. (1977). A study of recovery from heat injury in the blowfly (*Calliphora erythrocephala*) using split-dose experiments. *Journal of Thermal Biology*, 2, 183–184.

Kearney, M., Porter, W.P., Williams, C., Ritchie, S. & Hoffmann, A.A. (2009a). Integrating biophysical models and evolutionary theory to predict climatic impacts on species'

ranges: the dengue mosquito Aedes aegypti in Australia. *Functional Ecology*, 23, 528–538.

Kearney, M., Shine, R. & Porter, W.P. (2009b). The potential for behavioral thermoregulation to buffer "cold-blooded" animals against climate warming. *Proc. Natl. Acad. Sci. U.S.A.*, 106, 3835–3840.

Kearney, M.R., Isaac, A.P. & Porter, W.P. (2014). microclim: Global estimates of hourly microclimate based on long-term monthly climate averages. *Sci Data*, 1, 140006.

Kearney, M.R. & Maino, J.L. (2018). Can next-generation soil data products improve soil moisture modelling at the continental scale? An assessment using a new microclimate package for the R programming environment. *Journal of Hydrology*, 561, 662–673.

Kearney, M.R. & Porter, W.P. (2017). NicheMapR – an R package for biophysical modelling: the microclimate model. *Ecography*, 40, 664–674.

Kearney, M.R. & Porter, W.P. (2020). NicheMapR – an R package for biophysical modelling: the ectotherm and Dynamic Energy Budget models. *Ecography*, 43, 85–96.

Kellermann, V., Overgaard, J., Hoffmann, A.A., Flojgaard, C., Svenning, J.-C. & Loeschcke, V. (2012). Upper thermal limits of Drosophila are linked to species distributions and strongly constrained phylogenetically. *Proceedings of the National Academy of Sciences*, 109, 16228–16233.

Kellermann, V., Van Heerwaarden, B. & Sgrò, C.M. (2017). How important is thermal history? Evidence for lasting effects of developmental temperature on upper thermal limits in *Drosophila melanogaster*. *Proc. R. Soc. B.*, 284, 20170447.

King, A.M. & MacRae, T.H. (2015). Insect Heat Shock Proteins During Stress and Diapause. *Annual Review of Entomology*, 60, 59–75.

Kingsolver, J.G., Arthur Woods, H., Buckley, L.B., Potter, K.A., MacLean, H.J. & Higgins, J.K. (2011). Complex Life Cycles and the Responses of Insects to Climate Change. *Integrative and Comparative Biology*, 51, 719–732.

Kingsolver, J.G. & Buckley, L.B. (2017). Evolution of plasticity and adaptive responses to climate change along climate gradients. *Proceedings of the Royal Society B: Biological Sciences*, 284, 20170386.

Kingsolver, J.G., Diamond, S.E. & Buckley, L.B. (2013). Heat stress and the fitness consequences of climate change for terrestrial ectotherms. *Functional Ecology*, 27, 1415–1423.

Kingsolver, J.G., Higgins, J.K. & Augustine, K.E. (2015). Fluctuating temperatures and ectotherm growth: distinguishing non-linear and time-dependent effects. *Journal of Experimental Biology*, 218, 2218–2225.

Klinges, D.H., Duffy, J.P., Kearney, M.R. & Maclean, I.M.D. (2022). mcera5: Driving microclimate models with ERA5 global gridded climate data. *Methods Ecol Evol*, 13, 1402–1411.

Kraemer, M.U.G., Reiner, R.C., Brady, O.J., Messina, J.P., Gilbert, M., Pigott, D.M., *et al.* (2019). Past and future spread of the arbovirus vectors Aedes aegypti and Aedes albopictus. *Nat Microbiol*, 4, 854–863.

Lehmann, T., Dao, A., Yaro, A.S., Adamou, A., Kassogue, Y., Diallo, M., *et al.* (2010). Aestivation of the African Malaria Mosquito, Anopheles gambiae in the Sahel. *Am J Trop Med Hyg*, 83, 601–606.

Lenoir, J. & Svenning, J.-C. (2015). Climate-related range shifts – a global multidimensional synthesis and new research directions. *Ecography*, 38, 15–28.

Lindquist, S. & Craig, E.A. (1988). THE HEAT-SHOCK PROTEINS. *Annual Review of Genetics*, 22, 631–677.

Lutterschmidt, W.I. & Hutchison, V.H. (1997). The critical thermal maximum: history and critique. *Can. J. Zool.*, 75, 1561–1574.

Ma, C.-S., Ma, G. & Pincebourde, S. (2021). Survive a Warming Climate: Insect Responses to Extreme High Temperatures. *Annu. Rev. Entomol.*, 66, 163–184.

Ma, G., Hoffmann, A.A. & Ma, C.-S. (2015). Daily temperature extremes play an important role in predicting thermal effects. *Journal of Experimental Biology*, jeb.122127.

McMillan, D.M., Fearnley, S.L., Rank, N.E. & Dahlhoff, E.P. (2005). Natural temperature variation affects larval survival, development and Hsp70 expression in a leaf beetle. *Functional Ecology*, 19, 844–852.

Medlock, J.M. & Leach, S.A. (2015). Effect of climate change on vector-borne disease risk in the UK. *The Lancet Infectious Diseases*, 15, 721–730.

Mordecai, E.A., Caldwell, J.M., Grossman, M.K., Lippi, C.A., Johnson, L.R., Neira, M., *et al.* (2019). Thermal biology of mosquito-borne disease. *Ecology Letters*, 22, 1690–1708.

Mordecai, E.A., Ryan, S.J., Caldwell, J.M., Shah, M.M. & LaBeaud, A.D. (2020). Climate change could shift disease burden from malaria to arboviruses in Africa. *The Lancet Planetary Health*, 4, e416–e423.

Olson, D.M., Dinerstein, E., Wikramanayake, E.D., Burgess, N.D., Powell, G.V.N., Underwood, E.C., *et al.* (2001). Terrestrial Ecoregions of the World: A New Map of Life on Earth. *BioScience*, 51, 933.

Parmesan, C. (2006). Ecological and Evolutionary Responses to Recent Climate Change. *Annual Review of Ecology, Evolution, and Systematics*, 37, 637–669.

Parmesan, C. & Yohe, G. (2003). A globally coherent fingerprint of climate change impacts across natural systems. *Nature*, 421, 37–42.

Peck, L.S., Clark, M.S., Morley, S.A., Massey, A. & Rossetti, H. (2009). Animal temperature limits and ecological relevance: effects of size, activity and rates of change. *Functional Ecology*, 23, 248–256.

Pinsky, M.L., Eikeset, A.M., McCauley, D.J., Payne, J.L. & Sunday, J.M. (2019). Greater vulnerability to warming of marine versus terrestrial ectotherms. *Nature*, 569, 108–111.

Pinsky, M.L., Worm, B., Fogarty, M.J., Sarmiento, J.L. & Levin, S.A. (2013). Marine Taxa Track Local Climate Velocities. *Science*, 341, 1239–1242.

Pörtner, H.O., Bennett, A.F., Bozinovic, F., Clarke, A., Lardies, M.A., Lucassen, M., *et al.* (2006). Trade-Offs in Thermal Adaptation: The Need for a Molecular to Ecological Integration. *Physiological and Biochemical Zoology*, 79, 295–313.

Ramalho, Q., Vale, M.M., Manes, S., Diniz, P., Malecha, A. & Prevedello, J.A. (2023). Evidence of stronger range shift response to ongoing climate change by ectotherms and high-latitude species. *Biological Conservation*, 279, 109911.

Reisen, W. (1995). Effect of Temperature on Culex tarsalis (Diptera: Culicidae) from the Coachella and San Joaquin Valleys of California. *Journal of Medical Entomology*, 32, 636–645.

Rezende, E.L., Castañeda, L.E. & Santos, M. (2014). Tolerance landscapes in thermal ecology. *Functional Ecology*, 28, 799–809.

Ruybal, J.E., Kramer, L.D. & Kilpatrick, A.M. (2016). Geographic variation in the response of Culex pipiens life history traits to temperature. *Parasites & Vectors*, 9, 116.

Ryan, S.J., Carlson, C.J., Mordecai, E.A. & Johnson, L.R. (2019). Global expansion and redistribution of Aedes-borne virus transmission risk with climate change. *PLOS Neglected Tropical Diseases*, 13, e0007213.

Ryan, S.J., Carlson, C.J., Tesla, B., Bonds, M.H., Ngonghala, C.N., Mordecai, E.A., *et al.* (2021). Warming temperatures could expose more than 1.3 billion new people to Zika virus risk by 2050. *Global Change Biology*, 27, 84–93.

Scheffers, B.R., Evans, T.A., Williams, S.E. & Edwards, D.P. (2014). Microhabitats in the tropics buffer temperature in a globally coherent manner. *Biology Letters*, 10, 20140819.

Schiffer, M., Hangartner, S. & Hoffmann, A.A. (2013). Assessing the relative importance of environmental effects, carry-over effects and species differences in thermal stress resistance: a comparison of Drosophilids across field and laboratory generations. *Journal of Experimental Biology*, 216, 3790–3798.

Shocket, M.S., Anderson, C.B., Caldwell, J.M., Childs, M.L., Couper, L.I., Han, S., *et al.* (2021). Environmental drivers of vector-borne disease. In: *Population Biology of Vector-borne Diseases*. Oxford University Press, pp. 85–118.

Shocket, M.S., Ryan, S.J. & Mordecai, E.A. (2018). Temperature explains broad patterns of Ross River virus transmission. *eLife*, 7, e37762.

Shocket, M.S., Verwillow, A.B., Numazu, M.G., Slamani, H., Cohen, J.M., El Moustaid, F., *et al.* (2020). Transmission of West Nile and five other temperate mosquito-borne viruses peaks at temperatures between 23°C and 26°C. *eLife*, 9, e58511.

Sørensen, J.G., Kristensen, T.N. & Loeschcke, V. (2003). The evolutionary and ecological role of heat shock proteins. *Ecology Letters*, 6, 1025–1037.

Sunday, J., Bennett, J.M., Calosi, P., Clusella-Trullas, S., Gravel, S., Hargreaves, A.L., *et al.* (2019). Thermal tolerance patterns across latitude and elevation. *Phil. Trans. R. Soc. B*, 374, 20190036.

Sunday, J.M., Bates, A.E., Kearney, M.R., Colwell, R.K., Dulvy, N.K., Longino, J.T., *et al.* (2014). Thermal-safety margins and the necessity of thermoregulatory behavior across latitude and elevation. *Proc. Natl. Acad. Sci. U.S.A.*, 111, 5610–5615.

Terblanche, J.S., Deere, J.A., Clusella-Trullas, S., Janion, C. & Chown, S.L. (2007). Critical thermal limits depend on methodological context. *Proceedings of the Royal Society B: Biological Sciences*, 274, 2935–2943.

Tesla, B., Demakovsky, L.R., Mordecai, E.A., Ryan, S.J., Bonds, M.H., Ngonghala, C.N., *et al.* (2018). Temperature drives Zika virus transmission: evidence from empirical and mathematical models. *Proceedings of the Royal Society B: Biological Sciences*.

Thomson, R.C.M. (1938). The Reactions of Mosquitoes to Temperature and Humidity. *Bulletin of Entomological Research*, 29, 125–140.

Trpiš, M. (1972). Dry season survival of Aedes aegypti eggs in various breeding sites in the Dar es Salaam area, Tanzania. *Bulletin of the World Health Organization*, 47, 433–437.

Urban, M.C. (2015). Accelerating extinction risk from climate change. *Science*, 348, 571–573.

VanDerWal, J., Murphy, H.T., Kutt, A.S., Perkins, G.C., Bateman, B.L., Perry, J.J., *et al.* (2013). Focus on poleward shifts in species' distribution underestimates the fingerprint of climate change. *Nature Clim Change*, 3, 239–243.

Vasseur, D.A., DeLong, J.P., Gilbert, B., Greig, H.S., Harley, C.D.G., McCann, K.S., *et al.* (2014). Increased temperature variation poses a greater risk to species than climate warming. *Proceedings of the Royal Society B: Biological Sciences*, 281, 20132612.

Verhulst, N.O., Brendle, A., Blanckenhorn, W.U. & Mathis, A. (2020). Thermal preferences of
subtropical Aedes aegypti and temperate Ae. japonicus mosquitoes. *Journal of Thermal*
Biology, 91, 102637.

Vezzani, D. & Albicócco, A.P. (2009). The effect of shade on the container index and pupal
productivity of the mosquitoes Aedes aegypti and Culex pipiens breeding in artificial
containers. *Medical and Veterinary Entomology*, 23, 78–84.

Villena, O.C., Ryan, S.J., Murdock, C.C. & Johnson, L.R. (2022). Temperature impacts the
environmental suitability for malaria transmission by Anopheles gambiae and Anopheles
stephensi. *Ecology*, 103, e3685.

Vinagre, C., Dias, M., Cereja, R., Abreu-Afonso, F., Flores, A.A.V. & Mendonça, V. (2019).
Upper thermal limits and warming safety margins of coastal marine species – Indicator
baseline for future reference. *Ecological Indicators*, 102, 644–649.

Waite, H.R. & Sorte, C.J.B. (2022). Negative carry-over effects on larval thermal tolerances
across a natural thermal gradient. *Ecology*, 103, e03565.

Wood, S. (2017). *Generalized Additive Models*. 2nd edn. Boca Raton.

Woods, H.A., Dillon, M.E. & Pincebourde, S. (2015). The roles of microclimatic diversity and of
behavior in mediating the responses of ectotherms to climate change. *Journal of Thermal*
Biology, What sets the limit? How thermal limits, performance and preference in
ectotherms are influenced by water or energy balance, 54, 86–97.

Woods, H.A., Kingsolver, J.G., Fey, S.B. & Vasseur, D.A. (2018). Uncertainty in geographical
estimates of performance and fitness. *Methods in Ecology and Evolution*, 9, 1996–2008.