Dysfunction from loss of essential outer membrane functions causes drug hypersensitization in

Acinetobacter baumannii overexpressing multidrug efflux pumps

Efrat Hamami[1], Wenwen Huo[1], Katherine Neal[3], Isabelle Neisewander[1], Edward Geisinger[1,2], Ralph R. Isberg[1,*]

[1]Department of Molecular Biology and Microbiology, Tufts University School of Medicine, 150 Harrison Ave., Boston, MA 02111, USA.

[2]Department of Biology, Northeastern University, Boston, MA, 02115, USA

[3]Department of Biochemistry, Curry College, Milton, MA, USA.

[*]Corresponding Author

Abstract

Elevated expression of Resistance-Nodulation-Cell Division (RND) drug transporters is commonly observed in clinical isolates of *Acinetobacter baumannii*, a nosocomial pathogen associated with multidrug-resistant (MDR) infections. We describe here a CRISPRi platform directed toward identifying hypomorphic mutations that preferentially changed resistance levels to the fluoroquinolone antibiotic ciprofloxacin in RND pump overproducers. An sgRNA library including single and double nucleotide mutations directed against essential genes of *A. baumannii* was constructed and introduced into multiple strain backgrounds, allowing strain-specific, titratable knockdown efficiencies to be analyzed. Other than NusG depletions, there were few candidates in the absence of drug treatment that showed lowered fitness specifically in strains overexpressing the RND efflux pumps AdeAB, AdeIJK, or AdeFGH. In the presence of ciprofloxacin, the hypomorphs causing hypersensitivity were predicted to result in outer membrane dysfunction, with the AdeFGH overproducer appearing particularly sensitive to disruptions. Increased ciprofloxacin susceptibility in two pump hyper-expressers was observed after depletions of a monovalent cation-proton antiporter protein and a nucleoid-associated protein, the latter of which was shown to promote pump hyper-activation. On the other hand, depletions of translation-associated proteins as well as components of the proton-pumping ATP synthase resulted in fitness benefits in the presence of the drug for at least two pump-overproducing strains. Therefore, pump overproduction exacerbated stress caused by defective outer membrane integrity, while the efficacy of drug resistance in efflux pump overproducers was enhanced by slowed translation or defects in the proton-pumping ATP synthase.

Introduction

Acinetobacter baumannii is a Gram-negative nosocomial pathogen that causes ventilator-associated pneumonia, bacteremia, and urinary tract infections (1-3). The difficulty in combating diseases caused by this often multidrug-resistant (MDR) organism has stimulated the demand for new antimicrobials and improved therapeutic strategies (1, 3). Among the various determinants of resistance to currently available drugs, hyper-expression of drug efflux pumps is commonly associated with MDR across multiple species (4-9). The Resistance-Nodulation-Division (RND) class of efflux pumps provides the most clinically relevant strategy for extruding antibiotics in *A. baumannii,* with the AdeABC, AdeIJK, and AdeFGH efflux pumps frequently associated with MDR (6-9). Therefore, strategies that target isolates overexpressing these pumps or which hypersensitize these strains to currently available antibiotics are of great interest.

For Gram-negative organisms, the primary route to fluoroquinolone resistance is *via* acquisition of mutations in the active sites of the major type II topoisomerases in the cell, DNA gyrase and topoisomerase IV (10). In sequenced *A. baumannii* clinical strains, resistance mutations in both *gyrA* and *parC* are found, encoding the A subunits of gyrase and topo IV, respectively (10). Based on MIC testing and sequencing of isolates, clinical strains also commonly have a third mutation in one of the regulatory genes that activate an RND antibiotic efflux pump. Mutations in these genes, encoded by *adeS*, *adeN*, or *adeL* upregulate the AdeAB(C), AdeIJK, or AdeFGH efflux pumps respectively, and result in high fluoroquinolone resistance (FQR) (7, 11-15).

Previous work identified mutations that caused strain-specific vulnerabilities of pump overproduction in nonessential *A. baumannii* genes. These mutations hindered growth of strains overexpressing efflux pumps or synergized with the fluoroquinolone ciprofloxacin (CIP) to

hypersensitize the overproducers to the drug (16). Based on that work, lesions in nonessential

genes predicted to alter envelope integrity sensitize efflux pump overproducing strains to CIP

treatment. Furthermore, the strain with the activating mutation in *adeL* (causing *adeFGH*

overexpression) appeared to result in the most significant physiological consequences, with a

broader mutation spectrum found to reduce fitness, and a more robust transcriptional shift was

apparent relative to the other efflux pump overproducers. These effects may be due to either

special features of this pump or extreme overexpression of the *adeFGH* locus relative to other

pump-overexpressing strains (16).

Most effective antimicrobials interfere with proteins essential for growth, making them

desirable targets for identifying mutations that preferentially sensitize strains overproducing

efflux pumps. To approach this problem, we used CRISPRi to target essential genes with

sgRNAs. The library design takes advantage of a previously developed strategy that generates

multiple hypomorphic alleles with varying fitness defects in each essential gene, enabling the

discovery of depletions that show strain-specific aggravation of drug sensitivity (17-19).

Depletions that cause drug hypersensitivity were found to predominantly target proteins involved

in maintaining outer membrane (OM) integrity. Interestingly, interference of translation and the

assembly of the proton pumping F_1F_0 ATPase provided the FQR strains with a fitness advantage

in the presence of drug. Other recent studies have also used CRISPRi to identify mutations in *A.*

baumannii that have altered sensitivities to a large panel of antibiotics (20, 21).

Results

    Development of a CRISPRi strategy to uncover hypomorphs with reduced fitness in FQR strains. In our previous study (16), we identified transposon insertion mutations that reduce the fitness of efflux pump overproducers or that specifically hypersensitize these variants to ciprofloxacin (CIP). Except for very rare exceptions (22), insertion mutations could only be analyzed in nonessential genes. This interfered with our ability to identify proteins encoded by the essential gene set that could similarly sensitize pump overproducers. To overcome this shortcoming, we devised a CRISPRi screening strategy to identify depletion variants with decreased fitness in the presence of drug or fitness defects in a subset of strains, but still allowed viability of the bacterium in the absence of drug, as previously described (Fig. 1) (17, 18). We applied a pool design similar to that used for eukaryotic cells (17). Using sequencing data of a previously generated dense transposon bank in the ATCC 17978UN strain background (16, 22, 23), the insertion landscape was inspected to identify essential genes based on open reading frames (ORFs) having significantly reduced or no insertions (Materials and Methods (24, 25)). After comparison to other previously published *A. baumannii* studies, a list of 406 essential genes was generated (16, 23, 26, 27).

    A single 24-nucleotide sgRNA was designed for each essential gene by identifying the PAM site on the noncoding strand most proximal to the translational start. For each perfect sgRNA, we devised a set of 23 mismatch sgRNAs that differed from the original guide sequence by 1 or 2 nucleotides at varying locations across the sequence (17). We also created five shuffle sgRNAs comprising the same nucleotides in a disordered fashion to use as negative controls, anticipating that they would have a neutral effect on fitness. An additional collection of approximately 200 shuffle sgRNAs was created with minimal anticipated homology to the

genome as well, called pool shuffles. The final library consisted of approximately 12,000

sgRNAs that had a size and complexity similar to a previously constructed model pool (Fig. 1)

(17). To facilitate cloning, the pYDE007 gRNA delivery vector was converted into a

GoldenGate-compatible plasmid to allow efficient construction of the sgRNA library via

insertion of duplex target oligonucleotides from the pool into the site generated by a Type II

restriction enzyme (see Materials and Methods). Control of CRISPRi-dCas9 targeted depletions

was provided by an anhydrotetracycline inducible dCas9 introduced into the genomes of the WT

and FQR pump overexpressing strains (using a Tn7 strategy originally devised by Peters *et al*.

(28) and modified by Bai *et al*. (23); Materials and Methods). The FQR strain panel had the

gyrAparC background with either the *adeS*, *adeN,* or *adeL* pump overproducing mutations (GPS,

GPN, and GPL respectively). These strains were then transformed in parallel with the CRISPRi

sgRNA-pDNA bank, ensuring at least 10^6 colonies were pooled for each pool.

Mismatched sgRNAs result in a gradient of fitness defects. Pools harbored in all strain

backgrounds were cultured in LB broth for 7-8 generations in the absence or presence of CIP or

anhydrotetracycline (aTc) to induce dCas9 transcription (Materials and Methods; Fig. 1), with

cultures harvested before (timepoint T1) and after bank outgrowth (timepoint T2). Plasmid DNA

was extracted at both timepoints and the sgRNA region amplified from the extracted plasmid

DNA, barcoded, and prepared for sequencing (Materials and Methods). Reads were processed to

determine the abundance of each unique sgRNA at both time points in all conditions. The

sgRNA reads showed a normal distribution (Supp. Fig. 1A) that was maintained after

introduction into the WT *A. baumannii* strain (Supp. Fig. 1B). A similar distribution was

observed from the shuffle sgRNAs after aTc induction (Supp. Fig. 1C), consistent with

nontargeting behavior as expected for negative controls. Fitness was calculated after

stoichiometric depletion of target genes across replicate cultures as previously described

(Materials and Methods (10, 22, 29), with fitness values normalized to a small set of shuffle

sgRNAs that had no statistically significant effect on fitness after dCas9 induction. As

anticipated, most shuffle guides showed no change in fitness upon dCas9 induction (Fig. 2A),

whereas the majority of sgRNAs with perfect target homology showed diminished fitness upon

dCas9 induction (73% of perfect guides reducing fitness based on FDR <0.05, fitness < 0.96

after induction compared to control, with fitness at least 10% lower than uninduced cultures; Fig.

2B). Improved fitness was observed by guides harboring mismatched nucleotides closest to the

3' PAM site critical for dCas9 targeting (34% of these guides with mismatched nucleotides most

proximal to the PAM site reducing fitness, based on FDR <0.05 and fitness < 0.93 after

induction, with fitness lowered by at least 10% from uninduced cultures; Fig. 2C) (17, 18). This

pattern can be seen by focusing on sgRNA sets for individual genes such as *gyrA*, encoding

subunit A of DNA gyrase, which showed a broad range of fitness defects dependent on the

mismatch site (Fig. 2D, E).

The mild effects observed by the remaining 27% of perfect-match sgRNAs could be due

to a variety of factors, one of which may be the relative distance of the target sequence from the

translation start codon. A distinct relationship was difficult to establish except when comparing

fitness yielded by guides targeting sites 25-40 nucleotides away from the target codon start site

versus those recruited to sites only 1-10 nucleotides away (Supp. Fig. 2A-D). As many sgRNAs

with high fitness likely resulted from poor target depletion (ex. Supp. Fig. 2E), we determined if

continued growth for additional generations (14-16 generations vs. 7-8) could amplify fitness

defects. We found that continued growth expanded the number of perfect sgRNAs causing a

significant defect (a 35-38% boost in guides with fitness lowered by at least 0.5; Supp. Fig. 3)

without a substantial loss of fitness for the shuffle guides (Supp. Fig. 4). These data indicate that

the defects observed in hypomorphs can be aggravated by challenging the sgRNA banks for

longer periods of time, consistent with the target proteins being subjected to turnover and

dilution through bacterial division.

Identification of hypomorphs with significantly altered fitness in the FQR strain

backgrounds. To uncover factors that shape fitness of FQR pump hyper-expressing strains,

multiple replicates (n=10) of libraries in four strain backgrounds (WT, GPS, GPN, and GPL)

were cultured in rich LB broth in the presence of aTc inducer (for dCas9 expression), and fitness

scores of hypomorphs were compared between the drug susceptible WT and FQR strains. To

identify candidates important for maintaining FQR strain fitness under fluoroquinolone pressure,

the libraries were also grown in the presence of sub-MIC levels of CIP in parallel using

concentrations that resulted in 20-30% growth inhibition across the strains. The statistical

significance of each depletion (per unique sgRNA) was calculated as a function of the change in

fitness between conditions or the difference in fitness between strains, determined by multiple t-

test comparisons using an FDR or q value <5% (Materials and Methods) (30). Discoveries

showing a significant fitness shift between each comparison were sorted *in silico* to identify

sgRNAs yielding fitness scores that are at least 20% higher (knockdown imparting a fitness

benefit or alleviated effect) or at least 20% lower (knockdown causing a fitness defect or

aggravated effect). Among these, targets that were represented by 3 or more guides from their

sgRNA set were identified as top hits. Targets were identified which fulfilled these criteria

during CIP treatment (vs no treatment, or untreated (untx)) and preferentially in the FQR strain

(vs WT). Among these, top candidates consisted of targets that were identified in more than one

FQR strain background.

In the absence of antibiotics, there were few candidates with three or more guides that

passed this statistical bar, further supporting the Tn-seq results indicating that *A. baumannii* is

quite tolerant of pump overproduction (Supp. Fig. 5A-C) (16). This was particularly true of the

GPS strain, indicating that overproduction of the AdeAB pump is well tolerated by *A. baumannii*

(Supp. Fig. 5A vs. B and C). Expanding the criteria for fitness scores in the FQR strains that

differ from WT by at least 10% identified a total of 28 targets (originally 1) that reduced GPS

fitness relative to WT, including proteins involved in secretion across the inner membrane (*secY,*

secA, ffh), DNA replication, the electron transport chain, and the TCA cycle (Supp. Fig. 5D-E,

Supp. Fig. 6A). Doing so also uncovered 9 targets that diminished fitness of all three FQR strains

(Supp. Fig. 5D) and 29 in at least two FQR strains (Supp. Fig. 5E).

Unexpectedly, outgrowth for additional generations revealed a smaller set of fitness

determinants in GPS when compared to WT, only three of which overlap with those identified

after 7-8 generations of growth (n=5 here, Supp. Fig 6B vs. Supp. Fig. 6A). However, it is

notable that depletion of factors involved in competing processes resulted in opposing

phenotypes after 14-16 generations. For example, depletion of the transcription termination

factor Rho improved fitness of the GPS mutant relative to WT, while depletion of the anti-

terminator NusG showed stronger fitness defects in GPS (*nusG* and *rho*; Supp. Fig. 6B).

Importantly, when considering hypomorphs whose fitness varied by at least 20%, repression of

anti-transcription termination factor *nusG* was the single target identified as lowering fitness of

more than one pump hyper-expresser in the absence of drug pressure. In fact, knockdown of

nusG showed profound reductions in fitness in all three FQR strains when compared to WT

(Supp. Fig. 5A-D). In *E.coli*, NusG-dependent antitermination is tightly linked to regulatory

control of ribosomal RNA transcription (31, 32).

Depletion of either translation-associated or F_1F_0 ATPase assembly proteins

increases CIP tolerance of pump overproducers. Using criteria similar to those used to

identify strain-specific fitness differences in the absence of drug, we identified 8 targets with a

fitness advantage and 10 candidates with a fitness defect in pump overproducers after exposure

to CIP (Supp. Fig. 7). Analysis of pump overproducer and WT depletion derivatives (Figs 3A-F;

Supp. Fig. 7A) revealed a clear pattern: the GPL background (Fig. 3C, 3F) generated the most

depletions that selectively hypersensitized it to drug relative to WT, with few targets

differentiating the GPS and WT backgrounds (compare Fig. 3D to 3E, F). This is consistent with

a previous study analyzing mutations in nonessential genes (16). Furthermore, depletions leading

to dysfunction in cell envelope integrity were the most prominent candidates for CIP synergy,

supporting the previous Tn-seq study (16).

Depletion of *lpxC* and *lpxD* genes, encoding LOS biogenesis proteins, uniformly caused

fitness defects relative to WT in the presence of drug, even for the robust *adeS* mutant (Fig. 3,

Supp. Fig. 7B). With GPN and GPL, the spectrum of depletions that caused fitness defects was

expanded to include other loci involved in LOS biosynthesis (*kdtA/waaA*, *kdsD/kpsF*) as well as

a large cluster of genes involved in LOS transport to the surface (*lptA-C*, *lptG*) (Fig. 3, Supp. Fig.

7B). These data emphasize that OM integrity maintenance is required for FQR survival in the

presence of drug, with factors controlling the composition of LOS being particularly important.

Of note is the fact that depletions of BamD had weaker effects on the GPS pump overproducer

than the other two FQR strains, consistent with our observation that there were fewer

hypomorphs interfering with composition of the OM that reduced GPS fitness (Fig. 3, Supp. Fig.

7B). Notably, activation of the *adeS*-regulated system does not result in overexpression of its

outer membrane protein component, which would likely be linked to hypersensitizing defects in

OM biogenesis (33, 34).

Our most surprising result is that depletion of loci associated with ribosome biogenesis

and assembly, as well as translation, strongly increased fitness of at least two FQR strains

relative to WT, consistent with slower translational rates increasing fitness of efflux

overproducers (Fig. 3 vs. Supp. Fig 7A, Supp. Fig. 7C). Supporting this hypothesis are

depletions of *spoT* and *obgE* that similarly increased fitness of the GPL pump overproducer (see

Supp. Data 4). These are predicted to prevent degradation of ppGpp-associated alarmones,

resulting in translational stalling (35, 36). There was also a clear demonstration of increased

fitness exhibited by depletions of the F_1F_0 proton-pumping ATPase/synthase in the presence of

CIP, tying fitness to the overall proton gradient of the cell (Fig. 3, Supp. Fig. 7C). As the

function of the antibiotic efflux pumps is to antiport drugs in exchange for protons, decreased

F_1F_0 ATP synthase activity may remove an important competitor for protons or facilitate

cytoplasmic pH homeostasis.

High fitness FQR hypomorphs sensitized to CIP are largely impaired in envelope

integrity. We next focused on guides showing minimal fitness defects in the absence of

antibiotic treatment, allowing condition-specific effects to be detected throughout a large

dynamic range. This set of sgRNAs, called neutral or enhancer guides, were categorized as those

resulting in fitness scores that averaged around 1.00 (neutral) or 1.06 (enhancer) during dCas9

induction in the WT background. They demonstrated no substantial change in fitness between

growth in the presence or absence of inducer (fitness score differences within 0.05) (Fig. 4A). Of

284 targets that comprise the neutral and enhancer guide list (shuffles excluded and given

representation by at least 3 unique sgRNAs), 13 candidates were identified that showed

significantly lower fitness in the FQR background compared to WT in the presence of CIP (Fig.

4B, Supp. Data 4). These 13 candidates are predominantly proteins that shape outer membrane

integrity, with WaaA and Lpt proteins being top discoveries when restricting guides to those

with inefficient knockdown (Fig. 4B).

High dependence of GPL on the function of outer membrane biogenesis proteins to

maintain fitness in presence of CIP stress. Of the 13 FQR mutants that are sensitized to CIP,

WaaA, LolA, PhaF, IhfB, LptC, LptG, and LpxC were selected for experimental verification.

The sgRNAs were chosen based on their strong fitness defects during CIP exposure in GPL and

one other FQR strain (Fig. 5B-D vs Fig. 5F-H), as well as having minimal effects on WT fitness

in the absence of drug (Fig. 5A). We observed the strongest fitness deficiencies in the GPL strain

background (Fig. 5H), so we created individual sgRNA knockdowns of the top discoveries in

GPL and evaluated their growth alongside those in the WT strain background. Multiple sgRNAs

were tested per target (highlighted in red, blue, or orange spots, with their respective fitness

defects during CIP exposure shown in Fig. 6A). These hypomorphs were cultured in the presence

and absence of CIP and their growth rates compared to a strain harboring a shuffled non-

targeting sgRNA ("NC", NC_159), growing for the same time periods as performed in the screen

(Materials and Methods). The doubling times of hypormophs were then compared to those of the

control to detect altered growth rates caused by depletions (Fig. 6B, with the sgRNAs marked the

same way as in Fig. 6A for reference). Depletion of WaaA, LolA, PhaF, LptG, IhfB, LptC, and

LpxC all significantly slowed growth of GPL in comparison to WT during CIP exposure (Fig.

6B).

Notably, not all tested sgRNAs yielded the expected phenotype in the GPL background

(Supp. Fig. 8A), but this appeared to be linked to poor knockdown, as indicated by qRT-PCR

(Supp. Fig. 8B) or poor growth in the absence of the drug. For this reason, additional replicates

of LptC and LpxC hypomorphs were tested. Depletions of LpxC and LptC resulted in heightened

CIP sensitivity in GPL, with statistical significance only observed for LpxC (Supp. Fig. 9A).

Finally, the top 13 candidate hypomorphic guide RNAs were introduced into the GP, GPS, GPN,

and GPL strain backgrounds and then cultured in the presence or absence of CIP to determine

allele specificity. Again, the GPL strain background showed an appreciably heightened

sensitivity to CIP upon depletion of all targets of interest, but none significantly altered the

growth rate of GPS (Fig. 6C). On the other hand, GPN which showed more fitness dependencies

than GPS (Fig. 5G vs Fig. 5F; Fig. 5H) exhibited significantly stunted growth compared to GP

upon depletion of *waaA*, *ihfB*, and *lpxC* (Fig. 6D). The fact that there were fewer sensitivity loci

and much smaller defects than identified for GPL, is consistent with our previous analysis of

nonessential genes targeted by transposon mutagenesis. (16).

As depletions of LptC and LptG resulted in strain-specific fitness defects in the presence

of ciprofloxacin, we tested whether we could mimic genetic hypersensitization through the

addition of zosurabalpin (ZOS), a drug targeting the LptB$_2$FGC LPS transporter subcomplex

(37). The WT, GP, and GPL strains were grown in LB broth in the presence or absence of CIP

and/or ZOS, monitoring growth for 12 hours (Fig. 7A-C). ZOS in combination with CIP

dampened the growth of the GPL strain, with the effect stronger than that observed for either

drug alone (Fig. 7C). This aggravating effect was not observed in the WT strain or GP parent

strain (Fig. 7A, B), nor in the GPL strain lacking the AdeFGH efflux pump that is upregulated by

the *adeL* mutation (compare Fig. 7F to 7D, E). LOS-targeting drugs colistin (COL) and

polymyxin B (PB) exhibited even stronger effects: COL or PB in combination with CIP

diminished growth of GPL (Fig. 7I, L). This effect was not detected in GP (Fig. 7G, J) nor in

GPL lacking the AdeFGH pump (Fig. 7H). These results are consistent with the effect of Lpt,

WaaA, and LpxC depletions (Fig. 6B) and illustrates that identification of mutations that cause

aggravating effects can identify drug targets that allow aggravating effects on bacterial growth to

be identified.

It is anticipated that depletion of Lpt proteins, as well as other proteins involved in

shaping the OM, causes disruption of membrane integrity. To test for OM dysfunction,

hypomorphs were cultured in microtiter plates in the presence of vancomycin (VANC) or

sodium dodecyl sulfate (SDS), and their growth was monitored in the absence of any other drug

addition (Fig. 8A, B). As expected, depletion of WaaA, LpxC, LolA, LptC, and LptG yielded

increased sensitivity to the addition of either vancomycin or sodium dodecyl sulfate, consistent

with an impaired membrane barrier in these hypomorphs (Fig. 8A, B). To confirm that the

defects were the consequence of efflux pump overexpression, Δ*adeFGH* introduced into the GPL

was challenged in the presence or absence CIP (Fig. 8C). In the presence of an intact efflux

pump, the doubling time escalated for hypomorphs in the GPL background, with growth rates

significantly higher than those observed in the parent GP strain (Fig. 8C). Deletion of the

AdeFGH efflux pump reversed this effect, consistent with the vulnerability associated with the

adeL activation allele being attributed to pump overproduction (Fig. 8C).

The phenotype of *ihfB* hypomorphs did not fit the model of OM dysfunction, and given

its documented role in modulating transcription, it seemed likely that depletion of *ihfB* altered

expression of key resistance determinants. Furthermore, IhfB has been shown to control

expression of the *Salmonell* AcrAB pump activator gene *robA* (38-41). Therefore, transcript

levels of *adeG*, encoding the RND protein of the AdeFGH efflux pump, and the *adeL* regulator

were determined after IhfB depletion. Depletion of IhfB resulted in a steep decline in *adeG*

transcript levels and a corresponding increase in *adeL* expression (Fig. 8D). Therefore, reduction

of pump gene expression resulting from IhfB depletion is positively correlated with decrease of

GPL fitness in the presence of CIP.

In addition to drug sensitizers, we identified hypomorphs that conferred a fitness

advantage for efflux pump overexpressers in the presence of CIP. These included strains

depleted of inner membrane-associated F_1F_0 ATPase/synthase subunits (*atpI, atpF* in all FQR

strains; multiple other *atp* genes in GPL) (Fig. 3D-F). Some studies describe the bidirectionality

of the F_1F_0 ATPase/synthase complex, having the ability to both synthesize and hydrolyze ATP

using proton motive force (PMF) to drive ATP synthesis or reversibly hydrolyze ATP to pump

protons out of the cytosol. ATPase activity, however, was found to be limited and structurally

regulated in bacterial species like *A. baumannii*, *E. coli*, and *M. smegmatis*, albeit by different

means (42-45). Our results show that the protein plays an important role in maintaining pH

homeostasis, as examination of the intracellular pH (pHi) during AtpF depletion showed a

significant decrease in cytosolic pH (Fig. 8E, F), linking lowered ATPase activity to the failure

to remove poreons from the cytoplasm.

Discussion

Hyperexpression of RND drug efflux pumps often drives MDR, making them excellent

targets for antimicrobial therapy. As essential proteins provide particularly attractive targets for

the development of new antimicrobials, we depleted the expression of essential genes to identify

hypomorphs in strains harboring fluoroquinolone efflux pumps that are specifically sensitized to

the addition of ciprofloxacin (CIP). This approach showed that depletion of transcripts encoding

proteins involved in LOS synthesis and transport, lipoprotein maturation and trafficking to the

outer membrane all reduced the fitness of pump overproducers in the presence of antibiotic (CIP)

treatment. In particular, CIP sensitivity was specifically observed in the GPL (AdeFGH pump

hyperexpressed) strain, dependent upon the presence of the efflux pump.

BamD, which is required to support fitness of efflux pump overproducers in the presence

of CIP, is an outer membrane component of the Bam complex essential for the assembly and

proper insertion of outer membrane proteins (OMPs) (46). Interestingly, we previously

demonstrated that transposon knockouts of SurA, a non-essential periplasmic chaperone

involved in OMP assembly, also reduce the fitness of GPL strains in the presence of CIP (16, 46-

48). Depletions of BamD may interfere with the proper assembly or insertion of the pump outer

membrane factor (OMF) or other OMPs that maintain the outer membrane barrier (34).

Alternatively, a partially assembled Bam complex may be specifically disruptive to the cell

envelope, resulting in the combined defects of periplasmic accumulation of precursors and loss

of OMP content, leading to OM dysfunction. Perhaps directly related to this defect, targeting of

genes associated with lipoprotein maturation and trafficking (*lnt* and *lolA*) also showed

sensitization of efflux pump overproducers. BamD is one of multiple Bam lipoprotein

components, consistent with the proposition that depletions of *lnt* or *lolA* could directly

phenocopy defects in the Bam complex.

One noteworthy result was that depletions of both LOS synthesis (*waaA*, *kds*, and *lpx*

loci) as well as LOS transport (*lpt* loci) sensitized pump hyperexpressers to the presence of CIP.

This somewhat differs from two earlier reports regarding drug hypersensitivity, which showed

fitness defects specifically resulting from defects in LOS transport (21, 49). Decreased levels of

Lpt proteins produce a defective cell envelope that results in accumulation of material in the

periplasmic space (50, 51). Presumably, the accumulation of LOS precursors inserted in the inner

membrane in the absence of Lpt function causes a general sensitivity to most clinically effective

antibiotics (21). In contrast, focusing specifically on strains encoding efflux pumps with OM

components broadened the spectrum of sensitizing targets to LOS synthesis (*lpx*), and possibly

any factor that disrupts OM biogenesis. In fact, reduced fitness by a larger spectrum of defects

was predicted by our previous results, which showed that insertion mutations in *lpsB,* which

truncates LOS, also diminishes the fitness of pump-overexpressing strains during CIP treatment

(16, 22).

Targeting proteins that control OM composition and LOS dynamics may diminish the

integrity of the OM and facilitate drug entry (Fig. 8A, B), but this alone does not explain why

fitness is more severely compromised in the high-level FQR pump hyper-expressers than in the

WT or GP strains. One explanation is that there is direct binding of LOS to the RND transporter

complex, and the absence of this lipid results in destabilization of the outer membrane

component of the transporter. It is known that LPS can bind OMPs and that OMP-interfacing

lipids may mediate protein self-association, contributing to outer membrane integrity (52, 53).

By this model, proteins that shape the OM support assembly of the pump complex and thereby

resilience in the presence of drug. Overall, the targets revealed by this work indicate that the

FQR pump hyper-expressers are more vulnerable to changes in the outer membrane composition

than strains lacking these pumps. Components that support the integrity of the permeability

barrier, as well as stabilize the RND drug transporters, are likely to be important for survival

during drug treatment. Consistent with these results is the demonstration that RND pump

overexpressing strains are hypersensitive to Zosurabalpin, which disrupts the outer membrane

(49).

In addition to targets that are associated with OM biogenesis, a pair of loci were identified

that had no clear connection to supporting envelope integrity. IhfB is part of the heterodimeric

integration host factor complex that is nucleoid-associated and has the ability to locally and

globally modulate chromosome structure and influence gene expression (38-41). Depletion of

IhfB resulted in reduced expression of *adeG* accompanied by increased expression of the pump

regulator gene *adeL*, consistent with low pump concentrations leading to CIP sensitization.

Although it is unclear how a deficit in IhfB causes this defect, this result opens the door to

targeting global regulators as a strategy for combatting drug resistance. The other sensitization

locus not easily linked to OM dysfunction was PhaF, which is part of the monovalent

cation/proton antiporter PhaACDEFG complex (54-56). The antiporter may function to control

ion or pH homeostasis in the FQR strains to compensate for increased drug-proton antiport

activity of the RND efflux pumps. In its absence, there may be suboptimal functioning of the

AdeFGH pump leading to increased drug susceptibility.

We also identified that strains depleted of inner membrane-associated F_1F_0

ATPase/synthase subunits (*atpI, atpF* in all FQR strains; multiple other *atp* genes in GPL) and of

translation-associated factors (*fmt*, *rimM*, *der/engA*, *rnpA*) enhanced fitness of the pump hyper-

expressers in CIP. AtpF encodes the ATPase B subunit that connects the F_1 catalytic domain to

the F_0 membrane-spanning proton translocator, linking proton translocation to ATP generation

by F_1 (44, 45). AtpI, on the other hand, functions as a poorly conserved non-structural chaperone,

supporting the formation of the F_0 c-ring and complex stability (57, 58). Depletion of the specific

subset of components leading to drug tolerance may be a consequence of their limited lethality

after depletion as compared to other components.

There are two clear models for how depletion of F_1F_0 ATP synthase components could

provide a fitness advantage in the presence of the drug. First, lowering the cytosolic pH could

simply reduce the efficacy of the antibiotic (59, 60). Secondly, similar to the F_1F_0 ATP synthase,

RND pumps utilize the proton motive force generated by the electron transport chain, and

reducing ATPase synthase concentrations on the membrane may facilitate kinetic linkage of the

efflux pump to the respiratory chain. This model is consistent with our previous observation that

there were significantly reduced numbers of viable transposon insertion disruptions in the

cytochrome b_o (3) ubiquinol oxidase complex in the GPL strain (16). Therefore, the oxidase

complex could provide the protons necessary for efflux pump activity. In fact, depletion of the

ubiquinol oxidase complex as well as other systems involving the electron transport chain

reduced fitness of GPN and GPL even in the absence of drug in this study, indicating that the

oxidase may play an important homeostatic role even in the absence of known substrates (Supp.

Fig. 5E). Therefore, the ATP synthase and the ubiquinol oxidase appear to play opposing roles in

supporting fitness of pump overproducing strains.

Interestingly, in our RNA-seq experiments, we found that *atpE* transcription was

significantly downregulated in the GPL efflux overproducer when compared to the GP strain

grown in the absence of drug (16). Although depletion of this critical F_0 component was not

shown to protect from drug action, it does emphasize the similarity between drug stress and

stress related to overproduction of the AdeFGH pump. Therefore, manipulation of the proton

motive force may be a critical determinant in protecting against both pump overproduction and

The ATP synthase depletion phenotype after CIP treatment mirrors that of translation

apparatus component depletions, which also result in increased fitness. Perhaps allowing efflux

overproducers to enter an ATP-depleted state of dormancy protects against both pump

overproduction and drug stress, and this is mirrored by translational disruption. Both lowered

ATP and reduced translation have been associated with persister cell formation, allowing

bacteria to tolerate stress experienced during drug treatment (61, 62). Interestingly, pump

hyperexpression has also been documented to contribute to the development of persistence (63),

most notably in the case of the AdeFGH efflux pump (64). Therefore, either depressed ATP

synthesis or translation could support the fitness of pump overexpressing isolates, allowing

survival in the presence of high antibiotic levels, compensating for stress imparted by elevated

levels of transporter. This may be one factor in explaining why translation inhibitors have

occasionally been linked to antagonism of fluoroquinolone action (62, 65-68). Future efforts will

be directed to exploring these hypotheses in hopes of better understanding the hidden

susceptibilities of pump over-expression and identifying proteins that help support the viability

of the MDR state.

Acknowledgements

This work is supported by NIAID grants U01AI124302, U19AI158076, R21AI128328, and

U19AI142780. We thank Megan W. Tse (Broad Institute of Massachusetts Institute of

Technology and Harvard) for LptC and LptG hypomorphs, Defne Surujon (Boston College) for a

sam-to-wig file conversion script, Yunfei Dai (Northeastern University) for communications

regarding dCas9 strain generation, sgRNA design, and TRANSIT, and for the pYDE009 and

pYDE007 plasmids, Rebecca Batorsky (Tufts Technology Services) for aiding with TRANSIT

accessibility, and Albert Tai (Tufts Genomics Core Facility) for communications regarding

sequencing and library preparation.

Materials and Methods

Bacterial strains and growth conditions. Bacterial strains used in this work are listed in

Supplementary Data 1. All *Acinetobacter baumannii* strains are derived from ATCC 17978UN.

Cultures were grown in lysogeny broth (LB) (10 g/L tryptone, 5 g/L yeast extract, 10 g/L NaCl)

or on LB agar plates (LB supplemented with 15g/L agarose). Superoptimal broth with catabolite

repression (SOC) was used for recovery following all electroporations of *A. baumannii.*

Electroporations were performed using 0.1cm gap cuvettes and Bio Rad Gene Pulser, with the

settings 200 Ω, 25 µF, and 1.8 kV. For growth in broth culture, bacteria were aerated at 37°C on

a platform shaker, roller drum, or shaking in 96-well plates (Costar) with growth monitored by

measuring absorbance via a spectrophotometer at 600nm (A_{600}) with Genesys 30 (Thermo

Scientific) or Epoch 2 or SynergyMx (Biotek) plate readers. Cultures were first incubated to

early post-exponential phase in LB broth before diluted to a starting A_{600} of 0.003 in 5mL LB

broth or in 200µl LB (96-well plates) supplemented with aTc for dCas9 induction (when

hypomorphs are evaluated) and drug where applicable. Growth was monitored every 15 minutes

(plate reader) or evaluated after 170min (dCas9 induced but not drug treated) and 230min (dCas9

induced and drug treated). LB was supplemented with antibiotics, as needed, at the following

concentrations: Anhydrotetracycline (aTc): 100ng/mL, Carbenicillin (Carb): 50/100µg/mL,

Gentamicin (Gent): 10µg/mL, Chloramphenicol (Cm): 10/25µg/mL, Vancomycin (VANC): 64

µg/mL, sodium dodecyl sulfate (SDS): 0.05%, Ciprofloxacin (CIP): varying concentrations

(Sigma), or Zosurabalpin (ZOS): 2.5µm (MedChemExpress).

Molecular cloning and mutant construction. Primers (IDT or GENEWIZ) and

plasmids used throughout this study are listed in Supplementary Data 1. The dCas9 construction

is tetracycline (Tet)-inducible and chromosomally embedded at the Tn7 site in the four strains of

interest. The plasmids and techniques used to create these strains are as previously described (23,

28). Briefly, the mobile Tet-inducible dCas9 was introduced into the four *A. baumannii* recipient

strains of interest via four parental mating and Tn7 transposition. The recipient *A. baumannii*

strains, the HB101 RK600Cm^R helper, EGE187 DH5α λpir (pTNS3) encoding the T7

transposase, and the YDE009 DH5α (pUC18T-miniTn7T-GentR with *tetR-P$_{Tet}$::dCas9-rrnB*T1-

T7Te) donor strain were grown overnight in LB broth with appropriate antibiotics. The following

day the cultures were normalized to A$_{600}$ of 2.0 and 100µl of each of the three *E. coli* cultures

were combined with 100µl of each *A. baumannii* recipient culture in 600µl LB broth, prewarmed

to 37^{o}C. The mixture was washed twice in 37^{o}C LB broth, then resuspended in 25µl of fresh

broth and spotted on pre-warmed LB agar plates. These plates were incubated at 37^{o}C for

approximately 5 hours then resuspended in 1mL sterile phosphate buffered saline (PBS) before

plating and overnight selection on Vogel-Bonner minimal medium (VBM) containing 10µg/mL

of Gent to select against E. coli strains and the parental Gent^S *A. baumannii*. Exconjugants were

colony-purified on VBM containing 10µg/mL Gent, and screened on LB agar with 100µg/mL

Carb and on LB agar with 10µg/mL Gent. The insertion at the *att*Tn7 site downstream the *glmS*

gene was confirmed by colony PCR using the primers glmS2-F1 and Tn7R-R1.

The vector pYDE007 used to introduce the sgRNAs is a shuttle plasmid previously

described (23) containing the origin of replication of pBR322 and an origin of replication from a

natural *Acinetobacter* isolate pWH1277 (23) to allow for propagation in both *E. coli* and *A.*

baumannii respectively. It encodes for the β-lactamase gene (*bla*) on the backbone and contains a

sequence to be replaced by an sgRNA sequence upstream of the dCas9 handle and *S. pyogenes*

transcriptional terminator, with the fusion driven by the constitutive synthetic promoter pJ23119.

For efficient cloning of sgRNAs, the vector was made GoldenGate™-compatible by first

mutating the single BsaI Type II restriction site present in the *bla* gene and then introducing two

BsaI sites between the promoter and dCas9 handle to allow for directional sgRNA oligo

insertion. Using this backbone, all 24nt sgRNA oligos were constructed with additional 4nt

overhangs that allowed for directional cloning. The sgRNA bank was designed by loosely

following a previous strategy for stoichiometric knockdowns (17), and the vector and library

design were used for construction of the sgRNA-plasmid library by contract (GENEWIZ).

Identification of essential genes in *Acinetobacter baumannii*. Essential genes of *A.*

baumannii ATCC 17978UN were determined using transposon mutagenesis sequence data in the

WT strain (16, 22), and the data from that study was subjected to TRANSIT Gumbel analysis to

identify the essential genes (23-25). Briefly, sequence mapped SAM files were converted to wig

files with a custom python script bowtie_to_wig2.py provided by Defne Surujon from the Tim

van Opijnen lab. The wig files were then used as input, using the TRANSIT Gumbel analysis

pipeline (25) with default settings. Cross referencing our list of essential genes to those similarly

identified in other studies in the same strain or the more recently isolated AB5075 clinical strain

revealed 406 genes exhibiting a high likelihood of essentiality, given their classification as

essential in at least one of the other published analyses as well (23, 26, 27).

sgRNA library design. The 24nt sgRNAs were principally designed (to target each

essential gene) with the antisense strand sequence using a PAM site on the antisense strand that

is as close as possible to the translational start site. The sgRNA pool was then expanded with an

additional 23 mismatch sgRNAs that differ from each targeting sgRNA by 1 or 2 nucleotides,

along with 5 shuffle sgRNAs that comprise the same nucleotides as the original target-

homologous sgRNA but are shuffled to minimize the likelihood of off-target effects. An

additional 216 "pool shuffle" sgRNAs of an array of nucleotides with the GC content of the

targeted genes designed to minimize homology to any genomic sequence, bringing the final

sgRNA pool to 12,000 sgRNAs total (Supp. Data 2). Mismatches were employed to introduce

sgRNAs with reduced knockdown efficiencies, while the shuffles served as negative controls

(anticipated to provide a neutral effect on fitness). A BLAST search did reveal that up to 12bp

overlap with the genome was unavoidable among some members of the control pool. These

mismatches and shuffle sgRNAs were designed as previously described (17), using the python

script 'final sgRNA dirty pool design.ipynb' (GitHub).

Preparation of pools with sgRNA library. The sgRNA-pDNA library was introduced

into the FQR pump hyper-expressers and WT strain with chromosomally encoded Tet-inducible

dCas9 via electroporation. About 50µl electrocompetent cells were transformed with 23-46ng of

the pDNA, the resuspended mixture diluted 1/10, and 200µl plated on five large pre-warmed

SOC agar plates overlain with mixed cellulose ester filters (0.45µm pore size, 137mm diameter,

Advantec). The mixture was spread on the filters using beads. This was repeated for a total of 3

transformations per strain background. These transformants were allowed to recover at 37^{o}C for

an hour on SOC agar plates before the filters were transferred to selective LB agar plates

containing 50µg/mL Carb and incubated overnight at 37^{o}C for about 13.5-14 hours. The

following day, filters were pooled to acquire somewhere around 250,000-350,000 CFUs,

estimated based on using image analysis of previous pools (ImageJ). Pooling entailed washing

colonies off with 100mL sterile PBS with gentle shaking in a sterile round bottom 3L beaker.

Glycerol was added to a final concentration of 10-20% and 1mL aliquots were placed into

cryotubes for -80°C storage until needed.

Identification of hypomorphs with fitness defects. Following previous protocols for

determining fitness using Tn-Seq pools (10, 22), each vial of *A. baumannii* P_{Tet}::*dCas9*

harboring the sgRNA array was thawed on ice, diluted into 25mL LB broth to an A_{600}=0.1, and

revived by incubation at 37°C with shaking for 50min-60min until they reached around A_{600}=0.2.

Each culture was then diluted to an A_{600}=0.003 in 10mL LB broth in the absence or presence of

100ng/mL aTc for dCas9 induction and the fluoroquinolone ciprofloxacin. Cultures were

incubated for approximately 7-8 generations on the drum wheel at 37°C, for about 170 (no drug

treatment) or 230 (CIP treated) minutes, until densities reached A_{600}= 0.5-0.895. The cultures

were harvested both before (T1) and after (T2) dCas9 induction and drug treatment to allow

sgRNA fitness analyses. Sample pellets were stored at -20°C until all were collected. Plasmid

was extracted from each sample using the QIAprep Spin Miniprep Kit (Qiagen) following

manufacturer's protocol.

sgRNAs were amplified from harvested plasmid DNA using primers For2_sgRNA_Pool

and Rev_sgRNA_Pool at 0.6µM in a 50µl PCR reaction with 10µl 5x Q5 buffer, 0.5µl of Q5 hot

start polymerase, 1µl 100mM dNTPs (NEB), 100ng of the pDNA, and UltraPure water (Thermo

Fisher Scientific). The thermocycler settings were as follows with a heated lid: 98°C for 1min, 14

cycles of 98°C for 10sec, 70°C for 20sec, and 72°C for 1min, then finally 72°C for 2min.

Barcodes were then introduced with a second round PCR using a legacy Nextera kit of adaptors

(Illumina, compatible with FC-121-1031, FC-121-1030, FC-121-1012, FC-121-1011). The

second PCR was performed using 1µl of the first PCR product, 4µl UltraPure water, 7.5µl 2x

NEBNext High Fidelity Master Mix (NEB), and 1.25µl of each barcoded Nextera N5x and N7x

index primers per reaction. The thermocycler was set up to hold a heated lid while running at

72^{o}C for 3min, 98^{o}C for 2:45min, and 8 cycles of 98^{o}C for 15sec, 65^{o}C for 30sec, and 72^{o}C for

1:30min. The desired amplified product is 328nt in length. 5µl samples of the PCR2 products

were fractionated on a 2% agarose Tris-Acetate-EDTA (TAE) gel using Sybr Safe dye

(Invitrogen) alongside a set amount of 100bp DNA ladder (NEB) and run for 1hr at about 90V

for quantification. Samples were imaged and concentrations measured using a BioRad

quantitative gel doc system, with quantitation performed using Image J, or dedicated Bio-Rad

software so that approximately 100ng of each sample was pooled for sequencing. The samples

were PCR column purified (Qiagen) and their final concentrations determined via Nanodrop

prior to submission to Tufts University Core Facility of Genomics for sequencing. The pools

were sequenced via Illumina HiSeq 2500 SE100 using a legacy Nextera primer mixture.

The 5' sequence preceding the 24nt sgRNA was trimmed using Cutadapt (69). Reads

were assigned per sgRNA using Python script as described (18). To calculate fitness of the

hypomorphs, read counts for each sgRNA were first normalized to the total assigned reads per

sample. Any counts of 0 were given a read count of 1 for downstream fitness calculations.

Fitness yields from each sgRNA was determined as described (29, 70). Fitness values were

normalized to a set of 25 shuffle sgRNAs serving as negative/neutral controls by dividing each

individual sgRNA's fitness score by the average fitness of those 25 sgRNAs in each sample.

These shuffle sgRNAs were chosen as follows: The median read counts were identified for all

shuffle sgRNAs in each replicate sample tested (dataset was from a preliminary study

comprising of 10 replicate cultures of WT harboring the sgRNA library grown in rich media

without dCas9 induction, with dCas9 induction, and the same in the GPS strain background).

These median values were then expanded to a range of read counts (25-75[th] percentile). Shuffle

sgRNAs that fell in this range across all replicates and growth conditions in both the WT and

GPS strain backgrounds were identified. Among these sgRNAs, those yielding mean fitness

values closest to the median and which do so in at least 2 of the 4 conditions were selected for

downstream normalization. Normalized fitness values of each sgRNA were then compared

between dCas9 induced and uninduced conditions, between drug treated and untreated samples,

and between the FQR and WT strains. Multiple t-test analyses were employed to identify

significant differences in fitness due to growth condition or strain background. The change in

fitness generated by each sgRNA due to induction, due to drug treatment, and the fitness

difference of each hypomorph between strains was calculated. Discoveries of fitness scores that

significantly differ in these comparisons were determined by the two-stage step-up method of

Benjamini, Krieger, and Yekutieli with $Q < 5\%$ (30). Findings characterized as top discoveries

were targets represented by 3 or more sgRNAs. Those with <10 cumulative reads across

replicate cultures at T1 prior to outgrowth as well as ribosomal genes were excluded. Data

pertaining to the CIP screen are shown in Supplementary Data 4. All sequence data is also

deposited in the SRA database under BioProject number PRJNA1067098.

pHi Assay. This assay was performed as previously described (16). The AtpF

hypomorphs (and shuffle controls) evaluated in this work were grown to late exponential phase

before dilution to A_{600} of 0.01 in 8mL LB + 100ng/mL aTc. These cultures were incubated to

mid-exponential growth phase (A_{600}=0.510-0.574). Approximately 5mL were pelleted for

downstream pHi evaluation.

qRT-PCR. Cultures were grown in LB broth to early post-exponential phase before

dilution into 5mL LB with 100ng/mL aTc to a starting A_{600} of 0.003 (0.01 in 3mL for AtpF

depletion). Cultures were harvested after 170min reaching A_{600} of 0.134 to 0.904 (average

0.466). The following steps are as previously described (16): RNA was extracted using RNeasy

(Qiagen). RNA was treated with ezDNase before cDNA synthesis with the SuperScript VILO

cDNA kit (Thermo Fisher). Targets were amplified using PowerUp SYBR Green Master Mix

(Applied Biosystems) and CT values determined using the StepOnePlus Real-Time PCR system

(Applied Biosystems). The efficiency of primer pairs used for target amplification was evaluated

using a standard curve of cDNA at varying dilutions and consequently verified to be about 94-

108% for each pair used. Three biological replicates were evaluated for each strain or

hypomorph and three technical replicates examined per biological sample during target

amplification. To test for gDNA contamination, a set of control samples were tested in parallel

which were not reverse transcribed during the cDNA synthesis reactions. Transcript levels of

each specified target were compared between each hypomorph and the control strain harboring a

non-targeting shuffle sgRNA, NC_159, after normalization to levels of 16S.

Author Contributions

R.R.I and E.H conceptualized and devised the study. W.H and E.H designed the sgRNA library.

K.N, I.N, and E.H ran the preliminary screens with WT and GPS in the absence of drug. E.H

executed the CIP screen and hit validations. W.H tested depletion efficiency of select high fitness

perfect sgRNAs and provided bioinformatic support. E.G shared all strains, vectors, and

methodologies for CRISPRi set up. E.H and R.R.I interpreted the results and wrote the

manuscript.

Figure legends

Figure 1: Identification of hypomorphs with reduced fitness in FQR strains. Diagram for

experimental approach and sgRNA library design (Materials & Methods). sgRNAs were

designed to target 406 essential genes in *A. baumannii* ("perfect" guides). For each perfect guide,

a set of 23 guides with 1 to 2 mismatched nucleotides were created to achieve titratable

knockdown efficiencies as well as 5 sequence-shuffled guides as negative controls. Additional

pool shuffle sgRNAs were made to generate a library of 12,000 guides. Cultures were harvested

before (T1) and after (T2) outgrowth with or without CIP and the dCas9 inducer

anhydrotetracycline (aTc). Plasmid was extracted and prepared for sequencing to identify

hypomorphs with altered fitness under tested conditions. Hypomorph fitness scores, normalized

to scrambled guides, were compared between control and experimental groups to determine

essential gene targets that generate altered fitness of high-level FQR strains relative to WT in

either the presence or absence of CIP. FQR: fluoroquinolone resistant. CIP: ciprofloxacin. Untx:

untreated. Ind: dCas9 induced using anhydrotetracycline (aTc).

Figure 2: The sgRNAs with perfect sequence identity to essential genes lower fitness of *A.

***baumannii*.** Volcano plots are displayed to show statistical significance as a function of the mean

change in fitness scores that result from dCas9 induction (100ng/mL aTc, n=10). All symbols

refer to unique sgRNAs (light gray). The horizontal dotted line (A-D): FDR or Q=5%. (A)

Displayed are shuffle sgRNAs that serve as negative controls (black). (B) The sgRNAs with

perfect sequence identity to their target (black). (C) The sgRNAs with mismatched nucleotides

(nt) located most proximal to the 3' PAM site (black; nucleotides 23 or 24 of the 24nt sgRNA, -1

or 0 nt from PAM site respectively). (D, E): The sgRNAs designed for *gyrA* (black). S: Shuffle

sgRNAs. P: Perfect guide. Positions of mismatched nucleotides in relation to the 3' PAM site

noted, with numbering starting at 0 for the nucleotide immediately upstream the PAM site. (D)

Display of mean fitness difference of *gyrA* sgRNAs after induction vs. uninduced. (E) Display of

gyrA data, showing average fitness scores normalized to fitness in the absence of dCas9 inducer

for the perfect sgRNA and sgRNAs with single and double mismatches as well as shuffle

sgRNAs (± SEM).

Figure 3: Opposing effects of translation/ATP generation and LOS biogenesis on the fitness

of FQR strains relative to WT in the presence of CIP. (A-C) Statistical significance plotted as

a function of mean fitness score differences (n=10) between CIP and Untx conditions (top) or

FQR and WT strains during CIP treatment (bottom), of the FQR strain backgrounds GPS (A, D),

GPN (B, E), or GPL (C, F). Gray: Statistically significant sgRNAs passing the FDR or Q=5%

(30). 16 prominent targets from the CIP CRISPRi screen are highlighted. Dotted lines: selection

parameters setting fitness shifts by ≥0.2 to identify candidates. CIP: ciprofloxacin treated (50-

65µg/mL for the FQR strains or 0.11-0.12µg/mL for WT). Untx: untreated. WT: wildtype. GPS:

gyrAparCadeS. GPN: *gyrAparCadeN*. GPL: *gyrAparCadeL*.

Figure 4: sgRNAs with minimal effects in the absence of drug that decrease fitness in the

presence of CIP. (A) Statistical significance plotted as a function of mean fitness comparing

growth in the presence and absence of dCas9 inducer (100ng/mL aTc) for 7-8 generations in the

WT strain background (n=10). Gray: all sgRNAs. Blue: sgRNAs displaying neutral fitness

effects (scores between -0.05 and 0.05). Red: sgRNAs revealing a fitness enhancement (scores of

0.05 or higher). (B) Heat map of neutral and enhancer sgRNAs (as defined in A) that

significantly diminish FQR fitness in the presence of CIP. These pass the FDR or Q=5% (n=10)

via multiple t-test analyses comparing hypomorphs incubated with and without drug (first 4

columns) and between strain backgrounds (FQR-WT) during CIP treatment (last 3 columns).

Candidates were considered if at least 3 sgRNAs had fitness score decreased by $\geq$0.095 and at

least one sgRNA did so in more than one FQR strain. Gene numbers are originally preceded by

"ACX60_" for *A. baumannii* strain ATCC 17978UN (accession NZ_CP012004.1). 0: Perfect

sgRNA. FQR: fluoroquinolone resistant. CIP: ciprofloxacin. Untx: untreated. WT: wildtype.

GPL: *gyrAparCadeL*. GPN: *gyrAparCadeN*. GPS: *gyrAparCadeS*.

Figure 5: FQR strain hypomorphs that show high fitness levels in the absence of drug and

are hypersensitive to CIP show defects in membrane integrity. (A-H) Statistical significance

is plotted as a function of average fitness score differences for hypomorphs comparing dCas9

induced and uninduced conditions (A-D) or comparing CIP and Untx conditions (E-H) in the

WT (A, E), GPS (B, F), GPN (C, G), or GPL (D, H) strain backgrounds (n=10). Gray: All

sgRNAs. Prominent targets from the screen (7) are highlighted, of which most depletions had

little effect on fitness in the absence of drug but showed significantly lowered fitness in the FQR

strains during CIP exposure. Dotted lines: selection parameters defining either fitness defect or

high fitness. CIP: ciprofloxacin. Untx: untreated. WT: wildtype. GPS: *gyrAparCadeS*. GPN:

gyrAparCadeN. GPL: *gyrAparCadeL*.

Figure 6: GPL and GPN rely on proteins involved in LOS biogenesis to sustain growth

under CIP stress. (A) Mean fitness score changes imparted by each sgRNA during CIP

exposure as compared to untreated conditions as derived from the CIP CRISPRi screen (n=10).

Dotted line represents the mean of scrambled controls exerting no change in fitness. Light gray:

all sgRNAs designed for the listed target (mismatched and shuffled guides included). Data points

in red, blue, or orange highlight individual sgRNAs used to validate their respective targets'

impact on FQR fitness, as labeled in B. (B-D) These individual sgRNAs were constructed and

hypomorphs in all strain backgrounds were cultured in the presence or absence of CIP. Doubling

time differences in untreated and CIP treated conditions are superimposed. (B) Mean ± SEM

(n=3) growth rates (in min) of hypomorphs are compared to those harboring a shuffle control

sgRNA (NC_159). Unpaired t test analyses (individual variance each row) were performed on

pre-selected pairs following the two-stage set-up of Benjamini, Krieger, and Yekutieli with a

false discovery rate of 5% (30). Significant differences in doubling times yielded by targeted

depletions between strains during CIP exposure are shown: *, $p<0.05$; **, $p<0.02$; ***, $p<0.01$;

****, $p<0.001$. (C) Mean ± SEM (n≥3) growth rates are shown (in min) for the listed

hypomorphs. Two-way Anova with Tukey's multiple comparisons test reveals significant

variations in growth rates between GP and GPL or GPS during CIP exposure: ****, $p<0.0001$.

(D) Two-way Anova analyses using pre-selected pairs with Šídák's multiple comparisons test

(n=3): **, $p=0.0013$; ***, $p=0.0002$; ****, $p<0.0001$. CIP: ciprofloxacin (0.1 µg/mL for WT, 13

µg/mL for GP, 50 µg/mL for GPS, 45 µg/mL for GPN, and 60 µg/mL for GPL). Untx: untreated.

WT: wildtype. GP: *gyrAparC*. GPL: *gyrAparCadeL*. GPN: *gyrAparCadeN*. GPS:

gyrAparCadeS.

Figure 7: Zosurabalpin, Colistin, and Polymyxin B amplify Ciprofloxacin selectively in the GPL strain. (A-L) Growth of the indicated strains in LB at 37°C is displayed as the mean ± SEM (n≥3) after Y values were log transformed. Strains were incubated in the presence or absence of 2.5µM ZOS (A-F), COL or PB at 0.2 µg/mL, together with CIP at 0.11 µg/mL (A), 13 µg/mL (B), 15 µg/mL (D, E, G, H, J, K), 60 µg/mL (C, I, L), or 70 µg/mL (F). CIP: Ciprofloxacin. ZOS: Zosurabalpin. COL: Colistin. PB: Polymyxin B. WT: wildtype. GP: *gyrAparC*. GPL: *gyrAparCadeL*.

Figure 8: Depletion of LOS and lipoproteins results in defective envelope integrity that is exacerbated by the overexpression of the AdeL-dependent pump. (A-B) Growth of the indicated hypomorphs in the absence of CIP is displayed as the mean ± SEM (n=4) after Y values were log transformed. All strains were grown in presence or absence of 64 µg/mL VANC (A) or 0.05% SDS (B). (C) Mean growth rates (in min) ± SEM (n=3) for hypomorphs growing in the absence or presence of CIP with superimposed values. Statistical significance determined by a two-way Anova with pre-selected pairs and Šídák's multiple comparisons test: ****, p<0.0001. (D) Mean fold change ($2^{-\Delta\Delta CT}$) ± SEM (n=3) in transcript levels of each target in GPL during *ihfB* depletion with *ihfB*_P sgRNA relative to those observed with a non-targeting shuffle sgRNA (NC_159). (E) Mean ± SEM (n=3) pHi values are displayed. Cultures were incubated in LB with dCas9 inducer (100ng/mL aTc) to mid-exponential phase. Ordinary one-way Anova with Šídák's multiple comparisons test: *, p=0.0308; **, p=0.0021. (F) Mean fold change ($2^{-\Delta\Delta CT}$) ± SEM (n=3) in transcript levels of *atpF* during depletion using sgRNAs AtpF_5 (empty dark gray bars) and AtpF_20 (filled light gray bars) compared to those observed using a non-targeting shuffle sgRNA (NC₁). The values inside the bars: mean transcript levels ($2^{-\Delta\Delta CT}$)

relative to the shuffle control; values outside the bars: knockdown efficiency ($2^{\Delta\Delta CT}$) compared

to shuffle control. Untx: untreated. CIP: Ciprofloxacin (10 µg/mL for GP and GPLΔ*adeFGH* or

50 µg/mL for GPL). ZOS: Zosurabalpin. VANC: Vancomycin. SDS: Sodium dodecyl sulfate.

WT: Wildtype. GP: *gyrAparC*. GPL: *gyrAparCadeL*.

Supplementary Figure 1: Mapping of sgRNAs shows a normal distribution of reads.

Histograms of the mean reads for all sgRNAs (A) from the original pDNA library prior to

introduction into *A. baumannii*; (B) after introduction in the *A. baumannii* ATCC 17978UN

(WT) strain background at T1 prior to dCas9 induction (n=3); and (C) all shuffle sgRNAs in the

WT strain background at T2 after dCas9 induction (50ng/mL aTc, n=3). Reads were normalized

to total mapped reads and log transformed. Reads are binned by increments of 10.

Supplementary Figure 2: Lack of relationship between target location and fitness. (A-C)

Statistical significance (n=10) plotted as a function of fitness differences of hypomorphs in the

presence and absence of dCas9 induction (100ng/mL aTc) in the WT strain background. Gray:

all sgRNAs. Black: sgRNAs overlapping start codon (A). sgRNAs that from either the 3' or 5'

end of the sgRNA are 1-10nt (B) or 25-40nt (C) away from the first nucleotide of the start codon.

(B, C) Red: sgRNAs downstream the start codon. Blue: sgRNAs upstream the start codon. nt:

nucleotides. Dotted line: FDR or Q=5%. (D) Distribution of fitness differences (bin of 0.1) vs.

fraction of perfect sgRNAs from each category falling in each fitness range. (E) Expression

efficiency of four targets achieved by perfect sgRNAs that showed minimal effects on fitness are

depicted alongside that of *gyrA* that shows a strong fitness defect. Efficiency is noted as the

percent fold change in transcript levels observed by hypomorphs that were incubated with dCas9

inducer against those grown in parallel without inducer.

Supplementary Figure 3: Outgrowth of banks for 14-16 generations enhances fitness

defects. (A, B) Statistical significance (n=5) plotted as a function of fitness scoring differences

after growth for 7-8 generations compared to 14-16 generations in the presence (Black) or

absence (Gray) of dCas9 induction (100ng/mL aTc) in WT (A) or GPS (B) strain backgrounds.

(C-F) Statistical significance is plotted as a function of fitness changes after growth for 7-8

generations (C, E) or 14-16 generations (D, F) of all hypomorphs upon dCas9 induction. Black:

perfect guides. Dark gray: statistically significant sgRNAs passing the FDR or Q=5% via

multiple t-test analyses. Light gray: sgRNAs not statistically significant. Dotted red box to

highlight shifts in fitness that are observed between conditions. aTc: anhydrotetracycline. WT:

Wildtype. GPS: *gyrAparCadeS*.

Supplementary Figure 4: Outgrowth of banks for 14-16 generations does not substantially

affect fitness of the shuffle guides. (A-D) Statistical significance (n=5) plotted as a function of

fitness differences after growth for 7-8 generations (A, C) compared to 14-16 generations (B, D)

of all hypomorphs upon dCas9 induction. Black: perfect guides. Dark gray: statistically

significant sgRNAs passing the FDR or Q=5% via multiple t-test analyses. Light gray: sgRNAs

not statistically significant. Red: shuffle sgRNAs. Dotted blue box to visualize shifts in fitness

observed between conditions. WT: Wildtype. GPS: *gyrAparCadeS*.

Supplementary Figure 5: Depletion of a limited number of essential genes results in RND

pump overproducer-dependent fitness defects. (A) Statistical significance plotted as a

function of the difference in fitness scores for hypomorphs in GPS (A), GPN (B), or GPL (C) vs.

WT in the absence of drug after growth for 7-8 generations (n=10). Of the statistically significant

sgRNAs passing the FDR or Q=5% via multiple t-test analyses (Gray), the noted targets show

significant fitness score differences at least 20% from WT (limited to targets that are represented

by 3 or more sgRNAs in this category, with reads of each sgRNA being at least 10 at T1,

excluding ribosomal proteins). Black: sgRNAs not statistically significant. Dotted lines: selection

parameters wherein substantial changes in fitness are imparted by sgRNAs. (D, E) Fitness in

each strain background generated by sgRNAs of the indicated targets. Targets were identified as

in A-C, however, the targets displayed fitness score differences of at least 10% from WT in all

three FQR strains (D) or in two FQR strains (E). Dotted line marks neutral fitness value of 1.

Untx: untreated. WT: Wildtype. GPS: *gyrAparCadeS*. GPL: *gyrAparCadeL*. GPN:

gyrAparCadeN.

Supplementary Figure 6: Continued depletion through 14-16 generations in the absence of

drug reveals a smaller set of fitness determinants tied to AdeAB overexpression than that

identified after 7-8 generations. (A-B) Scatter plots of fitness resulting from indicated sgRNAs.

Targets were identified after comparing fitness of hypomorphs in WT (black circles) vs. GPS

(blue squares) after 7-8 generations (A, n=10) or 14-16 generations (B, n=5). The respective

sgRNAs showed significant fitness score differences of at least 10% between strains, passing the

FDR or Q=5% via multiple t-test analyses. Targets had to be represented by 3 or more sgRNAs,

reads of each sgRNA at T1 were at least 10, and with ribosomal proteins excluded from analysis.

Dotted line marks a neutral fitness value of 1. Red boxes highlight targets whose sgRNAs in

GPS yielded fitness scores that differed by at least 20% from WT. WT: Wildtype. GPS:

gyrAparCadeS.

Supplementary Figure 7: Identification of sgRNAs that alter fitness of FQR strains relative

to WT in presence of CIP. (A) Statistical significance plotted as a function of fitness score

differences between hypomorphs grown in CIP and Untx conditions for 7-8 generations in the

WT strain background (n=10). Gray: Statistically significant sgRNAs passing the FDR or Q=5%

via multiple t-test analyses. Black circles: sgRNAs not statistically significant. 16/18 of the top

targets identified as causing significantly altered fitness in the FQR strains compared to the WT

strain under CIP stress are highlighted as described for Figure 3, but here in the WT strain

background to depict the abated dependence on these proteins for fitness in WT. Dotted lines:

selection parameters wherein substantial changes in fitness are imparted by sgRNAs. (B-C)

Scatter plots exhibit the mean fitness generated by individual sgRNAs (n=10) corresponding to

the 18 top FQR fitness determinants as seen in the WT (black), GPS (gray), GPN (red), or GPL

(blue) strains after CIP treatment. These sgRNAs lower fitness of more than one FQR strain in

the presence of CIP (B) or provide more than one FQR strain with a fitness advantage during

CIP exposure (C). All unique sgRNAs are marked by circles, while the individual sgRNAs

causing fitness modifications that varied significantly between FQR and WT strains (and

between CIP vs Untx conditions) are labeled by empty squares. The dotted line marks the neutral

fitness value of 1. CIP: ciprofloxacin. Untx: untreated. WT: wildtype. GPL: *gyrAparCadeL.*

GPN: *gyrAparCadeN*. GPS: *gyrAparCadeS*.

Supplementary Figure 8: Fluctuations in hypomorph growth rates explained by varied

levels of transcript depletion. (A) Mean ± SEM (n=3) growth rate differences (in min) between

hypomorphs and a shuffle non-targeting control sgRNA (NC_159). Doubling time differences in

untreated and CIP-treated conditions are superimposed. Unpaired t test analyses (individual

variance each row) were performed on pre-selected pairs following the two-stage set-up of

Benjamini, Krieger, and Yekutieli with a false discovery rate of 5% (30). Target depletions

yielding significant differences in doubling times between GPL and WT during CIP exposure are

shown: *, p<0.05; **, p<0.02; ***, p<0.01; ****, p<0.001. (B) Mean fold change ($2^{\Delta\Delta CT}$) ±

SEM (n=3) in transcript levels for each target as compared to a non-targeting shuffle sgRNA

control (NC_159) in the WT (light gray) and GPL (dark gray) strain backgrounds. Values are

negative to display lowered transcript levels resulting from the targeted depletions. CIP:

ciprofloxacin (50 µg/mL for GPL and 0.09 µg/mL for WT). Untx: untreated. WT: wildtype.

GPL: *gyrAparCadeL.*

Supplementary Figure 9: Variability during LpxC and LptC depletion. (A) Mean ± SEM

(n=9) growth rate differences (in min) between hypomorphs and a shuffle non-targeting control

sgRNA (NC_159). Unpaired t test analyses (individual variance each row) were performed on

pre-selected pairs following the two-stage set-up of Benjamini, Krieger, and Yekutieli with a

false discovery rate of 5% (30). Target depletions yielding significant differences in doubling

times between GPL and WT during CIP exposure are shown: ****, p=0.000007. (B) Mean fold

change ($2^{\Delta\Delta CT}$) ± SEM (n=9) in transcript levels for each target as compared to a non-targeting

shuffle sgRNA control (NC_159) in the WT (black) and GPL (gray) strain backgrounds.

Transcript levels were evaluated via qRT-PCR in samples harvested from cultures grown in the

absence of drug (Untx). Values are negative to portray lowered transcript levels resulting from

the targeted depletions, and the mean values are noted above the x-axis. (A-B) Color of each data

point represents a biological replicate to better relate repression and growth rate results. CIP:

ciprofloxacin (50 µg/mL for GPL and 0.09 µg/mL for WT). Untx: untreated. WT: wildtype.

GPL: *gyrAparCadeL.*

References

1.   CDC. 2019. Antibiotic Resistance Threats in the United States. Atlanta, GA: US

Department of Health and Human Services, CDC.

2.   CDC. 2022. COVID-19: U.S. Impact on Antimicrobial Resistance, Special Report 2022.

Atlanta, GA: US Department of Health and Human Services, CDC.

3.   Tacconelli E, Carrara E, Savoldi A, Harbarth S, Mendelson M, Monnet DL, Pulcini C,

Kahlmeter G, Kluytmans J, Carmeli Y, Ouellette M, Outterson K, Patel J, Cavaleri M,

Cox EM, Houchens CR, Grayson ML, Hansen P, Singh N, Theuretzbacher U, Magrini N,

Group WHOPPLW. 2018. Discovery, research, and development of new antibiotics: the

WHO priority list of antibiotic-resistant bacteria and tuberculosis. Lancet Infect Dis

18:318-327.

4.   Sun J, Deng Z, Yan A. 2014. Bacterial multidrug efflux pumps: mechanisms, physiology

and pharmacological exploitations. Biochem Biophys Res Commun 453:254-67.

5.   Poole K. 2004. Efflux-mediated multiresistance in Gram-negative bacteria. Clin

Microbiol Infect 10:12-26.

6.   Coyne S, Courvalin P, Perichon B. 2011. Efflux-mediated antibiotic resistance in

890   *Acinetobacter* spp. Antimicrob Agents Chemother 55:947-53.

891   7.   Pagdepanichkit S, Tribuddharat C, Chuanchuen R. 2016. Distribution and expression of

892   the Ade multidrug efflux systems in *Acinetobacter baumannii* clinical isolates. Can J

893   Microbiol 62:794-801.

894   8.   Rumbo C, Gato E, Lopez M, Ruiz de Alegria C, Fernandez-Cuenca F, Martinez-Martinez

895   L, Vila J, Pachon J, Cisneros JM, Rodriguez-Bano J, Pascual A, Bou G, Tomas M,

896   Spanish Group of Nosocomial I, Mechanisms of A, Resistance to A, Spanish Society of

897   Clinical M, Infectious D, Spanish Network for Research in Infectious D. 2013.

898   Contribution of efflux pumps, porins, and beta-lactamases to multidrug resistance in

899   clinical isolates of *Acinetobacter baumannii*. Antimicrob Agents Chemother 57:5247-57.

900   9.   Fernando D, Zhanel G, Kumar A. 2013. Antibiotic resistance and expression of

901   resistance-nodulation-division pump- and outer membrane porin-encoding genes in

902   *Acinetobacter* species isolated from Canadian hospitals. Can J Infect Dis Med Microbiol

903   24:17-21.

904   10.   Geisinger E, Vargas-Cuebas G, Mortman NJ, Syal S, Dai Y, Wainwright EL, Lazinski D,

905   Wood S, Zhu Z, Anthony J, van Opijnen T, Isberg RR. 2019. The Landscape of

906   Phenotypic and Transcriptional Responses to Ciprofloxacin in *Acinetobacter baumannii*:

907   Acquired Resistance Alleles Modulate Drug-Induced SOS Response and Prophage

908   Replication. mBio 10.

909   11.   Naidu V, Bartczak A, Brzoska AJ, Lewis P, Eijkelkamp BA, Paulsen IT, Elbourne LDH,

910   Hassan KA. 2023. Evolution of RND efflux pumps in the development of a successful

911   pathogen. Drug Resist Updat 66:100911.

12. Magnet S, Courvalin P, Lambert T. 2001. Resistance-nodulation-cell division-type efflux

pump involved in aminoglycoside resistance in *Acinetobacter baumannii* strain BM4454.

Antimicrob Agents Chemother 45:3375-80.

13. Coyne S, Rosenfeld N, Lambert T, Courvalin P, Perichon B. 2010. Overexpression of

resistance-nodulation-cell division pump AdeFGH confers multidrug resistance in

Acinetobacter baumannii. Antimicrob Agents Chemother 54:4389-93.

14. Yoon EJ, Chabane YN, Goussard S, Snesrud E, Courvalin P, De E, Grillot-Courvalin C.

2015. Contribution of resistance-nodulation-cell division efflux systems to antibiotic

resistance and biofilm formation in *Acinetobacter baumannii*. mBio 6.

15. Rosenfeld N, Bouchier C, Courvalin P, Perichon B. 2012. Expression of the resistance-

nodulation-cell division pump AdeIJK in *Acinetobacter baumannii* is regulated by AdeN,

a TetR-type regulator. Antimicrob Agents Chemother 56:2504-10.

16. Hamami E, Huo W, Hernandez-Bird J, Castaneda A, Bai J, Syal S, Ortiz-Marquez JC,

van Opijnen T, Geisinger E, Isberg RR. 2025. Identification of determinants that allow

maintenance of high-level fluoroquinolone resistance in *Acinetobacter baumannii*. mBio

16:e0322124.

17. Jost M, Santos DA, Saunders RA, Horlbeck MA, Hawkins JS, Scaria SM, Norman TM,

Hussmann JA, Liem CR, Gross CA, Weissman JS. 2020. Titrating gene expression using

libraries of systematically attenuated CRISPR guide RNAs. Nat Biotechnol 38:355-364.

18. Hawkins JS, Silvis MR, Koo BM, Peters JM, Osadnik H, Jost M, Hearne CC, Weissman

JS, Todor H, Gross CA. 2020. Mismatch-CRISPRi Reveals the Co-varying Expression-

Fitness Relationships of Essential Genes in *Escherichia coli* and *Bacillus subtilis*. Cell

Syst 11:523-535 e9.

19.   Rousset F, Cui L, Siouve E, Becavin C, Depardieu F, Bikard D. 2018. Genome-wide
936   CRISPR-dCas9 screens in *E. coli* identify essential genes and phage host factors. PLoS
937   Genet 14:e1007749.

938   20.   Ward RD, Tran JS, Banta AB, Bacon EE, Rose WE, Peters JM. 2024. Essential gene
939   knockdowns reveal genetic vulnerabilities and antibiotic sensitivities in *Acinetobacter*
940   *baumannii*. mBio 15:e0205123.

941   21.   Tran JS, Ward RD, Iruegas-Lopez R, Ebersberger I, Peters JM. 2025. Chemical genomics
942   informs antibiotic and essential gene function in *Acinetobacter baumannii*. PLoS Genet
943   21:e1011642.

944   22.   Geisinger E, Mortman NJ, Dai Y, Cokol M, Syal S, Farinha A, Fisher DG, Tang AY,
945   Lazinski DW, Wood S, Anthony J, van Opijnen T, Isberg RR. 2020. Antibiotic
946   susceptibility signatures identify potential antimicrobial targets in the *Acinetobacter*
947   *baumannii* cell envelope. Nat Commun 11:4522.

948   23.   Bai J, Dai Y, Farinha A, Tang AY, Syal S, Vargas-Cuebas G, van Opijnen T, Isberg RR,
949   Geisinger E. 2021. Essential Gene Analysis in *Acinetobacter baumannii* by High-Density
950   Transposon Mutagenesis and CRISPR Interference. J Bacteriol 203:e0056520.

951   24.   DeJesus MA, Ambadipudi C, Baker R, Sassetti C, Ioerger TR. 2015. TRANSIT--A
952   Software Tool for Himar1 TnSeq Analysis. PLoS Comput Biol 11:e1004401.

953   25.   DeJesus MA, Zhang YJ, Sassetti CM, Rubin EJ, Sacchettini JC, Ioerger TR. 2013.
954   Bayesian analysis of gene essentiality based on sequencing of transposon insertion
955   libraries. Bioinformatics 29:695-703.

956   26.   Gallagher LA, Ramage E, Weiss EJ, Radey M, Hayden HS, Held KG, Huse HK,

957   Zurawski DV, Brittnacher MJ, Manoil C. 2015. Resources for Genetic and Genomic

958   Analysis of Emerging Pathogen *Acinetobacter baumannii*. J Bacteriol 197:2027-35.

959   27.   Wang N, Ozer EA, Mandel MJ, Hauser AR. 2014. Genome-wide identification of

960   *Acinetobacter baumannii* genes necessary for persistence in the lung. mBio 5:e01163-14.

961   28.   Peters JM, Koo BM, Patino R, Heussler GE, Hearne CC, Qu J, Inclan YF, Hawkins JS,

962   Lu CHS, Silvis MR, Harden MM, Osadnik H, Peters JE, Engel JN, Dutton RJ, Grossman

963   AD, Gross CA, Rosenberg OS. 2019. Enabling genetic analysis of diverse bacteria with

964   Mobile-CRISPRi. Nat Microbiol 4:244-250.

965   29.   van Opijnen T, Bodi KL, Camilli A. 2009. Tn-seq: high-throughput parallel sequencing

966   for fitness and genetic interaction studies in microorganisms. Nat Methods 6:767-72.

967   30.   Benjamini Y, Krieger, A. M., & Yekutieli, D. 2006. Adaptive linear step-up procedures

968   that control the false discovery rate. Biometrika 93:491-507.

969   31.   Yakhnin AV, Bubunenko M, Mandell ZF, Lubkowska L, Husher S, Babitzke P, Kashlev

970   M. 2023. Robust regulation of transcription pausing in *Escherichia coli* by the ubiquitous

971   elongation factor NusG. Proc Natl Acad Sci U S A 120:e2221114120.

972   32.   Baniulyte G, Singh N, Benoit C, Johnson R, Ferguson R, Paramo M, Stringer AM, Scott

973   A, Lapierre P, Wade JT. 2017. Identification of regulatory targets for the bacterial Nus

974   factor complex. Nat Commun 8:2027.

975   33.   Leus IV, Weeks JW, Bonifay V, Smith L, Richardson S, Zgurskaya HI. 2018. Substrate

976   Specificities and Efflux Efficiencies of RND Efflux Pumps of *Acinetobacter baumannii*.

977   J Bacteriol 200.

34.   Sugawara E, Nikaido H. 2014. Properties of AdeABC and AdeIJK efflux systems of

Acinetobacter baumannii compared with those of the AcrAB-TolC system of *Escherichia*

coli. Antimicrob Agents Chemother 58:7250-7.

35.   Raskin DM, Judson N, Mekalanos JJ. 2007. Regulation of the stringent response is the

essential function of the conserved bacterial G protein CgtA in *Vibrio cholerae*. Proc Natl

Acad Sci U S A 104:4636-41.

36.   Diez S, Ryu J, Caban K, Gonzalez RL, Jr., Dworkin J. 2020. The alarmones (p)ppGpp

directly regulate translation initiation during entry into quiescence. Proc Natl Acad Sci U

S A 117:15565-15572.

37.   Pahil KS, Gilman MSA, Baidin V, Clairfeuille T, Mattei P, Bieniossek C, Dey F, Muri D,

Baettig R, Lobritz M, Bradley K, Kruse AC, Kahne D. 2024. A new antibiotic traps

lipopolysaccharide in its intermembrane transporter. Nature 625:572-577.

38.   Chen S, Hu M, Hu A, Xue Y, Wang S, Liu F, Li C, Zhou X, Zhou J. 2022. The

integration host factor regulates multiple virulence pathways in bacterial pathogen

Dickeya zeae MS2. Mol Plant Pathol 23:1487-1507.

39.   Fan Z, Fu T, Li Z, Du B, Cui X, Zhang R, Feng Y, Zhao H, Xue G, Cui J, Yan C, Gan L,

Feng J, Xu Z, Yu Z, Tian Z, Ding Z, Chen J, Chen Y, Yuan J. 2023. The role of

integration host factor in biofilm and virulence of high-alcohol-producing *Klebsiella*

pneumoniae. Microbiol Spectr 11:e0117023.

40.   Nicolau SE, Lewis K. 2022. The Role of Integration Host Factor in *Escherichia coli*

Persister Formation. mBio 13:e0342021.

41.   Friedman DI. 1988. Integration host factor: a protein for all reasons. Cell 55:545-54.

42. Sielaff H, Duncan TM, Borsch M. 2018. The regulatory subunit epsilon in *Escherichia*

coli F(O)F(1)-ATP synthase. Biochim Biophys Acta Bioenerg 1859:775-788.

43. Guo H, Courbon GM, Bueler SA, Mai J, Liu J, Rubinstein JL. 2021. Structure of

mycobacterial ATP synthase bound to the tuberculosis drug bedaquiline. Nature 589:143-

147.

44. Saw WG, Le KCM, Shin J, Kwek JHM, Wong CF, Ragunathan P, Fong TC, Muller V,

Gruber G. 2023. Atomic insights of an up and down conformation of the *Acinetobacter*

baumannii F(1) -ATPase subunit epsilon and deciphering the residues critical for ATP

hydrolysis inhibition and ATP synthesis. FASEB J 37:e23040.

45. Demmer JK, Phillips BP, Uhrig OL, Filloux A, Allsopp LP, Bublitz M, Meier T. 2022.

Structure of ATP synthase from ESKAPE pathogen *Acinetobacter baumannii*. Sci Adv

8:eabl5966.

46. Xu Q, Guo M, Yu F. 2023. beta-Barrel Assembly Machinery (BAM) Complex as Novel

Antibacterial Drug Target. Molecules 28.

47. Sklar JG, Wu T, Kahne D, Silhavy TJ. 2007. Defining the roles of the periplasmic

chaperones SurA, Skp, and DegP in *Escherichia coli*. Genes Dev 21:2473-84.

48. Fenn KL, Horne JE, Crossley JA, Bohringer N, Horne RJ, Schaberle TF, Calabrese AN,

Radford SE, Ranson NA. 2024. Outer membrane protein assembly mediated by BAM-

SurA complexes. Nat Commun 15:7612.

49. Tse MW, Zhu M, Peters B, Hamami E, Chen J, Davis KP, Nitz S, Weller J, Warrier T,

Hunt DK, Morales Y, Kawate T, Gaulin JL, Come JH, Hernandez-Bird J, Huo W,

Neisewander I, Yu X, Aceves-Salvador JA, Kiessling LL, Hung DT, Mecsas J, Aldridge

BB, Isberg RR, Blainey PC. 2025. Large-scale combination screens reveal small-

molecule sensitization of antibiotic-resistant gram-negative ESKAPE pathogens. Proc

Natl Acad Sci U S A 122:e2402017122.

50.   Ruiz N, Gronenberg LS, Kahne D, Silhavy TJ. 2008. Identification of two inner-

membrane proteins required for the transport of lipopolysaccharide to the outer

membrane of *Escherichia coli*. Proc Natl Acad Sci U S A 105:5537-42.

51.   Sperandeo P, Cescutti R, Villa R, Di Benedetto C, Candia D, Deho G, Polissi A. 2007.

Characterization of lptA and lptB, two essential genes implicated in lipopolysaccharide

transport to the outer membrane of *Escherichia coli*. J Bacteriol 189:244-53.

52.   Arunmanee W, Pathania M, Solovyova AS, Le Brun AP, Ridley H, Basle A, van den

Berg B, Lakey JH. 2016. Gram-negative trimeric porins have specific LPS binding sites

that are essential for porin biogenesis. Proc Natl Acad Sci U S A 113:E5034-43.

53.   Webby MN, Oluwole AO, Pedebos C, Inns PG, Olerinyova A, Prakaash D, Housden NG,

Benn G, Sun D, Hoogenboom BW, Kukura P, Mohammed S, Robinson CV, Khalid S,

Kleanthous C. 2022. Lipids mediate supramolecular outer membrane protein assembly in

bacteria. Sci Adv 8:eadc9566.

54.   Dzioba-Winogrodzki J, Winogrodzki O, Krulwich TA, Boin MA, Hase CC, Dibrov P.

2009. The *Vibrio cholerae* Mrp system: cation/proton antiport properties and

enhancement of bile salt resistance in a heterologous host. J Mol Microbiol Biotechnol

16:176-86.

55.   Ito M, Guffanti AA, Wang W, Krulwich TA. 2000. Effects of nonpolar mutations in each

of the seven *Bacillus subtilis* mrp genes suggest complex interactions among the gene

products in support of Na(+) and alkali but not cholate resistance. J Bacteriol 182:5663-

70.

56.   Kosono S, Haga K, Tomizawa R, Kajiyama Y, Hatano K, Takeda S, Wakai Y, Hino M,

1047   Kudo T. 2005. Characterization of a multigene-encoded sodium/hydrogen antiporter

1048   (sha) from *Pseudomonas aeruginosa*: its involvement in pathogenesis. J Bacteriol

1049   187:5242-8.

1050   57.   Koumandou VL, Kossida S. 2014. Evolution of the F0F1 ATP synthase complex in light

1051   of the patchy distribution of different bioenergetic pathways across prokaryotes. PLoS

1052   Comput Biol 10:e1003821.

1053   58.   Liu J, Hicks DB, Krulwich TA. 2013. Roles of AtpI and two YidC-type proteins from

1054   alkaliphilic *Bacillus pseudofirmus* OF4 in ATP synthase assembly and nonfermentative

1055   growth. J Bacteriol 195:220-30.

1056   59.   Dalhoff A, Stubbings W, Schubert S. 2011. Comparative *In Vitro* Activities of the Novel

1057   Antibacterial Finafloxacin against Selected Gram-Positive and Gram-Negative Bacteria

1058   Tested in Mueller-Hinton Broth and Synthetic Urine. Antimicrobial Agents and

1059   Chemotherapy 55:1814-1818.

1060   60.   Dalhoff A, Schubert S, Vente A. 2017. Pharmacodynamics of Finafloxacin,

1061   Ciprofloxacin, and Levofloxacin in Serum and Urine against TEM- and SHV-Type

1062   Extended-Spectrum-ß-Lactamase-Producing *Enterobacteriaceae* Isolates from Patients

1063   with Urinary Tract Infections. Antimicrobial Agents and Chemotherapy

1064   61:10.1128/aac.02446-16.

1065   61.   Manuse S, Shan Y, Canas-Duarte SJ, Bakshi S, Sun WS, Mori H, Paulsson J, Lewis K.

1066   2021. Bacterial persisters are a stochastically formed subpopulation of low-energy cells.

1067   PLoS Biol 19:e3001194.

62.   Shan Y, Brown Gandt A, Rowe SE, Deisinger JP, Conlon BP, Lewis K. 2017. ATP-

Dependent Persister Formation in *Escherichia coli*. mBio 8.

63.   El Meouche I, Dunlop MJ. 2018. Heterogeneity in efflux pump expression predisposes

antibiotic-resistant cells to mutation. Science 362:686-690.

64.   Huo W, Busch LM, Hernandez-Bird J, Hamami E, Marshall CW, Geisinger E, Cooper

VS, van Opijnen T, Rosch JW, Isberg RR. 2022. Immunosuppression broadens

evolutionary pathways to drug resistance and treatment failure during *Acinetobacter*

baumannii pneumonia in mice. Nat Microbiol 7:796-809.

65.   Bollenbach T, Quan S, Chait R, Kishony R. 2009. Nonoptimal microbial response to

antibiotics underlies suppressive drug interactions. Cell 139:707-18.

66.   Chait R, Craney A, Kishony R. 2007. Antibiotic interactions that select against

resistance. Nature 446:668-71.

67.   Cokol M, Kuru N, Bicak E, Larkins-Ford J, Aldridge BB. 2017. Efficient measurement

and factorization of high-order drug interactions in *Mycobacterium tuberculosis*. Sci Adv

3:e1701881.

68.   Ocampo PS, Lazar V, Papp B, Arnoldini M, Abel zur Wiesch P, Busa-Fekete R, Fekete

G, Pal C, Ackermann M, Bonhoeffer S. 2014. Antagonism between bacteriostatic and

bactericidal antibiotics is prevalent. Antimicrob Agents Chemother 58:4573-82.

69.   Martin M. 2011. Cutadapt removes adapter sequences from high-throughput sequencing

reads. 2011 17:3.

70.   van Opijnen T, Camilli A. 2013. Transposon insertion sequencing: a new tool for

systems-level analysis of microorganisms. Nat Rev Microbiol 11:435-42.

1090

A. baumannii
harboring 12,000
sgRNA library

Target sgRNA
x 406 essential
gene targets

1hr

T1

Uninduced | Ind, Untx | Ind, CIP

7-8
gens

T2

Evaluate the fitness
of hypomorphs

Determine essential
gene targets that
aggravate RND pump
overexpression &
sensitize high-level
FQR strains to CIP

1 "perfect" guide   5'-agttcgtcctcaatggcaatcggt-3'

23 mismatched   5'-agttcgtcctcaatggcaatcg**at**-3'

5'-agttcgtcctcaa**g**ggcaatcggt-3'

5 shuffles   5'-gtatccgtgtgggttaaccatacc-3'

+ additional pool shuffle sgRNAs

Fig 1

Fig 2

Fig 3

Fig 4

Fig 5

Fig 6

Fig 7

A Vancomycin sensitivity in GPL

B SDS sensitivity in GPL

C

D IhfB Depletion

E

F *atpF*

Fig 8

A

B

C

Supp Fig 1

Number of Reads

Supp Fig 2

Supp Fig 3

Supp Fig 4

Supp Fig 5

Supp Fig 6

Supp Fig 7

Supp Fig 8

Supp Fig 9