1

The intrinsically disordered N-terminus of SUMO1 is an intramolecular inhibitor

of SUMO1 interactions

Sebastian M. Richter[1,6], Fan Jin[2,3,6], Tobias Ritterhoff[4,6], Aleksandra Fergin[5], Eric

Maurer[1], Andrea Frank[1], Michael Daube[5], Alex Hajnal[5], Rachel Klevit[4], Frauke Gräter[2,3],

Annette Flotho[1,*] and Frauke Melchior[1,*]

[1] Zentrum für Molekulare Biologie der Universität Heidelberg (ZMBH),

DKFZ - ZMBH Alliance, Heidelberg, Germany

[2] Heidelberg Institute for Theoretical Studies, Heidelberg, Germany.

[3] Interdisciplinary Center for Scientific Computing, Heidelberg University, Heidelberg,

Germany.

[4] Department of Biochemistry, University of Washington, Seattle, WA 98195, USA.

[5] Department of Molecular Life Sciences, University of Zürich, Zürich, Switzerland

[6] contributed equally

* Corresponding authors

Email: f.melchior@zmbh.uni-heidelberg.de

a.flotho@zmbh.uni-heidelberg.de

Phone: +49-6221 546790

Running Title:

SUMO´s N-terminus controls SIM interactions

Key words:

SUMMARY

Ubiquitin-related proteins of the SUMO family are reversibly attached to thousands of

proteins in eukaryotic cells. Many SUMO substrates, effectors and enzymes carry short

motifs (SIMs) that mediate low affinity interactions with SUMO proteins. How specificity

is achieved in target selection, SUMO paralogue choice and SUMO-dependent

interactions is largely unknown. A unique but poorly understood feature of SUMO

proteins is their intrinsically disordered N-terminus. We reveal a function for N-termini of

human, *C. elegans,* and yeast SUMO proteins as intramolecular inhibitors of SUMO-

SIM interactions. Mutational analyses, NMR spectroscopy, and Molecular Dynamics

simulations indicate that SUMO's N-terminus can inhibit SIM binding by fast and fuzzy

interactions with SUMO's core. Deletion of the *C. elegans* SUMO1 N-terminus leads to

p53-dependent apoptosis during germline development, indicating an important role in

DNA damage repair. Our findings reveal a mechanism of disorder-based autoinhibition

that contributes to the specificity of SUMOylation and SUMO-dependent interactions.

INTRODUCTION

SUMOylation, the reversible attachment of small ubiquitin-related modifier SUMO to target proteins, serves as a molecular switch that regulates protein functions. It is particularly important in pathways that contribute to chromatin organization and function including transcription, replication, DNA damage repair, nucleocytoplasmic transport, and chromosome segregation[1-4]. At least a thousand proteins are subject to this essential protein modification under physiological conditions, and many more in severe stress (e.g., [5,6]). The mechanism of SUMOylation resembles that of Ubiquitylation: Isopeptide bond formation between the C-terminal di-glycine motif of SUMO and the ε-amino group of a lysine sidechain in target proteins is catalyzed by an enzymatic trio of the SUMO-specific E1 activating heterodimer Aos1/Uba2 (SAE1/SAE2), the SUMO E2 conjugating enzyme Ubc9, and usually one of several SUMO E3 ligases. SUMO-specific isopeptidases reverse the modification. One striking difference to the Ubiquitin system is the small number of enzymes involved. Over 600 E3 ligases are known for Ubiquitylation but fewer than ten have been identified for SUMOylation, amongst them the Siz/PIAS RING E3 ligases, ZNF451, and the nucleoporin RanBP2[1,7]. Similarly, nearly one hundred isopeptidases with specificity for Ubiquitin are known, yet only eight enzymes appear to act on mammalian SUMO[8,9]. On the other hand, many organisms have more than one SUMO protein: Whereas *S. cerevisiae, D. melanogaster, and C. elegans* harbor only one SUMO gene, the weed *A. thaliana* expresses 8 SUMO genes (reviewed in [10]) and the human genome encodes five SUMO paralogues, three of which are ubiquitously expressed, SUMO1 and the twins SUMO2 and SUMO3 (reviewed in [7]). SUMO2 and 3 are virtually identical but share only about 50% identity with SUMO1. The two classes of SUMO proteins differ in expression levels, response to cellular stress, and their ability to form SUMO chains. The N-termini of human SUMO2 and 3 as well as *S. cerevisiae* Smt3 contain at least one lysine residue, on which SUMO chains can

assemble[11-13], whereas the human SUMO1 and *C. elegans* SMO-1 N-termini are

devoid of lysine (see also below). Although many proteins can be modified *in vitro* and

in cells with both classes of SUMO proteins, a significant number of proteins is

preferentially or exclusively modified with one or the other SUMO paralogue at

endogenous expression levels in cells (e.g. [14]).

How paralogue-specific modification and effector binding are brought about is only

partially understood. Both events require SUMO-binding (reviewed, e.g. in [7]). A

SUMO-binding motif in the target may help to recruit the Ubc9 - SUMO thioester. If

this motif prefers a specific SUMO paralogue, the target is also preferentially modified

with this paralogue as in USP25 and BLM[15,16]. SUMO-binding motifs in downstream

effectors help to recruit specific SUMOylated proteins. Examples are the recruitment

of Srs2 to SUMOylated PCNA[17] and the recruitment of the ubiquitin ligase RNF4 to

polySUMOylated PML[18,19].

To date, at least three different classes of SUMO interactions have been identified, of

which the best studied class I interactions involve <u>S</u>UMO <u>I</u>nteraction <u>M</u>otifs (SIMs)[20].

Known examples for class II interactions with a surface of SUMO opposite of the SIM

binding site are Dpp9 and Ubc9 [7,21-24], while ZZ domains have been shown to either

bind SUMO's core as in the case of CBP[25] or the absolute N-terminus as for HERC2[26].

SIMs typically consist of a four amino acid patch with three hydrophobic residues,

most commonly valine or isoleucine, and a variable at position 3 ([P/I/L/V/M]-[I/L/V/M]-

x- [I/L/V/M]), followed by a stretch of negatively-charged amino acids and/or serine

residues[27]. In some cases, the orientation of the SIM can be reversed. A distinct

subclass of SIMs is characterized by a [V/I/L/F/Y]-[I/V]-D-L-T consensus motif[28]. The

hydrophobic amino acids in the core of SIMs insert as a β-strand into the hydrophobic

groove formed by the β2-strand and the α1-helix of SUMO (SIM-binding groove)[29,30].

An exception is TDP2, which binds the SIM-binding area of SUMO2/3 via five

separate binding elements, two β-sheets and three surface loops, forming a so-called

"split SIM"[31]. As is common for binding modules that recognize Ubiquitin or Ubls[32],

SUMO - SIM interactions are usually of low affinity (K_D between 1 - 100 µM)[20].

Biological relevance comes about by combining multiple SIMs (as in RNF4),

combining a SIM with a second interaction module specific for a given substrate (as is

true for Srs2, which recognizes both SUMO and PCNA), or through posttranslational

modifications in proximity to the SIM that increase the affinity for SUMO (e.g.,

phosphorylation in proximity to a SIM in PIAS1[33]). How paralogue-specific interactions

are established in SUMO - SIM interactions remains enigmatic. Most of the amino

acids in the SIM-binding groove that contribute to SUMO - SIM interactions by NMR

are conserved between SUMO1 and SUMO2/3[34-38]. Initially it was proposed that a

patch of negative amino acids adjacent to the SIM is required for SUMO1 - but

dispensable for SUMO2/3 - interaction[29]. However, USP25[15] and ATF7IP[38] exhibit a

high preference for SUMO2/3 despite a patch of negative amino acids adjacent to

their SIM. Attempts to convert SUMO2/3 into a SUMO1 interacting protein by

mutagenesis have thus far been unsuccessful (e.g., [38]). A defining feature of SUMO

proteins is the intrinsically disordered N-terminus[39], whose function is only partly

understood[7,40-42]. While N-termini of the SUMO2/3 family contain a SUMOylation

consensus motif (ΨKxE, where Ψ is a bulky hydrophobic amino acid) used for SUMO

chain formation[11-13], this motif is lacking in SUMO1 proteins. In yeast, polymeric

SUMO chains contribute to chromatin regulation[40,43] but the N-terminus itself is

dispensable for essential SUMO functions[41]. Here, we reveal an unexpected function

for the intrinsically disordered N-termini as intramolecular inhibitors of SUMO - SIM

interactions.

RESULTS

Intrinsically disordered N-termini of SUMO1 proteins inhibit protein interactions

All SUMO orthologs from yeast to human carry an intrinsically disordered N-terminal

extension[39], typically in the range of 13 - 23 amino acids, as shown in Figure 1A for

SUMO1 (adapted from [39]) and as indicated by boxed residues in the primary

structures of the four SUMO proteins used in this study, human SUMO1, human

SUMO2, *C. elegans* SMO-1, and *S. cerevisiae* Smt3 (Fig. 1B). Previously, we had

performed pull-down screens that allowed us to identify paralogue - specific SUMO-

binding partners[15]. In these experiments, we also included a SUMO1 variant lacking

the first 19 amino acids as we hypothesized that the N-terminus contributes to

interactions. To our surprise, the contrary was the case - the pulldown with

SUMO1ΔN19 yielded more proteins than the pull-down with SUMO1 (Fig. S1A + S1B).

To test whether SUMO's disordered N-terminus indeed prevents protein interactions,

we conducted pull-down assays with validated binding partners and wild-type SUMO1,

the N-terminally truncated variants SUMO1ΔN19 and SUMO1ΔN10, wild-type

SUMO2, and SUMO2ΔN13 (Fig. 1C - F). All SUMO variants were untagged and

immobilized on CNBr-activated Sepharose in random orientation. Binding partners

included two proteins with preference for SUMO2/3, the canonical SIM-containing

protein USP25[15], and TDP2, which has an unusual "split SIM"[31], the SUMO1-specific

SIM-containing RanBP2 fragment RanBP2ΔFG[44,45] and Ubc9, which binds SUMO1

and SUMO2/3 via a region distinct from the SIM binding groove[46]. Indeed, deletion of

the complete SUMO1 N-terminus dramatically increased interaction with USP25 (Fig.

1C + D). Similarly, interaction with TDP2 was enhanced when SUMO1's N-terminus

was removed, while the interaction with Ubc9 was unaffected. Some increase was

observed when half the SUMO1 N-terminus was deleted. As shown in Figure 1E and

1F, deletion of SUMO1's N-terminus allowed USP25 to bind SUMO1 as efficiently as

wt SUMO2. Importantly, deletion of SUMO2's N-terminus (ΔN13) did not increase

binding any further, even for the SUMO1-specific RanBP2, suggesting that SUMO1's -

but not SUMO2′s - N-terminus inhibits the interaction. To exclude artefacts due to the

SUMO immobilization, we repeated pulldown experiments with C-terminally GST-

tagged SUMO variants immobilized on Glutathione beads (Fig. S1C + S1D). Again,

USP25 bound to ΔN19 SUMO1 as efficiently as to SUMO2, and much better than to

SUMO1.

To address whether the inhibitory role of the unstructured N-terminus is a unique

feature of mammalian SUMO1 or whether it may be conserved in evolution, we turned

to yeast and *C. elegans*. The single yeast SUMO protein Smt3 shares about 45%

sequence identity with both SUMO1 and SUMO2/3. While hsSUMO1 can complement

loss of Smt3, hsSUMO2 and 3 fail to do so, due to incompatibility of these human

SUMO proteins with some of the yeast SUMO enzymes[41,47]. Like SUMO2/3, Smt3 has

SUMOylation consensus sites in its unstructured N-terminal region that are utilized for

chain formation (reviewed in [48]), yet loss of the N-terminal lysines or the Smt3 N-

terminus is tolerated without detectable growth defects[41]. The single *C. elegans*

SUMO protein SMO-1 shares about 67% sequence identity with SUMO1 and 48%

identity with SUMO2/3. Intriguingly, its N-terminus contains no lysine residues and can

thus not contribute to canonical SUMO chain formation. As shown in Figures 1I an 1J,

deleting the N-terminus in Smt3 and SMO-1 indeed enhanced interaction with SIM-

containing model proteins, the SUMO E3 ligases *S.c.* Siz1[49], and *C.e.* GEI-17[50],

respectively. These results suggest that the inhibitory effect of SUMO1's N-terminus is

indeed an evolutionarily conserved trait.

SUMO1's disordered N-terminus interferes with SIM-dependent functions

One explanation for increased interaction upon deletion of SUMO1's N-terminus is that

it usually occludes the SUMO1 SIM-binding groove, a feature apparently unique to

SUMO1. If so, its removal should lead both to enhanced non-covalent interaction (Fig.

2A, C-F), and to enhanced SUMOylation (Fig. 2B, G-L).

Indeed, a SIM-deficient USP25 (USP25 VI91,92AA[15]) lost binding not only to SUMO2

but also to SUMO1 ΔN19 (Fig. 2C + D), indicating that the interaction with truncated

SUMO1 also depends on a bona fide SIM. Moreover, mutating two residues in

SUMO1's SIM-binding groove, Val38 and Lys39 (Fig. 2E + F) led to a significant loss

of interaction with USP25 and with RanBP2.

We next turned to SUMOylation. As reported previously, USP25 was SUMOylated

more efficiently with SUMO2 compared to SUMO1. However, SUMO1ΔN19 can

SUMOylate USP25 as efficiently as SUMO2 (Fig. 2G). Consistent with binding assays

(Fig. 1E-F), N-terminal truncation of SUMO2 did not further enhance SUMOylation

efficiency. As a control, we used the SIM-independent SUMO target RanGAP1 (Fig.

2H), which was equally modified with all SUMO1 and SUMO2 variants. Similar results

were obtained when we compared *C. elegans* wild-type SMO-1 and SMO-1ΔN12 for

their ability to modify our model human substrates (Fig. 2I + J). In SUMOylation

assays with USP25 SIM mutants (Fig. 2K) and with SUMO1ΔN19 variants that carried

mutations in the SIM-binding groove (Fig. 2L), SUMOylation of USP25 was abolished,

indicating that the increase in SUMOylation upon deletion of SUMO1's N-terminus

depends on a SIM-dependent interaction.

Taken together, our findings indicate that SUMO1's intrinsically disordered N-terminus

inhibits both SIM-dependent interactions and SIM-mediated SUMOylation. This offers

an unanticipated explanation for the apparent SUMO2 preference of SIM-dependent

SUMO binding partners such as USP25.

NMR experiments indicate physical interaction between the N-terminus

and the SIM-binding groove of SUMO1

To gain structural insights into the mechanism by which SUMO1's N-terminus

prevents SUMO - SIM interactions, we turned to nuclear magnetic resonance (NMR).

To assess motions in the nanosecond/sub-nanosecond time regime, ^{1}H-^{15}N-

heteronuclear Nuclear Overhauser effect (hNOE) measurements were made on

SUMO1. Residues 21-93 exhibit uniform positive hNOEs, characteristic of stable,

globular domains and thus consistent with the known folded structure of the core (Fig.

3A). The C-terminal residues of SUMO1 have hNOEs near or below 0, a hallmark of

highly flexible regions and consistent with the high degree of flexibility commonly

observed in Ubl proteins near their C-terminal Gly-Gly motif[39,51]. Residues 4-20 in the

N-terminus of SUMO1 displayed lower hNOE values than those of the folded core

consistent with their being more flexible. However, while residues 5-9 showed large

negative hNOEs consistent with high flexibility, residues 10-20 have hNOE values

close to 0. This is suggestive of a split in the SUMO1 N-terminus into a highly flexible

region (residues 5-9) and a region of intermediate flexibility (residues 10-20) whose

dynamics are slower but not as slow as the folded core. Residue 4 has a hNOE value

close to 0 hinting at slower dynamics in the very N-terminal part of the protein.

Unfortunately, residues 1-3 were not available for hNOE analysis, so no conclusions

can be drawn about this sub-region of the N-terminus. The slower-than-expected

dynamics of residues 10-20 indicate that this region spends time associated with the

structured core. We speculate that the interactions could occlude the SIM-binding

groove of SUMO1.

225
To investigate this further, we compared HSQC-TROSY spectra of full-length SUMO1

and the ΔN19 variant (Fig. 3B) to assess how the presence and absence of the N-

terminus affects the structured core domain. Substantial chemical shift perturbations

(CSPs) were observed for peaks in the core domain (CSPs above the 80th percentile

ranging from 0.032 to 0.119 ppm). If the N-terminus were flexible and structurally

independent from the core domain, the spectra would be expected to be virtually

identical. The CSPs appear as three clusters rather than randomly throughout the

sequence: (1) residues 21-24 are directly adjacent to the flexible N-terminus and are

thus expected to be affected when the latter is deleted, (2) residues 32-54 are in the

SIM-binding groove, and (3) residues from the 70s to the 80s (dubbed here the "70/80

region"). Together with the hNOE results, the observations strongly suggest that the

N-terminus interacts transiently with (at least) two specific regions of the core.

MD simulations provide insight into SUMO1 N-terminus - core

interactions

Given the timescale of the motions of the SUMO1 N-terminus as determined by

hNOE, we turned to all-atom molecular dynamics (MD) simulations to gain further

insights into the molecular mechanism by which SUMO1's N-terminus prevents SUMO

- SIM interactions. For extensive sampling of the conformational space accessible to

the disordered N-terminal region, we initiated the MD simulation with different

orientations of the N-terminus from the NMR structure for SUMO1 (see methods,

Table S1). We used an IDP-adequate force field, and the simulations were validated

by monitoring the Cα root-mean-square-deviation (RMSD) of the SUMO core against

the initial structures (< 3 Å) (Fig. S2). In addition, chemical shifts back-calculated from

MD simulations were in good accordance with the experimentally derived ones (Fig.

S3).

In MD simulations, we observed the disordered N-terminus to interact reversibly with

the SUMO core on the microsecond MD time scale, as measured by changes of the

binding area between the N-terminus and the core (Fig. S4A + S4B and Table S1).

The N-terminus of SUMO1 engaged with the core for about 65% of the simulation time

resulting in an average minor shielding of the total core area of only 1.4% (Fig. S5A +

S5B). Thus, the MD simulation, like the NMR experiments, points to a highly dynamic

interaction of the SUMO1 N-terminus with the core. Moreover, MD simulations

recapitulated findings by NMR by indicating that the SIM binding groove (residues

His35, Lys37 and Lys39 primarily affected) and the 70/80 region are temporarily

occupied by the SUMO1 N-terminus (Fig. 3C). Taken together, we conclude that MD

simulations are a suitable predictive *in silico* tool to study the N-terminus/core

interaction further.

MD simulations reveal significant differences of N-termini - core interactions across SUMO proteins

We next performed similar simulations for SUMO2, SMO-1 and Smt3. Again, we observed transient interactions of disordered N-termini of these paralogues with their respective cores on the microsecond MD time scale (Figs. 3D - G and S4C – S4E). However, comparison of all four SUMO proteins also revealed clear differences in average binding times and binding areas, with SMO-1 showing the longest binding time and largest binding area, followed by SUMO1, Smt3, and SUMO2 (Fig. 3G, Fig. S5A – S5D and Table S1). The most striking differences exist for the interaction with the SIM-binding groove, while interactions with the 70/80 region are rather comparable (Fig. 3C - F, compare Fig. S5B and 3G). These observations are consistent with a role of SUMO1 N-termini as intramolecular inhibitors of SUMO - SIM interactions, and show that this inhibitory effect is evolutionarily conserved in animals. The N-terminus of yeast Smt3, which is evolutionarily neither a SUMO1 nor SUMO2 protein (see [41] and above), shows an intermediate effect, while SUMO2's N-terminus does not seem to inhibit SUMO - SIM interactions, all in congruence with our biochemical data (see Figs. 1 and 2).

Interaction between SUMO's disordered N-termini and the SIM binding groove is highly dynamic

Intrinsically disordered proteins exhibit particularly fast binding kinetics when binding does not involve conformational changes such as secondary structure formation[52,53]. SUMO's N-termini form neither in the bound nor unbound states secondary structures in MD simulations consistent with PSIPRED secondary structure predictions[54] indicating low local secondary structure propensities (Fig. S6A). Moreover, conformational adaption of the N-termini upon binding was very minor (Fig. S6B). We thus wanted to test if the lack of structure also entails fast binding and unbinding of the

intrinsically disordered N-terminus to the SIM-binding groove. For this, we resorted to

the MD simulations. We assumed a simple two-state model of bound and unbound

states and determined the dwell times of all binding and unbinding events as

measured by changes in binding area between the N-terminus of SUMO and the SIM

binding groove (Fig. 3H and Fig. S5E). The MD-calculated off-rates k_{off} for SUMO1,

SUMO2, SMO-1, and yeast Smt3 range between 0.25 and 0.50 ns^{-1}, thus are

extremely high and hardly differ among the SUMOs. The on-rates vary by more than

an order of magnitude, with the lowest on-rate for SUMO2 of 0.02 ns^{-1} and the highest

on-rate for SMO-1 at 0.3 ns^{-1}. While k_{on} of SUMO2 and yeast Smt3 are significantly

lower than k_{off} resulting in an overall preference for the unbound state, the difference

between on- and off rates is much less pronounced for SUMO1 (0.15 versus 0.31 ns^{-1})

and for SMO-1 k_{on} is higher than k_{off} (0.31 versus 0.24 ns^{-1}), signifying a preference for

the bound state. In consequence, of the four N-termini, those of SUMO1 and SMO-1

possess the highest dwell time in the SIM-bound state in our simulations.

307 Acidic residues in SUMO's N-termini mediate the inhibitory effect on SIM-

308 dependent interactions

SUMO1 residues that interacted most prominently in MD simulations with SUMO1's N-

terminus included His35, Lys37, Lys39, Lys46, and Arg54 (see Fig. 1B, 3C and 4A),

suggestive of electrostatic interactions being vital for the interaction. All SUMO

proteins have negatively-charged amino acids in their N-termini, albeit at different

positions (see Fig. 1B, 4A, 4B and S6C). Most striking in SUMO1's N-terminus were

two neighboring negatively-charged residues, Glu11 and Asp12. Indeed, mutating

these residues to Lys individually or jointly to disrupt the N-terminus-SIM-binding

groove interaction strongly increased the interaction of SUMO1 with USP25 and with

TDP2 (Fig. 4C + 4D) as well as SIM-dependent SUMOylation (Fig. 4G + H).

Next, we tested whether our findings with human SUMO1 also hold true for *C. elegans*

SMO-1. Indeed, mutating Asp3 and Asp4 to Lys (DD3,4KK) in SMO-1 strengthens

interaction with the SIM-containing SUMO E3 ligase GEI-17 (Fig. 4E + F). Taken

together, our findings suggest that the intrinsically disordered N-termini of SUMO1

proteins inhibit SIM-dependent interactions and SUMOylation due to electrostatic

interactions of N-terminal acidic residues with basic residues in SUMO's core.

The SUMO1 N-terminus shields the SIM-binding groove via ionic

interactions

To substantiate interpretations from the biochemical assays described above, we

compared the HSQC-TROSY NMR spectrum of a [15]N-labeled SUMO1 ED11,12KK

variant with that of wild-type SUMO1. Strikingly, there are large CSPs in the entire N-

terminus and in the region of the core directly adjacent to the N-terminus (Fig. 5A).

The N-terminal CSPs are more widespread than would be predicted for amino acid

changes in an IDR, suggesting that the mutant N-terminus experiences a different

chemical environment to the wild-type. There are small, yet notable CSPs in the SIM-

binding groove. This finding confirms that the ED11,12KK mutations disrupt the

interaction between the N-terminus and the SIM-binding groove and stem. Consistent

with this notion, MD simulations of SUMO1 ED11,12KK reveal that the mutations

significantly decrease interaction with the SIM-binding groove (Fig. 5 B - D). Similar

behavior was observed comparing wt *C. elegans* SMO-1 with SMO-1 DD3,4KK (Fig.

4B, 5E - G). The slight increase in interactions with the 70/80 region seen in the MD

simulations of both mutants, however, can not be supported by the CSPs (Fig. 5A),

and is potentially caused by the known overstabilization of salt bridges in canonical

protein force fields[55]. Together, our findings indicate that the SUMO1 N-terminus acts

as a cis inhibitor of SUMO - SIM interactions due to its ability to mask the SIM binding

groove via dynamic ionic interactions.

345
346

The inhibitory function of SUMO1's N-terminus can be partially transplanted to SUMO2

The inhibitory effect of SUMO1's N-terminus seems largely driven by ionic interactions with residues flanking the SIM-binding groove (Fig. 4A + B). SIM-binding grooves are flanked by basic residues in all SUMO paralogues, raising the question of whether its effect could be transplanted onto SUMO2. We created several SUMO1/SUMO2 chimeras and tested them for interaction with USP25 in pulldown assays (Fig. 5H - I). Intriguingly, a SUMO2 chimera with SUMO1's N-terminus ($S1_N$-$S2_C$) showed reduced interaction. Moreover, a SUMO1 core that carries the SUMO2 N-terminus ($S2_N$-$S1_C$) bound USP25 more efficiently than wt SUMO1, albeit not as efficiently as SUMO2. In conclusion, SUMO N-termini are transplantable elements that contribute to the differences of SUMO proteins in SIM - dependent interactions.

Phosphorylation may regulate the inhibitory effect of SUMO's intrinsically disordered N-termini

Based on the ionic nature of the interaction, it is conceivable that posttranslational modifications that add or remove charges may influence inhibition. Numerous SUMO residues have been identified by proteomics as acetylated, phosphorylated, or methylated (summarized on phosphosite.org), some of which have been validated (e.g., pS2 in SUMO1[56], or AcK32 in SUMO2[57]).

To explore these ideas further, we turned again to MD simulations. We tested an array of modifications for SUMO1 (Fig. 6A - C and S7A – E), of which phosphorylation of Ser9 alone or in combination with Thr10 significantly increases the residence time of the SUMO1 N-terminus on the SIM-binding groove predicting that phosphorylation at these sites may further inhibit SIM-dependent interactions. Whether these predictions hold true and translate into a functional outcome needs to be tested experimentally in the future using post-translationally modified forms of SUMO.

For SUMO2 (Fig. 6D - F and S7F – S7J), MD simulations indicate a significant increase

in the overall residence time on the SIM-binding groove upon phosphorylation of Thr12.

Could the increased interaction of the N-terminus with the SIM-binding groove predicted

by MD simulation lead to a reduction of SIM-dependent interactions? In the absence of

quantitatively phosphorylated SUMO2, we turned to phospho-mimetic variants. Indeed,

mutating Thr12 in SUMO2's N-terminus either to Asp or Glu significantly decreased the

interaction of SUMO2 with USP25 (Fig. 6G + H). Together, these findings allow

speculation that the N-terminus of SUMO2 may act as an inhibitor of SUMO - SIM

interactions upon posttranslational modification(s).

In conclusion, we reveal a function for SUMOs' intrinsically disordered N-termini as

intramolecular regulators of SUMO-dependent interactions. N-termini of human and *C.*

elegans SUMO1 proteins are clearly inhibitory and other SUMO N-termini may acquire

such a function upon posttranslational modification of the N-terminus (Figure 6I).

Deletion of the flexible N-terminus in *C. elegans* SMO-1 reduces germ cell

survival

Finally, we wanted to address the biological significance of our findings. The N-terminus

of yeast Smt3 only shows an intermediate effect on SIM binding (our findings above),

serves as an acceptor for SUMO chains, and is dispensable for normal growth[40,41]. We

thus turned to *C. elegans* that expresses the single SUMO protein SMO-1. Its N-

terminus is not only a strong inhibitor of SIM interactions (our findings described

above), it is also devoid of lysines and can thus not contribute to canonical SUMO chain

formation.

We created an in-frame deletion allele by CRISPR/Cas9 genome editing in *C. elegans*

smo-1 (*smo-1(zh156)*, referred to as SMO-1ΔN12), removing 11 amino acids at the N-

terminus but leaving the start methionine*.* SMO-1ΔN12 animals were viable and

displayed no obvious anatomical defects or embryonic lethality. We could also not

detect significant differences in SUMO levels and overall SUMOylation in total animal

extracts (Fig. S8). However, SMO-1ΔN12 mutant hermaphrodites had a smaller brood

size (i.e. the average number of progeny per animal) than wild-type controls suggesting

a defect in germline development or oocyte fertilization (Fig. 7B). The *C. elegans*

sumoylation pathway is important for germ cell development and specifically, for

chromosome congression in germ cells at the pachytene stage of meiosis I[50,58,59].

Whole-mount DAPI staining revealed that the gonads of SMO-1ΔN12 mutants

contained a smaller pachytene region with fewer germ cell nuclei but a similar density

of nuclei (i.e. number of nuclei per area) as wild-type controls (Fig. 7A, C - E). This

phenotype could be due to decreased germ cell proliferation or to increased germ cell

apoptosis. Under physiological conditions, around half of the germ cells at the

pachytene stage undergo apoptosis, and the rate of apoptosis can be further increased

by introducing environmental stress such as DNA damage[60]. We therefore monitored

germ cell apoptosis in SMO-1ΔN12 mutants. First, we examined the CED-1::GFP

reporter, which outlines apoptotic corpses that are engulfed by the somatic sheath

cells[61], and observed an increased number of CED-1::GFP corpses in adult SMO-

1ΔN12 animals 24 and 48 hours post L4 stage despite the overall reduced number of

pachytene cells (Fig. 7F + 7G). As a second assay, we performed live-staining with

SYTO12, a dye that accumulates in germ cells undergoing apoptosis[61]. Also SYTO12

staining revealed an elevated number of apoptotic germ cells in adult SMO-1ΔN12

mutants 29 hours post L4 stage (Fig. 7H). To determine whether the increased levels of

apoptosis in the SMO-1ΔN12 mutant are due to elevated DNA damage, we used the

cep-1(gk138) p53 loss-of-function allele, which causes resistance to DNA-damage-

induced germ cell apoptosis. SMO-1ΔN12; *cep-1(gk138)* double mutants showed levels

of apoptosis comparable to WT or *cep-1(gk138)* single mutants (Fig. 7G)*.* Thus*,* the

increased apoptosis levels in SMO-1ΔN12 mutants depend on the DNA damage

response.

Together, these data indicate that the flexible N-terminus of *C. elegans* SMO-1 is

required for normal germline development and in particular for the survival of germ cells

entering the pachytene stage of meiosis.

DISCUSSION

SUMO N-termini are Intrinsically Disordered Regions

Intrinsically disordered proteins and protein regions (IDP/IDR) have been largely

overlooked until the 1990s even though they constitute 40% of the proteome of

eukaryotes[62] and fulfill many important biological functions including signal

transduction[63,64]. Both their mechanism of action and biological functions vary greatly.

For example, IDP/IDRs may function as (auto)inhibitors of enzymes or act as assembly

platforms for multi-subunit complexes (reviewed in [65]).

Unstructured N-terminal regions of 13 - 23 amino acids are a hallmark of all SUMO

proteins (Fig. 1B, Table S1) and distinguishes them from Ubiquitin and most other Ubl

relatives. These feature the expected amino acid composition of IDRs[66], namely a

relatively low content of hydrophobic amino acids along with a higher ratio of polar or

charged amino acids (Figure 4A + 4B, and S6). The disordered and flexible nature of

SUMO N-termini function has remained rather enigmatic. Some SUMO N-termini

engage in chain formation (e.g., [67,68]). Additionally, SUMO N-termini have been

suggested to act as entropic bristles that may prevent aggregation, but SUMO

aggregation has only been observed upon extended incubation at very high

temperature[69]. Here, we discovered a rapid nanosecond equilibrium between "open"

SUMO conformations available for interactions with SIMs and "closed" ones that are

cis-inhibited by the N-terminal region. The N-terminal region remains fully disordered in

the bound state during our MD simulations and is thus a classic example of intrinsic

disorder irrespective of the binding state.

456   **SUMO N-termini form fuzzy complexes with SUMOs' SIM binding grooves**

457   How specificity is achieved by an IDR at the molecular level is an important question.

458   Upon binding to their biological partners, IDRs often undergo a conformational

459   transition to a folded state, also referred to as coupled binding and folding[70]. A well-

460   known example is the cell cycle inhibitor p27, which folds around cyclin / Cdk

461   complexes[71]. Nevertheless, increasing evidence demonstrates that at least some IDRs

462   remain unstructured upon interaction with partners, which is referred to as formation of

463   a "fuzzy complex"[72]. Interactions in such a complex are mediated by transient

464   hydrophobic and electrostatic interactions in the fuzzy region, which can provide non-

465   specific contacts or specific interactions with multiple motifs[53]. Examples include the

466   fuzzy complex of transcriptional activator Gcn4 and its partner Gal11/Med15[73], ultrafast

467   interaction between FG-repeat containing nucleoporins and nuclear transport

468   receptors[74], or the fuzzy intramolecular binding between the IDR and folded domain of

469   Src kinase[75] or galectin[76].The interaction mode of SUMOs' N-termini with the SUMO

470   core is in line with the fuzzy complex model. First, we observed in MD simulations a

471   very fast (ns) rate of the binding / unbinding process, and secondly, the bound state of

472   the SUMO N-terminus and the SUMO core is transient and dynamic. Although the N-

473   terminus hovers specifically over the SIM binding groove, it does not assume a specific

474   stable configuration but rather forms a "ligand cloud" (Fig. S6B)[77]. Despite this fuzzy

475   nature, the N-terminus can compete efficiently with SIM-containing proteins for the SIM

476   binding groove, as supported by both experiments and simulations.

477

478   **SUMO N-termini are cis-inhibitors of SIM-dependent interactions**

479   The N-termini of recombinant SUMO1, SMO-1 and Smt3 act as inhibitors of SIM

480   dependent interactions (this study,[42]), and potentially also of interactions with other

481   SUMO surfaces, namely the 70/80 region. A recent study by Lussier-Price et al. also

482   reported on a paralogue-specific auto-inhibition of SIM binding by SUMO1's N-

483   terminus[42]. Similar to our study in the outcome but with a different experimental

approach, they show strongly reduced K_Ds for the phosphorylated SIMs of PML and

Daxx upon deletion of SUMO1's but not SUMO2's N-terminus and shielding of a

hydrophobic area on SUMO1 by the N-terminus that can be competed off by a

phosphorylated SIM peptide of PML.

Inhibition via a module within the same protein (cis-inhibition) plays an important role in

regulation[78]. IDRs are frequently involved in cis-inhibition, possibly as a way to

economize genome/protein resources[65]. A consequential benefit of this mechanism is

the ease of regulation via posttranslational modifications[79]. This is likely the case for

SUMOs' N-termini because the interaction of SUMOs' IDRs with the SIM-binding

grooves is largely mediated by electrostatics, which may be strengthened or weakened

by a host of posttranslational modifications that are emerging for SUMO proteins both in

the IDR (e.g., [56,68] or flanking the SIM binding groove[34,57]). Here we provide the first

evidence for this idea from MD simulations and pulldown assays (Fig. 6 and Fig. S7).

An important implication from our analyses is that the SUMO2 N-terminus may become

a cis-inhibitor upon phosphorylation of Thr12.

SUMO N-termini add control to SIM-dependent interactions

SIM-dependent interactions contribute to all aspects of the SUMO pathway yet how

specificity can be ensured with a single SUMO binding module is still unclear.

Increasing evidence suggests regulation of SUMO - SIM interactions at multiple levels.

For example, acetylation of lysine residues surrounding the SIM binding groove can

alter the affinity for specific binding partners (see, e.g., [57,80]), while phosphorylation of

residues in a given SIM may increase the affinity for SUMO and change SUMO

paralogue preferences (e.g., [33,35]). Our findings add yet another layer of regulation:

whether SUMO on a given target is available for interaction with a downstream effector

can be controlled via SUMO's intrinsically disordered N-terminus (model in Fig. 6I). Our

analysis of *C. elegans* SMO-1 points to a specific function of the SUMO N-terminus in

germ cell survival (Fig. 7). Whereas complete loss of SMO-1 function causes pleitropic

effects in the germline and soma, affecting a variety of cell fates, deletion of SMO-1's

N-terminus reveals increased p53-dependend germ cell apoptosis, suggesting a

specific role of the SMO-1 N-terminus in controlling DNA damage-induced apotosis.

These findings are consistent with previous reports showing that the SUMO system in

C. elegans plays an essential role in sensing environmental stress and maintaining

genome stabiblity[81]. Taken together, our findings offer an intriguing explanation for the

presence and strict evolutionary conservation of SUMOs' intrinsically disordered N-

termini: they seem to equip SUMO with an environment-sensing and regulatory

capacity of SIM-dependent processes including SIM-dependent modification and

downstream effector binding.

Limitations of the study

Here we assigned an important function to SUMO1 and Smo-1 N-termini as inhibitors of

SUMO-dependent interactions, and provide an explanation for paralogue preferences

of SUMO targets such as USP25. An exciting question that we discuss is whether the

inhibitory potential of different SUMO - N-termini can be tuned by posttranslational

modifications of the N-termini or the SUMO core (model in Fig. 6I). While our MD

simulations and *in vitro* studies with selected mutants point in this direction, we have

not been able to generate quantitatively acetylated and/or phosphorylated SUMO

variants to test this hypothesis. An important question for future studies will be the

identification of pathways and protein targets for which this specialized function of

SUMO's N-terminus is most relevant. Although Smt3's N-terminus can be deleted

without detectable effect on growth[41], at least in standard laboratory conditions, we are

not aware of any eukaryotic organism with a SUMO protein that lacks the flexible N-

terminus. In light of our findings *in C. elegans,* were deletion of Smo's N-terminus

increases p53-dependent cell death, we speculate that stress response pathways that

provide populations with evolutionary advantages are the right places to look for.

STAR+METHODS

Detailed methods are provided in the online version of this paper

and include the following:

KEY RESOURCES TABLE

Lead contact and materials availability statement

METHOD DETAILS

SUMO nomenclature

Expression constructs

Protein expression and purification

In vitro binding assays

In vitro SUMOylation assays

Quantification and Statistical Analysis

Structure preparation and Molecular Dynamics (MD) simulation system setup

Energy minimization, equilibration and molecular dynamics simulations

Analyses of molecular dynamics simulation

STAR+METHODS

Key Resources Table

Contact for Reagent and Resource Sharing

Further information and requests for reagents may be directed to, and will be fulfilled by

the corresponding author Frauke Melchior (f.melchior@zmbh.uni-heidelberg.de).

REAGENT or RESOURCE	SOURCE	IDENTIFIER
Antibodies		
rabbit anti-USP25	Meulmeester et al.[15]	N/A
goat anti-RanGAP1	Hutten et al.[82]	N/A
mouse anti-SMO-1	Pelisch et al.[58]	AB_2618393
Bacterial and Virus Strains		
E. coli BL21 (DE3)	Stratagene	N/A

C. elegans Strains		
AH6024	This work	N/A
WS4116	Zhou et al.[83]	N/A
AH6119	This work	N/A
AH5891	Hajnal lab	N/A
AH6143	This work	N/A
VC172	CGC[84]	N/A
Chemicals, Peptides, and Recombinant Proteins		
Cyanogen bromide-activated-Sepharose 4B	Sigma-Aldrich	C9142
SUMO1 wt	This work	N/A
SUMO1ΔN10	This work	N/A
SUMO1ΔN19	This work	N/A
SUMO1ΔN19-VK38,39AA	This work	N/A
SUMO1-ED11,12KK	This work	N/A
SUMO1-E11K	This work	N/A
SUMO1-D12K	This work	N/A
SUMO1-Gst	This work	N/A
SUMO1ΔN19-Gst	This work	N/A
SUMO1-V38A,K39A-Gst	This work	N/A
SUMO1ΔN19-VK38,39AA-Gst	This work	N/A
SUMO2 wt	This work	N/A
SUMO2ΔN13	This work	N/A
SUMO2-IK33,34AA	This work	N/A
SUMO2-T12D	This work	N/A
SUMO2-T12E	This work	N/A
SUMO2-Gst	This work	N/A
$S1_N$-$S2_C$-Gst	This work	N/A
$S2_N$-$S1_C$-Gst	This work	N/A
Smt3 wt	This work	N/A
Smt3ΔN18	This work	N/A
SMO-1 wt	This work	N/A
SMO-1ΔN12	This work	N/A
SMO-1 DD3,4KK	This work	N/A
USP25 wt	This work	N/A
USP25-VI91,92AA	This work	N/A
Ovalbumin	Sigma Aldrich	N/A
Gst	This work	N/A
TDP2 wt	This work	N/A
Ubc9 wt	This work	N/A
RanBP2ΔFG (aa 2553–2838)	This work	N/A
RanGAP1 wt	This work	N/A
Siz1(313-508)	This work	N/A
6xHis-TEV-GEI-17(133-509)	This work	N/A
Recombinant DNA		
pET11a-SUMO1 wt	Pichler et al. 2002[44]	N/A
pET11a-SUMO1 ΔN10	This work	N/A
pET11a-SUMO1 ΔN19	This work	N/A
pET11a-SUMO1 ΔN19-V38A, K39A	This work	N/A
pET11a-SUMO1-ED11,12KK	This work	N/A
pET11a-SUMO1-E11K	This work	N/A
pET11a-SUMO1-D12K	This work	N/A

pET23-SUMO1-Gst	This work	N/A
pET23-SUMO1ΔN19-Gst	This work	N/A
pET23-SUMO1-V38A, K39A-Gst	This work	N/A
pET23-SUMO1ΔN19-V38A, K39A-Gst	This work	N/A
pET11a-SUMO2 wt	Meulmeester et al. 2008[15]	N/A
pET11a-SUMO2 ΔN13	This work	N/A
pET11a-SUMO2-IK33,34AA	Meulmeester et al. 2008[15]	N/A
pET11a-SUMO2-T12D	This work	N/A
pET11a-SUMO2-T12E	This work	N/A
pET23-SUMO2-Gst	This work	N/A
pET23-S1$_N$-S2$_C$-Gst	This work	N/A
pET23-S2$_N$-S1$_C$-Gst	This work	N/A
pET11a-Smt3 wt	This work	N/A
pET11a-Smt3 ΔN18	This work	N/A
pET11a-SMO-1 wt	This work	N/A
pET11a-SMO-1ΔN12	This work	N/A
pET11a-SMO-1 DD3,4KK	This work	N/A
pET11a-His-TEV-USP25	Meulmeester et al. 2008[15]	N/A
pET11a-His-TEV-USP25-VI91,92AA	Meulmeester et al. 2008[15]	N/A
pGEX-6P-3-TDP2	This work	N/A
pET23a-Ubc9	Pichler et al. 2002[44]	N/A
pGEX3X-RanBP2ΔFG (aa 2553–2838)	Pichler et al. 2002[44]	N/A
pET11a-hRanGAP1	Werner et al. 2012[85]	N/A
pET21-Siz1(313-508)	Johnson and Gupta 2001[86]	N/A
6xHis-TEV-GEI-17(133-509)	Pelisch et al. 2014[58]	N/A

Software and Algorithms		
Microsoft Excel for Mac 2011 Vers.14.3.6	Microsoft Corporation	N/A
ImageJ 10.2	Wayne Rasband, National Institutes of Health, USA	N/A
Serial Cloner Version 2.6.1	Franck Perez	N/A
Clustal Omega multiple sequence alignment (web tool)	The European Bioinformatics Institute	N/A
PyMol v1.7.4.5 Edu enhanced for MacOS X	Schrödinger LLC.	https://www.pymol.
LAS-400 2.1	Fujifilm	N/A
Pytms	Warnecke et al. 2014[87]	https://pymolwiki.o
GROMACS v. 2016.3	Van der Spoel et al. 2005[88]	http://www.gromac
UCSF Chimera	Pettersen et al. 2004[89]	https://www.cgl.uc
MODELLER	Webb and Sali 2014[90]	https://salilab.org/

Method Details

SUMO nomenclature

The nomenclature for mammalian SUMO2 and SUMO3 is used inconsistently. Like

many colleagues in the SUMO field, we follow the nomenclature as introduced by [91].

Their assignment was consistent with the original description of mammalian SUMO

genes (reviewed in [3]). According to this, mature SUMO2 (Smt3A) is 92 amino acids

long, mature SUMO3 (Smt3B) consists of 93 amino acids.

Expression constructs

Bacterial expression constructs for untagged SUMO1(ΔC4) and SUMO2(ΔC11),

USP25, USP25-V91A, I92A[15], GST-RanBP2 ΔFG, Ubc9, SAE1 (Aos1), SAE2 (Uba2)[44]

and RanGAP1[85] were previously described. All constructs but one were based on

human cDNA sequences. The Ubc9 construct was derived from mouse cDNA, but the

encoded mouse protein is identical to human Ubc9. The coding sequence of mature

full-length *S. cerevisiae* Smt3 (Smt3ΔC3) and *C. elegans* SMO-1 (SMO-1ΔC1) were

PCR-amplified and cloned into the NdeI and BamHI sites of pET11a. The coding

sequence of mature full-length human SUMO1 and SUMO2 was PCR-amplified and

Gibson-cloned into the SacI and NotI sites of pET23a; GST was fused to the C

terminus of SUMO by PCR-amplification of GST and cloning into the NotI and XhoI

sites. SUMO1ΔN19 and SUMO2ΔN13 were PCR-amplified and cloned into the NdeI

and BamHI sites of pET11a. Other N-terminal deletions and amino acid mutations of

SUMO-, Smt3-, and SMO-1 plasmids were performed using the QuikChange site-

directed mutagenesis method (Agilent) or by Gibson DNA assembly. For the

SUMO1/SUMO2 hybrid constructs, the SUMO1 and SUMO2 N- and C-termini (SUMO1

N: aa 1-20, C: aa 21-97; SUMO2 N: aa 1-15, C: aa 16-92) were PCR-amplified from

the C-terminal GST fusion constructs including roughly half of the vector backbone

each and fused in the desired combinations by Gibson DNA assembly. The coding

sequence of human TDP2 was PCR-amplified from the pENTR221-TDP2 plasmid

(genomics and proteomics core facility, DKFZ) and cloned into the BamHI and XmaI

sites of the pGEX-6P-3 vector. The expression plasmid for the catalytic fragment of

Siz1(313-508)-pET21 was a kind gift by Dr. Erica S. Johnson[86]. 6xHis-TEV-GEI-

17(133-509) was a gift from Ronald Hay[58] (Addgene plasmid #87056;

http://n2t.net/addgene:87056 ; RRID:Addgene_87056). The sequences of all

expression constructs used in this work were verified by Sanger sequencing.

Protein expression and purification

SUMO proteins

Untagged SUMO1-, SUMO2-, Smt3- and SMO-1 variants were essentially purified

according to an established protocol[92]. *E. coli* BL21 (DE3) cells were transformed with

the respective pET11a-SUMO plasmid and a single colony was grown over night in LB

medium supplemented with ampicillin. The cells were centrifuged for 5 min at 3900 g

and resuspended in fresh LB medium supplemented with ampicillin or MOPS medium

supplemented with (^{15}N)ammonium chloride and ampicillin (for NMR). When the main

culture reached an OD_{600} of 0.6 - 0.8, expression of SUMO variants was induced with 1

mM IPTG and expressed for 3 - 4 h at 37°C or overnight at 16°C . Cells were harvested

by centrifugation for 10 min at 7200 g, resuspended in 50 mM Tris–HCl pH 8.0, 50 mM

NaCl, 1 mM DTT, supplemented with 1 µg/ml of each aprotinin, leupeptin, pepstatin A

and flash frozen in N2(l). After thawing, the cell suspension was lysed using an

EmulsiFlex-C5 (Avestin) in the presence of DNase I and cleared by ultra-centrifugation

for at least 45 min at 100,000 g and 4 °C. The supernatant was incubated with 10 ml

DEAE- or Q-Sepharose for at least 1 h at 4 °C, the sepharose material was sedimented

and the supernatant was concentrated using a centrifugal filter unit with 5 kDa MWCO.

The protein was then purified by gel filtration over a Superdex200 column equilibrated

in Transport Buffer (20 mM HEPES-KOH pH 7.3, 110 mM KAcO, 2 mM Mg(AcO)2, 1

mM EGTA, 1 mM DTT) supplemented with 1 µg/ml of each aprotinin, leupeptin,

pepstatin A (TB+++). In some but not all SUMO preparations, an additional gel filtration

run in 50 mM Tris pH 7.5, 1M LiCl, 150 mM NaCl, 1 mM DTT, 1 µg/ml of each aprotinin,

leupeptin, pepstatin A was included before the gel filtration in TB+++. SUMO containing

fractions were pooled, concentrated and applied to a Superdex75 gel filtration column

equilibrated in TB+++. SUMO containing fractions were pooled, concentrated, flash-

frozen in N2(l) and stored at -80 °C.

All SUMO-GST proteins were expressed from the respective pET23a-SUMO-GST

plasmids and lyzed in 50 mM Tris-HCl pH 8.0, 100 mM NaCl, 1 mM EDTA, 1 mM

EGTA, 1 mM CaCl2, 1 mM MgCl2, 1 mM DTT, 1 µg/ml of each aprotinin, leupeptin,

pepstatin A, Pefa in the presence of DNAase I as above, and the protein in the

supernatant after ultracentrifugation was purified by pull-down with glutathione agarose

equilibrated in GST lysis buffer. Bound protein was washed with GST wash buffer (50

mM Tris-HCl pH 8.0, 300 mM NaCl, 1 mM EDTA, 1 mM DTT, 1 µg/ml of each aprotinin,

leupeptin, pepstatin A) and eluted with GST wash buffer supplemented with 20 mM

glutathione. SUMO-GST containing fractions were pooled, concentrated and applied to

a Superdex75 gel filtration column equilibrated in TB+++. SUMO-GST containing

fractions were pooled, diluted 1:2 with 50 mM Tris pH 7.5, bound to a MonoQ column

equilibrated in 50 mM Tris pH 7.5. Bound protein was eluted with a gradient of 50 mM –

1 M NaCl in 50 mM Tris pH 7.5. SUMO-GST containing fractions were pooled,

concentrated, flash-frozen in N2(l) and stored at -80 °C.

GST-tagged TDP2 and RanBP2ΔFG

For expression of N-terminally GST-tagged TDP2 and RanBP2ΔFG, *E. coli* BL21 (DE3) cells were transformed with the pGEX-6P-3-TDP2 and the pGEX-3X-RanBP2ΔFG plasmid, respectively, and a single colony was grown over night in LB medium supplemented with ampicillin. The cells were centrifuged for 5 min at 3900 g and resuspended in fresh LB medium supplemented with ampicillin. When the main culture reached an OD600 of 0.6 - 0.8, it was shifted to 16 °C, expression was induced with 0.5 mM IPTG and expressed for 21 h at 16 °C. Cells were harvested by centrifugation for 10 min at 7200 g, resuspended in GST lysis buffer and flash frozen in N2(l). After thawing, the cell suspension was lysed using an EmulsiFlex-C5 and cleared by ultra-centrifugation for at least 45 min at 100,000 g and 4 °C. The GST-tagged protein in the supernatant was then purified by pull-down with glutathione-sepharose equilibrated in GST lysis buffer. For this, the supernatant was incubated with 10 ml glutathione sepharose / l main culture for at least 1 h at 4 °C. The bead material was then collected in a column and washed with 50 CV GST lysis buffer. Elution was performed with GST elution buffer. Protein-containing fractions were pooled, concentrated and applied to a Superdex200 gel filtration column equilibrated in TB+++. Protein containing fractions were pooled, concentrated, flash frozen in N2 (l) and stored at - 80 °C.

USP25

His-TEV-USP25 was transformed into *E. coli* BL21 (DE3) and a single colony was grown over night in LB medium supplemented with ampicillin. The cells were centrifuged for 5 min at 3900 g, resuspended in fresh LB medium supplemented with ampicillin and grown at 37 °C. When the main culture reached an OD600 of 0.6 - 0.8, it was shifted to 30 °C, USP25 expression was induced with 0.5 mM IPTG and expressed for 4 h. Cells were harvested by centrifugation for 10 min at 7200 g, resuspended in 50 mM Na-phosphate pH 8.0, 300 mM NaCl, 10 mM imidazole, 1 mM DTT, 1 μg/ml of each aprotinin, leupeptin, pepstatin A and flash-frozen in N2 (l). The cells were lysed using an EmulsiFlex-C5. The supernatant after 100,000 g ultracentrifugation was incubated with Ni-NTA Sepharose for 2 h at 4 °C and eluted with 50 mM Na-phosphate pH 8.0, 300 mM NaCl, 200 mM imidazole, 1 mM DTT, 1 μg/ml of each aprotinin, leupeptin, pepstatin A. Protein-containing fractions were pooled, concentrated and applied to a Superdex200 gel filtration column equilibrated in TB+++. USP25 in the eluate was pooled, concentrated, flash-frozen in N2(l) and stored at -80 °C.

RanGAP1

Untagged RanGAP1 was expressed and purified according to an established
protocol[93]. *E. coli* BL21 (DE3) cells were transformed with pET11a-RanGAP1 and a
single colony was grown over night in LB medium supplemented with ampicillin. The
cells were centrifuged for 10 min at 5000 g and resuspended in fresh LB medium
supplemented with ampicillin, 1 mM MgCl2 and 0.1 % glucose. When the main culture
reached an OD600 of 0.8 - 0.9, RanGAP1 was induced with 0.5 mM IPTG and
expressed for 3 h at 37 °C. Cells were harvested by centrifugation for 15 min at 5000 g,
resuspended in 50 mM Tris–HCl pH 8.0, 100 mM NaCl, 1 mM EDTA, 1 mM DTT,
supplemented with 1 μg/ml of each aprotinin, leupeptin, pepstatin A and flash frozen in
N2(l). After thawing, the suspension was supplemented with 1 mg/ml lysozyme,
incubated for 1 h on ice and lysed by sonication (from this point all buffers
supplemented 1 μg/ml of each aprotinin, leupeptin, pepstatin A). The pellet after
centrifugation for 30 min at 20,000 g was washed by resuspension using a glas
douncer and centrifugation successively in 50 mM Tris–HCl pH 8.0, 1% Triton X-100,
and in 50 mM Tris–HCl pH 7.4, 2 M urea. The final pellet was solubilized in 50 mM
Tris–HCl pH 7.4, 8 M urea. The supernatant was dialyzed against 50 mM Tris–HCl pH
7.4, 100 mM NaCl with two buffer exchanges, centrifuged at 100,000 g for 1 h, applied
to a Q–Sepharose (GE Healthcare) column equilibrated with 50 mM Tris–HCl pH 7.4,
150 mM NaCl, washed with 50 mM Tris–HCl pH 7.4, 300 mM NaCl and eluted with 50
mM Tris–HCl pH 7.4, 1 M NaCl. Protein fractions were then applied to a HiPrep- 16/60
Sephacryl S-300 HR (GE Healthcare) column equilibrated in TB+++ and RanGAP1-
containing fractions were concentrated, pooled, flash-frozen in N2 (l) and stored at -80
°C.

SUMO E1 activating enzyme (SAE1-SAE2 heterodimer)
Expression and purification of the SAE1-SAE2 heterodimer followed an established
protocol[44]. *E. coli* BL21 (DE3) cells were co- transformed with pET28a-Aos1 (His-SAE1)
and pET11d-Uba2 (SAE2) and grown for 18 h at 37 °C in 500 ml LB medium
supplemented with kanamycin and ampicillin. After addition of 1.5 l of LB medium
supplemented with kanamycin and ampicillin, protein expression was induced with 1
mM IPTG and performed for 6 h at 25 °C. Cells were harvested by centrifugation for 15
min at 5000 g, resuspended in E1 lysis buffer (50 mM Na-phosphate pH 8.0, 300 mM
NaCl, 10 mM imidazole) and flash-frozen in N2(l).
After thawing, the respective cell suspensions were supplemented with 1 μg/ml of each
aprotinin, leupeptin, pepstatin A and 1 mM DTT. Cells were lysed using an EmulsiFlex-
C5. The supernatant after ultracentrifugation at 100,000 g for 1 h was incubated with
Ni-NTA Sepharose equilibrated with E1 lysis buffer for 1 h at 4 °C, washed thoroughly

with E1 wash buffer (E1 lysis buffer with 20 mM imidazole) and eluted with E1 elution
buffer (E1 lysis buffer with 250 mM imidazole). Protein-containing fractions were
pooled, concentrated and applied to a Superdex200 gel filtration column equilibrated in
TB+++. Eluate fractions were checked for the presence of both proteins, His-SAE1 and
SAE2, by SDS-PAGE. The respective fractions were pooled and applied to a MonoQ
5/50 GL column (GE Healthcare) equilibrated in 50 mM Tris–HCl, pH 7.5, 50 mM NaCl,
2 mM DTT, 1 μg/mL of each aprotinin, leupeptin, pepstatin. The E1-complex was eluted
by applying a linear gradient from 50 to 500 mM NaCl in 20 column volumes. Collected
fractions were analyzed by SDS-PAGE and complex containing fractions were pooled,
dialyzed against TB+++ and flash-frozen in N2(l).

SUMO E2 conjugating enzyme Ubc9

Expression and purification of Ubc9 followed an established protocol[92]. *E. coli* BL21
(DE3) cells were transformed with pET23a-Ubc9 and a single colony was grown over
night at 37 °C in LB medium supplemented with ampicillin, 1 mM MgCl2 and 0.1 %
glucose. The cells were centrifuged for 10 min at 5000 g and resuspended in fresh LB
medium supplemented with ampicillin, 1 mM MgCl2 and 0.1 % glucose. When the main
culture reached an OD600 of 0.6, Ubc9 expression was induced with 1 mM IPTG and
expressed for 3 h at 37 °C. Cells were harvested by centrifugation for 15 min at 5000 g,
resuspended in Ubc9 lysis buffer (50 mM Na-phosphate buffer, pH 6.5 supplemented
with 1 mM DTT, 1 μg/ml of each aprotinin, leupeptin, pepstatin A) and flash-frozen in
N2(l). After thawing, the lysate was cleared by 100,000 g ultracentrifugation at 4 °C and
applied to an SP-Sepharose column equilibrated in Ubc9 lysis buffer. The column
washed extensively using Ubc9 lysis buffer and Ubc9 eluted with Ubc9 lysis buffer
supplemented with 300 mM NaCl. Protein-containing fractions were applied to a
Superdex75 gel filtration column equilibrated in TB+++. Ubc9 in the eluate was pooled,
concentrated, flash-frozen in N2(l) and stored at -80 °C.

SIM-containing fragment of the SUMO E3 ligase *S. c.* Siz1 (aa 313-508)

Siz1(313-508)-pET21 was transformed into *E. coli* BL21 (DE3) and a single colony was
grown over night in LB medium supplemented with ampicillin. The cells were
centrifuged for 5 min at 3900 g, resuspended in fresh LB medium supplemented with
ampicillin and grown at 37 °C. When the main culture reached an OD600 of 0.6 - 0.8,
Siz1 expression was induced with 1 mM IPTG and expressed for 4 h. Cells were
harvested by centrifugation for 10 min at 7200 g, resuspended in 50 mM Na-phosphate
pH 8.0, 300 mM NaCl, 10 mM imidazole, 1 mM DTT, 1 μg/ml of each aprotinin,

leupeptin, pepstatin A and flash-frozen in N2 (l). After thawing, the cells were lysed

using an EmulsiFlex-C5. The supernatant after 100,000 g ultracentrifugation was

incubated with Ni-NTA Sepharose for 2 h at 4 °C and eluted with 50 mM Na-phosphate

pH 8.0, 300 mM NaCl, 200 mM imidazole, 1 mM DTT, 1 μg/ml of each aprotinin,

leupeptin, pepstatin A. Protein-containing fractions were pooled, concentrated and

applied to a Superdex75 gel filtration column equilibrated in TB+++. Siz1 in the eluate

was pooled, concentrated, flash-frozen in N2 (l) and stored at -80 °C.

SIM-containing fragment of the SUMO E3 ligase *C. e.* GEI-17 (aa 133-509)

Rosetta2 (DE3) cells transformed with 6xHis-TEV-GEI-17 were grown in LB medium

supplemented with ampicillin, 1 mM MgCl$_2$, 0.1% glucose at 37°C. Expression was

induced with 0.1 mM IPTG for 16 h at 20°C. The cells were harvested, resuspended in

lysis buffer (50 mM Tris-HCl pH 7.5, 500 mM NaCl, 10 mM imidazole, 0.1% Triton-

X100, 1 mM DTT, 1 μg/ml of each aprotinin, leupeptin, pepstatin A, Pefa bloc) and

flash-frozen in N2 (l). After thawing, the cells were lysed using an EmulsiFlex-C5. After

100,000 g ultracentrifugation the supernatant was passed over a Ni-NTA sepharose

column at 4 °C, the column was first washed with lysis buffer, then with 50 mM Tris-HCl

pH7.5, 500 mM NaCl, 30 mM Imidazole, 1 mM DTT, 1 μg/ml of each aprotinin,

leupeptin, pepstatin A. Bound protein was eluted with 50 mM Tris-HCl pH 7.5, 150 mM

NaCl, 200 mM imidazole, 1 mM DTT, 1 μg/ml of each aprotinin, leupeptin, pepstatin A.

Protein-containing fractions were pooled, concentrated and applied to a Superdex200

gel filtration column equilibrated in TB+++. GEI-17-containing elution fractions were

pooled, concentrated, flash-frozen in N2 (l) and stored at -80 °C.

***In vitro* binding assays**

For the preparation of SUMO sepharose, SUMO1, SUMO2, Smt3 or SMO-1

concentrations were determined thoroughly by absorption at 280 nm using the protein-

specific absorption coefficients, and by Bradford assay. The concentration

measurements were verified by SDS-PAGE followed by Coomassie staining.

Importantly, we only compared SUMO proteins prepared in the same way, either

including or excluding the additional LiCl purification step in one experiment. Equal

amounts of the untagged SUMO proteins were coupled to Cyanogen bromide-

activated-Sepharose in Carbonate buffer (0.2 M pH 8.9) at a concentration of 1 mg

protein per ml beads[15]. To block remaining coupling sites, beads were subsequently

incubated with 100 mM Tris-HCl pH 8.0. Beads were washed several times with PBS

supplemented with 500 mM NaCl then with PBS only. Beads were stored in Transport

Buffer (TB: 20 mM HEPES/KOH pH7.3, 110 mM potassium acetate, 2 mM magnesium
acetate, 0.5 mM EGTA, 1 mM DTT and protease inhibitors) supplemented with 1 mM
sodium azide.

Equal amounts of the SUMO-GST variants were bound to Glutathione sepharose in
PBS supplemented with 0.05% Tween20 at a concentration of 1 mg/ml for 1 h at 4°C.
Beads were washed several times with PBS supplemented with 0.05% Tween 20.

The coupling efficiency of all SUMO beads was always controlled by comparing the
protein content of the coupling reaction before and after coupling.

For binding assays, 10 μg of recombinant binding partner per 300 μl reaction was
incubated with 10 - 20 μl (USP25, TDP2, RanBP2, Siz1, GEI-17) or 5 - 7.5 μl (Ubc9)
SUMO beads in TB supplemented with 0.05% Tween 20, and 0.2 mg/ml ovalbumin for
at least 1 h at 4 °C. Beads were washed three times with TB or TB supplemented with
0.05% Tween 20 and eluted with 2x SDS sample buffer. The samples were analyzed
by SDS-PAGE followed by Coomassie staining. The gels were scanned using the gel
documentation system LAS-4000 (GE Healthcare) and the respective protein
containing bands quantified with ImageJ.

***In vitro* SUMOylation assays**

In vitro SUMOylation assays with purified recombinant proteins were essentially
performed as described previously[92]. Reactions containing USP25 were prepared with
100 nM USP25, 50 nM SAE1/SAE2 (E1 enzyme), 500 nM Ubc9 (E2 enzyme), 5 μM
SUMO or SMO-1 variants, and 1 or 5 mM ATP in TB supplemented with 0.05% Tween
20 and 0.2 mg/ml ovalbumin and incubated at 30 °C for up to 80 minutes. Reactions
containing RanGAP1 were prepared with 500 nM RanGAP1, 10 nM SAE1/SAE2, 10
nM Ubc9, 5 μM SUMO or SMO-1 variants, and 1 or 5 mM ATP and incubated at 30 °C
for up to 30 minutes. Reactions were stopped by adding SDS sample buffer and
incubation at 95 °C for 5 min. Samples were analyzed by SDS-PAGE followed by
immunoblotting.

Quantification and Statistical Analysis of biochemical experiments

The amount of SUMO binding partners in the eluates of SUMO or Smt3 binding
reactions were quantified relative to the respective Ubc9 or SUMO-GST signal, if
applicable, using ImageJ[94]. Data were analyzed using a paired Student's t-test in Excel.
Error bars represent one standard deviation. The number of independent experiments
(n) is given in the respective figure legend. Asterisks indicate p-values: *: P ≤ 0.05; **: P
≤ 0.01; ***: P ≤ 0.001; n.s.: not significant.

 NMR spectroscopy

NMR spectra were recorded on a 500 MHz Bruker Avance II (University of Washington)

at 295 K in 25 mM sodium phosphate pH 7.0, 150 mM NaCl, 10% D2O. ^{1}H/^{15}N-HSQC-

TROSY and hNOE experiments were collected with 370 µM labeled protein (700 µM for

SUMO1 ΔN19). Datasets were processed using NMRPipe/NMRDraw[95], and visualized

with NMRView[96]. Chemical shift perturbations observed by 2D HSQC-TROSY NMR

were quantified in parts per million with the equation $\Delta\delta(^1H/^{15}N) = \{[\delta(^1H) - \delta(^1H)_0]^2 +$

$0.04[\delta(^{15}N) - \delta(^{15}N)_0]^2\}^{1/2}$.

Structure preparation and Molecular Dynamics (MD) simulation system setup

The initial structures for wild type SUMO1 were selected from NMR structures with the

PDB codes 1A5R[39] and 2N1V[97], and for mature wild type SUMO2 (92 aa) from an NMR

structure with the PDB code 2RPQ[38] without any preference for the distance between

the N-terminal tail and the main body of SUMO.

The initial structure for yeast Smt3 was truncated from the X-ray structure of yeast

Smt3 in complex with ULP1 (PDB code: 1EUV[98]). The missing N-terminal and C-

terminal parts were modelled by MODELLER[90]. The initial structure for C. elegans

SMO-1 were randomly selected from NMR structures with the PDB code 5XQM[99]. The

mutants of SUMO were prepared using UCSF Chimera[89] based on the wild type

structures. Structures for a) SUMO1 that was acetylated on Lys23 (ACK23), b) SUMO1

that was phosphorylated on Ser2/Ser9/Ser9Thr10 (SUMO1 pS2/pS9/pS9pT10), and c)

SUMO2 that was phosphorylated/acetylated on Lys7 (SUMO2 ACK7), Thr12 (SUMO2

pT12) and Lys11Thr12 (SUMO2 ACK11pT12) were prepared by using PyTMs[87], a

PyMOL[100] plugin. The forcefield parameters of phosphorylated residues and acetylated

Lys were reported previously[101,102].

Energy minimization, equilibration and molecular dynamics simulations

Molecular Dynamics simulations were performed using the GROMACS 2016.3

package[88]. The system was described by the Amber99sb*-ILDN force field[103]. Each

SUMO structure was immersed in an explicit TIP4PD[104] truncated dodecahedron water

box, with at least 2.5 nm between periodic replicas of the protein (about 940 - 1540

nm^3). The system was neutralized by adding ions, and extra NaCl was added to

represent a solution with an ionic strength of 0.15 M to mimic physiological conditions.

The systems contained 125 to 200k atoms and were minimized using the steepest

descent minimization approach. After the minimization, the system was equilibrated in

the NVT ensemble for 200 ps with all heavy atom position-restrained with a force

constant of 239 kcal/mol. The temperature was maintained at 310 K using a V-rescale

thermostat with a coupling constant of 0.1 ps. Further equilibration was conducted in
the NPT ensemble with all heavy atoms position-restrained for 200 ps and Cα atoms
position-restrained for 1 ns, where the pressure was maintained at 1 atmosphere using
a Parrinello-Rahamn barostat with the coupling constant set to 2.0 ps.

All equilibrations were performed with a time step of 1 fs. For the production run, the
thermostat and barostat settings were the same as for the NPT run. To enable 2 fs time
steps, all bonds were constrained to equilibration length using the LINCS algorithm[105].
A real-space cutoff of 10 Å was used for the electrostatic and Lennard-Jones forces.
Snapshots from each trajectory were saved to disk every 50 ps. The simulations
performed in this work are listed in Table S1.

872 Analyses of molecular dynamics simulation

The change of solvent-accessible surface area (SASA) was used to quantify the
binding and unbinding process between the N-terminus and the SUMO SIM binding
groove or the complete SUMO core (Fig. S3 and S9). In case of the SIM binding
groove, the 'unbound' state includes all conformations with the N-termini not interacting
with the SUMO core and interacting with the core outside of the SIM-binding motif.
Kinetic rate constants were obtained by fitting the integrated dwell binding/unbinding
times. Six residues in the N-terminus adjacent to the SUMO core were not included for
calculating the contact area between the N-terminus and the SUMO core or the SIM
binding groove, respectively, to leave enough space (about 2 nm) for the SASA
calculation probe (radius of the probe is 0.14 nm) (Fig. S3A). A threshold (0.3 nm^2) was
used in binding event determinations with the SUMO core or the SIM binding groove.
The percentage of binding time of the N-terminus with the SIM binding groove or the
SUMO core was calculated as the ratio of MD snapshots with binding areas (ΔSASA)
larger than 0.3 nm^2 along MD simulations. The analyses were using GROMACS
utilities, and the first 100 ns of all production trajectories were regarded as equilibration
and not used in final analyses. The MD back-calculated chemical shifts were calculated
by using SHIFTX2[106]. The structures in the figures were prepared using PyMol[100].

891 General methods and maintenance of *C. elegans*

892 The Caenorhabditis elegans strains were maintained at 20°C on the standard NGM
893 (Nematode Growth Medium) plates seeded with E. coli bacterial strain OP50 as a food
894 source[107]. The derivate of Bristol strain N2 served as a wild-type reference. To generate
895 double mutants, standard crossing methods were used. The list of used strains in this
896 study is provided in the Key Resources Table and including the genotype in Table S2.

CRISPR/Cas9 genome editing

Genome editing was performed according to the co-CRISPR strategy described by[108]. An oligonucleotide corresponding to a target sequence near the smo-1 translational start site (sgRNA #1: 5` GCCGATGATGCAGCTCAAGC 3`) was cloned into the plasmid pMW46 (derivate of pDD162 from Addgene). The deletion of the eleven amino acids ADDAAQAGDNA at the SMO-1 N-terminus was achieved using the oligonucleotide pAF64 as repair template: CTC TAC CTC TCT CCT TCT ATC TCT TTT TCT CTT TTC AAA TCT AAT TTC GTT TCA GAG ACT CCC GCT ATA AAC GAT GGA ATA CAT CAA GAT CAA GGT CGT TGG ACA GGT AAT TTG ACT GGA AAT TCG CCG CGA ATT TGT TAA TAA TTC CC. The following constructs with indicated final concentration were microinjected into young adult N2 hermaphrodites: dpy-10 sgRNA PJA58 – 25 ng/µl, dpy-10 donor template oligonucleotide AF-ZF-827 – 0.5 nM, smo-1 sgRNA #1 pAF25 – 75 ng/µl, smo-1 donor template oligonucleotide pAF64 – 0.5 nM and the transformation marker myo-2::mCherry pCFJ90 – 2.5 ng/µl. Two to three days post injection, 100 F1 recombinants showing a Rol phenotype were transferred to separate NGM plates. Animals containing the desired 33bp smo-1(zh156) deletion were identified by PCR amplification using the primers OAF381 (CAC TCG TGT GAG TTG CAT TCT CCA TAG) and OAF383 (CAC GGA AGT GCA CTT CGT TGC TGT C) followed by sequencing. The smo-1(zh156) mutant was back-crossed twice to N2.

SUMO protein levels by immunoblotting

Worms were collected in H_2O supplemented with 20 mM N-ethylmaleimide. For each sample, 50 worms in a volume of 25 ml were mixed with 2x SDS sample buffer, boiled for 5 minutes and stored at -80~C until analysis. The samples were then reboiled, cooled to RT and treated with DNAse for 10 – 15 minutes. Half of each sample was separated by SDS PAGE and analyzed by immunoblotting using anti-SUMO 6F2. SUMO 6F2 was obtained from the DSHB, where it had been deposited by Pelisch, F. / Hay, R.T. (DSHB Hybridoma Product SUMO 6F2). The immunoblots were quantified with Fiji[109] and statistical analysis was performed using GraphPad Prism.

Brood size assay

The brood size assay was performed according to [110]. The L4 hermaphrodites were selected into individual NGM plates seeded with E. coli strain OP50 and incubated at 20°C. The animals were transferred to a fresh plate every 24h until the end of their self-reproduction period (around fifth day of the adult life). The number of the progeny from

each plate was counted. Total brood size is a sum of daily collected data. The
experiment was replicated three times, and each experiment included 10 animals of
wild-type and the mutant smo-1(zh156) genotype. The hermaphrodites, which were lost
or died due to transfer to a new plate, were excluded from the analysis.

DAPI and SYTO12 staining

L4 hermaphrodites (50-60 animals per genotype) were selected to fresh NGM plate
containing OP50 bacteria and incubated at 20°C for 24 h. The adult animals were
transferred in M9 buffer to a siliconized Eppendorf tube, spun down for 1min at 3000
rpm, washed with 750 µl of PBS and spun down. Washed animals were incubated in
750 µl ice-cold methanol at -20°C for 5 min and washed twice with 750µl of PBS-T. The
animals were incubated in 750 µl DAPI solution (0.1% in PBS) for 15 min and washed
twice with 750 µl of PBS-T before mounting in Mowiol. To stain corpses in adult
hermaphrodites germline, the SYTO 12 staining protocol was used according to[60]. We
scored SYTO 12 positive germ cells per gonad arm of the adult hermaphrodites (29
hours post L4). Three biological replicates of the assay were performed.

Microscopy and image processing

For Nomarski and fluorescent imaging, live animals were mounted on 4% agarose pads
and immobilized by 20 mM tetramisole hydrochloride solution in M9 buffer, unless
stated otherwise. To count DAPI-stained nuclei and for SYTO 12 staining, a Leica
DM6000 B microscope equipped with Nomarski and fluorescence optics and the Leica
Application Suite X software was used. The animals expressing CED-1::GFP were
imaged on a Leica DMRA microscope controlled by custom MatLab script and
equipped with a beam splitter and two Hamamatsu ORCA-flash4.0L cameras to
simultaneously acquire z-stacks in the DIC and GFP channels was used. Images were
analyzed and quantified with Fiji software[109]. The L4 hermaphrodites of the strains
WS4116 and AH6119 containing the CED-1::GFP reporter were picked to fresh NGM
plate seeded with OP50 bacteria and incubated for 24h or 48h at 20°C before being
prepared for image analysis. CED-1::GFP-labeled corpses were counted by inspecting
the z-stacks recorded across the pachytene zone. DAPI-stained nuclei in the pachytene
region were manually counted across z-stacks in fixed and stained animals, and the
area corresponding to the pachytene region in each gonad arm was identified through
the morphology of the DAPI-stained nuclei.
The statistical analysis was performed by using GraphPad Prism as indicated in the
figure legend.

REFERENCES FOR STAR METHODS

973 Flotho, A., Werner, A., Winter, T., Frank, A.S., Ehret, H. and Melchior, F. (2012).
974 Recombinant reconstitution of sumoylation reactions in vitro. Methods Mol Biol. *832*,
975 93-110.
976
977 Hess, B. (2008). P-LINCS: A Parallel Linear Constraint Solver for Molecular
978 Simulation. J Chem Theory Comput. *4*, 116–122.
979
980 Hutten, S., Flotho, A., Melchior, F. and Kehlenbach, R.H. (2008) The Nup358-
981 RanGAP complex is required for efficient importin alpha/beta-dependent nuclear
982 import. Mol. Biol. Cell *10*, 2300-2310.
983
984 Johnson, E.S. and Gupta, A.A. (2001) An E3-like factor that promotes SUMO
985 conjugation to the yeast septins. Cell *106*, 735–744.
986
987 Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J.L., Dror, R.O. and
988 Shaw, D.E. (2010). Improved side-chain torsion potentials for the Amber ff99SB
989 protein force field. Proteins *78*, 1950-1958.
990
991 Mahajan, R., Delphin, C., Guan, T., Gerace, L. and Melchior, F. (1997) A small
992 ubiquitin related polypeptide involved in targeting RanGAP1 to nuclear pore complex
993 protein RanBP2. Cell *88*, 97-107.
994
995 Melchior, F. (2000). SUMO-1 - Non-Classical Ubiquitin. Annu. Rev. Cell Dev. Biol. *16*,
996 591-626.
997
998 Meulmeester, E., Kunze, M., Hsiaoh, H.H., Urlaub, H., Melchior, F. (2008)
999 Mechanism and consequences of paralog specific sumoylation of USP25. Mol.
1000 Cell *30*, 610-619.
1001
1002 Pettersen E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M., Meng,
1003 E.C., Ferrin, T.E. (2004). UCSF Chimera--a visualization system for exploratory
1004 research and analysis. J Comput Chem. *25*, 1605–1612.
1005
1006 Piana, S., Donchev, A.G., Robustelli, P. and Shaw, D.E. (2015). Water dispersion
1007 interactions strongly influence simulated structural properties of disordered protein
1008 states. J Phys Chem B., *119*, 5113–5123.
1009
1010 Pichler, A., Gast, A., Seeler, J.S., Dejean, A. and Melchior, F. (2002) The nucleoporin
1011 RanBP2 is a SUMO1 E3 Ligase. Cell *108*, 109-120.
1012
1013 Saitoh, H. and Hinchey, J. (2000). Functional heterogeneity of small ubiquitin-related
1014 protein modifiers SUMO-1 versus SUMO-2/3. J Biol Chem *275*, 6252–58.
1015
1016 Van Der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A.E., Berendsen, H.J.
1017 (2005). GROMACS: fast, flexible, and free. J Comput Chem. *26*, 1701–1718.
1018

1019   Warnecke, A., Sandalova, T., Achour, A. and Harris, R.A. (2014). PyTMs: a useful
1020   PyMOL plugin for modeling common post-translational modifications. BMC
1021   Bioinformatics. *15*, 370.

1022
1023   Webb, B., and Sali, A. (2014). Comparative Protein Structure Modeling Using
1024   MODELLER. Curr Protoc Bioinformatics *47*, 5.6.1-32.

1025
1026   Werner, A., Flotho, A. and Melchior, F. (2012). The RanBP2/RanGAP1*SUMO1/Ubc9
1027   Complex Is a Multisubunit SUMO E3 Ligase. Mol Cell *46*, 287–298.
1028
1029

1030

ACKNOWLEDGEMENTS

We acknowledge Ana Cristina Laranjeira for generating the C. elegans strain AH5891,

and thank Klaus "Gerry" Meese for excellent technical support, and former and current lab

members for constructs, reagents and helpful discussions. S.M.R. acknowledges the

Heidelberg Biosciences International Graduate School (HBIGS). F.J. is grateful for

funding from the BIOMS program of Heidelberg University. F.G. acknowledges funding

by the Klaus Tschira Foundation. F.M. and F.G. were members of the Cluster

CellNetworks.

1039

AUTHOR CONTRIBUTIONS

S.M.R., A.Fr. and A.Fl. designed and carried out most biochemical experiments, F.J.

designed and carried out all Molecular Dynamics simulations, T.R. designed and

carried out all NMR experiments, A.Fe. designed and carried out *C. elegans* in vivo

experiments. E.M. carried out some biochemical experiments. M.D. carried out some *C.*

elegans in vivo experiments. F.G. guided simulations, R.K. guided NMR experiments,

A.H. guided *C.elegans* in vivo experiments, F.M. and A. Fl. guided the overall design of

the project. T.R., F.J., S.M.R., A.Fe., A.H., R.K., F.G., A.Fl. and F.M. wrote the

manuscript.

1049

DECLARATION OF INTERESTS

1051   The authors declare no competing financial interests.

1052

1053

FIGURE LEGENDS

Figure 1 N-terminal deletions of SUMO1 proteins, but not SUMO2, greatly increase binding of SIM-containing proteins.

(A) Overlay of 5 selected states of a SUMO1 solution structure (2n1v) using Pymol. Whereas the SUMO core and C-terminus (dark grey) remain stable in all states, the N-terminus (light grey) shows a high degree of flexibility. (B) Partial sequence alignment of human SUMO1, SUMO2, *C. elegans* SMO-1*,* and *S. cerevisiae* Smt3 using the Clustal Omega web tool. The flexible N-terminal extension is indicated by a box, the SIM-binding groove is highlighted grey. (C) and (E) Pulldown of recombinant Usp25, Ubc9, TDP2 or RanBP2 with SUMO1 or SUMO2 analyzed by SDS-PAGE and Coomassie staining. (D) The amounts of Usp25 in the eluates of SUMO pull-downs in (C) were quantified relative to Ubc9 (n = 4). (F) The amounts of Usp25 in the eluates of SUMO pull-downs in (E) were quantified relative to Ubc9 (n = 6). (G) Pulldown of recombinant *S. cerevisiae* Siz1 or hUbc9 with Smt3 analyzed by SDS-PAGE and Coomassie staining. (H) The amounts of Siz1 were quantified relative to hUbc9 (n = 4). (I) Pulldown of recombinant *C. elegans* GEI-17 or hUbc9 with Ovalbumin (Ova) or Smo-1 beads analyzed by SDS-PAGE and Coomassie staining. (J) The amounts of GEI-17 were quantified relative to hUbc9 (n = 3). Error bars represent one standard deviation. Asterisks indicate p-values: n.s.: P > 0.05; **: P ≤ 0.01; ***: P ≤ 0.001

Figure 2 N-terminal deletions of SUMO1 augment binding and SUMOylation of Usp25 by enhancing SUMO-SIM interactions.

(A) If not occupied by the N-terminus, the SIM-binding groove of SUMO1 can aid complex formation between SUMOylated proteins and their SIM-containing potential interaction partners. (B) SIMs in SUMOylation targets can promote SUMO-modification by stabilizing complex formation with SUMO-loaded Ubc9 termed "SIM-dependent SUMOylation". (C) and (E) Pulldown of recombinant Usp25 wt, Usp25-V91A,I92A (SIM mutant), or RanBP2 with glutathione beads loaded with the indicated SUMO (S)-GST variants, analyzed by SDS-PAGE and Coomassie staining. (D) and (F) The amounts of Usp25 and RanBP2 in the eluates of SUMO pull-downs in (C) and (E) were quantified relative to SUMO-GST (n = 3). (G) - (L) *In vitro* SUMOylation reactions of recombinant Usp25 wt or -V91A,I92A, and RanGAP1 with the indicated human SUMO1, SUMO2 or *C. elegans* SMO-1 mutants, analyzed by SDS-PAGE and immunoblotting. Error bars represent one standard deviation. Asterisks indicate p-values: **: P ≤ 0.01; ***: P ≤ 0.001;

Figure 3 NMR experiment and Molecular Dynamics simulations reveal that the N-termini of SUMO1 and SMO-1 transiently occupy the SIM binding groove.

(A) ^{1}H-^{15}N hNOE values for SUMO1. The x-axis shows SUMO1 residues in a non-contiguous fashion; residues not shown were either prolines (residues 8, 58 and 77) or could not be analyzed due to insufficient data quality (residues 1-3, 24, 29, 41 and 97). Colored backgrounds show areas with indicated dynamics. (B) HSQC-TROSY experiments comparing full-length SUMO1 wt and the ΔN19 variants. Histogram of CSPs of (^{15}N)SUMO1 ΔN19 compared to full-length (^{15}N)SUMO1; the resonance for Met19 could not be assigned confidently (shown as off-scale in histogram - marked with *); residues 1-18 are displayed only for clarity with a value of 0; Pro8, Pro58, Pro77 are not shown. The SIM binding groove and 70/80 region are marked (black bars). (C-F) Binding time, averaged over 8 1μs-trajectories, of the SUMO N-terminus to other residues for SUMO1, SUMO2, SMO-1 and yeast Smt3. (G) Average binding time of the SUMO N-terminus to the SIM binding groove for SUMO1, SUMO2, SMO-1 and yeast Smt3. Data points (n=8): average along single 1 μs trajectories; bars: average over all 8 trajectories, errors bars: standard errors of the mean of the 8 averages. Asterisks indicate p-values: *: P ≤ 0.05; **: P ≤ 0.01; ***: P ≤ 0.001. (H) Kinetic rate constants for binding/unbinding processes between the N-terminus of SUMO1, SUMO2, SMO-1 and yeast Smt3 and the SIM binding groove.

Figure 4 Mutating negatively charged residues in the N-terminus of SUMO1 abolishes the inhibitory effect of the N-terminus.

The SIM binding groove, encompassing β2-strand and α-1 helix, and N-terminus were shown as non-transparent. The positively/negatively charged residues were colored in blue/red, respectively. The charged residues in N-terminus and around the SIM binding groove were shown as ball and stick. The residues related with post-translational modifications or mutations were highlighted in the figure with their names. (A) human SUMO1; (B) *C. elegans* SMO-1; (C) Pulldown of recombinant Usp25, Ubc9 or TDP2 with the indicated SUMO1 beads analyzed by SDS-PAGE and Coomassie staining. (D) The amounts of Usp25 in the eluates of SUMO pull-downs in (C) were quantified relative to Ubc9 (n = 6). (E) Pulldown of recombinant *C. elegans* GEI-17 or hUbc9 with Ovalbumin (Ova) or indicated SMO-1 beads analyzed by SDS-PAGE and Coomassie staining. (F) The amounts of GEI-17 were quantified relative to hUbc9 (n = 3). (G) and (H) *In vitro* SUMOylation of recombinant Usp25 and RanGAP1 with human SUMO1 wt or the indicated mutants, analyzed by SDS-PAGE and immunoblotting. Error bars represent one standard deviation. Asterisks indicate p-values: *: P ≤ 0.05; **: P ≤ 0.01; ***: P ≤ 0.001

Figure 5 Negatively charged residues in the N-terminus of SUMO1 mediate the

inhibitory effect on SIM-dependent interactions

(A) HSQC-TROSY experiments comparing full-length SUMO1 wt and the ED11,12KK

mutant. Histogram of CSPs of (^{15}N)SUMO1-ED11,12KK compared to full-length

(^{15}N)SUMO1; the resonances for Thr10, Lys12, Gly14 and Glu20 of the mutant could

not be assigned confidently (shown as off-scale in histogram - marked with *); Pro8,

Pro58 and Pro77 are not shown. The SIM binding groove (black bars) as well as the

folded core of the protein (grey background) are marked. (B-C) Binding time and (D)

Average binding time of the SUMO N-terminus to the SIM binding groove for wt

(identical to Fig. 3C) and ED11,12KK variant of SUMO1. (E-F) Binding time and (G)

Average binding time of the N-terminus of wt (identical to Fig. 3E) and DD3,4KK of

SMO-1 to residues of the SUMO core. Data points: average along single 1 μs

trajectories; bars: average over 8 trajectories, errors bars: standard errors of the mean

of the 8 averages. (H) Pulldown of recombinant Usp25 with GST-tagged

SUMO1/SUMO2 chimeras, analyzed by SDS-PAGE and Coomassie staining. (I) The

amounts of Usp25 in the eluates of SUMO pull-downs in (H) were quantified relative to

the SUMO-GST proteins (n = 3). Asterisks indicate p-values: *: P ≤ 0.05; **: P ≤ 0.01

1146

Figure 6 Post-translation modifications of SUMO's intrinsically disordered N-

termini may regulate SIM-dependent interactions.

(A) Average binding time of the SUMO N-terminus to the SIM binding groove for

SUMO1 wt (identical to Fig. 3C) and pS2, pS9, and pS9pT10 SUMO1 variants. (B) and

(C) Binding time of the N-terminus of SUMO1 wt (identical to Fig. 3C) and SUMO1-pS9

to individual residues of the SUMO core. (D) Average binding time of the SUMO N-

terminus to the SIM binding groove for SUMO2 wt and the pT12, AcK7 and AcK11pT12

SUMO2 variants. (E) and (F) Binding time of the N-terminus of SUMO2 wt (identical to

Fig. 3D) and SUMO2-pT12 to individual residues of the SUMO core. Data points:

average along single 1 μs trajectories; bars: average over 8 trajectories, errors bars:

standard errors of the mean of the 8 averages. (G) Pulldown of recombinant Usp25 or

Ubc9 with the indicated SUMO1 beads, analyzed by SDS-PAGE and Coomassie

staining. (H) The amounts of Usp25 in the eluates of SUMO pull-downs in (G) were

quantified relative to Ubc9 (n = 4). Error bars represent one standard deviation.

Asterisks indicate p-values: *: P ≤ 0.05; **: P ≤ 0.01; ***: P ≤ 0.001. (I) Model of the

intrinsically disordered region of SUMO serving as a regulatable cis - inhibitor for SIM -

dependent interactions through post-translation modification.

1164

Figure 7: Deletion of the flexible N-terminus in C. elegans SMO-1 reduces germ cell survival.

(A) Whole-mount DAPI staining of the wild-type and of SMO-1ΔN12 (smo-1(zh156)) animals 24 hours post L4 stage. (B) Brood size, (C) number of germ cell nuclei at the pachytene stage per gonad arm, and (D) area of the pachytene region in wild-type and SMO-1ΔN12 animals. (E) Average number of pachytene stage germ cell nuclei per area. (F) Apoptotic corpses outlined by CED-1::GFP staining in wild-type and SMO-1ΔN12 mutants 24 and 48 hours post L4. The engulfed corpses are indicated with white arrowheads. The left panels show the DIC images overlayed with CED-1::GFP signal (average intensity projections of the outer gonad layers) and the right panels the CED-1::GFP signal alone. (G) Number of CED-1::GFP-positive apoptotic corpses per gonad arm in wild-type and SMO-1ΔN12 animals with or without *cep-1(gk148)* background 24 and 48 hours post L4 stage. (H) Number of SYTO12 positive germ cells per gonad arm in wild-type and SMO-1ΔN12 animals 29 hours post L4 stage. Bars indicate average values, dots individual values and error bars the standard deviations. The numbers in brackets indicate the numbers of animals scored. If the values were normally distributed (in B - E, H), statistical significance was calculated with an unpaired parametric t-test, or else (in G) a Kruskal-Wallis test was used. Asterisks indicate the p-values as * p≤0.05; ** p≤ 0.01; *** p≤ 0.001; **** p≤0.001, n.s. for not significant. The scale bars in (A & F) represent 20μm.

REFERENCES

1.   Flotho, A., and Melchior, F. (2013). Sumoylation: a regulatory protein modification in health and disease. Annu Rev Biochem *82*, 357-385. 10.1146/annurev-biochem-061909-093311.

2.   Gareau, J.R., and Lima, C.D. (2010). The SUMO pathway: emerging mechanisms that shape specificity, conjugation and recognition. Nat Rev Mol Cell Biol *11*, 861-871. 10.1038/nrm3011.

3.   Melchior, F. (2000). SUMO--nonclassical ubiquitin. Annu Rev Cell Dev Biol *16*, 591-626. 10.1146/annurev.cellbio.16.1.591.

4.   Vertegaal, A.C.O. (2022). Signalling mechanisms and cellular functions of SUMO. Nat Rev Mol Cell Biol *23*, 715-731. 10.1038/s41580-022-00500-y.

5.   Golebiowski, F., Matic, I., Tatham, M.H., Cole, C., Yin, Y., Nakamura, A., Cox, J., Barton, G.J., Mann, M., and Hay, R.T. (2009). System-wide changes to SUMO modifications in response to heat shock. Sci Signal *2*, ra24. 10.1126/scisignal.2000282.

6.   Hendriks, I.A., and Vertegaal, A.C. (2016). A comprehensive compilation of SUMO proteomics. Nat Rev Mol Cell Biol *17*, 581-595. 10.1038/nrm.2016.81.

7.   Pichler, A., Fatouros, C., Lee, H., and Eisenhardt, N. (2017). SUMO conjugation - a mechanistic view. Biomol Concepts *8*, 13-36. 10.1515/bmc-2016-0030.

8.   Nayak, A., and Muller, S. (2014). SUMO-specific proteases/isopeptidases: SENPs and beyond. Genome Biol *15*, 422. 10.1186/s13059-014-0422-2.

9.   Hickey, C.M., Wilson, N.R., and Hochstrasser, M. (2012). Function and
regulation of SUMO proteases. Nat Rev Mol Cell Biol *13*, 755-766. 10.1038/nrm3478.
10.   Novatchkova, M., Budhiraja, R., Coupland, G., Eisenhaber, F., and Bachmair, A.
(2004). SUMO conjugation in plants. Planta *220*, 1-8. 10.1007/s00425-004-1370-y.
11.   Bencsath, K.P., Podgorski, M.S., Pagala, V.R., Slaughter, C.A., and Schulman,
B.A. (2002). Identification of a multifunctional binding site on Ubc9p required for Smt3p
conjugation. J Biol Chem *277*, 47938-47945. 10.1074/jbc.M207442200.
12.   Hendriks, I.A., Lyon, D., Young, C., Jensen, L.J., Vertegaal, A.C., and Nielsen,
M.L. (2017). Site-specific mapping of the human SUMO proteome reveals co-
modification with phosphorylation. Nat Struct Mol Biol *24*, 325-336.
10.1038/nsmb.3366.
13.   Tatham, M.H., Jaffray, E., Vaughan, O.A., Desterro, J.M., Botting, C.H.,
Naismith, J.H., and Hay, R.T. (2001). Polymeric chains of SUMO-2 and SUMO-3 are
conjugated to protein substrates by SAE1/SAE2 and Ubc9. J Biol Chem *276*, 35368-
35374. 10.1074/jbc.M104214200.
14.   Becker, J., Barysch, S.V., Karaca, S., Dittner, C., Hsiao, H.H., Berriel Diaz, M.,
Herzig, S., Urlaub, H., and Melchior, F. (2013). Detecting endogenous SUMO targets in
mammalian cells and tissues. Nat Struct Mol Biol *20*, 525-531. 10.1038/nsmb.2526.
15.   Meulmeester, E., Kunze, M., Hsiao, H.H., Urlaub, H., and Melchior, F. (2008).
Mechanism and consequences for paralog-specific sumoylation of ubiquitin-specific
protease 25. Mol Cell *30*, 610-619. 10.1016/j.molcel.2008.03.021.
16.   Zhu, J., Zhu, S., Guzzo, C.M., Ellis, N.A., Sung, K.S., Choi, C.Y., and Matunis,
M.J. (2008). Small ubiquitin-related modifier (SUMO) binding determines substrate
recognition and paralog-selective SUMO modification. J Biol Chem *283*, 29405-29415.
10.1074/jbc.M803632200.
17.   Pfander, B., Moldovan, G.L., Sacher, M., Hoege, C., and Jentsch, S. (2005).
SUMO-modified PCNA recruits Srs2 to prevent recombination during S phase. Nature
436, 428-433. 10.1038/nature03665.
18.   Lallemand-Breitenbach, V., Jeanne, M., Benhenda, S., Nasr, R., Lei, M., Peres,
L., Zhou, J., Zhu, J., Raught, B., and de The, H. (2008). Arsenic degrades PML or PML-
RARalpha through a SUMO-triggered RNF4/ubiquitin-mediated pathway. Nat Cell Biol
10, 547-555. 10.1038/ncb1717.
19.   Tatham, M.H., Geoffroy, M.C., Shen, L., Plechanovova, A., Hattersley, N.,
Jaffray, E.G., Palvimo, J.J., and Hay, R.T. (2008). RNF4 is a poly-SUMO-specific E3
ubiquitin ligase required for arsenic-induced PML degradation. Nat Cell Biol *10*, 538-
546. 10.1038/ncb1716.
20.   Kerscher, O. (2007). SUMO junction-what's your function? New insights through
SUMO-interacting motifs. EMBO Rep *8*, 550-555. 10.1038/sj.embor.7400980.
21.   Capili, A.D., and Lima, C.D. (2007). Structure and analysis of a complex
between SUMO and Ubc9 illustrates features of a conserved E2-Ubl interaction. J Mol
Biol *369*, 608-618. 10.1016/j.jmb.2007.04.006.
22.   Duda, D.M., van Waardenburg, R.C., Borg, L.A., McGarity, S., Nourse, A.,
Waddell, M.B., Bjornsti, M.A., and Schulman, B.A. (2007). Structure of a SUMO-
binding-motif mimic bound to Smt3p-Ubc9p: conservation of a non-covalent ubiquitin-
like protein-E2 complex as a platform for selective interactions within a SUMO pathway.
J Mol Biol *369*, 619-630. 10.1016/j.jmb.2007.04.007.
23.   Liu, Q., Jin, C., Liao, X., Shen, Z., Chen, D.J., and Chen, Y. (1999). The binding
interface between an E2 (UBC9) and a ubiquitin homologue (UBL1). J Biol Chem *274*,
16979-16987. 10.1074/jbc.274.24.16979.
24.   Pilla, E., Moller, U., Sauer, G., Mattiroli, F., Melchior, F., and Geiss-Friedlander,
R. (2012). A novel SUMO1-specific interacting motif in dipeptidyl peptidase 9 (DPP9)
that is important for enzymatic regulation. J Biol Chem *287*, 44320-44329.
10.1074/jbc.M112.397224.
25.   Diehl, C., Akke, M., Bekker-Jensen, S., Mailand, N., Streicher, W., and
Wikstrom, M. (2016). Structural Analysis of a Complex between Small Ubiquitin-like
Modifier 1 (SUMO1) and the ZZ Domain of CREB-binding Protein (CBP/p300) Reveals

a New Interaction Surface on SUMO. J Biol Chem *291*, 12658-12672.
10.1074/jbc.M115.711325.
26.   Liu, J., Xue, Z., Zhang, Y., Vann, K.R., Shi, X., and Kutateladze, T.G. (2020).
Structural Insight into Binding of the ZZ Domain of HERC2 to Histone H3 and SUMO1.
Structure *28*, 1225-1230 e1223. 10.1016/j.str.2020.07.003.
27.   Vogt, B., and Hofmann, K. (2012). Bioinformatical detection of recognition
factors for ubiquitin and SUMO. Methods Mol Biol *832*, 249-261. 10.1007/978-1-61779-
474-2_18.
28.   Sun, H., and Hunter, T. (2012). Poly-small ubiquitin-like modifier (PolySUMO)-
binding proteins identified through a string search. J Biol Chem *287*, 42071-42083.
10.1074/jbc.M112.410985.
29.   Hecker, C.M., Rabiller, M., Haglund, K., Bayer, P., and Dikic, I. (2006).
Specification of SUMO1- and SUMO2-interacting motifs. J Biol Chem *281*, 16117-
16127. 10.1074/jbc.M512757200.
30.   Song, J., Durrin, L.K., Wilkinson, T.A., Krontiris, T.G., and Chen, Y. (2004).
Identification of a SUMO-binding motif that recognizes SUMO-modified proteins. Proc
Natl Acad Sci U S A *101*, 14373-14378. 10.1073/pnas.0403498101.
31.   Schellenberg, M.J., Lieberman, J.A., Herrero-Ruiz, A., Butler, L.R., Williams,
J.G., Munoz-Cabello, A.M., Mueller, G.A., London, R.E., Cortes-Ledesma, F., and
Williams, R.S. (2017). ZATT (ZNF451)-mediated resolution of topoisomerase 2 DNA-
protein cross-links. Science *357*, 1412-1416. 10.1126/science.aam6468.
32.   Husnjak, K., and Dikic, I. (2012). Ubiquitin-binding proteins: decoders of
ubiquitin-mediated cellular functions. Annu Rev Biochem *81*, 291-322.
10.1146/annurev-biochem-051810-094654.
33.   Stehmeier, P., and Muller, S. (2009). Phospho-regulated SUMO interaction
modules connect the SUMO system to CK2 signaling. Mol Cell *33*, 400-409.
10.1016/j.molcel.2009.01.013.
34.   Cappadocia, L., Mascle, X.H., Bourdeau, V., Tremblay-Belzile, S., Chaker-
Margot, M., Lussier-Price, M., Wada, J., Sakaguchi, K., Aubry, M., Ferbeyre, G., and
Omichinski, J.G. (2015). Structural and functional characterization of the
phosphorylation-dependent interaction between PML and SUMO1. Structure *23*, 126-
138. 10.1016/j.str.2014.10.015.
35.   Chang, C.C., Naik, M.T., Huang, Y.S., Jeng, J.C., Liao, P.H., Kuo, H.Y., Ho,
C.C., Hsieh, Y.L., Lin, C.H., Huang, N.J., et al. (2011). Structural and functional roles of
Daxx SIM phosphorylation in SUMO paralog-selective binding and apoptosis
modulation. Mol Cell *42*, 62-74. 10.1016/j.molcel.2011.02.022.
36.   Escobar-Cabrera, E., Okon, M., Lau, D.K., Dart, C.F., Bonvin, A.M., and
McIntosh, L.P. (2011). Characterizing the N- and C-terminal Small ubiquitin-like
modifier (SUMO)-interacting motifs of the scaffold protein DAXX. J Biol Chem *286*,
19816-19829. 10.1074/jbc.M111.231647.
37.   Namanja, A.T., Li, Y.J., Su, Y., Wong, S., Lu, J., Colson, L.T., Wu, C., Li, S.S.,
and Chen, Y. (2012). Insights into high affinity small ubiquitin-like modifier (SUMO)
recognition by SUMO-interacting motifs (SIMs) revealed by a combination of NMR and
peptide array analysis. J Biol Chem *287*, 3231-3240. 10.1074/jbc.M111.293118.
38.   Sekiyama, N., Ikegami, T., Yamane, T., Ikeguchi, M., Uchimura, Y., Baba, D.,
Ariyoshi, M., Tochio, H., Saitoh, H., and Shirakawa, M. (2008). Structure of the small
ubiquitin-like modifier (SUMO)-interacting motif of MBD1-containing chromatin-
associated factor 1 bound to SUMO-3. J Biol Chem *283*, 35966-35975.
10.1074/jbc.M802528200.
39.   Bayer, P., Arndt, A., Metzger, S., Mahajan, R., Melchior, F., Jaenicke, R., and
Becker, J. (1998). Structure determination of the small ubiquitin-related modifier SUMO-
1. J Mol Biol *280*, 275-286. 10.1006/jmbi.1998.1839.
40.   Bylebyl, G.R., Belichenko, I., and Johnson, E.S. (2003). The SUMO
isopeptidase Ulp2 prevents accumulation of SUMO chains in yeast. J Biol Chem *278*,
44113-44120. 10.1074/jbc.M308357200.

41.   Newman, H.A., Meluh, P.B., Lu, J., Vidal, J., Carson, C., Lagesse, E., Gray,
J.J., Boeke, J.D., and Matunis, M.J. (2017). A high throughput mutagenic analysis of
yeast sumo structure and function. PLoS Genet *13*, e1006612.
10.1371/journal.pgen.1006612.
42.   Lussier-Price, M., Wahba, H.M., Mascle, X.H., Cappadocia, L., Bourdeau, V.,
Gagnon, C., Igelmann, S., Sakaguchi, K., Ferbeyre, G., and Omichinski, J.G. (2022).
Zinc controls PML nuclear body formation through regulation of a paralog specific auto-
inhibition in SUMO1. Nucleic Acids Res *50*, 8331-8348. 10.1093/nar/gkac620.
43.   Srikumar, T., Lewicki, M.C., Costanzo, M., Tkach, J.M., van Bakel, H., Tsui, K.,
Johnson, E.S., Brown, G.W., Andrews, B.J., Boone, C., et al. (2013). Global analysis of
SUMO chain function reveals multiple roles in chromatin regulation. J Cell Biol *201*,
145-163. 10.1083/jcb.201210019.
44.   Pichler, A., Gast, A., Seeler, J.S., Dejean, A., and Melchior, F. (2002). The
nucleoporin RanBP2 has SUMO1 E3 ligase activity. Cell *108*, 109-120.
45.   Tatham, M.H., Kim, S., Jaffray, E., Song, J., Chen, Y., and Hay, R.T. (2005).
Unique binding interactions among Ubc9, SUMO and RanBP2 reveal a mechanism for
SUMO paralog selection. Nat Struct Mol Biol *12*, 67-74. 10.1038/nsmb878.
46.   Knipscheer, P., van Dijk, W.J., Olsen, J.V., Mann, M., and Sixma, T.K. (2007).
Noncovalent interaction between Ubc9 and SUMO promotes SUMO chain formation.
EMBO J *26*, 2797-2807. 10.1038/sj.emboj.7601711.
47.   Takahashi, Y., Iwase, M., Konishi, M., Tanaka, M., Toh-e, A., and Kikuchi, Y.
(1999). Smt3, a SUMO-1 homolog, is conjugated to Cdc3, a component of septin rings
at the mother-bud neck in budding yeast. Biochem Biophys Res Commun *259*, 582-
587. 10.1006/bbrc.1999.0821.
48.   Vertegaal, A.C. (2010). SUMO chains: polymeric signals. Biochem Soc Trans
38, 46-49. 10.1042/BST0380046.
49.   Takahashi, Y., and Kikuchi, Y. (2005). Yeast PIAS-type Ull1/Siz1 is composed
of SUMO ligase and regulatory domains. J Biol Chem *280*, 35822-35828.
10.1074/jbc.M506794200.
50.   Pelisch, F., Tammsalu, T., Wang, B., Jaffray, E.G., Gartner, A., and Hay, R.T.
(2017). A SUMO-Dependent Protein Network Regulates Chromosome Congression
during Oocyte Meiosis. Mol Cell *65*, 66-77. 10.1016/j.molcel.2016.11.001.
51.   Wand, A.J., Urbauer, J.L., McEvoy, R.P., and Bieber, R.J. (1996). Internal
dynamics of human ubiquitin revealed by 13C-relaxation studies of randomly
fractionally labeled protein. Biochemistry *35*, 6116-6125. 10.1021/bi9530144.
52.   Borgia, A., Borgia, M.B., Bugge, K., Kissling, V.M., Heidarsson, P.O.,
Fernandes, C.B., Sottini, A., Soranno, A., Buholzer, K.J., Nettels, D., et al. (2018).
Extreme disorder in an ultrahigh-affinity protein complex. Nature *555*, 61-66.
10.1038/nature25762.
53.   Sharma, R., Raduly, Z., Miskei, M., and Fuxreiter, M. (2015). Fuzzy complexes:
Specific binding without complete folding. FEBS Lett *589*, 2533-2542.
10.1016/j.febslet.2015.07.022.
54.   Jones, D.T. (1999). Protein secondary structure prediction based on position-
specific scoring matrices. J Mol Biol *292*, 195-202. 10.1006/jmbi.1999.3091.
55.   Ahmed, M.C., Papaleo, E., and Lindorff-Larsen, K. (2018). How well do force
fields capture the strength of salt bridges in proteins? Peerj *6*. ARTN e4967
10.7717/peerj.4967.
56.   Matic, I., Macek, B., Hilger, M., Walther, T.C., and Mann, M. (2008).
Phosphorylation of SUMO-1 occurs in vivo and is conserved through evolution. J
Proteome Res *7*, 4050-4057. 10.1021/pr800368m.
57.   Ullmann, R., Chien, C.D., Avantaggiati, M.L., and Muller, S. (2012). An
acetylation switch regulates SUMO-dependent protein interaction networks. Mol Cell
46, 759-770. 10.1016/j.molcel.2012.04.006.
58.   Pelisch, F., Sonneville, R., Pourkarimi, E., Agostinho, A., Blow, J.J., Gartner, A.,
and Hay, R.T. (2014). Dynamic SUMO modification regulates mitotic chromosome

assembly and cell cycle progression in Caenorhabditis elegans. Nat Commun *5*, 5485.
10.1038/ncomms6485.
59.   Reichman, R., Shi, Z., Malone, R., and Smolikove, S. (2018). Mitotic and Meiotic
Functions for the SUMOylation Pathway in the Caenorhabditis elegans Germline.
Genetics *208*, 1421-1441. 10.1534/genetics.118.300787.
60.   Gumienny, T.L., Lambie, E., Hartwieg, E., Horvitz, H.R., and Hengartner, M.O.
(1999). Genetic control of programmed cell death in the Caenorhabditis elegans
hermaphrodite germline. Development *126*, 1011-1022. 10.1242/dev.126.5.1011.
61.   Schumacher, B., Hanazawa, M., Lee, M.H., Nayak, S., Volkmann, K., Hofmann,
E.R., Hengartner, M., Schedl, T., and Gartner, A. (2005). Translational repression of C.
elegans p53 by GLD-1 regulates DNA damage-induced apoptosis. Cell *120*, 357-368.
10.1016/j.cell.2004.12.009.
62.   Ward, J.J., Sodhi, J.S., McGuffin, L.J., Buxton, B.F., and Jones, D.T. (2004).
Prediction and functional analysis of native disorder in proteins from the three kingdoms
of life. J Mol Biol *337*, 635-645. 10.1016/j.jmb.2004.02.002.
63.   Dunker, A.K., Brown, C.J., Lawson, J.D., Iakoucheva, L.M., and Obradovic, Z.
(2002). Intrinsic disorder and protein function. Biochemistry *41*, 6573-6582.
64.   Uversky, V.N. (2002). Natively unfolded proteins: a point where biology waits for
physics. Protein Sci *11*, 739-756. 10.1110/ps.4210102.
65.   Oldfield, C.J., and Dunker, A.K. (2014). Intrinsically disordered proteins and
intrinsically disordered protein regions. Annu Rev Biochem *83*, 553-584.
10.1146/annurev-biochem-072711-164947.
66.   Tompa, P. (2002). Intrinsically unstructured proteins. Trends Biochem Sci *27*,
527-533.
67.   Ulrich, H.D. (2008). The fast-growing business of SUMO chains. Mol Cell *32*,
301-305. 10.1016/j.molcel.2008.10.010.
68.   Gartner, A., Wagner, K., Holper, S., Kunz, K., Rodriguez, M.S., and Muller, S.
(2018). Acetylation of SUMO2 at lysine 11 favors the formation of non-canonical SUMO
chains. EMBO Rep *19*. 10.15252/embr.201846117.
69.   Grana-Montes, R., Marinelli, P., Reverter, D., and Ventura, S. (2014). N-terminal
protein tails act as aggregation protective entropic bristles: the SUMO case.
Biomacromolecules *15*, 1194-1203. 10.1021/bm401776z.
70.   Gianni, S., Dogan, J., and Jemth, P. (2016). Coupled binding and folding of
intrinsically disordered proteins: what can we learn from kinetics? Curr Opin Struct Biol
36, 18-24. 10.1016/j.sbi.2015.11.012.
71.   Lacy, E.R., Filippov, I., Lewis, W.S., Otieno, S., Xiao, L., Weiss, S., Hengst, L.,
and Kriwacki, R.W. (2004). p27 binds cyclin-CDK complexes through a sequential
mechanism involving binding-induced protein folding. Nat Struct Mol Biol *11*, 358-364.
10.1038/nsmb746.
72.   Tompa, P., and Fuxreiter, M. (2008). Fuzzy complexes: polymorphism and
structural disorder in protein-protein interactions. Trends Biochem Sci *33*, 2-8.
10.1016/j.tibs.2007.10.003.
73.   Brzovic, P.S., Heikaus, C.C., Kisselev, L., Vernon, R., Herbig, E., Pacheco, D.,
Warfield, L., Littlefield, P., Baker, D., Klevit, R.E., and Hahn, S. (2011). The acidic
transcription activator Gcn4 binds the mediator subunit Gal11/Med15 using a simple
protein interface forming a fuzzy complex. Mol Cell *44*, 942-953.
10.1016/j.molcel.2011.11.008.
74.   Milles, S., Mercadante, D., Aramburu, I.V., Jensen, M.R., Banterle, N., Koehler,
C., Tyagi, S., Clarke, J., Shammas, S.L., Blackledge, M., et al. (2015). Plasticity of an
ultrafast interaction between nucleoporins and nuclear transport receptors. Cell *163*,
734-745. 10.1016/j.cell.2015.09.047.
75.   Arbesu, M., Maffei, M., Cordeiro, T.N., Teixeira, J.M., Perez, Y., Bernado, P.,
Roche, S., and Pons, M. (2017). The Unique Domain Forms a Fuzzy Intramolecular
Complex in Src Family Kinases. Structure *25*, 630-640 e634. 10.1016/j.str.2017.02.011.
76.   Lin, Y.H., Qiu, D.C., Chang, W.H., Yeh, Y.Q., Jeng, U.S., Liu, F.T., and Huang,
J.R. (2017). The intrinsically disordered N-terminal domain of galectin-3 dynamically

mediates multisite self-association of the protein through fuzzy interactions. J Biol
Chem *292*, 17845-17856. 10.1074/jbc.M117.802793.

77.   Jin, F., Yu, C., Lai, L., and Liu, Z. (2013). Ligand clouds around protein clouds: a
scenario of ligand binding with intrinsically disordered proteins. PLoS Comput Biol *9*,
e1003249. 10.1371/journal.pcbi.1003249.

78.   Pufall, M.A., and Graves, B.J. (2002). Autoinhibitory domains: modular effectors
of cellular regulation. Annu Rev Cell Dev Biol *18*, 421-462.
10.1146/annurev.cellbio.18.031502.133614.

79.   Wright, P.E., and Dyson, H.J. (2015). Intrinsically disordered proteins in cellular
signalling and regulation. Nat Rev Mol Cell Biol *16*, 18-29. 10.1038/nrm3920.

80.   Mascle, X.H., Gagnon, C., Wahba, H.M., Lussier-Price, M., Cappadocia, L.,
Sakaguchi, K., and Omichinski, J.G. (2019). Acetylation of SUMO1 Alters Interactions
with the SIMs of PML and Daxx in a Protein-Specific Manner. Structure.
10.1016/j.str.2019.11.019.

81.   Zilio, N., Eifler-Olivi, K., and Ulrich, H.D. (2017). Functions of SUMO in the
Maintenance of Genome Stability. Adv Exp Med Biol *963*, 51-87. 10.1007/978-3-319-
50044-7_4.

82.   Hutten, S., Flotho, A., Melchior, F., and Kehlenbach, R.H. (2008). The Nup358-
RanGAP complex is required for efficient importin alpha/beta-dependent nuclear import.
Mol Biol Cell *19*, 2300-2310. 10.1091/mbc.E07-12-1279.

83.   Zhou, Z., Hartwieg, E., and Horvitz, H.R. (2001). CED-1 is a transmembrane
receptor that mediates cell corpse engulfment in C-elegans. Cell *104*, 43-56. Doi
10.1016/S0092-8674(01)00190-8.

84.   Barstead, R., Moulder, G., Cobb, B., Frazee, S., Henthorn, D., Holmes, J.,
Jerebie, D., Landsdale, M., Osborn, J., Pritchett, C., et al. (2012). Large-Scale
Screening for Targeted Knockouts in the Caenorhabditis elegans Genome. G3-Genes
Genom Genet *2*, 1415-1425. 10.1534/g3.112.003830.

85.   Werner, A., Flotho, A., and Melchior, F. (2012). The
RanBP2/RanGAP1*SUMO1/Ubc9 complex is a multisubunit SUMO E3 ligase. Mol Cell
46, 287-298. 10.1016/j.molcel.2012.02.017.

86.   Johnson, E.S., and Gupta, A.A. (2001). An E3-like factor that promotes SUMO
conjugation to the yeast septins. Cell *106*, 735-744. 10.1016/s0092-8674(01)00491-3.

87.   Warnecke, A., Sandalova, T., Achour, A., and Harris, R.A. (2014). PyTMs: a
useful PyMOL plugin for modeling common post-translational modifications. BMC
Bioinformatics *15*, 370. 10.1186/s12859-014-0370-6.

88.   Van der Spoel, D., Lindahl, E., Hess, B., Groenhof, G., Mark, A.E., and
Berendsen, H.J.C. (2005). GROMACS: Fast, flexible, and free. Journal of
Computational Chemistry *26*, 1701-1718. 10.1002/jcc.20291.

89.   Pettersen, E.F., Goddard, T.D., Huang, C.C., Couch, G.S., Greenblatt, D.M.,
Meng, E.C., and Ferrin, T.E. (2004). UCSF Chimera--a visualization system for
exploratory research and analysis. J Comput Chem *25*, 1605-1612. 10.1002/jcc.20084.

90.   Webb, B., and Sali, A. (2014). Comparative Protein Structure Modeling Using
MODELLER. Curr Protoc Bioinformatics *47*, 5 6 1-32. 10.1002/0471250953.bi0506s47.

91.   Saitoh, H., and Hinchey, J. (2000). Functional heterogeneity of small ubiquitin-
related protein modifiers SUMO-1 versus SUMO-2/3. J Biol Chem *275*, 6252-6258.
10.1074/jbc.275.9.6252.

92.   Flotho, A., Werner, A., Winter, T., Frank, A.S., Ehret, H., and Melchior, F.
(2012). Recombinant reconstitution of sumoylation reactions in vitro. Methods Mol Biol
832, 93-110. 10.1007/978-1-61779-474-2_5.

93.   Mahajan, R., Delphin, C., Guan, T., Gerace, L., and Melchior, F. (1997). A small
ubiquitin-related polypeptide involved in targeting RanGAP1 to nuclear pore complex
protein RanBP2. Cell *88*, 97-107.

94.   Schneider, C.A., Rasband, W.S., and Eliceiri, K.W. (2012). NIH Image to
ImageJ: 25 years of image analysis. Nat Methods *9*, 671-675. 10.1038/nmeth.2089.

1485 95. Delaglio, F., Grzesiek, S., Vuister, G.W., Zhu, G., Pfeifer, J., and Bax, A. (1995).
1486 NMRPipe: a multidimensional spectral processing system based on UNIX pipes. J
1487 Biomol NMR *6*, 277-293. 10.1007/BF00197809.
1488 96. Johnson, B.A., and Blevins, R.A. (1994). NMR View: A computer program for
1489 the visualization and analysis of NMR data. J Biomol NMR *4*, 603-614.
1490 10.1007/BF00404272.
1491 97. Naik, M.T., Chang, C.C., Naik, N.M., Kung, C.C., Shih, H.M., and Huang, T.H.
1492 (2011). NMR chemical shift assignments of a complex between SUMO-1 and SIM
1493 peptide derived from the C-terminus of Daxx. Biomol NMR Assign *5*, 75-77.
1494 10.1007/s12104-010-9271-4.
1495 98. Mossessova, E., and Lima, C.D. (2000). Ulp1-SUMO crystal structure and
1496 genetic analysis reveal conserved interactions and a regulatory element essential for
1497 cell growth in yeast. Mol Cell *5*, 865-876. 10.1016/s1097-2765(00)80326-3.
1498 99. Surana, P., Gowda, C.M., Tripathi, V., Broday, L., and Das, R. (2017). Structural
1499 and functional analysis of SMO-1, the SUMO homolog in Caenorhabditis elegans.
1500 PLoS One *12*, e0186622. 10.1371/journal.pone.0186622.
1501 100. Schrodinger, LLC (2015). The PyMOL Molecular Graphics System, Version 1.8.
1502 101. Papamokos, G.V., Tziatzos, G., Papageorgiou, D.G., Georgatos, S.D., Politou,
1503 A.S., and Kaxiras, E. (2012). Structural role of RKS motifs in chromatin interactions: a
1504 molecular dynamics study of HP1 bound to a variably modified histone tail. Biophys J
1505 *102*, 1926-1933. 10.1016/j.bpj.2012.03.030.
1506 102. Homeyer, N., Horn, A.H., Lanig, H., and Sticht, H. (2006). AMBER force-field
1507 parameters for phosphorylated amino acids in different protonation states:
1508 phosphoserine, phosphothreonine, phosphotyrosine, and phosphohistidine. J Mol
1509 Model *12*, 281-289. 10.1007/s00894-005-0028-4.
1510 103. Lindorff-Larsen, K., Piana, S., Palmo, K., Maragakis, P., Klepeis, J.L., Dror,
1511 R.O., and Shaw, D.E. (2010). Improved side-chain torsion potentials for the Amber
1512 ff99SB protein force field. Proteins *78*, 1950-1958. 10.1002/prot.22711.
1513 104. Piana, S., Donchev, A.G., Robustelli, P., and Shaw, D.E. (2015). Water
1514 dispersion interactions strongly influence simulated structural properties of disordered
1515 protein states. J Phys Chem B *119*, 5113-5123. 10.1021/jp508971m.
1516 105. Hess, B. (2008). P-LINCS: A Parallel Linear Constraint Solver for Molecular
1517 Simulation. J Chem Theory Comput *4*, 116-122. 10.1021/ct700200b.
1518 106. Han, B., Liu, Y., Ginzinger, S.W., and Wishart, D.S. (2011). SHIFTX2:
1519 significantly improved protein chemical shift prediction. J Biomol NMR *50*, 43-57.
1520 10.1007/s10858-011-9478-4.
1521 107. Brenner, S. (1974). The genetics of Caenorhabditis elegans. Genetics *77*, 71-
1522 94. 10.1093/genetics/77.1.71.
1523 108. Arribere, J.A., Bell, R.T., Fu, B.X., Artiles, K.L., Hartman, P.S., and Fire, A.Z.
1524 (2014). Efficient marker-free recovery of custom genetic modifications with
1525 CRISPR/Cas9 in Caenorhabditis elegans. Genetics *198*, 837-846.
1526 10.1534/genetics.114.169730.
1527 109. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch,
1528 T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-
1529 source platform for biological-image analysis. Nat Methods *9*, 676-682.
1530 10.1038/nmeth.2019.
1531 110. Chawla, D.G., Shah, R.V., Barth, Z.K., Lee, J.D., Badecker, K.E., Naik, A.,
1532 Brewster, M.M., Salmon, T.P., and Peel, N. (2016). Caenorhabditis elegans
1533 glutamylating enzymes function redundantly in male mating. Biol Open *5*, 1290-1298.
1534 10.1242/bio.017442.
1535

Figure 3