Discovery of giant viruses as past and present infections of zoosporic fungi

Keywords: giant virus, fungi, endogenous viral elements (EVEs), NCLDV, Nucleocytoviricota, mycovirus, zoosporic fungi

Jillian M. Myers[1], Frederik Schulz[2], Saleh Rahimlou[1], Vikas Yadav[3], Kevin R. Amses[1◊], D. Rabern Simmons[1◊], Sheng Sun[3], Michelle Orozco-Quime[1], Joseph Heitman[3], Jason E. Stajich[4✉], Timothy Y. James[1,5✉]

Author Affiliations:
[1]Ecology and Evolutionary Biology, University of Michigan, Ann Arbor, MI, USA
[2]DOE Joint Genome Institute, Lawrence Berkeley National Laboratory, Berkeley, CA, USA
[3]Molecular Genetics and Microbiology, Duke University School of Medicine, Durham, NC, USA
[4]Microbiology and Plant Pathology, University of California Riverside, Riverside, CA, USA
[5]University of Michigan Herbarium, Ann Arbor, MI, USA
◊current affiliations: University of Pennsylvania Perelman School of Medicine, Philadelphia, PA, USA (KRA), Botany and Plant Pathology, Purdue University, West Lafayette, Indiana, USA (DRS)
✉email: tyjames@umich.edu; jason.stajich@ucr.edu

Abstract

Giant viruses of the phylum *Nucleocytoviricota* have emerged as particularly notable due to their increasingly recognized impacts on eukaryotic genome evolution. Their origins are hypothesized to predate or coincide with the diversification of eukaryotes, and they have been detected in hosts that span the eukaryotic tree of life. But surprisingly, such viruses have not been definitively found in Kingdom Fungi, though genomic and metagenomic work suggests a putative association. Here we report both "viral fossils" and active infection by giant viruses in fungi, particularly in the zoosporic phyla Blastocladiomycota and Chytridiomycota. The recovered viral genomes span up to 350 kb, encode over 300 genes, and form a monophyletic family-level clade within the *Nucleocytoviricota* related to orders *Imitervirales* and *Algavirales*, which we name *Mycodnaviridae*. We observed variation in infection status among the isolates including apparent active infection and transcriptionally-suppressed states, suggesting that viral activation may be constrained to certain life stages of the host. Our experimental findings add to the limited natural virus-host systems available in culture for the study of giant viruses. These viruses may have shaped the early evolution of these fungal lineages by the shuttling of genes between major kingdoms and domains of life. Our findings expand the known host range of *Nucleocytoviricota* into a new kingdom that contains many model species. *Mycodnaviridae* have a global distribution, which invites inquiry into the implications of these infections for host traits, host genome evolution, and the metabolic impacts to ecosystems.

Main

In the entire Fungal Kingdom, only a handful of DNA viruses are known–all small circular single-stranded (CRESS) viruses (Li et al. 2020; Yu et al. 2010). This apparent rarity of DNA viruses in fungi likely results from taxonomic biases in searches, rather than a true biological phenomenon. The study of mycoviruses has historically focused on their application for biological control of fungal pathogens, particularly in agricultural contexts, and as a result most research has involved viruses infecting fungi in the Ascomycota—one of two phyla comprising the subkingdom Dikarya, which contains the majority of described fungal taxa. We previously demonstrated that by expanding the taxonomic scope of considered fungal hosts to include lineages outside of the Dikarya the known RNA mycoviral diversity was greatly increased (Myers et al. 2020). The current fungal taxonomic division includes eight phyla of zoosporic fungi that reproduce with motile spores and represent a particularly understudied paraphyletic assemblage that includes the last common ancestor of fungi (Hyde et al. 2024). The presence of DNA viruses in these fungi has been almost entirely unexplored (but see Clemons et al. 2023), and indeed, the finding of giant viruses, which we report here, was a serendipitous discovery resulting from a large genomic sequencing effort across these fungal phyla (Amses et al. 2022).

While only a few active DNA viruses in fungi have been documented, previous studies have suggested putative associations between *Nucleocytoviricota* and fungi through the identification of fungal homologs in metagenomic-assembled giant viruses (Schulz et al. 2018, Schulz et al. 2020, Bhattacharjee et al. 2023) and giant virus EVEs in fungal genomes (Gallot-Lavallée and Blanc 2017, Gong et al. 2020, Zhao et al. 2023, Nobre 2024). These links vary in their robustness but nonetheless the cumulative evidence suggests a rich evolutionary history between fungi and giant viruses. Despite these tantalizing findings, active giant viral infections in fungi have not previously been evidenced nor have novel clades that include active giant viruses in fungi been characterized.

Results

Zoosporic fungi contain giant virus genes and genomes with global distribution

We identified homologs of the giant virus major capsid protein (MCP; NCVOG0022) in the genomes of 17/141 fungi with HMM searches (Fig S1, Table S1). The distribution of MCP homologs is restricted to non-Dikaryotic lineages, and, particularly, the zoosporic fungi; no MCPs were found by hmmsearch in non-zoosporic lineages (n= 40). We further searched all fungal genomes in Mycocosm (Grigoriev et al. 2014; n = 2,543) for MCP homologs by blastp, which resulted in one hit to a non-zoosporic fungal species: *Dimargaris cristalligena* (phylum Zoopagomycota), a mycoparasite. Among the zoosporic lineages, MCPs were found in Chytridiomycota (n=12), Monoblephidomycota (n=2), and Blastocladiomycota (n=3). Blastocladiomycota are a particularly poorly sampled group, and thus our genomic searches

77   were limited (n=7). However, MCPs appear to be common in the most highly sampled genus
78   within this phylum, *Allomyces.* Based on available sequences, we designed primers targeting
79   the MCP gene of *Allomyces* and found 30/58 unique isolates were positive for capsid by PCR.
80
The contigs containing MCP genes vary widely among the fungal isolates in gene content, GC
composition, intergenic distance, and other metrics, suggesting numerous independent
endogenization events (Fig 1 A). In three isolates (*Podochytrium* sp. JEL797, *Gaertneriomyces*
semiglobifer Barr43, and *Fimicolochytrium jonesii* JEL569), MCP was the only gene of viral
provenance on contigs which otherwise contain primarily fungal genes and have GC content
and intron density similar to other host contigs (Fig S2). In other isolates, contigs with MCPs
appear as mosaics of virus, prokaryote, and fungal genes, which include multiple additional
giant virus orthologous genes (GVOGs), and have markedly greater gene density compared to
host contigs, consistent with reference giant virus genomes (Fig 1 A, B; Fig S2). These giant
virus-like contigs range up to nearly 350 kb and are suggestive of either very recent
endogenization or proviral state. Interestingly, in five isolates the MCP contig contains a gene
encoding an adintovirus-like integrase.
93
94   Based on MCP data previously extracted from global metagenomics datasets, we assessed the
95   global distribution of these novel fungal MCPs. Environmental MCP homologs with bidirectional
96   best blast hits to fungal MCPs have a global distribution (Fig 1 C). The *Allomyces* isolates tested
97   by PCR for MCP have worldwide provenance, thus also suggestive of a long history of viral
98   infection.
99
The most contiguously assembled viral genomes were found from two strains of *Allomyces.* A.
arbusculus Burma1F mycodnavirus 1 includes two contigs of total size 345,111 bp, encoding
314 genes with likely provenance, based on homology, assigned as follows: 10% fungi, 14%
bacteria, 20% virus, 4% putatively *Mycodnaviridae*-specific, and 52% unknown/ORFan (Fig 1
B). The isolate *A. javanicus* California12 has multiple large contigs (ranging between 80-288kb)
with giant virus signatures (Fig S3). Interestingly, when binned by tetramer frequency, the viral
contigs in *A. javanicus* California12 form two bins suggesting the presence of two distinct
viruses. The two bins are roughly similar in size (212,591 bp and 287,844 bp) and show some
duplication of hallmark GVOGs: each bin contains copies of MCP, D5-like helicase-primase,
DNA PolB, mRNA capping enzyme large subunit, and superfamily II helicase. The larger bin
contains an additional five GVOGs (YqaJ recombinase, RNA ligase, disulfide (thiol)
oxidoreductase, transcription initiation factor IIB, and A32-like packaging ATPase). *A. javanicus*
strains are believed to be hybrids (Emerson 1941) and the two viruses could have originated
from the two parents of this diploid sporophyte.
114

Fig 1. Genomic features of *Mycodnaviridae* (A) Boxplots of genome statistics for the fungal genomes
found to contain MCP genes (n=17): mean values across all contigs ("Fungal mean", purple), contigs
containing at least one major capsid protein (MCP) gene ("MCP contigs", pink), and MCP contigs that met
the criteria set for orthology analysis (ViralRecall score >2, viral region length > 70%, intron density >25%
different from host; "*Mycodnaviridae*", teal). (B) The genome of Allomyces arbusculus Burma1F
mycodnavirus 1 is a mosaic of bacteria; viral, and eukaryotic genes. Contigs are represented by the
outermost ring in the circularized genome plot. From outside to inside: ORFs on the forward strand, ORFs
on the reverse strand (both color coded by provenance), GC skew. (C) Global distribution of the
Mycodnaviridae-like sequences based on the major capsid protein. Colored circles correspond to
metagenomic MCP sequences with best blast hit (BBH) to *Mycodnaviridae* identified in this study and
circles with a black outline indicate geographic location of *Mycodnaviridae* and hosts described in this
study.

Mycodnaviridae is a monophyletic clade of fungal giant viruses

A phylogeny of representative giant viruses including the 7 fungal samples with at least 4/7 hallmark *Nucleocytoviricota* genes (GVOG7 marker set; Aylward et al. 2021) show that fungal giant viruses form a monophyletic clade together with three giant virus metagenome-assembled genomes (GVMAGs) of the previously proposed *Imitervirales* family IM_02 and in a sister position to *Imitervirales* IM_18 (Fig 2A). Our phylogenetic analysis showed that less complex phylogenetic models (WAG) and reconstruction algorithms (FastTree, IQ-tree2 -fast) place the IM_02-IM_18 cluster basal to other *Imitervirales*. However, with more complex phylogenetic models (IQtree2 with LG+I+G4, LG4X) the IM_02-IM_18 cluster was split off from *Imitervirales* and groups sister to *Pandoravirales* and *Algavirales*. The monophyletic grouping of the fungal giant viruses with IM_02 was well supported across supermatrix-based species trees and single protein trees for each of the markers in the GVOG7 set. Both IM_02 and IM_18 previously consisted exclusively of GVMAGs but since the new genomes added can be connected with fungal hosts, we dub IM_02 *Mycodnaviridae* (Fig 2 A). Our analysis also suggests that IM_02 together with IM_18 is not part of the order *Imitervirales* but may represent a novel order in the *Nucleocytoviricota*. Further detailed phylogenomic analysis after improved taxon sampling will be required to provide additional evidence for such a relationship. The hosts of IM_02 and IM_18 are unknown, but the clustering of these clades with giant viruses from fungi strongly suggest a fungal association.

We investigated whether a core suite of genes exist that functionally define the *Mycodnaviridae* by conducting orthology analysis on the most complete viral genomes–five have at least 9 of the 20 core GVOGs present in the majority of the *Nucleocytoviricota* (Yutin et al. 2009). This resulted in 27 gene clusters shared by all five (Fig 2 B) though this is likely an underestimate due to variation in genome completeness, which is suggested by variation in viral genome assembly size. In addition to GVOGs such as MCPs, VLTF3, and YqaJ-like recombinase, the core set is comprised of gene orthogroups involved in DNA biosynthesis such as DNA helicase, ribonucleotide reductase, and proliferating cell nuclear antigen, which are also GVOGs found in other large and giant viruses. Other shared gene clusters include a cysteine protease, protein with N-acetylglucosaminyltransferase activity (possibly involved in cell wall modification), six genes of unknown function, and five novel genes apparently unique to these viruses.

162 **Fig 2. Phylogenetic position of the *Mycodnaviridae* and core genes**. (A) Position of the
163 *Mycodnaviridae* clade in the species tree of the *Nucleocytoviricota*. Red branches indicate
164 *Mycodnaviridae* genomes. The tree is rooted at Pokkesviricetes. (B) Venn diagram indicating the number
165 of orthologous gene clusters shared between the 5 most complete *Mycodnaviridae* genomes.

Allomyces virus is integrated in the nuclear genome, with the capacity for productive infection

One critical question was whether these giant virus contigs are merely relics of past infections
that have been maintained as EVEs in fungal genomes, or whether they are capable of
replicative infection. We have accumulated evidence that supports both scenarios. In *Allomyces*
arbusculus Burma1F, long-read whole-genome sequencing by Oxford Nanopore revealed the
viral sequence integrated into a chromosome that harbors telomere repeats on both ends
(contig 03: 1397320-1743578; (Figure 3 A & B). Nanopore-based genome assembly also
revealed 13 clusters of LTR-transposons, only one per contig, that potentially represent
centromeres similar to those observed in the dikaryotic fungus *Cryptococcus neoformans*
(Yadav 2018, Janbon 2014). Furthermore, 26 telomere repeat-containing ends were identified
across all contigs, which combined with 13 putative centromeres, indicate Burma1F harbors 13
chromosomes. With interest in whether the viral genes are expressed and, if so, whether
expression occurs throughout host tissue or is constrained to some structures or life stages, we
sequenced RNA of *A. arbusculus* Burma1F at three different stages. We found almost no
evidence of expression from the viral genes, with the exception of genes in the flanking region
(Fig 3 B). Interestingly, CpG methylation was not detected in these viral regions per the
Nanopore-generated data. These data suggest active silencing of the viral genes by the host,
and the expression data help define the border of the integrated virus. The mechanism of
silencing employed by the host is as yet unknown.

An additional distinguishing feature between viruses and EVEs is read coverage of the viral
contigs because sequencing during viral replication is expected to result in extremely high
coverage of viral reads. Overall, we found no clear pattern between coverage of fungal contigs
containing MCPs and coverage of other genomic contigs, with one exception. We sequenced
genomes of an additional 9 strains of *Allomyces;* one isolate, *Allomyces arbusculus* Cali8,
included 8 contigs (total 309,231 bp) that averaged nearly 50-fold higher coverage than the rest
of the genome (Fig 3 C). Of those, two contigs include MCPs and a combined 7 additional core
GVOGs as well as a viral-like integrase. These results strongly indicate an active infection by a
giant virus in this strain of *Allomyces.* Other sequenced *Allomyces* strains had contigs similarly
rich with giant virus hallmark genes but lacking differences in coverage from the genome-wide
median estimated by read depth. It is plausible that in these strains the conditions for triggering
viral release were not met under laboratory culture conditions, and thus the virus remained in a
proviral state. Given the high occurrence of MCPs we suspect that additional *Allomyces* strains
contain provirus that is perhaps capable of activating and forming an infectious propagule,
though the triggering mechanism for viral release is unknown.
202

203
Figure 3. *Allomyces* viruses are integrated into the nuclear genome, with complex patterns of expression/suppression and phenotypic effects. (A) Assembly of *A. arbusculus* Burma1F shows the virus integrated into the host nuclear genome (Ctg 03) (B) Mapping of Nanopore reads (genome coverage), RNAseq reads, and CpG methylation extracted from Nanopore sequencing reads across the virus-harboring contig of *A. arbusculus* Burma1F, showing silencing of the integrated virus unrelated to methylation. (C) Plots of the GC content and genomic read coverage for assemblies of *A. anomalus* BEA2 (upper) and *A. arbusculus* Cali8 (lower). Contigs that include a major capsid protein (MCP) are shaded blue and size of data points reflect contig length. Cali8 includes two contigs with both MCPs and very high coverage. (D) Rolling average of normalized read coverage of a non-viral (upper) and viral (lower) genome segment in three isolates of Cali8. (E) Cali8-N gametophytes display normal phenotype with both male and female gametangia while Cali8-V nearly exclusively produces malformed and unproductive male gametangia (F) Cali8-N and Cali8-V grown at 37 C and 22C. At both temperatures Cali8-V had dramatically reduced growth rate.

In an initial effort to assess functional impacts of giant virus presence, strains of *Allomyces* with
and without detectable MCP genes were grown on agar and liquid Emerson's YpSS media to
test if the presence of MCP influenced growth rate of the fungal strain. No significant differences
in growth rate (t(10) = -1.10, p = 0.30) and biomass accumulation (t(10) = -0.06, p = 0.95) were
observed (Fig S4, A & B). This is likely due to the absence of expression of virus genes when
present in the fungal chromosome (as in Fig 3 B). We also compared the thermal maximum of
Allomyces strains with and without MCP grown on solid YpSS media. We found no significant
difference (t(10) = -0.25, p = 0.80; Fig S4 C) depending on presence of MCP, however, one of
the strains, Cali8, showed sectoring at 42 C. With archived samples of Cali8, we generated
meiospores by germinating resting sporangia. These spores mostly showed abortive growth,
however, two single spore isolates with divergent morphologies were obtained (Cali8-V and
Cali8-N) (Fig 3 E). Cali8-V mycelium was consistent with a gametophyte stage, however, the
strain primarily produced male gametangia that were enlarged, fewer in number than Cali8-N,
and no gametes could be released. In comparison, the Cali8-N mycelium appeared normal, and
growth rates of Cali8-N at both 22 C and 37 C were much higher than Cali8-V (Fig 3 E & F).
These two strains were then subjected to whole genome sequencing in an effort to discern
whether viral titer was responsible for the altered morphology. After normalization, average
coverage of the viral region in the sector demonstrating abnormal hyphal morphology was 25
times greater than the sector with normal growth; this region in the originally sequenced
Allomyces arbusculus Cali8 strain (polysporous isolate) had 490 times greater coverage (Fig 3
D). These results highlight that viral release is complicated and may be linked with
environmental and host lifecycle triggers as well as the complex life history of this fungus. These
results also indicate that the host possesses mechanisms for avoiding viral release and/or
restricting the spread of viral particles.

Transmission electron microscopy reveals virus-like particles

Arguably the presence of active viruses is most directly evidenced by images of the viral
particles themselves. We took two approaches to this end: we searched the literature on
zoospore ultrastructure for unrecognized or mis-identified electron-dense bodies that could be of
viral origin and also prepared zoospores of *Allomyces* for transmission electron microscopy
(TEM). Both approaches resulted in electron micrographs of giant virus-like particles with similar
morphologies (VLPs, Fig 4).
248
The literature on zoosporic fungi ultrastructure is rich with unidentified bodies with uncertain
functions. Zoospores are relatively simple, single cells, and yet are home to unique organelles,
vacuoles, lipid bodies, and inclusions that have proven difficult to assign a functionality. Indeed,
some of these appear remarkably virus-like. Letcher and Powell (2018) describe a chytrid
isolate *Polyphlyctis willoughbyi* PL AUS 26, writing "occurring sparsely in the peripheral
cytoplasm were electron-opaque hexagonal crystals ~0.1 µm in the broadest dimension" (Fig 4
C-E). Simmons et al (2009) describe the chytrid *Clydaea vesicula*, which is so named for the
"aggregation of vesicles with electron-opaque contents" within the zoospore cytoplasm (Fig 4 F
& G). Our in-house transmission electron microscopy of *A. javanicus* California12 zoospores
also shows cytoplasmic VLPs (Fig 4 A & B). The VLPs are round to hexagonal, 150-175 nm
diameter with an approximately 100 nm granular core, and appear in low abundance in

zoospores. Taken together, it is striking that the VLPs of these three zoosporic fungi converge
on a shared morphology, which may represent that of the *Mycodnaviridae*. We report these
findings with caution, however, as VLPs have not yet been isolated and sequenced directly.
263

Fig 4. Cross-sections of zoospores including virus-like particles (VLPs) visualized by
transmission electron microscopy. (A) *A. javanicus* California12 zoospore cross-section at 2500x
containing a VLP (arrow) near the nuclear cap (nc), and (B) the VLP at 15000x magnification. (C) Cross-
section of *Polyphlyctis willoughbyi* PL AUS 26 zoospore containing "hexagonal inclusions", magnified in
(D & E). (F) *Clydaea vesicula* cross-section with electron-dense vesicles (Ves, arrow), magnified in (G).
Images reused with permission (C & E, Elsevier license number 5995480828397; F & G, Elsevier license
number 5995480495911).

Discussion

In this paper, we report the discovery of *Mycodnaviridae*, a new family of *Nucleocytoviricota* with
close phylogenetic affinities to *Algavirales* and *Imitervirales*. The three phyla of fungi in which
we have found evidence of giant viruses—Blastocladiomycota, Chytridiomycota, and
Zoopagomycota—have life cycle stages similar to some algae and amoebae, which may
facilitate infection by viruses with persistent infection strategies, such as the phaeoviruses in
Algavirales. In both algae and zoosporic fungi there exist life-stages (spores and gametes)
consisting of flagellated single-cells lacking a protective cell wall. The phaeoviruses have
evolved stable coexistence with their hosts through a persistent viral strategy whereby these
spores or gametes are infected by virus particle(s) that then integrate into the host genome

(Müller and Parodi 1993). Development proceeds with one copy of the virus integrated into the
DNA of each cell (Meints et al. 2008). Thalli often appear symptom-free until some unknown
mechanism triggers virus emergence from the genome. Reproductive organs may develop
irregularly as they become factories for viral replication (Henry and Meints 1992; Stevens et al.
2014). Very recently, another integrated viral genome was identified within a green alga and
was shown to actively produce viral particles (Erazo-Garcia 2025). We hypothesize that fungal
giant viruses in zoosporic lineages share similar replication cycle characteristics.

Genome coverage data, transcriptional patterns, and the presence of an integrase gene in
Mycodnaviridae are consistent with this hypothesized replication cycle. In most genomes viral
contigs have similar coverage to the host average, as would be expected for a provirus
integrated into a single-celled spore/gamete that propagates the organism. Genome coverage
data in one isolate of *Allomyces* (Cali8), however, suggests active replication at the time of initial
sequencing. Subcultures of the isolate are consistent with either viral endogenization into the
nuclear genome or near-complete loss of the virus. In another isolate of *Allomyces* (Burma1F),
transcriptional patterns suggest that the provirus may be silenced in the host genome,
consistent with transcriptional patterns of phaeoviruses integrated in their algal hosts (Cock et
al. 2010). The reason that we failed to observe any viral effect on fungal host phenotypes in this
isolate may be related to this absence of viral gene expression.
299
Electron micrographs of putatively infected zoospores show virus-like particles (VLPs) with
similar morphology to the phaeoviruses: round to hexagonal VLPs of 150-175 nm diameter with
an approximately 100 nm granular core. VLPs typically appear in low abundance in zoospores,
consistent with expectations that zoospores are recipients of infective viruses that migrate to the
nucleus and, upon zoospore encystment, integrate into the host genome. The apparent near-
restriction of *Mycodnaviridae* to zoosporic lineages is consistent with this hypothesis for the
replicative cycle, though a mechanism triggering viral release remains unknown.

Like all cellular organisms, the evolution of the fungi has been impacted by interactions with
viruses. Giant viruses are hypothesized to have originated with the diversification of eukaryotes,
and metagenomic sequence data demonstrates their abundance in the environment. The lack of
evidence of active infection by giant viruses in fungi before now is therefore surprising, though
previous reports have shown their association (Zhao et al. 2023; Bhattacharjee et al. 2023;
Gong 2020; Schulz et al. 2020; 2018). Most interestingly, the EVEs reported by others are of
distinct and distant relation to the clade of giant fungal viruses we report here. Phylogenetic
reconstruction of the DNA polymerase B (PolB) using data from four such studies as well as
Mycodnaviridae shows at least five unique clades of giant virus PolBs associated with Fungi
(Supplemental Figure S5). Taken together, these data suggest that fungi have historically and,
perhaps, currently been hosts to multiple distinct lineages of giant viruses.

It is intriguing to consider the implications of a possible ancient association between fungi and
giant viruses in relation to fungal evolution because viral gene acquisition can be an impetus for
evolutionary innovation (Frank and Feschotte 2017). In the fungi, a transcription factor involved
in cell cycle regulation that is, surprisingly, unique to fungi is hypothesized to have been
acquired by the fungal ancestor through lateral gene transfer from a virus (Medina et al. 2016).

Phylogenetic evidence points to the source as either a giant virus or phage-infected bacterium. We searched MCP-containing contigs but found no homologs containing the diagnostic viral-like KilA-N domain of this cell-cycle protein, though this does not preclude giant viruses as the original source. Other evolutionary innovations resulting from giant virus association could involve novel metabolic functions. Especially in GVMAGs, a vast diversity of metabolic genes has been found (Schulz et al. 2020, Moniruzzaman et al. 2020) including many involved in carbon and nitrogen metabolism. Assessing the genetic and functional impact of giant virus infection on fungal metabolism should be a future research priority. To understand the impact of the viruses, we need to resolve when and under what conditions the viruses are expressed during the life cycle.

Notably, the most contiguous viral genomes were found in *Allomyces* spp., from cultures that were recovered from filter paper stocks after 30 years of storage. This fact demonstrates desirable aspects of this fungus that bolster motivation for its revival as a model system. *Allomyces* has historically, and currently though with much less frequency, been a model for studying biological phenomena such as anisogamy (Olsen 1984; Phadke 2018), sensorimotor systems including phototaxis and chemotaxis (Machlis 1969; Swafford and Oakley 2018), and even cellular differentiation (Cantino and Lovett 1964). The development of *Allomyces* as a model for studying giant viruses has enormous potential. Much of what is known about giant virus biology has been learned from protist-virus interactions or metagenomics where the host is unknown. The development of additional laboratory model systems of giant viruses and their natural hosts could advance our understanding of virus-host interactions, including immune response, viral "reprogramming" of host cells, metabolic impacts, and their ecosystem-level consequences.

The discovery of the first giant viruses in fungi that are restricted primarily to the zoosporic fungi inspire numerous questions that beg further investigation. How is viral expression and virion production related to the host life cycle, and vice-versa? What are the metabolic impacts of infection, and how important are they on the global scale? What defense systems do fungi employ against giant viruses, and do they explain why Dikarya fungi are not infected? To what extent have giant viruses shaped genome evolution in fungi? Giant virus model systems in zoosporic fungi may be the key to exposing these latent revelations.

Data Availability

Mycodnaviridae genome assemblies and alignment for the *Nucleocytoviricota* phylogeny were deposited in Deep Blue Data, the University of Michigan's institutional data repository (DOI: https://doi.org/10.7302/p5ay-ng61). Fungal genome assemblies, and RNAseq data are available in NCBI BioProject PRJNA1257431.

Acknowledgements

Cultures and DNAs were obtained from the Collection of Zoosporic Eufungi at the University of Michigan. Next-generation sequencing was performed at the Advanced Genomics Core at the

University of Michigan. Transmission electron microscopy was conducted at the University of Michigan's Department of Molecular, Cell, and Developmental Biology Imaging Core; we heartily thank Gregg Sobocinski for his technical assistance with sectioning and training on the Jeol. We also thank Dr. Pete Letcher for contributing previously prepared grids and blocks of embedded zoospores for TEM. We thank Mia Sinks for technical assistance with *Allomyces* cultures. This project was funded in part by the National Science Foundation grants DEB-1929738 and DEB-2403677 to TYJ. KRA was supported by the following National Institutes of Health training grant: "Michigan Predoctoral Training in Genetics" [T32GM007544]. JES, TYJ, and JH are fellows and co-director (JH) of the CIFAR program Fungal Kingdom: Threats & Opportunities, and these studies were supported in part by a CIFAR Catalyst Award. The work conducted by the U.S. Department of Energy Joint Genome Institute (https://ror.org/04xm1d337), a DOE Office of Science User Facility, is supported by the Office of Science of the U.S. Department of Energy operated under Contract No. DE-AC02-05CH11231. Efforts by SS, VY, and JH were supported in part by NIH/NIAID R01 grants AI39115-27 and AI050113-20.

Competing Interests

FS is the co-founder and CEO of SampleX, a decentralized marketplace for biological samples, and co-founder and CTO of BioKEA, a consulting company for AI in biology. These affiliations did not influence the design, execution, or interpretation of the research presented in this manuscript.

Author Contributions

JES, TYJ, and JMM conceived of and planned the study. Culturing and DNA preparation was conducted by MOQ and JMM. RNA preparation, PCR, transcriptome assembly, viral content identification, and orthology analysis was conducted by JMM. Giant virus phylogeny and distribution analyses were performed by FS. Genome assembly and annotation was conducted by JMM, KRA, and SR. Transcriptome analysis was conducted by SR. Nanopore sequencing DNA extraction, as well as genome assembly and analysis of strain Burma1F was conducted by VY, SS and JH. Summary statistics were conducted by JES and JMM. Growth experiments were performed by TYJ. Transmission electron microscopy was performed by JMM and DRS. Manuscript was written by JMM with contributions from TYJ and FS. All authors read and approved the final manuscript.

References

Amses, Kevin R., D. Rabern Simmons, Joyce E. Longcore, Stephen J. Mondo, Kensuke Seto, Gustavo H. Jerônimo, Anne E. Bonds, et al. 2022. "Diploid-Dominant Life Cycles Characterize the Early Evolution of Fungi." *Proceedings of the National Academy of Sciences* 119 (36): e2116841119. https://doi.org/10.1073/pnas.2116841119.

Aylward, Frank O., Mohammad Moniruzzaman, Anh D. Ha, and Eugene V. Koonin. 2021. "A Phylogenomic Framework for Charting the Diversity and Evolution of Giant Viruses." *PLOS Biology* 19 (10): e3001430. https://doi.org/10.1371/journal.pbio.3001430.

Bhattacharjee, Ananda S., Frederik Schulz, Tanja Woyke, Beth N. Orcutt, and Joaquín Martínez Martínez. 2023. "Genomics Discovery of Giant Fungal Viruses from Subsurface Oceanic Crustal Fluids." *ISME Communications* 3 (1): 1–9. https://doi.org/10.1038/s43705-022-00210-8.

Cantino, Edward, and Lovett. 1964. "Non-Filamentous Aquatic Fungi: Model Systems for Biochemical Studies of Morphological Differentiation." In *Advances in Morphogenesis*, 3:33–93. Elsevier. https://doi.org/10.1016/B978-1-4831-9950-4.50005-9.

Clemons, Rebecca, Mark Yacoub, Evelyn Faust, L. Felipe Toledo, Thomas S. Jenkinson, Erik Kalinka, Lillian K. Fritz-Laylin, Timothy Y. James, and Jason E. Stajich. 2023. "DNA Virus BdDV-1 of the Amphibian Pathogen Batrachochytrium Dendrobatidis Is Associated with Hypervirulence." *bioRxiv*, March, 2023.03.16.532857. https://doi.org/10.1101/2023.03.16.532857.

Cock, J. Mark, Lieven Sterck, Pierre Rouzé, Delphine Scornet, Andrew E. Allen, Grigoris Amoutzias, Veronique Anthouard, et al. 2010. "The Ectocarpus Genome and the Independent Evolution of Multicellularity in Brown Algae." *Nature* 465 (7298): 617–21. https://doi.org/10.1038/nature09016.

Emerson, R. (1941) An Experimental Study of the Life Cycles and Taxonomy of Allomyces. Lloydia, 4, 77-144.

Erazo-Garcia, Maria P, Uri Sheyn, Zachary Barth, Rory Craig, Petronella Wessman et al. 2025. "Cryptic infection of a giant virus in a unicellular green alga". *Science* 0, eads6303 https://doi.org/ 10.1126/science.ads6303

Frank, JA, and Feschotte. 2017. "Co-Option of Endogenous Viral Sequences for Host Cell Function." *Current Opinion in Virology* 25 (August): 81–89. https://doi.org/10.1016/j.coviro.2017.07.021.

Gallot-Lavallée, L, and G Blanc. 2017. "A Glimpse of Nucleo-Cytoplasmic Large DNA Virus Biodiversity through the Eukaryotic Genomics Window." Viruses 9 (1). https://doi.org/10.3390/v9010017.

Gong, Zhen, Yu Zhang, and Guan-Zhu Han. 2020. "Molecular Fossils Reveal Ancient Associations of dsDNA Viruses with Several Phyla of Fungi." *Virus Evolution* 6 (1). https://doi.org/10.1093/ve/veaa008.

Grigoriev, Igor V., Roman Nikitin, Sajeet Haridas, Alan Kuo, Robin Ohm, Robert Otillar, Robert Riley, et al. 2014. "MycoCosm Portal: Gearing up for 1000 Fungal Genomes." *Nucleic Acids Research* 42 (D1): D699–704. https://doi.org/10.1093/nar/gkt1183.

Henry, Eric C., and Russel H. Meints. 1992. "A PERSISTENT VIRUS INFECTION IN FELDMANNIA (PHAEOPHYCEAE)1." *Journal of Phycology* 28 (4): 517–26. https://doi.org/10.1111/j.0022-3646.1992.00517.x.

Hyde, KD, MT Noorabadi, V Thiyagaraja, MQ He, PR Johnston, et al. 2024. The 2024 Outline of *Fungi* and fungus-like taxa. *Mycosphere.* 15(1): 5146-6239. https://doi.org/10.5943/mycosphere/15/1/25

Janbon, Guilhem, Kate L. Ormerod, Damien Paulet, Edmond J. Byrnes III, Vikas Yadav, Gautam Chatterjee, Nandita Mullapudi, et al. 2014. "Analysis of the Genome and Transcriptome of *Cryptococcus neoformans* var. *grubii* Reveals Complex RNA

444   Expression adn Microevolution Leading to Virulence Attenuation. *PLoS Genetics* 10(4):
445   e100426. https://doi.org/10.1371/journal.pgen.1004261

446 Letcher, Peter and Martha Powell. 2018. "Morphology, zoospore ultrastructure, and
447   phylogenetic position of *Polyphlyctis willoughbyi,* a new species in *Chytridiales*
448   *(Chytridiomycota).*" *Fungal Biology* 122: 1171-1183.
449   https://doi.org/10.1016/j.funbio.2018.08.003

450 Li, Pengfei, Shuangchao Wang, Lihang Zhang, Dewen Qiu, Xueping Zhou, and Lihua Guo.
451   2020. "A Tripartite ssDNA Mycovirus from a Plant Pathogenic Fungus Is Infectious as
452   Cloned DNA and Purified Virions." *Science Advances* 6 (14): eaay9634.
453   https://doi.org/10.1126/sciadv.aay9634.

454 Machlis, Leonard. 1969. "Fertilization-Induced Chemotaxis in the Zygotes of the Watermold
455   Allomyces." *Physiologia Plantarum* 22 (2): 392–400. https://doi.org/10.1111/j.1399-
456   3054.1969.tb07391.x.

457 Medina, Edgar M., Jonathan J. Turner, Raluca Gordân, Jan M. Skotheim, and Nicolas E.
458   Buchler. 2016. "Punctuated Evolution and Transitional Hybrid Network in an Ancestral
459   Cell Cycle of Fungi." eLife. eLife Sciences Publications Limited. May 10, 2016.
460   https://doi.org/10.7554/eLife.09492.

461 Meints, Russel H., Richard G. Ivey, Amy M. Lee, and Tae-Jin Choi. 2008. "Identification of Two
462   Virus Integration Sites in the Brown Alga Feldmannia Chromosome." *Journal of Virology*,
463   February. https://doi.org/10.1128/jvi.01983-07.

464 Moniruzzaman, Mohammad, Carolina A. Martinez-Gutierrez, Alaina R. Weinheimer, and Frank
465   O. Aylward. 2020. "Dynamic Genome Evolution and Complex Virocell Metabolism of
466   Globally-Distributed Giant Viruses." *Nature Communications* 11 (1): 1–11.
467   https://doi.org/10.1038/s41467-020-15507-2.

468 Moniruzzaman, Mohammad, Alaina R. Weinheimer, Carolina A. Martinez-Gutierrez, and Frank
469   O. Aylward. 2020. "Widespread Endogenization of Giant Viruses Shapes Genomes of
470   Green Algae." *Nature* 588 (7836): 141–45. https://doi.org/10.1038/s41586-020-2924-2.

471 Müller, D. G., and Elisa Parodi. 1993. "Transfer of a Marine DNA Virus fromEctocarpus
472   toFeldmannia (Ectocarpales, Phaeophyceae): Aberrant Symptoms and Restitution of the
473   Host." *Protoplasma* 175 (3): 121–25. https://doi.org/10.1007/BF01385009.

474 Myers, J. M., A. E. Bonds, R. A. Clemons, N. A. Thapa, D. R. Simmons, D. Carter-House, J.
475   Ortanez, et al. 2020. "Survey of Early-Diverging Lineages of Fungi Reveals Abundant
476   and Diverse Mycoviruses." *mBio* 11 (5): e02027-20. https://doi.org/10.1128/mBio.02027-
477   20.

478 Nobre SVA, de Andrade GAK, Metz GF, Lucini F, de Albuquerque MP, Victória FC. Antarctica's
479   hidden mycoviral treasures in fungi isolated from mosses: A first genomic approach. J
480   Basic Microbiol. 2024 Jul;64(7):e2300671. doi: 10.1002/jobm.202300671. Epub 2024
481   May 12. PMID: 38736205.

482 Olsen, Lauritz. 1984. *Allomyces: A Different Fungus*. Opera Botanica 73. Council for Nordic
483   Publications in Botany.

484 Phadke, Sujal S., Shawn M. Rupp, and Melissa A. Wilson Sayres. 2018. "Evolution of
485   Anisogamy in the Early Diverging Fungus, Allomyces." *bioRxiv*, November, 230292.
486   https://doi.org/10.1101/230292.

487 Saito, Makoto, Peiyu Xu, Guilhem Faure, Samantha Maguire, Soumya Kannan, Han Altae-Tran,
488   Sam Vo, AnAn Desimone, Rhiannon K. Macrae, and Feng Zhang. 2023. "Fanzor Is a
489   Eukaryotic Programmable RNA-Guided Endonuclease." *Nature* 620 (7974): 660–68.
490   https://doi.org/10.1038/s41586-023-06356-2.

491 Schulz, Frederik, Lauren Alteio, Danielle Goudeau, Elizabeth M. Ryan, Feiqiao B. Yu, Rex R.
492   Malmstrom, Jeffrey Blanchard, and Tanja Woyke. 2018. "Hidden Diversity of Soil Giant
493   Viruses." *Nature Communications* 9 (1): 1–9. https://doi.org/10.1038/s41467-018-07335-
494   2.

Schulz, Frederik, Simon Roux, David Paez-Espino, Sean Jungbluth, David A. Walsh, Vincent J.
Denef, Katherine D. McMahon, et al. 2020. "Giant Virus Diversity and Host Interactions
through Global Metagenomics." *Nature* 578 (7795): 432–36.
https://doi.org/10.1038/s41586-020-1957-x.
Simmons, D. Rabern, TY James, AF Meyer, and JE Longcore. 2009. "*Lobulomycetales,* a new
order in the *Chytridiomycota*". Mycological Research 113: 450-460.
doi:10.1016/j.mycres.2008.11.019
Starrett, Gj, Mj Tisza, Nl Welch, Ak Belford, A Peretti, Dv Pastrana, and Cb Buck. 2021.
"Adintoviruses: A Proposed Animal-Tropic Family of Midsize Eukaryotic Linear dsDNA
(MELD) Viruses." *Virus Evolution* 7 (1). https://doi.org/10.1093/ve/veaa055.
Stevens, Kim, Karen Weynberg, Christopher Bellas, Sonja Brown, Colin Brownlee, Murray T.
Brown, and Declan C. Schroeder. 2014. "A Novel Evolutionary Strategy Revealed in the
Phaeoviruses." *PLOS ONE* 9 (1): e86040. https://doi.org/10.1371/journal.pone.0086040.
Swafford, Andrew J. M., and Todd H. Oakley. 2018. "Multimodal Sensorimotor System in
Unicellular Zoospores of a Fungus." *Journal of Experimental Biology* 221 (2).
https://doi.org/10.1242/jeb.163196.
Yadav, Vikas, Sheng Sun, R. Blake Billmyre, and Kaustav Sanyal. 2018. "RNAi is a critical
determinant of centromere evolution in closely related fungi." *Proceedings of the*
National Academy of Sciences 115 (12): 3108-3113.
https://doi.org/10.1073/pnas.1713725115
Yu, Xiao, Bo Li, Yanping Fu, Daohong Jiang, Said A. Ghabrial, Guoqing Li, Youliang Peng, et
al. 2010. "A Geminivirus-Related DNA Mycovirus That Confers Hypovirulence to a Plant
Pathogenic Fungus." *Proceedings of the National Academy of Sciences* 107 (18): 8387–
92. https://doi.org/10.1073/pnas.0913535107.
Yutin, Natalya, Yuri I. Wolf, Didier Raoult, and Eugene V. Koonin. 2009. "Eukaryotic Large
Nucleo-Cytoplasmic DNA Viruses: Clusters of Orthologous Genes and Reconstruction of
Viral Genome Evolution." *Virology Journal* 6: 223. https://doi.org/10.1186/1743-422X-6-
223.
Zhao, Hongda, Ruixuan Zhang, Junyi Wu, Lingjie Meng, Yusuke Okazaki, Hiroyuki Hikida, and
Hiroyuki Ogata. 2023. "A 1.5-Mb Continuous Endogenous Viral Region in the Arbuscular
Mycorrhizal Fungus Rhizophagus Irregularis." *Virus Evolution* 9 (2).
https://doi.org/10.1093/ve/vead064.
527
528

Methods

Fungal strains, nucleic acid preparation, and sequencing:

We initially searched 135 genomes in this study including 131 fungi, 92 of which are zoosporic *(Table S1)*. For full details about DNA preparation and sequencing of these isolates see (Amses et al. 2022). An additional nine strains of *Allomyces* spp. were whole-genome sequenced from DNA archived in the CZEUM collection (Simmons et al. 2020). Library preparation and sequencing was performed by the Advanced Genomics Core at the University of Michigan by NovaSeq S4 300 cycle on 4% of a flowcell.

Allomyces javanicus California12 and *Allomyces arbusculus* Burma1F were grown in PmTG liquid media for 7-10 days at 22℃ with shaking at 180 rpm. Tissue was subsampled with sterile tweezers and immediately frozen and ground to a powder in liquid nitrogen. Approximately 50 mg was apportioned to 1.5mL tubes, 1 mL of TRIzol applied, and stored at -20℃ until further processing. For *A. arbusculus* Burma1F, the remaining tissue was decanted, rinsed once with sterile water, and replenished with sterile water. After overnight incubation at 22℃, tissue was decanted, rinsed once with sterile water, subsampled and processed as described above, then transferred to a sterile tea strainer and submerged in 50 mL sterile water for two hours on the benchtop. The tea strainer was removed, and the remaining liquid observed under a microscope for zoospore presence, followed by centrifugation at 4000x g and 4℃ for 12 minutes. The zoospore pellet was decanted, 1mL of TRIzol added, ground with a pestle, then stored at -20℃. The RNA extraction for all samples was completed per manufacturer protocol. Total RNAs were quality-checked on a Tapestation and had RIN values between 6.8-10. Samples were Poly-A enriched, and libraries were prepared and sequenced by the University of Michigan Advanced Genomics Core by NovaSeq S4 300 cycle on 1% of the flowcell.

Genome and transcriptome assembly, annotation, and statistics:

Transcriptome data were quality filtered by removing reads with Phred score <20 using Fastx-toolkit (Hannon 2010), then assembled *de novo* using Trinity 2.12 (Grabherr et al. 2011). Hisat2 v.2.2.1 (Kim et al. 2019) was used for the RNAseq analysis: the hisat2-build function was used to build an index from genome assemblies, then RNA sequencing reads were aligned to the indexed assemblies using default parameter settings. Alignments were visualized with SeqMonk v.1.48.1 (RRID:SCR_001913).

Paired-end genomic data of *Allomyces* strains were quality-filtered by trimmomatic (Bolger, Lohse, and Usadel 2014) with a sliding window of 4bp and threshold of Phred 20, then assembled by Spades 3.15.5 (Prjibelski et al. 2020) and annotated with both Prodigal v. 2.6.3 (Hyatt et al. 2010) and Augustus 3.2.1 (Stanke et al. 2006) trained with *Rhizopus oryzae.*

For the genome plots of *Allomyces arbusculus* Burma 1F *and Allomyces javanicus* California12, we used Prodigal v. 2.6.3 (Hyatt et al. 2010) for gene prediction of soft-masked viral contigs with inputs from Funannotate v. 1.8.16 (Palmer and Stajich 2023) which is a fungal gene annotation

573   tool and GeneMarkS (http://exon.gatech.edu/genemark/index.html) using the viral option for
574   input sequence type. GC skew was calculated in a 5-kb sliding window with a 1-kb step size.
575   Circos v. 0.54 (Krzywinski et al. 2009) was used to construct the circular genome visualization.
576
577   <u>Identification of viral content:</u>
578
We used the program ViralRecall (Aylward and Moniruzzaman 2021) to identify putative viral
genes, viral contigs, and viral regions within contigs. We first ran ViralRecall 1.0 on all
assemblies with both stringent (-s 15 -m 30 -v 10) and relaxed (-s 15 -m 2 -v 2) parameters–the
latter for poorer quality genome assemblies–and identified isolates containing MCP homologs
with e-value <$1e^{-10}$. ViralRecall searches gene predictions generated using Prodigal (Hyatt et al.
2010); we also performed hmmsearches (HMMER3 v3.1b2; S. Eddy 2009) with the ViralRecall
2.0 GVOG hmm on a subset of our Augustus-predicted genomes. We found no differences in
the number of MCPs called, and so continued characterizations for the isolates that these initial
ViralRecall results predicted at least one MCP homolog. For each MCP-containing contig, we
calculated GC content, intron density per the Augustus annotation, and performed BLAST
searches (Altschul et al. 1990) on both gene prediction datasets.
590
To aid in distinguishing viral genomes from within the fungal assemblies, we searched the 16
prodigal annotated genomes previously identified to contain MCPs with HMMER3 (Eddy 2009)
as in Schulz et al. 2020 using hmms of a subset of the nucleocytoplasmic viral orthologous
genes (NCVOGs)— the 20 NCVOGs most likely to have been vertically inherited (Yutin et al.
2009). We made individual gene trees including all hits with e-values less than 1e-10 by aligning
gene sequences with MAFFT version 7 using the E-INS-i algorithm (Katoh and Standley 2013),
trimming the resulting alignments with the -automated1 method in TrimAl (Capella-Gutiérrez
2009), and reconstructing trees with the approximately maximum-likelihood approach
implemented in FastTree (Price, Dehal, and Arkin 2009) with 100 bootstrap replicates, rooting at
the midpoint. We also ran the latest version of ViralRecall (2.0; Aylward and Moniruzzaman
2021) on these 16 assemblies and used these results in combination with genome statistics to
discern putatively viral contigs, which we classified as belonging to a viral genome if they met
the following criteria: ViralRecall score >2, viral region length (per ViralRecall) >70% of the total
contig length, and intron density > 25% different from the host average. One isolate appeared to
contain two distinct NCLDVs; we used MetaBAT2 (Kang et al. 2019) on the entire genome of
Allomyces javanicus California12 under the default settings to separate the two NCLDV
genomes.

We conducted BLASTP searches of the hybrid proteomes against nr to identify homologs (e-
value <1e-10). Because many symbiont genes are likely misidentified as their hosts' in the
databases, we labeled a gene as "viral" or "bacterial" if any of the top hits were to a virus or
bacterium. Some genes have homologs only in the fungi which we have shown also encode
MCP; we labeled these "putative FNCLDV".

To assess prevalence of fungal giant virus-like MCP throughout the fungal kingdom, we used
blastp to search all unmasked fungal genomes in Mycocosm (Grigoriev et al. 2014) using MCP
of *Allomyces arbusculus* Burma1F as query (total 2,543 genomes; conducted on Dec 30, 2023).

616
<u>Giant virus phylogeny, orthology analysis, and KilA-N domain search:</u>
To build the giant virus species tree the nsgtree pipeline was used (https://github.com/NeLLi-
team/nsgtree) on genome representatives of the viral phylum Nucleocytoviricota (Aylward et al.
2021). In brief, 7 giant virus orthologous groups (GVOGs; Aylward et al. 2021) were identified
using hmmsearch, extracted GVOGs were aligned with MAFFT (version 7.31; Katoh and
Standley 2013), trimmed with trimal (-gt 0.1, v1.4; Capella-Gutiérrez, Silla-Martínez, and
Gabaldón 2009) and concatenated. To limit the amount of missing data we removed genomes
that had less than 4 out of 7 GVOGs. A species tree was built from the supermatrix alignment
using IQ-Tree (version 2.03); (Minh et al. 2020; Kalyaanamoorthy et al. 2017) with LG+I+F+G4
and ultrafast bootstrap replications (Hoang et al. 2018) and visualized in iToL (Letunic and Bork
2021).
628
To analyze the global distribution of *Mycodnaviridae* we used the MCP as a query for a blastp
using diamond (--query-cover 30 --subject-cover 30; Buchfink, Xie, and Huson 2015) against all
MCP that we previously extracted (Schulz et al. 2020) from publicly available IMG
metagenomes (Chen et al. 2023). All blastp hits were then extracted and used as query for
another blastp against a database consisting of all MCP extracted from representative giant
virus genomes (Aylward et al. 2021) and the initial *Mycodnaviridae* MCPs. We considered as
bidirectional best blastp hits (BBH) the environmental MCP that had the best hit based on
bitscore to the initial *Mycodnaviridae* MCPs. Geolocation of the BBHs was then used to plot the
distribution of MCPs related to *Mycodnaviridae* on a world map projection.
To search for homologs of the viral KilA-N domain in *Mycodnaviridae* genomes, the source data
from figure 6 of Medina et al. 2016 (219 protein sequences) was downloaded. Sequences were
aligned using mafft (version 7.31; Katoh and Standley 2013), and an hmmprofile made with
HMMER hmmbuild (Eddy 2009), which was then used to hmmsearch the fungal giant virus
genomes.
Orthofinder 2.5.2 (Emms and Kelly 2019) was used to identify orthologous gene clusters from
the prodigal predictions of the most complete viral genomes, determined as the viral genomes
containing the greatest number (at least nine) of the NCVOGs20. Visualization of the shared
orthologous gene clusters was by adaptation of
https://commons.wikimedia.org/wiki/File:Symmetrical_5-set_Venn_diagram.svg. Sequences of
orthologous gene clusters were fed into EggNOG-mapper 5.0 (Huerta-Cepas et al. 2019) with
default parameters but enabling SMART annotations.
Allomyces capsid PCR and growth experiment:
651
Primers were designed from an alignment of the MCP sequences of *A. javanicus* California12
and *A. arbusculus* Burma1F (F 5' TACACCAACTTTGCGATGGA3' R 5'
GTAGCTCGTTGACCCGACAT3'). DNA was provided by the Collection of Zoosporic Eufungi at
the University of Michigan. A standard PCR was performed using GoTaq (Promega) and the
products visualized by agarose gel electrophoresis for absence/presence. Burma1F was used

as a positive control, and Burma3-35 served as a negative control based on absence of the
gene in the sequenced genome.
659
We chose 6 strains that tested positive for MCP (BEA2, Burma1F, Cali8, Cali12, FijiF1,
NorthCaro2), and 6 testing negative (ATCC10983, Burma3-35, CubaS7, DJ02, DJ07, Mex58) to
determine phenotypic effects of the gene's presence. Strains were grown in triplicate on
Emerson's YpSS media (Emerson 1958). The inoculum for each plate was made by growing
strains for 48 hours on fresh YpSS and removing a small circular inoculum by punching discs
with the back of a glass pipette. After 72 hrs of growth at 35 C, linear growth was measured for
each plate with two perpendicular radii. We also estimated biomass accumulation of the same
strains using liquid YpSS. Triplicate 250 ml flasks with 100 ml of media were inoculated with 4
discs cut using the back of glass pipettes. The strains were grown at 35 C for 48 hours with
shaking at 120 rpm. The mycelium was harvested using miracloth filtration, transferred to pre-
weighed filter papers, dried completely, and dry weight was measured. A final experiment
measured thermal maxima of strains by inoculation of YpSS plates and placing them at multiple
temperatures: 38, 40, 42, 44, and 46 C. Duplicate plates were prepared from 9 day old YpSS
plates using discs excised using glass pipettes. Plates were scored for evidence of growth,
where < 2 mm of growth after 1 week was considered negative.
675
<u>Nanopore Sequencing and Genome Assembly:</u>

To obtain High Molecular Weight DNA for long read sequencing, the strain was first grown on
YpSS solid medium for 3 weeks. A block (3mm x 3mm) from the edge of the growing culture
was then excised and immersed in DS buffer to induce zoospore release. The spore suspension
was used to inoculate 100 ml fresh YpSS liquid medium, which was then incubated for 36 hours
at 33 °C in a floor shaker. After incubation, the cells were collected with strainer, washed once
with PBS, frozen at -80 °C, and lyophilized. The frozen-dry cells were broken into powder using
4 mm glass beads and vortex at the maximum speed. Genomic DNA was then extracted using
a modified CTAB protocol, and DNA quality was inspected with CHEF gel electrophoresis as
previously described (https://doi.org/10.1073/pnas.2416656121).

Purified DNA was processed for library preparation and sequencing on an Oxford Nanopore
MinION Mk1C device using R10 flow cells (FLO-MIN114) with MinKNOW UI (v23.07.12)
following the sequencing kit guidelines. The generated Pod5 data was subjected to base calling
using Dorado (v0.6.0) followed by genome assembly using Canu (v2.1.1). The resulting genome
assembly was analyzed for the presence of the capsid sequence using BLASTn analysis and
transposons were identified using Repeatmasker (v4.0.7; http://www.repeatmasker.org),
supplemented with Dfam (v3.3), RepBaseRepeatMaskerEdition-20181026 libraries, and
RepBase EMBL database (v26.04; Bao 2015, Storer 2021). CpG methylation analysis was
performed using the nanopore Modkit tool (v0.2.5-rc1).

<u>Transmission Electron Microscopy:</u>
696
Plate cultures were flooded with WC media for 2 hours. Recovered liquid containing discharged
zoospores was fixed with equal volume 3% glutaraldehyde in 0.01M Sym-collidine, incubated

for two hours at room temperature, and then overnight at 4℃. Zoospores were pelleted and
washed thrice with 0.05% Sym-collidine. The pellet was secondarily fixed with 1% osmium
tetroxide at room temperature for 30 minutes, followed by two washes each with 0.05% sym-
collidine and water. Samples were embedded in 1% noble agar, cut into small blocks (~1mm^3)
washed twice with 10% acetone, and incubated in 2% uranyl acetate at 4℃ overnight. Blocks
were serially dehydrated with increasing concentrations of 10%, 30%, 50%, 75%, 95% acetone,
followed by two washes with 100% acetone. Samples were infiltrated with a 3:1 acetone:Epon-
Araldite resin, and incubated for 1 hr, followed by incubation in 1:1 acetone:resin for 1 hr, and
again in 1:3 acetone:resin. Samples were then set in 100% resin, vacuum centrifuged overnight,
and polymerized at 60℃ for 48 hours. Embedded material was sectioned at 80nm with a
diamond knife, sections collected on 200 mesh formvar/carbon nickel thin bar square mesh,
stained with lead citrate, and visualized on a Jeol 1400plus transmission electron microscope at
the University of Michigan's Department of Molecular, Cell, and Developmental Biology Imaging
Core. *Polyphlyctis willoughbyi* PL AUS 21 was previously prepared (see Letcher and Powell
2018), stored, and imaged on the Jeol 1400plus transmission electron microscope at the
University of Michigan's Department of Molecular, Cell, and Developmental Biology Imaging
Core.
716

References

Altschul, Stephen, Warren Gish, Webb Miller, Eugene Myers, and David Lipman. 1990. "Basic Local Alignment Search Tool." *Journal of Molecular Biology* 215 (3): 403–10. https://doi.org/10.1016/S0022-2836(05)80360-2.

Amses, Kevin R., D. Rabern Simmons, Joyce E. Longcore, Stephen J. Mondo, Kensuke Seto, Gustavo H. Jerônimo, Anne E. Bonds, et al. 2022. "Diploid-Dominant Life Cycles Characterize the Early Evolution of Fungi." *Proceedings of the National Academy of Sciences* 119 (36): e2116841119. https://doi.org/10.1073/pnas.2116841119.

Aylward, Frank O., and Mohammad Moniruzzaman. 2021. "ViralRecall—A Flexible Command-Line Tool for the Detection of Giant Virus Signatures in 'Omic Data." *Viruses* 13 (2): 150. https://doi.org/10.3390/v13020150.

Aylward, Frank O., Mohammad Moniruzzaman, Anh D. Ha, and Eugene V. Koonin. 2021. "A Phylogenomic Framework for Charting the Diversity and Evolution of Giant Viruses." *PLOS Biology* 19 (10): e3001430. https://doi.org/10.1371/journal.pbio.3001430.

Bao W, KK Kojima, O Kohany. 2015. "Repbase Update, a database of repetitive elements in eukaryotic genomes." *Mob DNA* 6(11) doi: 10.1186/s13100-015-0041-9

Bolger, A, M Lohse, and B Usadel. 2014. "Trimmomatic: A Flexible Trimmer for Illumina Sequence Data." *Bioinformatics (Oxford, England)* 30 (15). https://doi.org/10.1093/bioinformatics/btu170.

Buchfink, Benjamin, Chao Xie, and Daniel H. Huson. 2015. "Fast and Sensitive Protein Alignment Using DIAMOND." *Nature Methods* 12 (1): 59–60. https://doi.org/10.1038/nmeth.3176.

Capella-Gutiérrez, Salvador, José M. Silla-Martínez, and Toni Gabaldón. 2009. "trimAl: A Tool for Automated Alignment Trimming in Large-Scale Phylogenetic Analyses." *Bioinformatics* 25 (15): 1972. https://doi.org/10.1093/bioinformatics/btp348.

Chen, I.-Min A., Ken Chu, Krishnaveni Palaniappan, Anna Ratner, Jinghua Huang, Marcel Huntemann, Patrick Hajek, et al. 2023. "The IMG/M Data Management and Analysis System v.7: Content Updates and New Features." *Nucleic Acids Research* 51 (D1): D723–32. https://doi.org/10.1093/nar/gkac976.

Eddy, Sean. 2009. "A New Generation of Homology Search Tools Based on Probabilistic Inference." *Genome Informatics. International Conference on Genome Informatics* 23 (1). https://pubmed.ncbi.nlm.nih.gov/20180275/.

Emerson, Ralph. 1958. "Mycological Organization." *Mycologia* 50 (5): 589–621. https://doi.org/10.2307/3756168.

Emms, David M., and Steven Kelly. 2019. "OrthoFinder: Phylogenetic Orthology Inference for Comparative Genomics." *Genome Biology* 20 (1): 1–14. https://doi.org/10.1186/s13059-019-1832-y.

Grabherr, Manfred G., Brian J. Haas, Moran Yassour, Joshua Z. Levin, Dawn A. Thompson, Ido Amit, Xian Adiconis, et al. 2011. "Full-Length Transcriptome Assembly from RNA-Seq Data without a Reference Genome." *Nature Biotechnology* 29 (7): 644–52. https://doi.org/10.1038/nbt.1883.

Grigoriev, Igor V., Roman Nikitin, Sajeet Haridas, Alan Kuo, Robin Ohm, Robert Otillar, Robert Riley, et al. 2014. "MycoCosm Portal: Gearing up for 1000 Fungal Genomes." *Nucleic Acids Research* 42 (D1): D699–704. https://doi.org/10.1093/nar/gkt1183.

Hoang, Diep Thi, Olga Chernomor, Arndt von Haeseler, Bui Quang Minh, and Le Sy Vinh. 2018. "UFBoot2: Improving the Ultrafast Bootstrap Approximation." *Molecular Biology and Evolution* 35 (2): 518–22. https://doi.org/10.1093/molbev/msx281.

Huerta-Cepas, Jaime, Damian Szklarczyk, Davide Heller, Ana Hernández-Plaza, Sofia K. Forslund, Helen Cook, Daniel R. Mende, et al. 2019. "eggNOG 5.0: A Hierarchical, Functionally and Phylogenetically Annotated Orthology Resource Based on 5090

Organisms and 2502 Viruses." *Nucleic Acids Research* 47 (D1): D309–14. https://doi.org/10.1093/nar/gky1085.

Hyatt, Doug, Gwo-Liang Chen, Philip F. LoCascio, Miriam L. Land, Frank W. Larimer, and Loren J. Hauser. 2010. "Prodigal: Prokaryotic Gene Recognition and Translation Initiation Site Identification." *BMC Bioinformatics* 11 (1): 1–11. https://doi.org/10.1186/1471-2105-11-119.

Kalyaanamoorthy, Subha, Bui Quang Minh, Thomas K. F. Wong, Arndt von Haeseler, and Lars S. Jermiin. 2017. "ModelFinder: Fast Model Selection for Accurate Phylogenetic Estimates." *Nature Methods* 14 (6): 587–89. https://doi.org/10.1038/nmeth.4285.

Kang, Dongwan D., Feng Li, Edward Kirton, Ashleigh Thomas, Rob Egan, Hong An, and Zhong Wang. 2019. "MetaBAT 2: An Adaptive Binning Algorithm for Robust and Efficient Genome Reconstruction from Metagenome Assemblies." *PeerJ* 7. https://doi.org/10.7717/peerj.7359.

Katoh, Kazutaka, and Daron M. Standley. 2013. "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability." *Molecular Biology and Evolution* 30 (4): 772. https://doi.org/10.1093/molbev/mst010.

Kim, Daehwan, Joseph M. Paggi, Chanhee Park, Christopher Bennett, and Steven L. Salzberg. 2019. "Graph-Based Genome Alignment and Genotyping with HISAT2 and HISAT-Genotype." *Nature Biotechnology* 37 (8): 907–15. https://doi.org/10.1038/s41587-019-0201-4.

Krzywinski, Martin, Jacqueline Schein, İnanç Birol, Joseph Connors, Randy Gascoyne, Doug Horsman, Steven J. Jones, and Marco A. Marra. 2009. "Circos: An Information Aesthetic for Comparative Genomics." *Genome Research* 19 (9): 1639. https://doi.org/10.1101/gr.092759.109.

Letunic, Ivica, and Peer Bork. 2021. "Interactive Tree Of Life (iTOL) v5: An Online Tool for Phylogenetic Tree Display and Annotation." *Nucleic Acids Research* 49 (W1): W293–96. https://doi.org/10.1093/nar/gkab301.

Medina, Edgar M., Jonathan J. Turner, Raluca Gordân, Jan M. Skotheim, and Nicolas E. Buchler. 2016. "Punctuated Evolution and Transitional Hybrid Network in an Ancestral Cell Cycle of Fungi." eLife. eLife Sciences Publications Limited. May 10, 2016. https://doi.org/10.7554/eLife.09492.

Minh, Bui Quang, Heiko A. Schmidt, Olga Chernomor, Dominik Schrempf, Michael D. Woodhams, Arndt von Haeseler, and Robert Lanfear. 2020. "IQ-TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era." *Molecular Biology and Evolution* 37 (5): 1530–34. https://doi.org/10.1093/molbev/msaa015.

Nobre SVA, de Andrade GAK, Metz GF, Lucini F, de Albuquerque MP, Victória FC. Antarctica's hidden mycoviral treasures in fungi isolated from mosses: A first genomic approach. J Basic Microbiol. 2024 Jul;64(7):e2300671. doi: 10.1002/jobm.202300671. Epub 2024 May 12. PMID: 38736205.

Palmer, Jonathan, and Jason E. Stajich. 2023. "Funannotate," August. https://doi.org/10.5281/zenodo.1134477.

Katoh, Kazutaka, and Daron M. Standley. 2013. "MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability." *Molecular Biology and Evolution* 30 (4): 772. https://doi.org/10.1093/molbev/mst010.

Price, Mn, Ps Dehal, and Ap Arkin. 2009. "FastTree: Computing Large Minimum Evolution Trees with Profiles Instead of a Distance Matrix." *Molecular Biology and Evolution* 26 (7). https://doi.org/10.1093/molbev/msp077.

Prjibelski, Andrey, Dmitry Antipov, Dmitry Meleshko, Alla Lapidus, and Anton Korobeynikov. 2020. "Using SPAdes De Novo Assembler." *Current Protocols in Bioinformatics* 70 (1): e102. https://doi.org/10.1002/cpbi.102.

817   Schulz, Frederik, Simon Roux, David Paez-Espino, Sean Jungbluth, David A. Walsh, Vincent J.
818   Denef, Katherine D. McMahon, et al. 2020. "Giant Virus Diversity and Host Interactions
819   through Global Metagenomics." *Nature* 578 (7795): 432–36.
820   https://doi.org/10.1038/s41586-020-1957-x.
821   Simmons, D. Rabern, Anne E. Bonds, Buck T. Castillo, Rebecca A. Clemons, Alex D. Glasco,
822   Jillian M. Myers, Natasha Thapa, et al. 2020. "The Collection of Zoosporic Eufungi at the
823   University of Michigan (CZEUM): Introducing a New Repository of Barcoded
824   Chytridiomyceta and Blastocladiomycota Cultures." *IMA Fungus* 11 (1): 1–22.
825   https://doi.org/10.1186/s43008-020-00041-z.
826   Stanke, M, O Keller, I Gundez, A Hayes, Waack, and K Morgenstern. 2006. "AUGUSTUS: Ab
827   Initio Prediction of Alternative Transcripts." *Nucleic Acids Research* 34 (Web Server
828   issue). https://doi.org/10.1093/nar/gkl200.
829   Storer J, R Hubley, J Rosen, TJ Wheeler, AF Smit. 2021. "The Dfam community resource of
830   transposable element families, sequence models, and genome annotations." *Mob DNA*
831   12(1):2. doi: 10.1186/s13100-020-00230-y
832   Yutin, Natalya, Yuri I. Wolf, Didier Raoult, and Eugene V. Koonin. 2009. "Eukaryotic Large
833   Nucleo-Cytoplasmic DNA Viruses: Clusters of Orthologous Genes and Reconstruction of
834   Viral Genome Evolution." *Virology Journal* 6: 223. https://doi.org/10.1186/1743-422X-6-
835   223.