Unveiling the Impact of Vernalization on Seed Oil Content and Fatty Acid Composition in Rapeseed

Through Simulated Shorter Winters

İrem Çağlı[1], Büşra Elif Kıvrak[2], Osman Altunbaş[3], Çağla Sönmez[1,2*]

[1] Department of Biotechnology, Institute of Graduate Studies, Konya Food and Agriculture

University, 42080 Konya, Türkiye

[2] Department of Molecular Biology and Genetics, Faculty of Agriculture and Natural Sciences, Konya

Food and Agriculture University, 42080 Konya, Türkiye

[3] Strategic Products Research and Development Center (SARGEM), Konya Food and Agriculture

University, 42080 Konya, Türkiye

ORCIDs

İrem Çağlı 0000-0002-4029-6622

Büşra Elif Kıvrak 0000-0002-2835-3495

Osman Altunbaş 0000-0003-1840-628X

Çağla Sönmez 0000-0002-5864-6379

[*] Corresponding author: Çağla Sönmez

e.mail: cagla.altun@gmail.com

Abstract

Climate change is leading to warmer winters world-wide with an increasing number of extreme

events every year. Plants are majorly impacted by the escalating effects of global warming. In this

study, we set up an experimental model to simulate warmer and shorter winters under laboratory

conditions. Winter and spring varieties of rapeseed (*Brassica napus* L.) were subjected to diverse

vernalization scenarios including three and four weeks-long vernalization as well as vernalization

interruptions by one week-long devernalization at warm temperatures. The aim of the study was to

assess the effects of the 'vernalization models' on *BnaFLC* (*BnaFLCA02*, *BnaFLCA10* and *BnaFLCC02*)

expression, some yield related traits, a set of genes involved in fatty acid synthesis and seed oil

content and fatty acid composition. A notable difference in vernalization responsiveness was

observed in *BnaFLCA02*, *BnaFLCA10*, and *BnaFLCC02* between the late-flowering winter variety,

Darmor, the early-flowering winter variety Bristol and the spring variety, Helios, after a three-week

vernalization period. Our findings unveil a robust correlation between vernalization and seed oil

content, as well as fatty acid composition in rapeseed. While the expression levels of fatty acid

synthesis-related genes, including *BnaFAD2*, *BnaFAD5*, *BnaFATB*, *BnaMCOA* (*AAE13*), and *BnaWD40*,

exhibited significant changes under cold conditions in leaves, the expression levels of the same

genes in developing seeds did not exhibit a strong correlation with vernalization, flowering time, or

oil and fatty acid contents in seeds. Our results suggest that vernalization plays a role in seed oil

biosynthesis beyond its impact on flowering time.

Keywords: Climate Change; *Brassica napus*; Flowering time; FLC; Oleic acid

1. Introduction

Rapeseed (*Brassica napus* L.) is an economically important oilseed crop valued for its high seed oil

content ranging from 35% to 50%. The adaptability of both winter and spring rapeseed varieties to

diverse environmental conditions significantly contributes to its widespread cultivation (Marjanovic-

Jeromela et al., 2019).

The winter varieties need a prolonged cold period, known as vernalization, to transition from the

vegetative to the generative growth phase in a timely manner. This vernalization process allows

plants to monitor environmental signals, ensuring optimal flowering and subsequent seed

development under favorable conditions for reproductive success. Extensive molecular analyses

conducted on *Arabidopsis thaliana* have revealed the intricate nature of vernalization responses in

plants. Key determinants of vernalization-induced flowering time in Arabidopsis include the

Flowering Locus C (*FLC*) and *FRIGIDA* (*FRI*) genes, which exhibit antagonistic effects on the

vernalization response in plants (Napp-Zinn, 1979; Koornneef et al., 1991; Clarke and Dean, 1994;

Koornneef et al., 1994; Michaels and Amasino, 1999; Shindo et al., 2005).

The specific role of FRI in vernalization is to directly bind to the *FLC* mRNA, keeping *FLC* gene

expression high (Michaels and Amasino, 2001; Geraldo et al., 2009). FLC acts as a transcriptional

repressor and regulates the expression levels of many genes associated with flowering time such as

FLOWERING LOCUS T (*FT*) and *SUPPRESSOR OF CO* (*SOC1*), which are involved in floral transition

(Whittaker and Dean, 2017). Consequently, *FLC* functions to inhibit flowering. During vernalization,

FLC gene expression gradually decreases from the second or third week onwards, and this reduction

is mediated by an epigenetic mechanism that involves the Polycomb Repressive Complex 2 (PRC2)

and nucleation (Bastow et al., 2004; De Lucia et al., 2008; Swiezewski et al., 2009). Additionally, a

chromatin loop formed by the 5' and 3' regions of the *FLC* gene is disrupted during the early stages of

vernalization (Crevillen et al., 2013).

Brassica napus (2n = 4x = 38; genome AACC) is an allotetraploid belonging to the Brassicaceae family

like *A. thaliana* (Chalhaub et al., 2014). Many genes related to vernalization and flowering time are

conserved between Arabidopsis and *B. napus* (Schiessl et al., 2014). Unlike the single *FLC* copy in *A.*

thaliana, *B. napus* genome has nine copies of the *FLC* gene (Zou et al., 2012). Specifically, *BnaFLCA02*,

BnaFLCC02, and *BnaFLCA10* exhibit differential expression in vernalization-dependent versus

vernalization-independent varieties and are downregulated by vernalization (Schiessl et al., 2019).

While *BnaFLCA02* and *BnaFLCC02* expression delay flowering in spring varieties, they do not prevent

it (Tudor et al., 2020). Genome-wide association study (GWAS) of 188 different Brassica accessions

revealed that natural variation in flowering time is primarily determined by *BnaFLCA10* or *BnaFLCA3a*

accounting for 15% and *BnaFLCC02* accounting for 23% of the variance (Raman et al., 2016). QTL-seq

analysis further identified *BnaFLCA02* accounting for flowering time variation in winter rapeseed

cultivars with distinct vernalization requirements (Tudor et al., 2020). Although various studies

highlighted the relative contributions of each *FLC* gene to the vernalization response and the

flowering time, a modeling approach revelead that the total expression levels of nine *FLC* paralogs

collectively contribute to the vernalization response in different rapeseed varieties (Calderwood et

al., 2020).

Climate change poses a significant threat to plant production, leading to considerable crop losses.

(Lesk et al., 2016). Nationwide yield data from the United Kingdom demonstrated a strong

correlation between a 1 °C increase in early winter temperature and a substantial reduction in winter

rapeseed yield (Brown et al., 2019). Recent evidence suggests that vernalization occurs in early

autumn in the field, driven not only by prolonged cold but also by the absence of warmth (Hepworth

et al., 2018; O'Neill et al., 2019).

Growth temperature emerges as a crucial factor influencing rapeseed yield, seed oil content, and

fatty acid composition (Deng and Scarth, 1998; Marjanovic-Jeromela 2019). Triacylglycerols (TAGs)

constitute 94.1-99.1% of the total oil content in *B. napus* seeds, with oleic acid (C18:1), a

monounsaturated 18-carbon fatty acid, being pivotal for oil quality and health (Shahidi, 2005; Dar et

al., 2017). Breeding strategies target the development of high oleic acid varieties (Maher et al.,

2007). While the impact of temperature on seed yield and quality is well-established, studies

addressing the integration of temperature cues at the molecular level in developing rapeseed seeds,

especially the role of epigenetic factors in response to abiotic stress during early development, are

limited. Fatty acid desaturase 2 (FAD2) catalyzes the conversion of oleic acid (C18:1) to linoleic acid

(C18:2) (Miquel and Browse, 1992). Mutants of *fad2* result in seeds with higher oleic acid content in

both Arabidopsis and *B. napus* (Okuley et al., 1994; Hu et al., 2006). Four copies of *BnaFAD2* have

been identified (Lee et al., 2013). Additionally, a GWAS analysis linked FATTY ACID DESATURASE 5

(FAD5) and Fatty acyl-ACP thioesterase B (FATB) to variations in fatty acid composition in winter

rapeseed (Gacek et al., 2017).

Few studies have explored the relationship between vernalization and membrane fatty acid

composition in seedlings (Thomson & Zalik, 1973; Dubert et al., 1992; Skoczowski & Filek, 1994).

Winter wheat embryos exposed to different vernalization-devernalization cycles exhibited altered

fatty acid composition in germinated seedlings, correlating with changes in the timing of floral

induction (Dubert et al., 1992). However, the impact of a vernalization cycle temporarily interrupted

by warm temperatures (>15 °C) during early growth on the fatty acid composition of seeds remains

unknown.

In this study, we set up a series of experimental models to simulate shortened winters that include

cold temperatures (4 °C), interrupted by one-week-long period of warmer temperature (22 °C) under

laboratory conditions. These simulations, referred to as vernalization models, were implemented to

investigate their impact on seed oil content and fatty acid composition in both winter and spring

varieties of rapeseed. Additionally, we assessed the expression levels of three *BnaFLC* genes

(*BnaFLCA02*, *BnaFLCA10*, and *BnaFLCC02*) in seedlings under these vernalization regimes. Moreover,

we examined the expression patterns of various genes associated with fatty acid synthesis in both

seedlings and developing seeds. The overarching goal of this study was to assess the correlations

between vernalization, flowering time and several yield-related traits, and the oil content and fatty

acid composition of seeds of rapeseed.

2. Materials and Methods

2.1 Plant material, seed sowing and germination:

The seeds of *Brassica napus* winter varieties, Darmor and Bristol, and spring variety, Helios were

kindly provided by the Germplasm Resource Information Network (GRIN, Czech Republic). Darmor

has been used as the reference variety for the whole genome sequencing of *B. napus* (Chalhoub et

al., 2014). Both Bristol and Helios were grown in Central Anatolia, and their suitability for cultivation

in Türkiye has been confirmed (Başalma, 2004; Öztürk et al., 2008). Darmor and Bristol have been

designated as late and early flowering, respectively (https://www.grin-global.org).

The experiments were conducted for three consecutive years between 2021-2023. Two seeds were

sown per pot (dimensions 11x11x18 cm) filled with a soil mixture of peat and perlite (1:1) moistened

with the Hoagland solution (Hoagland and Arnon, 1950). The pots were placed in a growth room

with a 12-hour day/night cycle, at 22 °C, 50-55% humidity, and a light intensity of 150 µmol m^{-2} s^{-1}.

Plants were irrigated with fifty milliliters of Hoagland solution or distilled water at two-three day

intervals to maintain soil moisture.

2.2 The vernalization treatment:

The germinated seedlings that reached the four-five leaf stage (BBCH scale 14-15) were moved to a

vernalization chamber (Nüve TK 252) at 4 °C; short days. The nonvernalized (NV) samples were left in

the growth room at 22 °C; short days. The different vernalization models are illustrated in Fig. 1. At

least twenty plants from each variety were subjected to vernalization for three weeks (3w); four

weeks (4w); six weeks (6w); two weeks vernalization, one week devernalization at 22 °C followed by

four weeks vernalization (2+4w); 8 weeks (8w) and 2 weeks vernalization, one week devernalization

at 22 °C followed by 6 weeks vernalization (2+6w). For devernalization, plants were transferred to the

growth room at 22 °C; short days. Previous studies have shown that the expression levels of *BnaFLC*

genes in winter rapeseed varieties cannot be detected after 8 weeks of vernalization (Schiessl et al.,

2019).

2.3 Plant growth conditions post-vernalization:

Plants were transplanted to soil (slightly alkaline, clay loam textured and fertilized with diammonium

phosphate before planting) outdoors on April 26, 2021, and on April 25, 2022 when the night

temperature reached 10 °C and above and to a semi-controlled greenhouse with indoor heating but

no cooling system on January 11, 2023. 150 W LED lamps (Philips Xitanium) provided supplementary

lighting when required in the greenhouse. On October 11, 2022, Helios and Bristol seeds were

directly sown in soil as winter planting. The highest and lowest daily temperatures were recorded

and are given in Fig. 2A-D. The soil was irrigated every two to three days. Flowering time was

recorded as the number of days after the day of transplantation when the first flower appeared or

days after sowing in 2023 field plants. Yield associated physiological parameters including plant

height (cm), the number of lateral branches (n), the number of pods on the main stem (n), the

number of seeds per pod (n), and thousand seed weight (g) were measured.

2.4 RNA extraction and analysis:

Leaf tissue was collected from at least three plants per vernalization model after the vernalization

was completed and immediately frozen in liquid nitrogen. Siliques on the main stem that reached

BBCH growth scale 80 (i.e. beginning of ripening, seeds green, filling pod cavity) were collected,

developing seeds were separated and immediately frozen in liquid nitrogen. Tissue was homogenized

and RNA extraction was performed as described in Queseda et al. (2003). Total RNA was DNase

treated with the Turbo DNA-free kit (Thermo Fisher Scientific). For quantitative real-time RT-PCR,

one µg total RNA was reverse-transcribed with the iScript cDNA synthesis kit (Bio-Rad) according to

the manufacturer's instructions. Six µL cDNA (diluted 1:5 with distilled water) was used per qRT-PCR

reaction in a 15-µL reaction volume using SYBR Green (BIO-RAD, iTaq Universal SYBR Green

Supermix) and a BIO-RAD CFX96 real time PCR instrument. Expression was normalized to Ubiquitin

using the comparative cT method.

Primers for *B. napus* genes including *BnaFAD2.A5* and *BnaFAD2.C5*, *BnaFAD5* (*BnaA05g23670D*),

BnaFATB (*BnaA05g23790D*), *BnaMCOA* (*AAE13*; BnaA05g23520D), *BnaWD40* (*BnaA05g24090*),

BnaFLCA02, BnaFLCA10, BnaFLCC02, and *Ubiquitin* as internal control were either designed using the

Primer3Plus program (http://www.bioinformatics.nl/cgibin/primer3plus/primer3plus.cgi) or

according to the relevant reference indicated in Supplementary Table S1. The gene sequences are

available at http://www.genoscope.cns.fr/brassicanapus/.

2.5 Total lipid extraction and gravimetric lipid determination:

Seeds were harvested from at least three plants per vernalization model and dried. Total lipid

extraction was done as described by Li et al. (2006). Briefly, two mL of isopropanol was added to 100

mg seeds in a glass tube with a Teflon© cap. The tubes were incubated at 85 °C in a water bath for

10 minutes. The seeds were then homogenized using a glass-glass homogenizer. Three mL of hexane

was added to the sample and vortexed. Following a five-minute incubation at room temperature, 2.5

mL of 15% w/v sodium sulfate was added, leading to phase separation. The upper phase was

transferred to a new glass tube and evaporated under nitrogen. The total oil content (%) was

calculated gravimetrically.

2.6 Determination of fatty acid composition of seeds:

The fatty acid composition of *B. napus* seeds was determined by gas chromatography (GC). The lipid

extracts were subjected to transesterification with 0.5 mL of 2 N potassium hydroxide as catalyst. 10

mL of n-hexane was added to the samples and mixed. The mixture was centrifuged at 5000 rpm for

15 minutes, and the upper phase was collected for GC analysis using SHIMADZU (GCMS-QP2020)

instrument with a column (RESTEK; Rxi-5 Sil MS 30 Meter 0.25 mm ID 0.25 µm df). The column was

operated at 250 °C and 34.17 psi (split flow: 207.1 mL/min, total flow: 215.9 mL/min). Fatty acid

methyl esters were quantified as percentages by comparing peak areas with standards.

2.7 Statistical analysis:

All experiments were carried out with at least three biological replicates and results are expressed as

mean ± standard error (SE). The mean values of data were analyzed by one-way analysis of variance

(ANOVA) and unpaired t-tests (Mann-Whitney U test) was employed for group comparisons.

Statistical significance was determined at the 5% level ($p < 0.05$). Correlations among different

parameters were calculated using Pearson's correlation coefficient (r). The statistical analyses were

performed using GraphPad Prism 9 software.

3. Results

3.1 Expression profiles of *BnaFLC* genes and flowering time under different vernalization models:

We addressed the effect of the vernalization models (NV, 3w, 4w, 6w, 2+4w, 8w and 2+6w) on the

expression levels of three *BnaFLC* genes; *BnaFLCA02*, *BnaFLCA10* and *BnaFLCC02* as well as the

flowering time in rapeseed winter varieties Darmor and Bristol and spring variety Helios. Initially, the

expression levels of *BnaFLC* genes in the nonvernalized (NV) leaf samples of the three varieties were

within a comparable range (Fig. 3). Notably, after three weeks of vernalization (3w), a significant

disparity in the downregulation rates of *FLC* genes was observed in Darmor, Bristol, and Helios.

Darmor, characterized as a late-flowering winter variety, exhibited only a marginal reduction in

BnaFLCA02 and *BnaFLCC02* expression levels and a three-fold reduction in *BnaFLCA10* expression

(Fig. 3A-C). In contrast, Bristol, an early-flowering winter variety, displayed a higher reduction rate in

all three *BnaFLC* genes compared to Darmor, yet not as pronounced as the sudden decrease

observed in Helios (Fig. 3). Specifically, *BnaFLCA02* and *BnaFLCC02* levels in Helios were reduced

approximately five-fold, whereas *BnaFLCA10* showed an eleven-fold reduction after three weeks of

vernalization (3w) (Fig. 2G-I). All three *FLC* genes were undetected in Helios after six weeks of

vernalization (6w). Despite being significantly lower than the nonvernalized (NV) state, Darmor and

Bristol still displayed expression of *BnaFLCA02*, *BnaFLCA10*, and *BnaFLCC02* after eight weeks of

vernalization (8w). *BnaFLCA10* emerged as the most responsive to vernalization in all three varieties.

Notably, Darmor's *BnaFLCC02* expression level was slightly higher in plants exposed to eight-week

vernalization with a one-week interruption (2+6w) compared to those exposed to eight-week

vernalization without any interruption (8w) (Fig. 3C). Conversely, in Bristol, 2+6w *BnaFLCC02*

expression was higher than 8w expression (Fig. 3F).

In the initial two years (2021 and 2022) of the study, seedlings were transplanted into soil and grown

under field conditions post-vernalization. Bristol and Helios were previously grown in Central

Anatolia, while there is no information available about the cultivation of Darmor in Türkiye (Başalma,

2004; Öztürk et al., 2008). Darmor exhibited severe growth defects and extreme sensitivity to pests

in 2021, with only a few plants from the 6w, 8w, and 2+6w vernalization models flowering. Plants

from other vernalization models (NV, 3w, 4w, 2+4w) did not flower 160 days after transplantation

(Fig. 3A-C). Consequently, we determined that Darmor is unsuitable for planting in the soil and

weather conditions of Konya, Türkiye, and discontinued its cultivation in the following years of the

study.

In 2021 and 2023, plants from all vernalization models of Bristol flowered. However, in 2022, Bristol

NV plants did not flower, while the rest flowered significantly earlier than in the 2021 field season.

Although flowering was observed, plants from 3w and 4w vernalization models failed to complete

their growth cycle in 2022. Across all three years, there was no statistically significant difference in

flowering time between 6w and 2+4w or 8w and 2+6w vernalization models in Bristol (Fig. 3D-F).

Helios NV plants consistently flowered earlier than Bristol NV plants over the three years (Fig. 3G-I).

Throughout the 2021 and 2022 field seasons, flowering times of plants in all vernalization models of

Helios were similar. In 2023, under greenhouse conditions, while flowering times were longer

compared to previous years, a faster rate of reduction in flowering time between NV and 3w

vernalization models was observed than in 2021 and 2022. A slight increase in mean flowering times

of plants in the 2+4w vernalization model compared to 6w was noted across all three years (~3-4

days in 2021 and 2022; ~8 days in 2023). There was no meaningful difference between 2+6w and 8w

in 2021 and 2023, with a slight increase in 2+6w compared to 8w (~5 days) in 2022. These results

were not statistically significant, possibly due to the variation among flowering times of plants and

the small sample size (3≤n<10).

During the 2022-2023 growing season, alongside transplanting the seedlings in the greenhouse,

Bristol and Helios seeds were directly sown in soil in October 2022. The plants from the Bristol and

Helios varieties flowered in early April 2023; 179±2 and 182±4 days after sowing, respectively.

We analyzed the relationship between vernalization, represented by vernalization models NV, 3w,

4w, 6w and 8w expressed as days of vernalization, and the expression levels of the *BnaFLC* genes, as

well as the annual flowering time. The Pearson correlation coefficients (r) between each pair of

variables are presented in Fig. 6. As anticipated, a robust negative correlation between vernalization

and flowering times was observed consistently across all three years for both Bristol and Helios

varieties ($p < 0.05$). The expression levels of *BnaFLCA02*, *BnaFLCA10*, and *BnaFLCC02* of both Bristol

and Helios varieties exhibited a negative correlation with vernalization and a positive correlation with

flowering time consistent with the role of FLC in vernalization response.

3.2 Yield related physiological traits of plants under different vernalization models:

We performed a comprehensive analysis of various yield-related traits in rapeseed plants, including

parameters such as plant height, number of lateral branches, number of pods on the main stem,

number of seeds per pod, and thousand seed weight. This investigation spanned three consecutive

years and involved the evaluation of Bristol and Helios varieties under different vernalization models.

In 2023, the plant height of both Bristol and Helios varieties cultivated in the greenhouse reached its

maximum, surpassing the heights of field-grown plants in both 2021 and 2022. (Fig. S1A-B).

Intriguingly, the 2+6w vernalization model resulted in the shortest plants during the same year ($p <$

0.05). Conversely, during the 2021 planting season, the tallest plants were observed in the Bristol

variety under the 2+6w vernalization model ($p < 0.05$). Notably, a substantial variability in

physiological traits was discerned across the three years, underscoring the influence of multifactorial

events, possibly attributable to specific growth conditions prevailing in each year. An exception to

this trend was noted in the number of lateral branches in the Helios variety, which exhibited a subtle

yet consistent decline with increasing vernalization time, irrespective of the planting year (Fig. S1D).

In the 2023 planting season, the vernalization models 2+4w and 2+6w in both Bristol and Helios

varieties yielded plants with a reduced number of pods on the main stem compared to those

subjected to 6w and 8w vernalization, respectively, in greenhouse conditions. This observation

suggests an interplay between vernalization conditions and pod development.

We calculated the Pearson correlation coefficients (r) to gain insight into the interplay among

vernalization, flowering time, and yield-related physiological traits (Fig. 6). In 2021, there was a

positive correlation between the number of pods on the main stem and a negative correlation

between the seed number per pod and the duration of vernalization in Bristol ($p < 0.05$) (Fig. 6A). In

2022, Bristol NV, 3w, and 4w plants initiated flowering but failed to progress into fully developed,

healthy plants. Consequently, our analysis for this specific year excluded physiological traits other

than plant height (Fig. 6B). In 2023, although a positive correlation was noted between the number

of pods on the main stem and the vernalization model in Bristol, the result was not statistically

significant (Fig. 6C). Turning to Helios, a robust positive correlation between thousand seed weight

and the days of vernalization (i.e., vernalization model) was evident in 2022 ($p < 0.001$), but this

association was not observed in other years (Fig. 6E). On the contrary, the number of lateral

branches displayed a robust positive correlation with flowering time in both 2021 and 2022, and a

subtle positive correlation in 2023 (Fig. 6D-F). These findings suggest that multiple variables, beyond

vernalization and flowering time, contribute to the variability in yield-related physiological traits.

3.3 Effects of vernalization models on seed oil and oleic acid contents in rapeseed:

We assessed the impact of vernalization models—specifically, variations in duration and the

presence or absence of an interruption during vernalization—on the seed oil content and fatty acid

composition of rapeseed varieties Bristol and Helios (Fig. 4).

In Bristol, during both 2021 and 2022, the 2+6w vernalization model exhibited the highest oil content

(% w/w) at 21% and 41%, respectively. Notably, during the same two years, a significant difference in

seed oil content was observed between the 8w and 2+6w vernalization models ($p < 0.05$),

underscoring the influence of vernalization interruption by warm temperatures on seed oil content.

This pattern was similarly reflected in the percentage of oleic acid content (Fig. 4B). Across all three

years, seeds from plants vernalized for 2+6w consistently demonstrated higher oleic acid content

compared to those vernalized for 8w.

In 2023, the 4w vernalization model exhibited the highest oil content at 40% (Fig. 4A). There was no

significant difference in seed oil content (% w/w) between the 8w and 2+6w vernalization models in

2023. The relationship between seed oil content (%) and flowering time was notably strong in 2021 (r

= 0.999; $p < 0.05$) and 2023 (r = 0.839; $p > 0.05$) (see Fig. 6A and C). Furthermore, a robust positive

correlation between flowering time and percentage of oleic acid content was observed in 2023 (r =

0.894; $p < 0.05$).

In Helios, seeds from non-vernalized (NV) plants exhibited the lowest oil content across all three

years. In 2022, NV plants were adversely affected by the weather conditions and pests, leading to the

production of small, unhealthy-looking seeds and a consequently, a low seed oil yield (8%).

Nevertheless, even in the other two years, seeds from NV plants displayed lower oil content than

those of the other vernalization models (Fig. 4C).

6w plants had the highest seed oil contents across all three years, surpassing those of 2+4w plants.

Similarly, the seed oil contents of 8w plants in 2021 and 2022 exceeded those of 2+6w plants ($p <$

0.05), though the difference was not statistically significant in 2023. Oleic acid contents (%) showed

significant differences between the 6w and 2+4w, as well as the 8w and 2+6w vernalization models in

2022. A similar trend was evident in 2021 and 2023, albeit with more subtle differences.

We calculated the relative amounts of saturated (SFA), monounsaturated (MUFA), and

polyunsaturated (PUFA) fatty acids for each vernalization model throughout the 2021-2023 period

(refer to the supplementary file). Remarkably, there was a significant increase in MUFA during

interrupted vernalization models (2+4w and 2+6w) compared to uninterrupted vernalization (6w and

8w, respectively) across all three years in the Bristol variety. A similar effect was observed in Helios

during the 2021 and 2022 field seasons, albeit with a more subtle impact.

The relationship between vernalization duration, flowering time, and seed oil content in Helios

remained robust across all field seasons (Fig. 6D-F). The percentage of seed oil content exhibited a

positive correlation with increasing vernalization duration in 2021 (r = 0.825; $p > 0.05$), 2022 (r =

0.944; $p < 0.05$), and 2023 ($r = 0.884$; $p < 0.05$). Oleic acid content (%) also displayed a positive

correlation with vernalization duration in 2021 ($r = 0.790$; $p > 0.05$) and 2023 ($r = 0.921$; $p < 0.05$).

This interplay between oil content and vernalization was further highlighted by the strong negative

correlation between gene expression levels of *BnaFLCA02*, *BnaFLCA10*, and *BnaFLCC02* and seed oil

content in Helios across all three years (Fig. 6D-F). In summary, our findings underscore the profound

impact of vernalization on seed oil content and fatty acid composition in rapeseed.

3.4 The expression profiles of the fatty acid synthesis-related genes in leaf tissues across the

vernalization models:

We investigated the expression profiles of several genes related to fatty acid synthesis, including

BnaFAD2, *BnaFAD5*, *BnaFATB*, *BnaMCOA* (*AAE13*), and *BnaWD40*, in the leaf tissue of Darmor,

Bristol, and Helios varieties under various vernalization models. In NV samples, the expression levels

of all five genes were higher in Helios compared to Darmor and Bristol (Fig. S3). Notably, there was

no uniform response to vernalization in gene expression levels among different varieties.

For instance, in Darmor, *BnaFAD2*, responsible for catalyzing the conversion of oleic acid (C18:1) to

linoleic acid (C18:2), exhibited an increase in expression upon cold exposure until six weeks (6w and

2+4w) of vernalization, after which a sudden drop occurred at 8w. Conversely, in Helios, there was an

abrupt decline in *BnaFAD2* levels after three weeks (3w) of cold exposure, which persisted after 6w

and 8w. Interestingly, interrupting the cold exposure after 2w with a week of warm temperature,

followed by an additional four or six weeks of cold, caused a further decrease in expression levels of

BnaFAD2 (Fig. S3A).

The relative increase in expression levels of *BnaFAD5*, *BnaFATB*, *BnaMCOA*, and *BnaWD40* between

the 6w vs 2+4w and 8w vs 2+6w vernalization models was evident in Helios (Fig. S3B-E). In Darmor,

BnaFAD5 expression was suppressed throughout all vernalization models (Fig. S3B). Similarly, in

Bristol, the expression of *BnaFATB*, which encodes Fatty acyl-ACP thioesterase B, remained

consistently low under cold conditions compared to NV (Fig. S3C). Notably, *BnaMCOA/AAE13*,

encoding a malonyl-CoA synthetase, showed minimal expression in Darmor and Bristol NV samples,

with its expression increasing upon cold exposure in Bristol (Fig. S3D).

The candidate gene *BnaWD40* (BnaA05g24090), encoding a protein from the Transducin/WD40

repeat-like superfamily, is implicated in the regulation of oleic acid (Gacek et al., 2017). *BnaWD40*

was downregulated upon exposure to cold in Darmor at three weeks (3w), Bristol at four weeks (4w),

and Helios at six weeks (6w), as illustrated in Fig. S3E.

Overall, *BnaFAD2*, *BnaFAD5*, *BnaFATB*, *BnaMCOA*, and *BnaWD40* were differentially regulated upon

cold exposure in rapeseed varieties.

3.5 The expression profiles of the fatty acid synthesis-related genes in developing seeds across the

vernalization models:

We studied the expression levels of *BnaFAD2*, *BnaFAD5*, *BnaFATB*, *BnaMCOA*, and *BnaWD40* in

developing seeds annually between 2021 and 2023, assessing their potential correlation with seed

oil content and fatty acid composition across different vernalization models in Bristol and Helios. As

mentioned earlier, Darmor plants exhibited suboptimal growth under field conditions in this study,

preventing the examination of gene expression profiles in developing seeds of this variety.

The relative expression patterns of *BnaFAD2*, *BnaFAD5*, *BnaFATB*, *BnaMCOA*, and *BnaWD40* in

developing seeds across all three years displayed high variability in both Bristol and Helios (Fig. 5).

This stochasticity in gene expression was further evident in the statistical correlation analysis. For

instance, in 2021, a negative correlation between *BnaFATB* expression and vernalization duration was

observed in Bristol (r = -0.997; $p < 0.05$) (Fig. 5A). However, in 2023, a positive correlation was

identified (r = 0.776; $p > 0.05$) (Fig. 5C). Similarly, in 2021, Bristol *BnaFAD2* expression was positively

correlated with *BnaFLCA02* (r = 1.000; $p < 0.01$), *BnaFLCA10* (r = 0.997; $p < 0.05$), and *BnaFLCC02* (r =

0.940; $p > 0.05$), whereas this correlation was not observed in 2023 (see Fig. 5A and C).

The negative correlation between Bristol *BnaFAD2* expression and oleic acid content (%) in 2021 (r = -

0.801; $p > 0.05$) and 2023 (r = -0.847; $p > 0.05$), was not observed in Helios (see Fig. 5).

In summary, our findings indicate the absence of consistent correlations between the expression

patterns of fatty acid-related genes that were included in this study in developing rapeseed seeds

and vernalization, flowering time, oil content, oleic acid content.

Acknowledgements

This study was funded by TUBITAK grant number 120Z278 to Ç. Sönmez. B. E. Kıvrak was funded by

TUBITAK 2209-A undergraduate fellowship program.

References

Basalma D. (2004) Comparison of Winter Rapeseed (*Brassica napus* ssp. oleifera L.) Varieties in terms

of yield and yield components under Ankara conditions. Journal of Agricultural Sciences, *10*(2), 211-

217.

Bastow, R., Mylne, J. S., Lister, C., Lippman, Z., Martienssen, R. A., & Dean, C. (2004). Vernalization

requires epigenetic silencing of FLC by histone methylation. *Nature*, *427*(6970), 164-167.

Brown, J. K., Beeby, R., & Penfield, S. (2019). Yield instability of winter oilseed rape modulated by

early winter temperature. *Scientific reports*, *9*(1), 6953.

Calderwood, A., Lloyd, A., Hepworth, J., Tudor, E. H., Jones, D. M., Woodhouse, S., ... & Morris, R. J.

(2021). Total FLC transcript dynamics from divergent paralogue expression explains flowering

diversity in Brassica napus. *New Phytologist*, *229*(6), 3534-3548.

Chalhoub, B., Denoeud, F., Liu, S., Parkin, I. A., Tang, H., Wang, X., ... & Corréa, M. (2014). Early

allopolyploid evolution in the post-Neolithic *Brassica napus* oilseed genome. *Science*, *345*(6199),

950-953

Clarke, J. H., & Dean, C. (1994). Mapping FRI, a locus controlling flowering time and vernalization

response in Arabidopsis thaliana. *Molecular and General Genetics MGG*, *242*(1), 81-89

Crevillen, P., Sonmez, C., Wu, Z., & Dean, C. (2013). A gene loop containing the floral repressor FLC is

disrupted in the early phase of vernalization. *The EMBO journal*, *32*(1), 140-148.

Dar, A. A., Choudhury, A. R., Kancharla, P. K., & Arumugam, N. (2017). The FAD2 gene in plants:

occurrence, regulation, and role. *Frontiers in plant science*, *8*, 1789.

De Lucia, F., Crevillen, P., Jones, A. M., Greb, T., & Dean, C. (2008). A PHD-polycomb repressive

complex 2 triggers the epigenetic silencing of FLC during vernalization. *Proceedings of the National*

Academy of Sciences, *105*(44), 16831-16836.

Deng, X., & Scarth, R. (1998). Temperature effects on fatty acid composition during development of

low-linolenic oilseed rape (Brassica napus L.). *Journal of the American Oil Chemists' Society*, *75*(7),

759-766.

Dubert, F., Filek, M., Marcinska, I., & Skoczowski, A. (1992). Influence of warm intervals on the effects

of vernalization and the composition of phospholipid fatty acids in seedlings of winter wheat. *Journal*

of Agronomy and Crop Science, *168*(2), 133-141.

Gacek, K., Bayer, P. E., Bartkowiak-Broda, I., Szala, L., Bocianowski, J., Edwards, D., & Batley, J. (2017).

Genome-wide association study of genetic control of seed fatty acid biosynthesis in Brassica

napus. *Frontiers in Plant Science*, *7*, 2062.

Geraldo, N., Bäurle, I., Kidou, S. I., Hu, X., & Dean, C. (2009). FRIGIDA delays flowering in Arabidopsis

via a cotranscriptional mechanism involving direct interaction with the nuclear cap-binding

complex. *Plant Physiology*, *150*(3), 1611-1618.

Hepworth, J., Antoniou-Kourounioti, R. L., Bloomer, R. H., Selga, C., Berggren, K., Cox, D., ... &

Howard, M. (2018). Absence of warmth permits epigenetic memory of winter in Arabidopsis. *Nature*

communications, *9*(1), 1-8.

Hoagland, D.R., and D.I. Arnon. 1950. The water culture methods for growing plants without soil.

Calif. Agric. Exp. Stn. Circular Number 347 (Revised January 1950), College of Agriculture University of

California, Berkeley, CA.

Hu, X., Sullivan-Gilbert, M., Gupta, M., & Thompson, S. A. (2006). Mapping of the loci controlling

oleic and linolenic acid contents and development of fad2 and fad3 allele-specific markers in canola

(Brassica napus L.). *Theoretical and Applied Genetics*, *113*, 497-507.

Koornneef, M., Hanhart, C. J., & Van der Veen, J. H. (1991). A genetic and physiological analysis of late

flowering mutants in Arabidopsis thaliana. *Molecular and General Genetics MGG*, *229*(1), 57-66.

Koornneef, M., Blankestijn-de Vries, H., Hanhart, C., Soppe, W., & Peeters, T. (1994). The phenotype

of some late-flowering mutants is enhanced by a locus on chromosome 5 that is not effective in the

Landsberg erecta wild-type. *The Plant Journal*, *6*(6), 911-919

Lee, K. R., Sohn, S. I., Jung, J. H., Kim, S. H., Roh, K. H., Kim, J. B., ... & Kim, H. U. (2013). Functional

analysis and tissue-differential expression of four FAD2 genes in amphidiploid Brassica napus derived

from Brassica rapa and Brassica oleracea. *Gene*, *531*(2), 253-262.

Lesk, C., Rowhani, P., & Ramankutty, N. (2016). Influence of extreme weather disasters on global crop

production. *Nature*, *529*(7584), 84-87.

Li, Y., Beisson, F., Pollard, M., & Ohlrogge, J. (2006). Oil content of Arabidopsis seeds: the influence of

seed anatomy, light and plant-to-plant variation. *Phytochemistry*, *67*(9), 904-915.

Maher, L., Burton, W., Salisbury, P., Debonte, L., & Deng, X. (2007, March). High Oleic, low linolenic

(HOLL) specialty canola development in Australia. In *Proceedings of the 12th International Rapeseed*

Congress (Vol. 5, pp. 22-24).

Marjanović-Jeromela, A., Terzić, S., Jankulovska, M., Zorić, M., Kondić-Špika, A., Jocković, M., ... &

Nagl, N. (2019). Dissection of Year Related Climatic Variables and Their Effect on Winter Rapeseed

(*Brassica napus* L.) Development and Yield. *Agronomy*, *9*(9), 517.

Michaels, S. D., & Amasino, R. M. (2001). Loss of FLOWERING LOCUS C activity eliminates the late-

flowering phenotype of FRIGIDA and autonomous pathway mutations but not responsiveness to

vernalization. *The Plant Cell*, *13*(4), 935-941.

Miquel M, Browse J (1992) Arabidopsis mutants deficient in polyunsaturated fatty-acid synthesis—

biochemical and genetic-characterization. *Journal of Biological Chemistry* 267:1502–1509

Napp-Zinn, K. (1979). On the genetical basis of vernalization requirement in Arabidopsis thaliana (L.)

Heynh. *La Physiologie de la Floraison*, 217-220.

Okuley, J., Lightner, J., Feldmann, K., Yadav, N., Lark, E., & Browse, J. (1994). Arabidopsis FAD2 gene

encodes the enzyme that is essential for polyunsaturated lipid synthesis. *The Plant Cell*, *6*(1), 147-

158.

O'Neill, C. M., Lu, X., Calderwood, A., Tudor, E. H., Robinson, P., Wells, R., ... & Penfield, S. (2019).

Vernalization and Floral Transition in Autumn Drive Winter Annual Life History in Oilseed

Rape. *Current Biology*.

Öztürk, Ö., Rahim, A. D. A., & Akinerdem, F. (2008). Determination of appropriate planting time for

spring rapeseed varieties under Konya conditions. Selçuk Journal of Agricultural Sciences, *22*(46), 6-

17.

Raman, H., Raman, R., Coombes, N., Song, J., Prangnell, R., Bandaranayake, C., ... & Balasubramanian,

S. (2016). Genome-wide association analyses reveal complex genetic architecture underlying natural

variation for flowering time in canola. *Plant, Cell & Environment*, *39*(6), 1228-1239.

Schiessl, S., Samans, B., Hüttel, B., Reinhard, R., & Snowdon, R. J. (2014). Capturing sequence

variation among flowering-time regulatory gene homologs in the allopolyploid crop species Brassica

napus. *Frontiers in plant science*, *5*, 404.

Schiessl, S. V., Quezada-Martinez, D., Tebartz, E., Snowdon, R. J., & Qian, L. (2019). The vernalisation

regulator FLOWERING LOCUS C is differentially expressed in biennial and annual Brassica

napus. *Scientific reports*, *9*(1), 1-15.

Shahidi F., 2005. Bailey's Industrial Oil and Fat Products, Volume 6, Wiley-Interscience, New Jersey,

USA.

Shindo, C., Aranzana, M. J., Lister, C., Baxter, C., Nicholls, C., Nordborg, M., & Dean, C. (2005). Role of

FRIGIDA and FLOWERING LOCUS C in determining variation in flowering time of Arabidopsis. *Plant*

physiology, *138*(2), 1163-1173.

Skoczowski, A., & Filek, M. (1994). Changes in fatty acids composition of subcellular fractions from

hypocotyls of winter rape growing at 2 C or 20 C. *Plant Science*, *98*(2), 127-133.

Swiezewski, S., Liu, F., Magusin, A., & Dean, C. (2009). Cold-induced silencing by long antisense

transcripts of an Arabidopsis Polycomb target. *Nature*, *462*(7274), 799-802.

Thomson, L. W., & Zalik, S. (1973). Lipids in rye seedlings in relation to vernalization. *Plant*

physiology, *52*(3), 268-273.

Tudor, E. H., Jones, D. M., He, Z., Bancroft, I., Trick, M., Wells, R., ... & Dean, C. (2020). QTL-seq

identifies BnaFT. A02 and BnaFLC. A02 as candidates for variation in vernalization requirement and

response in winter oilseed rape (Brassica napus). *Plant biotechnology journal*, *18*(12), 2466-2481.

Quesada, V., Macknight, R., Dean, C., & Simpson, G. G. (2003). Autoregulation of FCA pre-mRNA

processing controls Arabidopsis flowering time. *The EMBO Journal*, *22*(12), 3142-3152.

Whittaker, C., & Dean, C. (2017). The FLC locus: a platform for discoveries in epigenetics and

adaptation. *Annual Review of Cell and Developmental Biology*, *33*, 555-575.

Zou, X., Suppanz, I., Raman, H., Hou, J., Wang, J., Long, Y., ... & Meng, J. (2012). Comparative analysis

of FLC homologues in Brassicaceae provides insight into their role in the evolution of oilseed rape.

Figure legends:

Fig. 1 Vernalization model chart. Each box represents one week. Vernalization and devernalization

took place at 4 °C and 22 °C, respectively. NV; nonvernalized, 3w; three weeks-vernalization, 4w; four

weeks-vernalization, 6w; six weeks-vernalization, 2+4w; two weeks-vernalization, one week-

devernalization, four weeks-vernalization, 8w; eight weeks-vernalization, 2+6w; two weeks-

vernalization, one week-devernalization, six weeks-vernalization.

Fig. 2 Daily maximum (Tmax) and minimum (Tmin) temperature data for Meram, Konya, Türkiye for

the years 2021-2023. Tmax and Tmin starting on (Day 0) **A.** April 26, 2021; **B.** April 25, 2022; **C.**

January 11, 2023 in the greenhouse; **D.** March 1, 2023 in the field.

Fig. 3 *BnaFLC* expression and flowering times of rapeseed plants under the vernalization models. **A.**

BnaFLCA02; **B.** *BnaFLCA10*; **C.** *BnaFLCC02* relative expression levels in Darmor, **D.** *BnaFLCA02*; **E.**

BnaFLCA10; **F.** *BnaFLCC02* relative expression levels in Bristol, **G.** *BnaFLCA02*; **H.** *BnaFLCA10*; **I.**

BnaFLCC02 relative expression levels in Helios. Histograms show mean values ± SEM for two

independent PCR amplifications on three biological replicates. The left panel y axis shows the relative

expression after normalization to *Ubiquitin* expression. The right panel y axis shows the flowering

time as the days after transplantation when the first flower was seen. DNF: did not flower. * $p <$

0.05; ** $p < 0.01$; *** $p < 0.001$

Fig. 4 Seed oil content (% w/w) and % oleic acid content of **A** and **B** Bristol; **C** and **D** Helios under the

vernalization models between 2021-2023. Histograms show mean values ± SEM for three biological

replicates.

Fig. 5 The expression profiles of the fatty acid synthesis-related genes in developing seeds under the

vernalization models between 2021-2013. Relative expression levels of *BnaFAD2* in **A.** Bristol, **B.**

Helios; *BnaFAD5* in **C.** Bristol, **D.** Helios; *BnaFATB* in **E.** Bristol, **F.** Helios; *BnaMCOA* in **G.** Bristol, **H.**

Helios; *BnaWD40* in **I.** Bristol, **J.** Helios are shown as histograms of mean values ± SEM for three

biological replicates. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

Fig. 6 Heat maps of Pearson correlation coefficients ($-1 \leq r \leq 1$) between the vernalization models

(NV, 3w, 4w, 6w and 8w) and the flowering time, *BnaFLC* gene expressions, fatty acid-related gene

expressions, seed oil content, oleic acid content and the yield-related traits of **A.** Bristol in 2021; **B.**

Bristol in 2022; **C.** Bristol in 2023; **D.** Helios in 2021; **E.** Helios in 2022; **F.** Helios in 2023. VM:

vernalization model, FT: flowering time, PH: plant height, LB: lateral branches, TSW: thousand seed

weight, NOPOTMS: number of pods on the main stem, SNPP: seed number per pod. * $p < 0.05$; ** p

< 0.01; *** $p < 0.001$

Fig. S1 Yield-related physiological traits under the vernalization models between 2021-2023. Plant

height (cm) of **A.** Bristol, **B.** Helios; lateral branches (n) of **C.** Bristol, **D.** Helios; number of pods on the

main stem of **E.** Bristol, **F.** Helios; seed number per pod of **G.** Bristol, **H.** Helios; thousand seed weight

(g) of **I.** Bristol, **J.** Helios. Histograms show mean values ± SEM for at least three biological replicates.

Fig. S2 The comparison of Bristol and Helios varieties grown in the field by direct sowing in Fall 2022.

A. Yield-related traits; plant height (PH), lateral branches (LB), thousand seed weight (TSW), number

of pods on the main stem (NOPOTMS), and seed number per pod (SNPP) are shown as histograms of

mean values ± SEM for at least three biological replicates. **B.** The expression profiles of the fatty acid

synthesis-related genes in developing seeds, and **C.** Seed oil content (% w/w) and oleic acid content

(%) are shown as histograms of mean values ± SEM for three biological replicates.

Fig. S3 The expression profiles of the fatty acid synthesis-related genes in leaves of seedlings under

the vernalization models in Darmor, Bristol and Helios. Relative expression levels of **A.** *BnaFAD2*; **B.**

BnaFAD5; **C.** *BnaFATB*; **D.** *BnaMCOA*; **E.** *BnaWD40* are shown as histograms of mean values ± SEM for

three biological replicates. * $p < 0.05$; ** $p < 0.01$; *** $p < 0.001$

NV	22 °C	22 °C	22 °C	22 °C	22 °C	22 °C	22 °C	22 °C	22 °C
3w	4 °C	4 °C	4 °C	22 °C	22 °C	22 °C	22 °C	22 °C	22 °C
4w	4 °C	4 °C	4 °C	4 °C	22 °C	22 °C	22 °C	22 °C	22 °C
6w	4 °C	4 °C	4 °C	4 °C	4 °C	4 °C	22 °C	22 °C	22 °C
2+4w	4 °C	4 °C	22 °C	4 °C	4 °C	4 °C	4 °C	22 °C	22 °C
8w	4 °C	4 °C	4 °C	4 °C	4 °C	4 °C	4 °C	4 °C	22 °C
2+6w	4 °C	4 °C	22 °C	4 °C	4 °C	4 °C	4 °C	4 °C	4 °C

1
2

3   Fig. 1

5   Fig. 2

3   Fig. 3

2   Fig. 4

2   Fig. 5

2   Fig. 6

1   Fig. S1

3   Fig. S2

5   Fig. S3