Sensory neurons regulate stimulus-dependent humoral immunity.

Diane Aguilar[1,2*], Fengli Zhu[1,2*], Antoine Millet[1,2], Nicolas Millet[2,3], Patrizia Germano[4,5,7], Joseph
Pisegna[5,6,8], Omid Akbari[9], Taylor A. Doherty[10], Marc Swidergall[2,3,11], Nicholas Jendzjowsky[1,2,11]

[1] Division of Respiratory and Critical Care Medicine and Physiology, Harbor-UCLA Medical
Center, Torrance, CA 90502, USA
[2]The Lundquist Institute for Biomedical Innovation at Harbor-UCLA, Torrance, CA 90502, USA
[3] Division of Infectious Disease, Harbor-UCLA Medical Center Torrance, CA 90502, USA
[4]Research Service, Veterans Affairs Greater Los Angeles Healthcare System, Los Angeles, CA
90073, USA.
[5]CURE/Digestive Diseases Research Center, Department of Medicine, University of California,
Los Angeles, CA 90073, USA.
[6]Division of Gastroenterology, Hepatology and Parenteral Nutrition, VA Greater Los Angeles
Healthcare System and Department of Medicine, Los Angeles, CA 90073, USA.
[7]Division of Human Genetics, David Geffen School of Medicine at UCLA, Los Angeles, CA
90095, USA.
[8]Division of Pulmonary and Critical Care, Veterans Affairs Greater Los Angeles Healthcare
System, Los Angeles, CA 90073, USA.
[9]Department of Molecular Microbiology and Immunology, Keck School of Medicine, University of
Southern California, Los Angeles, CA 90033, USA.
[10]Division of Allergy and Immunology, Department of Medicine, University of California San
Diego, Veterans Affairs San Diego Healthcare System, La Jolla, CA 92093, USA
[11]David Geffen School of Medicine, Los Angeles, CA 90073, USA

25
26 *Authors contributed equally
27

*Corresponding Author: Nicholas Jendzjowsky, PhD
Rm 209 Martin Research Building,
Respiratory Medicine and Exercise Physiology, The Lundquist Institute for Biomedical
Innovation at Harbor-UCLA Medical Center
Division of Respiratory and Critical Care Medicine and Physiology, David Geffen School of
Medicine, UCLA
1124 W Carson Street
Torrance, CA
90502-2006
Office:323-457-1663
Email: nicholas.jendzjowsky@lundquist.org

Graphical Abstract

Abstract

Sensory neurons sense pathogenic infiltration, informing immune coordination of host defense. However, sensory neuron-immune interactions have been predominantly shown to drive innate immune responses. Humoral memory, whether protective or destructive, is acquired early in life - as demonstrated by both early exposure to streptococci and allergic disease onset. Our study further defines the role of sensory neuron influence on humoral immunity in the lung. Using a murine model of *Streptococcus pneumonia* pre-exposure and infection and a model of allergic asthma, we show that sensory neurons are required for B-cell and plasma cell recruitment and antibody production. In response to *S. pneumoniae*, sensory neuron depletion resulted in a larger bacterial burden, reduced B-cell populations, IgG release, and neutrophil stimulation. Conversely, sensory neuron depletion reduced B-cell populations, IgE, and asthmatic characteristics during allergen-induced airway inflammation. The sensory neuron neuropeptide released within each model differed. With bacterial infection, vasoactive intestinal polypeptide (VIP) was preferentially released, whereas substance P was released in response to asthma. Administration of VIP into sensory neuron-depleted mice suppressed bacterial burden and increased IgG levels, while VIP1R deficiency increased susceptibility to bacterial infection. Sensory neuron-depleted mice treated with substance P increased IgE and asthma, while substance P genetic ablation resulted in blunted IgE, similar to sensory neuron-depleted asthmatic mice. These data demonstrate that the immunogen differentially stimulates sensory neurons to release specific neuropeptides that specifically target B-cells. Targeting sensory neurons may provide an alternate treatment pathway for diseases involved with insufficient and/or aggravated humoral immunity.

Abstract word count: 245

Keywords: sensory neurons, vagus, B-cell, plasma cell, substance P, vasoactive intestinal polypeptide, humoral immunity, immunoglobulin, *Streptococcus pneumoniae, Alternaria alternata*

Introduction

With each breath, the lung risks exposure to atmospheric infectious agents and allergens. Sensory neurons work alongside immune cells and influence host defense against infection and injury[1–7], but appear to exacerbate allergy[8–12]. Stimulation of sensory neurons by cytokines[4,5,8,12,13], proteases[11,14–17], PAMPs[1,1–3] and immunoglobulins[9,18–20] lead to neuronal depolarization. The release of neuropeptides by sensory neurons in response to infection or allergy[21–23], is a necessary cofactor in immune coordination[21,24]. The unique interplay between neurons and immune cells is pathology-dependent. A growing body of work demonstrates immune-stimulus-dependent neuropeptide release and, therefore, differential coordination of varying cell types[1–3,7,25–28]. Such differences speak to the dichotomous neuro-immune response, which can elicit a pro- or anti-inflammatory state[21,29].

Most current evidence involving sensory neuron and immune cell crosstalk involves cellular-mediated immunity. For example, sensory neurons inhibit bacterial clearance and fungal infection by suppressing Th1 and Th17-mediated cytokine release and, therefore, suppress neutrophil-mediated clearance of infection in response to acute infection in naïve mice[2,7]. Given that many infectious diseases are recurring, prolonged, or stem from commensal microbes becoming invasive[30,31], further inquiry into the neural influence on humoral immunity in response to infection is warranted.

Conversely, during allergy and asthma, vasoactive intestinal polypeptide (VIP)[8] and substance P[11] release, activate ILC2s[8] and dendritic cells[11] to increase eosinophilic infiltration[8] and mast cell degranulation[11]. However, the catalytic event that drives allergic symptoms[32] is a dysregulated increase of allergen-specific IgE and its downstream effects. Given the very high prevalence of allergic disorders, there is a great need to further understand the role of sensory neurons in regulating humoral immunity, which may also lead to future therapeutic targets.

B-cells are responsible for immune memory by class switch recombination, a process of VDJ recombination where heavy chains are cleaved from IgM to produce IgG, IgA, IgE, or IgD depending on the immune effector response required[33]. Early investigations have shown that B-cells respond to exogenous neuropeptides *in vitro* and can steer the predominant release of specific immunoglobulin[34–38]. For example, B-cells express neurotransmitter receptors for, substance P[34], vasoactive intestinal peptide[39,40], noradrenaline, and neuropeptide Y[41,42]. *In vitro*, some of these neurotransmitters appear to increase immunoglobulin release[34–38]. Sensory neurons also mediate B-cell development in the mucosa[39,43] and promote B-cell class switching in the spleen[44] and lung[10]. It was shown that sensory neuron ablation reduced IgE release subsequent to a reduction of B220+ cells in the lung, in response to house dust mite asthma and calcipitrol skin allergy[10]. Substance P was demonstrated to be responsible for IgE class switch recombination *in vitro,* and recent *in vivo* results support substance P-mediated class switch[10]. Our study investigates whether sensory neuron neuropeptide release can be a global effector for infection and allergy and whether the interplay between sensory neurons and B-cells is antigen-dependent.

We analyzed how sensory neurons traffic and stimulate B-cells in response to *Streptococcus pneumoniae* pulmonary infection and *Alternaria alternata*-induced allergic airway inflammation. We show that the release of sensory neuropeptides differs between our model of *S. pneumoniae* pre-exposure and infection and in an *A. alternata*-induced asthma model. Sensory neuron depletion suppressed select sensory neuropeptides in the lungs of each model and, as a

result, reduced pulmonary B-cell and plasma cell immunoglobulin release. Neuronal depletion
also exacerbated infection but improved asthma, proving the necessity of sensory neurons to
stimulate humoral immunity. Experiments using neuropeptide knockout mice for respective
neuropeptides support neuron depletion data. Supplementation of neuropeptides back into
sensory neuron-depleted mice in each model either improved bacterial clearance or worsened
asthma features due to a restoration of immunoglobulin production and release- thus supporting
the sufficiency of sensory neurons to regulate humoral immunity. In summary, sensory neurons
modulate B-cell production of immunoglobulins by variatïons in neurotransmitter release. The
release of neuropeptides is altered by the immune stimulus, which appears to provide a way to
steer the humoral response.

Results

TRPV1+ neurons mediate survival and bacterial clearance in pneumonia. Transient
receptor potential vanilloid 1+ (TRPV1) neurons have been demonstrated to influence T-cell-
mediated coordination of immunity by the release of specific neuropeptides[2,3,8]. The TRPV1 ion
channel responds to capsaicin, protons, prostaglandins, lipids and heat stimuli[45]. TRPV1+
neurons have been found to coordinate cellular immunity and influence T-cell stimulation of
neutrophils in acute *Staphylococcus aureus* infection[2] and eosinophils and mast cells in allergic
asthma[8]. Thus, we tested how TRPV1+ sensory neurons play a role in humoral immunity.

Using an established model of sensory neuron chemical ablation with resiniferatoxin (RTX)[2,46,47],
we suppressed TRPV1-containing neurons similarly in the vagal (nodose/jugular) ganglion by
>85% compared to TRPV1-DTR mice injected with the diphtheria toxin directly into the vagal
ganglia as assessed by qPCR and TRPV1 immunohistochemistry (**Supplementary Figure 1**).
RTX, administered by subcutaneous injection, induces global ablation of TRPV1+ neurons, thus
blocking sensory neuron vesicle emission of neuropeptides and rendering them
ineffective/atrophied[2,46,47]; we confirm the suppression of neuropeptide transcripts in the vagal
ganglia with RTX by qPCR (**Supplementary Figure 1**). Although there are indications that
dorsal root ganglia (distal site, primarily innervate skin and visceral organs[48]) provide some
innervation of the lungs[49], recent tracer studies have demonstrated sparse innervation with an
overwhelming majority of lung sensory neuron innervation provided by the vagal ganglia[50].

To investigate humoral immunity in response to *S. pneumoniae* lung infection, we aimed to
replicate the colonization and infection patterns observed in humans, considering that mice are
not naturally colonized with *S. pneumoniae*[51,52]. We established a model of *S. pneumoniae*
infection, which involved exposure of mice with a low inoculum on Day 0, and a subsequent
exposure to a high infectious dose of bacteria on Day 9 (**Figure 1a**). Following the low
inoculating dose, both sensory neuron intact and RTX mice showed a similar amount of bacteria
remaining 16hrs after the low inoculum (**Figure 1b**). Both groups were able to clear the bacteria
prior to the infectious dose (**Figure 1b**). Compared to naïve mice, our *S. pneumoniae* model
showed an initially increased bacterial burden, which was not different from RTX mice, 16hrs
after infection (**Figure 1b**). Forty-eight hours after the last infectious dose of *S. pneumoniae*,
sensory neuron intact mice almost completely cleared their bacterial burden, while RTX treated
mice exhibited an elevated bacterial burden in comparison to sensory intact mice and naïve
mice and this persisted 6 days after the final infection(**Figure 1b**). We further assessed the role
of sensory neurons in cross-protection against a lethal pneumococcal strain. Sensory neuron
ablation resulted in a higher bacterial burden and reduced survival when pre-exposed to
serotype 19F and infected with serotype 3 (**Figure 1c, d**). Additionally, in response to a single

high dose of *S. pneumoniae,* in both sensory neuron intact and RTX ablated mice, bacterial
burden and IgG remained unchanged, and when exposed to a single infectious dose of
serotype 3, survival was unchanged (**Supplementary Figure 2**); similar to previous
investigations[2]. Bacterial clearance between RTX-treated and targeted vagal TRPV1 ablation
with direct diphtheria toxin (DTX) injection into the vagal ganglia of TRPV1-DT receptor (DTR)
mice[53] was similar (**Supplementary Figure 3**). Consequently, our findings suggest that vagal-
specific sensory neurons are required to control *S. pneumoniae* during the course of lung
infection in our "recall" model.

Next, we determined whether nociceptors regulated pro-inflammatory-cytokine production
before lung infection clearance. Post-infection, RTX-treated mice showed reduced inflammatory
cytokine levels, including IL-1β, IL-6, and TNFα (**Figure 1e, Supplementary Figure 4**).
Furthermore, cytokines involved in B-cell stimulation (BAFF, TNFSF13, IL-2, and IL-5) and
chemokines involved in B-cell attraction (CXCL13, CCL5, and CCL7) were reduced in RTX mice
compared to control mice (**Figure 1e, Supplementary Figure 4**).

TRPV1+ neurons regulate B-cells.

Given that cytokines and chemokines for B-cell recruitment (**Figure 1e**) were reduced, we
assessed immunoglobulin levels in lung homogenates. Immunoglobulins were reduced in RTX
mice compared to sensory neuron intact mice after 16hrs of infection (**Figure 2b**). *S.*
pneumoniae serotype 19F specific IgG was also reduced in RTX as the IC50 calculated from
lung homogenate dilutions was higher in RTX compared to Veh (**Figure 2b**). The reduction in
immunoglobulins was directly due to a reduction in isotype-switched memory, resident memory
B-cells, plasma cells, and plasmablasts (**Figure 2c-g, Supplementary Figure 5-7**). It is
important to note that RTX did not affect B-cell populations immediately before pre-exposure or
after first exposure (**Figure 2c-g**). The reduction in B-cell populations remained significant at
48h (**Figure 2c-g, Supplementary Figure 6**), in direct accord with an augmented bacterial
burden in RTX mice (**Figure 1b,c**). As B-cells were diminished in RTX following infection, in
accord with immunoglobulins, our data are consistent with a role for sensory neurons with
adaptive immunity. In contrast, T-cells were unaffected by sensory neuron ablation
(**Supplementary Figure 8**).

To confirm that bacterial clearance in response to infection after pre-exposure relies on intact B-
cells, we compared bacterial burden from our sensory neuron intact and depleted mice to mice
lacking mature B-cells (µMT mice). Like RTX mice, µMT mice had a significantly increased
bacterial burden at 48hrs after our recall model of *S. pneumoniae* (**Figure 2h**),

Neutrophil depletion in sensory neuron intact mice significantly increased the bacterial burden to
a level of RTX mice (**Figure 2i, Supplementary Figure 9**). Neutrophil depletion did not further
augment bacterial burden in RTX mice, showing that the interaction between B-cells and
neutrophils may be an important mechanism of bacterial clearance in our recall model (**Figure
2i, Supplementary Figure 9**). To confirm the B-cell-neutrophil axis, naïve neutrophils were
incubated with decomplemented serum harvested from sensory neuron intact and RTX, which
underwent our pre-exposure and infection model, as well as naïve mice. Neutrophils that were
incubated with serum from sensory neuron intact mice had the highest level of bacterial killing
(**Figure 2j**). Therefore, the main effect of sensory neurons in response to infection is to
coordinate B-cell homing and immunoglobulin production to enhance neutrophil-mediated *S.*
pneumoniae clearance.

TRPV1+ neuron neuropeptide release of VIP stimulates B-cell immunoglobulin release. B-cells express select neuropeptide receptors[54]. Therefore, we reasoned that neurons would stimulate B-cells within their proximity. Using a sensory neuron reporter mouse (Vglutcre-tdTomato[50]), we showed that B220+ cells were localized within <20um of sensory neurons in the lungs 48h after the final infectious dose of *S. pneumoniae* (**Figure 3a**). In comparison, infected mice treated with RTX, showed a more dispersed distribution of B220+ cells, extending further from sensory neurons. Of note, whereas RTX depletes TRPV1 positive neurons, sensory neurons have a wide array of genetic composition[50,55], and therefore, some nerves remain following TRPV1 depletion (**Figure 3a**). Thus, it was likely that sensory neuropeptides released by TRPV1+ neurons were suppressed with RTX. Therefore, we measured known sensory neurotransmitters with noted immunological effects (CGRP[2], Substance P[10,11,56,56–58], VIP[8,40,59,60], NPY[35,42,61]) in lung homogenates. We show pre-exposure and infection increased NPY and VIP (**Figure 3b, c**). In comparison to sensory neuron intact mice, we demonstrate that RTX suppresses NPY, VIP, and CGRP (**Figure 3d, e**). The suppressed neuropeptide concentrations in the lung were associated with reduced cDNA expression in the vagal ganglion (**Supplementary Figure 1**). The neuropeptide concentrations measured from lungs are inversely correlated with bacterial burden and positively correlated with immunoglobulins (**Supplementary Figure 10**). Therefore, we hypothesized that NPY or VIP released from sensory neurons increase B-cell immunoglobulin release, to stimulate neutrophil-mediated bacterial clearance. Neuropeptides bind to specific receptors. We show that VIP1R and NPY1R are both present on B-cells and plasma cells, which demonstrates the likelihood of stimulation by sensory neuropeptides (**Supplementary Figure 11**).

We isolated splenic B-cells and cultured them with NPY or VIP. VIP stimulation increased IgG secretion (**Figure 3i**) and surface-bound levels in CD138+ plasma cells (**Figure 3f, g, Supplementary Figure 12**). However, neurotransmitter stimulation did not affect the proliferation of B-cells and plasma cells, assessed by Ki67+ intracellular staining (**Figure 3h**). Given the high expression of VIP1R on B-cells, we tested the effects of VIP stimulation in WT and VIP1R-/- isolated B-cells. In VIP1R-/- cultured B-cells, VIP and other neuropeptides did not increase bound and released IgG, demonstrating that VIP1R is the main receptor responding to VIP stimulation (**Figure 3j-l**). We then isolated B-cells from lungs of naïve, and pre-exposed and infected mice. VIP again increased bound and released IgG (**Figure 3m-o**). However, we noted that pre-exposed and infected mice had a suppressed response to all conditions. Therefore, we tested whether this suppression was due to B-cell exhaustion. B-cells harvested from pre-exposed and infected mice, which were then re-stimulated in culture for 96h, showed an augmentation of CD11c and reduction of CD23, to a greater extent than lung B-cells harvested and cultured from naïve mice (**Figure 3p, q**), indicative of B-cell exhaustion[62–64].

We supplemented sensory neuron intact and RTX sensory-depleted mice with VIP to test a functional outcome of VIP release in response to *S. pneumoniae* (**Figure 4a, Supplementary Figure 13**). VIP supplementation rescued RTX mice by suppressing their bacterial burden at 48h post-infection (**Figure 4n**). Furthermore, B memory, plasma cells, plasmablasts, and lung IgG were increased with VIP supplementation in RTX mice compared to RTX mice supplemented with PBS (**Figure 4b-g**). Finally, *VIP1R*[−/−] had an altered accumulation of select B-cell populations (**Figure 4h-k**), reduced immunoglobulins and cytokines (**Figure 4m, Supplementary Figure 14**), and increased bacterial burden in comparison to WT littermates in response to pre-exposure and infection (**Figure 4o**). In addition, when neutrophils were ablated with a neutralizing antibody, *VIP1R*[−/−] mice did not further increase the bacterial burden in their

lungs following pre-exposure and infection (**Figure 4p**). These data show that VIP stimulation of
B-cells through the VIP1 receptor increases B-cell neutrophil interaction through an increased
release of immunoglobulins.

In order to rule out non-neuronal sources of VIP, we stained for VIP in T-cells, macrophages and
neutrophils which others have identified as potential non-neuronal sources of VIP[65–67]. RTX did
not suppress VIP expression in leukocytes (**Supplementary Figure 15**). Neuroendocrine cells
within the lungs also have the potential to release neuropeptides[68]. However, VIP fluorescence
was not diminished with RTX in the epithelium where pulmonary neuroendocrine cells reside
(**Supplementary Figure 15).** It is possible that these alternate sources of neuropeptides may
contribute to B-cell stimulation. However, we note that in conjunction with the reduction of VIP in
RTX lung homogenates (which selectively ablate neurons, **Figure 3b**), the suppressed
expression of VIP transcripts in vagal ganglia in RTX compared to vehicle mice
(**Supplementary Figure 1)** and a maintained amount of VIP-positive lymphocytes and
neuroendocrine cells, the predominant source of neuropeptides in response to *S. pneumoniae*
is likely lung TRPV1+ sensory neurons.

Neural stimulation of immunoglobulins exacerbates asthma features. Given that
immunoglobulins and humoral immunity are important mechanisms for bacterial clearance, we
then asked whether an overproduction of immunoglobulins as a result of neural stimulation
could result in immune dysfunction. We hypothesized that this would be neuropeptide driven
but, likely mediated by an alternate peptide as previous data supports[10,58]. *A. alternata* causes a
swift and distinct rise of IgE in response to four inoculations (**Figure 5a**) as demonstrated
previously[69]. *A. alternata*-induced asthma increased airway hyper-responsiveness and goblet
cell metaplasia, which are reduced in sensory neuron depleted mice (**Supplementary Figure
16b-e**) where airway hyper-responsiveness was correlated with IgE (**Supplementary Figure
16c**). Specific B-cell recruitment of cytokines is suppressed with RTX sensory neuron ablation
(**Supplementary Figure 17**). RTX also suppressed B-cells, immunoglobulins, and mast cells in
asthmatic mice (**Figure 5c-h, Supplementary Figure 18**). *A. alternata* specific IgE was also
reduced in RTX mice as the IC50 calculated from lung homogenate dilutions was higher in RTX
compared to Veh mice(**Figure 5b**). NPY and Substance P were increased in the lungs during *A.
alternata*-induced asthma, in comparison to naïve mice, while decreased with RTX
(**Supplementary Figure 19a-d**), VIP was not increased in response to *A. alternata*-induced
lung inflammation (**Supplementary Figure 19d**). Substance P receptors (Neurokinin 1 and 2)
were not present in our isolated naïve B-cells (**Supplementary Figure 11**) but CDNA was
detectable in lung B-cells. Substance P also binds to mas-related g protein-coupled receptors[70],
where cDNA and protein were present on all B-cells (**Supplementary Figure 11**). NPY and
Substance P increased bound and released IgE in naïve splenic B-cells (**Supplementary
Figure12, 19e-h**). To test whether substance P could stimulate B-cells, we supplemented
substance P into *A. alternata* mice (**Figure 6a, Supplementary Figure 20**). We show that
substance P supplementation increased B-cell resident populations in the lungs of RTX sensory
neuron-depleted mice treated with *A. alternata* (**Figure 6b-f**). Further, IgE was increased when
A. alternata RTX mice were supplemented with substance P (**Figure 6l**). This was in line with
the removal of substance P, as substance P knockout mice (unable to produce substance P/
have receptors) showed reduced levels of IgE (**Figure 6m**), lung-resident B-cell populations
(**Figure 6g-k**), as well as cytokines and other immunoglobulins (**Supplementary Figure 21**) in
response to *A. alternata*-induced lung inflammation compared to WT mice with *A. alternata-
induced* lung inflammation. To account for alternate sources of substance P, we stained for

substance P in T-cells, eosinophils, and mast cells, which could be sources of substance P[65–67].
RTX did not suppress substance P expression in leukocytes (**Supplementary Figure 22**). We
note that substance P fluorescence was not diminished with RTX in the epithelium where
neuroendocrine cells are located (**Supplementary Figure 22**). In conjunction with the reduction
of substance P in RTX lung homogenates (**Supplementary Figure 19**), the suppressed
expression of substance P transcripts in vagal ganglia in RTX compared to vehicle mice
(**Supplementary Figure 1**) and a maintained amount of substance P positive leukocytes and
epithelium, we believe the predominant source of substance P in response to *A. alternata* are
lung sensory nerves. It is still possible that these alternate sources of neuropeptides, including
leukocytes and neuroendocrine cells, could contribute to B-cell stimulation. In summary, these
data demonstrate that substance P exacerbates asthmatic features in mice by increasing B-cell
production of IgE.

Discussion

Models of repeated immune stimulation show that sensory neurons may stimulate humoral
immunity[10]. In contrast, sensory neurons appear to suppress innate responses to select single-
dose models of bacterial and fungal infection[1–7]. Recent data show that sensory neurons
modulate B-cell populations in the lungs[10] and contribute to B-cell maturation in the mucosa[39,43].
Select neuropeptides, released by sensory neurons, can induce IgE, IgG, and IgA release *in*
vitro[34–38]. Therefore, we reasoned that sensory neurons could play critical roles in humoral
immunity[10,71–73]. The differential roles of neurons with innate and adaptive immunity may explain
why sensory neurons can produce divergent results depending on the underlying immune
stimulus, which is likely dependent on the timing and type of neuropeptide released.

Sensory neurons are able to sense pathogenic infiltration directly through PAMPs or indirectly
by binding cytokines and proteases[21–23]. Upon stimulation, sensory neurons release
neuropeptides directly into the immediate vicinity[21–23]. The type and amount of neuropeptide
released is likely dependent on the neuron type stimulated, which would dictate how it senses
the environment and which neuropeptides would be released[55,74]. Depletion of sensory neurons
reduced B-cell populations, IgG, and neutrophil activity, which subsequently augmented
bacterial burden in our recall model. This humoral regulation was due to the release of VIP,
likely from sensory neurons and the resulting stimulation of B-cells and plasma cells via the
VIP1 receptor. We demonstrate that B-cells and plasma cells have VIP1 receptors, which
stimulate PKCs that are critical to the development and release of immunoglobulins[75,76]. These
data are confirmed with VIP1 receptor knockout mice, highlighting the essential role of VIP1
receptor signaling in the proper functioning and immunoglobulin-secreting capability of B-cells
and plasma cells. While RTX treatment suppresses neuropeptide gene transcripts in sensory
neurons and reduces neuropeptide levels in lung homogenates, sensory neuropeptides may still
activate the release of neuropeptides from other sources, such as lymphocytes and
neuroendocrine cells. This potential alternative source of neuropeptide release means that our
current analysis might underestimate the overall stimulation of B-cells by neuropeptides, further
underscoring the importance of neuropeptide stimulation of B-cells.

Our results are consistent with early *in vitro* data which show that VIP can stimulate B-cell
immunoglobulin release[34–38]. In the context of bacterial lung infection, our results contrast with
the finding that sensory neurons suppress immunity in response to acute infection[2]. Specifically,
sensory neuron release of CGRP suppressed bacterial clearance of acute pulmonary
Staphylococcus aureus infection by dampening γδT cell-neutrophil stimulation[2]. When we used

a single dose of *S. pneumoniae*, we saw that sensory neuron depletion had a similar bacterial
burden and IgG as sensory neuron intact mice 48 hours following the single challenge as well
as survival in response to a lethal strain. In response to pre-exposure and infection, CGRP did
not significantly increase. We suggest that CGRP may be necessary for early innate
neuroimmune interaction, which is superseded by VIP upon re-infection, or that CGRP may
have an alternate role for B-cell maturation in germinal centers[77]. While humans are colonized
with pneumococcal strains at birth, mouse models appear to require pre-exposure prior to
infection in order to engage humoral memory and mimic the human condition[30,31,51,52]. Therefore,
our model of pre-exposure and infection may provide data that is more applicable to the
translation to clinical practice.

B-cells can become dysregulated and exacerbate diseases. In the context of allergen-induced
lung inflammation, we show that sensory neurons become hypersensitized and stimulate B-
cells. This is in direct accord with many investigations demonstrating that hypersensitization of
sensory neurons leads to augmented cytokine release and subsequent eosinophil trafficking by
T-cells or innate resident lymphocytes[8,9,28]. Our study extends these findings by showing that
sensory neuron depletion reduces B-cell-attracting cytokines and, as a result, reduces B-cell
recruitment. Mast cells were also reduced in sensory neuron-depleted mice. It is likely that the
combined reduction in B-cells and IgE release also reduced mast cell stimulation. Conversely,
mast cells also harbor neuropeptide receptors[78], and sensory neuron ablation could directly
reduce mast cell stimulation.

Substance P released by sensory neurons regulates B-cell stimulation in our asthma model. We
and others show the ability of substance P to induce class switch recombination[10]. We
demonstrate that mas-related gene protein-coupled receptors that are stimulated by substance
P[11] are ubiquitously expressed by B-cells with tachykinin receptor transcripts detectable in
pulmonary B-cells. Our data significantly extend previous data by showing that the chemical
ablation of neurons reduced B-cell populations and IgE levels. Further, when substance P was
supplemented in asthmatic sensory neuron-depleted mice, IgE was once again restored. Finally,
genetic ablation of the ability to produce substance P completely prevented IgE production in
response to *A. alternata.* Increased substance P production has been associated with an
asthmatic phenotype. Specifically, increased substance P-expressing neurons in asthma[79]
augmented mucus secretion and goblet cell hyperplasia[80–82], which contributes to overall airway
hyperresponsiveness[8], as we show. Tachykinins such as substance P, therefore, contribute to
asthma induction via direct airway stimulation and through IgE release.

While the direct effects of neuropeptides on B-cells have been demonstrated, recent findings
show that the lymph node innervation by sensory neurons is restricted to the macrophage-rich
medulla, not the lymphocyte-rich cortex[71,77,83]. Therefore, sensory neurons or neural circuits are
unlikely to recruit B-cells from lymph nodes or secondary lymphoid tissues. In line with this
reasoning, we show no changes in B-cells in the spleens or bone marrow in response to
sensory neuron depletion. Our data demonstrate that the effects of sensory neuropeptides are
confined to the lungs and produce these effects locally. Lung tissue-resident B-cells establish
themselves after primary infection or colonization[84,85], and are poised to release antibodies[86,87]
upon antigen encounter with a challenge infection. Our data demonstrate that B-cells cluster
around neurons in lungs and that the changes to B-cell numbers were only in the lungs. Further,
CD62L negative B resident cells were prominent in our model and reduced with sensory neuron
ablation, demonstrating that tissue-resident B-cells were employed upon infection after pre-

exposure. In contrast to the lengthy time course required for germinal center-mediated B-cell
maturation, our rapid antibody responses were consistent with tissue-resident extra-follicular B-
cell antibody production[88,89]. We suggest that neuropeptides act to assist in the maturation and
development of tissue-resident B-cells.

Our data provides a new role of sensory neuropeptides acting as B-cell stimulants and
highlights the humoral influence of sensory neurons. This study demonstrates that sensory
neurons are critical in regulating pulmonary humoral immunity and the outcomes of bacterial
lung infections and asthma. Targeting neuro-immunological communication through VIP and
substance P or other potentially uncovered molecular mechanisms may be an effective
approach to enhance host protection or reduce allergic responses by influencing
immunoglobulin production through accessory pathways.

METHODS

Mice. All animal experiments were approved by The Lundquist Institute at Harbor UCLA
Institutional Animal Care and Use Committee protocol # 32183. Mice were housed in a specific-
pathogen-free animal facility at The Lundquist Institute. C57BL/6J, Tac-/- (B6.Cg-
Tac1tm1Bbm/J), and μMT (B6.129S2-Ighmtm1Cgn/J) mice were purchased from Jackson
Laboratories. Vglut-TdTomato mice were provided by X. Sun (UCSD- crossed from Vglut2cre-
and RosaTdTomato originally attained from JAX B6J.129S6(FVB)-Slc17a6tm2(cre)Lowl/MwarJ
and B6.Cg-Gt(ROSA)26Sortm14(CAG-tdTomato)Hze/J). TRPV1-DTR mice were obtained from
Dr. Isaac Chiu, with MTA provided by M Hoon (NIH). VIP1R-/- were kindly donated by Joseph
Pisegna, Patrizia Germano, and James Waschek (UCLA, MGI: 177616). At the start of
experiments, mice was 5-8 weeks old. Age-matched male and female mice were used for
experiments.

***Streptococcus pneumoniae* cultures**. *S. pneumoniae* strain 49619 serotype 19F and strain
6303 serotype 3 were purchased from ATCC. Cultures were originally grown on Tryptic Soy
Agar (BD Biosciences 236950) supplemented with 5% defibrinated sheep's blood (Remel
R54012) and cultured for 24h, then re-cultured for an additional 24h in 5% CO_2 at 37°C.
Colonies were then picked for shape and hemolytic activity and cultured in Todd Hewitt Broth
supplemented with 2% Yeast Extract (Fisher 50489152) or Brain Heart infusion media (BD
DF0037178) in 5% CO_2 at 37°C. Cultures were adjusted to OD 600 1.0 and resuspended in
PBS prior to inoculation.

Infection model. Colonies were checked for OD, then spun at 4000 rpm, 10 min, and washed
in PBS. On day 0, mice were infected with 10^4 CFU of Serotype 19F in 50 μl of PBS,
intranasally. Then on day 9, mice were infected with 10^8 CFU of serotype 19F in 50 μl of PBS
intranasally. Bacterial burden were assessed in naïve mice (prior to any inoculation),
immediately following the first inoculum (Day 1), prior to the second infectious dose (Day 8), 16
hours after the infectious dose (Day 10), 48 hours after the infectious dose (Day 11) and 6 days
after the final infectious dose (Day 15). In a separate set of experiments serotype cross-
protection was tested. Mice were infected with 10^4 CFU of serotype 19F in 50 μl of PBS,
intranasally. Then on day 9, mice were infected with 10^6 CFU of serotype 3 in 50 μl of PBS
intranasally., Survival and bacterial burden (at 48h post-infection) were assessed. Control
animals were intranasally infused with 50μl PBS only.

Allergy model. *A. alternata* extract was purchased from CiteQ Laboratories (09.01.26) and
dissolved to yield a final concentration of 25ug/ml. Then mice were inoculated intranasally with

427 50ul on days 0, 3, 6, and 9 where experiments took place 16 or 48 hours later[90]. All control mice
428 received PBS in the same volume intranasally.

429 **Sensory neural ablation.** For chemical ablation of TRPV1+ neurons, mice 5 weeks of age
430 were treated with RTX (Adipogen 502053716) as previously described[2,4,7]. Mice were injected
431 subcutaneously in the dorsal flank on consecutive days with three increasing doses of RTX (30,
432 70, and 100 µg/kg) dissolved in 2% DMSO with 0.15% Tween 80 in PBS. Control mice received
433 vehicle.

434 For targeted TRPV1+ neuron depletion, we injected 20 ng DT in 200 nl PBS containing 0.05%
435 Retrobeads (LumaFluor INC) into nodose/jugular/petrosal VG with a nanoinjector (Drummond
436 Scientific Company)[91]. Mice were anesthetized with 87.5mg/kg ketamine and 12.5mg/kg
437 xylazine. The vagal ganglion was exposed after a midline incision in the neck (~1.5 cm in
438 length). DT was gently injected at a rate of 5nl/s. Once completed, the injection needle
439 remained in the ganglia for 2-3 minutes in order to reduce the spillover of DT. This process was
440 repeated for the vagal ganglion on the other side of the body.

441 **Bacterial burden determination.** Lungs were weighed and then homogenized in 2ml sterile
442 PBS with a tissue homogenizer. Homogenates were serially diluted on tryptic soy agar plates
443 with 5% sheep's blood and 10ug/ml neomycin (Fisher Biochemicals BP2669-5). The bacterial
444 CFU were enumerated after overnight incubation at 37°C and 5% CO_2.

445 **Neurotransmitter dosing.** Vasoactive intestinal peptide (200ng in 50 µl PBS) was delivered
446 intranasally at 48, 24, and 0h before and 24h after (for 48h harvest only) final infection with 10^8
447 CFU of *S. pneumoniae*. Control mice received PBS at the same time points. When mice
448 received bacteria on the same day, VIP was combined with the bacterial inoculum.

449 Substance P (1000ng in 50uL PBS) was delivered intranasally on Days 5,6,7,8,9 of the *A.*
450 *alternata* model. On days where *A. alternata* was delivered, Substance P was mixed with *A.*
451 *alternata.*

452 **Neutrophil depletion.** For neutrophil depletion, we followed an established protocol[92]. Mice
453 were injected i.p. with 150 µg of anti-A18 (clone 1A8, BioXCell BE-0075-1) (in 200 µl) per
454 mouse 48, 24, 0h before lung infection and 24h after lung infection (48h analysis only). Control
455 mice received 150 µg of rat IgG isotype control (2A3 BioXCell).

456 **Neutrophil bacteriocidal assay.** Mouse neutrophils were purified as described previously[93]. In
457 brief, bone marrow cells were flushed from femurs and tibias of 8 week old C57BL/6J mice
458 using sterile RPMI 1640 medium supplemented with 10% FBS and 2 mM EDTA onto a 50 mL
459 screw top Falcon tube fitted with a 100 µm filter. Mouse neutrophils were purified from bone
460 marrow cells using negative magnetic bead selection (MoJo Sort 480057, BioLegend) according
461 to the manufacturer's instructions. Bone marrow-enriched neutrophils had >98% purity and
462 >93% viability. Neutrophil killing of S*. pneumoniae* 19F was determined by CFU enumeration. *S.*
463 *pneumoniae* 19F was incubated with 50% decomplemented (20 min; 65°C) serum in PBS from
464 naïve, primary infected, and primary infected mice treated with RTX for 30 min on ice. Bacteria
465 were washed with PBS, and $1x10^3$ serum-coated *S. pneumoniae* 19F were incubated with
466 $1x10^5$ BM-neutrophils for 30 min. Neutrophils were lysed with 0.02% Triton X-100 in ice-cold
467 water for 5 minutes, diluted, and the remaining bacterial cells were quantitatively cultured.

Flexivent lung function. Mice anesthetized with ketamine/xylazine (87.5/12.5mg/kg), paralyzed with pancuronium bromide (MP Biomedicals ICN15605350), and subject to methacholine challenge with 200µl of doubling doses of methacholine (MP Biomedicals ICN19023110) dissolved in PBS for 20 breaths. Single and double compartment model Respiratory resistance was measured with the Flexivent system (SCIREQ) as previous[91]. Both absolute and percentage change values from PBS were calculated.

Tissue collection. For flow cytometry, mice were euthanized by Isoflurane inhalation. The lungs were then dissected and flushed with PBS, coarse dissected, and incubated at 37°C for 45min in 1.75mg/ml collagenase IV (C4-22-1g Sigma) in PBS; then washed, macerated through a 21 gauge needle and filtered through a 70 µm mesh filter, then treated for flow cytometry. Spleens were macerated through a 70 µm mesh filter and then treated for flow cytometry. Bone marrow was flushed with RPMI media supplemented with 2mM EDTA and 10% FBS, filtered through a 70 µm mesh filter, then treated for flow cytometry.

For cytokine analysis, lungs were dried, weighed, then macerated in 2ml PBS supplemented with 25ul protease inhibitors (consisting of AEBSF HCl (100 mM), Aprotinin (80 µM), Bestatin (5 mM), E-64 (1.5 mM), Leupeptin (2 mM) and Pepstatin (1 mM) Tocris 5500), spun at 8000g 4 min 4C and then the supernatant was snap frozen and stored t -80°C until analysis.

B-cell isolation. B-cells were isolated from spleens or lungs (below) using the EasySep Mouse B-cell negative selection kit. CD138+ cells (for qPCR, below) were isolated using the EasySep Mouse CD138 positive selection kit.

Quantitative real-time PCR. RNA was isolated from tissues or cells using the inTron Easy spin Total RNA extraction Kit (Boca Scientific 17221), which was reverse transcribed to cDNA with a Tetro cDNA Synthesis Kit (Tetro biosystems NC1352749). Relative gene expression was determined by quantitative real-time PCR on a QuantStudio 3 System (Thermo Fisher Scientific) with TaqMan Fast Advanced Master Mix (Thermo Fisher Scientific 4444557) with the TaqMan probe sets (ThermoScientific Applied Biosystems). Expression values relative to HPRT were calculated by Δ ΔCTT method. All taqman probes utilized were purchased from Thermofisher Vip1R Mm00449214_m1; Vip2R Mm01238618_g1; Tac1R Mm00436892_m1; Tac2R Mm01175997_m1; NPY1R Mm00650798_g1; NPY2R Mm01956783_s1; NPY4R Mm00435894_s1; NPY5R Mm02620267_s1; NPY6R Mm00440546_s1; MRGPRA Mm01984314_s1; MRGPRB2 Mm01956240_s1; MRGPRG Mm01701870_s1; RAMP1 Mm00489796_m1; Tac1 Mm01166996_m1; Calca Mm00801463_g1; VIP Mm00660234_m1; NPY Mm01410146_m1; TRPV1 Mm01246300_m1; HPRT Mm03024075_m1).

Cytokine, neurotransmitter, and immunoglobulin analysis. Cytokine and immunoglobulin levels in lung homogenates were measured through Luminex multiplex assay using R&D discovery or Thermo kits; analytes were measured with the Luminex 200 system. Enzyme-linked immunosorbent assay (ELISA) kits were used according to the manufacturer's instructions (R&D systems, Raybiotech, or Thermofisher) and analyzed with a BioTek Synergy H1 plate analyzer. Luminex and ELISA kits are all commercially available and listed in the supplementary materials list.

Specific immunoglobulin analysis. For specific IgG determination, crude S. pneumoniae (12.53µg/ml ATCC 49619) extract was incubated in 96 well plates overnight at 4°C, then blocked with goat serum for 1h at 4°C. Then log dilutions of lung homogenates (from groups above) were incubated at RT for 2 hours. Goat anti-mouse IgG HRP conjugated secondary

antibody (Abcam ab205719, 1:1000) was then added and incubated for 2 hours RT. The ELISA
was developed with 0.5% TMB solution (Fisher AAJ61325AP) and stopped with 2N H2SO4
(Fisher 828016). For specific IgE determination, *A. alternata* (100µg/ml CiteQ 09.01.26) was
incubated as above and developed as above, except Goat anti-mouse IgE HRP secondary
(Thermofisher PA1-84764, 1:1000) was used. OD450 for each dilution was used with a 4-point
sigmoidal curve to determine IC50 for each sample in each group.

Flow cytometry. Red blood cells were lysed with ACK lysing buffer (Gibco A10492-01), treated
with Fc Block (Biolegend Trustain 101320), and resuspended in FACS buffer (HBSS-Gibco
10010-023 with 2% FBS Gibco. Incubations with antibody cocktails were conducted at 4°C for
60 min, and samples were subjected to two washes and resuspension in FACS buffer (HBSS
+2% FBS Sigma F8192). For intracellular staining, cells were fixed/permeabilized with BD
cytofix/cytoperm kit (554714), washed, and stained overnight at 4°C. Flow cytometry was
conducted on a Symphony A5 flow cytometer (BD). Data were collected with BD DIVA software,
and files were analyzed with FlowJo (Treestar, version 10.0.8r1). A live-cell stain (APC-Cy7,
Invitrogen) was used to exclude dead cells. Gating strategies are provided in **Supplementary**
Figure 5. Positive staining and gates for each fluorescent marker were defined by comparing
full stain sets with fluorescence minus one (FMO) control stain sets. All antibodies were used at
1:100. Antibodies used: Biolegend-CD11c (Clone N418), Ly6g (Clone 1A8), CD117 (Clone 2B8),
Siglec F (Clone 1RNM44N Invitrogen), F4/80 (Clone BM8), CD3 (BUV805), CD3 (Clone17As),
CD11c (Clone N48), CD4 (Clone GK1.5), CD8 (Clone SK1), T-bet (Clone 4B10), GATA3 (Clone
16E10A23), B220 (Clone RA3-6B2), CD19 (Clone 6D5), CD62L (Clone MEL-14), CD44 (Clone
IM7), CD138 (Clone 281-2), IgG1 (Clone RMG1-1), IgE (Clone RME-1) CD45 (Clone 30-F11),
MRGPRX (Clone K125H4); BDbiosciences- IgM (Clone IL/41), IgD (Clone 11-26C.2a), RorγT
(Clone Q31378; Invitrogen- CD38 (Clone 90), Ki67 (Clon SolA15); AlomoneLabs- VIP1R (Clone
AB_2341081), Tac1 (polyclonal Proteintech), VIP (polyclonal Proteintech).

All in vivo cells were gated on leukocytes, singlets, live, CD45. Mast cells were
CD117+/MHCII+. Netrophils were Ly6g+, Eosinophils were SiglecF+/CD11b+. T cells were
divided into CD4+ or CD8+ fractions and intracellularly stained for Tbet (Th1), GATA3 (Th2) or
RORgammaT (Gamma delta). B-cell lineage were B220+/CD19+. These were subdivided based
on B Memory (CD44+/CD38+), B Resident Memory (CD44+/CD38+/IgD-/CD62L-), Isotype
switched (IgM-/IgD-). Plasma cells were CD138+,plasmablasts were CD138+/B220-/CD19-.

For assessment of of either intracellular neuropeptides or neuropeptide receptors, cells were
assessed from CD45+, live singlets and VIP or SP was stained intracellularly on CD3+,
CD117+, Siglec F+, Ly6G+, or F4/80+ cells. VIP1R, MRGPRA1 was assessed on B220+/CD19+
or CD138+ cells.

For in vitro cultures cells were divided into B220+/CD19+ or CD138+. Then based on these
gates assessed for IgG1 or IgE positivity. Ki67 was assessed from either CD138+/IgG1+ or
CD138+/IgE+ gates. Exhaustion was assessed as CD11c+/CD23- from CD138+ and
B220+/CD19+ gates.

Immunofluorescence, histology, and microscopy. The lungs were perfused with PBS for
immunostaining, followed by 4% paraformaldehyde (PFA) in PBS. Lungs were dissected and
postfixed overnight in 4% PFA/PBS at 4 °C, incubated at 4 °C with 30% sucrose/PBS for two
days, and stored in 0.1% sodium azide in PBS until cutting. Lungs were embedded in optimal
cutting temperature compound (OCT, Tissue-Tek, PA), and 50-µm cryosections were cut at -

20°C and then blocked for 4 h in PBS with 10% donkey serum, 2% bovine serum albumin (BSA)
and 0.8% Triton X-100. Sections were immunostained with the following antibodies: B220
(Biolegend 103225; 1:100 FITC), CD45 (Invitrogen 14045182 1:200), E-cadherin
(Thermoscientific MA512547 1:200), VIP (Thermoscientific PA578224, 1:200), Substance P
(Thermoscientific PA5106934 1:200) with appropriate secondary antibodies (Goat anti mouse
Cy3 Jackson Immuno 115-165-003; Goat anti rat Alexa 647 Jackson Immuno 112-605-003;
Goat anti rabbit Alexa 488 Abcam AB150077), washed and mounted with prolong antifade
diamond (Thermofisher P36961).

Vagal (nodose/jugular) and dorsal root ganglia were dissected, fixed in 4%PFA for 1 hour,
cryopreserved in 30% sucrose overnight (all 4C), then placed in OCT, and 14µm sections were
cut and placed on gelatin-coated slides. Sections were stained with TRPV1 (Alomone labs
Clone- AB_2313819 1:200) followed by incubation with secondary antibody and secondary
(Goat Anti Rabbit IgG H&L, Abcam). All sections were mounted with a Prolong diamond antifade
compound (Thermofisher P36961).

For histology, lungs were embedded in paraffin and cut on a microtome (Leica) 5µm and stained
with hematoxylin (Epredia 6765001) and eosin (Epredia 6766007), gram stain (Sigma HT90T-
1KT) or periodic acid, Schiff's reagent (Sigma395B-1kit). Blood smears were stained with
Giemsa solution (AB150670). Additionally, to confirm neutrophil counts in select experiments,
Giemsa stain (Abcam AB150670) was used on lysed blood smears. Histological slides were
mounted with permount toluene solution.

Sections were either imaged with a Leica SP8 confocal microscope or Leica Thunder system.
Data were imaged and analyzed with Leica LASx software. For quantification of the proximity of
B-cells to nerves, LASx online calipers were used to measure the distance from cell-to-nerve in
3D from the z-stack.

B-cell culture. Spleens or lungs were used to isolate B-cells under sterile conditions. B-cells
were isolated using the EasySep Mouse B-cell separation kit (StemCell Technologies 19854A).
Cells were then treated in B-cell culture media made using 438 mL RPMI 1640 (Gibco 11835-
030), 2.5mL 1 M HEPES buffer (Sigma H3662), 5 mL GlutaMAX (Thermo Fisher Scientific
35050061), 5 mL penicillin/streptomycin stock (Sigma P0781), 30 mL heat-inactivated fetal
bovine serum stock (Gibco A38400-01), and 454 µL β-mercaptoethanol (Gibco 21985-023) and
vacuum filter sterilized (0.22 µm filter)[94]. Cells were then treated with either media + IL4
(20ng/ml, Sigma Aldrich I1020-5UG) + LPS (10µg/ml, L6529-1MG Sigma Aldrich),
media+IL4+LPS+NPY (50ng/ml, Tocris 1153), media+IL4+LPS+VIP (50ng/ml, Tocris 1911) or,
media+IL4+LPS+Substance P (50ng/ml, 1156 Tocris). After 96 hours media was stored at -80°C
until immunoglobulin analysis. Cells were stained and analyzed with a Symphony A5 flow
cytometer and data were processed with FlowJo (Treestar 10.881).

Sample size and statistical analysis. We used animal numbers between 5 and 12 mice per
experimental group/genotype for bacterial burden studies. For measurement of cytokine,
neuropeptide, and immunoglobulins, 5-8 mice per group/genotype were used. For flow
cytometry 5-10 mice per group/genotype were used. For in vitro experiments, at least 3 animals
were pooled into one sample and 5-7 replicates were used. Bacterial burden data were
analyzed with the two-way RM ANOVA (time x group) with either Tukey or Holm-Sidak posthoc
test; FACS and cytokines were compared with ANOVA with Tukey or Holm-Sidak posthoc tests,
two-tailed unpaired t-tests for parametric analyses, or Mann–Whitney test for nonparametric

analyses. Correlations were run with Pearson correlation. Data were plotted in Prism
(GraphPad).

Acknowledgments

NJ is supported by NIH grant R21AI159221, R56AI175328, UCLA CTSI UL1TR001881-01 and
T32KT4708 of the Regents of the University of California Tobacco-Related Diseases Research
Program. T.A.D. is supported by NIH AI171795 and Veterans Affairs BLR&D BX005073.N.M.
and D.A. are supported by a California Institute for Regenerative Medicine Stem Cell Biology
Training Grant EDUC4-12837. The content is solely the responsibility of the authors and does
not necessarily represent the official views of the National Institutes of Health. We are grateful to
Dr. Isaac Chiu (Harvard) for providing TPV1-DTR mice as per the direction of Dr. Mark Hoon
(NIH). We are grateful to Dr. Xin Sun for providing Vglut-Tdtomato mice. Figures 1a, 2g, 2h, 4a,
4p, 5a, 6a and Supplementary Figures 2a, 16a were made using Biorender.

Author contributions

613   D.A. and F.Z. contributed to experimental design, conducted experiments and analyzed data.
614   A.M. and N.M. contributed to experimental execution and data analysis. P.G. J.P., T. A.D, O.A.
615   assisted with experimental design and analysis. M.S contributed to study design and manuscript
616   preparation. N.J. designed the study, conducted experiments, analyzed data, prepared the
617   manuscript and figures. All authors agree on the manuscript.

Conflict of interest statement

619   D.A, M.S. and N.J. declare the following competing interest. U.S. Patent Application Serial
620   Number 63/492,846. Methods of using sensory neuron neurotransmitters to enhance humoral
621   immunity. The remaining authors do not declare competing interests.

     Figure legends

Figure 1. Sensory neurons are required for successful clearance of *S. pneumoniae*
following pre-exposure and infection. a) Model of *S. pneumoniae* pre-exposure and infection.
Mice received escalating doses of resiniferatoxin (RTX) or Vehicle (Veh) starting 21 days prior
to inoculation. 10^4 CFU of *S. pneumoniae (Serotype 19F ATCC49619)* in 50µl PBS was
delivered on day 0 to expose mice, then 10^8 CFU in 50µl PBS was delivered on Day 9. Bacterial
burden was enumerated on Day 10, 11 and 15. Red asterisk and dotted line denote the
separate set of experiments assessing cross-protection between serotypes: 10^4 CFU of *S.*
pneumoniae (Serotype 19F ATCC49619) was delivered on day 0 to expose mice, then 10^6 CFU
(Serotype 3, ATCC6303) in 50µl PBS was delivered on Day 9, survival was assessed as was
bacterial burden on Day 11. **b)** Bacterial recovery from lungs in untreated sensory neuron intact
mice (Naïve), untreated sensory neuron ablated mice (Naive RTX), pre-exposed and infected
sensory neuron intact (Veh) or pre-exposed and infected sensory neuron ablated mice (RTX).
Naïve n=6, Naïve RTX: n=5, Veh: n=5-13, RTX: n=5-7. Two-way ANOVA with Newman-Keuls
post hoc test. Data were pooled from three independent experiments. Gram stain shows *S.*
pneumoniae infection (purple and green arrows point to example of gram-positive space) was
greater in RTX compared to Veh. Scale bar=20µm. **c)** Veh mice had reduced bacterial burden in
response to primary inoculation with serotype 19F and infection with serotype 3 (Veh n=7, RTX
n=6). Data were compared with two-sided t-test. **d**) 90% of Veh mice survived infection with
serotype 3 compared to 20% of RTX mice after inoculation with 19F (Veh n=8, RTX n=7). Data
were compared with Log-rank (Mantel-Cox) test. **e)** Quantification of B cell activating factor
(BAFF), APRIL13 (TNFSF13), TNFα, IL-1β, IL-2, IL-6, IL-5, CXCL13, CCL5, CCL7 in sensory
neuron intact (Veh n=7) and sensory neuron ablated (RTX, n=6) mice 16h following the final 10^8
CFU dose of *19F.* Samples were analyzed with Luminex bead assay. IL-6 was re-run with
ELISA. Two-sided t-test. Data were pooled from 2 independent experiments.

648
Figure 2. Sensory neuron ablation reduces, immunoglobulins, B-cells and disrupts
neutrophil effector function after *S. pneumoniae*. a) In response to our pre-exposure and
infection model (as in Figure 1a) **b)** IgA, IgM, IgG1, IgG2a, IgG2b, IgG3 were reduced with
sensory neuron ablation 16h (Day 10) after final infection. Veh: n=7, RTX: n=7. Isotyping
Luminex was used to analyze lung homogenates. Two-sided T-test. IgG specific to *S.*
pneumoniae was tested with crude *Sp* extract. Serial dilutions were used to model the IC50
with a 4-point log model to determine differences between Veh (n=8) and RTX (n=6). Two-sided
T-test. Data were collected from two independent experiments. **c)** B-memory
B220+/CD19+/CD44+/CD38+ cells, **d)** B resident memory B220+/CD19+/CD44+/CD38+/IgD-
/CD62l- cells, **e)** Isotype switched B220+/CD19+/IgM-/IgD- cells, **f)** Plasma CD138+ cells and,
g) Plasmablast CD138+/B220-/CD19- cells were downregulated by sensory neuron ablation at
16 and 48h after infection. Only B memory and plasma cells were different between Veh and
RTX on Day 8. Two-way ANOVA with Holm-Sidak test. Veh: n=5-7, RTX n=5-7. **h)** µMT mice
had reduced burden at 16h in comparison to sensory neuron intact mice (Veh). Sensory neuron
ablated (RTX) mice demonstrated a bacterial burden similar to mice with immature B cells
(µMT) 48h after infection. Sensory neuron intact mice (Veh) reduced their bacterial burden at
48h. Veh n=13, RTX n=5, uMT infected n=7. Data from four independent experiments. Two-way
ANOVA with Holm-Sidak tests. **i)** Ly6G A18 neutrophil depletion antibody or isotype control was
delivered before, during and after the final infection dose of *S. pneumoniae.* Bacterial burden
(Log CFU/ g lung mass) from lung homogenates was greater with neutrophil depletion in
sensory neuron intact infected mice compared to isotype control 48h (Day 11) after infection.
Sensory neuron depleted mice (RTX) did not further increase bacterial burden with neutrophil
depletion demonstrating that B-cell neutrophil interaction was disrupted. One-way ANOVA with
Holm-Sidak post-hoc test. Veh(Iso): n=12, Veh(Ly6g): n=8, RTX(iso): n=10, RTX(Ly6g):n=9.
Data were pooled from two independent experiments. **j)** Naïve neutrophils were incubated with
decomplemented serum harvested from naïve, Veh or RTX mice (pre-exposed and infected)

and plated in S pneumoniae 19F coated wells (5000 CFU). Neutrophils were lysed 30 minutes
later and plated on blood agar plates and CFU were enumerated 24h later. One-way ANOVA
with Holm-Sidak post-hoc, n=6 per group.
678

Figure 3. Vasoactive intestinal peptide attracts B-cells and regulates immunoglobulin
production and release. a) B220+ cells cluster around sensory nerves from lungs of sensory
neuron intact mice (Veh). In RTX sensory neuron depleted mice, B220+ cells are further
dispersed. 30 cells (10 cells per sample) from n=3 mice for Veh & RTX. Two-sided t-test. **b-e)**
Prominent sensory neuron neuropeptides were increased in lung homogenates 16 hours after
pre-exposure and infection with *S. pneumoniae*. **b)** VIP was increased with infection and
suppressed with sensory neuron depletion. **c)** NPY was increased with infection and
suppressed with sensory neuron depletion. **d)** Substance P did not significantly increase
following infection with *S pneumoniae.* **e)** CGRP did not significantly increase following infection
with *S pneumoniae.* Lung homogenates analyzed with ELISA. PBS: n=6, Veh: n=7, RTX: n=7.
One-way ANOVA with Tukey's post hoc test. Data from 2 independent experiments. **f-i)** B cells
were isolated from spleens of naïve mice and cultured in media with IL4+LPS and
supplemented with VIP or NPY. Cells and media were sampled after 96 hours of incubation. **f)**
VIP did not increase IgG bound to B220/CD19+ cells. **g)** VIP significantly increased bound IgG
in CD138+ plasma cells. **h)** Proliferation as assessed by intracellular Ki67 staining was not
affected by neuropeptides (from CD138+ gate). **i)** Secreted IgG was increased by VIP and NPY.
n=5-10. One-way ANOVA and Tukey's post-hoc test. Data from 2 independent experiments. **J-l)**
B-cells were isolated from the spleens of VIP1 receptor knockout (VIP1R-/-) and WT littermates
and cultured for 96h. **j)** B220/CD19+ cells were not significantly affected by VIP1R deletion. **k)**
VIP1R-/- CD138+ cells did not significantly upregulate IgG in response to VIP and other
neuropeptides. **l)** IgG released into the media was not increased with VIP or other peptides with
VIP1R genetic deletion; VIP increased WT B-cell IgG release. Two-way ANOVA with Holm-
Sidak post-hoc test. N=4-7. **m-p)** B-cells were isolated from Lungs of naïve or *S. pneumoniae*
pre-exposed and infected mice. **m, n)** VIP increased bound IgG in B220+/CD19+ and CD138+
cells. This effect was greater in naïve cells. **o)** released IgG was increased with VIP in both
naïve and pre-exposed and infected cells. **p,q)** Pre-exposed and infected cells show higher
markers of B cell exhaustion after culture. Two-way ANOVA with Holm-Sidak post-hoc. Naïve
n=4-7, S pneumoniae n=4-7.
707

Figure 4. Vasoactive intestinal peptide supplementation improves bacterial clearance,
memory B cells, plasma cells and IgG in response to pre-exposure and infection with *S.*
***pneumoniae*. a)** Vasoactive intestinal peptide (10ng/µl) or vehicle (PBS) was delivered as per
the schematic in addition to the model of *S. pneumoniae* pre-exposure and infection with
serotype 19F*.* Cells were gated on Live/CD45+ cells and total populations assessed with
counting beads. **b)** B memory cells, **c)** B resident memory, **d)** Isotype switched, **e)** Plasma cells,
f) plasmablasts were reduced with sensory neuron ablation. Supplementation with VIP
increased B-cell populations. VIP supplementation did not further increase B-cell populations in
sensory neuron intact mice. **g)** IgG was reduced with RTX as previous but increased with the
supplementation of VIP to RTX mice. **h)** VIP 1 receptor knockout (VIP1R-/-) mice had reduced B
memory cells compared WT mice, **i)** B resident memory and, **j)** Isotype switched B-cells were
similar in VIP1R-/- and WT mice, **k)** Plasma cells, and l) plasmablasts were reduced in VIP1R-/-
vs WT mice. Two-sided T-test. WT n=5, VIP1R-/- n=5. **m)** VIP1R-/- mice had reduced IgG
following pre-exposure and infection compared to WT littermates. **n)** VIP supplementation
suppressed bacterial burden in RTX mice. No effect of VIP supplementation was observed in
Veh mice. **o)** VIP1R-/- mice have increased bacterial burden following pre-exposure and
infection with *S. pneumoniae* compared to WT littermates*.* Veh (PBS): n=8, Veh (VIP): n=7-8,

RTX(PBS): n=6-7, RTX(VIP): n=6-7. One-way ANOVA with Holm sidak post hoc test. Data from
3 independent experiments. WT n=8-11, VIP1R-/- n=5-7. Two-sided t-test. Data from 3
independent experiments. **p**) Ly6g neutralizing antibody or isotype was given to WT or VIP1R-/-
mice. Bacterial burden (Log CFU/ g tissue mass) from lung homogenates was greater with
neutrophil depletion in WT mice compared to isotype control. VIP1R-/- mice did not further
increase bacterial burden with neutrophil depletion demonstrating that VIP significantly
stimulates B-cells to increase neutrophil-mediated bacterial clearance. One-way ANOVA with
Holm-Sidak post-hoc test. WT(Iso): n=6, WT(Ly6g): n=5, VIP1R-/-(Iso): n=4, VIP1R-/-
(Ly6g):n=5. Data were pooled from two independent experiments.
734

Figure 5. Sensory neuron ablation reduces B-lymphocytes, immunoglobulins and, mast
cells in *A. alternata* treated mice. a) *A. alternata* induced asthma was elicited as per
Cavagnero et al. 25µg of *A. alternata* extract in 50µl PBS was delivered as per the schematic.
Experiments took place 16hrs later. **b)** IgE, IgG1, IgG3 (n=6, RTX: n=7) were reduced with
sensory neuron ablation after final *A. alternata dose.* Isotyping Luminex was used to analyze
lung homogenates. *A. alternata* extract with lung homogenates and IC50 modeling was used to
determine IgE specific to *A. alternata* (Veh n=6, RTX n=4) Two-sided t-test. Data were collected
from two independent experiments. **c**) B-Memory B220+/CD19+/CD44+/CD38+ cells, **d**) B
resident memory B220+/CD19+/CD44+/CD38+/IgD-/CD62l- cells, **e**) Isotype switched
B220+/CD19+/IgM-/IgD- cells, **f**) Plasma CD138+ cells and, **g**) Plasmablasts CD138+/B220-
/CD19- cells and **h**) mast cells CD117+ were downregulated by sensory neuron ablation 16h
after *A. alternata* asthma induction. Cells were gated on live/CD45+ cells and counting bead
calculation determined total cell population. Red gate is the measured cell population. Veh:
n=10, RTX: n=7. Two-sided t-test. Data from 3 independent experiments.
749

Figure 6. Substance P increases B-cells and IgE in mice treated with *A. alternata*. a)
Substance P (100ng/µl) or vehicle (PBS) was delivered in conjunction with *A. alternata* extract
as per the schematic, in accordance with our *A. alternata* model of asthma. Cells were gated on
Live/CD45+ cells and total populations assessed with counting beads. **b**) B memory cells, **c**) B
resident memory, **d**) Isotype switched, **e**) Plasma cells, **f**) plasmablasts were reduced with
sensory neuron ablation. Supplementation with Substance P increased B-cell populations.
Substance P supplementation did not further increase B-cell populations in sensory neuron
intact mice. Veh (PBS): n=5, Veh (Sub P): n=6, RTX(PBS): n=6, RTX(SubP): n=6. One-way
ANOVA with Holm sidak post hoc test. Data from 3 independent experiments. Tac1-/- (lack
Tac1 gene, unable to produce substance P, have receptors) had similar B memory (**g**), but
reduced B Resident Memory (**h**), Isotype switched (**i**), plasma cells (**j**), and plasmablast cell (**k**)
compared to WT mice receiving *A. alternata* induction of asthma. WT n=6, Tac1-/- n=8,
Student's unpaired t-test. Data from 2 independent experiments. **l**) Supplementation with
substance P increased IgE in *A. alternata* mice treated with RTX to a level consistent with Veh.
Substance P did not further increase IgE in Veh mice (All groups n=6). One-way ANOVA with
Holm sidak post hoc test. Data from 2 independent experiments. **m**) Substance P knockout
mice (Tac-/- n=7) had reduced IgE compared to WT *A. alternata* mice (n=8). Two-sided t-test.
Data from 2 independent experiments.
768

772 **References**

773 1. Chiu, I. M. *et al.* Bacteria activate sensory neurons that modulate pain and

774   inflammation. *Nature* 501, **52–57 (2013).**

775 2. Baral, P. *et al.* Nociceptor sensory neurons suppress neutrophil and γδ T cell

776   responses in bacterial lung infections and lethal pneumonia. *Nat Med* 24, **417–426**

777   **(2018).**

778 3. Pinho-Ribeiro, F. A. *et al.* Blocking Neuronal Signaling to Immune Cells Treats

779   Streptococcal Invasive Infection. *Cell* 173, **1083-1097.e22 (2018).**

780 4. Riol-Blanco, L. *et al.* Nociceptive sensory neurons drive interleukin-23-mediated

781   psoriasiform skin inflammation. *Nature* 510, **157–161 (2014).**

782 5. Ramirez, V. T. *et al.* Sensory Nociceptive Neurons Contribute to Host Protection

783   During Enteric Infection With Citrobacter rodentium. *The Journal of Infectious*

784   *Diseases* 221, **1978–1988 (2020).**

785 6. März, P. *et al.* Sympathetic neurons can produce and respond to interleukin 6.

786   *Proceedings of the National Academy of Sciences* 95, **3251–3256 (1998).**

787 7. Kashem, S. W. *et al.* Nociceptive Sensory Fibers Drive Interleukin-23 Production from

788   CD301b+ Dermal Dendritic Cells and Drive Protective Cutaneous Immunity. *Immunity*

789   43, **515–526 (2015).**

790 8. Talbot, S. *et al.* Silencing Nociceptor Neurons Reduces Allergic Airway Inflammation.

791   *Neuron* 87, **341–354 (2015).**

792 9. Crosson, T. *et al.* FcεR1-expressing nociceptors trigger allergic airway inflammation.

793   *J Allergy Clin Immunol* 147, **2330–2342 (2021).**

794 10. Mathur, S. *et al.* Nociceptor neurons promote IgE class switch in B cells. *JCI Insight* 6,

795   e148510 **(2021).**

11. Perner, C. *et al.* Substance P Release by Sensory Neurons Triggers Dendritic Cell

Migration and Initiates the Type-2 Immune Response to Allergens. *Immunity* 53, 1063-

1077.e7 (2020).

12. Oetjen, L. K. *et al.* Sensory Neurons Co-opt Classical Immune Signaling Pathways to

Mediate Chronic Itch. *Cell* 171, 217-228.e13 (2017).

13. Jendzjowsky, N. G. *et al.* PKCε stimulation of TRPV1 orchestrates carotid body

responses to asthmakines. *J Physiol* 599, 1335–1354 (2021).

14. Chen, Y., Yang, C. & Wang, Z. J. Proteinase-activated receptor 2 sensitizes transient

receptor potential vanilloid 1, transient receptor potential vanilloid 4, and transient

receptor potential ankyrin 1 in paclitaxel-induced neuropathic pain. *Neuroscience*

193, 440–451 (2011).

15. Serhan, N. *et al.* House dust mites activate nociceptor-mast cell clusters to drive type

2 skin inflammation. *Nat. Immunol.* 20, 1435–1443 (2019).

16. Vellani, V. *et al.* Protease activated receptors 1 and 4 sensitize TRPV1 in nociceptive

neurones. *Mol. Pain* 6, 61 (2010).

17. Voisin, T. *et al.* The CysLT2R receptor mediates leukotriene C4-driven acute and

chronic itch. *Proc Natl Acad Sci U S A* 118, e2022087118 (2021).

18. Rijnierse, A. *et al.* Immunoglobulin-free light chains mediate antigen-specific

responses of murine dorsal root ganglion neurons. *J. Neuroimmunol.* 208, 80–86

(2009).

19. Cui, H. *et al.* Neuronal FcεRIα directly mediates ocular itch via IgE-immune complex

in a mouse model of allergic conjunctivitis. *Journal of Neuroinflammation* 19, 55

(2022).

20. Andoh, T. & Kuraishi, Y. Direct action of immunoglobulin G on primary sensory

neurons through Fc gamma receptor I. *FASEB J.* 18, 182–184 (2004).

21. Udit, S., Blake, K. & Chiu, I. M. Somatosensory and autonomic neuronal regulation of

the immune response. *Nat Rev Neurosci* 23, **157–171 (2022).**

22. Donnelly, C. R., Chen, O. & Ji, R.-R. How Do Sensory Neurons Sense Danger Signals?

Trends Neurosci 43, **822–838 (2020).**

23. Crosson, T. *et al.* Profiling of how nociceptor neurons detect danger – new and old

foes. *Journal of Internal Medicine* 286, **268–289 (2019).**

24. Veiga-Fernandes, H. & Freitas, A. A. The S(c)ensory Immune System Theory. *Trends*

Immunol 38, **777–788 (2017).**

25. Yissachar, N. *et al.* An Intestinal Organ Culture System Uncovers a Role for the

Nervous System in Microbe-Immune Crosstalk. *Cell* 168, **1135-1148.e12 (2017).**

26. Lai, N. Y. *et al.* Gut-Innervating Nociceptor Neurons Regulate Peyer's Patch Microfold

Cells and SFB Levels to Mediate Salmonella Host Defense. *Cell* 180, **33-49.e22 (2020).**

27. Blake, K. J. *et al.* Staphylococcus aureus produces pain through pore-forming toxins

and neuronal TRPV1 that is silenced by QX-314. *Nat Commun* 9, **(2018).**

28. Cardoso, V. *et al.* Neuronal regulation of type 2 innate lymphoid cells via neuromedin

U. *Nature* 549, **277–281 (2017).**

29. Foster, S. L., Seehus, C. R., Woolf, C. J. & Talbot, S. Sense and Immunity: Context-

Dependent Neuro-Immune Interplay. *Front Immunol* 8, **(2017).**

30. Wilson, R. *et al.* Protection against Streptococcus pneumoniae lung infection after

nasopharyngeal colonization requires both humoral and cellular immune responses.

Mucosal Immunology 8, **627–639 (2015).**

31. Thevaranjan, N. *et al.* Streptococcus pneumoniae Colonization Disrupts the Microbial

Community within the Upper Respiratory Tract of Aging Mice. *Infect Immun* 84, **906–**

916 (2016).

32. Xiong, H., Dolpady, J., Wabl, M., Curotto de Lafaille, M. A. & Lafaille, J. J. Sequential

class switching is required for the generation of high affinity IgE antibodies. *Journal*

of Experimental Medicine 209, **353–364 (2012).**

33. Geha, R. S., Jabara, H. H. & Brodeur, S. R. The regulation of immunoglobulin E class-

switch recombination. *Nat Rev Immunol* 3, **721–732 (2003).**

34. Pascual, D. W., Xu-Amano, J. C., Kiyono, H., McGhee, J. R. & Bost, K. L. Substance P

acts directly upon cloned B lymphoma cells to enhance IgA and IgM production. *J*

Immunol 146, **2130–2136 (1991).**

35. Tan, E. M. S. *et al.* Neuropeptide Y receptor 1 is expressed by B and T lymphocytes

and mast cells in infantile haemangiomas. *Acta Paediatr* 106, **292–297 (2017).**

36. McGillis, J. P., Humphreys, S., Rangnekar, V. & Ciallella, J. Modulation of B

lymphocyte differentiation by calcitonin gene-related peptide (CGRP). I.

Characterization of high-affinity CGRP receptors on murine 70Z/3 cells. *Cell Immunol*

150, **391–404 (1993).**

37. Schratzberger, P. *et al.* Similar involvement of VIP receptor type I and type II in

lymphocyte chemotaxis. *J Neuroimmunol* 87, **73–81 (1998).**

38. Kimata, H., Yoshida, A., Ishioka, C. & Mikawa, H. Differential effect of vasoactive

intestinal peptide, somatostatin, and substance P on human IgE and IgG subclass

production. *Cell Immunol* 144, **429–442 (1992).**

39. Kimata, H. & Fujimoto, M. Vasoactive intestinal peptide specifically induces human

IgA1 and IgA2 production. *Eur J Immunol* 24, **2262–2265 (1994).**

40. Zeng, H.-T. *et al.* Vasoactive intestinal peptide alleviates food allergy via restoring

regulatory B cell functions. *Immunobiology* 224, **804–810 (2019).**

41. Courties, G. *et al.* Glucocorticoids Regulate Bone Marrow B Lymphopoiesis After

Stroke. *Circ Res* 124, **1372–1385 (2019).**

42. Bedoui, S. *et al.* Differential effects of neuropeptide Y (NPY) on leukocyte subsets in

the blood: mobilization of B-1-like B-lymphocytes and activated monocytes. *J*

Neuroimmunol 117, **125–132 (2001).**

43. Mora, J. R. & von Andrian, U. H. Differentiation and homing of IgA-secreting cells.

Mucosal Immunol 1, **96–109 (2008).**

44. Zhang, X. *et al.* Brain control of humoral immune responses amenable to behavioural

modulation. *Nature* 581, **204–208 (2020).**

45. Julius, D. & Basbaum, A. I. Molecular mechanisms of nociception. *Nature* 413, **203–**

210 (2001).

46. Mishra, S. K. & Hoon, M. A. Ablation of TrpV1 neurons reveals their selective role in

thermal pain sensation. *Mol Cell Neurosci* 43, **157–163 (2010).**

47. Kissin, I. & Szallasi, A. Therapeutic targeting of TRPV1 by resiniferatoxin, from

preclinical studies to clinical trials. *Curr Top Med Chem* 11, **2159–2170 (2011).**

48. Usoskin, D. *et al.* Unbiased classification of sensory neuron types by large-scale

single-cell RNA sequencing. *Nat. Neurosci.* 18, **145–153 (2015).**

49. Kim, S.-H. *et al.* Mapping of the Sensory Innervation of the Mouse Lung by Specific

Vagal and Dorsal Root Ganglion Neuronal Subsets. *eNeuro* 9, **(2022).**

50. Su, Y. *et al.* Identification of lung innervating sensory neurons and their target

specificity. *Am J Physiol Lung Cell Mol Physiol* 322, **L50–L63 (2022).**

51. Borsa, N., Di Pasquale, M. & Restrepo, M. I. Animal Models of Pneumococcal

pneumonia. *International Journal of Molecular Sciences* 20, **4220 (2019).**

52. Gogos, A. & Federle, M. J. Modeling Streptococcus pyogenes Pharyngeal

Colonization in the Mouse. *Frontiers in Cellular and Infection Microbiology* 9, **(2019).**

53. Pogorzala, L. A., Mishra, S. K. & Hoon, M. A. The Cellular Code for Mammalian

Thermosensation. *J Neurosci* 33, **5533–5541 (2013).**

54. Morgan, D. & Tergaonkar, V. Unraveling B cell trajectories at single cell resolution.

Trends in Immunology 43, **210–229 (2022).**

55. Kupari, J., Häring, M., Agirre, E., Castelo-Branco, G. & Ernfors, P. An Atlas of Vagal

Sensory Neurons and Their Molecular Specialization. *Cell Reports* 27, **2508-2523.e4**

(2019).

56. Neil, G. A., Blum, A. & Weinstock, J. V. Substance P but not vasoactive intestinal

peptide modulates immunoglobulin secretion in murine schistosomiasis. *Cell*

Immunol 135, **394–401 (1991).**

57. Pascual, D. W., Xu-Amano, J. C., Kiyono, H., McGhee, J. R. & Bost, K. L. Substance P

acts directly upon cloned B lymphoma cells to enhance IgA and IgM production. *J*

Immunol 146, **2130–2136 (1991).**

58. Bost, K. L. & Pascual, D. W. Substance P: a late-acting B lymphocyte differentiation

cofactor. *Am J Physiol* 262, **C537-545 (1992).**

59. Schratzberger, P. *et al.* Similar involvement of VIP receptor type I and type II in

lymphocyte chemotaxis. *J Neuroimmunol* 87, **73–81 (1998).**

60. Kimata, H. Vasoactive intestinal peptide differentially modulates human

immunoglobulin production. *Adv Neuroimmunol* 6, **107–115 (1996).**

61. Bedoui, S. *et al.* NPY modulates epinephrine-induced leukocytosis via Y-1 and Y-5

receptor activation in vivo: sympathetic co-transmission during leukocyte

mobilization. *J Neuroimmunol* 132, **25–33 (2002).**

62. Rubtsov, A. V. *et al.* Toll-like receptor 7 (TLR7)-driven accumulation of a novel CD11c[+]

B-cell population is important for the development of autoimmunity. *Blood* 118, **1305–**

1315 (2011).

63. Hao, Y., O'Neill, P., Naradikian, M. S., Scholz, J. L. & Cancro, M. P. A B-cell subset

uniquely responsive to innate stimuli accumulates in aged mice. *Blood* 118, **1294–**

1304 (2011).

64. Gao, X. & Cockburn, I. A. The development and function of CD11c+ atypical B cells -

insights from single cell analysis. *Front Immunol* 13, 979060 (2022).

65. O'Dorisio, M. S., O'Dorisio, T. M., Cataland, S. & Balcerzak, S. P. Vasoactive intestinal

polypeptide as a biochemical marker for polymorphonuclear leukocytes. *J Lab Clin*

Med 96, 666–672 (1980).

66. Delgado, M. VIP: a very important peptide in T helper differentiation. *Trends in*

Immunology 24, 221–224 (2003).

67. Smalley, S. G. R., Barrow, P. A. & Foster, N. Immunomodulation of innate immune

responses by vasoactive intestinal peptide (VIP): its therapeutic potential in

inflammatory disease. *Clin Exp Immunol* 157, 225–234 (2009).

68. Kuo, C. S. *et al.* Neuroendocrinology of the lung revealed by single-cell RNA

sequencing. *eLife* 11, e78216 (2022).

69. Doherty, T. A. *et al.* STAT6 regulates natural helper cell proliferation during lung

inflammation initiated by Alternaria. *Am J Physiol Lung Cell Mol Physiol* 303, L577-

588 (2012).

70. Gour, N. & Dong, X. The MRGPR family of receptors in immunity. *Immunity* 57, 28–39

(2024).

71. Huang, S. *et al.* Lymph nodes are innervated by a unique population of sensory

neurons with immunomodulatory potential. *Cell* 184, 441-459.e25 (2021).

72. Zhang, X. *et al.* Brain control of humoral immune responses amenable to behavioural

modulation. *Nature* 581, 204–208 (2020).

73. Mina-Osorio, P. *et al.* Neural signaling in the spleen controls B-cell responses to

blood-borne antigen. *Mol Med* 18, 618–627 (2012).

74. Häring, M. *et al.* Neuronal atlas of the dorsal horn defines its architecture and links

sensory input to transcriptional cell types. *Nat Neurosci* 21, 869–880 (2018).

75. Fukao, S., Haniuda, K., Tamaki, H. & Kitamura, D. Protein kinase Cδ is essential for

the IgG response against T-cell-independent type 2 antigens and commensal

bacteria. *eLife* 10, e72116 (2021).

76. Saijo, K., Mecklenbräuker, I., Schmedt, C. & Tarakhovsky, A. B Cell Immunity

Regulated by the Protein Kinase C Family. *Annals of the New York Academy of*

Sciences 987, 125–134 (2003).

77. Wu, M. *et al.* Innervation of nociceptor neurons in the spleen promotes germinal

center responses and humoral immunity. *Cell* 187, 2935-2951.e19 (2024).

78. Undem, B. J. & Taylor-Clark, T. Mechanisms underlying the neuronal-based

symptoms of allergy. *J. Allergy Clin. Immunol.* 133, 1521–1534 (2014).

79. Ollerenshaw, S. L., Jarvis, D., Sullivan, C. E. & Woolcock, A. J. Substance P

immunoreactive nerves in airways from asthmatics and nonasthmatics. *The*

European respiratory journal 4, 673–82 (1991).

80. Sponchiado, M. *et al.* Overexpression of Substance P in pig airways increases

MUC5AC through an NF-kβ pathway. *Physiol Rep* 9, e14749 (2021).

81. Talbot, S. *et al.* Vagal sensory neurons drive mucous cell metaplasia. *Journal of*

Allergy and Clinical Immunology 0, (2020).

82. Choi, J. Y. *et al.* Substance P stimulates human airway submucosal gland secretion

mainly via a CFTR-dependent process. *J Clin Invest* 119, 1189–1200 (2009).

83. Hanes, W. M. *et al.* Neuronal Circuits Modulate Antigen Flow Through Lymph Nodes.

Bioelectron Med 3, 18–28 (2016).

84. Etesami, N. S. *et al.* B cells in the pneumococcus-infected lung are heterogeneous

and require CD4+ T cell help including CD40L to become resident memory B cells.

Front Immunol 15, 1382638 (2024).

85. Millet, N., Solis, N. V. & Swidergall, M. Mucosal IgA Prevents Commensal Candida

albicans Dysbiosis in the Oral Cavity. *Front Immunol* 11, 555363 (2020).

86. Allie, S. R. *et al.* The establishment of resident memory B cells in the lung requires

local antigen encounter. *Nat Immunol* 20, 97–108 (2019).

87. Allie, S. R. & Randall, T. D. Resident Memory B Cells. *Viral Immunol* 33, 282–293

(2020).

88. Lam, J. H. & Baumgarth, N. The Multifaceted B Cell Response to Influenza Virus. *J

Immunol* 202, 351–359 (2019).

89. Rothaeusler, K. & Baumgarth, N. B-cell fate decisions following influenza virus

infection. *Eur J Immunol* 40, 366–377 (2010).

90. Cavagnero, K. J. *et al.* Unconventional ST2- and CD127-negative lung ILC2

populations are induced by the fungal allergen Alternaria alternata. *J Allergy Clin

Immunol* 144, 1432-1435.e9 (2019).

91. Tränkner, D., Hahne, N., Sugino, K., Hoon, M. A. & Zuker, C. Population of sensory

neurons essential for asthmatic hyperreactivity of inflamed airways. *PNAS* 111,

11515–11520 (2014).

92. Ghasemlou, N., Chiu, I. M., Julien, J.-P. & Woolf, C. J. CD11b+Ly6G- myeloid cells

mediate mechanical inflammatory pain hypersensitivity. *Proc Natl Acad Sci U S A* 112,

E6808-6817 (2015).

93. Swidergall, M. *et al.* EphA2 Is a Neutrophil Receptor for Candida albicans that

Stimulates Antifungal Activity during Oropharyngeal Infection. *Cell Rep* 28, 423-

433.e5 (2019).

94. Chiu, H. *et al.* Reduced eIF4E function impairs B-cell leukemia without altering normal

B-lymphocyte function. *iScience* 24, (2021).

Figure 1.

Figure 2.

Figure 3.

Figure 4.

Figure 5.

Figure 6.

b B Memory
B220+/CD19+/CD44+/
CD38+ cells

p = 0.0003
p = 0.3396
p < 0.0001
p = 0.0344

Cell (#)

Veh (PBS) | Veh (SubP) | RTX (PBS) | RTX (SubP)

c B Resident Memory
B220+/CD19+/CD44+/
CD38+/IgD-/CD62L- cells

p < 0.0001
p = 0.2075
p < 0.0001
p = 0.0018

Veh (PBS) | Veh (SubP) | RTX (PBS) | RTX (SubP)

d Isotped switched
B220+/CD19+-
IgM-/IgD- cells

p = 0.0696
p = 0.9949
p = 0.0004
p = 0.0975

Veh (PBS) | Veh (SubP) | RTX (PBS) | RTX (SubP)

e Plasma
CD138+ cells

p = 0.8407
p = 0.9793
p = 0.0075
p = 0.0435

Veh (PBS) | Veh (SubP) | RTX (PBS) | RTX (SubP)

f Plasmablast
B220^{low}/CD19+/CD138+ cells

p = 0.9521
p = 0.9611
p = 0.0230
p = 0.0697

Veh (PBS) | Veh (SubP) | RTX (PBS) | RTX (SubP)

g B Memory
B220+/CD19+/CD44+/
CD38+ Cells

p = 0.9416

Cell (#)

WT | Tac1-/-

h B Resident Memory
B220+/CD19+/CD44+/
CD38+/IgD-/CD62L- cells

p = 0.0130

WT | Tac1-/-

i Isotype switched
B220+/CD19+/
IgM-/IgD- cells

p = 0.0121

WT | Tac1-/-

j Plasma
CD138+ cells

p = 0.0001

WT | Tac1-/-

k Plasmablast
B220low/CD19+/
CD138+ cells

p = 0.0032

WT | Tac1-/-

l IgE

p < 0.0001
p < 0.0001

pg/g tissue

Veh (PBS) | Veh (SubP) | RTX (Veh) | RTX (SubP)

m IgE

p < 0.0001

WT | Tac1-/-