The recombination landscape of introgression in yeast

Enrique J. Schwarzkopf[1], Nathan Brandt[1], Caiti Smukowski Heil*[1]

Affiliation:

[1] Department of Biological Sciences, North Carolina State University, Raleigh, NC

*Corresponding author: cheil@ncsu.edu

Abstract

Meiotic recombination is an evolutionary force that acts by breaking up genomic linkage,

increasing the efficacy of selection. Recombination is initiated with a double-strand break which

is resolved via a crossover, which involves the reciprocal exchange of genetic material between

homologous chromosomes, or a non-crossover, which results in small tracts of non-reciprocal

exchange of genetic material. Crossover and non-crossover rates vary between species,

populations, individuals, and across the genome. In recent years, recombination rate has been

associated with the distribution of ancestry derived from past interspecific hybridization

(introgression) in a variety of species. We explore this interaction of recombination and

introgression by sequencing spores and detecting crossovers and non-crossovers from two

crosses of the yeast *Saccharomyces uvarum*. One cross is between strains which each contain

introgression from their sister species, *S. eubayanus*, while the other cross has no introgression

present. We find that the recombination landscape is significantly different between *S. uvarum*

crosses, and that some of these differences can be explained by the presence of introgression

in one cross. Crossovers are reduced and non-crossovers are increased in heterozygous

introgression compared to syntenic regions in the cross without introgression. This translates to

reduced allele shuffling within introgressed regions, and an overall reduction of shuffling on

most chromosomes with introgression compared to the syntenic regions and chromosomes

without introgression. Our results suggest that hybridization can significantly influence the

recombination landscape, and that the reduction in allele shuffling contributes to the initial

purging of introgression in the generations following a hybridization event.

Keywords: introgression, hybridization, recombination, crossover, non-crossover, yeast,

Saccharomyces

Introduction

Recombination is the exchange of genetic material between homologous chromosomes during

meiosis and is a staple of eukaryotic sexual reproduction. While the processes involved in

recombination are largely conserved (Arter & Keeney, 2023), recombination rates vary between

sexes, populations, and species (Smukowski & Noor, 2011; Stapley et al., 2017).

Recombination rates also vary along the genome, with conflicting patterns of enriched or

depleted recombination in promoter regions and punctate or dispersed recombination

depending on the species (Auton et al., 2013; Rockman & Kruglyak, 2009; Singhal et al., 2015;

Smukowski Heil et al., 2015). These patterns in recombination can affect pairing of alleles after

meiosis–in other words, the shuffling of alleles–in a population. Much of the evolutionary

advantage of recombination is understood to originate from its role in shuffling alleles, which

increases the number of different allele combinations segregating in a population. The increase

in allele combinations can reduce selection interference–the effect that genetically linked sites

have on the evolutionary fate of either beneficial or deleterious alleles (Felsenstein, 1974; Hill &

Robertson, 1966; McDonald et al., 2016; McGaugh et al., 2012).

How much allele decoupling is produced by recombination will depend on the type of

recombination event. Each recombination event begins with the severing of both strands of a

sister chromatid of one of the homologous chromosomes in what is referred to as a meiotic

double-strand break (DSB) (Keeney, 2001). A meiotic DSB is repaired as a crossover (CO),

which results in the reciprocal exchange of genetic information between homologous

chromosomes, or as a non-crossover (NCO) gene conversion –where a small segment

(typically 100-2000 bp) of a chromosome is replaced by a copy of its homolog (Chovnick et al.,

1971; Hilliker et al., 1994; Jeffreys & May, 2004; Judd & Petes, 1988). COs generally produce

more allele shuffling, and therefore degrade linkage faster than NCOs. However, NCOs can

occur in regions where COs are typically suppressed, like centromeres and inversions (Korunes

& Noor, 2019; Mancera et al., 2008; Miller et al., 2016; Schaeffer & Anderson, 2005; Shi et al.,

2010; Talbert & Henikoff, 2010; Wijnker et al., 2013). NCOs are also crucial to reducing linkage

within coding regions and, unlike COs, result in 3:1 allele ratio in the meiotic product at

heterozygous sites, potentially changing allele frequencies (Korunes & Noor, 2017).

Variation in the number and distribution of COs and NCOs, and their respective associated

effects on linkage, have important implications for molecular evolution. Recombination has long

been appreciated to play a role in the distribution of various genomic features including

nucleotide diversity. Nucleotide diversity has a positive correlation with recombination rate in a

number of species, interpreted to result from selective sweeps and background selection

removing genetic variation in regions of low recombination (Begun & Aquadro, 1992;

Charlesworth et al., 1993; Smith & Haigh, 1974). Similarly, recombination breaking up genetic

associations is particularly notable in the context of interspecific hybridization. In first-generation

(F_1) hybrids, the hybrid genome is heterozygous for each parental species. If the hybrids then

back-cross to one of the parental species, recombination will produce genomes that are a

mosaic of genetic information from the two species with a minor contribution from the species

that was not backcrossed to (introgression) (Aguillon et al., 2022). When each population has

evolved alleles that are deleterious when present in the background of the other population (the

Dobzhansky-Muller hybrid incompatibility model) we expect introgressed regions with low rates

of recombination to be quickly purged from the population, as the accumulation of incompatible

alleles incurs a steep fitness cost. In contrast, when introgressed regions have high

recombination rates, the break up of genetic associations will reduce selective interference

between the incompatible alleles and their surrounding haplotypes, allowing for neutral and

beneficial alleles brought in with the introgression to escape the fate of neighboring

incompatibilities (Barton & Bengtsson, 1986; Butlin, 2005; Moran et al., 2021; Nachman, M. &

Payseur, B., 2012; Schumer et al., 2018; Veller et al., 2023). This theory is supported

empirically through enrichment of introgressed segments in regions of higher recombination in a

number of organisms including Mimulus, maize, butterflies, swordtail fish, stickleback, and

humans (Brandvain et al., 2014; Calfee et al., 2021; Edelman et al., 2019; Martin et al., 2019;

Ravinet et al., 2018; Schumer et al., 2018).

This positive correlation between introgressed ancestry and recombination is emerging as a

nearly ubiquitous pattern (though see (Dagilis & Matute, 2023; Duranton & Pool, 2022; Pool,

2015)), however, it is unclear how these observations relate to the known effect of sequence

divergence on DSB repair. Introgression, particularly between highly diverged species, can have

low sequence similarity with the genomic region it is replacing. A DSB in a region of low

sequence similarity will recruit mismatch repair proteins, which ensure COs are occurring

between homologous chromosomes and at equivalent positions to prevent ectopic

recombination (Harfe & Jinks-Robertson, 2000; Hunter et al., 1996). Mismatch repair proteins

typically reduce the frequency of CO events as sequence divergence increases (Borts & Haber,

1987; Chen & Jinks-Robertson, 1999; Cooper et al., 2021; Martini et al., 2011). Given that

heterozygous introgression will have divergent sequences, we expect a decrease in COs, and

possibly an increase in NCOs as DSBs fail to be repaired as COs in heterozygous introgression.

To help us understand this interaction of introgression and recombination, and identify patterns

in CO and NCO in closely related populations, we utilized the budding yeast *Saccharomyces*.

Yeasts provide an excellent opportunity to study DSB repair, as we can readily isolate and

collect all four meiotic products of a given meiosis and detect both CO and more elusive NCO

events (Figure 1A) (Brion et al., 2017; Gerton et al., 2000; Liu et al., 2018, 2019; Mancera et al.,

2008). Recombination rates vary between strains of *S. cerevisiae* (Cubillos et al., 2011; Raffoux

et al., 2018) and between *S. cerevisiae* and its sister species *S. paradoxus* (Liu et al., 2019;

Tsai et al., 2010). Strains of different *Saccharomyces* species have often hybridized with other

species and carry introgressed DNA from these events (Albertin et al., 2018; Almeida et al.,

2014; Bendixsen et al., 2022; D'Angiolo et al., 2020; Langdon et al., 2019; Stelkens &

Bendixsen, 2022; Tellini et al., 2023).

Figure 1: (A) Visual representation of how our crosses were conducted. Haploid yeast from two

genetically distinct parental strains are mated to produce a heterozygous diploid. Meiosis is

induced and the resulting meiotic products (tetrads) are manually dissected, and each haploid

meiotic product is grown mitotically to obtain enough material for DNA extraction and whole

genome sequencing. We then call SNPs on the resulting sequences and retain loci with fixed

differences between parents. These loci are then coded as 1 or 0 depending on the parent of

origin and the CrossOver software detects COs and NCOs. (B) A schematic of how

introgression likely arose in the strains sampled from fermentation environments. These

introgressions are likely due to S. eubayanus *hybridizing with* S. uvarum *at some point in the*

past, resulting in F1 hybrids that then potentially crossed with other S. uvarum *individuals for*

some number of generations. Eventually, the S. eubayanus *ancestry was degraded in the*

population of S. uvarum *until the introgressions we observe today remained, potentially*

segregating in the population. A similar process likely happened in each of the parental strains

we utilized, but with different introgressions remaining in each strain. We crossed haploid

individuals from two parental strains, resulting in a diploid that is heterozygous for each

introgression.

In this study, we look at patterns of recombination and introgression by crossing two pairs of

Holarctic *Saccharomyces uvarum* strains. One pair of strains was isolated from natural

environments in North America and the other pair was isolated from European fermentation

environments (Almeida et al., 2014). The *S. uvarum* strains isolated from European

fermentation environments each carry introgression from their sister species, *Saccharomyces*

eubayanus, which is approximately 6% divergent from *S. uvarum* (Almeida et al., 2014;

Langdon et al., 2020; Nespolo et al., 2020). The diploid F_1 genome of these strains is

heterozygous for nine different introgressions which make up approximately 10% of the genome

(Figure 1B). The strains from the North American cross do not carry *S. eubayanus*

introgression, thus allowing us to assess the impact of introgression on the recombination

landscape. We obtained whole genome sequencing data from individual meiotic events from the

first offspring generation of each cross and used this data to detect CO and NCO events along

the genome (Figure 1A). From these maps, we aim to understand (i) how patterns of CO and

NCO differ between closely related strains, (ii) how regions of introgression differ in their CO

and NCO patterns, and (iii) how these different patterns affect shuffling of alleles locally and at

the chromosome level. Understanding these objectives will provide us novel insights into how

introgression impacts the recombination landscape.

Results

The recombination landscape differs dramatically between closely related crosses

We isolated and sequenced products of 48 meioses (192 haploid spores) for two crosses of *S.*

uvarum, a cross between strains isolated from North America (natural cross) and a cross

between strains isolated from Europe (fermentation cross). We detected COs and NCOs across

the 16 nuclear chromosomes of *S. uvarum*. Genomewide, we found significantly more COs on

average in the natural cross (82.54 COs/meiosis, SE 1.5; 0.72 cM/kb) than in the fermentation

cross (63.66 COs/meiosis, SE 1.9; 0.55 cM/kb; Wilcoxon rank sum test, p-value<<0.001) and

significantly fewer NCOs in the natural cross (25.48 NCOs/meiosis, SE 0.99) than the

fermentation cross (34.85 NCOs/meiosis, SE 8.92; Wilcoxon rank sum test, p-value=0.001448).

The number of COs per meiosis in the natural and fermentation crosses are slightly higher than

those of particular strains of *S. paradoxus* (54.8) and *S. cerevisiae* (76.5) respectively (Liu et al.,

2019). Additionally, the average number of NCOs in the natural cross is lower than those of *S.*

cerevisiae (46.4) and *S. paradoxus* (26.9), while the average number of NCOs in the

fermentation cross is in between these values (Figure 2A; Tables S4 & S5; Liu et al., 2019). It's

important to note that there is known strain variability in CO and NCO counts per meiosis for *S.*

cerevisiae (CO: 90.5, 76.3, 73; NCO: 46.6, 46.4, 27; Mancera et al., 2008; Liu et al., 2019;

Martini et al., 2011), similar to differences between strains of *S. uvarum* reported here.

To further explore the differences in recombination landscapes between our crosses, we split

the genome into 20kb, non-overlapping windows, and obtained CO, NCO, and marker counts

for each region. We downsampled markers in the introgression of the fermentation cross and

found a significant effect of marker density on NCO count and tract length for most of the

regions (Figures S2-10, Table S10). Because of how correlated marker density, NCO count,

and introgression are, disentangling their relationships is complicated. For this reason, we

applied an especially stringent correction for marker density–we used a previously published

simulation based method for correction (Liu et al., 2019; Wijnker et al., 2013) and took the

average NCO tract length (550bp) from introgression, where our marker density is highest, as

the correction factor. For each window across the genome, we established our expected

probability of detecting an NCO of that length and divided our observed NCO count by our

probability of detecting an NCO event (see Methods). After the correction we found modest–but

significant–genomewide correlation between our crosses for both COs (Spearman's correlation:

0.27; p<0.0001) and NCOs (Spearman's correlation: 0.13; p=0.0018).

Figure 2: (A) Barplot depicting the number of COs and NCOs detected per meiosis in S. uvarum

crosses (green: natural cross; pink: fermentation cross). The error bars represent the standard

error around the mean. NCO counts are corrected for marker resolution.The counts for S.

paradoxus *and* S. cerevisiae *are represented by arrows and taken from Liu et al. (2019). (B)*

Mean CO/kb and (C) NCO/kb by cross and introgression (0 denotes intervals without

introgression; 1 denotes introgression present in the fermentation cross. While the natural cross

does not contain introgression, the region where introgression is present in the fermentation

cross was compared to its syntenic region in the natural cross). NCO counts are corrected for

marker resolution. Error bars represent the standard error around the mean. (D) S. uvarum

chromosomes containing introgressions split into 20kb, non-overlapping windows. CO, NCO,

and SNP counts are reported for both crosses (fermentation and natural). Shaded regions

denote introgressed regions. CO counts are smoothed when the true location of the CO split

could be in one of multiple windows. NCO counts are corrected for marker resolution.

We hypothesized that the low correlations between crosses might be impacted by the presence

of heterozygous introgression from *S. eubayanus* in the fermentation cross. To explore this

possibility, we separated the 20kb windows into introgressed and non-introgressed windows

(based on whether they overlapped with an introgressed region). We will refer to introgression in

the fermentation cross as "introgression" and use the term "introgressed region" to refer

generally to the syntenic region, regardless of which cross we are focusing on. We calculated

Spearman's correlations of COs, NCOs, and marker counts between the two crosses for each

chromosome. We find positive correlation coefficients between CO counts between the

fermentation and natural crosses for all chromosomes when looking at non-introgressed

regions–though only chromosome 11 was significant (Table S6). For introgressed regions, we

found no significant CO correlations between crosses and no consistency in the direction of the

correlations, which is consistent with the hypothesis that CO landscapes are changed in

introgression (Table S7). There were no significant correlations among NCOs. This was likely

affected by the fact that markers used to detect NCOs are differently distributed between the

crosses, and the small size of NCO tracts means that the marker resolution of a given region

will affect the reported location of NCOs. This is because NCOs are reported to be at the

midpoint of the markers that identify it; more dense markers will give a more precise location,

which may conflict with the location reported by less dense markers. We find that introgressions

tend to have lower CO counts and higher NCO counts in the fermentation cross when compared

to syntenic regions in the natural cross (Wilcoxon rank sum test; CO p-value<10^{-5}, NCO p-

value<10^5). The fermentation cross has fewer COs than the natural cross overall, but the

difference is greater in the introgressed regions (Figure 2B-C). Additionally, introgressed regions

in the natural cross show significantly more COs and NCOs (Wilcoxon rank sum test; CO p-

value=0.007, NCO p-value=0.006), indicating that these regions may be predisposed to having

more recombination events even in the absence of introgression.

To further explore and test possible explanations for the patterns of COs and NCOs in

introgressed regions, we constructed linear models for CO and NCO counts. Our model of CO

counts showed a significant positive effect of the interactions between natural cross and

introgression (whether a genomic window is from the natural cross and whether that window is

in an introgressed region), as well as a significant negative effect of introgression on the number

of COs, and a significant positive effect of GC content on CO count (Table 1). These results are

consistent with increased COs in GC-rich regions and reduced COs in introgressions. Our

model of NCO counts showed a similar positive effect of GC content on NCO counts, but

showed an opposite significant coefficient for introgression and no significant effect of the

interaction between introgression and cross–which was removed from the final model (Table 2).

This indicates that GC content still plays an important role in localizing NCOs, and supports our

findings that patterns of NCOs in introgressed regions are opposite to those of COs.

Furthermore, the lack of a significant effect of cross echoes our finding of increased NCOs in

introgressed regions in the natural cross.

Table 1: Coefficients of gaussian generalized linear model modeling CO counts per 20kb

window.

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	-8.9355	1.4713	-6.0732	1.70e-09
Introgression	-3.6343	0.6293	-5.7755	9.84e-09
GC	37.9733	3.7037	10.2528	< 2e-16
Introgression:crossnatural	5.2378	0.8595	6.0941	1.49e-09

259   *Table 2: Coefficients of gaussian generalized linear model modeling NCO counts per 20kb*

260   *window.*

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	-11.4763	4.7810	-2.4004	0.0165
Introgression	4.3490	1.4937	2.9115	0.0037
GC	45.2894	12.0353	3.7631	0.0002

Reduced diploid sequence similarity helps explain non-crossover repair of DSBs in introgression

One possible explanation for an increase of NCOs in introgression is that the reduced sequence

similarity is biasing DSBs in the region to be repaired as NCOs rather than COs. We would

therefore expect to see NCOs to be negatively correlated with sequence similarity. To evaluate

the relationship of sequence similarity to the CO and NCO landscapes in introgressions, we

measured diploid sequence similarity (the proportion of bases that are expected to match when

we sample one base from each of the two parental strains), NCO depth, and CO count in 101

bp sliding windows with 50 bp overlaps along each of the introgressions. Mismatch repair

proteins in *Saccharomyces* seem to suppress COs with very little mismatch in small regions

(~350 bp), which informed our window size (Chen & Jinks-Robertson, 1999). We counted the

number of NCO tracts that intersect with each window as a measurement of NCO depth, and

simply counted the CO events in a given window (Figure 3). We then ran Spearman's

correlations and a loess regression along each introgression and found a weak, but often

significant ($p<<0.001$) correlation between NCOs and sequence similarity in the introgression

(Table 3), suggesting that repair of double strand breaks is biased towards NCOs when

sequence similarity is low. From the loess regression, we can observe an increase in NCO as

sequence similarity reduces until about 0.9-0.8 sequence similarity, at which point NCOs level

out or reduce (Figure S1). However, this effect is very weak with respect to the NCO counts,

and at low levels of sequence similarity the uncertainty of the regression line is very large. This

is primarily driven by the number of windows with no NCOs .

Table 3: Spearman's correlations of NCOs to sequence similarity *in introgression in the*

fermentation cross. A cutoff p-value of 0.001 was selected by calculating the Bonferroni

corrected α=0.05 for nine comparisons (0.0011) and rounding down.

Chromosome	Start	End	Corr	p<0.001?
4	866500	983774	-0.1686	TRUE
6	1	65500	-0.0394	FALSE
7	1	53500	-0.2435	TRUE

9	158500	298500	-0.1460	TRUE
10	234500	288500	0.0073	FALSE
10	301500	428500	-0.1097	TRUE
13	26500	103500	-0.1982	TRUE
14	18500	586500	-0.2300	TRUE
15	367500	434500	-0.2328	TRUE

Figure 3: CO, NCO, and sequence similarity *in 101bp sliding windows with 50bp overlaps of*

fermentation cross introgression. CO counts are shown in blue, the depth of NCO tracts are

shown in orange, and the proportion of expected homologous bases between the two

fermentation strains is shown in black.

The low CO count in introgression leaves us unable to investigate effects of sequence similarity

on CO counts except on chromosomes 7 and 14, where we found significantly higher sequence

similarity around COs than around NCOs (Table 4). The introgressions on these two

chromosomes are unique in that they contain a highly homologous portion of sequence and

therefore contain enough COs for us to have power to detect differences between CO and NCO

neighborhoods (Figure 3).

Table 4: Welch two sample t-test results for differences in sequence similarity *between CO-*

adjacent regions and NCO-adjacent regions per introgression.

Introgression	CO mean sequence similarity	CO SE	NCO mean sequence similarity	NCO SE	p-value
Chromosome 4	0.9467	0.0148	0.9248	0.0029	0.2835
Chromosome 6	0.9663	0.0203	0.9784	0.0020	0.5594
Chromosome 7	1.0000	0.0000	0.9838	0.0018	<0.0001
Chromosome 9	0.9650	0.0102	0.9312	0.0037	0.0084
Chromosome 10 (1)	0.9671	0.0092	0.9476	0.0066	0.0419
Chromosome 10 (2)	0.9400	0.0074	0.9276	0.0040	0.1575
Chromosome 13	0.9700	0.0161	0.9304	0.0025	0.1292
Chromosome 14	0.9893	0.0026	0.9415	0.0018	<0.0001
Chromosome 15	0.9317	0.0093	0.9252	0.0031	0.5088

Introgression decreases allele shuffling locally and at the chromosome level

Because NCOs still play a small role in shuffling alleles along the chromosome, we were

interested in whether the increase in NCOs of the fermentation cross would supplement the lost

shuffling from the suppression of COs in the introgressions. To test this hypothesis, we used the

measure r, which accounts for the number and positioning of recombination events to estimate

the probability that a randomly chosen pair of loci shuffles their alleles in a gamete (Veller et al.,

2019). We calculated the average r per chromosome and for each introgressed region for each

of the two crosses. We observed high levels of shuffling at the chromosome level when

compared to humans. The intra-chromosomal component of r in humans is 0.0135 in females

and 0.0177 in males (Veller et al., 2019), while our measurements for chromosomes varied

between 0.216 and 0.426. We find that most chromosomes do not have a significantly different

amount of allele shuffling between the two crosses, even though the natural cross generally has

more COs (Table S8; Bonferroni-adjusted $\alpha = 0.00313$). However, of the six chromosomes with

significantly different r values, all of them showed more shuffling in the natural cross, and five of

the six (chromosomes 4, 9, 10, 14, and 15) contained introgressed regions (Figure 4). Lower r

is not observed when introgressions are small and near telomeres, while even a small

introgression near the center of the chromosome can lead to a large reduction in r (as is the

case for chromosome 15). Chromosome 12 was the only chromosome without an introgressed

region to have significantly different shuffling between crosses, and it also showed more

shuffling in the natural cross. While it is unclear what potential mechanism is mediating the

difference in shuffling on chromosome 12, we note that the rDNA locus on chromosome 12 is

known to differ dramatically in repeat content across strains of *S. cerevisiae* (22–227 copies)

(Sharma et al., 2022), and we speculate that differences in rDNA copy number between strains

in our crosses could impact shuffling. All of the introgressed regions showed significantly more

shuffling in the natural cross, indicating that the large increase of NCOs in the introgressions

does not make up for the loss of shuffling from the depletion of COs (Table S9; Bonferroni-

adjusted $\alpha = 0.00556$). This finding indicates that an introgression that is segregating in a

population will incur a shuffling cost in heterozygous individuals on top of any other evolutionary

effects the introgression may have.

Figure 4: Average r for each chromosome (A) and for each introgressed region (B). Asterisks

indicate a significant difference in chromosome r-bar between crosses. All introgressed regions

had a significant difference in r. Error bars indicate standard error around the mean.

Discussion

Our study is motivated by understanding recombination rate variation within a species and

uncovering potential genetic factors underlying this variation. To investigate this question, we

crossed two pairs of *S. uvarum* strains, one pair isolated from natural environments and one

pair from fermentation environments, and explored the distribution of CO and NCO events from

both crosses. We detected more COs and fewer NCOs in our natural cross when compared to

our fermentation cross, within a similar range of COs and NCOs per meiosis as previous studies

in *S. cerevisiae* and *S. paradoxus* (Liu et al., 2019; Mancera et al., 2008). Our findings

demonstrate significant differences in recombination rate between closely related strains,

adding to a body of literature across plants, animals, and fungi that recombination rate can

evolve rapidly (Bauer et al., 2013; Danguy des Déserts et al., 2021; Kong et al., 2014; McGaugh

et al., 2012; Petit et al., 2017; Raffoux et al., 2018; Samuk et al., 2019; Sandor et al., 2012;

Schreiber et al., 2022).

We hypothesized that these differences in the recombination landscape between our crosses

were in part influenced by introgression, given that heterozygous introgression creates

sequence divergence, and that COs in regions of heterozygosity are known to be curtailed in

yeast and other organisms (Chen & Jinks-Robertson, 1999; Cooper et al., 2021; Martini et al.,

2011). We therefore explored the relationship between introgressions and the differences in CO

and NCO counts between crosses. We modeled CO and NCO locations, correcting for GC

content (a well characterized driver of recombination events (Kiktev et al., 2018; Marsolier-

Kergoat & Yeramian, 2009)) and found that the distribution of COs and NCOs we observed

could be partly explained by introgressions. While we are limited in our interpretations by only

comparing two crosses (one cross with heterozygous introgression and one without

introgression), these results are in line with findings in inversions, where heterozygotes show

sharp decreases in COs, but the presence of NCOs in the inverted region (Crown et al., 2018;

Korunes & Noor, 2019). However, unlike heterozygous inversions where an increase in COs is

observed on freely recombining chromosomes (the inter-chromosomal effect), we do not see an

increase in COs on the borders flanking introgression or on chromosomes without introgression.

Relatedly, our results also differ from observations in Arabidopsis, in which COs are elevated in

regions with more heterozygosity at the cost of COs in adjacent homologous regions (Ziolkowski

et al., 2015; Ziolkowski & Henderson, 2017). These contrasting outcomes are likely in part due

to the difference between intraspecific levels of polymorphism in Arabidopsis studies compared

to the high level of inter-specific sequence divergence in our heterozygous introgression,

although it also appears that mismatch repair gene MSH2 has functionally diverged between

Arabidopsis and Saccharomyces (Blackwell et al., 2020; Dluzewska et al., 2023; Szymanska-

Lejman et al., 2023). Regardless, it is curious that we do not find evidence for compensation of

the depletion of COs due to introgression; the underlying mechanism responsible requires

further study.

The introgressions present in the fermentation cross strains provided a unique opportunity to

analyze NCO patterns, as the highly increased marker density lends the ability to detect NCOs

with smaller tract lengths. We find evidence that DSB repair is biased towards NCOs when

sequence similarity is low, with a weak but significant effect of increasing NCOs with decreasing

sequencing similarity to a peak at sequence similarity between 0.8-0.9. Our data are thus

aligned with previous work in *S. cerevisiae* and other systems, in which *MSH2* and *SGS1* act as

anti-recombinases and disassemble D-loops with mismatched sequences, resulting in NCOs

instead of COs (Borts & Haber, 1987; Chen & Jinks-Robertson, 1999; Martini et al., 2011;

Mazina et al., 2004; Myung et al., 2001). By downsampling markers in the introgressed regions,

we found an increase in the number of NCOs and a decrease in their tract length as we

increased our resolution (Figures S2-S10). Our expectation was to find a marker density at

which NCO counts and their tract lengths would no longer increase and decrease respectively.

However, we didn't find such a density, despite in some cases exploring marker densities that

included all the markers in the introgression. This leaves us with the uncertainty of whether

NCOs in the introgressions are more frequent and smaller than in other regions, or if we may be

underestimating NCO counts and overestimating their tract lengths in cases where marker

resolution is lower.

One likely effect of the CO reduction we observed in introgressions is a reduction in allele

shuffling at the regional and chromosomal level. While NCOs can increase local shuffling, they

likely have a much weaker effect on the likelihood of two random alleles being shuffled than

COs do. We find this is the case for our two crosses, where despite a large number of NCOs in

introgressions, the amount of shuffling (as measured by r) is significantly lower in the

fermentation cross. This loss of shuffling translates to frequently lower r in the fermentation

cross at the chromosome level for chromosomes containing introgressions. The exceptions

being small introgressions near the telomeres, which is consistent with the expectation that COs

near the center of chromosomes generate much more shuffling of alleles than terminal COs

(Veller et al., 2019). Our findings indicate that reducing COs, especially near the center of

chromosomes, has a cost to shuffling that is not compensated by the increase of NCOs that we

observe. If the benefit of recombination is its ability to generate new combinations of alleles,

then the loss of shuffling resulting from being heterozygous for divergent DNA sequences may

come at an additional cost beyond the possibilities of genetic incompatibilities between

hybridizing species. This cost is likely higher as divergence increases and as the length of

divergent sequences is greater, as is the case with early generation hybrids (Dagilis & Matute,

2023). Ultimately, if sequence divergence is too high, the resultant failure to recombine can

become a postzygotic reproductive barrier (Bozdag et al., 2021; Hunter et al., 1996; Rogers et

al., 2018).

The shuffling cost to introgression that we identify in our crosses may play an important role in

the fate of introgression in the generations following hybridization. When heterozygotes for an

introgression are formed, the reduction in shuffling inside the introgression will increase the

likelihood that the introgression is purged from the population. This is because it will likely be

inherited in its entirety and will carry the fitness cost of incompatibilities combined with a cost of

shuffling. This cost is incurred because the reduction of COs in the introgression will reduce

shuffling of alleles on either side of it and will vary in its intensity depending on the location and

size of the introgression. In generations immediately following hybridization, introgressions will

be much larger and are therefore expected to be more costly (although this likely depends on a

number of factors including time since divergence). These predictions are consistent with

modeling and empirical data on the purging of introgression in Drosophila and humans in the

first generations following hybridization (Veller et al., 2023).

As to longer term dynamics of recombination and selection, we predict that the excess NCOs

detected in heterozygous introgressions should begin to erode the divergence between the

sequences, increasing homology and slowly reducing the cost of the introgression. This

hypothesis posits that recombination can act to remove the larger, more deleterious regions of

an introgression quickly while whittling away slightly deleterious alleles that may be linked to any

beneficial regions of an introgression. While our current study doesn't capture longer term

patterns of recombination or the landscape of recombination in introgressions that would lead to

introgressions preferentially remaining in high-CO regions, it's notable that introgressed regions

coincide with high-recombination regions in the natural cross, indicating that these regions may

be primed to retain introgressions after a hybridization event–possibly due to the increased

shuffling allowing for a larger fraction of the introgression to remain (Veller et al., 2023).

Finally, we note that *Saccharomyces* typically reproduce asexually, with only infrequent sexual

cycles (Magwene et al., 2011; Ruderfer et al., 2006; Zeyl & Otto, 2007). When they do mate,

they often mate within a tetrad resulting in increased homozygosity. For example, each diploid

progenitor of the parents of our fermentation cross was homozygous for introgression across

the genome, meaning that recombination would neither break up nor aid in purging the

introgression in isolated populations of each parent. This suggests that the fate of introgressions

in this species is perhaps more loosely tied to recombination patterns than it would be in an

obligately sexually reproducing species.

Despite some limitations to interpretation, this study provides a unique view of the early

dynamics of hybridization and the role of recombination in the presence of introgression. By

focusing not only on the distribution of recombination events but on their specific role in shuffling

alleles, we can more closely connect the physical process of recombination to its role among

other evolutionary forces.

Methods

Strain and library construction

S. uvarum strains (UCD61-137, yHCT78 , GM14, and DBVPG7787) were obtained from the

Portuguese Yeast Culture Collection and from Chris Hittinger (Table S1) (Almeida et al., 2014).

All four *S. uvarum* strains had their *HO* locus replaced with a kanMX marker using a modified

version of the high-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method.

Briefly, the kanMX marker was amplified from plasmid pCSH2 with homology to genomic DNA

flanking the HO ORF with primers CSH239

(GGTGGAAAACCACGAAAAGTTAGAACTACGTTCAGGCAAAgacatggaggcccagaatac) and

CSH241 (GTGACCGTATTGGTACTTTTTTTGTTACCTGTTTTAGTAGcagtatagcgaccagcattc).

For each strain, overnight cultures were inoculated in 25 mL of YPD at an OD of ~ 0.0005 and

incubated at room temperature on a shaker for ~24 hours until the cultures reached an OD

between 0.6 and 0.9. Subsequently, 1 ug of the template DNA was transformed with a heat

shock temperature of 37°C for 45 minutes. The transformed cells were allowed to recover in

liquid YPD for 4 hours before being plated onto G418 selective plates and incubated at room

temperature for 2 days.

Single colonies were selected from the transformation plates, restreaked onto G418 plates and

allowed to grow at room temperature for 2 days. Single colonies from those plates were then

inoculated into 2 mL of YPD + G418 and incubated in a roller drum at room temperature

overnight. From those cultures, 250 uL was used to inoculate 2 mL of sporulation media (1%

potassium acetate, 0.1 % yeast extract, 0.05% dextrose) and incubated at room temperature for

3 to 5 days. Strains were confirmed to have the ho::KanMX via tetrad dissection on a Singer

SporPlay+ microscope (Singer Instruments). Plates with tetrads were incubated at room

temperature for 2 days and then replica plated to test for proper segregation of the kanMX

marker and mating type within individual tetrads.

Crosses between strains UCD61-137 and yHCT78 (natural cross), and between strains GM14

and DBVPG7787 (fermentation cross) were set up by micromanipulation of single MATa and

MATx cells using a Singer SporPlay+. The plates were incubated at room temperature for 2

days and then replica plated to mating type tester strains to test for potential diploids. Identified

diploids were then sporulated by growing a culture of the cross in 2 mL YPD + G418 at room

temperature overnight. From those cultures, 250 uL were used to inoculate 2 mL of sporulation

media and incubated at room temperature for 3 to 5 days. Sporulated cultures were dissected

on 3 YPD plates (24 tetrads per plate) using a Singer SporPlay+. Fifty of the fully viable tetrads

were selected and had all their spores inoculated into YPD (200 spores total) and incubated at

room temperature. The DNA was extracted from these cultures using a modified version of the

Hoffman-Winston DNA Prep (Hoffman & Winston, 1987). The DNA concentration was then

measured using SYBR green, and 150 ng of each sample's DNA was used to prepare a

sequencing library using an Illumina DNA Prep Kit, modified to use half the normal amounts of

reagents. Libraries were pooled and run on an Illumina NovaSeq 500 with 150bp paired end

reads.

Calling SNPs

We scored SNPs from parents and offspring using the *S. uvarum* reference genome (Scannell

et al., 2011) and custom scripts that invoked bwa (v0.7.17), samtools (v1.12), bcftools (v1.13),

picard tools (v2.25.6), and gatk (v4.2.0.0) (Danecek et al., 2021; Li & Durbin, 2009; McKenna et

al., 2010). The custom scripts are available in the github repository: *schwarzkopf/CO-NCO.* We

joint genotyped parents and offspring with default filters for gatk with the exception of the QUAL

filter, which was set as < 100 for parents and < 30 for offspring. We further filtered variants by

requiring they be fixed differences between the two parental strains. We kept a total of 24,574

markers for the natural cross and 74,619 markers for the fermentation cross. We utilized

LUMPY to identify structural variants in the parent strains that were greater than 5000 bp and

verified calls using the Integrative Genomics Viewer (Layer et al., 2014; Robinson et al., 2011).

We identified three amplifications in strain GM14 (one of the fermentation cross parents) that

were absent in other strains (Table S2).

Generating CO/NCO maps

We generated "seg" files by coding tetrad variants by their parental origin. These seg files were

the input for CrossOver (v6.3) from the ReCombine suite of programs, which we used to detect

COs and NCOs (Anderson et al., 2011). We then filtered to remove non-crossovers with fewer

than three associated markers and split the genome into 20kb windows. In each window we

counted crossovers, non-crossovers, and markers. We established regions of introgressions

through visual inspection of marker density in the fermentation cross (introgressions showed

more divergence between fermentation strains) and confirmed them using the findings of

Almeida et al. (2014). We found nine heterozygous introgressions on chromosomes 4, 6, 7, 9,

10, 10, 13, 14, and 15 respectively that we included in further analyses (Table S3). We excluded

two additional introgressions due to poor mapping (chromosome 13:0-17,000; chromosome 16:

642,000-648,000). We analyzed the effect of marker resolution on NCO detection by randomly

removing markers from each introgression and running CrossOver and its downstream counting

process as described above. For each introgression, we downsampled to multiple percentiles of

the distribution of non-introgressed region marker counts per kb (10^{th}: 0.05; 25^{th}: 0.75; 50^{th}:

1.95; 75^{th}: 2.95; 90^{th}: 4.45; 95^{th}: 7.87; 99^{th}: 14.01) and the median marker count per kb for

introgressed (3.225) and non-introgressed (1.35) regions. We repeated each downsampling

amount 20 times for each introgression and extracted average CO counts, NCO counts, and

NCO tract lengths. To account for the difference in number of markers in introgressed vs non-

introgressed windows and their effect on NCO detection, we applied a previously published

simulation-based method (Liu et al., 2019; Wijnker et al., 2013). We chose the average NCO

tract length of the fermentation cross's introgressed regions–550 bp–and for each window

randomly placed an NCO event of that length 10,000 times to establish our expected probability

of detecting an NCO of that length. We then divided our observed NCO count by our probability

of detecting an NCO event. Additionally, because COs that occurred in large regions devoid of

markers would be called in the middle of the empty windows, we split CO counts in regions with

multiple consecutive marker-free windows evenly between the empty windows. With these

corrected maps, we calculated spearman correlations between crosses using R (v4.1.0, R Core

Team 2021). Additionally, we modeled NCO and CO count as a function of introgression,

introgression by cross, and GC content using a gaussian generalized linear model in R (v4.1.0,

R Core Team 2021).

Sequence similarity

We calculated diploid sequence similarity between the two fermentation cross strains in 101bp

windows with 50bp overlaps. At each nucleotide position in the window, we counted fixed

differences as zero diploid sequence similarity, invariant sites between strains as full diploid

sequence similarity (1), and polymorphic sites in either or both strains as half diploid sequence

similarity (0.5). We then averaged these sequence similarity values across the window. This

measure represents the probability that both strains will have the same nucleotide base at a

given position. We used this measure of fine-scale sequence similarity to determine how

sequence similarity related to NCO counts in introgressed regions. For this, we used Loess

regressions and Spearman's correlations on each of the introgressed regions comparing

sequence similarity to NCO count, both implemented in R (v4.1.0, R Core Team 2021). We then

focused on each recombination event (CO or NCO) and compared the sequence similarity

100bp up and downstream of CO breakpoints and 100bp up and downstream of NCO tracts.

We then used Welch's two sample t-tests to compare CO and NCO sequence similarity in each

introgression.

560 r

We use r, a measure genetic shuffling defined in Veller et al. (2019) to measure how much

shuffling occurs in each chromosome for each cross. Our data provides parental origin for each

fixed difference between parental strains. We assume that all loci between pairs of markers that

come from the same parent are also from that parent. We also assume that when a pair of

successive markers come from different parents, the location of the change from one parental

origin to the other happens at the midpoint between our markers. With this in mind, we counted

the number of bases that come from one parent and divided by the chromosome size to obtain

the proportion of the chromosome that was inherited from said parent (p) and used the formula

from Veller et al. (2019): $r = 2p(1-p)$. We calculated r for each full chromosome and each

introgressed regions in every gamete from both crosses. We then averaged across gametes to

obtain average r values. We then compared average r between crosses in each chromosome

or introgressed region using Welch two sample t-tests and correcting for multiple tests using a

Bonferroni correction in R (v4.1.0, R Core Team 2021).

Data Access

Sequences for the parental strains can be found on NCBI SRA (SRR1119189, SRR1119180

SRR1119199, SRR1119200) (Almeida et al., 2014). Sequencing of the tetrads is deposited at

NCBI SRA under Project PRJNA1061120. Scripts are available in the github repository:

ejschwarzkopf/CO-NCO.

Acknowledgements

We are grateful to members of the Heil lab, Mohamed Noor, Nathan Layman, Mark Smithson,

and three anonymous reviewers for comments on this manuscript. We thank Chris Hittinger and

the Portuguese Yeast Culture Collection for *S. uvarum* strains. This work was supported by NIH

R35GM142849 to CSH.

References

Blackwell, Alexander & Dluzewska, Julia & Szymanska-Lejman, Maja & Desjardins,

Stuart & Tock, Andrew & Kbiri, Nadia & Lambing, Christophe & Lawrence, Emma &

Bieluszewski, Tomasz & Rowan, Beth & Higgins, James & Ziolkowski, Piotr &

Henderson, Ian. (2020). MSH2 shapes the meiotic crossover landscape in relation to

interhomolog polymorphism in Arabidopsis. *The EMBO journal. 39.*

10.15252/embj.2020104858.

Martini, Emmanuelle & Borde, Valérie & Legendre, Matthieu & Audic, Stéphane &

Regnault, Beatrice & Soubigou, Guillaume & Dujon, Bernard & Llorente, Bertrand.

(2011). Genome-Wide Analysis of Heteroduplex DNA in Mismatch Repair–Deficient

Yeast Cells Reveals Novel Properties of Meiotic Recombination Pathways. *PLoS*

genetics. 7. e1002305. 10.1371/journal.pgen.1002305.

Aguillon, S. M., Dodge, T. O., Preising, G. A., & Schumer, M. (2022). Introgression. *Current*

Biology, *32*(16), R865–R868. https://doi.org/10.1016/j.cub.2022.07.004

Albertin, W., Chernova, M., Durrens, P., Guichoux, E., Sherman, D. J., Masneuf-Pomarede, I., &

Marullo, P. (2018). Many interspecific chromosomal introgressions are highly prevalent

in Holarctic Saccharomyces uvarum strains found in human-related fermentations.

Yeast, *35*(1), 141–156. https://doi.org/10.1002/yea.3248

Almeida, P., Gonçalves, C., Teixeira, S., Libkind, D., Bontrager, M., Masneuf-Pomarède, I.,

Albertin, W., Durrens, P., Sherman, D. J., Marullo, P., Todd Hittinger, C., Gonçalves, P.,

& Sampaio, J. P. (2014). A Gondwanan imprint on global diversity and domestication of

wine and cider yeast *Saccharomyces uvarum. Nature Communications*, *5*, 4044.

https://doi.org/10.1038/ncomms5044

Anderson, C. M., Chen, S. Y., Dimon, M. T., Oke, A., DeRisi, J. L., & Fung, J. C. (2011).

ReCombine: A Suite of Programs for Detection and Analysis of Meiotic Recombination in

Whole-Genome Datasets. *PLoS ONE*, *6*(10).

615   https://doi.org/10.1371/journal.pone.0025509

616 Arter, M., & Keeney, S. (2023). Divergence and conservation of the meiotic recombination

617   machinery. *Nature Reviews Genetics*, 1–17. https://doi.org/10.1038/s41576-023-00669-

618   8

619 Auton, A., Li, Y. R., Kidd, J., Oliveira, K., Nadel, J., Holloway, J. K., Hayward, J. J., Cohen, P.

620   E., Greally, J. M., Wang, J., Bustamante, C. D., & Boyko, A. R. (2013). Genetic

621   Recombination Is Targeted towards Gene Promoter Regions in Dogs. *PLOS Genetics*,

622   *9*(12), e1003984. https://doi.org/10.1371/journal.pgen.1003984

623 Barton, N., & Bengtsson, B. O. (1986). The barrier to genetic exchange between hybridising

624   populations. *Heredity*, *57*(3), Article 3. https://doi.org/10.1038/hdy.1986.135

625 Bauer, E., Falque, M., Walter, H., Bauland, C., Camisan, C., Campo, L., Meyer, N., Ranc, N.,

626   Rincent, R., Schipprack, W., Altmann, T., Flament, P., Melchinger, A. E., Menz, M.,

627   Moreno-González, J., Ouzunova, M., Revilla, P., Charcosset, A., Martin, O. C., & Schön,

628   C.-C. (2013). Intraspecific variation of recombination rate in maize. *Genome Biology*,

629   *14*(9), R103. https://doi.org/10.1186/gb-2013-14-9-r103

630 Begun, D. J., & Aquadro, C. F. (1992). Levels of naturally occurring DNA polymorphism

631   correlate with recombination rates in D. melanogaster. *Nature*, *356*(6369), 519–520.

632   https://doi.org/10.1038/356519a0

633 Bendixsen, D. P., Frazão, J. G., & Stelkens, R. (2022). Saccharomyces yeast hybrids on the

634   rise. *Yeast*, *39*(1–2), 40–54. https://doi.org/10.1002/yea.3684

635 Blackwell, A. R., Dluzewska, J., Szymanska-Lejman, M., Desjardins, S., Tock, A. J., Kbiri, N.,

636   Lambing, C., Lawrence, E. J., Bieluszewski, T., Rowan, B., Higgins, J. D., Ziolkowski, P.

637   A., & Henderson, I. R. (2020). MSH2 shapes the meiotic crossover landscape in relation

638   to interhomolog polymorphism in Arabidopsis. *The EMBO Journal*, *39*(21), e104858.

639   https://doi.org/10.15252/embj.2020104858

640 Borts, R. H., & Haber, J. E. (1987). Meiotic Recombination in Yeast: Alteration by Multiple

Heterozygosities. *Science, 237*(4821), 1459–1465.

https://doi.org/10.1126/science.2820060

Bozdag, G. O., Ono, J., Denton, J. A., Karakoc, E., Hunter, N., Leu, J.-Y., & Greig, D. (2021).

Breaking a species barrier by enabling hybrid recombination. *Current Biology*, *31*(4),

R180–R181. https://doi.org/10.1016/j.cub.2020.12.038

Brandvain, Y., Kenney, A. M., Flagel, L., Coop, G., & Sweigart, A. L. (2014). Speciation and

Introgression between Mimulus nasutus and Mimulus guttatus. *PLOS Genetics*, *10*(6),

e1004410. https://doi.org/10.1371/journal.pgen.1004410

Brion, C., Legrand, S., Peter, J., Caradec, C., Pflieger, D., Hou, J., Friedrich, A., Llorente, B., &

Schacherer, J. (2017). Variation of the meiotic recombination landscape and properties

over a broad evolutionary distance in yeasts. *PLOS Genetics*, *13*(8), e1006917.

https://doi.org/10.1371/journal.pgen.1006917

Butlin, R. K. (2005). Recombination and speciation. *Molecular Ecology*, *14*(9), 2621–2635.

https://doi.org/10.1111/j.1365-294X.2005.02617.x

Calfee, E., Gates, D., Lorant, A., Perkins, M. T., Coop, G., & Ross-Ibarra, J. (2021). Selective

sorting of ancestral introgression in maize and teosinte along an elevational cline.

bioRxiv, 2021.03.05.434040. https://doi.org/10.1101/2021.03.05.434040

Charlesworth, B., Morgan, M. T., & Charlesworth, D. (1993). The effect of deleterious mutations

on neutral molecular variation. *Genetics*, *134*(4), 1289–1303.

https://doi.org/10.1093/genetics/134.4.1289

Chen, W., & Jinks-Robertson, S. (1999). The role of the mismatch repair machinery in

regulating mitotic and meiotic recombination between diverged sequences in yeast.

Genetics, *151*(4), 1299–1313.

Chovnick, A., Ballantyne, G. H., & Holm, D. G. (1971). Studies on gene conversion and its

relationship to linked exchange in Drosophila melanogaster. *Genetics*, *69*(2), 179–209.

https://doi.org/10.1093/genetics/69.2.179

Cooper, T. J., Crawford, M. R., Hunt, L. J., Marsolier-Kergoat, M.-C., Llorente, B., & Neale, M. J. (2021). *Mismatch repair disturbs meiotic class I crossover control* (p. 480418). bioRxiv. https://doi.org/10.1101/480418

Crown, K. N., Miller, D. E., Sekelsky, J., & Hawley, R. S. (2018). Local Inversion Heterozygosity Alters Recombination throughout the Genome. *Current Biology: CB*, *28*(18), 2984-2990.e3. https://doi.org/10.1016/j.cub.2018.07.004

Cubillos, F. A., Billi, E., Zörgö, E., Parts, L., Fargier, P., Omholt, S., Blomberg, A., Warringer, J., Louis, E. J., & Liti, G. (2011). Assessing the complex architecture of polygenic traits in diverged yeast populations. *Molecular Ecology*, *20*(7), 1401–1413. https://doi.org/10.1111/j.1365-294X.2011.05005.x

Dagilis, A. J., & Matute, D. R. (2023). The fitness of an introgressing haplotype changes over the course of divergence and depends on its size and genomic location. *PLOS Biology*, *21*(7), e3002185. https://doi.org/10.1371/journal.pbio.3002185

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A., Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

D'Angiolo, M., De Chiara, M., Yue, J.-X., Irizar, A., Stenberg, S., Persson, K., Llored, A., Barré, B., Schacherer, J., Marangoni, R., Gilson, E., Warringer, J., & Liti, G. (2020). A yeast living ancestor reveals the origin of genomic introgressions. *Nature*, *587*(7834), Article 7834. https://doi.org/10.1038/s41586-020-2889-1

Danguy des Déserts, A., Bouchet, S., Sourdille, P., & Servin, B. (2021). Evolution of Recombination Landscapes in Diverging Populations of Bread Wheat. *Genome Biology and Evolution*, *13*(8), evab152. https://doi.org/10.1093/gbe/evab152

Dluzewska, J., Dziegielewski, W., Szymanska-Lejman, M., Gazecka, M., Henderson, I. R., Higgins, J. D., & Ziolkowski, P. A. (2023). MSH2 stimulates interfering and inhibits non-interfering crossovers in response to genetic polymorphism. *Nature Communications*,

14, 6716. https://doi.org/10.1038/s41467-023-42511-z

Duranton, M., & Pool, J. E. (2022). Interactions Between Natural Selection and Recombination Shape the Genomic Landscape of Introgression. *Molecular Biology and Evolution*, *39*(7), msac122. https://doi.org/10.1093/molbev/msac122

Edelman, N. B., Frandsen, P. B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R. B., García-Accinelli, G., Van Belleghem, S. M., Patterson, N., Neafsey, D. E., Challis, R., Kumar, S., Moreira, G. R. P., Salazar, C., Chouteau, M., Counterman, B. A., Papa, R., Blaxter, M., Reed, R. D., … Mallet, J. (2019). Genomic architecture and introgression shape a butterfly radiation. *Science*, *366*(6465), 594–599. https://doi.org/10.1126/science.aaw2090

Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics*, *78*(2), 737–756.

Gerton, J. L., DeRisi, J., Shroff, R., Lichten, M., Brown, P. O., & Petes, T. D. (2000). Global mapping of meiotic recombination hotspots and coldspots in the yeast Saccharomyces cerevisiae. *Proceedings of the National Academy of Sciences*, *97*(21), 11383–11390. https://doi.org/10.1073/pnas.97.21.11383

Harfe, B. D., & Jinks-Robertson, S. (2000). Dna Mismatch Repair and Genetic Instability. *Annual Review of Genetics*, *34*(1), 359–399. https://doi.org/10.1146/annurev.genet.34.1.359

Hill, W. G., & Robertson, A. (1966). The effect of linkage on limits to artificial selection. *Genetical Research*, *8*(3), 269–294.

Hilliker, A. J., Harauz, G., Reaume, A. G., Gray, M., Clark, S. H., & Chovnick, A. (1994). Meiotic gene conversion tract length distribution within the rosy locus of Drosophila melanogaster. *Genetics*, *137*(4), 1019–1026. https://doi.org/10.1093/genetics/137.4.1019

Hoffman, C. S., & Winston, F. (1987). A ten-minute DNA preparation from yeast efficiently releases autonomous plasmids for transformation of Escherichia coli. *Gene*, *57*(2–3),

267–272. https://doi.org/10.1016/0378-1119(87)90131-4

Hunter, N., Chambers, S. R., Louis, E. J., & Borts, R. H. (1996). The mismatch repair system contributes to meiotic sterility in an interspecific yeast hybrid. *The EMBO Journal*, *15*(7), 1726–1733. https://doi.org/10.1002/j.1460-2075.1996.tb00518.x

Jeffreys, A. J., & May, C. A. (2004). Intense and highly localized gene conversion activity in human meiotic crossover hot spots. *Nature Genetics*, *36*(2), Article 2. https://doi.org/10.1038/ng1287

Judd, S. R., & Petes, T. D. (1988). Physical Lengths of Meiotic and Mitotic Gene Conversion Tracts in Saccharomyces Cerevisiae. *Genetics*, *118*(3), 401. https://doi.org/10.1093/genetics/118.3.401

Keeney, S. (2001). Mechanism and control of meiotic recombination initiation. *Current Topics in Developmental Biology*, *52*, 1–53. https://doi.org/10.1016/s0070-2153(01)52008-6

Kiktev, D. A., Sheng, Z., Lobachev, K. S., & Petes, T. D. (2018). GC content elevates mutation and recombination rates in the yeast *Saccharomyces cerevisiae*. *Proceedings of the National Academy of Sciences*, *115*(30), E7109–E7118. https://doi.org/10.1073/pnas.1807334115

Kong, A., Thorleifsson, G., Frigge, M. L., Masson, G., Gudbjartsson, D. F., Villemoes, R., Magnusdottir, E., Olafsdottir, S. B., Thorsteinsdottir, U., & Stefansson, K. (2014). Common and low-frequency variants associated with genome-wide recombination rate. *Nature Genetics*, *46*(1), 11–16. https://doi.org/10.1038/ng.2833

Korunes, K. L., & Noor, M. A. F. (2017). Gene conversion and linkage: Effects on genome evolution and speciation. *Molecular Ecology*, *26*(1), 351–364. https://doi.org/10.1111/mec.13736

Korunes, K. L., & Noor, M. A. F. (2019). Pervasive gene conversion in chromosomal inversion heterozygotes. *Molecular Ecology*, *28*(6), 1302–1315. https://doi.org/10.1111/mec.14921

Langdon, Q. K., Peris, D., Baker, E. P., Opulente, D. A., Nguyen, H.-V., Bond, U., Gonçalves,

P., Sampaio, J. P., Libkind, D., & Hittinger, C. T. (2019). Fermentation innovation

through complex hybridization of wild and domesticated yeasts. *Nature Ecology &*

Evolution, 1–11. https://doi.org/10.1038/s41559-019-0998-8

Langdon, Q. K., Peris, D., Eizaguirre, J. I., Opulente, D. A., Buh, K. V., Sylvester, K., Jarzyna,

M., Rodríguez, M. E., Lopes, C. A., Libkind, D., & Hittinger, C. T. (2020). Postglacial

migration shaped the genomic diversity and global distribution of the wild ancestor of

lager-brewing hybrids. *PLOS Genetics*, *16*(4), e1008680.

https://doi.org/10.1371/journal.pgen.1008680

Layer, R. M., Chiang, C., Quinlan, A. R., & Hall, I. M. (2014). LUMPY: A probabilistic framework

for structural variant discovery. *Genome Biology*, *15*(6), R84. https://doi.org/10.1186/gb-

2014-15-6-r84

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics (Oxford, England)*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Liu, H., Huang, J., Sun, X., Li, J., Hu, Y., Yu, L., Liti, G., Tian, D., Hurst, L. D., & Yang, S.

(2018). Tetrad analysis in plants and fungi finds large differences in gene conversion

rates but no GC bias. *Nature Ecology & Evolution*, *2*(1), Article 1.

https://doi.org/10.1038/s41559-017-0372-7

Liu, H., Maclean, C. J., & Zhang, J. (2019). Evolution of the Yeast Recombination Landscape.

Molecular Biology and Evolution, *36*(2), 412–422.

https://doi.org/10.1093/molbev/msy233

Magwene, P. M., Kayıkçı, Ö., Granek, J. A., Reininga, J. M., Scholl, Z., & Murray, D. (2011).

Outcrossing, mitotic recombination, and life-history trade-offs shape genome evolution in

Saccharomyces cerevisiae. *Proceedings of the National Academy of Sciences*, *108*(5),

1987–1992. https://doi.org/10.1073/pnas.1012544108

Mancera, E., Bourgon, R., Brozzi, A., Huber, W., & Steinmetz, L. M. (2008). High-resolution

mapping of meiotic crossovers and non-crossovers in yeast. *Nature*, *454*(7203), 479–485. https://doi.org/10.1038/nature07135

Marsolier-Kergoat, M.-C., & Yeramian, E. (2009). GC Content and Recombination: Reassessing the Causal Effects for the Saccharomyces cerevisiae Genome. *Genetics*, *183*(1), 31–38. https://doi.org/10.1534/genetics.109.105049

Martin, S. H., Davey, J. W., Salazar, C., & Jiggins, C. D. (2019). Recombination rate variation shapes barriers to introgression across butterfly genomes. *PLOS Biology*, *17*(2), e2006288. https://doi.org/10.1371/journal.pbio.2006288

Martini, E., Borde, V., Legendre, M., Audic, S., Regnault, B., Soubigou, G., Dujon, B., & Llorente, B. (2011). Genome-Wide Analysis of Heteroduplex DNA in Mismatch Repair–Deficient Yeast Cells Reveals Novel Properties of Meiotic Recombination Pathways. *PLOS Genetics*, *7*(9), e1002305. https://doi.org/10.1371/journal.pgen.1002305

Mazina, O. M., Mazin, A. V., Nakagawa, T., Kolodner, R. D., & Kowalczykowski, S. C. (2004). Saccharomyces cerevisiae Mer3 helicase stimulates 3'-5' heteroduplex extension by Rad51; implications for crossover control in meiotic recombination. *Cell*, *117*(1), 47–56. https://doi.org/10.1016/s0092-8674(04)00294-6

McDonald, M. J., Rice, D. P., & Desai, M. M. (2016). Sex speeds adaptation by altering the dynamics of molecular evolution. *Nature*, *531*(7593), 233–236. https://doi.org/10.1038/nature17143

McGaugh, S. E., Heil, C. S. S., Manzano-Winkler, B., Loewe, L., Goldstein, S., Himmel, T. L., & Noor, M. A. F. (2012). Recombination Modulates How Selection Affects Linked Sites in Drosophila. *PLOS Biology*, *10*(11), e1001422. https://doi.org/10.1371/journal.pbio.1001422

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K., Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Research, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Miller, D. E., Smith, C. B., Kazemi, N. Y., Cockrell, A. J., Arvanitakis, A. V., Blumenstiel, J. P., Jaspersen, S. L., & Hawley, R. S. (2016). Whole-Genome Analysis of Individual Meiotic Events in Drosophila melanogaster Reveals That Noncrossover Gene Conversions Are Insensitive to Interference and the Centromere Effect. *Genetics*, *203*(1), 159–171. https://doi.org/10.1534/genetics.115.186486

Moran, B. M., Payne, C., Langdon, Q., Powell, D. L., Brandvain, Y., & Schumer, M. (2021). The genomic consequences of hybridization. *eLife*, *10*, e69016. https://doi.org/10.7554/eLife.69016

Myung, K., Datta, A., Chen, C., & Kolodner, R. D. (2001). SGS1, the Saccharomyces cerevisiae homologue of BLM and WRN, suppresses genome instability and homeologous recombination. *Nature Genetics*, *27*(1), 113–116. https://doi.org/10.1038/83673

Nachman, M. & Payseur, B. (2012). Recombination rate variation and speciation: Theoretical predictions and empirical results from rabbits and mice. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *367*(1587), 409–421. https://doi.org/10.1098/rstb.2011.0249

Nespolo, R. F., Villarroel, C. A., Oporto, C. I., Tapia, S. M., Vega-Macaya, F., Urbina, K., Chiara, M. D., Mozzachiodi, S., Mikhalev, E., Thompson, D., Larrondo, L. F., Saenz-Agudelo, P., Liti, G., & Cubillos, F. A. (2020). An Out-of-Patagonia migration explains the worldwide diversity and distribution of Saccharomyces eubayanus lineages. *PLOS Genetics*, *16*(5), e1008777. https://doi.org/10.1371/journal.pgen.1008777

Petit, M., Astruc, J.-M., Sarry, J., Drouilhet, L., Fabre, S., Moreno, C. R., & Servin, B. (2017). Variation in Recombination Rate and Its Genetic Determinism in Sheep Populations. *Genetics*, *207*(2), 767–784. https://doi.org/10.1534/genetics.117.300123

Pool, J. E. (2015). The Mosaic Ancestry of the Drosophila Genetic Reference Panel and the D. melanogaster Reference Genome Reveals a Network of Epistatic Fitness Interactions.

Molecular Biology and Evolution, *32*(12), 3236–3251.

https://doi.org/10.1093/molbev/msv194

Raffoux, X., Bourge, M., Dumas, F., Martin, O. C., & Falque, M. (2018). Role of Cis, Trans, and

Inbreeding Effects on Meiotic Recombination in Saccharomyces cerevisiae. *Genetics*,

210(4), 1213–1226. https://doi.org/10.1534/genetics.118.301644

Ravinet, M., Yoshida, K., Shigenobu, S., Toyoda, A., Fujiyama, A., & Kitano, J. (2018). The

genomic landscape at a late stage of stickleback speciation: High genomic divergence

interspersed by small localized regions of introgression. *PLOS Genetics*, *14*(5),

e1007358. https://doi.org/10.1371/journal.pgen.1007358

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &

Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology*, *29*(1), 24–26.

https://doi.org/10.1038/nbt.1754

Rockman, M. V., & Kruglyak, L. (2009). Recombinational Landscape and Population Genomics

of Caenorhabditis elegans. *PLOS Genetics*, *5*(3), e1000419.

https://doi.org/10.1371/journal.pgen.1000419

Rogers, D. W., McConnell, E., Ono, J., & Greig, D. (2018). Spore-autonomous fluorescent

protein expression identifies meiotic chromosome mis-segregation as the principal cause

of hybrid sterility in yeast. *PLOS Biology*, *16*(11), e2005066.

https://doi.org/10.1371/journal.pbio.2005066

Ruderfer, D. M., Pratt, S. C., Seidel, H. S., & Kruglyak, L. (2006). Population genomic analysis

of outcrossing and recombination in yeast. *Nature Genetics*, *38*(9), Article 9.

https://doi.org/10.1038/ng1859

Samuk, K., Manzano-Winkler, B., Ritz, K. R., & Noor, M. A. F. (2019). *Natural selection shapes*

variation in genome-wide recombination rate in Drosophila pseudoobscura [Preprint].

Genetics. https://doi.org/10.1101/787382

Sandor, C., Li, W., Coppieters, W., Druet, T., Charlier, C., & Georges, M. (2012). Genetic

variants in REC8, RNF212, and PRDM9 influence male recombination in cattle. *PLoS*

Genetics, *8*(7), e1002854. https://doi.org/10.1371/journal.pgen.1002854

Scannell, D. R., Zill, O. A., Rokas, A., Payen, C., Dunham, M. J., Eisen, M. B., Rine, J.,

Johnston, M., & Hittinger, C. T. (2011). The Awesome Power of Yeast Evolutionary

Genetics: New Genome Sequences and Strain Resources for the Saccharomyces sensu

stricto Genus. *G3: Genes, Genomes, Genetics*, *1*(1), 11–25.

https://doi.org/10.1534/g3.111.000273

Schaeffer, S. W., & Anderson, W. W. (2005). Mechanisms of Genetic Exchange Within the

Chromosomal Inversions of Drosophila pseudoobscura. *Genetics*, *171*(4), 1729–1739.

https://doi.org/10.1534/genetics.105.041947

Schreiber, M., Gao, Y., Koch, N., Fuchs, J., Heckmann, S., Himmelbach, A., Börner, A., Özkan,

H., Maurer, A., Stein, N., Mascher, M., & Dreissig, S. (2022). Recombination Landscape

Divergence Between Populations is Marked by Larger Low-Recombining Regions in

Domesticated Rye. *Molecular Biology and Evolution*, *39*(6), msac131.

https://doi.org/10.1093/molbev/msac131

Schumer, M., Xu, C., Powell, D. L., Durvasula, A., Skov, L., Holland, C., Blazier, J. C.,

Sankararaman, S., Andolfatto, P., Rosenthal, G. G., & Przeworski, M. (2018). Natural

selection interacts with recombination to shape the evolution of hybrid genomes.

Science, *360*(6389), 656–660. https://doi.org/10.1126/science.aar3684

Sharma, D., Denmat, S. H.-L., Matzke, N. J., Hannan, K., Hannan, R. D., O'Sullivan, J. M., &

Ganley, A. R. D. (2022). A new method for determining ribosomal DNA copy number

shows differences between *Saccharomyces cerevisiae* populations. *Genomics*, *114*(4),

110430. https://doi.org/10.1016/j.ygeno.2022.110430

Shi, J., Wolf, S. E., Burke, J. M., Presting, G. G., Ross-Ibarra, J., & Dawe, R. K. (2010).

Widespread Gene Conversion in Centromere Cores. *PLoS Biology*, *8*(3), e1000327.

https://doi.org/10.1371/journal.pbio.1000327

Singhal, S., Leffler, E. M., Sannareddy, K., Turner, I., Venn, O., Hooper, D. M., Strand, A. I., Li,

Q., Raney, B., Balakrishnan, C. N., Griffith, S. C., McVean, G., & Przeworski, M. (2015).

Stable recombination hotspots in birds. *Science (New York, N.Y.)*, *350*(6263), 928–932.

https://doi.org/10.1126/science.aad0843

Smith, J. M., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical*

Research, *23*(1), 23–35.

Smukowski, C. S., & Noor, M. a. F. (2011). Recombination rate variation in closely related

species. *Heredity*, *107*(6), 496–508. https://doi.org/10.1038/hdy.2011.44

Smukowski Heil, C. S., Ellison, C., Dubin, M., & Noor, M. A. F. (2015). Recombining without

Hotspots: A Comprehensive Evolutionary Portrait of Recombination in Two Closely

Related Species of Drosophila. *Genome Biology and Evolution*, *7*(10), 2829–2842.

https://doi.org/10.1093/gbe/evv182

Stapley, J., Feulner, P. G. D., Johnston, S. E., Santure, A. W., & Smadja, C. M. (2017).

Variation in recombination frequency and distribution across eukaryotes: Patterns and

processes. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1736), 20160455. https://doi.org/10.1098/rstb.2016.0455

Stelkens, R., & Bendixsen, D. P. (2022). The evolutionary and ecological potential of yeast

hybrids. *Current Opinion in Genetics & Development*, *76*, 101958.

https://doi.org/10.1016/j.gde.2022.101958

Szymanska-Lejman, M., Dziegielewski, W., Dluzewska, J., Kbiri, N., Bieluszewska, A., Poethig,

R. S., & Ziolkowski, P. A. (2023). The effect of DNA polymorphisms and natural variation

on crossover hotspot activity in Arabidopsis hybrids. *Nature Communications*, *14*(1), 33.

https://doi.org/10.1038/s41467-022-35722-3

Talbert, P. B., & Henikoff, S. (2010). Centromeres Convert but Don't Cross. *PLOS Biology*, *8*(3),

e1000326. https://doi.org/10.1371/journal.pbio.1000326

Tellini, N., De Chiara, M., Mozzachiodi, S., Tattini, L., Vischioni, C., Naumova, E., Warringer, J.,

Bergström, A., & Liti, G. (2023). *Ancient and recent origins of shared polymorphisms in yeas.* https://doi.org/10.21203/rs.3.rs-2573222/v1

Tsai, I. J., Burt, A., & Koufopanou, V. (2010). Conservation of recombination hotspots in yeast. *Proceedings of the National Academy of Sciences.* https://doi.org/10.1073/pnas.0908774107

Veller, C., Edelman, N. B., Muralidhar, P., & Nowak, M. A. (2023). Recombination and selection against introgressed DNA. *Evolution*, *77*(4), 1131–1144. https://doi.org/10.1093/evolut/qpad021

Veller, C., Kleckner, N., & Nowak, M. A. (2019). A rigorous measure of genome-wide genetic shuffling that takes into account crossover positions and Mendel's second law. *Proceedings of the National Academy of Sciences of the United States of America*, *116*(5), 1659–1668. https://doi.org/10.1073/pnas.1817482116

Wijnker, E., Velikkakam James, G., Ding, J., Becker, F., Klasen, J. R., Rawat, V., Rowan, B. A., de Jong, D. F., de Snoo, C. B., Zapata, L., Huettel, B., de Jong, H., Ossowski, S., Weigel, D., Koornneef, M., Keurentjes, J. J., & Schneeberger, K. (2013). The genomic landscape of meiotic crossovers and gene conversions in Arabidopsis thaliana. *eLife*, *2*, e01426. https://doi.org/10.7554/eLife.01426

Zeyl, C. W., & Otto, S. P. (2007). A short history of recombination in yeast. *Trends in Ecology & Evolution*, *22*(5), 223–225. https://doi.org/10.1016/j.tree.2007.02.005

Ziolkowski, P. A., Berchowitz, L. E., Lambing, C., Yelina, N. E., Zhao, X., Kelly, K. A., Choi, K., Ziolkowska, L., June, V., Sanchez-Moran, E., Franklin, C., Copenhaver, G. P., & Henderson, I. R. (2015). Juxtaposition of heterozygous and homozygous regions causes reciprocal crossover remodelling via interference during Arabidopsis meiosis. *eLife*, *4*, e03708. https://doi.org/10.7554/eLife.03708

Ziolkowski, P. A., & Henderson, I. R. (2017). Interconnections between meiotic recombination and sequence polymorphism in plant genomes. *New Phytologist*, *213*(3), 1022–1029.

927   https://doi.org/10.1111/nph.14265

930