The recombination landscape of introgression in yeast

Enrique J. Schwarzkopf[1], Nathan Brandt[1], Caiti Smukowski Heil*[1]

Affiliation:

[1] Department of Biological Sciences, North Carolina State University, Raleigh, NC

*Corresponding author: cheil@ncsu.edu

Abstract

Meiotic recombination is an important evolutionary force that acts by breaking up genomic

linkage, thereby increasing the efficacy of selection. Meiotic recombination is initiated with a

double-strand break which is resolved via a crossover, which involves the reciprocal exchange

of genetic material between homologous chromosomes, or a non-crossover, which results in

small tracts of non-reciprocal exchange of genetic material. While the meiotic process is largely

conserved, crossover and non-crossover rates vary between species, populations, individuals,

and across the genome. In recent years, recombination is observed to be positively associated

with the distribution of ancestry derived from past interspecific hybridization (introgression) in a

variety of species. This trend has been interpreted to signify that introgression carries genetic

incompatibilities that are selected against, such that introgression is enriched in regions of high

recombination. However, recombination is well known to be suppressed in divergent sequence

to prevent non-homologous recombination. Since introgressed DNA is often divergent, we

sought to explore this interaction of recombination and introgression by sequencing spores and

detecting crossover and non-crossover events from two crosses of the budding yeast

Saccharomyces uvarum. One cross is between strains isolated from natural environments, and

the other cross is between strains from fermentation environments, in which each strain

contains introgression from their sister species, *S. eubayanus*. We find that the recombination

landscape is significantly different between *S. uvarum* crosses, and that most of these

differences can be explained by the presence of heterozygous introgression in the fermentation

cross. Crossovers are significantly reduced and non-crossovers are increased in heterozygous

introgression compared to syntenic regions in the natural cross without introgression. This

translates to reduced allele shuffling within introgressed regions, and an overall reduction of

shuffling on most chromosomes with introgression compared to the syntenic regions and

chromosomes without introgression. Our results indicate that recent hybridization can

significantly influence the recombination landscape, and suggest that the reduction in allele

shuffling contributes to the initial purging of introgressed ancestry in the generations following a

hybridization event.

Keywords: introgression, hybridization, recombination, crossover, non-crossover, yeast,

Saccharomyces

Introduction

Recombination is the exchange of genetic material between homologous chromosomes during meiosis and is a staple of eukaryotic sexual reproduction. While the processes involved in recombination are largely conserved (Arter & Keeney, 2023), recombination rates vary between sexes, populations, and species (Smukowski & Noor, 2011; Stapley et al., 2017). Recombination rates also vary along the genome, with conflicting patterns of enriched or depleted recombination in promoter regions and punctate or dispersed recombination depending on the species (Auton et al., 2013; Rockman & Kruglyak, 2009; Singhal et al., 2015; Smukowski Heil et al., 2015). These patterns in recombination can affect pairing of alleles after meiosis–in other words, the shuffling of alleles–in a population. Much of the evolutionary advantage of recombination is understood to originate from its role in shuffling alleles, which increases the number of different allele combinations segregating in a population. The increase in allele combinations can reduce selection interference–the effect that genetically linked sites have on the evolutionary fate of either beneficial or deleterious alleles (Felsenstein, 1974; Hill & Robertson, 1966; McDonald et al., 2016; McGaugh et al., 2012).

How much allele decoupling is produced by recombination will depend on the type of recombination event. Each recombination event begins with the severing of both strands of a sister chromatid of one of the homologous chromosomes in what is referred to as a double-strand break (DSB) (Keeney, 2001). The distribution of DSBs is influenced by a variety of factors, many of which are organism-dependent, but often include decreases of DSBs near telomeres and centromeres as well as increases in genomic regions enriched for GC content, CpG sites, and depleted of methylation (CpG islands) and near promoter regions (de Massy, 2013; Lam & Keeney, 2015; Pan et al., 2011; Zelkowski et al., 2019). When a DSB occurs, the homologous strand is recruited to repair the break, and during this process genetic information

is exchanged. The most evident and widely studied resolution of a DSB is a crossover (CO),

where all the genetic information on one side of the DSB from one homologous chromosome is

spliced with all the genetic information on the other side of the DSB from the other homologous

chromosome. DSB resolution can also involve gene conversions–or non-crossovers (NCOs)–

which result in one small segment (typically 100-2000 bp) of a homologous chromosome's

genetic information being copied onto the other (Chovnick et al., 1971; Hilliker et al., 1994;

Jeffreys & May, 2004; Judd & Petes, 1988). DSBs can be resolved through both of these

processes in a single instance, and even more than one NCO can occur at a single break point.

Each possible resolution can produce a variety of genetic patterns at the site of a DSB. COs

generally produce more allele shuffling, and therefore degrade linkage faster than NCOs,

because they recombine all loci from one side of the event with all loci on the other side.

However, NCOs can occur in regions where COs are typically suppressed, like centromeres

and inversions (Korunes & Noor, 2019; Mancera et al., 2008; Miller et al., 2016; Schaeffer &

Anderson, 2005; Shi et al., 2010; Talbert & Henikoff, 2010; Wijnker et al., 2013). NCOs are also

crucial to reducing linkage within coding regions and, unlike COs, result in 3:1 allele ratio in the

meiotic product at heterozygous sites, potentially changing allele frequencies (Korunes & Noor,

2017).

The number of DSBs that occur per meiosis varies by species, as does the number of DSBs

that are repaired as COs or NCOs (de Massy, 2013; Korunes & Noor, 2017). In Arabidopsis and

maize, several hundred DSBs are resolved into only a handful of COs, whereas in

Saccharomyces cerevisiae the CO:NCO ratio is close to 2:1 (Choi et al., 2018; He et al., 2017;

Mancera et al., 2008). How CO:NCO resolution may be evolving is not well understood, but

evidence from *Saccharomyces* suggests that the DSB landscape is well conserved across

species, while recombination rates significantly differ between sister species *S. cerevisiae* and

S. paradoxus (Lam & Keeney, 2015; Liu et al., 2019). This suggests that alternate resolution of

DSBs may be important in altering recombination rates between closely related species. This is

further supported by the finding that several genes involved in the CO vs. NCO decision of DSB

resolution show evidence of rapid evolution and directional selection in mammals (Dapper &

Payseur, 2019), and that some of these genes are linked to variation in recombination rates

within and between species (Brand et al., 2018; Kong et al., 2008; Murdoch et al., 2010; Yang et

al., 2015).

Variation in the number and distribution of COs and NCOs, and their respective associated

effects on linkage, have important implications for molecular evolution. Recombination has long

been appreciated to play a role in the distribution of various genomic features including

nucleotide diversity. Nucleotide diversity has a positive correlation with recombination rate in a

number of species, interpreted to result from selective sweeps and background selection

removing genetic variation in regions of low recombination (Begun & Aquadro, 1992;

Charlesworth et al., 1993; Smith & Haigh, 1974). Similarly, recombination breaking up genetic

associations is particularly notable in the context of interspecific hybridization. In first-generation

(F_1) hybrids, each set of homologous chromosomes is composed of sister chromatids carrying

genetic information from one of the parents. Since no recombination has physically separated

any alleles on the chromosome at the hybrid F_1 stage, each parent's genetic contribution is

perfectly linked. If the hybrids then back-cross to one of the parental populations, recombination

will produce genomes that are a mosaic of genetic information from the two populations

(introgression) (Aguillon et al., 2022). Recombination drives the pattern of fragmentation in the

introgressed regions over time, and therefore plays an important role in the distribution of

introgressed DNA in a population (Barton & Bengtsson, 1986; Butlin, 2005; Moran et al., 2021;

Nachman Michael W. & Payseur Bret A., 2012; Schumer et al., 2018; Veller et al., 2023). When

each population has evolved alleles that are deleterious when present in the background of the

other population (the Dobzhansky-Muller hybrid incompatibility model) we expect introgressed

regions with low rates of recombination to be quickly purged from the population, as the

accumulation of incompatible alleles incurs a steep fitness cost. In contrast, when introgressed

regions have high recombination rates, the break up of genetic associations will reduce

selective interference between the incompatible alleles and their surrounding haplotypes,

allowing for neutral and beneficial alleles brought in with the introgression to escape the fate of

neighboring incompatibilities (Moran et al., 2021; Schumer et al., 2018). This theory is

supported empirically through enrichment of introgressed segments in regions of higher

recombination in a number of organisms including Mimulus, maize, butterflies, swordtail fish,

stickleback, and humans (Brandvain et al., 2014; Calfee et al., 2021; Edelman et al., 2019;

Martin et al., 2019; Ravinet et al., 2018; Schumer et al., 2018).

This positive correlation between introgressed ancestry and recombination is emerging as a

nearly ubiquitous pattern (though see (Dagilis & Matute, 2023; Duranton & Pool, 2022; Pool,

2015)), however, it is unclear how these observations relate to the known effect of sequence

divergence on DSB resolution. Introgression, particularly between highly diverged species, can

have low sequence homology with the genomic region it is replacing. A DSB in a region of low

homology will recruit mismatch repair proteins, which ensure COs are occurring between

homologous chromosomes and at equivalent positions to prevent ectopic recombination (Harfe

& Jinks-Robertson, 2000; Hunter et al., 1996). Mismatch repair proteins reduce the frequency of

CO events as sequence divergence increases (Chen & Jinks-Robertson, 1999; Cooper et al.,

2021; L. Li et al., 2006; Welz-Voegele & Jinks-Robertson, 2008). Given that heterozygous

introgression will have divergent sequences, we expect a decrease in COs, and possibly an

increase in NCOs as DSBs fail to be resolved as COs in introgressed regions .

To help us understand this interaction of introgression and recombination, and identify patterns

in CO and NCO in closely related populations, we utilized the budding yeast *Saccharomyces*.

Yeasts provide an excellent opportunity to study DSB resolution, as we can readily isolate and

collect all four meiotic products of a given meiosis and detect both CO and more elusive NCO

events (Figure 1A) (Brion et al., 2017; Gerton et al., 2000; Liu et al., 2018, 2019; Mancera et al.,

2008). Recombination rates vary between strains of *S. cerevisiae* (Cubillos et al., 2011; Raffoux

et al., 2018) and between *S. cerevisiae* and its sister species *S. paradoxus* (Liu et al., 2019;

Tsai et al., 2010). Strains of different *Saccharomyces* species have often hybridized with other

species and carry introgressed DNA from these events (Albertin et al., 2018; Almeida et al.,

2014; Bendixsen et al., 2022; D'Angiolo et al., 2020; Langdon et al., 2019; Stelkens &

Bendixsen, 2022; Tellini et al., 2023).

In this study, we look at patterns of recombination and introgression at the population level by

crossing two pairs of Holarctic *Saccharomyces uvarum* strains. One pair of strains was isolated

from natural environments in North America and the other pair was isolated from European

fermentation environments (Almeida et al., 2014). The *S. uvarum* strains isolated from

European fermentation environments each carry introgression from their sister species,

Saccharomyces eubayanus, which is approximately 6% divergent from *S. uvarum* (Almeida et

al., 2014; Langdon et al., 2020; Nespolo et al., 2020). The diploid F_1 genome of these strains is

heterozygous for nine different introgressions which make up approximately 10% of the genome

(Figure 1B). The strains from the North American cross do not carry *S. eubayanus*

introgression, thus allowing us to assess the impact of introgression on the recombination

landscape. We obtained whole genome sequencing data from individual meiotic events from the

first offspring generation of each cross and used this data to detect CO and NCO events along

the genome (Figure 1A). From these maps, we aim to understand (i) how patterns of CO and

NCO differ between closely related strains, (ii) how regions of introgression differ in their CO

and NCO patterns, and (iii) how these different patterns affect shuffling of alleles locally and at

the chromosome level. Understanding these objectives will provide us novel insights into how

introgression impacts the recombination landscape.

Methods

Strain and library construction

S. uvarum strains (UCD61-137, yHCT78 , GM14, and DBVPG7787) were obtained from the

Portuguese Yeast Culture Collection and from Chris Hittinger (Table S1) (Almeida et al., 2014).

All four *S. uvarum* strains had their *HO* locus replaced with a kanMX marker using a modified

version of the high-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method.

Briefly, the kanMX marker was amplified from plasmid pCSH2 with homology to genomic DNA

flanking the HO ORF with primers CSH239

(GGTGGAAAACCACGAAAAGTTAGAACTACGTTCAGGCAAAgacatggaggcccagaatac) and

CSH241 (GTGACCGTATTGGTACTTTTTTTGTTACCTGTTTTAGTAGcagtatagcgaccagcattc).

For each strain, overnight cultures were inoculated in 25 mL of YPD at an OD of ~ 0.0005 and

incubated at room temperature on a shaker for ~24 hours until the cultures reached an OD

between 0.6 and 0.9. Subsequently, 1 ug of the template DNA was transformed with a heat

shock temperature of 37°C for 45 minutes. The transformed cells were allowed to recover in

liquid YPD for 4 hours before being plated onto G418 selective plates and incubated at room

temperature for 2 days.

Single colonies were selected from the transformation plates, restreaked onto G418 plates and

allowed to grow at room temperature for 2 days. Single colonies from those plates were then

inoculated into 2 mL of YPD + G418 and incubated in a roller drum at room temperature

overnight. From those cultures, 250 uL was used to inoculate 2 mL of sporulation media (1%

potassium acetate, 0.1 % yeast extract, 0.05% dextrose) and incubated at room temperature for

3 to 5 days. Strains were confirmed to have the ho::KanMX via tetrad dissection on a Singer

SporPlay+ microscope (Singer Instruments). Plates with tetrads were incubated at room

temperature for 2 days and then replica plated to test for proper segregation of the kanMX

marker and mating type within individual tetrads.

Crosses between strains UCD61-137 and yHCT78 (natural cross), and between strains GM14

and DBVPG7787 (fermentation cross) were set up by micromanipulation of single MATa and

MATx cells using a Singer SporPlay+. The plates were incubated at room temperature for 2

days and then replica plated to mating type tester strains to test for potential diploids. Identified

diploids were then sporulated by growing a culture of the cross in 2 mL YPD + G418 at room

temperature overnight. From those cultures, 250 uL were used to inoculate 2 mL of sporulation

media and incubated at room temperature for 3 to 5 days. Sporulated cultures were dissected

on 3 YPD plates (24 tetrads per plate) using a Singer SporPlay+. Fifty of the fully viable tetrads

were selected and had all their spores inoculated into YPD (200 spores total) and incubated at

room temperature. The DNA was extracted from these cultures using a modified version of the

Hoffman-Winston DNA Prep (Hoffman & Winston, 1987). The DNA concentration was then

measured using SYBR green, and 150 ng of each sample's DNA was used to prepare a

sequencing library using an Illumina DNA Prep Kit, modified to use half the normal amounts of

reagents. Libraries were pooled and run on an Illumina NovaSeq 500 with 150bp paired end

reads.

Calling SNPs

We scored SNPs from parents and offspring using the *S. uvarum* reference genome (Scannell

et al., 2011) and custom scripts that invoked bwa (v0.7.17), samtools (v1.12), bcftools (v1.13),

picardtools (v2.25.6), and gatk (v4.2.0.0) (Danecek et al., 2021; H. Li & Durbin, 2009; McKenna

et al., 2010). The custom scripts are available in the github repository: *ejschwarzkopf/CO-NCO.*

We joint genotyped parents and offspring with default filters for gatk with the exception of the

QUAL filter, which was set as < 100 for parents and < 30 for offspring. We further filtered

variants by requiring they be fixed differences between the two parental strains. We kept a total

of 24,574 markers for the natural cross and 74,619 markers for the fermentation cross. We

utilized LUMPY to identify structural variants in the parent strains that were greater than 5000

bp and verified calls using the Integrative Genomics Viewer (Layer et al., 2014; Robinson et al.,

2011). We identified three amplifications in strain GM14 (one of the fermentation cross parents)

that were absent in other strains (Table S2).

Generating CO/NCO maps

We generated "seg" files by coding tetrad variants by their parental origin. These seg files were

the input for CrossOver (v6.3) from the ReCombine suite of programs, which we used to detect

COs and NCOs (Anderson et al., 2011). We then filtered to remove non-crossovers with fewer

than three associated markers and split the genome into 20kb windows. In each window we

counted crossovers, non-crossovers, and markers. We established regions of introgressions

through visual inspection of marker density in the fermentation cross (introgressions showed

more divergence between fermentation strains) and confirmed them using the findings of

Almeida et al. (2014). We found nine heterozygous introgressions on chromosomes 4, 6, 7, 9,

10, 10, 13, 14, and 15 respectively that we included in further analyses (Table S3). We excluded

two additional introgressions due to poor mapping (chromosome 13:0-17,000; chromosome 16:

642,000-648,000). To account for the difference in number of markers in introgressed vs non-

introgressed windows and their effect on NCO detection, we applied a previously published

simulation-based method (Liu et al., 2019; Wijnker et al., 2013). Choosing an average NCO

tract length of 2kb, we simulated 1000 NCO events per window to establish our expected

probability of detecting an NCO event in that window. We then divided our observed NCO count

by our probability of detecting an NCO event. COs that occurred in large regions devoid of

markers would be called in the middle of the empty windows. We decided to deal with this by

splitting CO counts in regions with multiple consecutive windows without markers evenly

between the empty windows. With these corrected maps, we calculated spearman correlations

between crosses using R (v4.1.0, R Core Team 2021). Additionally, we modeled NCO and CO

count as a function of introgression, introgression by cross, and GC content using a gaussian

generalized linear model in R (v4.1.0, R Core Team 2021).

Homology

We calculated homology between the two fermentation cross strains in 51bp windows with 25bp

overlaps. At each nucleotide position in the window, we counted fixed differences as zero

homology, invariant sites between strains as full homology (1), and polymorphic sites in either or

both strains as half homology (0.5). We then averaged these homology values across the

window. This measure represents the probability that both strains will have the same nucleotide

base at a given position. We used this measure of fine-scale homology to determine how

homology related to NCO counts in introgressed regions. For this, we used Loess regressions

and Spearman's correlations on each of the introgressed regions comparing homology to NCO

count, both implemented in R (v4.1.0, R Core Team 2021). We then focused on each

recombination event (CO or NCO) and compared the homology 100bp up and downstream of

CO breakpoints and 100bp up and downstream of NCO tracts. We then used Welch's two

sample t-tests to compare CO and NCO homology in each introgression.

283   *r*

We use r, a measure genetic shuffling defined in Veller et al. (2019) to measure how much

shuffling occurs in each chromosome for each cross. Our data provides parental origin for each

fixed difference between parental strains. We assume that all loci between pairs of markers that

come from the same parent are also from that parent. We also assume that when a pair of

successive markers come from different parents, the location of the change from one parental

origin to the other happens at the midpoint between our markers. With this in mind, we counted

the number of bases that come from one parent and divided by the chromosome size to obtain

the proportion of the chromosome that was inherited from said parent (p) and used the formula

from Veller et al. (2019): $r = 2p(1 - p)$. We calculated r for each full chromosome and each

introgressed regions in every gamete from both crosses. We then averaged across gametes to

obtain average r values. We then compared average r between crosses in each chromosome

or introgressed region using Welch two sample t-tests and correcting for multiple tests using a

Bonferroni correction in R (v4.1.0, R Core Team 2021).

Data availability

Sequences for the parental strains can be found on NCBI SRA (SRR1119189, SRR1119180

SRR1119199, SRR1119200) (Almeida et al., 2014). Sequencing of the tetrads is deposited at

NCBI SRA under Project PRJNA1061120.

Results

The recombination landscape differs dramatically between closely related crosses

We isolated and sequenced products of 48 meioses (192 haploid spores) for two crosses of *S.*

uvarum, a cross between strains isolated from North America (natural cross) and a cross

between strains isolated from Europe (fermentation cross). We detected COs and NCOs across

the 16 nuclear chromosomes of *S. uvarum*. Genomewide, we found significantly more COs on

average in the natural cross (82.54 COs/meiosis, SE 1.5; 0.72 cM/kb) than in the fermentation

cross (63.66 COs/meiosis, SE 1.9; 0.55 cM/kb) but no significant difference when comparing the

average number of NCOs between the natural cross (60.44 NCOs/meiosis, SE 2.7) and the

fermentation cross (68.1 NCOs/meiosis, SE 12.47). The number of COs per meiosis in the

natural and fermentation crosses are slightly higher than those of *S. paradoxus* (54.8) and *S.*

cerevisiae (76.5) respectively, but both crosses have NCO averages higher than those of *S.*

cerevisiae (46.4) and *S. paradoxus* (26.9) (Figure 2A; Tables S4 & S5; Liu et al. 2019). The

significant difference in COs per meiosis between our fermentation and natural crosses was

unexpected, as we predicted that the recombination landscape would be more similar within a

species (*S. uvaum*) than between species *S. cerevisiae* and *S. paradoxus*.

To further explore the differences in recombination landscapes between our crosses, we split

the genome into 20kb, non-overlapping windows, and obtained CO, NCO, and marker counts

for each region (Figure 2D). We corrected for differences in marker resolution between crosses

(see Methods). We found modest, but significant genomewide correlation between our crosses

for both COs (Spearman's correlation: 0.273; p<0.0001) and NCOs (Spearman's correlation:

0.1456; p<0.001). Again, these correlations were unexpectedly lower than those reported by Liu

et al. (2019) between *S. cerevisiae* and *S. paradoxus* (0.48 for COs and 0.17 for NCOs).

We hypothesized that the low correlations between crosses might be impacted by the presence

of heterozygous introgression from *S. eubayanus* in the fermentation cross. To explore this

possibility, we separated the 20kb windows into introgressed and non-introgressed windows

(based on whether they overlapped with an introgressed region). We will refer to introgression in

the fermentation cross as "introgression" and use the term "introgressed region" to refer

generally to the syntenic region, regardless of which cross we are focusing on. We calculated

Spearman's correlations of COs, NCOs, and marker counts between the two crosses for each

chromosome. We find positive correlations between CO counts for all chromosomes when

looking at non-introgressed regions (though some were not significant; Table S6). For

introgressed regions, we found no significant CO correlations between crosses, which is

consistent with the hypothesis that CO landscapes are changed in introgression (Table S7).

There was one significant, highly positive correlation among NCOs (chromosome 1), but all

other correlations were not significant. This was likely affected by the fact that markers used to

detect NCOs are very differently distributed between the crosses, and the small size of NCO

tracts means regions with more markers tend to reveal more NCOs.

We find that introgressions tend to have lower CO counts and higher NCO counts in the

fermentation cross when compared to syntenic regions in the natural cross. The fermentation

cross has fewer COs than the natural cross overall, but the difference is greater in the

introgressed regions (Figure 2B-C). To further explore and test possible explanations for the

patterns of COs and NCOs in introgressed regions, we constructed linear models for CO and

NCO counts. Our model of CO counts showed a significant positive effect of the interactions

between natural cross and introgression (whether a genomic window is from the natural cross

and whether that window is in an introgressed region), as well as a significant negative effect of

introgression on the number of COs, and a significant positive effect of GC content on CO count

(Table 3). These results are consistent with increased COs in GC-rich regions and reduced

COs in introgressions. Our linear model of NCO counts showed a similar positive effect of GC

content on NCO counts, but showed opposite significant coefficients for the other two

explanatory variables (Table 4). This indicates that GC content still plays an important role in

localizing NCOs, and supports our findings that patterns of NCOs in introgressed regions are

opposite to those of COs.

Reduced sequence homology helps explain non-crossover resolution of DSBs in introgressions

One possible explanation for an increase of NCOs in introgressions is that the reduced

homology is biasing DSBs in the region to be resolved as NCOs rather than COs. We would

therefore expect to see NCOs to be negatively correlated with homology. To evaluate the

relationship of homology to the CO and NCO landscapes in introgressions, we measured

homology, NCO depth, and CO count in 101 bp sliding windows along each of the

introgressions. Mismatch repair proteins in *Saccharomyces* seem to suppress COs with very

little mismatch in small regions (~350 bp), which informed our window size (Chen & Jinks-

Robertson, 1999; Datta et al., 1997). We counted the number of NCO tracts that intersect with

each window as a measurement of NCO depth, and simply counted the CO events in a given

window. We measured homology in the sliding windows by measuring the proportion of bases

that are expected to match between the two parent strains (Figure 3). We then ran Spearman's

correlations and a loess regression along each introgression and found a weak, but often

significant ($p \ll 0.001$) correlation between NCOs and homology in the introgression (Table 1),

suggesting that resolution of double strand breaks is biased towards NCOs when homology is

low. From the loess regression, we can observe an increase in NCO as homology reduces until

about 0.9-0.8 homology, at which point NCOs level out or reduce. However, this effect is very

weak with respect to the NCO counts, and at low levels of homology the uncertainty of the

regression line is very large. This is primarily driven by the number of windows with no NCOs

(Figure 4).

The low CO count in introgressed regions leaves us unable to investigate effects of homology

on CO counts except on chromosomes 7 and 14, where we found significantly higher homology

around COs than around NCOs (Table 2). The introgressions on these two chromosomes are

unique in that they contain a highly homologous portion of sequence and therefore contain

enough COs for us to have power to detect differences between CO and NCO neighborhoods

(Figure 3).

Introgression decreases allele shuffling locally and at the chromosome level

Because NCOs still play a small role in shuffling alleles along the chromosome, we were

interested in whether the increase in NCOs of the fermentation cross would supplement the lost

shuffling from the suppression of COs in the introgressions. To test this hypothesis, we used the

measure r, which accounts for the number and positioning of recombination events to estimate

the probability that a randomly chosen pair of loci shuffles their alleles in a gamete (Veller et al.,

2019). We calculated the average r per chromosome and for each introgressed region for each

of the two crosses. We observed high levels of shuffling at the chromosome level when

compared to humans. The intra-chromosomal component of r in humans is 0.0135 in females

and 0.0177 in males (Veller et al., 2019), while our measurements for chromosomes varied

between 0.216 and 0.426. We find that most chromosomes do not have a significantly different

amount of allele shuffling between the two crosses, even though the natural cross generally has

more COs (Table S8; Bonferroni-adjusted $\alpha = 0.00313$). However, of the six chromosomes with

significantly different r values, all of them showed more shuffling in the natural cross, and five of

the six (chromosomes 4, 9, 10, 14, and 15) contained introgressed regions (Figure 5).

Chromosome 12 was the only chromosome without an introgressed region to have significantly

different shuffling between crosses, and it also showed more shuffling in the natural cross. All of

the introgressed regions showed significantly more shuffling in the natural cross, indicating that

the large increase of NCOs in the introgressions does not make up for the loss of shuffling from

the depletion of COs (Table S9; Bonferroni-adjusted $\alpha = 0.00556$). This finding indicates that an

introgression that is segregating in a population will incur a shuffling cost in heterozygous

individuals on top of any other evolutionary effects the introgression may have.

Discussion

Our study is motivated by understanding recombination rate variation within a species and

uncovering potential genetic factors underlying this variation. To investigate this question, we

crossed two pairs of *S. uvarum* strains, one pair isolated from natural environments and one

pair from fermentation environments, and explored the distribution of CO and NCO events from

both crosses. We found a similar range of COs per meiosis as previous studies in *S. cerevisiae*

and *S. paradoxus* (Liu et al., 2019; Mancera et al., 2008). The number of NCOs detected,

however, was considerably higher, despite having less sequence differences between strains in

each of our crosses and therefore lower genomic resolution for detection. We also detected

more COs and fewer NCOs in our natural cross when compared to our fermentation cross.

We hypothesized that these differences in the recombination landscape were influenced by

introgression, given that heterozygous introgression creates sequence divergence, and that

COs in regions of lower homology are known to be curtailed (Chen & Jinks-Robertson 1999, Li

et al. 2006, Weltz-Voegele & Jinks-Robertson 2008, Cooper et al. 2021). We therefore explored

the relationship between introgressions and the differences in CO and NCO counts between

crosses. Our correlations between crosses, though mostly not significant, pointed to more CO

distribution differences in introgressed regions than in the rest of the genome. When we then

modeled CO and NCO locations, correcting for GC content (a well characterized driver of

recombination events (Kiktev et al., 2018; Marsolier-Kergoat & Yeramian, 2009)), we found that

the distribution of COs and NCOs we observed was well explained by introgressions. While we

are limited in our interpretations by only comparing two crosses (one cross with heterozygous

introgression and one without introgression), these results are in line with findings in inversions,

where heterozygotes show sharp decreases in COs and an increase in NCOs in the inverted

region (Crown et al., 2018; Korunes & Noor, 2019). However, unlike heterozygous inversions

where an increase in COs is observed on freely recombining chromosomes (the inter-

chromosomal effect), we do not see an increase in COs outside of introgression or on

chromosomes without introgression.

One likely effect of this CO reduction is a reduction in shuffling at the regional and chromosomal

level. While NCOs can increase local shuffling, they likely have a much weaker effect on the

likelihood of two random alleles being shuffled than COs do. We find this is the case for our two

crosses, where despite a large number of NCOs in introgressions, the amount of shuffling (as

measured by r) is significantly lower in the fermentation cross. This loss of shuffling translates

to frequently lower r in the fermentation cross at the chromosome level for chromosomes

containing introgressions. Lower r is not observed when introgressions are small and near

telomeres, while even a small introgression near the center of the chromosome can lead to a

large reduction in r (as is the case for chromosome 15). This is consistent with the expectation

that COs near the center of chromosomes generate much more shuffling of alleles than terminal

COs (Veller et al., 2019). Our findings indicate that reducing COs, especially near the center of

chromosomes, has a cost to shuffling that is not compensated by the increase of NCOs that we

observe. If the benefit of recombination is its ability to generate new combinations of alleles,

then the loss of shuffling resulting from being heterozygous for divergent DNA sequences may

come at an additional cost beyond the possibilities of genetic incompatibilities between

hybridizing species. This cost is likely higher as divergence increases and as the length of

divergent sequences is greater, as is the case with early generation hybrids (Dagilis & Matute,

2023). Ultimately, if sequence divergence is too high, the resultant failure to recombine can

become a postzygotic reproductive barrier (Bozdag et al., 2021; Hunter et al., 1996; Rogers et

al., 2018).

The shuffling cost to introgression that we identify in our crosses may play an important role in

the fate of introgression in the generations following hybridization. When heterozygotes for an

introgression are formed, the reduction in shuffling inside the introgression will increase the

likelihood that the introgression is purged from the population. This is because it will likely be

inherited in its entirety and will carry the fitness cost of incompatibilities combined with a cost of

shuffling. This cost is incurred because the reduction of COs in the introgression will reduce

shuffling of alleles on either side of it and will vary in its intensity depending on the location and

size of the introgression. In generations immediately following hybridization, introgressions will

be much larger and are therefore expected to be more costly (although this likely depends on a

number of factors including time since divergence). These predictions are consistent with

modeling and empirical data on the purging of introgression in Drosophila and humans in the

first generations following hybridization (Veller et al., 2023).

As to longer term dynamics of recombination and selection, we predict that the excess NCOs

detected in heterozygous introgressions should begin to erode the divergence between the

sequences, increasing homology and slowly reducing the cost of the introgression. This

hypothesis posits that recombination can act to remove the larger, more deleterious regions of

an introgression quickly while whittling away slightly deleterious alleles that may be linked to any

beneficial regions of an introgression. However, in our current study, we are limited by only

observing one generation of sexual reproduction between two pairs of strains. This means that

we don't capture longer term patterns of recombination or the landscape of recombination in

introgressions that are segregating in a population. Consequently, our findings do not reflect the

effects of long-term selection (and hypothesized degradation of sequence divergence through

NCO) that would lead to introgressions preferentially remaining in high-CO regions, as has been

observed in other organisms. Furthermore, *Saccharomyces* typically reproduce asexually, with

only infrequent sexual cycles (Magwene et al., 2011; Ruderfer et al., 2006; Zeyl & Otto, 2007).

When they do mate, they often mate within a tetrad resulting in increased homozygosity. For

example, each diploid progenitor of the parents of our fermentation cross was homozygous for

introgression across the genome, meaning that recombination would neither break up nor aid in

purging the introgression in isolated populations of each parent. This suggests that the fate of

introgressions in this species is perhaps more loosely tied to recombination patterns than it

would be in an obligately sexually reproducing species.

Despite some limitations to interpretation, this study provides a unique view of the early

dynamics of hybridization and the role of recombination in the presence of introgression. By

focusing not only on the distribution of recombination events but on their specific role in shuffling

alleles, we can more closely connect the physical process of recombination to its role among

other evolutionary forces.

Acknowledgements

We are grateful to members of the Heil lab, Mohamed Noor, Nathan Layman, and Mark

Smithson for comments on this manuscript. We thank Chris Hittinger and the Portuguese Yeast

Culture Collection for *S. uvarum* strains. This work was supported by NIH R35GM142849 to

CSH.

Figures and Tables

Figure 1: (A) Visual representation of each of our crosses. Yeast from each of the parental

strains of a cross are induced to go through meiosis to generate haploids of each mating type.

Subsequently, they are mated, and their diploid offspring are induced to enter meiosis. The

resulting tetrads are manually dissected, and each haploid meiotic product is grown mitotically

to obtain enough material for DNA extraction and whole genome sequencing. We then call

SNPs on the resulting sequences and retain loci with fixed differences between parents. These

loci are then coded as 1 or 0 depending on the parent of origin and the CrossOver software

detects COs and NCOs. (B) The introgressions we observe in our crosses are due to S.

eubayanus *hybridizing with* S. uvarum*, resulting in F1 hybrids that then potentially crossed with*

other S. uvarum *individuals for some number of generations. Eventually, the* S. eubayanus

ancestry was degraded in the population of S. uvarum *until the introgressions we observe today*

remained, potentially segregating in the population. A similar process likely happened in each of

the parental strains we utilized, but with different introgressions remaining in each strain. We

then crossed individuals from each strain that were homozygous for the introgression, resulting

in offspring that were heterozygous for each introgression. It's important to note that due to the

life cycle of Saccharomyces*, mitotic recombination likely played an important role in the*

breakdown of introgressions.

Figure 2: (A) Barplot depicting the number of COs and NCOs detected per meiosis in S. uvarum

crosses (green: natural cross; pink: fermentation cross). The error bars represent the standard

534 *error around the mean. These values are not corrected by resolution. The counts for* S.

535 paradoxus *and* S. cerevisiae *are represented by arrows and taken from Liu et al. (2019). (B)*

536 *Mean CO/kb and (C) NCO/kb by cross and introgression (0 denotes intervals without*

537 *introgression; 1 denotes introgression present in the fermentation cross. While the natural cross*

538 *does not contain introgression, the region where introgression is present in the fermentation*

539 *cross was compared to its syntenic region in the natural cross). Error bars represent the*

540 *standard error around the mean. (D)* S. uvarum *chromosomes split into 20kb, non-overlapping*

541 *windows. CO, NCO, and SNP counts are reported for both crosses (fermentation and natural).*

542 *Shaded regions denote introgressed regions. CO counts are smoothed when the true location of*

543 *the CO split could be in one of multiple windows. NCO counts are corrected for marker*

544 *resolution.*

Figure 3: CO, NCO, and homology in 101bp sliding windows of fermentation cross introgressed

regions. CO counts are shown in blue, the depth of NCO tracts are shown in orange, and the

proportion of expected homologous bases between the two fermentation strains is shown in

black.

Figure 4: Average r for each chromosome (A) and for each introgressed region (B). Asterisks

indicate a significant difference in chromosome r-bar between crosses. All introgressed regions

had a significant difference in r. Error bars indicate standard error around the mean.

Table 1: Spearman's correlations of NCOs to homology in introgressed regions of the

fermentation cross.

Chromosome	Start	End	Corr	p<0.001?
4	866500	983774	-0.1562	TRUE
6	1	65500	-0.0409	FALSE
7	1	53500	-0.2102	TRUE
9	158500	298500	-0.1313	TRUE
10	234500	288500	-0.0236	FALSE
10	301500	428500	-0.0990	TRUE
13	26500	103500	-0.1943	TRUE
14	18500	586500	-0.2004	TRUE
15	367500	434500	-0.2076	TRUE

567   *Table 2: Welch two sample t-test results for differences in homology between CO-adjacent*

568   *regions and NCO-adjacent regions per introgression.*

Introgression	CO mean homology	CO SE	NCO mean homology	NCO SE	p-value
Chromosome 4	0.9467	0.0148	0.9248	0.0028	0.2760
Chromosome 6	0.9663	0.0203	0.9784	0.0017	0.5360
Chromosome 7	1.0000	0.0000	0.9838	0.0013	<0.0001
Chromosome 9	0.9650	0.0102	0.9312	0.0036	0.0109
Chromosome 10 (1)	0.9671	0.0092	0.9476	0.0058	0.1008
Chromosome 10 (2)	0.9400	0.0074	0.9276	0.0040	0.1991
Chromosome 13	0.9700	0.0161	0.9304	0.0024	0.1296
Chromosome 14	0.9893	0.0026	0.9415	0.0018	<0.0001
Chromosome 15	0.9317	0.0093	0.9252	0.0029	0.5648

579 *Table 3: Coefficients of gaussian generalized linear model modeling CO counts per 20kb*

580 *window.*

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	-8.9355	1.4713	-6.0732	1.70e-09
Introgression	-3.6343	0.6293	-5.7755	9.84e-09
GC	37.9733	3.7037	10.2528	< 2e-16
Introgression:crossnatural	5.2378	0.8595	6.0941	1.49e-09

584 *Table 4: Coefficients of gaussian generalized linear model modeling NCO counts per 20kb*

585 *window.*

	Estimate	Std. Error	t value	Pr(>|t|)
(Intercept)	-1.2261	1.4013	-0.8749	0.3818
Introgression	9.6354	0.5993	16.0770	<2e-16
GC	8.3792	3.5275	2.3754	0.0177
Introgression:crossnatural	-9.6414	0.8186	-11.7780	<2e-16

References

Aguillon, S. M., Dodge, T. O., Preising, G. A., & Schumer, M. (2022). Introgression. *Current Biology*, *32*(16), R865–R868. https://doi.org/10.1016/j.cub.2022.07.004

Albertin, W., Chernova, M., Durrens, P., Guichoux, E., Sherman, D. J., Masneuf-Pomarede, I., & Marullo, P. (2018). Many interspecific chromosomal introgressions are highly prevalent in Holarctic Saccharomyces uvarum strains found in human-related fermentations. *Yeast*, *35*(1), 141–156. https://doi.org/10.1002/yea.3248

Almeida, P., Gonçalves, C., Teixeira, S., Libkind, D., Bontrager, M., Masneuf-Pomarède, I., Albertin, W., Durrens, P., Sherman, D. J., Marullo, P., Todd Hittinger, C., Gonçalves, P., & Sampaio, J. P. (2014). A Gondwanan imprint on global diversity and domestication of wine and cider yeast *Saccharomyces uvarum*. *Nature Communications*, *5*, 4044. https://doi.org/10.1038/ncomms5044

Anderson, C. M., Chen, S. Y., Dimon, M. T., Oke, A., DeRisi, J. L., & Fung, J. C. (2011). ReCombine: A Suite of Programs for Detection and Analysis of Meiotic Recombination in Whole-Genome Datasets. *PLoS ONE*, *6*(10). https://doi.org/10.1371/journal.pone.0025509

Arter, M., & Keeney, S. (2023). Divergence and conservation of the meiotic recombination machinery. *Nature Reviews Genetics*, 1–17. https://doi.org/10.1038/s41576-023-00669-8

Auton, A., Li, Y. R., Kidd, J., Oliveira, K., Nadel, J., Holloway, J. K., Hayward, J. J., Cohen, P. E., Greally, J. M., Wang, J., Bustamante, C. D., & Boyko, A. R. (2013). Genetic Recombination Is Targeted towards Gene Promoter Regions in Dogs. *PLOS Genetics*, *9*(12), e1003984. https://doi.org/10.1371/journal.pgen.1003984

Barton, N., & Bengtsson, B. O. (1986). The barrier to genetic exchange between hybridising populations. *Heredity*, *57*(3), Article 3. https://doi.org/10.1038/hdy.1986.135

Begun, D. J., & Aquadro, C. F. (1992). Levels of naturally occurring DNA polymorphism correlate with recombination rates in D. melanogaster. *Nature, 356*(6369), 519–520. https://doi.org/10.1038/356519a0

Bendixsen, D. P., Frazão, J. G., & Stelkens, R. (2022). Saccharomyces yeast hybrids on the rise. *Yeast, 39*(1–2), 40–54. https://doi.org/10.1002/yea.3684

Bozdag, G. O., Ono, J., Denton, J. A., Karakoc, E., Hunter, N., Leu, J.-Y., & Greig, D. (2021). Breaking a species barrier by enabling hybrid recombination. *Current Biology, 31*(4), R180–R181. https://doi.org/10.1016/j.cub.2020.12.038

Brand, C. L., Cattani, M. V., Kingan, S. B., Landeen, E. L., & Presgraves, D. C. (2018). Molecular Evolution at a Meiosis Gene Mediates Species Differences in the Rate and Patterning of Recombination. *Current Biology: CB, 28*(8), 1289-1295.e4. https://doi.org/10.1016/j.cub.2018.02.056

Brandvain, Y., Kenney, A. M., Flagel, L., Coop, G., & Sweigart, A. L. (2014). Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. *PLOS Genetics, 10*(6), e1004410. https://doi.org/10.1371/journal.pgen.1004410

Brion, C., Legrand, S., Peter, J., Caradec, C., Pflieger, D., Hou, J., Friedrich, A., Llorente, B., & Schacherer, J. (2017). Variation of the meiotic recombination landscape and properties over a broad evolutionary distance in yeasts. *PLOS Genetics, 13*(8), e1006917. https://doi.org/10.1371/journal.pgen.1006917

Butlin, R. K. (2005). Recombination and speciation. *Molecular Ecology, 14*(9), 2621–2635. https://doi.org/10.1111/j.1365-294X.2005.02617.x

Calfee, E., Gates, D., Lorant, A., Perkins, M. T., Coop, G., & Ross-Ibarra, J. (2021). Selective sorting of ancestral introgression in maize and teosinte along an elevational cline. *bioRxiv*, 2021.03.05.434040. https://doi.org/10.1101/2021.03.05.434040

Charlesworth, B., Morgan, M. T., & Charlesworth, D. (1993). The effect of deleterious mutations on neutral molecular variation. *Genetics, 134*(4), 1289–1303.

https://doi.org/10.1093/genetics/134.4.1289

Chen, W., & Jinks-Robertson, S. (1999). The role of the mismatch repair machinery in

regulating mitotic and meiotic recombination between diverged sequences in yeast.

Genetics, *151*(4), 1299–1313.

Choi, K., Zhao, X., Tock, A. J., Lambing, C., Underwood, C. J., Hardcastle, T. J., Serra, H., Kim,

J., Cho, H. S., Kim, J., Ziolkowski, P. A., Yelina, N. E., Hwang, I., Martienssen, R. A., &

Henderson, I. R. (2018). Nucleosomes and DNA methylation shape meiotic DSB

frequency in Arabidopsis thaliana transposons and gene regulatory regions. *Genome*

Research, *28*(4), 532–546. https://doi.org/10.1101/gr.225599.117

Chovnick, A., Ballantyne, G. H., & Holm, D. G. (1971). Studies on gene conversion and its

relationship to linked exchange in Drosophila melanogaster. *Genetics*, *69*(2), 179–209.

https://doi.org/10.1093/genetics/69.2.179

Cooper, T. J., Crawford, M. R., Hunt, L. J., Marsolier-Kergoat, M.-C., Llorente, B., & Neale, M. J.

(2021). *Mismatch repair disturbs meiotic class I crossover control* (p. 480418). bioRxiv.

https://doi.org/10.1101/480418

Crown, K. N., Miller, D. E., Sekelsky, J., & Hawley, R. S. (2018). Local Inversion Heterozygosity

Alters Recombination throughout the Genome. *Current Biology: CB*, *28*(18), 2984-

2990.e3. https://doi.org/10.1016/j.cub.2018.07.004

Cubillos, F. A., Billi, E., Zörgö, E., Parts, L., Fargier, P., Omholt, S., Blomberg, A., Warringer, J.,

Louis, E. J., & Liti, G. (2011). Assessing the complex architecture of polygenic traits in

diverged yeast populations. *Molecular Ecology*, *20*(7), 1401–1413.

https://doi.org/10.1111/j.1365-294X.2011.05005.x

Dagilis, A. J., & Matute, D. R. (2023). The fitness of an introgressing haplotype changes over

the course of divergence and depends on its size and genomic location. *PLOS Biology*,

21(7), e3002185. https://doi.org/10.1371/journal.pbio.3002185

Danecek, P., Bonfield, J. K., Liddle, J., Marshall, J., Ohan, V., Pollard, M. O., Whitwham, A.,

Keane, T., McCarthy, S. A., Davies, R. M., & Li, H. (2021). Twelve years of SAMtools and BCFtools. *GigaScience*, *10*(2), giab008. https://doi.org/10.1093/gigascience/giab008

D'Angiolo, M., De Chiara, M., Yue, J.-X., Irizar, A., Stenberg, S., Persson, K., Llored, A., Barré, B., Schacherer, J., Marangoni, R., Gilson, E., Warringer, J., & Liti, G. (2020). A yeast living ancestor reveals the origin of genomic introgressions. *Nature*, *587*(7834), Article 7834. https://doi.org/10.1038/s41586-020-2889-1

Dapper, A. L., & Payseur, B. A. (2019). Molecular evolution of the meiotic recombination pathway in mammals. *Evolution*, *73*(12). https://doi.org/10.1111/evo.13850

Datta, A., Hendrix, M., Lipsitch, M., & Jinks-Robertson, S. (1997). Dual roles for DNA sequence identity and the mismatch repair system in the regulation of mitotic crossing-over in yeast. *Proceedings of the National Academy of Sciences of the United States of America*, *94*(18), 9757–9762.

de Massy, B. (2013). Initiation of Meiotic Recombination: How and Where? Conservation and Specificities Among Eukaryotes. *Annual Review of Genetics*, *47*(1), 563–599. https://doi.org/10.1146/annurev-genet-110711-155423

Duranton, M., & Pool, J. E. (2022). Interactions Between Natural Selection and Recombination Shape the Genomic Landscape of Introgression. *Molecular Biology and Evolution*, *39*(7), msac122. https://doi.org/10.1093/molbev/msac122

Edelman, N. B., Frandsen, P. B., Miyagi, M., Clavijo, B., Davey, J., Dikow, R. B., García-Accinelli, G., Van Belleghem, S. M., Patterson, N., Neafsey, D. E., Challis, R., Kumar, S., Moreira, G. R. P., Salazar, C., Chouteau, M., Counterman, B. A., Papa, R., Blaxter, M., Reed, R. D., … Mallet, J. (2019). Genomic architecture and introgression shape a butterfly radiation. *Science*, *366*(6465), 594–599. https://doi.org/10.1126/science.aaw2090

Felsenstein, J. (1974). The evolutionary advantage of recombination. *Genetics*, *78*(2), 737–756.

Gerton, J. L., DeRisi, J., Shroff, R., Lichten, M., Brown, P. O., & Petes, T. D. (2000). Global

696   mapping of meiotic recombination hotspots and coldspots in the yeast Saccharomyces

697   cerevisiae. *Proceedings of the National Academy of Sciences*, *97*(21), 11383–11390.

698   https://doi.org/10.1073/pnas.97.21.11383

699   Harfe, B. D., & Jinks-Robertson, S. (2000). Dna Mismatch Repair and Genetic Instability.

700   *Annual Review of Genetics*, *34*(1), 359–399.

701   https://doi.org/10.1146/annurev.genet.34.1.359

702   He, Y., Wang, M., Dukowic-Schulze, S., Zhou, A., Tiang, C.-L., Shilo, S., Sidhu, G. K., Eichten,

703   S., Bradbury, P., Springer, N. M., Buckler, E. S., Levy, A. A., Sun, Q., Pillardy, J.,

704   Kianian, P. M. A., Kianian, S. F., Chen, C., & Pawlowski, W. P. (2017). Genomic

705   features shaping the landscape of meiotic double-strand-break hotspots in maize.

706   *Proceedings of the National Academy of Sciences*, *114*(46), 12231–12236.

707   https://doi.org/10.1073/pnas.1713225114

708   Hill, W. G., & Robertson, A. (1966). The effect of linkage on limits to artificial selection.

709   *Genetical Research*, *8*(3), 269–294.

710   Hilliker, A. J., Harauz, G., Reaume, A. G., Gray, M., Clark, S. H., & Chovnick, A. (1994). Meiotic

711   gene conversion tract length distribution within the rosy locus of Drosophila

712   melanogaster. *Genetics*, *137*(4), 1019–1026.

713   https://doi.org/10.1093/genetics/137.4.1019

714   Hoffman, C. S., & Winston, F. (1987). A ten-minute DNA preparation from yeast efficiently

715   releases autonomous plasmids for transformation of Escherichia coli. *Gene*, *57*(2–3),

716   267–272. https://doi.org/10.1016/0378-1119(87)90131-4

717   Hunter, N., Chambers, S. R., Louis, E. J., & Borts, R. H. (1996). The mismatch repair system

718   contributes to meiotic sterility in an interspecific yeast hybrid. *The EMBO Journal*, *15*(7),

719   1726–1733. https://doi.org/10.1002/j.1460-2075.1996.tb00518.x

720   Jeffreys, A. J., & May, C. A. (2004). Intense and highly localized gene conversion activity in

721   human meiotic crossover hot spots. *Nature Genetics*, *36*(2), Article 2.

https://doi.org/10.1038/ng1287

Judd, S. R., & Petes, T. D. (1988). Physical Lengths of Meiotic and Mitotic Gene Conversion
Tracts in Saccharomyces Cerevisiae. *Genetics*, *118*(3), 401.
https://doi.org/10.1093/genetics/118.3.401

Keeney, S. (2001). Mechanism and control of meiotic recombination initiation. *Current Topics in
Developmental Biology*, *52*, 1–53. https://doi.org/10.1016/s0070-2153(01)52008-6

Kiktev, D. A., Sheng, Z., Lobachev, K. S., & Petes, T. D. (2018). GC content elevates mutation
and recombination rates in the yeast *Saccharomyces cerevisiae*. *Proceedings of the
National Academy of Sciences*, *115*(30), E7109–E7118.
https://doi.org/10.1073/pnas.1807334115

Kong, A., Thorleifsson, G., Stefansson, H., Masson, G., Helgason, A., Gudbjartsson, D. F.,
Jonsdottir, G. M., Gudjonsson, S. A., Sverrisson, S., Thorlacius, T., Jonasdottir, A.,
Hardarson, G. A., Palsson, S. T., Frigge, M. L., Gulcher, J. R., Thorsteinsdottir, U., &
Stefansson, K. (2008). Sequence Variants in the RNF212 Gene Associate with Genome-
Wide Recombination Rate. *Science*, *319*(5868), 1398–1401.
https://doi.org/10.1126/science.1152422

Korunes, K. L., & Noor, M. A. F. (2017). Gene conversion and linkage: Effects on genome
evolution and speciation. *Molecular Ecology*, *26*(1), 351–364.
https://doi.org/10.1111/mec.13736

Korunes, K. L., & Noor, M. A. F. (2019). Pervasive gene conversion in chromosomal inversion
heterozygotes. *Molecular Ecology*, *28*(6), 1302–1315. https://doi.org/10.1111/mec.14921

Lam, I., & Keeney, S. (2015). Nonparadoxical evolutionary stability of the recombination
initiation landscape in yeast. *Science*, *350*(6263), 932–937.
https://doi.org/10.1126/science.aad0814

Langdon, Q. K., Peris, D., Baker, E. P., Opulente, D. A., Nguyen, H.-V., Bond, U., Gonçalves,
P., Sampaio, J. P., Libkind, D., & Hittinger, C. T. (2019). Fermentation innovation

through complex hybridization of wild and domesticated yeasts. *Nature Ecology &*

Evolution, 1–11. https://doi.org/10.1038/s41559-019-0998-8

Langdon, Q. K., Peris, D., Eizaguirre, J. I., Opulente, D. A., Buh, K. V., Sylvester, K., Jarzyna,

M., Rodríguez, M. E., Lopes, C. A., Libkind, D., & Hittinger, C. T. (2020). Postglacial

migration shaped the genomic diversity and global distribution of the wild ancestor of

lager-brewing hybrids. *PLOS Genetics*, *16*(4), e1008680.

https://doi.org/10.1371/journal.pgen.1008680

Layer, R. M., Chiang, C., Quinlan, A. R., & Hall, I. M. (2014). LUMPY: A probabilistic framework

for structural variant discovery. *Genome Biology*, *15*(6), R84. https://doi.org/10.1186/gb-

2014-15-6-r84

Li, H., & Durbin, R. (2009). Fast and accurate short read alignment with Burrows-Wheeler

transform. *Bioinformatics (Oxford, England)*, *25*(14), 1754–1760.

https://doi.org/10.1093/bioinformatics/btp324

Li, L., Jean, M., & Belzile, F. (2006). The impact of sequence divergence and DNA mismatch

repair on homeologous recombination in Arabidopsis. *The Plant Journal: For Cell and*

Molecular Biology, *45*(6), 908–916. https://doi.org/10.1111/j.1365-313X.2006.02657.x

Liu, H., Huang, J., Sun, X., Li, J., Hu, Y., Yu, L., Liti, G., Tian, D., Hurst, L. D., & Yang, S.

(2018). Tetrad analysis in plants and fungi finds large differences in gene conversion

rates but no GC bias. *Nature Ecology & Evolution*, *2*(1), Article 1.

https://doi.org/10.1038/s41559-017-0372-7

Liu, H., Maclean, C. J., & Zhang, J. (2019). Evolution of the Yeast Recombination Landscape.

Molecular Biology and Evolution, *36*(2), 412–422.

https://doi.org/10.1093/molbev/msy233

Magwene, P. M., Kayıkçı, Ö., Granek, J. A., Reininga, J. M., Scholl, Z., & Murray, D. (2011).

Outcrossing, mitotic recombination, and life-history trade-offs shape genome evolution in

Saccharomyces cerevisiae. *Proceedings of the National Academy of Sciences*, *108*(5),

1987–1992. https://doi.org/10.1073/pnas.1012544108

Mancera, E., Bourgon, R., Brozzi, A., Huber, W., & Steinmetz, L. M. (2008). High-resolution

mapping of meiotic crossovers and non-crossovers in yeast. *Nature*, *454*(7203), 479–

485. https://doi.org/10.1038/nature07135

Marsolier-Kergoat, M.-C., & Yeramian, E. (2009). GC Content and Recombination: Reassessing

the Causal Effects for the Saccharomyces cerevisiae Genome. *Genetics*, *183*(1), 31–38.

https://doi.org/10.1534/genetics.109.105049

Martin, S. H., Davey, J. W., Salazar, C., & Jiggins, C. D. (2019). Recombination rate variation

shapes barriers to introgression across butterfly genomes. *PLOS Biology*, *17*(2),

e2006288. https://doi.org/10.1371/journal.pbio.2006288

McDonald, M. J., Rice, D. P., & Desai, M. M. (2016). Sex speeds adaptation by altering the

dynamics of molecular evolution. *Nature*, *531*(7593), 233–236.

https://doi.org/10.1038/nature17143

McGaugh, S. E., Heil, C. S. S., Manzano-Winkler, B., Loewe, L., Goldstein, S., Himmel, T. L., &

Noor, M. A. F. (2012). Recombination Modulates How Selection Affects Linked Sites in

Drosophila. *PLOS Biology*, *10*(11), e1001422.

https://doi.org/10.1371/journal.pbio.1001422

McKenna, A., Hanna, M., Banks, E., Sivachenko, A., Cibulskis, K., Kernytsky, A., Garimella, K.,

Altshuler, D., Gabriel, S., Daly, M., & DePristo, M. A. (2010). The Genome Analysis

Toolkit: A MapReduce framework for analyzing next-generation DNA sequencing data.

Genome Research, *20*(9), 1297–1303. https://doi.org/10.1101/gr.107524.110

Miller, D. E., Smith, C. B., Kazemi, N. Y., Cockrell, A. J., Arvanitakis, A. V., Blumenstiel, J. P.,

Jaspersen, S. L., & Hawley, R. S. (2016). Whole-Genome Analysis of Individual Meiotic

Events in Drosophila melanogaster Reveals That Noncrossover Gene Conversions Are

Insensitive to Interference and the Centromere Effect. *Genetics*, *203*(1), 159–171.

https://doi.org/10.1534/genetics.115.186486

Moran, B. M., Payne, C., Langdon, Q., Powell, D. L., Brandvain, Y., & Schumer, M. (2021). The
genomic consequences of hybridization. *eLife*, *10*, e69016.
https://doi.org/10.7554/eLife.69016

Murdoch, B., Owen, N., Shirley, S., Crumb, S., Broman, K. W., & Hassold, T. (2010). Multiple
loci contribute to genome-wide recombination levels in male mice. *Mammalian Genome*,
21(11), 550–555. https://doi.org/10.1007/s00335-010-9303-5

Nachman Michael W. & Payseur Bret A. (2012). Recombination rate variation and speciation:
Theoretical predictions and empirical results from rabbits and mice. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *367*(1587), 409–421.
https://doi.org/10.1098/rstb.2011.0249

Nespolo, R. F., Villarroel, C. A., Oporto, C. I., Tapia, S. M., Vega-Macaya, F., Urbina, K., Chiara,
M. D., Mozzachiodi, S., Mikhalev, E., Thompson, D., Larrondo, L. F., Saenz-Agudelo, P.,
Liti, G., & Cubillos, F. A. (2020). An Out-of-Patagonia migration explains the worldwide
diversity and distribution of Saccharomyces eubayanus lineages. *PLOS Genetics*, *16*(5),
e1008777. https://doi.org/10.1371/journal.pgen.1008777

Pan, J., Sasaki, M., Kniewel, R., Murakami, H., Blitzblau, H. G., Tischfield, S. E., Zhu, X., Neale,
M. J., Jasin, M., Socci, N. D., Hochwagen, A., & Keeney, S. (2011). A Hierarchical
Combination of Factors Shapes the Genome-wide Topography of Yeast Meiotic
Recombination Initiation. *Cell*, *144*(5), 719–731.
https://doi.org/10.1016/j.cell.2011.02.009

Pool, J. E. (2015). The Mosaic Ancestry of the Drosophila Genetic Reference Panel and the D.
melanogaster Reference Genome Reveals a Network of Epistatic Fitness Interactions.
Molecular Biology and Evolution, *32*(12), 3236–3251.
https://doi.org/10.1093/molbev/msv194

Raffoux, X., Bourge, M., Dumas, F., Martin, O. C., & Falque, M. (2018). Role of Cis, Trans, and
Inbreeding Effects on Meiotic Recombination in Saccharomyces cerevisiae. *Genetics*,

210(4), 1213–1226. https://doi.org/10.1534/genetics.118.301644

Ravinet, M., Yoshida, K., Shigenobu, S., Toyoda, A., Fujiyama, A., & Kitano, J. (2018). The

genomic landscape at a late stage of stickleback speciation: High genomic divergence

interspersed by small localized regions of introgression. *PLOS Genetics*, *14*(5),

e1007358. https://doi.org/10.1371/journal.pgen.1007358

Robinson, J. T., Thorvaldsdóttir, H., Winckler, W., Guttman, M., Lander, E. S., Getz, G., &

Mesirov, J. P. (2011). Integrative genomics viewer. *Nature Biotechnology*, *29*(1), 24–26.

https://doi.org/10.1038/nbt.1754

Rockman, M. V., & Kruglyak, L. (2009). Recombinational Landscape and Population Genomics

of Caenorhabditis elegans. *PLOS Genetics*, *5*(3), e1000419.

https://doi.org/10.1371/journal.pgen.1000419

Rogers, D. W., McConnell, E., Ono, J., & Greig, D. (2018). Spore-autonomous fluorescent

protein expression identifies meiotic chromosome mis-segregation as the principal cause

of hybrid sterility in yeast. *PLOS Biology*, *16*(11), e2005066.

https://doi.org/10.1371/journal.pbio.2005066

Ruderfer, D. M., Pratt, S. C., Seidel, H. S., & Kruglyak, L. (2006). Population genomic analysis

of outcrossing and recombination in yeast. *Nature Genetics*, *38*(9), Article 9.

https://doi.org/10.1038/ng1859

Scannell, D. R., Zill, O. A., Rokas, A., Payen, C., Dunham, M. J., Eisen, M. B., Rine, J.,

Johnston, M., & Hittinger, C. T. (2011). The Awesome Power of Yeast Evolutionary

Genetics: New Genome Sequences and Strain Resources for the Saccharomyces sensu

stricto Genus. *G3: Genes, Genomes, Genetics*, *1*(1), 11–25.

https://doi.org/10.1534/g3.111.000273

Schaeffer, S. W., & Anderson, W. W. (2005). Mechanisms of Genetic Exchange Within the

Chromosomal Inversions of Drosophila pseudoobscura. *Genetics*, *171*(4), 1729–1739.

https://doi.org/10.1534/genetics.105.041947

Schumer, M., Xu, C., Powell, D. L., Durvasula, A., Skov, L., Holland, C., Blazier, J. C.,

Sankararaman, S., Andolfatto, P., Rosenthal, G. G., & Przeworski, M. (2018). Natural

selection interacts with recombination to shape the evolution of hybrid genomes.

Science, *360*(6389), 656–660. https://doi.org/10.1126/science.aar3684

Shi, J., Wolf, S. E., Burke, J. M., Presting, G. G., Ross-Ibarra, J., & Dawe, R. K. (2010).

Widespread Gene Conversion in Centromere Cores. *PLoS Biology*, *8*(3), e1000327.

https://doi.org/10.1371/journal.pbio.1000327

Singhal, S., Leffler, E. M., Sannareddy, K., Turner, I., Venn, O., Hooper, D. M., Strand, A. I., Li,

Q., Raney, B., Balakrishnan, C. N., Griffith, S. C., McVean, G., & Przeworski, M. (2015).

Stable recombination hotspots in birds. *Science (New York, N.Y.)*, *350*(6263), 928–932.

https://doi.org/10.1126/science.aad0843

Smith, J. M., & Haigh, J. (1974). The hitch-hiking effect of a favourable gene. *Genetical*

Research, *23*(1), 23–35.

Smukowski, C. S., & Noor, M. a. F. (2011). Recombination rate variation in closely related

species. *Heredity*, *107*(6), 496–508. https://doi.org/10.1038/hdy.2011.44

Smukowski Heil, C. S., Ellison, C., Dubin, M., & Noor, M. A. F. (2015). Recombining without

Hotspots: A Comprehensive Evolutionary Portrait of Recombination in Two Closely

Related Species of Drosophila. *Genome Biology and Evolution*, *7*(10), 2829–2842.

https://doi.org/10.1093/gbe/evv182

Stapley, J., Feulner, P. G. D., Johnston, S. E., Santure, A. W., & Smadja, C. M. (2017).

Variation in recombination frequency and distribution across eukaryotes: Patterns and

processes. *Philosophical Transactions of the Royal Society B: Biological Sciences*,

372(1736), 20160455. https://doi.org/10.1098/rstb.2016.0455

Stelkens, R., & Bendixsen, D. P. (2022). The evolutionary and ecological potential of yeast

hybrids. *Current Opinion in Genetics & Development*, *76*, 101958.

https://doi.org/10.1016/j.gde.2022.101958

Talbert, P. B., & Henikoff, S. (2010). Centromeres Convert but Don't Cross. *PLOS Biology*, *8*(3),

879   e1000326. https://doi.org/10.1371/journal.pbio.1000326

880   Tellini, N., De Chiara, M., Mozzachiodi, S., Tattini, L., Vischioni, C., Naumova, E., Warringer, J.,

881   Bergström, A., & Liti, G. (2023). *Ancient and recent origins of shared polymorphisms in*

882   *yeas*. https://doi.org/10.21203/rs.3.rs-2573222/v1

883   Tsai, I. J., Burt, A., & Koufopanou, V. (2010). Conservation of recombination hotspots in yeast.

884   *Proceedings of the National Academy of Sciences*.

885   https://doi.org/10.1073/pnas.0908774107

886   Veller, C., Edelman, N. B., Muralidhar, P., & Nowak, M. A. (2023). Recombination and selection

887   against introgressed DNA. *Evolution*, *77*(4), 1131–1144.

888   https://doi.org/10.1093/evolut/qpad021

889   Veller, C., Kleckner, N., & Nowak, M. A. (2019). A rigorous measure of genome-wide genetic

890   shuffling that takes into account crossover positions and Mendel's second law.

891   *Proceedings of the National Academy of Sciences of the United States of America*,

892   *116*(5), 1659–1668. https://doi.org/10.1073/pnas.1817482116

893   Welz-Voegele, C., & Jinks-Robertson, S. (2008). Sequence Divergence Impedes Crossover

894   More Than Noncrossover Events During Mitotic Gap Repair in Yeast. *Genetics*, *179*(3),

895   1251–1262. https://doi.org/10.1534/genetics.108.090233

896   Wijnker, E., Velikkakam James, G., Ding, J., Becker, F., Klasen, J. R., Rawat, V., Rowan, B. A.,

897   de Jong, D. F., de Snoo, C. B., Zapata, L., Huettel, B., de Jong, H., Ossowski, S.,

898   Weigel, D., Koornneef, M., Keurentjes, J. J., & Schneeberger, K. (2013). The genomic

899   landscape of meiotic crossovers and gene conversions in Arabidopsis thaliana. *eLife*, *2*,

900   e01426. https://doi.org/10.7554/eLife.01426

901   Yang, F., Silber, S., Leu, N. A., Oates, R. D., Marszalek, J. D., Skaletsky, H., Brown, L. G.,

902   Rozen, S., Page, D. C., & Wang, P. J. (2015). TEX11 is mutated in infertile men with

903   azoospermia and regulates genome-wide recombination rates in mouse. *EMBO*

Molecular Medicine, *7*(9), 1198–1210. https://doi.org/10.15252/emmm.201404967

Zelkowski, M., Olson, M. A., Wang, M., & Pawlowski, W. (2019). Diversity and Determinants of

Meiotic Recombination Landscapes. *Trends in Genetics*, *35*(5), 359–370.

https://doi.org/10.1016/j.tig.2019.02.002

Zeyl, C. W., & Otto, S. P. (2007). A short history of recombination in yeast. *Trends in Ecology &*

Evolution, *22*(5), 223–225. https://doi.org/10.1016/j.tree.2007.02.005

912