Investigating the *cis*-Regulatory Basis of C$_3$ and C$_4$ Photosynthesis in Grasses at Single-Cell Resolution

John Pablo Mendieta[1], Xiaoyu Tu[2], Daiquan Jiang[3], Haidong Yan[1], Xuan Zhang[1], Alexandre P. Marand[1,4], Silin Zhong[3], Robert J. Schmitz[1,*]

[1]Department of Genetics, University of Georgia

[2]Joint Center for Single Cell Biology, School of Agriculture and Biology, Shanghai Jiao Tong University, Shanghai 200240, China

[3]State Key Laboratory of Agrobiotechnology, School of Life Sciences, The Chinese University of Hong Kong

Current Address

[4]Department of Molecular, Cellular, and Development Biology, University of Michigan

Abstract:

While considerable knowledge exists about the enzymes pivotal for C_4 photosynthesis, much less is known about the *cis*-regulation important for specifying their expression in distinct cell types. Here, we use single-cell-indexed ATAC-seq to identify cell-type-specific accessible chromatin regions (ACRs) associated with C_4 enzymes for five different grass species. This study spans four C_4 species, covering three distinct photosynthetic subtypes: *Zea mays* and *Sorghum bicolor* (NADP-ME), *Panicum miliaceum* (NAD-ME), *Urochloa fusca* (PEPCK), along with the C_3 outgroup *Oryza sativa*. We studied the *cis*-regulatory landscape of enzymes essential across all C_4 species and those unique to C_4 subtypes, measuring cell-type-specific biases for C_4 enzymes using chromatin accessibility data. Integrating these data with phylogenetics revealed diverse co-option of gene family members between species, showcasing the various paths of C_4 evolution. Besides promoter proximal ACRs, we found that, on average, C_4 genes have two to three distal cell-type-specific ACRs, highlighting the complexity and divergent nature of C_4 evolution. Examining the evolutionary history of these cell-type-specific ACRs revealed a spectrum of conserved and novel ACRs, even among closely related species, indicating ongoing evolution of *cis*-regulation at these C_4 loci. This study illuminates the dynamic and complex nature of CRE evolution in C_4 photosynthesis, particularly highlighting the intricate *cis*-regulatory evolution of key loci. Our findings offer a valuable resource for future investigations, potentially aiding in the optimization of C_3 crop performance under changing climatic conditions.

Introduction:

Photosynthesis is one of the most critical chemical reactions on the planet whereby CO_2 is metabolized into glucose. Plants have evolved numerous variations of photosynthesis. The most common type of photosynthesis uses the enzyme ribulose 1,5-biphosphate carboxylase oxygenase (RuBisCO) which combines CO_2 with a five carbon compound ribulose 1,5-biphosphate to create 3-phosphoglyceric acid. This three-carbon compound is then used in a redox reaction within the Calvin Benson cycle, where sucrose is made. The production of this three-carbon compound is what gives this type of photosynthesis, C_3, its name. However, although widely evolved and found in many crop plants, C_3 photosynthesis struggles to perform in hot, arid conditions. In non-ideal conditions, O2 can competitively bind the RuBisCO active site, causing the formation of a toxic intermediate, and reducing photosynthetic efficiency and plant performance (1). Due to increasing temperature caused by anthropogenic climate change, this reduction in photosynthetic capacity for key crop plants poses a major agricultural challenge (2). However, other types of photosynthesis have evolved in hotter conditions and offer a model to potentially alter key C_3 crop plants to be more efficient.

The C_4 photosynthetic pathway is an example of a modified style of photosynthesis that is able to perform in hot conditions. In brief, C_4 typically works by sequestering key photosynthetic enzymes into two different compartments in the leaf made up of different cell types. These two cell types/compartments are bundle sheath (BS) cells, which in C_4 plants generally form a concentric ring around the vasculature, and mesophyll (MS) cells, which make up large portions of the non-vascularized leaf internal cells (3). In the MS, CO_2 is imported, and converted to bicarbonate (HCO3-) by the enzyme carbonic anhydrase (CA). Bicarbonate is then converted to a four-carbon molecule oxaloacetate (OAA) by the O2-insensitive phosphoenolpyruvate carboxylase (PEPC). This OAA molecule made of a four-carbon compound (where C_4 derives its name) is finally converted into a stable metabolite, either malate or aspartate. This intermediate molecule is then transported to the BS where it undergoes a decarboxylation process, by one of three different types of decarboxylases, NAD-dependent malic enzyme (NAD-ME), NADP-dependent malic enzyme (NADP-ME), or phosphoenolpyruvate carboxykinase (PEPCK). This decarboxylation reaction releases a CO_2 molecule that enters into the Calvin Benson cycle. The generation and processing of intermediate molecules in cellular compartments allows for concentrated levels of CO_2 to interact with RuBisCO, reducing the inefficiencies mentioned above. Additional types of C_4 photosynthesis have been observed which don't rely on division of metabolites between MS and BS cell-types, but instead rely on using dimorphic chloroplast instead as in the species *Bienertia sinuspersici* (4,5). Current C_4 crops such as maize (*Zea mays*), sorghum (*Sorghum bicolor*), pearl millet (*Cenchrus americanus)*, foxtail millet (*Setaria italica*), and broomcorn millet (*Panicum miliaceum)* excel in their ability to operate in adverse conditions.

Although the evolution of C_4 photosynthesis is a complex process, there is tantalizing evidence that engineering C_3 crops to do C_4 photosynthesis might be possible. One piece of evidence that points to this is that C_4 photosynthesis has evolved independently 65 times in different lineages of plants (6). These results indicate that most plant lineages have the genetic material capable of evolving into C_4 photosynthesizers. The *Poaceae* lineage of grasses exemplifies this, as C_4 photosynthesis has evolved independently at least 18 times (7).

Interestingly, all of these species use the same core C_4 enzymes and steps, but many use
different decarboxylation enzymes as mentioned above (8–10). Furthering this hypothesis is the
fact that many C_4 related genes originally evolved from either C_3 photosynthetic genes or key
enzymes critical in core metabolism (11,12). For instance, PEPC is a key metabolism enzyme in
the glycolytic pathways of the Krebs Cycle, with some copies being important in guard cell
metabolism (13–15). Instead of novel gene content being the main driver of C_4 photosynthesis,
it's more likely due to the correct timing and compartmentalization of key enzymes into specific
cell types (16–18). This raises the question, how is gene expression of these key C_4 enzymes
regulated? Moreover, as C_4 has evolved multiple times convergently, have similar regulatory
networks and paradigms been co-opted to alter when and where these key genes are
expressed?
Cis-regulatory elements (CREs) are key players in gene regulation, as they both fine
tune expression and provide cell-type specificity (19–22). In brief, these regions operate as
binding sites for transcription factors (TFs). Transcription factors are proteins which are able to
alter transcription by binding DNA sequences and recruiting transcriptional machinery which
can either increase or decrease transcription (23). Thus TFs are able to significantly change
molecular phenotypes. Previous work has shown that CREs could be key players in the
transition to C_4 photosynthesis. This was demonstrated by taking C_4 genes from *Z. mays* and
transforming them into *Oryza sativa,* a C_3 species (24,25), which revealed that CREs from *Z.*
mays genes were able to drive cell-type-specific expression in MS in *O. sativa* (24,25).
Additional analyses have implicated CREs as drivers in the evolution of C_4 photosynthesis. In
the genus of plants *Flaveria,* which contains both C_4 and C_3 plants, one key difference in C_4
plants was a specific CRE driving gene expression in MS cells. This 41 bp motif named
Mesophyll expression module 1 is critical for cell-type-specific expression of *PEPC* in MS cells,
a critical first step in the C_4 pathway (19,26). Finally, four conserved non-coding sequences
were identified to be critical in MS-specific expression of *PEPC* in monocots (27). Furthermore,
a recent cross-species study examining the binding sites of GLK, a conserved TF regulating
photosynthetic genes, revealed that CREs can undergo rapid changes and result in diverse
gene expression patterns without the need of altering the TF itself (28). These findings show
that CREs are important genetic elements that plants use for the evolution of C_4 photosynthesis.
Although some CREs critical for cell-type-specific expression of key photosynthetic
genes have been identified, they've been restricted to those nearby the transcriptional start
sites. This is due, in part, to the challenge of identifying CREs genome wide, as well as
limitations in the isolation of BS and MS cells which is labor intensive and challenging. However,
a recent study used a multi-omic approach in *Z. mays* BS and MS cells and found CREs
genome-wide that might be critical in the cell-type-specific regulation of genes (29). One
example is the identification of a potential distal CRE ~40 kb upstream of *SULFATE*
TRANSPORTER4 (*ZmSFP4)*, a BS-specific sulfate transporter (29). These results highlight the
complexity of identifying loci involved in *cis* regulation. Identifying all CREs associated with C_4
loci is critical in enhancing our understanding of *cis* regulation of key C_4 genes, and would
greatly enhance attempts at engineering C_3 crops. During the evolution of C_4 photosynthesis, it's
unclear whether these CREs have been pre-established during evolution and co-opted for C_4
photosynthesis or if they evolved independently numerous times. Understanding the ways in

which *cis* regulation evolves to control timing and cell-type-specific expression of C_4
photosynthesis genes would greatly assist efforts in engineering C_3 plants to be more C_4 like.
125   To investigate the role of CREs and their potential contribution in controlling key C_4
genes, we used single-cell indexed Assay for Transposase Accessible Chromatin sequencing
(sciATAC-seq) to identify cell-type-specific CREs from five grass species representing diverse
C_4 subtypes, as well as an additional C_3 outgroup. We investigated the cell-type specificity of
both the core C_4 enzymes, and those which are unique to each photosynthetic subtype. Further,
we identify CREs of C_4 genes, and find previously unknown cell-type-specific CREs that might
be critical in C_4 gene expression. We find that some of these regulatory regions appear not just
conserved in a single C_4 subtype, but in all of the C_4 species we studied. Finally, we leverage
these data to find transcription factor binding motifs enriched in MS and BS cell types and use
these motifs to catalog these regulatory loci.
135

Results:

137 **Identification and Annotation of Cell Types in Diverse Species:**
138   To investigate CREs in BS and MS cells potentially important in C_4 photosynthesis, we
generated replicated sciATAC-seq libraries for four different C_4 species, comprising three
different C_4 subtypes NADP-ME (*Z. mays, S. bicolor*), NAD-ME (*Panicum miliaceum)*, and
PEPCK (*Urochloa fusca),* and a C_3 outgroup species (*O. sativa*) (**Figure 1A**). Libraries were
filtered for high-quality cells by first pseudo-bulking the sciATAC-seq libraries, and identifying
accessible chromatin regions (ACRs). Using these ACRs, per nuclei quality metrics were then
calculated such as fraction of reads in peaks, transcriptional start site enrichment, and total
integration events per nucleus (**Methods**). Nuclei found to have a high proportion of organellar
reads were also removed, with values being adjusted on a per library basis (**Methods**).
Clustering of cells was done on genomic bins, and with additional cells removed that had a high
correlation with *in-silico* generated doublets, and clusters were removed that were skewed
towards one replicate by greater than 75% (**Methods**)**.** After filtering on per nucleus quality
metrics, we identified 16,060 nuclei in *Z. mays*, 15,301 nuclei in *S. bicolor*, 7,081 nuclei in *P.*
miliaceum, 19,110 nuclei in *U. fusca,* and 5,952 nuclei in *O. sativa* (**Supplemental Figure 1,**
Supplemental Table 1).
153   Due to variation in genome size and content, cell-type annotation for each dataset was
done independently using the reference genome for each species (**Figure 1B**). We used
multiple approaches to annotate cell types. Orthologs of key marker genes from *Z. mays* and *O.*
sativa were identified using a phylogenetics based approach (**Methods**). This allowed for the
identification of marker genes for specific cell types in a cross species context. To gauge gene
activity of these marker genes, gene body chromatin accessibility was used as a proxy for
expression (**Figure 1D**) (21,30). Cell-type annotation was done manually taking into
consideration marker gene chromatin accessibility, marker enrichment in clusters, as well as
ontological relationships between cell types (**Supplemental Figure 2-19**). Due to the lack of
marker genes for many cell types in plants, as well as the challenge of annotating a broad
sample of species, we reduced resolution of our annotation across our datasets to ensure
accurate comparisons between variable species (**Figure 1B**).

Deeper exploration of the list of marker genes from *Z. mays* showed conservation of
gene body chromatin accessibility in markers for certain cell types (**Supplemental Table 2-3**) .
As expected, for the C₄ plants, *RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL*
SUBUNIT1 (RBCS1) and RIBULOSE BISPHOSPHATE CARBOXYLASE SMALL SUBUNIT2
(*RBCS2*) were enriched in BS cells compared to MS cells (**Figure 1C**), a pattern that was not
found in *O. sativa.* Additionally, *PEPC1* showed MS-specific chromatin accessibility in all of the
C₄ species sampled (**Figure 1D**). Additionally, we found conservation of marker genes like
SUCROSE TRANSPORTER 1 (*SUT1*) in companion cells and sieve elements, and *GLOSSY1*
(*GL1*) in epidermis cells, indicating that these historically described marker genes are likely
important in this diverse set of species. This analysis provides a first examination of core-C₄
marker genes' chromatin accessibility across a diverse sample of plant species at cell-type
resolution.
177

178
179
Figure 1: Annotation of cell types in diverse grass species at single-cell resolution **A)** A
phylogeny indicating the relationship of various C₃ and C₄ photosynthesizers sampled. In this
sample, two NADP-ME subtypes are represented, one NAD-ME subtype, a PEPCK subtype, as
well as a C₃ species. **B)** UMAP embedding showing the annotation for each species. A cell type
legend is below. **C)** Dotplots for various marker genes used to annotate each species. The y-
axis represents cell types, and the x-axis is a list marker genes used to annotate different cell
types. The size of each circle is proportional to the number of cells within that cell type that

showed chromatin accessibility of the marker. Color is z-score transformed values across
clusters of gene chromatin accessibility across the clusters. **D)** Screenshots of the *PEPC* locus
for all sampled species. For each screenshot, the top track shows the protein coding, the red
track is chromatin accessibility of MS cells, and the blue track is the chromatin accessibility of
the BS cells.
192
Chromatin Accessibility of Core C$_4$ Enzymes Shows Similar Cell-Type Bias, but Differing
Evolutionary Origins:
We measured the chromatin accessibility bias of the C$_4$-associated enzymes. Due to the
diverse nature of the plants sampled, and the C$_4$ photosynthetic subtypes, we separated
enzymes into core- and subtype-specific groups. This list comprised nine core C$_4$ enzymes, and
nine variable enzymes. These enzymes were assigned to one of these two groups based on if
they are found in all C$_4$ subtypes (core) or are specific to only one or two subtypes (variable).
One example of a core enzyme is carbonic anhydrase, which is used to generate bicarbonate
from CO$_2$, as well as for the regeneration of phosphoenolpyruvate from oxaloacetate in the BS
cells by means of PEPCK (**Figure 2A**). The list of gene families that we considered as core or
variable is found in (**Supplemental Table 4**).
To investigate the cell-type bias of these enzymes, we used chromatin accessibility of
the gene (gene body as well as 500 bp upstream of the transcriptional start site) (**Figure 2B**).
Cell-type bias was calculated as the log$_2$ fold change of BS/MS chromatin accessibility. To
identify core C$_4$ enzymes across these species, we used OrthoFinder, named and numbered the
enzyme models based off of their relatedness to *Z. mays* copies of known core C$_4$ genes (31).
Using only cell-type-specific chromatin accessibility data, we observed expected cell-type bias
with many orthologs of the maize MS-specific core C$_4$ genes showing MS-specific bias as
compared to BS (**Figure 2C**). For instance, in all C$_4$ species, *PEPCK*, which regenerates PEP
from OAA in BS cells, always showed a BS-specific bias (**Figure 2 A** & **C**). Additionally, *PEPC*,
which converts bicarbonate to OAA in MS cells, showed MS-specific bias for all species
sampled, except the C$_3$ outgroup *O. sativa* (**Figure 2A** & **C**). These results highlight the quality
of the data and the cell-type annotations for these single-cell datasets.
When analyzing these data in tandem with the phylogenetic trees, we noticed that some
of the key enzymes showed different cell-type specificity based on their evolutionary origin
(**Supplemental Figure 21-22**). For instance, for carbonic anhydrase in *P. miliaceum,* the
orthologs that showed the largest bias between MS and BS cell types were not the copies that
were the most evolutionary closely related to the *Z. mays* and *S. bicolor* cell-type-specific copies
(Here *PmCA1* and *PmCA2*). Rather, a copy found in a separate clade (*PmCA3*) showed the
most MS-specific bias (**Figure 2C**). This indicates that during the evolution of C$_4$, different sets
of carbonic anhydrases were likely co-opted for C$_4$. One challenge using chromatin accessibility
in this context, however, is the fact that neighboring gene models can occlude cell-type-specific
signals. For instance, in the *S. bicolor* copy of *RBCS1*, a BS-specific gene has a neighboring
gene model directly upstream which shares a promoter region making measurement of the cell-
type-specific bias of some loci challenging when using chromatin accessibility data
(**Supplemental Figure 23**).
One unexpected result from this analysis was the lack of cell-type-specific bias for
MALATE PHOSPHATE ANTIPORT 1 (DIC1), also known as

DICARBOXYLATE/TRICARBOXYLATE TRANSPORTER 1 (*DTC1*) in *Z. mays*. It has been
previously reported that *DIC1* had BS-specific expression bias in *Z. mays* as well as in *P.*
miliaceum (32–34)*.* However, there is not a clear signal based on the chromatin accessibility
data. This could indicate that some ACRs harbor multiple CREs active in different cell types that
are not obvious in chromatin accessibility data or that the cell-type-specificity observed is not
due to *cis*-regulation, possibly involving post-transcriptional processes (**Figure 2C**). Lastly, as
expected, there was very little bias in the C_3 outgroup (*O. sativa*). In total, 12/13 of the core C_4
enzymes showed cell-type-specific bias in *Z. mays*, 7/12 in *S. bicolor*, 16/21 in *P. miliaceum*,
11/13 in *U. fusca*, and finally 0/16 in *O. sativa*. These data demonstrate that chromatin-
accessibility data can be leveraged to investigate the cell-type regulation of C_4 genes while also
taking into consideration their evolutionary relationships in a cross species context.

247
248
Figure 2: Cell-type chromatin-accessibility bias for core enzymes in C_4 and C_3 species. **A**)
Schematic of the core C_4 enzymatic pathway. Core C_4 enzymes are defined as those which
maintain their cell-type-specificity in all C_4 subtypes sampled. The red and blue squares
represent MS and BS cells, respectively. Enzymes are labeled in bold, and transporters are
denoted by shapes. Intermediate molecules are indicated by non-bolded text. **B)** Screenshot of
PEPCK in *Z. mays*. Blue tracks correspond to BS chromatin accessibility and red tracks show
MS chromatin accessibility. Tracks are equally scaled to facilitate comparison. **C)** Heatmaps of

chromatin accessibility bias of the core C_4 enzymes. Values within each heatmap correspond to
Log2(BS/MS). Blue indicates increased BS chromatin accessibility and red indicates increased
MS chromatin accessibility. Each species column and subtype was clustered independently,
and genes were assigned as being MS- or BS-specific (top/bottom of heatmap) based on
literature. Enzyme copies were distinguished phylogenetically.
261
Key C_4 Subtype Enzymes Show Potential Convergent Evolution in Cell-type-specific
Bias:
We investigated the variable enzymes that give each C_4 subtype its unique properties by
focusing on two species (*S. bicolor* and *Z. mays*) from the *NADP-ME* subtype (**Figure 3A**). As
expected, chromatin accessibility bias was observed for enzymes previously reported as having
cell-type-specific expression patterns, similarly to the core C_4 enzyme set (29,35). Reassuringly,
one of the most biased enzymes identified was *NADP-ME*, the key enzyme of the redox step in
NADP-ME subtypes. More specifically, of the multiple copies of *NADP-ME* that exist in *Z. mays,*
we observed the expected cell-type bias for the known BS-specific copy, *ME3,* a key factor in C_4
(here *ZmNADP-ME1*) (**Figure 3B**). We noticed in *S. bicolor*, the BS-specific *NADP-ME* and the
MS-specific *NADP-malate dehydrogenase* (*NADP-MDH*) gene copies are recent tandem
duplications, each maintaining their respective cell-type specific chromatin accessibility (**Figure**
3B & **C**, **Supplemental Figure 22**). The malate transporters *DICARBOXYLIC ACID*
TRANSPORTER1/2 (DIT1/2) also demonstrated their expected cell-type-specific bias with *DIT1*
being MS specific and *DIT2* being BS specific in both species (**Figure 3B** & **C**). However, upon
further inspection of the phylogenies of the *DITs* in *S. bicolor,* we noticed a pattern where the
most BS-biased copy, *SbDIT4 (*Sobic.004G035500*)*, was phylogenetically more closely related
to the *ZmDIT1*. Something which has been previously reported (33,36). These results indicate
that over evolutionary time, even members of the same C_4 photosynthetic subtype, which likely
share a C_4 ancestor, can use different paralogous loci to achieve cell-type-specific expression.
This highlights that C_4 evolution is an ongoing process.
NAD-ME subtypes in *P. miliaceum* are interesting, as the intermediate molecule being
passed between MS and BS doesn't take the form of malate, but instead aspartate, alanine, and
oxaloacetate (**Figure 3D**). At least one copy of all of the key redox enzymes, NAD-ME and the
NAD-dependent malate dehydrogenase (NAD-MDH), show BS-biased chromatin accessibility
(**Figure 3E** & **F**). Interestingly, of the three copies of *NAD-MDH* analyzed, only two showed bias
for BS. Next, we evaluated two key enzymes associated with the generation of critical
intermediate metabolites, Aspartate aminotransferase (AspAT), and Alanine aminotransferase
(AlaAT). It has been reported that some AspAT have cell-type-specific expression patterns, with
the MS-specific copy of the protein being transported to the cytosol and the BS-specific copy
being transported to the mitochondria (**Figure 3E** & **F**) (37–39)**.** Of the four copies of AspAT we
examined, two (*PmAspAT3/4*) showed significant MS-specific bias, whereas the other two
copies (*PmAspAT1/2*) didn't show significant deviation towards BS (**Figure 3E**). This possibly
indicates differing levels of regulation for the AspAT copies that did not show the expected BS
bias, or missing copies of AspAT that we have not investigated. Within AlaAT, however, we
identified one copy, *PmAlaAT1,* showing MS-specific bias, and *PmAlaAT6* showing BS-specific
bias; something that has been previously hypothesized based on biochemical information (40).
Additionally, somewhat unexpectedly is that we didn't observe clear bias for sodium bile acid

symporters (*BASS*) and sodium:hydrogen antiporters (*NHD*) (**Figure 3E**). These two proteins
together form a functioning sodium bile acid symporter system, which balances the ratio of
sodium and is important in the transport of pyruvate into the chloroplast of MS cells (41)**.**
Although two copies of the *BASS* genes were MS biased, only a single copy of *NHD* was
slightly MS biased. Surprisingly, we do observe slight cell-type-specific chromatin accessibility
bias for malate transporter *DIT1/DIT2* in *P. miliaceum*. This is somewhat surprising, as malate is
not the main 4-carbon intermediate used by NAD-ME subtypes (10). This highlights the flexible
nature of *P. miliaceum* in terms of its C_4 photosynthetic style, as it has been implicated that it
can perform some of the metabolite shuttling as the NADP-ME subtype (10,42,43). The
potential flexibility of *P. miliaceum* in its style of C_4 makes it an extremely interesting species to
study, especially when considering that it doesn't share common C_4 ancestry with *Z. mays* or *S.*
bicolor. This lack of evolutionary relationship between *P. miliaceum* and *S. bicolor* and *Z. mays*
makes the comparison between *P.miliaceum* and its closer relative*U. fusca* all the more
valuable. These observations point to the complicated nature of some of these C_4
photosynthetic subtypes. While the obvious subtype-specific enzymes show expected
chromatin-accessibility bias, others do not.
Using the *PEPCK* subtype in *U. fusca,* we evaluated cell-type bias of enzymes that
operate as an intermediate between NAD-ME and NADP-ME subtypes (**Figure 3G**)**.** Copies of
NAD-ME and *PEPCK* showed significant BS bias (**Figure 3H** & **I**). Additionally, *NADP-MDH* was
significantly biased towards MS, reflecting its critical role in the regeneration of malate from
pyruvate (**Figure 3H**). We also observed one copy of *BASS,* which was heavily MS biased, as
well as the only copy of *NHD* being highly MS biased (**Figure 3G**) (44). Within the *BASS* family,
based on the phylogenies, it appears one clade of *BASS* genes was co-opted to be MS specific,
whereas the other clade remained somewhat BS specific. This potentially indicates that this co-
opted clade may have been predisposed for C_4 photosynthesis at the common ancestor of *P.*
miliaceum and *U. fusca.* Additionally, we also find one MS-biased and one BS-biased version of
AlaAT (**Figure 3H**).
Finally, when evaluating genes in the C_3 outgroup *O. sativa,* we only observed significant
chromatin accessibility bias for three of the 14 enzymes. This is expected given the overall lack
of enzymatic bias seen in C_3 species (**Figure 3K**). Interestingly though, we did find a single
instance where one copy of *AspAT* is BS specific, suggesting that this copy of *AspAT* might
slowly be co-opted into being more BS-specific (**Figure 3K**). Even more interesting is the slight
BS-specific bias of the rice *NAD-MDH*, a BS-specific enzyme in the *NAD-ME* subtypes. These
results show a series of complex evolutionary relationships where many different genes can be
co-opted into the C_4 pathway, and highlights the myriad ways in which C_4 evolution occurs.
335
336
337

338
Figure 3: Cell-type chromatin accessibility bias for variable C₄ genes associated with C₄
subtypes. **A/D/G/J)** Schematic of C₄ enzymatic pathways for various C₄ subtypes. The red and

blue squares represent MS and BS cells. Enzymes are labeled in bold, and transporters are
denoted by shapes. Intermediate molecules are indicated by non-bolded text. For clarity, core
enzymes have been removed. **B/E/H/K)** Heatmaps of chromatin accessibility bias in C_4 subtype
enzymes. Values within the heatmap correspond to Log2(BS/MS). Blue indicates increased BS-
chromatin accessibility and red indicates increased MS-chromatin accessibility. Genes were
labeled as being BS specific (blue) BS/MS specific (purple) or MS specific (red) based on
previous literature. **C/F/I)** Screenshot of various C_4 sub-type enzymes and their chromatin
accessibility profiles around the TSS. Blue tracks correspond to BS chromatin accessibility and
red tracks show MS chromatin accessibility. Tracks are equally scaled to facilitate comparison.
350
Cell-type-specific Accessible Chromatin Regions of Both Core- and Subtype-Specific
Enzymes:
Although measuring the gene body chromatin accessibility of C_4 enzymes is valuable, it
doesn't inform us about the cell-type-specific *cis*-regulatory environment controlling these
genes, as we only included 500 bp upstream in this initial analysis. To identify all potential CREs
important for regulation of C_4 enzymes, we identified cell-type-specific ACRs using a modified
entropy metric (**Methods; Supplemental Figure 33-34**). In short, cell-type-specific ACRs are
those which are unique to either a single cell-type or two or three cell-types in contrast to
broadly accessible ACRs which are accessible in many different cell-types. For each C_4
enzyme, in both the core and the non-core set, we identified ACRs around them. We only
considered ACRs to be potential regulators of a locus based on distance, with assigned ACRs
needing to be less than 200 kb away from the target enzyme, and requiring that no other gene
intervenes between the ACR and enzyme in question. In total, across all variable and core
enzymes and taking into consideration only C_4 species, we find that on average, C_4 genes have
between 2-3 cell-type-specific ACRs, with an additional 2-3 broadly-accessible ACRs (**Figure**
4A, Supplemental Table 5).
For all C_4 subtypes, the key redox enzymes all showed BS cell-type-specific ACRs,
potentially identifying critical CREs for proper cell-type-specific expression. For instance, in *Z.*
mays, NADP-ME1 had five BS-specific ACRs, in *S. bicolor, NADP-ME2* had five BS-specific
ACRs, in *P. miliaceum, NAD-ME1* had four BS-specific ACRs, and in *U. fusca, PEPCK,* had
three BS-specific ACRs (**Figure 4 A** & **C**). Additionally, of the MS-specific enzymes, we
consistently observed numerous cell-type-specific ACRs around the carbonic anhydrase family.
On average, there were 3.5 MS-specific ACRs for each copy of carbonic anhydrase across all of
the species. This likely reflects the fact that carbonic anhydrase is critical in the initial steps of
C_4, and also important in CO_2 sensing (45). We also noticed an intriguing pattern where
enzymes which were accessible in one cell type had cell-type-specific ACRs of the other cell
type. For instance, around *RBCS2,* a BS-specific enzyme, we found a series of MS-specific
ACRs (**Figure 4D**). On average, we found 2.5 BS-specific ACRs around *RBCS* and 1.5 MS-
specific ACRs. This contrasting pattern was observed in key photosynthetic enzymes in all of
the C_4 subtypes. This likely indicates that some of these ACRs contain CREs that negatively
regulate *RBCS* in MS, as cell-type-specific CRE usage has been implicated as being an
important driver in proper compartmentalization (46,47). The identification of ACRs around key
C_4 enzymes provides a detailed map about potential *cis*-regulators of these loci, which provides
the basis for future investigation into the direct function of each of these ACRs and how they

might be altering transcription in multiple different ways. These results show that there are likely
multiple ACRs important to cell-type specificity of these enzymes.
Traditionally, the field has focused on *cis*-regulation within a set distance from the
transcriptional start site, often 1-2 kb, which is thought to generally encompass the promoter
(48). However, we observed abundant distal cell-type-specific ACRs for many of these key
genes (**Figure 4B**). For instance, the average distance of an ACR to its C_4 enzyme is 10,080 bp
(*Z. mays*), 3,017 bp (*S.bicolor*), 4,260 bp (*P. miliaceum*), 2,358 bp (*U. fusca*), and 4,730 bp (*O.
sativa*), indicating that the *cis*-regulatory space for these enzymes is far greater than previously
appreciated, where a majority of the focus in the literature is on putative promoters. To test this,
we compared the identified ACRs to a series of previously reported cloned promoters. We found
that for *Zea mays* and *Sorghum bicolor* the ACR space identified includes significantly more
regions that are distal to their target gene (**Supplemental Figure 23C, Supplemental Table 6)**
(25,49,50).
The genome of *Z. mays* emphasizes this point, as the subtype-specific enzyme *NADP-ME* has
three cell-type-specific BS ACRs distal to the transcriptional start site, with the furthest being
34,336 bp away (**Figure 4C**). These distal ACRs provide critical regulatory loci to further
investigate. Interestingly, we found some enzyme/ACR pairs with opposite cell-type-specificity
(*i.e.* BS-specific enzyme, MS-specific ACR). Many of these ACRs were distally located. For
example, in *Z. mays,* the MS-specific ACR of *RBCS* was 36,171 bp upstream (**Figure 4D**).
When investigating ACRs around promoters, we were struck at how often cell-type-specific
ACRs occurred outside of the bounds of previously analyzed promoters. For example, in *PEPC*
in *P. miliaceum*, a recent analysis demonstrated that a series of conserved non-coding
sequences found between species were able to drive MS expression (27). When we looked at
chromatin accessibility data of the promoter fragment which was cloned from *PEPC*, we
identified many MS-specific ACRs within the cloned fragment, but an additional one upstream.
This results shows the advantage of using scATAC-seq data to identify candidate CREs for
certain genes, removing the guesswork of cloning fragments to investigate and providing a
detailed cell-type-specific regulatory map of the locus (**Figure 4E**). Thus, scATAC-seq greatly
improves the search space of the active CREs potentially driving cell-type-specific gene
expression patterns.
415

416
Figure 4: Investigating the number and distance of cell-type-specific ACRs around C$_4$ enzymes
across subtypes. **A)** Dot plots showing the number of cell-type-specific ACRs around each
enzyme. The x-axis indicates which cell type these enzymes are found in. The y-axis is counts
of ACRs. The graph is further subdivided with the top panel being broad ACRs, middle panel
BS-specific ACRs, and the bottom being MS-specific ACRs. Enzymes are labeled. **B)** Dotplots
showing the mean distance of cell-type-specific ACRs to their closest C$_4$ enzyme. The x-axis
indicates which cell type these enzymes are found in. The x-axis is the genomic distance to the
C$_4$ enzyme in question. If an enzyme had multiple cell-type-specific ACRs, the distance was
averaged (mean). **C)** Screenshot of *NADP-ME1* in *Z. mays*. Blue tracks correspond to BS

chromatin accessibility and red tracks show MS chromatin accessibility. Tracks are equally
scaled to facilitate comparison. All genes found within this window are shown. **D)** Screenshot of
RBCS2 in *Z. mays*. Blue tracks correspond to BS chromatin accessibility and red tracks show
MS chromatin accessibility. Tracks are equally scaled to facilitate comparison. All genes found
within this window are shown. **E)** Screenshot of *PEPC1* in *P. miliaceum.* The green fragment
represents the cloned promoter from Gupta et al 2020, which was identified by minimap2
alignment. Blue tracks correspond to BS chromatin accessibility and red tracks show MS
chromatin accessibility. Tracks are equally scaled to facilitate comparisons.

The Evolutionary Relationships of ACRs Associated with C_4 Genes is Complex and
Variable:
Next, we explored the evolutionary histories of these ACRs. Due to the fact that the C_4
subtypes come from different radiation events, (with *Z. mays* and *S. bicolor* likely sharing a C_4
ancestor and *U. fusca* and *P. miliaceum* sharing a different C_4 ancestor), we were curious to
evaluate if a majority of the ACR space around these genes were either novel, or shared among
these species. We implemented a pairwise sequence based approach by identifying sequence
conservation of ACRs between the study species using BLAST (**Methods)**. The majority of
important C_4 genes have both novel, and conserved ACRs. For example, PPDK, a MS-specific
enzyme, shares ~25% of its ACRs across all species examined including the *O. sativa* C_3
outgroup (**Figure 5A**). Interestingly, *RUBISCO ACTIVASE (RCA)*, a critical enzyme in
photosynthesis which removes inhibitory molecules from the RuBisCO active site, had novel
ACRs in all of the C_4 species examined, whereas *RCA* in the C_3 species *O. sativa* shared one
ACR with all of the C_4 species. This might indicate that each of the C_4 species gained regulatory
sequences at *RCA* or that *O. sativa* might have lost them (**Figure 5A**). Focusing on NADP-ME
revealed notable divergence in its associated ACRs, even among closely related species. For
example, in *Z. mays*, two out of nine ACRs linked to *NADP-ME1* were unique, lacking
counterparts in other species (**Figure 5A**). This is particularly striking given that *S. bicolor*,
belonging to the same C_4 subtype, diverged from *Z. mays* only 13 million years ago (51).
Similarly, in *S. bicolor*, the BS-specific *NADP-ME2* variant exhibited two out of five unique
ACRs. This pattern underscores the rapid and distinct evolutionary trajectories of ACRs in C_4
plants. A full list of gene families, and gene models, and their relative conservation is found in
Supplemental Figure 25A. Using this same approach to study all of the core class of C_4
enzymes did not reveal a generalizable pattern associated with gain or loss of ACRs around C_4
genes (**Supplemental Figure 25A**). Our findings not only confirm the dynamic evolution of *cis*-
regulatory sequences in C_4 enzymes but also align with existing research that highlights rapid
cis-regulatory changes among closely related species (48,52).
While investigating the ACRs around the C_4 genes is interesting, understanding how
cell-type specificity is achieved across C_4 subtypes is needed for efforts to engineer C_4
photosynthesis. When looking at just the cell-type-specific ACRs around key C_4 loci, we find a
similar pattern where there is a mix of both conserved and novel ACRs. For example, we
discovered that some of the MS-specific ACRs associated with *PPDK* and *PEPC* are highly
conserved in all of the studied species. Interestingly, the MS-specific ACRs around *PEPC* were
only found in the C_4 species, and not in the C_3 outgroup, *O.sativa* (**Figure 5B**). This indicates

that some of the CREs that allow *PEPC* expression in MS likely evolved after the split between
the most recent common ancestors. We also observed that *NADP-ME* possessed numerous
BS-specific ACRs that were conserved in all species, including *O. sativa* (**Figure 5B**).
Considering the fact that proper compartmentalization of *NADP-ME* in BS cells is only critical in
two of the three C_4 subtypes, this was surprising. However, in both *S. bicolor* and *Z. mays*, there
were novel BS-specific ACRs associated with each key *NADP-ME*. In *Z. mays,* one out of the
five BS-specific ACRs was novel to *Z. mays*, and in *S. bicolor* two out of the five were novel to
S. bicolor. Upon inspection of all the *NADP-ME* loci in genome browsers, we were struck by the
complexities and shuffling that occurred at these BS cell-type-specific ACRs (**Figure 5C**). These
results highlight that extensive *cis*-regulatory evolution is occurring in each of these species,
and in particular on a cell-type-specific level. Additionally, this may point to the fact that the
novel BS-specific ACRs found in *S. bicolor* and *Z. mays* may be more important for proper BS-
specific expression than the conserved regulatory elements.
482   Although binary classification of ACRs was useful to decipher larger scale patterns
between key enzymes, we next tested if larger segments of sequence were conserved around
some C_4 genes as compared to others. We profiled the relative amount of conserved sequence
at each of these ACRs, as alignment of sequence between species gives greater resolution
about important ACRs. One interesting observation from this analysis was the fact that the cell-
type-specific ACRs around *PEPCK* appear to be novel between *Z. mays* and *U. fusca* (**Figure
5D, Supplemental Figure 29-30**). This suggests that these regulatory loci emerged
independently, and yet are still likely important in cell-type-specific expression of *PEPCK*.
Additionally, around the *NAD-ME* loci in *P. miliaceum*, we found diverse evolutionary histories
with both copies *NAD-ME1* and *NAD-ME2* having both conserved and novel BS-specific ACRs
(one out of four ACRs were novel for *NAD-ME1*, and zero out of the two were conserved for
NAD-ME2) (**Figure 5D**). The ACRs from *NADP-ME1* are conserved in *U. fusca*, whereas all
three BS-specific ACRs are conserved in relation to *P. miliaceum.* Pointing to the fact that the
ACRs have likely maintained their cell-type specificity, and are likely critical drivers in the correct
expression of *NAD-ME* loci. These results highlight the dynamic evolution of cell-type-specific
ACRs around key C_4 loci, and that even closely related subtypes have evolved novel ACRs
potentially critical in terms of proper gene expression, as well as compartmentalization.
499
500
501
502

503
504
Figure 5: The evolutionary relationships of *cis*-regulatory regions around C4 genes is complex,
being composed of both novel and conserved ACRs. **A)** The proportion of all ACRs that are
conserved or novel for the following gene families *PPDK*, *RCA*, and *NADP-ME*. Purple bars
represent ACRs that have any sequence aligned to them from a different species, and gray
represents ACRs where sequences are not alignable. The number of ACRs in each locus is
labeled at the top of each column. **B)** The proportion of cell-type-specific ACRs that are
conserved and novel for the following gene families, *PPDK*, *PEPC*, and *NADP-ME*. Red bars
only consider MS-specific ACRs, and blue bars only consider BS-specific ACRs. **C)** Screenshot
of the conservation of BS-specific ACRs around *NADP-ME* across species. From top to bottom
the species are *Z. mays, S. bicolor, P. miliaceum, U. fusca, and O. sativa. NADP-ME* is
annotated in green for all species. Dashed bars between gene models represent the same gene
model, and yellow bars are conserved ACRs. Browser tracks are blue for BS, and red for MS.
Browser tracks are scaled within each species to allow for direct comparisons. **D)** The length of
ACRs that are conserved in a cross species context. Rows represent gene families, and
columns represent species. Each histogram is the number of ACRs within the loci of that gene
family. The x-axis is the length of the ACR that is conserved and the y-axis is the count. ACRs
are color coded according to the legend.
522

Identification of *de novo* TF-Binding Motifs from Cell-type-specific Chromatin Data
Reveals Rapid Sequence Diversification of ACRs

524
525

Leveraging the cell-type-resolved datasets, we identified *de novo* cell-type-specific TF motifs in
BS and MS ACRs (**Figure 6 A** & **B; Methods ; Supplemental Figure 31**). We selected the BS-
specific motifs based on motif similarity within C_4 species for BS, and motif similarity seen
across all species for MS. Additionally for the identification of BS specific motifs, we identified
motifs which didn't appear to have a corresponding motif in *O. sativa* (**Methods**). Reassuringly,
within the BS-specific motifs, we identified a DOF TF motif, which is a key driver in the switch to
C_4 photosynthesis (29,53,54). In brief, the DOF TFs have been implicated as being potential
drivers of proper gene expression in *Z. mays* C_4 genes, both as repressors and activators. For
example, *ZmDOF30* has been implicated as being important in driving BS specific gene
expression (29,53,54). In total we identified three BS-specific motifs, and four MS-specific *de*
novo motifs that are shared between the species sampled (**Figure 6 A** & **B ; Supplemental**
Figure 31). Using motif comparison tools, we were able to assign five out the of the six motifs to
a putative TF family, implicating potential novel regulators in BS-and MS-specific gene
expression (**Methods ; Supplemental Figure 29).** We surveyed the C_4 ACRs for the presence
and absence of these motifs to determine if they provide the information needed for cell-type
specificity. We additionally overlaid our BLAST results from the previous analysis in order to
explore the relationship between these motifs and conservation (**Figure 6C**). A substantial
number of motifs were present within the non-conserved regions of the ACRs. For instance, in
one MS-specific ACR associated with *ZmCA3*,12/13 MS-specific motifs were found in non-
conserved regions, suggesting these regions could be critical for driving the cell-type-specificity
of this locus (**Figure 6D).**

We expanded the analysis of BS- and MS-specific motifs in conserved and non-
conserved regions of ACRs across key loci in the C_4 species. On average the MS-specific
motifs are more conserved than the BS-specific motifs (**Figure 6E-F; Supplemental Figure 32**
). Agreeing with previous models of C_4 evolution where some motifs that are MS specific have
been co-opted to operate in C_4 photosynthesis (**Figure 6D**) (11). Interestingly, we noticed a
pattern where around *PPDK*, many of the MS-specific motifs appeared to be in non-conserved
sequences for all of our species sampled **(Figure 6E**). This pattern is further highlighted in both
NADPME, and *NADME* loci, where a majority of the BS-specific motifs occurred in non-
conserved ACR regions for *NADPME*. This pattern is more nuanced in the *NADME* ACRs, as *P.*
miliaceum and *U. fusca* share a significant amount of conserved sequence containing BS-
specific motifs in the ACRs, suggesting that the BS-specific regulatory changes associated with
these motifs are important (**Figure 6F**). These results highlight the capacity of genome-wide
single-cell *cis*-regulatory maps to pinpoint key TF motifs important for the evolution of cell-type
specificity.

561
562
563
564

565
Figure 6: Identification of cell-type-specific TF motifs reveal a complex relationship between
sequence conservation and motif prescnece. A subsample of MS- (**A**) and BS-specific (**B**) *de*
novo TF motifs identified. **Left)** *De novo* motifs were clustered by the correlation of their PWMs
and a correlation based tree was generated. **Right)** Representative PWMs from *de novo*
discovery. **C)** Screenshot of the *ZmCA3* locus. ACRs are color coded based on their cell-type
specificity. MS- and BS-chromatin accessibility tracks are equally scaled for comparison.
Sequence conservation is identified by the ACR having sequence homology to other *CA* ACRs
from a different species. **D)** An example of the conservation and motif landscape of one MS-
specific ACR at *ZmCA3*. Left, the location of the motifs in ACRs with MS- and BS-specific motifs
labeled. Orange highlighted regions correspond to the region of sequence conservation seen
above. Right**,** quantification of the motifs found in the ACR. X-axis is the motif count, and the y-
axis is the motif. **E)** The counts of TF motifs in conserved and non-conserved ACRs for three
different genes across all five species. Y-axis is the number of ACRs of a given type, and the x-
axis indicates the type of ACR. **F)** Odds ratio of four motifs when comparing their enrichment in
conserved versus non-conserved regions. A higher odds ratio indicates that the motif is more

often found in non-conserved regions within ACRs, whereas a lower odds ratio means the motif
is in conserved regions. The cell-type-specific motifs found in **A/B** are colored in red and blue,
respectively.
584
The DITs in the NADP-ME Subtypes Demonstrate Dynamic CRE Evolution
Upon analyzing the malate transporters *DICARBOXYLIC ACID TRANSPORTER's (DITs*
also known as the *DCTs)* we noticed the DITs in the NADP-ME subtypes showed an interesting
pattern where the copies of *DIT1* in *Z. mays* and *S. bicolor* showed MS-specific chromatin
accessibility, but the BS-specific copies of the *DITs* showed a more complex evolutionary
history (**Figure 3B**; **Figure 7A**). We generated a phylogeny with additional species, and found
that the BS-specific copy of *ZmDIT2* is related to two additional copies of *DITs* which are not
BS-specific in *S. bicolor* (Here *SbDIT2.2* and *SbDIT2.1*) (**Figure 7A**). *S. bicolor* has a BS-
specific copy of *SbDIT4,* which shares a clade with *ZmDIT1*. These results are consistent with
earlier studies that found similar patterns and gene expression profiles of these copies of the
DITs in *Z. mays* and *S. bicolor* (33,36,55). Although previous studies have documented changes
in cell-type-specific gene expression for the BS-specific copies of the DITs, the mechanisms
underlying these changes remain unclear. By using cell-type-specific ACRs, we explored if
expression changes are associated with changes in the number of cell-type-specific cis-
regulatory elements over evolutionary time.
To understand how cell-type specificity changed in these DITs due to changes in *cis-*
regulation, we compared the ACRs associated with the *DITs*, and mapped the TF-binding motifs
found within each ACR (**Methods**). For the MS-specific *DIT1s*, we focused on a MS-specific
ACR located at the 3' end of *DIT1* in *Z. mays* (**Figure 7B**). Upon comparing this ACR to *S.*
bicolor, we were struck that the sequence found in the *Z. mays* ACR was actually split in two in
S. bicolor, neither of which demonstrated cell-type specificity in *S. bicolor* (**Figure 7B ;**
Supplemental Figure 33). A closer inspection of motifs in these ACRs showed many MS-
specific motifs (**Figure 7B-C**). These motifs might promote MS-specific gene expression of this
locus. However, many *S. bicolor* MS-specific ACRs were not found in regions with any
homology to *Z. mays* (**Figure 7C**). These results point to the rapid change of candidate CREs
(cCRES) in this locus, and likely indicate that cCREs important in cell-type-specific gene
expression might not be only found in conserved regulatory regions (56). Rather, selection of
MS-specific gene expression is ongoing, and may yield significantly different regulatory
environments in relatively short evolutionary time scales.
Next, we examined the BS-specific *ZmDIT2* and its two orthologs *SbDIT2.1 and*
SbDIT2.2 , which are not BS specific (**Figure 7A, D**). The BS-specific ACR around *ZmDIT2* has
many DOF TF motifs (**Figure 7E**). These motifs are interesting, as expression changes within
the DOF TF family could be important in driving BS-specific gene expression in C_4 plants
(29,53,57). When comparing the BS-specific ACRs around *ZmDIT4* to the more closely related
copies of *SbDIT2.1* and *SbDIT2.2,* we found no conservation of these DOF TF motifs, and
rather a significant lack of BS-specific TF motifs (**Figure 7F**). Considering the fact that neither of
these *DIT* copies in *S. bicolor* show BS-specific expression, this result makes sense. Potentially
providing a model where the *ZmDIT4* locus either gained these cCREs allowing for this copy of
ZmDIT2 to have BS specific gene expression, or *S. bicolor* lost these BS-specific motifs, and
had a gain in *SbDIT4* specificity. In either scenario, it demonstrates the rapid pace of CRE

evolution, and how these regions might be altering cell-type-specific gene expression. These
results are in contrast to *SbDIT4*, where the ACRs around this locus are BS specific, and
contain BS-specific motifs identified in our previous analysis (**Figure 7F**). In total, these results
highlight the rapid rate of regulatory change around key C$_4$ loci, and highlight the fact that there
are likely key regulatory switches outside of conserved sequences. Finally, these results
emphasize the fast pace in which cell-type specificity changes in plants

Figure 7: A) Phylogenetic tree showing the evolutionary relationship of the *DITs* in the
monocots. *DITs* for *Z. mays* and *S. bicolor* are colored by their observed cell-type specificity,
with red being MS specific, and blue being BS specific. Additional species have been added to
increase resolution **B)** A screenshot of the *DIT1* between *Z. mays* (top) and *S. bicolor* (bottom).

Yellow boxes indicate ACR sequences with conserved homology **C/E/F)** Motif location of BS
and MS specific motifs in each ACR. The x-axis is the location within the ACR, and the y-axis is
the motif count. Yellow bars indicate regions of sequence homology. Within each histogram, the
x-axis is binned into 20bp regions for ease of graphing. Roman numerals in the top corner
highlight the corresponding ACR found in the screenshot. **(I-IX) top)** X-axis the genomic
coordinates of the given ACR. Yellow blocks denote the sequence homology as seen above. Y-
axis, the motif score as calculated by motifmatchR, higher scores indicate a more confident
motif. **bottom)** The count of each motif identified in the ACR. Note that BS and MS *de-novo*
identified motifs are in blue and red respectively. **D)** A screenshot of the BS specific DITs loci
between *Z. mays* (top) and *S. bicolor* (bottom). For the *S. bicolor* versions of the *DIT*s, *DIT4* is
colored blue for its observed BS specificity and *DIT2.1* and *DIT2.2* are colored green. Yellow
boxes indicate sequence homology.
650
651

Discussion:

Understanding the evolution of *cis*-regulation associated with C_4 photosynthesis has
been a long standing goal in the field of plant biology. In this study, we demonstrated the utility
of single-cell ATAC-seq data to investigate many aspects of the evolution of C_4 photosynthesis.
By identifying cell-type-specific chromatin accessibility from four C_4 species composed of three
different C_4 subtypes, as well as a single C_3 outgroup, we were able to compare and contrast
key genes and their ACRs which define and distinguish C_4 photosynthesis. We have shown that
by using gene-body chromatin accessibility data, we can measure cell-type-specific bias of both
core, and subtype-specific C_4 enzymes. When taken into consideration with the gene family
trees of many of these enzymes, we show diverse co-option of enzymes into the C_4 pathway.
Additionally, we identify cell-type-specific ACRs surrounding these key C_4 enzymes. We find
numerous cell-type-specific ACRs surrounding key C_4 enzymes, many of which fall outside of
the core promoter region. Additionally we find that around all of the C_4 enzymes there is a mix of
both conserved and novel cell-type-specific ACRs indicating that regulatory evolution of these
regions is ongoing. Finally, we use cell-type-specific ACRs to identify a series of *de-novo*
binding motifs which appear to be cell-type specific, and show that these motifs surround C_4
loci, and have a mixed relationship with conservation depending on the motif. This indicates that
cell-type-specific TF motifs are rapidly changing around C_4 loci.
Investigation of the CREs driving cell-type-specific expression of C_4 genes is
challenging. This often requires evaluation using transgenic plants, which limits the number of
CREs that can be tested. This has greatly hampered efforts at understanding how *cis*-regulation
of C_4 genes evolves, whether by co-option of existing CREs or emergence of new ones. Our
results show the complex nature of CRE evolution of C_4 genes, including those specific to C_4
subtypes. While we observe conservation of ACRs around many C_4 genes, we do see
interesting examples where the subtype-specific enzymes have evolved novel ACRs (*NAD-*
ME's in *P. miliaceum*, and *PEPCK* in *U. fusca*). These results support that there is likely a
combination of both co-opting pre-existing CREs, as well as evolving new ones to facilitate
proper expression and cell-type-specification of genes. This is further exemplified by the
analysis of the *DIT* family of transporters, where we show striking accumulation of cell-type-

specific TF motifs in non-conserved regions of ACRs between two closely related species. This
highlights that the regions of the genome promoting cell-type-specific gene expression are likely
found in both conserved, and novel regions. Another recent single-cell genomic study of the
evolution of CREs important for photosynthesis using a comparison between *O. sativa* and *S.*
bicolor reached similar conclusions (57). They frequently found different ACRs and TF motifs in
promoters of orthologous C_4 genes (57). Future efforts to assay these candidate CREs using
reporter assays, transgenesis and genome editing will be required. Additionally, expanding
these analysis outward to all genes associated with photosynthesis might provide valuable
insights into how genes in the Calvin-Benson cycle alter their regulation in their adaptation to C_4
photosynthesis. Fortunately, these high-resolution maps of cell-type-specific ACRs of these key
genes/species provide a strong foundation to build upon.
Although these studies provide a blueprint for the study of key candidate CREs
associated with C_4 enzymes, profiling cell-type-specific chromatin accessibility of additional
species would be greatly beneficial. Although *O. sativa* is an invaluable outgroup for this study,
additional more closely related C_3 species might make these comparisons simpler, and add
additional resolution. For instance the C_3 grass species *Dichanthelium oligosanthes* is more
closely related to *U. fusca* and *P. miliaceum* and has a recently completed reference genome
(58)*.* Adding more species would enable greater resolution in the comparison of cell-type-
specific ACRs, as the genetic distance between the species we examined and *O. sativa* make
identification of conserved and novel ACRs challenging. As an example, the ACRs associated
with *NAD-ME's* in *P. miliaceum* might be co-opted instead of novel, however, based on our
sampling, we cannot say.
Genome editing analysis of many of these ACRs would significantly advance which
ACRs, and more specifically which CREs within the ACRs are most important for cell-type-
specific expression (22). However, currently generating genome edits in monocots is
challenging, time consuming and expensive. Fortunately, improvements to transgenesis are
constantly improving making achieving these goals more likely in the future (59). It's also
important to consider that mutational analysis of CREs is not straightforward, often requiring
numerous editing events of the *cis*-regulatory landscape of each gene. Previous studies have
shown that deletions of many CREs produce variable molecular and morphological phenotypes,
further complicating our understanding of the *cis*-reglatory code (60–62). And finally, many
species, including *P. miliaceum* and *U. fusca* have to date never been transformed. This
highlights the need to continually improve transgenesis methods to help facilitate the molecular
dissection of CRE. In conclusion, this study provides a comprehensive map of cell-type-specific
ACRs around key C_4 genes, which reveals the dynamic evolution and diversity of *cis*-regulation
of C_4 genes.
717
718

Acknowledgments:

This research was funded by awards from the National Science Foundation (IOS-2134912 and
IOS-1856627) and the Office of Research to RJS and Hong Kong University Grant Council
(GRF 1409420) to SZ. APM and JPM were supported by the National Institutes of Health
(K99GM144742) and (T32GM142623), respectively. This research was additionally funded with

support from the NSFC (32100438 & 32370247) and Shanghai Jiao Tong University 2030
Initiative (to X.T.).

727 Methods:

Plant Growth Conditions and Sampling:

Seedlings of all five plant species, including maize (*Zea mays* B73), sorghum (*Sorghum bicolor*
BTx623), proso millet (*Panicum miliaceum* L. CGRIS 00000390), and browntop signalgrass
(*Urochloa fusca* LBJWC-52), along with the C_3 plant rice (*Oryza sativa* Nipponbare), were grown
under the conditions of 12:12 Light/Dark cycles at 30°C Light/22°C Dark and at 50% humidity.
The sampling of the C_4 species was timed to coincide with a specific developmental stage,
identified when the ligule of the third leaf became visible, marking the third leaf unfolding, yet
prior to the appearance of the fourth leaf. For the C_3 species, rice, 18-day-old leaves were used
to correspond with the equivalent stage of the C_4 species.

Library Preparation:

Nuclei isolation for the experiments was conducted using fresh seedlings of both the C_4 and C_3
species at their respective developmental stages. The methodology for nuclei extraction,
encompassing the buffer composition and the subsequent steps, was used with procedures
outlined for single-nucleus combinatorial indexing with transposed-based ATAC-seq library
construction, as detailed in a prior study (63).

Genomes:

The *Z. mays* genome version 5 was downloaded from MaizeGDB (64,65). The *O. sativa*
genome was downloaded from rice.uga.edu. The *S. bicolor* version v5.1 was downloaded and
used from Phytozome version 13, as well as the *U. fusca* genome version 1.1 (66). Finally the
P. miliaceum genome was downloaded from NCBI, bioproject number PRJNA431363 (43).

Barcode Correction Read Alignment and Mapping of Tn5 Insertions:

Read UMIs were processed using cutadapt (version 4.5) to identify UMIs (67). First, the index
adapter sequences were trimmed from the reads. Next, the well barcodes and Tn5 barcode
within the reads were identified, removed from the original sequencing read, and appended to
the read header. Finally, a shell script is used to integrate all barcode information from the
reads' headers and label them correspondingly in the paired-end sequencing fastq files. Reads
were aligned using BWA (version 0.7.17) (68). Reads were filtered using samtools (version
1.16.1) for mapping quality of >10 for *Z. mays* , *S. bicolor* , *U. fusca*, and *O. sativa*. *P. miliaceum*
required a greater threshold of 30 given its recent whole genome duplication event increasing
the rate of multi-mapping reads (69). Duplicate reads were removed using picard tools (version
2.25.0) (70). Single-base pair Tn5 integration events were mapped using the python script
`makeTn5bed.py` found in the GitHub utils directory
(https://github.com/Jome0169/Mendieta.C4_manuscript). Finally, for each barcode only unique
Tn5 integrations sites were used for analysis. So if a nuclei had the same identical fragments
multiple times, only a single event was considered.

Isolating High-Quality Cells:
Cells were filtered using Socrates (21). In short, Fraction of Reads in Peaks (FRiP) scores were
calculated for each cell by pseudo bulking the libraries and identifying peaks. For each
individual cell, FRiP was calculated by intersecting Tn5 integration events with peaks. Cells with
a FRiP score greater than 0.2 were used. Additionally, TSS enrichment was calculated by
looking at the number of Tn5 integrations around TSS. Cells that had a TSS enrichment greater
than 0.15 were used. Finally, cells were compared to a random sample of low quality cells which
did not pass filtering, representing the "background" of cells, and correlation was calculated
between passing cells and background cells using the corr package in R. Cells which had a
correlation lower than 0.3 percent as compared to background cells were used for further
analysis.
UMAP embeddings were then calculated for each species utilizing genomic bins (71).
For each dataset, bins of 500 bp were calculated. To reduce the size of features to cluster on,
bins had to show accessible chromatin in at least 0.005% of total cells (roughly 50~100 cells in
each species). Additionally, bins that were broadly accessible across greater than 10% of cells
in the given dataset were also discarded to remove regions of the genome which were
constitutively accessible and wouldn't facilitate clustering. Finally, regions of the genome which
were associated with either blacklist (21), or genes which were known to be related to cell cycle
and circadian rhythms were removed. The final resulting matrix, which represented cell
barcodes X genomic regions (here bins), were then put through the term-frequency inverse-
document-frequency (TF-IDF) algorithm to identify genomic regions more descriptive of the
entire dataset (30). The resulting matrix was then input into Singular Value Decomposition, and
clustering was then done on the remaining features with the number of principal components
(PCs) equaling 50, and any PC with a correlation to read depth greater than 0.5 removed (72)
(30). Clustering was done using the Louvain clustering algorithm in order to bin cells into similar
groups based off of the PCs calculated above, with parameters "res = 1.5, k.near = 30, m.dist =
.01" in order to set K nearest neighbors to 30, minimum louvain distance to .01 in euclidean
space (73). Using the UMAP embeddings, doublets were removed using the software Scrublet
as implemented in Socrates software (74). At random, 5,000 cells were used to generate *in-*
silico doublets, and cells which were scored as being likely doublets were removed. Adaptive
thresholds were set on a per library basis. The doublet rate from Scrublet was compared
against a mixed library where genotypes of *Z. mays* were mixed Mo17 and B73, and genotype
doublets were identified. We found that Scrublet, on average, removed more cells in a
conservative fashion than the birthday problem and genotype doublets identified, so we utilized
the Scrublet doublet scores to be conservative. For the P. *miliaceum* dataset, replicates were
found to integrate poorly in the UMAP embedding. Harmony (version 0.1.1) was used adjust
replicate overlap with parameters "theta = 2, nclust=4, and var = "sampleID" (75). After
integration, clusters which skewed greater than 75% towards one replicate were removed from
downstream analysis.
806
Identification of Putative Orthologs:
To annotate species with less marker gene information, we identified putative orthologs or
marker genes using OrthoFinder (version 2.5.4) (31). For each species, the primary protein
sequence of the transcript was used as input to Orthofinder. In the resulting orthofinder outputs,

the script "find_markers.orthofinder.py" was used to parse the resulting phylogenies and return
back putative orthologs (https://github.com/Jome0169/Mendieta.C4_manuscript). For all C_4
genes analyzed, each orthogroup was additionally annotated by hand in order to ensure
accurate assignment of nearest orthologs phylogenetically.
815
Annotation of Cell Types:
Cell types were annotated by calculating gene chromatin accessibility for marker genes in each
genome on a per cell basis. These values were then visualized on the UMAP embedding, and
clusters with numerous marker genes associated with the same cell-type were used as
evidence. Additionally, for each louvain cluster, enrichment of marker genes was calculated by
comparing the cluster average as compared to a random shuffle of random cells. The top five
most enriched markers were used in tandem with the UMAPs to ascertain cell-type identity. We
also tested the statistical significance of the marker gene using Presto, a modified Wilcoxon
rank-sum test in order to identify the most unique marker gene in each cluster (76). Additionally,
for specific clusters showing mixed signals from marker genes, sub-clustering was done by
isolating the cluster in question, and then re-clustering these cells on a new UMAP manifold.
The same steps were done to visualize marker genes, as well as test this enrichment, and
statistical significance. Finally, to bolster our set of marker genes across species, we used our
most confident cell-type annotation in *Z. mays* to *de novo* discover marker genes. To do so, we
utilized our gene-body-accessability metrics for each annotated cell-type, and ran DESeq2
(version 1.42.0) in a replicate aware fashion using all other cells as a null (77). Only statistically
significant markers were kept which had a fold change greater than 1.5, and a log fold standard
error of less than .6. OrthoFinder was used as mentioned above to find orthologs. To ensure
that we were comparing similar cell-types, we also took an orthogonal approach where we
compared the gene accessibility of the top 2000 most variable orthologs between our species. A
linear model was used for each species comparison where the mean gene accessibility was
taken into consideration, and the species was one-hot-encoded. Variation was calculated as the
average variation between both datasets. The resulting residuals were used to generate the
cell-type correlations.
840
Peak Identification:
To identify peaks, cells of the same annotation type were pseudo bulked in a replicate aware
fashion. Within each replicate MACS2 (version 2.2.9.1) was run with parameters "--nomodel --
keep-dup auto --extsize 150 --shift -75 --qvalue .05" and variable genome size flag '-g' (78).
Summits for each peak identified in each replicate were extended by 250 bp in either direction.
Only peaks which overlapped between replicates were used. To merge peaks from various cell
types and select peak boundaries, the p-value associated with each peak in each cell type was
compared by calculating the chromatin accessibility score for each peak per million, with those
peaks with the highest accessibility score being selected as the representative peak. This
method of identifying the most representative peaks across cell-types was inspired by previous
single cell ATAC-seq papers (30,79,80). Additionally, bigwigs were generated for each cell type
by normalizing each dataset to the number of reads/per million scaling factor. Implementation of
this algorithm is found in the script call_scACRs.py for ease of use and replication in other
experiments.

Identifying Cell-type-specific ACRs:

To identify cell-type specific ACRs, a modified bootstrapping method was used which drew inspiration from the modified entropy metrics found in (79). On a per ACR basis, Tn5 integrations per cell-type were summed and counts per million (CPM) normalized. These values were then converted to a probability by using the following equation (below, equation 1) where pi is the CPM value for the focal cell-type and qi is the total sum of all CPMs. From this probability statement, a modified shannon entropy metric was calculated, followed by a metric of specificity *Qpt*. For robust cell-type-specific ACR identification, the annotated cell-type was bootstrapped 5000 times, taking a sample of 250 cells from the cell population in question, and calculating both entropy and specificity scores. This was done to attempt to get a robust signal of specificity, which takes into consideration the variation in cell quality present in each cell-type annotation. To generate the null distribution of specificity scores, individual cell annotations were scrambled to generate an equal number of null cell-type classifications. For each null value, the entropy and specificity score were calculated. Finally to calculate a p-value, a non-parametric approach was used to identify how many of the real bootstraps fell outside of the null distribution using a one tailed test. ACRs which had a p-value of <0.001 were considered to be significant. ACRs were finally classified by the number of cell types they were specific to. ACRs specific to greater than three were classified as broadly accessible, less than or equal to three as cell type restricted, and a single cell-type as cell-type specific.

1) $pi \; = \; \frac{qi}{\Sigma(qi)}$

2) $Hp = -\Sigma pt log_2(pt)$

3) $Qpt \; = \; Hp \, - \, log_2(pt)$

Identifying Conserved ACRs Across Species

Since a majority of the C4 genes identified were not in synteny with one another, we took a gene family based approach to identify conserved and non-conserved ACRs associated with our C4 genes. In short, all ACRs within two gene models of a C4 gene are utilized for comparison. Sequences from the ACR were isolated using "bedtools getfasta" (version 2.31) (81). Then in a pairwise fashion each species had their ACRs from one C4 gene family compared to the corresponding genomic loci of the same gene family in a different species. Comparisons were made using Blastn (version 2.2.29) with the following parameters " -task blastn-short -evalue 1e-3 -max_target_seqs 4-word_size 7 -gapopen 5 -gapextend 2 -penalty -1 -reward 1 -dust no -outfmt 6" (82). The output blast files were further filtered requiring sequence alignment to be greater than 20 nts, and have an evalue of .001. This analysis and the detailed commands ran can be found in the following snakemake file titled "ID_syntenic_orthologous.ACRs.snake", and found in the snakemake directory in the associated github.

Identifying Cell-type-specific Motifs:

De-novo cell-type-specific motifs were identified by using XSTREME (version 5.5.3) of the MEME suite (version 5.5.5) package (83,84). In brief the sequences underlying the cell-type-specific ACRs were isolated, and equally matched null set of broadly-accessible ACRs were used the comparison for genomic enrichment. These null ACRs were matched in terms of GC

content, and were only allowed to be 5% different from the cell-type-specific set in question and generated using the script "gen_null_fa.py". Upon generation, motifs were analyzed using the universalmotifs package in R (version 3.18) (85). Motifs were first compared using HELL distance, and motifs which had a low correlation were discarded. In order to generate representative motifs, highly correlated motifs were merged using the function "merge_motifs" in found in the universalmotifs package. To identify the location of motifs, the R package motifmatchR were used, with a significant value cut off of .0005 (86).

Motifs Comparison:
In order to compare *de novo* identified motifs, position weight matrices were compared to using TomTom (version 5.5.5). Motifs were compared either against the non-redundant TF database for JASPAR plant TF binding motifs, or compared versus the consensus sequences found in Zenker et al 2024. The most significant motif was used to assign to potential TF families (87–89).

Data availability:
sciATAC-seq data for *Z. mays, S. bicolor, U. fusca,* and *P. miliceum* is found in NCBI under the following bioproject PRJNA1063172. Leaf data for *O.sativa* can be found under the following SRR bioproject PRJNA100757. All scripts used for processing and analyzing data in this manuscript can be found at the following github repository: https://github.com/Jome0169/Mendieta.C4_manuscript . Additionally, all datasets with both MS and BS specific accessibility profiles, their ACRs, as well as their BLASTN relationships can be found on the epigenome browser https://epigenome.genetics.uga.edu/PlantEpigenome. All datasets can be found under the sub-folder Mendieta_et_al.C4_project.

References:

1. Bowes G, Ogren WL, Hageman RH. Phosphoglycolate production catalyzed by ribulose diphosphate carboxylase. Biochem Biophys Res Commun. 1971 Nov 5;45(3):716–22.
2. Wheeler T, von Braun J. Climate Change Impacts on Global Food Security. Science. 2013 Aug 2;341(6145):508–13.
3. Hatch MD. C4 photosynthesis: a unique elend of modified biochemistry, anatomy and ultrastructure. Biochim Biophys Acta BBA - Rev Bioenerg. 1987 Jan 1;895(2):81–106.
4. Offermann S, Okita TW, Edwards GE. Resolving the Compartmentation and Function of C4 Photosynthesis in the Single-Cell C4 Species Bienertia sinuspersici. Plant Physiol. 2011 Apr 1;155(4):1612–28.
5. Akhani H, Barroca J, Koteeva N, Voznesenskaya E, Franceschi V, Edwards G, et al. Bienertia sinuspersici (Chenopodiaceae): A New Species from Southwest Asia and Discovery of a Third Terrestrial C4 Plant Without Kranz Anatomy. Syst Bot. 2005 Apr 1;30(2):290–301.
6. Sage RF. A portrait of the C_4 photosynthetic family on the 50th anniversary of its discovery: species number, evolutionary lineages, and Hall of Fame. J Exp Bot. 2016 Jul;67(14):4039–56.
7. Sage RF, Christin PA, Edwards EJ. The C4 plant lineages of planet Earth. J Exp Bot. 2011 May 1;62(9):3155–69.

8. Gowik U, Westhoff P. The Path from C3 to C4 Photosynthesis. Plant Physiol. 2011 Jan
1;155(1):56–63.
9. Grass Phylogeny Working Group II. New grass phylogeny resolves deep evolutionary
relationships and discovers C4 origins. New Phytol. 2012 Jan;193(2):304–12.
10. Rao X, Dixon RA. The Differences between NAD-ME and NADP-ME Subtypes of C4
Photosynthesis: More than Decarboxylating Enzymes. Front Plant Sci [Internet]. 2016 [cited
2023 May 22];7. Available from: https://www.frontiersin.org/articles/10.3389/fpls.2016.01525
11. Kajala K, Brown NJ, Williams BP, Borrill P, Taylor LE, Hibberd JM. Multiple Arabidopsis
genes primed for recruitment into C4 photosynthesis. Plant J. 2012;69(1):47–56.
12. Sheen J. C4 Gene Expression. Annu Rev Plant Physiol Plant Mol Biol. 1999;50(1):187–217.
13. Chollet R, Vidal J, O'Leary MH. PHOSPHO *ENOL* PYRUVATE CARBOXYLASE: A
Ubiquitous, Highly Regulated Enzyme in Plants. Annu Rev Plant Physiol Plant Mol Biol.
1996 Jun;47(1):273–98.
14. O'Leary MH. Phosphoenolpyruvate Carboxylase: An Enzymologist's View. Annu Rev Plant
Physiol. 1982 Jun;33(1):297–315.
15. Outlaw Jr William H. Kinetic Properties of Guard-Cell Phosphoenolpyruvate Carboxylase.
Biochem Physiol Pflanz. 1990 Jan 1;186(5):317–25.
16. Matsuoka M, Numazawa T. CIS-acting elements in the pyruvate, orthophosphate dikinase
gene from maize. Mol Gen Genet MGG. 1991 Aug 1;228(1):143–52.
17. Ku MS, Agarie S, Nomura M, Fukayama H, Tsuchida H, Ono K, et al. High-level expression
of maize phosphoenolpyruvate carboxylase in transgenic rice plants. Nat Biotechnol. 1999
Jan;17(1):76–80.
18. Nomura M, Sentoku N, Nishimura A, Lin JH, Honda C, Taniguchi M, et al. The evolution of
C4 plants: acquisition of cis-regulatory sequences in the promoter of C4-type pyruvate,
orthophosphate dikinase gene. Plant J. 2000;22(3):211–21.
19. Gowik U, Burscheidt J, Akyildiz M, Schlue U, Koczor M, Streubel M, et al. cis-Regulatory
Elements for Mesophyll-Specific Gene Expression in the C4 Plant Flaveria trinervia, the
Promoter of the C4 Phosphoenolpyruvate Carboxylase Gene[W]. Plant Cell. 2004 May
12;16(5):1077–90.
20. Kim ED, Dorrity MW, Fitzgerald BA, Seo H, Sepuru KM, Queitsch C, et al. Dynamic
chromatin accessibility deploys heterotypic cis/trans-acting factors driving stomatal cell-fate
commitment. Nat Plants. 2022 Dec;8(12):1453–66.
21. Marand AP, Chen Z, Gallavotti A, Schmitz RJ. A cis-regulatory atlas in maize at single-cell
resolution. Cell. 2021 May 27;184(11):3041-3055.e21.
22. Meng F, Zhao H, Zhu B, Zhang T, Yang M, Li Y, et al. Genomic editing of intronic enhancers
unveils their role in fine-tuning tissue-specific gene expression in Arabidopsis thaliana. Plant
Cell. 2021 Mar 25;33(6):1997–2014.
23. Spitz F, Furlong EEM. Transcription factors: from enhancer binding to developmental
control. Nat Rev Genet. 2012 Sep;13(9):613–26.
24. Matsuoka M, Kyozuka J, Shimamoto K, Kano-Murakami Y. The promoters of two
carboxylases in a C4 plant (maize) direct cell-specific, light-regulated expression in a C3
plant (rice). Plant J. 1994;6(3):311–9.
25. Matsuoka M, Tada Y, Fujimura T, Kano-Murakami Y. Tissue-specific light-regulated
expression directed by the promoter of a C4 gene, maize pyruvate,orthophosphate dikinase,
in a C3 plant, rice. Proc Natl Acad Sci. 1993 Oct 15;90(20):9586–90.
26. Gowik U, Schulze S, Saladié M, Rolland V, Tanz SK, Westhoff P, et al. A MEM1-like motif
directs mesophyll cell-specific expression of the gene encoding the C4 carbonic anhydrase
in Flaveria. J Exp Bot. 2017 Jan 1;68(2):311.
27. Gupta SD, Levey M, Schulze S, Karki S, Emmerling J, Streubel M, et al. The C4Ppc
promoters of many C4 grass species share a common regulatory mechanism for gene
expression in the mesophyll cell. Plant J. 2020;101(1):204–16.

28. Tu X, Ren S, Shen W, Li J, Li Y, Li C, et al. Limited conservation in cross-species
comparison of GLK transcription factor binding suggested wide-spread cistrome divergence.
Nat Commun. 2022 Dec 9;13(1):7632.
29. Dai X, Tu X, Du B, Dong P, Sun S, Wang X, et al. Chromatin and regulatory differentiation
between bundle sheath and mesophyll cells in maize. Plant J [Internet]. 2022 Jan 14 [cited
2022 Jan 14];n/a(n/a). Available from:
https://onlinelibrary.wiley.com/doi/abs/10.1111/tpj.15586
30. Cusanovich DA, Hill AJ, Aghamirzaie D, Daza RM, Pliner HA, Berletch JB, et al. A Single-
Cell Atlas of In Vivo Mammalian Chromatin Accessibility. Cell. 2018 Aug;174(5):1309-
1324.e18.
31. Emms DM, Kelly S. OrthoFinder: phylogenetic orthology inference for comparative
genomics. Genome Biol. 2019 Nov 14;20(1):238.
32. Taniguchi M, Sugiyama T. The Expression of 2-Oxoglutarate/Malate Translocator in the
Bundle-Sheath Mitochondria of Panicum miliaceum, a NAD-Malic Enzyme-Type C4 Plant, Is
Regulated by Light and Development. Plant Physiol. 1997 May 1;114(1):285–93.
33. Taniguchi Y, Nagasaki J, Kawasaki M, Miyake H, Sugiyama T, Taniguchi M. Differentiation
of Dicarboxylate Transporters in Mesophyll and Bundle Sheath Chloroplasts of Maize. Plant
Cell Physiol. 2004 Feb 15;45(2):187–200.
34. Tausta SL, Li P, Si Y, Gandotra N, Liu P, Sun Q, et al. Developmental dynamics of Kranz
cell transcriptional specificity in maize leaf reveals early onset of C4-related processes. J
Exp Bot. 2014 Jul;65(13):3543–55.
35. Borba AR, Reyna-Llorens I, Dickinson PJ, Steed G, Gouveia P, Górska AM, et al.
Compartmentation of photosynthesis gene expression in C4 maize depends on time of day.
Plant Physiol. 2023 Aug 9;kiad447.
36. Weissmann S, Huang P, Wiechert MA, Furuyama K, Brutnell TP, Taniguchi M, et al.
DCT4—A New Member of the Dicarboxylate Transporter Family in C4 Grasses. Genome
Biol Evol. 2021 Feb 1;13(2):evaa251.
37. Nomura M, Higuchi T, Katayama K, Taniguchi M, Miyao-Tokutomi M, Matsuoka M, et al.
The Promoter for C4-type Mitochondrial Aspartate Aminotransferase Does not Direct Bundle
Sheath-specific Expression in Transgenic Rice Plants. Plant Cell Physiol. 2005 May
1;46(5):743–53.
38. Taniguchi M, Kobe A, Kato M, Sugiyama T. Aspartate Aminotransferase Isozymes in
Panicum miliaceum L, an NAD-Malic Enzyme-Type C4 Plant: Comparison of Enzymatic-
Properties, Primary Structures, and Expression Patterns. Arch Biochem Biophys. 1995 Apr
20;318(2):295–306.
39. Taniguchi M, Sawaki H, Sasakawa H, Hase T, Sugiyama T. Cloning and sequence analysis
of cDNA encoding aspartate aminotransferase isozymes from Panicum miliaceum L., a C4
plant. Eur J Biochem. 1992;204(2):611–20.
40. Son D, Jo J, Sugiyama T. Purification and characterization of alanine aminotransferase from
Panicum miliaceum leaves. Arch Biochem Biophys. 1991 Aug 15;289(1):262–6.
41. Furumoto T, Yamaguchi T, Ohshima-Ichie Y, Nakamura M, Tsuchida-Iwata Y, Shimamura
M, et al. A plastidial sodium-dependent pyruvate transporter. Nature. 2011
Aug;476(7361):472–5.
42. Wang Y, Bräutigam A, Weber APM, Zhu XG. Three distinct biochemical subtypes of C4
photosynthesis? A modelling analysis. J Exp Bot. 2014 Jul 1;65(13):3567–78.
43. Zou C, Li L, Miki D, Li D, Tang Q, Xiao L, et al. The genome of broomcorn millet. Nat
Commun. 2019 Jan 25;10(1):436.
44. Washburn JD, Strable J, Dickinson P, Kothapalli SS, Brose JM, Covshoff S, et al. Distinct
C4 sub-types and C3 bundle sheath isolation in the Paniceae grasses. Plant Direct.
2021;5(12):e373.
45. Engineer C, Hashimoto-Sugimoto M, Negi J, Israelsson-Nordstrom M, Azoulay-Shemer T,

Rappel WJ, et al. CO2 sensing and CO2 regulation of stomatal conductance: advances and
open questions. Trends Plant Sci. 2016 Jan;21(1):16–30.

46. Bansal KC, Viret JF, Haley J, Khan BM, Schantz R, Bogorad L. Transient expression from
cab-m1 and rbcS-m3 promoter sequences is different in mesophyll and bundle sheath cells
in maize leaves. Proc Natl Acad Sci. 1992 Apr 15;89(8):3654–8.

47. Viret JF, Mabrouk Y, Bogorad L. Transcriptional photoregulation of cell-type-preferred
expression of maize rbcS-m3: 3' and 5' sequences are involved. Proc Natl Acad Sci. 1994
Aug 30;91(18):8577–81.

48. Lu Z, Marand AP, Ricci WA, Ethridge CL, Zhang X, Schmitz RJ. The prevalence, evolution
and chromatin signatures of plant regulatory elements. Nat Plants. 2019 Dec;5(12):1250–9.

49. Taniguchi M, Izawa K, Ku MSB, Lin JH, Saito H, Ishida Y, et al. The Promoter for the Maize
C4 Pyruvate,orthophosphate Dikinase Gene Directs Cell- and Tissue-Specific Transcription
in Transgenic Maize Plants. Plant Cell Physiol. 2000 Jan 1;41(1):42–8.

50. Heimann L, Horst I, Perduns R, Dreesen B, Offermann S, Peterhansel C. A Common
Histone Modification Code on C4 Genes in Maize and Its Conservation in Sorghum and
Setaria italica. Plant Physiol. 2013 May 1;162(1):456–69.

51. Paterson AH, Bowers JE, Bruggmann R, Dubchak I, Grimwood J, Gundlach H, et al. The
Sorghum bicolor genome and the diversification of grasses. Nature. 2009 Jan
29;457(7229):551–6.

52. Maher KA, Bajic M, Kajala K, Reynoso M, Pauluzzi G, West DA, et al. Profiling of
Accessible Chromatin Regions across Multiple Plant Species and Cell Types Reveals
Common Gene Regulatory Principles and New Control Modules. Plant Cell. 2018
Jan;30(1):15–36.

53. Yanagisawa S. Dof1 and Dof2 transcription factors are associated with expression of
multiple genes involved in carbon metabolism in maize. Plant J. 2000;21(3):281–8.

54. Perduns R, Horst-Niessen I, Peterhansel C. Photosynthetic Genes and Genes Associated
with the C4 Trait in Maize Are Characterized by a Unique Class of Highly Regulated Histone
Acetylation Peaks on Upstream Promoters. Plant Physiol. 2015 Aug 1;168(4):1378–88.

55. Emms DM, Covshoff S, Hibberd JM, Kelly S. Independent and Parallel Evolution of New
Genes by Gene Duplication in Two Origins of C4 Photosynthesis Provides New Insight into
the Mechanism of Phloem Loading in C4 Species. Mol Biol Evol. 2016 Jul;33(7):1796–806.

56. Yan H, Mendieta JP, Zhang X, Marand AP, Liang Y, Luo Z, et al. Evolution of cell-type-
specific accessible chromatin regions and the cis-regulatory elements that drive lineage-
specific innovation [Internet]. bioRxiv; 2024 [cited 2024 Jan 13]. p. 2024.01.08.574753.
Available from: https://www.biorxiv.org/content/10.1101/2024.01.08.574753v1

57. Swift J, Luginbuehl LH, Schreier T, Donald RM, Lee T, Nery J, et al. Single nuclei
sequencing reveals C4 photosynthesis is based on rewiring of ancestral cell identity
networks [Internet]. bioRxiv; 2023 [cited 2023 Oct 31]. p. 2023.10.26.562893. Available
from: https://www.biorxiv.org/content/10.1101/2023.10.26.562893v1

58. Studer AJ, Schnable JC, Weissmann S, Kolbe AR, McKain MR, Shao Y, et al. The draft
genome of the C3 panicoid grass species Dichanthelium oligosanthes. Genome Biol. 2016
Oct 28;17(1):223.

59. Chen Z, Debernardi JM, Dubcovsky J, Gallavotti A. The combination of morphogenic
regulators BABY BOOM and GRF-GIF improves maize transformation efficiency [Internet].
Plant Biology; 2022 Sep [cited 2024 Jan 2]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2022.09.02.506370

60. Ciren D, Zebell S, Lippman ZB. Extreme restructuring of cis-regulatory regions controlling a
deeply conserved plant stem cell regulator [Internet]. bioRxiv; 2023 [cited 2024 Jan 2]. p.
2023.12.20.572550. Available from:
https://www.biorxiv.org/content/10.1101/2023.12.20.572550v1

61. Liu L, Gallagher J, Arevalo ED, Chen R, Skopelitis T, Wu Q, et al. Enhancing grain-yield-

related traits by CRISPR–Cas9 promoter editing of maize CLE genes. Nat Plants. 2021
Mar;7(3):287–94.

62. Rodríguez-Leal D, Lemmon ZH, Man J, Bartlett ME, Lippman ZB. Engineering Quantitative
    Trait Variation for Crop Improvement by Genome Editing. Cell. 2017 Oct;171(2):470-480.e8.

63. Tu X, Marand AP, Schmitz RJ, Zhong S. A combinatorial indexing strategy for low-cost
    epigenomic profiling of plant single cells. Plant Commun. 2022 Jul 11;3(4):100308.

64. Hufford MB, Seetharam AS, Woodhouse MR, Chougule KM, Ou S, Liu J, et al. De novo
    assembly, annotation, and comparative analysis of 26 diverse maize genomes. Science.
    2021 Aug 6;373(6555):655–62.

65. Liu J, Seetharam AS, Chougule K, Ou S, Swentowsky KW, Gent JI, et al. Gapless assembly
    of maize chromosomes using long-read technologies. Genome Biol. 2020 May
    20;21(1):121.

66. Goodstein DM, Shu S, Howson R, Neupane R, Hayes RD, Fazo J, et al. Phytozome: a
    comparative platform for green plant genomics. Nucleic Acids Res. 2012 Jan;40(Database
    issue):D1178–86.

67. Martin M. Cutadapt removes adapter sequences from high-throughput sequencing reads.
    EMBnet.journal. 2011 May 2;17(1):10–2.

68. Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.
    Bioinformatics. 2009 Jul 15;25(14):1754–60.

69. Danecek P, Bonfield JK, Liddle J, Marshall J, Ohan V, Pollard MO, et al. Twelve years of
    SAMtools and BCFtools. GigaScience. 2021 Feb 1;10(2):giab008.

70. Picard Tools - By Broad Institute [Internet]. [cited 2024 Jan 4]. Available from:
    http://broadinstitute.github.io/picard/

71. McInnes L, Healy J, Melville J. UMAP: Uniform Manifold Approximation and Projection for
    Dimension Reduction [Internet]. arXiv; 2020 [cited 2022 Dec 19]. Available from:
    http://arxiv.org/abs/1802.03426

72. Stewart GW. On the Early History of the Singular Value Decomposition. SIAM Rev.
    1993;35(4):551–66.

73. Blondel VD, Guillaume JL, Lambiotte R, Lefebvre E. Fast unfolding of communities in large
    networks. J Stat Mech Theory Exp. 2008 Oct;2008(10):P10008.

74. Wolock SL, Lopez R, Klein AM. Scrublet: Computational Identification of Cell Doublets in
    Single-Cell Transcriptomic Data. Cell Syst. 2019 Apr;8(4):281-291.e9.

75. Korsunsky I, Millard N, Fan J, Slowikowski K, Zhang F, Wei K, et al. Fast, sensitive and
    accurate integration of single-cell data with Harmony. Nat Methods. 2019 Dec;16(12):1289–
    96.

76. Korsunsky I, Nathan A, Millard N, Raychaudhuri S. Presto scales Wilcoxon and auROC
    analyses to millions of observations [Internet]. Bioinformatics; 2019 May [cited 2023 Dec
    28]. Available from: http://biorxiv.org/lookup/doi/10.1101/653253

77. Love MI, Huber W, Anders S. Moderated estimation of fold change and dispersion for RNA-
    seq data with DESeq2. Genome Biol. 2014 Dec 5;15(12):550.

78. Zhang Y, Liu T, Meyer CA, Eeckhoute J, Johnson DS, Bernstein BE, et al. Model-based
    analysis of ChIP-Seq (MACS). Genome Biol. 2008;9(9):R137.

79. Zhang K, Hocker JD, Miller M, Hou X, Chiou J, Poirion OB, et al. A single-cell atlas of
    chromatin accessibility in the human genome. Cell. 2021 Nov 24;184(24):5985-6001.e19.

80. Domcke S, Hill AJ, Daza RM, Cao J, O'Day DR, Pliner HA, et al. A human cell atlas of fetal
    chromatin accessibility. Science [Internet]. 2020 Nov 13 [cited 2020 Nov 12];370(6518).
    Available from: https://science.sciencemag.org/content/370/6518/eaba7612

81. Quinlan AR, Hall IM. BEDTools: a flexible suite of utilities for comparing genomic features.
    Bioinformatics. 2010 Mar 15;26(6):841–2.

82. Camacho C, Coulouris G, Avagyan V, Ma N, Papadopoulos J, Bealer K, et al. BLAST+:
    architecture and applications. BMC Bioinformatics. 2009 Dec 15;10:421.

83. Grant CE, Bailey TL. XSTREME: Comprehensive motif analysis of biological sequence
datasets [Internet]. bioRxiv; 2021 [cited 2024 Jan 23]. p. 2021.09.02.458722. Available
from: https://www.biorxiv.org/content/10.1101/2021.09.02.458722v1
84. Bailey TL, Johnson J, Grant CE, Noble WS. The MEME Suite. Nucleic Acids Res. 2015 Jul
1;43(W1):W39–49.
85. Bioconductor [Internet]. [cited 2024 Jan 23]. universalmotif. Available from:
http://bioconductor.org/packages/universalmotif/
86. Bioconductor [Internet]. [cited 2024 Feb 1]. motifmatchr. Available from:
http://bioconductor.org/packages/motifmatchr/
87. Castro-Mondragon JA, Riudavets-Puig R, Rauluseviciute I, Lemma RB, Turchi L, Blanc-
Mathieu R, et al. JASPAR 2022: the 9th release of the open-access database of
transcription factor binding profiles. Nucleic Acids Res. 2022 Jan 7;50(D1):D165–73.
88. Rauluseviciute I, Riudavets-Puig R, Blanc-Mathieu R, Castro-Mondragon JA, Ferenc K,
Kumar V, et al. JASPAR 2024: 20th anniversary of the open-access database of
transcription factor binding profiles. Nucleic Acids Res. 2024 Jan 5;52(D1):D174–82.
89. Zenker S, Wulf D, Meierhenrich A, Becker S, Eisenhut M, Stracke R, et al. Transcription
factors operate on a limited vocabulary of binding motifs in *Arabidopsis thaliana* [Internet].
2023 [cited 2024 May 3]. Available from:
http://biorxiv.org/lookup/doi/10.1101/2023.08.28.555073