Invariant crossmodal equivalence between objects in rhesus monkeys

Elizabeth Cabrera-Ruiz,[1,2§†] Marlen Alva,[1†] Miguel Mata,[1] Mario Treviño,[3] José Vergara,[4] Tonatiuh Figueroa,[1] Javier Perez-Orive[2] & Luis Lemus[1*]

[1] Department of Cognitive Neuroscience, Instituto de Fisiología Celular, Universidad Nacional Autónoma de México, Mexico City, México.

[2] Instituto Nacional de Rehabilitación, "Luis Guillermo Ibarra Ibarra." Mexico City, México.

[3] Laboratorio de Plasticidad Cortical y Aprendizaje, Instituto de Neurociencias, Universidad de Guadalajara, Guadalajara, Jalisco, México.

[4] Department of Neuroscience, Baylor College of Medicine, Houston, TX, USA.

[§] Current address

[†] Contributed equally

[*] Correspondence: lemus@ifc.unam.mx

Author contributions

Conceptualization: LL. Data Curation: ECR, MM, TF. Formal Analysis: ECR, MA, MT, MM, JV & LL. Funding Acquisition: LL. Investigation: ECR, MM. Methodology: ECR & LL. Software: ECR, MA, TF, MT, & LL. Validation: JPO, MT, JV & LL. Visualization: ECR, MA & LL. Writing original draft: LL. Review & draft editing: ECR, MA, JPO, MT & JV.

Keywords

Crossmodal correspondence, Working memory, Multisensory, Macaque, Learning, Audiovisual.

Abbreviations

CC   Crossmodal categories

CME   Crossmodal equivalence

FA   False alarms

HR   Hit rate

NE   Nonequivalent

NHP   Nonhuman primates

JND   Just-noticeable difference

o   Learned sound

P   Picture

Pics   Pool of pictures on the touchscreen

PDF   Probability density function

PSE   Point of subjective equivalence

RT   Reaction times

S   Sound

STG   Superior temporal gyrus

STS   Superior temporal sulcus

Abstract

After hearing the words Little Red Riding Hood, most humans instantly visualize a girl

wearing a red hood in the woods. However, whether nonhuman primates also evoke such

visual imagery from sounds remains an open question. We sought to explore this from direct

behavioral measurements from two rhesus macaques trained in a delayed crossmodal

equivalence task. In each trial, they listened to a sound, such as a monkey vocalization or a

word, and three seconds later, selected a visual equivalent out of a pool of 2 to 4 pictures

appearing on a touchscreen. We propose two potential mechanisms for the brain to solve this

task: acoustic memory or visual imagery. In the first, sounds representations should remain in

working memory to discriminate later against the pictures on the screen. In the second,

listening to sounds could evoke visual representations of pictures that appear on the

touchscreen afterward. After analyzing the monkeys' choice accuracies and reaction times in

the task, we deduce that they experience visual imagery when listening to sounds. Therefore,

the ability of rhesus monkeys to perceive crossmodal equivalences between learned

categories poses rhesus monkeys as an ideal model organism for studying high-order

cognitive processes like semantics and conceptual thinking at the single-neuron level.

Introduction

Crossmodal correspondence, a natural animal ability, 'translates' information from one

sensory modality to another(Parise et al., 2016), for example by taking advantage of how they

both vary in a single physical dimension, e.g., lightness, roundness, texture, loudness, and

pitch(Walker et al., 2012; Ratcliffe et al., 2016). In lab settings, chimpanzees discriminate

visual shapes using sounds(Ravignani and Sonnweber, 2017), and trained macaques perform

crossmodal discriminations of shapes and textures(Weiskrantz and Cowey, 1975; Zhou and

Fuster, 2000), tones, and colors(Fuster et al., 2000), and flutter frequencies (Lemus et al.,

2010).

However, all these studies explored simple crossmodal correspondence along one physical

dimension such as shape or color, whereas in real life, we know that humans form complex

crossmodal equivalence (CME) between items of multiple and heterogeneous physical

attributes, for example, between words and the visual concept they evoke. Visualizing ideas

from words is a complex ability, facilitating the execution of intricate tasks, such as building

a cathedral. However, such a crucial evolutionary trait may have appeared first in an ancestor

common to other nonhuman primates (NHP). For instance, ethological studies implicitly

suggest CME in NHP(Ratcliffe et al., 2016). In a classical paper by Seyfarth and colleagues

(Seyfarth et al., 1980), vervet monkeys possibly evoked visual imagery because they looked

up, down, or ran into the trees after hearing conspecific alerts of leopards, eagles, and snakes.

       Neuroanatomical and behavioral similarities between humans and NHP indirectly

suggest that NHP can experience CME, for example, between voices and faces (Jordan et al.,

2005; Adachi and Hampton, 2011; Tyree et al., 2023). This mechanism could rely on the

integration of visual and auditory representations in multisensory areas of the brain (Gaffan

and Harrison, 1991; Calvert et al., 2001; Beauchamp et al., 2004; Khandhadia et al., 2021;

Diehl et al., 2022; Lemus and Lafuente, 2022), or from connecting neuronal representations

of monkey calls at the superior temporal gyrus (STG)(Leaver and Rauschecker, 2010;

Tsunada et al., 2011; Bodin and Belin, 2020; Bodin et al., 2021) with unitary activity

associated to a monkey face at the superior temporal sulcus (STS)(Tsao et al., 2003; Leopold

et al., 2006; Ohayon et al., 2012; Arcaro et al., 2017; Khandhadia et al., 2021). Regardless of

NHP being able to discriminate words phonetically(Melchor et al., 2021), it is unclear

whether their brains can encode complex sounds like words (Hickok and Poeppel, 2007; Yi et

al., 2019; Morán et al., 2021; Stephen et al., 2023) and associate them with visual

representations to experience CME.

Therefore, in the present study, we explore the ability of monkeys to perceive CME

between sounds, including words, and pictures rather than just triggering stereotyped

behaviors. Thus, we trained two rhesus macaques in a delayed CME task. In each trial, a

reference sound (S) appeared, and three seconds later, the monkeys had to match the sound

with its equivalent picture (P) from a pool of pictures (Pics) presented on a touchscreen.

Different hypotheses have been proposed about the perception of an object from different

modalities (Walker et al., 2012; Parise et al., 2016), we discuss two potential models for

solving the task: In the first, S elicits visual imagery in working memory to compare against

Pics. In the second, S identity persists in auditory working memory. Our results strongly

suggest that the monkeys experienced visual imagery during the CME task.

Results

We trained two rhesus monkeys in a CME task. In each trial, the monkeys listened to an S

and waited 3 seconds to find P out of Pics (**Fig 1A**, **S1 Video 1**; see methods). Solving the

task required the monkeys to hold information regarding S in working memory. We

hypothesized that such information would consist of visual imagery of P rather than a trace of

S. In other words, monkeys had to 'translate' S into P to discriminate Pics in the same

sensory modality. Therefore, we sought to find whether evoking learned visual information

about P predicted the animals' choice accuracy and reaction times (RT).

Fig 1. Delayed crossmodal equivalence task.

(A) Task events. In each trial, the monkey pressed a lever in response to a white cross on the

center of the touchscreen. After a variable period of 1-3 s, a reference 0.5s S was delivered.

Then, after a 3 s delay, Pics were presented simultaneously on the touchscreen, and the

animal released the lever to select the P equivalent to S. Correct responses were rewarded

with a drop of liquid. LP, lever press. **(B)** Sonograms and spectrograms of some S in the task,

such as two Spanish words shown in IPA nomenclature. Other sounds corresponded to a

monkey coo call and cow vocalization. **(C)** Pictures learned as equivalents of the S shown

above in **B**. Each CC consisted of an S-P pair.

The monkeys learned crossmodal equivalences in a few sessions.

We created eighteen crossmodal categories (CC) from combinations of six S and fourteen P

(**S2 Table of CCs**; See examples in **Fig 1BC**). The monkeys initially practiced the task in

trials with only two Pics, one equivalent to S and one nonequivalent (NE; see methods).

Then, we introduced more CCs and trials of 3 and 4 Pics. To analyze the monkeys' learning

process, we derived four learning parameters from fitting simple associative learning curves

to the performance in each CC throughout sessions: the hit rate (HR) in the first session (y0)

and sessions of statistical (γ), inertial (δ), and consolidated learning (λ), respectively (see

Methods).

The performance of monkey G at three CCs discriminated in trials of 2, 3, and 4 Pics

is shown in **Fig 2A**. Since the coo/monkey-face CC (left panel) was one of the first learned,

its y0, or initial performance (~300 trials) was close to chance level (intersection of the black

learning curve with the ordinates). Then, it took another eight sessions (~2700 trials) to reach

statistical learning (γ; **Fig 2A**, gray boxes, left edge) when the subjects began to solve the

task without being consistently correct. Then, inertial learning (δ) occurred after the 15th

session, where the change in performance from one session to another was insignificant, so

the performance is assumed to be constant. Our interpretation is that a transition from

sufficient performance (γ) to efficient (δ) behavior occurred before reaching the optimal

performance (λ).

Fig 2. Monkeys learn the crossmodal equivalence between sounds and pictures.

(**A**) Sigmoidal fits (black lines) of the behavioral accuracy of monkey G throughout learning

sessions. Blue lines, y' values of the first derivative of the sigmoidal fits. Each panel

corresponds to the accuracies in the CCs monkey, cow, and ['ro. xo] (the Spanish word for

red), discriminated in trials of 2, 3, and 4 Pics on the touchscreen, respectively. Grey boxes

delimit the sessions between the statistical learning γ and the inertial learning δ (see

methods). Dashed lines indicate the chance level at each number of Pics. (**B**) Same as in **A**,

but for the averaged accuracy of all CC performed by each monkey. (**C**) Boxplots of the

overall proportion of hits and false alarms (open) of sessions after δ.

However, the number of sessions for reaching γ and δ at new CCs decreased when the

monkeys learned the aim of the task. For example, monkey G took less than three sessions to

reach γ in trials of 3 and 4 Pics and less than ten to δ. The exception was the word [ro. xo]

(Spanish for red), where it spent ~14 sessions to δ. **S2 Table of CCs** shows all CCs and their

learning parameters. **Fig 2B** presents the mean of all CC learning for both monkeys. We

interpret the reduction in γ and δ as the monkeys solving the cognitive control of the motor

behavior required for the task (procedure memory), e.g., pressing and releasing the lever and

interacting correctly with the touchscreen, so that once this was done, the animals could focus

only on learning the CC associations.

Fig 2C shows examples of boxplots of the monkeys' overall accuracy after δ sessions,

i.e., after approaching the asymptote of learning the CC. In the case of monkey G, CCs were

presented in trials ranging from 2 to 4 Pics, whereas for monkey M, in 2 Pics (**S3 Table**

Overall performance). The open boxplots at the bottom of the figure correspond to false

alarms (FA), estimated as the percentage of times nonequivalent Pics were selected. The

overall HR of monkey G was 85.12 ± 9.11 (mean $\pm$ SD), and for monkey M, 87.07 ± 5.71

(mean $\pm$ SD). We also evaluated the performance of each category as a function of the angles

at which the correct visual cue was presented and found no bias (**S4 Figure**).

Regardless of the low proportion of FAs, they differed at every P so that, for each CC, there

was a sequence in FA distributions. For example, for monkey G, there often was a higher

proportion of FA for the picture 'red,' then for the cow, followed by the monkey, and, finally,

the human, suggesting a relationship between FAs and visual information, which we explore

further in the following sections. These results demonstrate that the monkeys achieved the

CME task proficiently after a few training sessions and with discriminations of 2 or more

Pics.

The RT distributions correlate with the number of visual categories.

We then explored the impact of Pics and each P on the RTs. As previously explained,

Monkey G discriminated in trials of 2 to 4 Pics and Monkey M in trials of 2 Pics. **Fig**

3A shows monkey G's RT distributions of hits for three S with all their P equivalents. The

RTs increased and dispersed more as a function of the number of Pics on the touchscreen

[Spearman rank correlation test between the number of Pics and RTs mean and STD, r (42) =

0.5828, p < 0.001 and r (42) = 0.5569, p < 0.001, respectively]. Similarly, for all CCs, the

higher the mean RT, the higher the variance (**Fig 3B**). These results indicate that the monkey

required more time to discriminate more Pics. Notably, no correlation existed between HR

and RTs (**Fig 3C,** Spearman rank correlation test).

Fig 3. RTs increase as a function of Pics.

(**A**) Probability frequency histograms showing RT increments of hits as a function of

increments in the number of Pics. Each panel corresponds to a CC. (**B**) Standard deviation of

RT distributions as a function of the distributions' mean. (**C**) RT did not affect accuracy.

By analyzing the RTs of different P in monkey G, which are all equivalent to a single

sound (e.g., four monkey faces equivalent to the same coo vocalization) (**Fig 4A**), we found

that the shape of the RT distributions were different for each P (p < 0.001, Kruskal-Wallis

test, p < 0.001 for all pairs of P and same S, post hoc Mann-Whitney U test, Bonferroni

corrected). Therefore, given that the S was the same, the differences in the RTs are due to the

P categories. The same happened in FA (**Fig 4A, insets**), where each distribution

corresponded to the selection of a P nonequivalent to several sounds (p < 0.001, Kruskal-

Wallis test, p < 0.01 for 71.43% for pairwise comparisons of coo, 76.19% for moo, and

82.14% for [si], Mann-Whitney U test, Bonferroni correction). In other words, the RT

distributions correlated with P categories while keeping S constant.

Fig 4. Reaction times in the CME task correlate with differences in visual categories.

(**A**) Cumulative density functions of RTs in various P, all equivalent to a single S. Each panel

shows RT distributions of correct responses to different P, all equivalent to a constant S.

Insets show RT of FAs produced when each P was selected, but the reference S was

nonequivalent. Notice how RT distributions resemble hits so that P selections could only

have biased RT. (**B**) RT distributions of monkey M during an experimental condition where

working memory was bypassed by presenting Pics since the beginning of the trial and

remained for S to be delivered as comparison. Notice how, again, hits and FA dispersed as P.

Insets show FAs of P nonequivalent to S (left), and S nonequivalent to P (right).

Our task requires four essential components: S, working memory, P, and Pics. Hence, given

that the auditory information remained the same, only three task components were left to

account for these variable RTs: visual information contained in Pics and P, i.e., the number of

pictures shuffled on the touchscreen in each trial, and visual imagery of P evoked during

working memory, respectively. In other words, the RTs would be a function of the visual

information. To disentangle between Pics and P, we evaluated monkey M in a condition

where Pics were constant. In each trial, we delivered Pics and remained on the touchscreen

for 1 – 3 seconds before S appeared (**S5 Video 2**). Again, the RTs correlated with the visual

rather than acoustic categories in hits and FAs (**Fig 4B**). Auditory memory was irrelevant in

this condition, and the monkeys were aware of the positions of the pictures from the

beginning of the trials. Therefore, S, invoking visual imagery, was the only one responsible

for the monkey's RTs.

The monkeys showed auditory perceptual invariance and RTs that followed the visual

categories.

So far, we have assumed that changes in RTs are not explained by S because they change

while keeping S constant. However, to verify our assumption was true, we presented the

monkeys with versions of S from emitters to which they had no previous exposure (**Fig**

5A). **Fig 5B** shows performance above chance in monkey G in 98.33% of the cases (paired-

sample t-test, $p < 0.05$; see **S6 Table Versions**). Notably, the RTs of hits at most versions

were dispersed in P categories rather than S versions (**Fig 5C, left panel**). The same was true

in FA when selecting a nonequivalent P (**Fig 5C, right panel**).

Fig 5. Monkeys show perceptual invariance to versions of S, reacting in times of P

categories.

(A) Spectrograms of some versions from different emitters of ['ro. xo] word. **(B)** Boxplots of

Monkey G accuracies at all S versions. Each color corresponds to each S category within

vertical line limits. Closed boxplots at the left of each category correspond to the accuracy

during the learned sound (o). Open boxplots correspond to versions of o. Closed boxplots at

the right of each group correspond to versions labeled on the x-axis, which were repeated

twice within 0.5 seconds. **(C)** The right panel shows that the RTs of hits of versions in **B,**

remain close to o's RTs (bold lines) and are distributed into the number of P categories. Left

panel: same as in right panel, but for FAs. **(D)** Same as in **C** but repeated S presented in thin

lines compared to o (bold lines). **(E)** Same as in **B**, but for monkey M. **(F)** Same as in **C**, but

for monkey M.

Meanwhile, monkey M performing in trials of 2 Pics, showed performance above

chance at 72.22 % of versions (5 to 7 versions per sound; paired-sample t-test., **Fig 5E**).

Here, RTs for hits and FAs were also dispersed in P categories (**Fig 5F**). These results

suggest that the monkeys experienced auditory constancy, leaving visual imagery as the

prevailing explanation for varying RTs during the CME task. Thus, we found that monkeys

could generalize, assigning auditory stimuli from different emitters to the learned S category.

Also, the RTs from the various emitters are explained by visual but not auditory properties.

Auditory changes involving certainty affect RTs.

After determining that acoustic versions did not affect RTs, we wanted to explore limitations

in the monkeys' auditory sensitivity favoring visual-based performance. To do so, we

presented two sets of auditory controls: acoustic repetitions and morphs. The repetitions

presented the same S twice in 500ms. The accuracy was best (**Fig 5B**, closed boxplots), and

the RTs were faster (**Fig 5C**, inset), suggesting that auditory certainty improved RTs.

Then, to estimate the effect of parametric modulations in S, we substituted S with

morphs between two S in gradual proportions (**Fig 6A**). The monkeys' accuracy in this

control set is shown in **Fig 6B.** The sigmoidal fit to the probability of finding each morph

proportion as equivalent to each learned P allowed us to derive the just noticeable difference

(JND) between morphs (see methods) and the proportion of morph where the monkeys

recognized an S 50% of the times (i.e., the point of subjective equivalence, PSE). When

fitting 2D Gaussian functions to the JNDs plotted against the PSEs of all morphing sets, it is

possible to see that recognizing the word [si] required more information than recognizing the

coo and moo (**Fig 6C**). Moreover, the monkeys performed better in morph repetitions, e.g.,

between the coo-coo to ['ro. xo] sounds (each morph were 500ms two-syllable sounds; **Fig**

6C, right panel). Overall, these results indicate that the monkeys could perceive subtle

differences in the morphs so that auditory deficits did not impair their crossmodal

discriminations.

Fig 6. Rhesus monkeys acutely recognize sounds from morphs.

(A) Spectrograms of a set of acoustic morphs between two learned sounds. **(B)** Sigmoidal fits

of the probability of the monkeys recognizing one S embedded in each morph percentage.

Each panel corresponds to the behavior of both monkeys at different morphing sets. Bold

lines: mean fit distribution. Shadowed fits: distributions of each of the sessions. **(C)** PSE as a

function of JND obtained from fits in **B. (D)** RTs of monkey G in each morph percentage and

set. **(E)** Same as in **D** but for monkey M.

In addition, the mean RTs of monkey G at each morph (**Fig 6D**) were higher as a function of

morph proportion, i.e., 50 %. However, the RT distributions differentiated into the number of

visual categories, regardless of the morphed S pairs. That fact was even more evident in

monkey M, where the RTs dispersed as a function of P across morphs (**Fig 6E**), except for

the P category "monkey," whose RTs were obtained at different morphing sets. In conclusion,

the monkeys' ability to detect little acoustic variations did not affect their visual-based

choices.

DISCUSSION

In our study, we trained two rhesus monkeys to perform a delayed crossmodal equivalence

task to recognize picture equivalents of sounds. We report three main findings from the

testing of the animals in different controls. First, the monkeys can learn associations between

visual and auditory objects as complex as faces, cartoons, and words (**Fig 2**). Second, they

perceive as invariant different versions of the learned sounds (**Fig 5**), which cannot be

explained by auditory performance (**Fig 6**), and third, sounds evoke visual representations in

their minds (**Fig 3-6**).

Rhesus macaques learn crossmodal equivalences across parametric processes.

In contrast to studies demonstrating the ability of monkeys to discriminate bimodally auditory, visual, or tactile stimuli with clear correspondence along one physical dimension (Weiskrantz and Cowey, 1975; Zhou and Fuster, 2000; Lemus et al., 2010; Nieder, 2012), for instance, the frequency of a tone with the frequency of a tactile vibration(Lemus et al., 2010), we show that they also discriminate perceptual objects of numerous and heterogeneous properties, as humans do between words and images. The fact that our monkeys learned numerous crossmodal associations should not be surprising since, in nature, monkeys recognize alarm calls about predators(Seyfarth et al., 1980). Moreover, there is evidence that neurons in the monkey's hippocampus respond invariantly to the face and voice of conspecifics in the colony(Tyree et al., 2023).

Additionally, in lab settings, monkeys have consistently proven their capabilities in bimodal discrimination tasks (Gaffan and Harrison, 1991; Fuster et al., 2000). Paradoxically, the mere training of monkeys in auditory tasks has been considered challenging (Ng et al., 2009; Scott et al., 2012). We also found it difficult since when we initially paired visual and auditory stimuli, the monkeys ignored the sounds. Thus, we trained them to detect the sounds first and do crossmodal associations afterwards. Once they grasped the task's objective, the learning time of CC categories was reduced significantly. Finally, we derived novel behavioral parameters from the performance throughout sessions to understand their learning process. The parameters, such as statistical and inertial learning derived from sigmoidal fits to performance, could help tailor individual monkey training.

Learned sounds evoke visual imagery in monkeys.

It has previously been reported that animals can imagine (Lai et al., 2023). However, whether

sounds evoke visual imagery in NHPs or set off stereotyped behaviors is unclear. We present

three results suggesting that sounds evoke visual imagery in monkeys. First, reaction times in

the CME task correlate with the number of picture categories. Second, because of the

monkeys' auditory perceptual invariance, the RTs did not disperse into new acoustic

categories but remained in distributions of visual categories. In addition, false alarms also

produced RTs that correlated with visual selections. Consequently, stereotyped reactions do

not explain the monkeys' behavior. Instead, their decisions rely on the discrimination of

visual images evoked by the sounds.

In our task, the monkeys had to listen to a sound and store the information necessary

for crossmodal comparisons in working memory. Therefore, in principle, they could use

acoustic memory to find its visual equivalent, as in studies showing parametric memory of

tactile and acoustic flutter, where discriminations of one-dimensional stimuli hardly predicted

the emergence of visual imagery (Lemus et al., 2009, 2010). However, we challenged the

auditory working memory strategy with a condition where the pictures appeared before the

sounds. This condition bypassed working memory because the touchscreen always projected

pictures. Once more, the RTs dispersed more with visual than with acoustic categories.

We verified the monkeys' sensitivity to slight auditory modulations by exposing them

to morphings of two sounds. The results show that the animals required a 10 to 20%

difference between sounds to recognize a category, so their acoustic perception was optimal.

However, their reaction times across morph proportions also corresponded with the number

of visual categories. Nevertheless, if auditory working memory remains a mechanism,

observing a picture on the screen would evoke an acoustic equivalent suitable for

discriminating with auditory memory. However, even in this scenario, a sound-to-visual

translation had to occur regardless of whether it occurred at the appearance of pictures.

Additionally, the bi-syllabic words ['ro. xo] and ['tʃaŋ.gi], carry more auditory

information than the coo and the moo vocalizations. Therefore, in the auditory working

memory scenario, the slowest RTs should have correlated with those words, but this was not

the case. We further assessed the effect of sound repetitions, e.g., coo-coo and [si]-[si],

finding that the performance was faster and more accurate than in single presentations, and

again, the RTs were grouped into visual categories. Another clue was that both monkeys

reacted faster to the "red" category, which consisted of only one visual attribute (even though

it is a bi-syllabic word in Spanish). At the same time, they spent more time discriminating the

monkey and human faces with similar attributes. In conclusion, the more complex the visual

discriminations, the longer it takes to achieve CME.

Similarly, sounds from different emitters are dispersed in a few visual categories

rather than in the numerous acoustic versions. In other words, like humans, monkeys

experience auditory perceptual invariance (Town et al., 2018)and make general relationships

with visual information. Although we only presented a limited number of visual equivalents,

visual imagery likely emerges during the delay period because, as suggested by previous

studies, discriminating auditory information is hard for macaques (Ng et al., 2009; Scott et

al., 2012), so visual imagery would present more naturally.

Multisensory areas like the prefrontal cortex or the STS, which store visual

objects(Beauchamp et al., 2004) and receive auditory inputs (Noesselt et al., 2007), are likely

to integrate acoustic and visual objects. We envisioned two alternative mechanisms to

achieve the CME: acoustic memory and visual imagery models. In the former, a sound would

activate its putative neural circuit at STG and remain active in working memory (**Fig 7A**).

Then, when its visual equivalent activates a visual representation at STS, it increases the STG

firing rate via a direct input to reach a CME threshold (**Fig 7B**). This model implies that

crossmodal equivalence translation only occurs during the comparison period.

Fig 7. Models for delayed CME in the brain.

(A) Auditory working memory model. When a sound elicits an STG circuit, it remains active

until a signal of the visual equivalent evoked at STS arrives to trigger a perceptual

equivalence to send to executive brain areas (dashed red arrows). **(B)** Firing rate of an STG

neuron across events of the CME task (bold line). The neuron responds to a sound (gray

dashed line) and remains active during the delay (green box). Since the neuron has no inputs

of nonequivalent visuals (NE; yellow dashed line), its firing rate remains constant. However,

when the equivalent P activates STG through direct input from STS, the firing rate increases

above the perceptual equivalence threshold (red dashed line). **(C)** Visual imagery model. The

arrival of a sound activates STG, which activates STS, evoking visual imagery of P. Visual

imagery remains until the actual sight of P creates perceptual equivalence, which is then

communicated to executive brain areas (dashed red arrow). Notice that NE P has no

interaction with the visual circuit activated by STG. **(D)** Same as in **B**, but for visual imagery

active in working memory.

Alternatively, in the visual imagery model (**Fig 7C**), once an STG circuit gets

activated by a sound, it immediately activates its visual equivalent at STS, which remains

active as visual imagery during the delay period until a picture on the touchscreen increases

its firing rate to equivalence threshold (**Fig 7D**). This model would explain the visual imagery

evoked from sounds. Our experiments demonstrate that the reaction times correlate with

evaluating visual information. Future recordings of the neural responses in areas such as STG

and STS can provide essential additional information (Cohen et al., 2007; Romanski, 2007;

Chandrasekaran et al., 2013; Huang and Brosch, 2016; Diehl et al., 2022).

Our findings are relevant because they prove that monkeys visualize faces out of

communication sounds. This crossmodal ability may have evolved in humans and monkeys

from a common ancestor to identify predators or rivals and understand other monkeys'

emotional states (Bodin et al., 2021; Diehl et al., 2022). Based on our results, we propose that

macaques present an ideal model for studying the neural basis of crossmodal equivalence.

This phenomenon is probably involved in other cognitive processes, such as the semantic

process in language and conceptual thinking.

Methods

Ethics statement

All experiments observed the Official Mexican Standard Recommendations for the Care and

Use of Laboratory Animals (NOM-062-ZOO-1999) and were approved by the Internal

Committee for the Use and Care of Laboratory Animals, Cell Physiology Institute, UNAM

(CICUAL, LLS200-2).

Subjects

Two adult rhesus monkeys (*Macaca mulatta*) participated in the experiments (a female,

monkey G, 10yr, 7 Kg, and a male, monkey M, 12yr, 12 kg). The animals inhabited single

cages in a room with constant temperature (22°C), filtered air, and day/light transitioning

lights. The animals had access to sources of enrichment, such as TV, toys, a recreating area,

and grooming with other animals through mesh sliding doors. They were restricted to water

for 12-15 hr. before experiments (Monday-Friday water intake > 20-30 ml/kg, ad libitum on

weekends). In addition to the primate pellets, they received 150 mg rations of fruits and

vegetables after the 2-3 hr. experimental sessions.

Experimental setup

The primates were trained to leave their cages and sit on a primate chair (Crist Instrument,

INC.) to be transferred to a soundproof booth adjacent to the vivarium for experiments.

Then, the chair was placed 30 cm from a touchscreen (ELO 2201L LED Display E107766,

HD wide-aspect ratio 22in LCD) and a spring lever below the touchscreen (ENV-610M, Med

Associates). Two speakers (Yamaha MSP5 Studio, 40 W, 0.050-40 kHz, and Logitech 12 W,

0.01-20 kHz) above the touchscreen delivered the sounds and a background noise at 45- and

65-dB SPL, respectively. The monkeys obtained liquid rewards through a stainless-steel

mouthpiece attached to the chair (Reward delivery system 5-RLD-E2-C Gravity feed

dispenser, Crist Instrument INC.).

Acoustic Stimuli

Sounds consisted of lab recordings of words, monkey vocalizations, and sounds downloaded

from free online libraries. All sounds were resized to 500ms, resampled to 44.1 kHz (cutoff

frequencies 0.1–20 kHz), and finally, normalized (RMS; Adobe Audition® 6.0). For the

perceptual invariance experiment, we replaced the sounds with versions of the same

categories and durations but with different emitters. We created acoustic morphs between two

sounds by interpolating the shortest trajectory between their fundamental and formant

frequencies at different proportions, i.e., from 10% to 90 % in steps of 10% (STRAIGHT

software package (Kawahara et al., 1999). Phonetic nomenclature of words in Spanish was

obtained using the Automatic Phonetic Transcriptionist by Xavier López Morrás

(http://aucel.com/pln/transbase.html).

Visual Stimuli

The visual stimuli consisted of black and white photographs of cows, monkeys, and human

faces, two cartoons, and a red oval circumscribed in ovals (200px/ sq-inch resolution),

presented equidistant from each other within a 4° radius from the center of the touchscreen.

The human face corresponds to the portray of the first author in this manuscript. Other

pictures were downloaded from free online sites and customize for the porpoise of the

experiments. Artwork of monkey in **Fig 1A** was created using a free online platform

(https://www.fotor.com/ai-art-generator).

Delayed crossmodal equivalence task

We trained two rhesus macaques in a delayed CME task. Each trial commenced with a 2°

aperture gray cross on the center of the touchscreen, indicating the monkey to press and hold

down the lever. Then, a 0.5 s sound followed by a 3 s delay period was delivered. Finally, a

pool of 2-4 simultaneous pictures appeared on the touchscreen, so the monkeys had to release

the lever and select the visual equivalent. The animals obtained a drop of liquid reward for

correct selections within 3 s of the appearance of the pictures. Sets of trials comprised ~300

trials of 2-4 CC presented intermingled. The task was programmed in LabVIEW 2014 (64-bit

SP1, National Instruments®).

Monkeys training

Our primary objective was to get the monkeys to perform the CME task as quickly and

efficiently as possible. Therefore, instructions included changes in stimuli, durations, and

rewards, sometimes within a single session, depending on the ongoing behavior of the

monkeys. Our approach was to train the animals to respond to a visual cue by pressing the

lever and interacting correctly with the touchscreen. Afterward, we introduced trials starting

with a sound and ending with two pictures. Only after they learned the first two CCs, we

presented the rest.

Behavioral measurements

Although our experiments did not aim to study the learning process of macaques, we

observed behavioral improvements across the sessions we sought to describe. First, we fit

learning curves to performance at each CC across sessions to evaluate the monkeys' learning

process. To establish a criterion for identifying discriminative choices within steady-state

behavior, we fitted learning curves to the monkeys' choice records using the Rescorla–

Wagner model of associative learning (Rescorla and Wagner, 1972). This model

conceptualizes learning through associations between conditioned (CS) and unconditioned

(US) stimuli. The extraction of the learning curves involved numerically solving the

following ordinary differential equation, as described before (Treviño, 2016):

$$\frac{dV}{dt} = \alpha\beta(\lambda - V) \qquad\qquad\qquad \text{Eq (1)}$$

This equation describes the evolution of the associative strength (V) towards a trained

CS as a function of the training trials (t). Optimized parameters from the model were $\alpha\beta$, the

product of the salience of the CS and the strength of the US (which were assumed to be fixed

during training, but see (Treviño, 2016), and the asymptote of learning (λ), which

corresponds to the maximum conditioning possible for the US. We then derived three

additional parameters from the extracted learning curves. Y0 described performance during

the first training session, corresponding to the fit intersection with the ordinates. Then, γ, or

statistical learning, was the first session at which performance statistically differed from

chance, i.e., it surpassed two standard deviations from chance level in a binomial distribution,

where p = chance level and n = mean number of trials per session. Meanwhile, the first

derivatives of the extracted learning curves were combined with user-defined thresholds to

identify the inertial (δ) experimental session at each CC where the monkey choices

approached the λ learning plateau, i.e., when the change in performance between two

consecutive sessions was equal or less than arbitrarily selected threshold of y' = 0.01 (Figure

2A, blue lines).

Reaction times (RTs) were the times from the appearance of the pictures to the lever

release. Choice accuracy was defined as the proportion of correct responses. False alarms

consisted of selections of nonequivalent pictures. Early responses and misses (not included in

this study) consisted of lever releases before or after the response window, leading to

aborting the trials.

To analyze biases toward a position on the screen for a specific P category, we

analyzed the difference between angles using the following quadrant approach. For a given P

angle (reference angle), we defined its reference 90°-quadrant as the area 45 degrees to the

left and 45 degrees to the right of that angle. We drew the rest of the 90°-quadrants relative to

the reference one. Then, using the multiple pairwise comparisons' results from the one-way

ANOVA (Tukey's HSD, $p<0.05$), we classified a quadrant as significant if at least half of its

angles significantly differed from the reference angle.

To evaluate the monkeys' ability to discriminate acoustic morphs performance was

fitted using a cumulative density function (CDF):

$$495 \qquad\qquad p(recognize[S]) = \frac{1}{2}\left[1 + \text{erf}\left(\frac{\%S-\mu}{\sqrt{2\sigma^2}}\right)\right] \qquad\qquad \text{Eq (2)}$$

where "p([S])" indicates the proportion of trials in which S category was recognized.

The μ, σ, and erf parameters correspond to the mean, standard deviation, and error function.

The mean of the Gaussian distribution is a measurement of the point of subjective equality

(PSE). This parameter lets us know the percentage of morphing the monkey chooses the

equivalent P 50% of the time. To calculate the percentage of morphing required to be

correctly recognized 75% of the time, also called just noticeable difference (JDN), we used

the following equation:

$$503 \qquad\qquad JND = \frac{norminv(0.75,\mu,\sigma) - norminv(0.25,\mu,\sigma)}{2} \qquad\qquad \text{Eq (3)}$$

where the *Norminv* MATLAB function, calculates normal inverse cumulative

distributions with mean "μ" and standard deviation "σ" from equation 1, at probability values

of 0.75 and 0.25. Finally, from different sessions, we created 2D-gaussian distributions (two

STD) of PSEs as a function of the JNDs of both monkeys.

Statistical analyses

Most analyses in this study comprised sessions after the δ session. Monkey M performed in

trials of two Pics and monkey G four Pics. Thus, performance above chance was greater than

0.5 for monkey M and 0.25 for monkey G. We only included RTs from sessions after δ

performance to compare RT distributions. We performed Spearman rank correlation tests to

find statistical differences between CCs RT distributions as a function of the number of Pics.

We performed a Kruskal-Wallis test to assess statistical differences among crossmodal

categories. In the case of null hypothesis rejection, a Mann-Whitney test was used to compare

cases of 2 and 3 P equivalent to one S and cases of a learned S with its versions. A

Bonferroni post hoc test was performed for multiple comparisons. Analyses were performed

using MATLAB R2022 (MathWorks).

Supporting Information

S1 Video 1. Monkey G performing in the delayed CME task.

S2 Table of CCs. Table presenting all crossmodal categories and the monkeys' parameters

obtained from their learning processes.

S3 Table Overall performance. Monkeys' performance in CCs most used.

S4 Figure. Monkeys' performance in numerous positions of P on the touchscreen. To assess

whether monkeys associated categories with specific regions on the screen, we analyzed the

performance of each CC as a function of the angles at which P was presented [one-way

ANOVA, False Discovery Rate (FDR) corrected for multiple comparisons using Benjamini-

Hochberg procedure (critical p-value = 0.0143)]. In monkey M, we did not find any bias

toward a particular angle ([si]: $F_{(11, 4.70215)} = 1.95$, $p = 0.034$; [coo]: $F_{(23, 30.6495)} =$

1.57, $p = 0.044$; [moo]: $F_{(23, 28.1022)} = 0.88$, $p = 0.632$). In monkey G, the performance of

the [si] category was not affected by P angles ($F_{(15, 130.086)} = 0.36$; $p = 0.988$), but we

found a significant effect for the categories [coo] ($F_{(15, 160.67)} = 1.97$; $p = 0.014$) and [moo]

($F_{(15, 150.619)} = 2.51$; $p = 0.001$). To make sense of these angle effects, we divided the

pairwise multiple comparisons (Tukey's HSD, $p<0.05$) into quadrants. We found such

differences between angles close to each other ($<90°$) and distributed across all the screen

quadrants. In other words, besides significant differences in the performance as a function of

the visual Target's angle, it did not show a systematic quadrant bias. With these finding, all

experiments presented here include presentations of pictures in only four quadrants.

S5 Video 2. Video of monkey M performing in a control condition of the CME task.

S6 Table Versions. Table of the performance of the monkeys during presentations of

different versions of the learned sounds.

Acknowledgments

We thank Francisco Pérez, Gerardo Coello, and Ana María Escalante of the computing

department of the IFC, Aurey Galván and Manuel Ortínez of the IFC workshop, and

Centenario 107 for providing invaluable working space and edible resources. Elizabeth

Cabrera Ruiz conducted this study to fulfill the requirements of Programa de Doctorado en

Ciencias Biomédicas of Universidad Nacional Autónoma de México and received a doctoral

scholarship from Consejo Nacional de Humanidades Ciencias y Tecnologías (245771). The

data in this work are part of her doctoral dissertation.

References

Adachi I, Hampton RR (2011) Rhesus monkeys see who they hear: Spontaneous cross-

modal memory for familiar conspecifics. PLoS One 6.

Arcaro MJ, Schade PF, Vincent JL, Ponce CR, Livingstone MS (2017) Seeing faces is

necessary for face-patch formation HHS Public Access Author manuscript. Nat

Neurosci 20:1404–1412.

Beauchamp MS, Lee KE, Argall BD, Martin A (2004) Integration of Auditory and Visual

Information about Objects in Superior Temporal Sulcus. Neuron 41:809–823.

Bodin C, Belin P (2020) Exploring the cerebral substrate of voice perception in primate

brains. Philos Trans R Soc B Biol Sci 375.

Bodin C, Trapeau R, Nazarian B, Sein J, Degiovanni X, Baurberg J, Rapha E, Renaud L,

Giordano BL, Belin P (2021) Functionally homologous representation of vocalizations in

the auditory cortex of humans and macaques. Curr Biol 31:4839-4844.e4.

Calvert G a, Hansen PC, Iversen SD, Brammer MJ (2001) Detection of audio-visual

integration sites in humans by application of electrophysiological criteria to the BOLD

567   effect. Neuroimage 14:427–438.

568 Chandrasekaran C, Lemus L, Ghazanfar AA (2013) Dynamic faces speed up the onset of

569   auditory cortical spiking responses during vocal detection. Proc Natl Acad Sci U S A

570   110:E4668-77 Available at:

571   http://www.pubmedcentral.nih.gov/articlerender.fcgi?artid=3845123&tool=pmcentrez&re

572   ndertype=abstract.

573 Cohen YE, Theunissen F, Russ BE, Gill P (2007) Acoustic features of rhesus vocalizations

574   and their representation in the ventrolateral prefrontal cortex. J Neurophysiol 97:1470–

575   1484 Available at: http://www.ncbi.nlm.nih.gov/pubmed/17135477.

576 Diehl MM, Plakke BA, Albuquerque R, Romanski LM (2022) Representation of Expression

577   and Identity by Ventral Prefrontal Neurons. Neuroscience 496:243–260 Available at:

578   https://doi.org/10.1016/j.neuroscience.2022.05.033.

579 Fuster JM, Bodner M, Kroger JK (2000) Cross-modal and cross-temporal association in

580   neurons of frontal cortex. Nature 405:347–351.

581 Gaffan D, Harrison S (1991) Auditory-visual associations, hemispheric specialization and

582   temporal-frontal interaction in the rhesus monkey. Brain 114:2133–2144.

583 Hickok G, Poeppel D (2007) The cortical organization of speech processing. Nat Rev

584   Neurosci 8:393–402.

585 Huang Y, Brosch M (2016) Neuronal activity in primate prefrontal cortex related to goal-

586   directed behavior during auditory working memory tasks. Brain Res 1640:314–327

587   Available at: https://pubmed.ncbi.nlm.nih.gov/26874071/ [Accessed March 14, 2023].

588 Jordan KE, Brannon EM, Logothetis NK, Ghazanfar AA (2005) Monkeys match the number

589   of voices they hear to the number of faces they see. Curr Biol 15:1034–1038 Available

590   at: http://www.cell.com/article/S0960982205004823/fulltext [Accessed March 13, 2023].

591 Kawahara H, Masuda-Katsuse I, De Cheveigné A (1999) Restructuring speech

592   representations using a pitch-adaptive time–frequency smoothing and an

593   instantaneous-frequency-based F0 extraction: Possible role of a repetitive structure in

594   sounds. Speech Commun 27:187–207.

Khandhadia AP, Murphy AP, Romanski LM, Bizley JK, Leopold DA (2021) Audiovisual

integration in macaque face patch neurons. Curr Biol 31:1826-1835.e3.

Lai C, Tanaka S, Harris TD, Lee AK (2023) Volitional activation of remote place

representations with a hippocampal brain-machine interface. Science 382:566–573.

Leaver AM, Rauschecker JP (2010) Cortical representation of natural complex sounds:

effects of acoustic features and auditory object category. J Neurosci 30:7604–7612.

Lemus L, Hernández A, Luna R, Zainos A, Romo R (2010) Do sensory cortices process

more than one sensory modality during perceptual judgments? Neuron 67:335–348.

Lemus L, Hernández A, Romo R (2009) Neural codes for perceptual discrimination of

acoustic flutter in the primate auditory cortex. Proc Natl Acad Sci U S A 106:9471–

9476.

Lemus L, Lafuente V De (2022) Look Who is Talking . Identities and Expressions in the

Prefrontal Cortex. Neuroscience 496:241–242 Available at:

https://doi.org/10.1016/j.neuroscience.2022.06.026.

Leopold DA, Bondar I V., Giese MA (2006) Norm-based face encoding by single neurons in

the monkey inferotemporal cortex. Nature 442:572–575.

Melchor J, Vergara J, Figueroa T, Morán I, Lemus L (2021) Formant-Based Recognition of

Words and Other Naturalistic Sounds in Rhesus Monkeys. Front Neurosci 15:1–10.

Morán I, Perez-Orive J, Melchor J, Figueroa T, Lemus L (2021) Auditory decisions in the

supplementary motor area. Prog Neurobiol 202:1–11.

Ng CW, Plakke B, Poremba A (2009) Primate auditory recognition memory performance

varies with sound type. Hear Res 256:64–74.

Nieder A (2012) Supramodal numerosity selectivity of neurons in primate prefrontal and

posterior parietal cortices. Proc Natl Acad Sci 109:11860–11865.

Noesselt T, Rieger JW, Schoenfeld MA, Kanowski M, Hinrichs H, Heinze HJ, Driver J (2007)

Audiovisual temporal correspondence modulates human multisensory superior

temporal sulcus plus primary sensory cortices. J Neurosci 27:11431–11441.

Ohayon S, Freiwald WA, Tsao DY (2012) What Makes a Cell Face Selective? The

Importance of Contrast. Neuron 74:567–581.

Parise C V., Spence C, Deroy O (2016) Understanding the correspondences: Introduction to

the special issue on crossmodal correspondences. Multisens Res 29:1–6.

Ratcliffe VF, Taylor AM, Reby D (2016) Cross-Modal Correspondences in Non-human

Mammal Communication. Multisens Res 74:49–91.

Ravignani A, Sonnweber R (2017) Chimpanzees process structural isomorphisms across

sensory modalities. Cognition 161:74–79.

Rescorla RA, Wagner AR (1972) A theory of Pavlovian conditioning: Variations in the

effectiveness of reinforcement and non-reinforcement. In: In Classical conditioning II

(Black A, Prokasy W, eds), pp 64–99. New York: Appleton-Century Crofts.

Romanski LM (2007) Representation and integration of auditory and visual stimuli in the

primate ventral lateral prefrontal cortex. Cereb Cortex 17:61–69.

Scott BH, Mishkin M, Yin P (2012) Monkeys have a limited form of short-term memory in

audition. Proc Natl Acad Sci U S A 109:12237–12241 Available at:

https://pubmed.ncbi.nlm.nih.gov/22778411/ [Accessed December 12, 2023].

Seyfarth RM, Cheney DL, Marler P (1980) Vervet monkey alarm calls: Semantic

communication in a free-ranging primate. Anim Behav 28:1070–1094.

Stephen EP, Li Y, Metzger S, Oganian Y, Chang EF (2023) Latent neural dynamics encode

temporal context in speech. Hear Res 437:1–13 Available at:

https://pubmed.ncbi.nlm.nih.gov/37441880/ [Accessed December 12, 2023].

Town SM, Wood KC, Bizley JK (2018) Sound identity is represented robustly in auditory

cortex during perceptual constancy. Nat Commun 9:1–15.

Treviño M (2016) Associative learning through acquired salience. Front Behav Neurosci

9:168673.

Tsao DY, Freiwald W a, Knutsen T a, Mandeville JB, Tootell RBH (2003) Faces and objects

in macaque cerebral cortex. Nat Neurosci 6:989–995.

Tsunada J, Lee JH, Cohen YE (2011) Representation of speech categories in the primate

auditory cortex. J Neurophysiol 105:2634–2646.

Tyree TJ, Metke M, Miller CT (2023) Cross-modal representation of identity in the primate

hippocampus. Science 382:417–423.

Walker L, Walker P, Francis B (2012) A common scheme for cross-sensory

correspondences across stimulus domains. Perception 41:1186–1192.

Weiskrantz L, Cowey A (1975) Cross modal matching in the rhesus monkey using a single

pair of stimuli. Neuropsychologia 13:257–261.

Yi HG, Leonard MK, Chang EF (2019) The Encoding of Speech Sounds in the Superior

Temporal Gyrus. Neuron 102:1096–1110.

Zhou YD, Fuster JM (2000) Visuo-tactile cross-modal associations in cortical somatosensory

cells. Proc Natl Acad Sci U S A 97:9777–9782.

A

			Sound	*delay*	
Cue	LP	0.5 - 1	0.5	3	release window

Time (s)

B

[si]   coo   moo   ['ro.xo]

Power (dB)
-20
-40
-60
-80
-100
-120
-140

Frequency (kHz)
10
5
0

Time (s)
0   0.5

C

A

coo

Cumulative density

Reaction times (s)

False alarms

moo

Cumulative density

Reaction times (s)

False alarms

B

Visual as reference

Cumulative density

Reaction times (s)

[tsangui]
coo
moo

False alarms by P

False alarms by S

[tsangui]
coo
moo

A

STS

[si]

coo

B

Acoustic memory

Firing Rate

COO

COO

NE   E

Time

C

STS

coo

D

Visual imagery

Firing Rate

COO

NE   E

Time