Title: Spatial characterization of interface dermatitis in cutaneous lupus reveals novel
chemokine ligand-receptor pairs that drive disease

Running head: Chemokine drivers of CLE interface dermatitis

Manuscript word, table and figure counts: 4,149 words, 6 figures, 7 supplemental tables, 9
supplemental figures

Authors: Saeed Shakiba[1*], Nazgol-Sadat Haddadi[1*], Khashayar Afshari[1], Janet E. Lubov[1],
Haya S. Raef[1], Robert Li[1], Ümmügülsüm Yildiz-Altay[1], Mridushi Daga[1], Maggi Ahmed Refat[1],
Evangeline Kim[1], Johanna Galindo de Laflin[1], Andressa Akabane[1], Priscilla Romano[1], Jane
Vongvirath[1], Shany Sherman[1], Elizabeth MacDonald[1], James P. Strassner[1], Liang Zhang[2],
Michael Leon[2], Christina E. Baer[3], Karen Dresser[4], Yan Liang[2], James B Whitley[5], Sladjana
Skopelja-Gardner[5], John E Harris[1], April Deng[4], Matthew D. Vesely[6], Mehdi Rashighi[1] & Jillian
Richmond[1**]

Institutions:

[1]UMass Chan Medical School, Dept of Dermatology, Worcester, MA, USA

[2]NanoString Technologies, Seattle, WA, USA

[3]UMass Chan Medical School, Sanderson Center for Optical Experimentation, Dept of
Microbiology and Physiological Systems, Worcester, MA, USA

[4]UMass Chan Medical School, Dept of Pathology, Worcester, MA, USA

[5]Dartmouth Hitchcock Medical Center, Dept of Medicine, Lebanon, NH, USA

[6]Yale University School of Medicine, Dept of Dermatology, New Haven, CT, USA

*Co-first authors

**Correspondence - for blister biopsies and patient info mehdi.rashighi@umassmed.edu, for all
other inquiries jillian.richmond@umassmed.edu

Acknowledgments: We thank those who donated tissue for this study. We thank Celia Hartigan
for clinical research support, and Jonathan Reichner for feedback on our manuscript. We thank
Alaina Ryan and the Comparative Pathobiology and Genomics Shared Resources ("CPGSR") in
Cummings School of Veterinary Medicine at Tufts University for running the protein DSP plate in
the MAX machine. We used Biorender.com for illustrations. SS used ChatGPT to correct English
grammar in the paragraphs he wrote for the first draft of the manuscript, and JMR edited for clarity
and brevity as a native English speaker.

Funding:
Lupus Research Alliance Target Identification in Lupus Award (JMR)

Lupus Research Alliance Diversity Research Supplement Award (JEL)

Dermatology Foundation Women's Health Career Development Award (JMR)

UMMS CCTS PPP grant (KA, JMR)

National Institute of Arthritis and Musculoskeletal and Skin Diseases Award Numbers
1R21AR079661-01 and 1R01AR08641-01A1 (SSG)

National Institute of Arthritis and Musculoskeletal and Skin Diseases Award K08 AR08777 (MDV)

National Institute of Arthritis and Musculoskeletal and Skin Diseases Award Numbers AR061437
and AR069114 (JEH)

Flow cytometry equipment used for this study is maintained by the UMass Chan Flow Cytometry
Core Facility. The UMass Chan Medical School Center for Clinical and Translational Research
was responsible for blood and biopsy collection and is supported by NIH Clinical and Translational
Sciences Award UL1TR000161. Protein DSP was supported by the Sanderson Center for Optical
Experimentation at UMass Chan (University of Massachusetts Medical School - UMass Chan -
SCOPE, RRID:SCR_022721) and the CPGSR at Tufts Cummings School of Veterinary Medicine.

The funders had no role in study design, data collection and analysis, decision to publish, or
preparation of the manuscript.

Conflict of Interest Disclosures: JMR is an inventor on patent application #63/478,900
"Diagnosis of skin diseases in veterinary and human patients" for CTCL. JEH & JMR are inventors
on patent application #62489191, "Diagnosis and Treatment of Vitiligo" which covers targeting IL-
15 and Trm for the treatment of vitiligo; and on patent application #15/851,651, "Anti-human
CXCR3 antibodies for the Treatment of Vitiligo" which covers targeting CXCR3 for the treatment
of vitiligo. JEH holds equity in Rheos Medicines and TeVido BioDevices; is a founder with equity
of Villaris Therapeutics, Aldena Therapeutics, NIRA Biosciences, Vimela Therapeutics, and Klirna
Therapeutics; has served as a consultant for Pfizer, Sanofi Genzyme, Incyte, Sun
Pharmaceuticals, LEO Pharma, Dermavant, Temprian Therapeutics, AbbVie, Janssen, Almirall,
Methuselah Health, Pandion, AnaptysBio, Avita, Aclaris Therapeutics, The Expert Institute,
BiologicsMD, Boston Pharma, Sonoma Biotherapeutics, Two Biotech, Admirx, Frazier
Management, 3rd Rock Ventrures, Gogen Therapeutics, Granular Therapeutics, Platelet
Biogenesis, BridgeBio, Merck, Matchpoint Therapeutics, and Klirna; has served as an investigator
for Pfizer, Sanofi Genzyme, Incyte, Sun Pharmaceuticals, LEO Pharma, Dermavant, Aclaris
Therapeutics, GSK, Celgene, Dermira, and EMD Serono. LZ, ML & YL are employees of
NanoString Technologies. MR is principal or co-investigator of studies sponsored by Pfizer,
Biogen, AbbVie, Incyte, LEO Pharma, Abeona Therapeutics, Dermavant, and Target RWE; and
MR provides consulting for Pfizer, Biogen, Incyte, Takeda, Inzen, ROME Therapeutics, Almirall,
Medicxi, Related Sciences, and VisualDx. Remaining authors declare that they have no
competing interests.

Contributor statement:
Conceptualization: JMR
Methodology: JMR, MR, SS, YL, CEB, JEH
Investigation: MAR, EK, JGDeL, JPS, AA, KA, LZ, ML, KD, EMacD, MDV
Validation: SS, UYA, MDV, SSher, MD, MR
Formal analysis: SS, NSH, KA, JEL, HSR, UYA, RL, JMR
Resources: AD, MR, CEB, JBW, SSG, JEH, MDV, JMR

Data Curation: SS, NSH, KA, JMR, JV, PR
Visualization: SS, NSH, KA, JEL, UYA, MD, JMR
Supervision: JMR, MR
Project administration: JMR
Funding acquisition: JMR
Writing - Original Draft: SS, NSH & JMR
Writing - Review & Editing: all authors

Data availability statement: Datasets related to this article have been deposited in National
Center for Biotechnology Information's Gene Expression Omnibus (GEO) *(72)* and are
accessible through GEO Series accession numbers GSE182302 for 96-plex immunoassay
data, and GSE182825 for spatial transcriptomics. Flow data has been deposited on
FlowRepository under accessions FR-FCM-Z4PL, FR-FCM-Z4PM, FR-FCM-Z4PN, FR-FCM-
Z4PQ, FR-FCM-Z4PX, FR-FCM-Z6UN, FR-FCM-Z7ZP, FR-FCM-Z7ZQ for blister biopsies and
blood immunophenotyping.

Ethics statement: This study was performed with internal review board (IRB) approved
protocols at the UMass Chan Medical School (H-14848 and H00021295 for fresh blood and skin
tissue; H00020503 for archival skin tissue), Yale University Institutional Review Board (Human
Investigative Committee no. 15010105235 for archival skin tissue), and/or Dartmouth Hitchcock
Medical Center (STUDY02001542 for PBMCs), and all samples were obtained with written
informed consent and were de-identified before use in experiments.

Bulleted statements:
• *What is already known about this topic?* Cutaneous lupus erythematosus (CLE) is a
rare, chronic autoimmune connective tissue disorder. Nonlesional skin was recently
reported to be enriched in CD14+CD16+ inflammatory myeloid cells.
• *What does this study add?* We mapped chemokine orchestrators of interface dermatitis
in lupus using spatial approaches on archival tissue and confirmed with fresh tissues, to
understand how keratinocytes, myeloid cells, T cells and other leukocytes form the
interface dermatitis reaction.
• *What is the translational message*? The translational message of our findings is that,
even on maintenance therapy, CLE patient skin is enriched for both T cells and antigen
presenting cells. Keratinocyte injury activates innate immune pathways that ultimately
induce chemokines. These chemokines recruit leukocytes, which begets additional
cascades of chemokines. Specifically, we show that the CD14+CD16+ inflammatory
myeloid cells respond to CXCL6, which is upregulated at the protein level in CLE
lesional interstitial skin fluid.
• *What are the clinical implications of this work*? Clinically, our findings have implications
for drug repurposing and future clinical trials. Namely, the CXCL6/CXCR1 inhibitor
reparixin and its derivatives, which are currently used in oncological applications, could
be tested in topical formulations for CLE.

Abstract:

Background Chemokines play critical roles in the recruitment and activation of immune cells in
both homeostatic and pathologic conditions. Here, we examined chemokine ligand-receptor

pairs to better understand the immunopathogenesis of cutaneous lupus erythematosus (CLE), a
complex autoimmune connective tissue disorder.

Objectives Our objectives were to (1) characterize the cellular and proteomic constitution of
interface dermatitis in CLE using blister biopsies, (2) map chemokine:ligand receptor pairs that
govern recruitment of immune cells to form interface dermatitis in CLE, and (3) perform
unbiased analyses in tandem on different clinical subtypes to identify novel genes and proteins
underlying discoid versus subacute CLE.

Methods We used suction blister biopsies to measure cellular infiltrates with spectral flow
cytometry in the interface dermatitis reaction, as well as 184 protein analytes in interstitial skin
fluid using 96-plex immunoassay targeted proteomics. Flow and 96-plex immunoassay data
concordantly demonstrated significant increases in T cells and antigen presenting cells (APCs).
We also performed spatial transcriptomics and spatial proteomics of punch biopsies using digital
spatial profiling (DSP) technology on CLE skin and healthy margin controls to examine discreet
locations within the tissue.

Results Spatial and 96-plex immunoassay data confirmed elevation of interferon (IFN) and IFN-
inducible CXCR3 chemokine ligands. Comparing involved versus uninvolved keratinocytes in
CLE samples revealed upregulation of essential inflammatory response genes in areas near
interface dermatitis, including *AIM2*. 96-plex immunoassay data confirmed upregulation of
Caspase 8, IL-18 which is the final product of AIM2 activation, and induced chemokines
including CCL8 and CXCL6 in CLE lesional samples. Chemotaxis assays using PBMCs from
healthy and CLE donors revealed that T cells are equally poised to respond to CXCR3 ligands,
whereas CD14+CD16+ APC populations are more sensitive to CXCL6 via CXCR1 and CD14+
are more sensitive to CCL8 via CCR2.

Conclusions Taken together, our data map a pathway from keratinocyte injury to lymphocyte
recruitment in CLE via AIM2-Casp8-IL-18-CXCL6/CXCR1 and CCL8/CCR2, and IFNG/IFNL1-
CXCL9/CXCL11-CXCR3, and identify potential novel biomarkers of disease.

Graphical abstract: Model of chemokine systems governing recruitment of immune cell subsets to form interface dermatitis in cutaneous lupus. A. Summary of fresh tissue blister biopsy studies demonstrating increases in HLADR+ cells in nonlesional lupus biopsies as assessed by flow cytometry, and increased CXCL6 and CCL8 as assessed by 96 plex immunoassay. Lesional lupus biopsies also demonstrated significant increases in T cells and CXCL9/10/11 production. B. Model of chemokine-directed formation of interface dermatitis in cutaneous lupus. 1. Whole transcriptome atlas (WTA) digital spatial profiling (DSP) revealed increased *AIM2* in keratinocytes proximal to inflammation, which is reported to be induced by environmental triggers including UV light and toxins. We also noted increased Caspase 8 (Casp8) and IL18 at the protein level, which can be induced downstream of AIM2. Chemokines including CXCL6 and CCL8 can be induced downstream of IL18, explaining how CCL8 and CXCL6 might be induced. 2. Recruitment of myeloid cell first responders by CCL8 and CXCL6. CD14+CD16+ myeloid cells, which were recently described in nonlesional lupus skin, express CXCR1 and migrate towards CXCL6, whereas CD14+CD16- myeloid cells express more CCR2 and migrate towards CCL8. 3. The CXCR3 ligands CXCL9/10/11 are expressed by keratinocytes, but more strongly in CD45+ immune cell and T cell regions of interest (ROIs). 4.

T cells express CXCR3 and migrate towards CXCL9 and to a greater extent CXCL11. 5. The
recruited HLADR+ myeloid populations and T cells contribute to formation of interface
dermatitis. Thus, we propose a model in which keratinocyte/myeloid crosstalk can reinforce
chemokine systems to optimally recruit lymphocytes and other immune cells to form interface
dermatitis. Created with Biorender.com.

Plain language summary: Lupus skin rashes arise during flares, after exposure to medications
or sunlight, or in response to other triggers of inflammation. To understand how white blood
cells enter the skin to cause these rashes, we used new technologies to look at proteins that
attract them into the skin. We found proteins that are expressed by skin cells in lesions that can
recruit specific types of white blood cells that are thought to be the initiators of skin rashes.
Once in the skin, these and other white blood cells can make additional proteins that bring in
more and more cells. We hope that our findings will be used to test new topical treatments for
lupus and other autoimmune skin rashes.

Main Text:

INTRODUCTION

Cutaneous lupus erythematosus (CLE) is a spectrum of clinically diverse autoimmune

connective tissue disorders with shared histopathological features including interface dermatitis

and lupus band reaction[1]. Chronic cutaneous lupus erythematosus (CCLE, most common sub-

entity discoid lupus erythematosus (DLE)), subacute cutaneous lupus (SCLE), and acute

cutaneous lupus erythematosus (ACLE) include the majority of cutaneous lupus clinical

subtypes, though there are other more rare entities which all have varying degrees of

association with systemic lupus erythematosus (SLE). In clinical practice, CLE can be refractory

to SLE treatments[2–4], but successful treatment of cutaneous disease may significantly decrease

the risk of systemic involvement[5].

A common feature of CLE is the presence of interface dermatitis, which refers to inflammation

and/or degenerative changes at the dermal-epidermal junction. The mechanisms by which

immune cells are recruited to form interface dermatitis in CLE are unknown. Previous studies

have found that interferon (IFN)-inducible chemokines are highly upregulated in CLE lesions [6,7].

One recent study characterized the associations of specific chemokine receptors and ligands in

T cell subsets in cutaneous versus systemic lupus, demonstrating different profiles in SLE, DLE

and SCLE[8]. The cellular sources of chemokine ligands binding to cognate receptors have not

been fully characterized.

To date, many CLE studies have focused on transcriptomics including bulk RNA sequencing

and single cell RNA sequencing[9], but overlooked proteomics approaches and cellular analyses.

Here, we used a blister biopsy technique[10] to sample interstitial skin fluid and cells from the

interface dermatitis reaction from CLE patients and compared these to healthy donors. Blister

fluid does not require enzymatic digestion for analysis, yielding high-quality liquid and cellular

biopsies. We used 96-plex immunoassay targeted proteomics to assess inflammatory

biomarkers in interstitial skin fluid, and flow cytometry to assess infiltrating immune cells and

their phenotypes. We demonstrate significant increases in T cells and HLA-DR+ cells in blister

fluid from patients compared to healthy controls. To understand how these immune cells might

enter the skin, we assessed chemokine ligand and receptor expression patterns. We confirmed

upregulation of CXCL9 and CXCL11 in patient blister fluid and skin biopsies, with expression of

the cognate receptor CXCR3 on T cells on both healthy and lupus blood donors. T cells from

healthy and lupus patients migrated towards CXCL9 and CXCL11 to varying degrees. We also

identified novel chemokine ligands that were elevated in lesional versus nonlesional lupus

blister fluid, namely CCL8 and CXCL6. CCL8 binds to the receptors CCR1, CCR2, CCR3 and

CCR5. We found that CD14+ monocytes expressed higher CCR2 than other populations of

myeloid cells and migrated towards CCL8. In contrast, CD14+CD16+ myeloid cells expressed

the CXCL6 receptor CXCR1 and migrated towards this ligand. We also performed matched

spatial transcriptomics and proteomics of DLE and SCLE archival biopsies to identify the

general cellular sources (keratinocytes vs immune cells) and locations of chemokines and other

immune proteins. We identified *AIM2* and *IFNG* in keratinocytes, which can lead to downstream

production of CXCL6, CCL8, and CXCL9/11 respectively. Notably, these ligands are not

expressed in healthy skin, meaning that immune cells are poised to respond to these ligands in

response to injury or insult such as UV light exposure. Further, spatial data identified unique

gene and protein expression signatures which may underlie differences in the discoid and

subacute clinical subtypes of CLE, in addition to unique expression of B7-H3 protein on SCLE T

cells. Importantly, we identify potential treatment targets in both immune cells and keratinocytes

that may be of use for future clinical trials, namely *IRAK*s and *CAPN*s and *TYK2*, as well as a

novel functional target CXCR1 which we demonstrate can be inhibited with reparixin.

MATERIALS AND METHODS

Study subjects - This study was performed with internal review board (IRB) approved protocols
at the UMass Chan Medical School (H-14848 and H00021295 for fresh blood and skin tissue;

H00020503 for archival skin tissue), Yale University Institutional Review Board (Human
Investigative Committee no. 15010105235 for archival skin tissue), and/or Dartmouth Hitchcock
Medical Center (STUDY02001542 for PBMCs), and all samples were obtained with written
informed consent and were de-identified before use in experiments. Inclusion/exclusion criteria
were as follows: subjects with a diagnosis of cutaneous lupus erythematosus by clinical exam
performed by a dermatologist were included. Subjects with recent onset of new lesions and
objective clinical signs of activity, specifically erythema and/or hair loss/alopecia, were recruited
to represent active disease. Patients on treatment were included in the study, as most lupus
patients must remain on maintenance therapy.

Statistics - Statistical analyses were performed using Prism software version 9 (GraphPad) for
flow cytometry, NPX software for 96-plex immunoassay, and GeoMX software for spatial
datasets. For comparisons across disease states and cell types, two-way ANOVAs with main
row effects comparisons were performed to identify highly significant biomarkers or cell types for
further analysis. Differences were considered significant at a P value of less than 0.05. DSP
statistical analyses using Linear Mixed Model (LMM) or unpaired T-tests were performed in
GeoMX software with the resulting p-values that were adjusted for multiple comparisons with
the Benjamini-Hochberg procedure (FDR, false discovery rate) and considering the size of
regions of interest (ROIs) and slide scanning batch effect. Box plots and volcano plots were
generated in GraphPad Prism version 9. P-value or FDR <0.05 were considered significant, with
P<0.01 as highly significant. We also used one-way ANOVA or two-way ANOVA for some
comparisons in a group of genes between healthy and disease ROIs to answer specific
hypotheses generated by -omics datasets.

RESULTS

Integrating spatial and -omics approaches for characterizing the inflammatory infiltrate in

cutaneous lupus interface dermatitis

We combined spatial studies of archival tissue with analyses of fresh tissue biopsies to validate

chemokine and other targets at the protein level (workflow **Fig 1A**). Blister biopsies were used

to sample interstitial skin fluid and cells from CLE patients and healthy donors (**Fig 1B, Table**

S1). 96-plex immunoassay targeted proteomics on the interstitial skin fluid confirmed previously

reported increased levels of IFNg, CXCL9 and CXCL11 in CLE skin biopsies (**Fig 1C**) [6,8]. 96-

plex immunoassay also identified higher levels of CXCL6 and hepatocyte growth factor (HGF),

which can increase CXCR3 expression on T cells [11], in CLE fluid compared to healthy controls,

previously not reported in CLE skin analyses. A 19 color flow panel was used to identify cell

types of interest by both UMAP and manual gating. Spectral flow cytometry analysis of blister

fluid cells revealed a predominantly lymphocytic infiltrate, with increases in myeloid populations

as compared to healthy (**Fig 1D**).

Spatial transcriptomics and proteomics on archival tissue biopsies was performed using

NanoString Digital Spatial Profiling (DSP, **Fig 1E, Table S2**). Spatial proteomics captured

enrichment of CD8 and CD14 as well as beta-2 microglobulin (B2M) and indoleamine 2,3-

dioxygenase (IDO) (**Fig 1F,** dataset QC in **Fig S1,** QC of regions of interest (ROIs) in **Fig S2**).

Spatial transcriptomics analysis using the Whole Transcriptome Atlas (WTA) of aggregate ROIs

recapitulated IFN signatures, and also harmonized with protein biomarkers identified in 96-plex

immunoassay and protein DSP (**Fig 1G**). We also validated our DSP WTA dataset using a

Cancer Transcriptome Atlas (CTA) dataset (**Table S3**), as well as a bulk RNA microarray

dataset previously generated from matched tissue blocks (**Fig S3 & 4**). These datasets had

many shared differentially expressed genes (DEGs), including *B2m, Cxcl9, Cxcl10* and other

IFN- and immune-related genes.

Characterization of the immune infiltrate in CLE interface dermatitis

We sought to characterize which cells comprised the interface dermatitis in CLE using

multimodal approaches. To do this, we performed spectral flow cytometry and 96-plex

immunoassay proteomics on blister biopsy fluid, and spatial transcriptomics and proteomics on

FFPE tissues to determine the cellular composition of CLE interface dermatitis. Quantification of

cellular infiltrates from blister biopsies revealed significant increases in HLA-DR+ antigen

presenting cell populations (p<0.0021 lesional vs nonlesional, p<0.0006 nonlesional vs healthy)

as well as CD8+ T cells (p<0.0166 lesional vs healthy, **Fig 2A**). Querying the 96-plex

immunoassay proteomics dataset for cell type specific proteins identified increased T cell (CD5,

CD8A) and APC markers (CD302) in both lesional and nonlesional blister fluid (**Fig 2B**). These

surface markers are likely shed due to the presence of increased matrix metalloproteinases

such as ADAM15 (**Fig 2B**). Additional significant protein biomarkers are presented in **Table S4**.

Next, we examined our cell type regions of interest (ROIs) in the spatial datasets. CD3+CD45+

masks were used to enrich for T cells, CD45+CD3- masks to enrich for other immune cells,

geometric autofluorescent ROIs to enrich for keratinocytes and SMA+ ROIs to enrich for

endothelium (**Fig 2C**). We were not able to capture many ROIs from endothelium in cross

section, therefore we focused our analyses on the T cell, immune cell and keratinocyte ROIs.

NanoString GeoMX software deconvolution of the immune cell ROIs in the WTA dataset

revealed increases in many immune cell types in the dermis including B cells, T cells, dendritic

cells and monocytes/macrophages as assessed by cell-type specific genes (**Fig 2D**).

Examination of protein-level expression of CD surface markers using the immune cell profiling

and immune cell typing protein modules confirmed CD20+ B cell, CD14+ and CD68+

monocyte/macrophage expression (**Fig 2E**). Taken together, the predominant cell types

comprising CLE interface dermatitis are HLADR+ myeloid cells and CD8+ T cells.

Examining differences in discoid versus subacute cutaneous lupus erythematosus

reveals unique gene and immune checkpoint expression

One of the remaining questions in the field of cutaneous lupus is what immunologic differences

underlie the pathogenesis of discoid lupus erythematosus (DLE) versus subacute cutaneous

lupus erythematosus (SCLE). To this end, we separated out the CLE subtypes and performed

both transcriptomic and proteomic analyses comparing the 2 entities, including donor as a

correction variable. Spatial transcriptomic analysis of DLE vs. SCLE skin detected 19 DEGs in

keratinocyte ROIs using a log2 fold change cutoff of +/- 1 and p value cutoff of 0.05 (**Fig 3A &**

B). Pathway analysis of these genes revealed terms related to G protein coupled receptors

(GPCRs) enriched in SCLE versus terms involved in RNA processing and interferon signaling in

DLE (**Fig 3C**). We also performed hypothesis testing by querying specific subsets of genes

across conditions. Examining calpains, which were recently described as a treatment target for

photosensitivity[12], revealed that *CAPN1* was enriched in SCLE, whereas *CAPN3* was enriched

in DLE (**Fig 3D**). Analysis of CD45+ ROIs detected 27 DEGs in DLE vs SCLE (**Fig 3E & F**).

Pathway analysis of these genes revealed keratinization was enriched in SCLE, whereas

interferon alpha/beta signaling and rRNA processing terms were enriched in DLE (**Fig 3G**). Of

note, the keratinization term includes immune-related genes such as calpains and cathepsins.

Examining IL-1 Receptor Associated Kinases (IRAKs), which were recently described as a

treatment target for skin inflammation[13], revealed *IRAK1* was enriched in DLE, whereas *IRAK4*

was enriched in SCLE immune cells (**Fig 3H**). Analysis of CD3+ ROIs detected 4 DEGs in DLE

vs SCLE (**Fig 3I & J**). Since so few genes were above a log2 fold change cutoff of 1, we

performed analysis of differentially expressed proteins in place of pathway analysis.

Interestingly, the immune inhibitory family member B7-H3 was uniquely upregulated in SCLE T

cells (**Fig 3K**). Examining Janus kinases (JAKs), which have been used for multiple

inflammatory skin disorders, revealed *TYK2* is enriched in both DLE and SCLE T cells

compared to healthy (**Fig 3L**). Taken together, these data suggest that nuances in immune cell

and keratinocyte signaling may underlie DLE versus SCLE pathogenesis, and that some

treatment targets such as B7-H3, calpains or IRAKs may be subtype specific, while others, like

TYK2, are shared.

Keratinocytes at the inflammatory and lesional sites shape chemotactic signals for

leukocyte recruitment

We next asked whether we could identify immune signaling pathways in keratinocytes that could

lead to the initiation of inflammation. We selected regions of interest above interface dermatitis

that exhibit hyperkeratosis and/or acantholysis in overlying keratinocytes. Comparing

keratinocytes that were above interface dermatitis (proximal) versus distal to an infiltrate (**Fig**

4A) revealed upregulation of key chemokines including *Cxcl9* and *Cxcl10* in keratinocytes

proximal to the immune infiltrate (**Fig 4B**). 96-plex immunoassay proteomic data in lesional and

nonlesional blister fluid was compared to the RNA spatial dataset, and demonstrated that

CXCL9 and CXCL11 proteins are significantly upregulated in lesional blister fluid (**Fig 4C**).

Additionally, increased levels of chemokine ligands CCL8 and CXCL6 in lesional versus

nonlesional blister fluid were identified (**Fig 4C**). We re-queried the WTA dataset to analyze

potential innate immune pathways that could lead to chemokine expression and found

increased *AIM2* expression (**Fig 4D**). Querying specific members of the AIM2 pathway in the

96-plex immunoassay dataset identified significant increases in caspase 8 [14] and the IL1 family

member IL18 [15] in lesional skin relative to both healthy and nonlesional samples (**Fig 4E & F**).

96-plex immunoassay also identified significant increases in IFNL1 and IFNG protein levels in

lesional CLE skin, which are upstream of CXCL9 and CXCL11 (**Fig 4G & H**). Of note, IL18

induces CXCL6 expression in murine lung[16], and CCL8 is optimally induced by combinations of

IL1, IFNG and dsRNA[17]. Taken together, these data suggest that keratinocyte injury, as

evidenced by hyperkeratosis and acantholysis visible on histology, can induce AIM2-caspase 8-

IL18 pathways and IFN pathways[18] to turn on chemokines including CCL8, CXCL6, CXCL9 and

CXCL11 for recruitment of immune cells in CLE.

Combining proteomic analysis of blister fluid with spatial transcriptomics of skin tissue

reveals cell-specific expression patterns of chemokine ligands in CLE skin

We[19–21] and others[9,22,23] have demonstrated upregulation of chemokine ligands in CLE by RNA

sequencing and microarray. 96-plex immunoassay proteomics identified significant increases in

many chemokine ligands at the protein level, including CXCR3 family, CCL8, and CXCL6

chemokines (**Fig 5A**). Chemokines were fairly stable over time, as determined by repeat

blistering of one of our donors approximately 6 months apart (**Fig S5**). Querying the WTA

dataset revealed cell-specific context expression of these ligands which differed by both ROI

type and disease state (**Fig 5B**). Specifically, *Ccl8* was expressed highly in the CD45+ ROI from

DLE biopsies, whereas *Ccl25* was expressed in SCLE biopsies, indicating a different cellular

context of these signals. In contrast, *Cxcl6* was expressed by both keratinocytes and immune

cells. *Cxcl9/10/11* were expressed mainly by the immune system, with the highest upregulation

in DLE, which fits with prior literature describing an IFNG "node" in DLE[9]. Taken together, these

data help to map cellular sources of chemokines, which in turn provides insights into the order in

which they are produced, whether by skin resident immune cells, keratinocytes or infiltrating

immune cells.

Functional chemotaxis assays recapitulate CXCR3- mediated T cell migration and reveal

CXCR1-sensitive CD14+CD16+ and CCR2-sensitive CD14+ populations

To determine the functional consequences of our data, blood was drawn from healthy and CLE

donors and used in chemotaxis assays, focusing on the ligands we identified in the 96-plex

immunoassay dataset (CCL8, CXCL6, CXCL9, CXCL11; **Table S5**). Flow cytometry was also

performed on input cells prior to migration to determine distribution of the relevant chemokine

receptors (antibody information & chemokine information in **Table S6 & Table S7**). CCL8 binds

to CCR1/2/3/5; CXCL6 binds to CXCR1/2, and CXCL9 and CXCL11 both bind to the receptor

CXCR3. CCR3 is typically expressed on eosinophils, so we did not assess this receptor, but

instead assessed CCR2/5 on monocytes which are the high affinity CCL8 receptors. Cells were

gated on live singlets and were identified with major lineage markers (e.g. CD3+ T cells, CD19+

B cells, CD56+ NK cells). Both lupus and healthy donor T cells expressed CXCR3 (**Fig 6A & B**)

and were able to migrate well towards CXCL11 (**Fig 6C**). However, lupus donor T cells only

exhibited a 2-fold increase in migration over media baseline in lupus patient samples, compared

to an average 5-fold increase in migration over media baseline in healthy samples (**Fig 6D**). We

hypothesize this may be due to maintenance therapy in lupus donors. Notably, CXCL9 and

CXCL11 are off in healthy skin, and turned on in lupus skin, exemplifying how T cells are poised

to respond to IFN-inducible chemokines.

We next assessed CXCL6 and its cognate receptor CXCR1[24], which has been reported to be

expressed on skin NK cells[25] and is important for their recruitment into tumors[26]. NK cells from

both healthy and lupus donors expressed CXCR1 (**Fig S6A**), though there was donor variability

for migratory capacity towards the CXCL6 ligand. Like T cells, NK cells migrated to CXCL9 and

CXCL11 (**Fig S6B & C**). We also assessed B cell migration, which we observed to be

dependent on CXCR5-CXCL13 (**Fig S6D-F**). These findings fit with current literature describing

NK cell and B cell migration pathways.

Last, we assessed chemokine expression on myeloid cells, which likely constitute the HLA-DR+

population we noted to be increased in blister biopsies. CD14+CD16+ myeloid cells were

recently identified in nonlesional CLE skin by Billi et al using 10X spatial transcriptomics and

were hypothesized to be the first responders to skin injury/inflammation in lupus[27]. We gated

live, single, CD45+CD3-CD19-CD56- cells (i.e. non-T/B/NK) followed by CD14 and CD16

expression (**Fig S7**) to confirm these cells are present in CLE blood and skin by flow cytometry

(**Fig S8**). We examined chemokine receptor expression in CD14+CD16-, CD14+CD16+ and

CD14-CD16+ myeloid cells and noted significant increases in CXCR1 expression on

CD14+CD16+ myeloid cells (**Fig 6E &F**). Total number of monocytes were higher at baseline in

healthy versus lupus patients, which we hypothesize may be due to maintenance therapy in

lupus patients. All CD14+ myeloid cells, but particularly CD14+CD16+ myeloid cells, migrated

towards CXCL6, providing a possible chemokine receptor:ligand pair that mediates their entry

into the skin (**Fig 6G**). To assess a potential inhibitor of this pathway, we treated healthy donor

PBMCs with the CXCR1/2 inhibitor reparixin (REPA) and assessed migration towards CXCL6.

REPA treatment reduced migration in all but one donor, who also paradoxically did not migrate

well towards CXCL6 (**Fig 6H**). Taken together, these data identify a potential inhibitor of

CD14+CD16+ cell entry into lupus skin, which could prevent new lesion formation.

In contrast to CXCR1, the CCL8 receptor CCR2 was enriched on CD14+CD16- myeloid cells

(**Fig 6I & J**), but not the CCR5 receptor (**Fig S9**). CD14+CD16- myeloid cells migrated towards

CCL8 (**Fig 6K**). To understand the potential effects of hydroxychloroquine (HCQ) on this

pathway, we pretreated healthy PBMCs and subjected them to chemotaxis assays. HCQ

treatment reduced CCL8 directed migration by CD14+CD16- monocytes (**Fig 6L**).

Taken together, these data exemplify how disease states can influence chemokine ligand levels

and influence timing and recruitment of specific immune cell populations to the tissue, namely

through keratinocytes following injury and subsequently through recruited immune cell

populations. We propose the model that keratinocyte injury induces activation of AIM2-related

pathways that turn on CXCL6 and CCL8 expression to recruit myeloid cells, which in turn

potentiate IFN-related signaling pathways to recruit T cells and other lymphocytes via CXCR3

ligands (**Graphical abstract**).

DISCUSSION

In this study, we sought to analyze chemokine orchestrators of T cell rich interface dermatitis in

CLE. Chemokines can be released by keratinocytes in response to UV light, infection, or other

environmental triggers[28]. These chemokines can then recruit immune cells to the skin, which can

lead to inflammation. One of the most highly upregulated chemokine families in CLE lesions is

the CXCR3 chemokine family[6,29,30]. CXCR3 binds to its interferon (IFN)-inducible ligands CXCL9,

CXCL10 and CXCL11 to mediate leukocyte migration and function[31]. IFN signaling has been

postulated to drive the pathogenesis of all subtypes CLE[32]. Further, CLE is a photosensitive

disorder, and UV light induces upregulation of IFN to amplify CXCR3 ligand production in

keratinocytes[18,33,34], linking the environmental insult to the recruitment of pathogenic immune cells

in the context of CLE. The interaction between CXCR3-expressing T-cells and its ligands CXCL9,

CXCL10 and CXCL11 have been associated with tissue damage in several CLE subtypes[8].

Examining chemokine ligand expression in our 96-plex immunoassay dataset and in our ROIs,

as well as chemokine receptor expression in our flow cytometry data, confirmed expression of the

CXCR3 family and also identified potential novel mediators of recruitment in CLE. Based on 96-

plex immunoassay and flow cytometry data, we hypothesize that CXCR1 is a key receptor for

CXCL6 on proinflammatory CD14+CD16+ APCs, and could be a viable treatment target for CLE,

especially given the role of CXCL6 in fibrosis which occurs in DLE and other forms of chronic

CLE[35] as well as in renal interstitial fibrosis[36]. CXCL6 is a particularly interesting target

immunologically, given that it may be upregulated by double stranded RNA or IL1β[37]. This is

important, as we know from work by Rodriguez-Pla et al. that IFN priming alone is not sufficient

to activate migratory programs in lupus monocytes[38], and from work by Billi et al. that

CD14+CD16+ monocyte/dendritic/myeloid cell populations are among the first cells recruited to

the skin in CLE[27]. Further studies are needed to understand epigenetic factors that result in

increased CXCR1 expression in CD14+CD16+ myeloid cells.

The 96-plex immunoassay dataset provides insight into novel protein biomarkers of CLE

beyond chemokine ligands that have interesting implications in immunopathogenesis, including

Flt3L, HGF, CD40 and CEACAM3. Flt3L is a dendritic cell growth and survival factor, and has

been previously described in SLE[39]. A recent study demonstrated generation of tolerogenic

conventional DCs in SLE patients undergoing mesenchymal stem cell therapy[40]. Thus, it is

possible that the upregulation of Flt3L in skin is a compensatory mechanism. HGF, which as

noted above can increase CXCR3 expression on T cells[11], is higher in SLE patients than

healthy controls[41] and in patients with higher disease activity[42]. HGF administration can prevent

lupus nephritis in mice[43], potentially indicating a compensatory mechanism for HGF in CLE.

CD40 overexpression causes autoimmune skin disease in mice[44], and increased CD40

expression has been reported in IHC of CLE lesions[45]. Genome wide differential expression

analysis identified CEACAM3 as a potential gene involved in the pathogenesis of lupus and

lupus nephritis[46], though its role in skin disease has yet to be elucidated.

We also found differences in DLE and SCLE that have not previously been appreciated by

using spatial technologies. For example, DLE lesional skin T cells express less B7-H3 than

SCLE lesional skin as assessed by protein DSP. B7-H3, also known as CD276, is an immune

checkpoint within the B7 molecular family that fine tunes immune responses[47] and is currently

being explored as a cancer immunotherapy target [48]. B7-H3 plays a pivotal role in mediating the

suppressive effects of DCs on T cells and downregulating Th1 responses[49]. Therefore, a

relative reduction in B7-H3 in DLE as compared to SCLE fits with previously described IFNG

signaling nodes identified in DLE by bulk RNA sequencing[9]. Further studies are needed to

assess the DEGs for DLE vs SCLE ROI comparisons that were identified in the WTA dataset.

In addition to insights into immune cell recruitment and potential novel biomarkers, our datasets

identify potential treatment targets. Reparixin and the related drug ladarixin are CXCR1

inhibitors that are currently in clinical trials for cancer[50]. It would be interesting to test a topical

formulation for CLE for prevention of new or spreading lesions. A calpain 1 inhibitor was

recently demonstrated to prevent photosensitivity in animal models of UVB exposure [12]. We

noted elevated *CAPN1* in SCLE keratinocytes, which makes this another attractive topical

treatment to test. TYK2 inhibitors are currently in clinical trials for SLE[51], and several case

reports have noted improvement in CLE[52]. We found elevated *TYK2* in T cell ROIs from both

DLE and SCLE patients. A small molecule inhibitor of IRAK4 was recently described as a

treatment target for psoriatic skin inflammation [13]. We identified elevated *IRAK4* in DLE, and

IRAK1 in SCLE immune cells. Given that both TYK2 and IRAKs were elevated in the immune

compartment ROIs, these treatments may require oral rather than topical administration for

highest efficacy. We hope that these data inspire future clinical trials repurposing drugs for CLE,

as well as studies testing topical versus oral routes of administration for more advanced

compounds.

While our study provides valuable insights into the immunopathogenesis of CLE, it is essential

to acknowledge the study's limitations. First, our research predominantly focuses on a specific

subset of chemokine ligand-receptor pairs and immune cell interactions that we were able to

measure with current methodologies. CXCR1 is a classical receptor for neutrophil recruitment to

inflamed tissues, and ligation of CXCR1 can enhance NET formation[53,54]. We were not able to

recover neutrophils from blister fluid, and opted to use Ficoll gradients, which remove the bulk of

granulocytes, to isolate PBMCs for chemotaxis assays. Future investigations should include

methodologies to specifically assess neutrophils and other granulocytes. Moreover, a

longitudinal approach with serial assessments would offer a more subtle understanding of

disease progression over time. Finally, our findings in different CLE subtypes and patient

populations should be examined further with larger patient cohorts. Future studies exploring

longitudinal analyses with more patients, and comparing myeloid cells in the skin of patients

with response or resistance to antimalarials, are warranted to confirm and expand upon these

findings. In-depth mechanistic studies and the development of targeted interventions based on

the identified chemokine and other pathways holds promise for improving therapeutic strategies

in CLE.

List of Supplementary Materials

Supplemental Materials & Methods

Fig S1. Quality Control for Digital Spatial Profiling (DSP) Whole Transcriptome Atlas (WTA) spatial
transcriptomics.

Fig S2. Quality Control and Pathway Analysis for Regions of Interest (ROIs).

Fig S3. Validation of DSP dataset using historical microarray dataset and a cancer transcriptome
atlas (CTA) dataset.

Fig S4. Validation of DLE dataset ROIs as compared to CTA.

Fig S5. Stability of chemokines in lesional and nonlesional samples over time.

Fig S6. Assessment of NK cell and B cell migration.

Fig S7. Flow gating strategy for assessing myeloid populations.

Fig S8. Examination of CD14+CD16+ cells in blister biopsies

Fig S9. CCR5 is not enriched on CD14 vs CD16 expressing myeloid cells.

Table S1. Blister biopsy and blood donation patient info & characterization (UMass Chan).

Table S2. Archival CLE biopsies used in WTA Digital Spatial Profiling (UMass Chan).

Table S3. Validation archival CLE biopsies used in CTA Digital Spatial Profiling (Yale).

Table S4. 96-plex immunoassay DEPs calculated by NPX software and 2-way ANOVA.

Table S5. Chemotaxis and chemokine receptor staining blood donor information (UMass Chan
and Dartmouth Hitchcock).

Table S6. Antibody information & RRIDs.

Table S7. Chemokine information.

521
522

Figures & Legends:

Fig. 1 Integration of high-resolution immune techniques with spatial approaches for studying cutaneous lupus erythematosus. (**A**) Study overview demonstrating workflow. We performed blister biopsies on cutaneous lupus erythematosus patients and healthy donors and analyzed the interstitial skin fluid using 96-plex immunoassay proteomics and the cell pellets using Aurora spectral flow cytometry. We also employed biobanked FFPE skin biopsies from DLE, SCLE and healthy margins for NanoString digital spatial profiling (DSP) for protein using 6 modules, and RNA using the whole transcriptome atlas (WTA). Created with Biorender.com. (**B**)

Sample photographs of blister biopsies (n=3 healthy and 4 CLE patients). (**C**) H&E images of the
slide set used for spatial transcriptomics and proteomics. (n=3 healthy margin controls, 5 SCLE
biopsies from 4 patients, and 4 DLE biopsies from 4 patients. * indicates tissue was not fully
seated in imaging area for WTA run). (**D**) Volcano plot of 96-plex immunoassay data from the
inflammation and neuroexploratory panels of all CLE blisters versus healthy controls (n=7 CLE
lesional, 7 CLE nonlesional blisters from 4 donors, and 3 healthy blisters). (**E**) Sample UMAP of
immune infiltrates in CLE lesional skin. (representative patient of 4 CLE patients) (**F**) Volcano plot
of DSP protein data for all CLE regions of interest (ROIs) versus healthy margin ROIs. (n=95 ROIs
assessed from 4 DLE, 3 SCLE and 3 healthy margin controls for protein DSP) (**G**) Volcano plot
of DSP RNA WTA data for all CLE ROIs versus healthy margin ROIs. Red dots are also
differentially expressed proteins (DEPs) in the 96-plex immunoassay protein dataset, and cyan
dots are also DEPs in the DSP protein dataset. (n=41 ROIs assessed from 4 DLE, 3 SCLE, and
2 healthy margin control biopsies for WTA DSP.) (FFPE, formalin fixed paraffin embedded; DLE,
discoid lupus erythematosus; SCLE, subacute cutaneous lupus erythematosus; WTA, whole
transcriptome atlas; DSP, digital spatial profiling; UMAP, Uniform Manifold Approximation and
Projection; DEP, differentially expressed protein; DEG, differentially expressed gene; ROI, region
of interest).

Fig. 2. Characterizing immune cell subsets in CLE interface dermatitis reveals significant increases in HLA-DR+ cells and CD8+ T cells. (**A**) Aurora flow cytometry for cell lineage

markers in CLE lesional and nonlesional skin compared to healthy blisters. Enumeration of
cellular infiltrates were normalized to 10,000 live singlet cells. One-way ANOVA with Tukey's post
hoc tests revealed significant increases in HLA-DR+ cells and CD8+ T cells compared to healthy
control blisters, with trends towards increased B cells, CD4+ T cells and NK cells. pDCs were
trending higher in nonlesional blister fluid. (n=3 healthy and 4 CLE patients) (**B**) 96-plex
immunoassay targeted proteomics for cell lineage markers in blister fluid. T cell markers (CD5,
CD8A) and antigen presenting cell markers (CD302) were also significantly elevated in CLE
blister fluid compared to healthy controls. NK marker CD244 was not significantly elevated,
matching flow cytometry findings. Note that shedding of surface CD molecules is likely due to the
presence of proteases such as ADAM15. (n=7 CLE lesional, 7 CLE nonlesional from 4 donors
and 3 healthy blisters) (**C**) Examples of Regions of interest (ROIs) chosen for spatial analysis of
FFPE CLE biopsies. We used cell masking approaches for CD3+ T cells, CD45+ immune cells,
geometric autofluorescent keratinocytes/epidermis and SMA+ endothelium. (**D**) Cell type
deconvolution in the CD45+ dermal ROIs from the WTA dataset was calculated using
NanoString's prediction module. We noted increases in T cells and B cells, as well as neutrophils
which were not assessed by flow cytometry due to neutrophil death upon extraction from skin.
(n=8 DLE, 6 SCLE and 2 healthy margin CD45+ ROIs) (**E**) Spatial proteomics was assessed for
the immune cell profiling and immune cell typing modules to predict the presence of immune cells.
Housekeeping proteins are depicted in the bottom module. (n=95 ROIs assessed from 4 DLE, 4
SCLE from 3 donors and 3 healthy margin controls for protein DSP)

Fig. 3. Examining differences in DLE versus SCLE skin biopsies reveals biological differences that may underlie pathogenesis and/or susceptibility to systemic disease. (**A**) Example keratinocyte/epidermal region of interest (ROI) cartoon. (**B**) Volcano plot of the keratinocyte ROI in DLE vs SCLE biopsies. (n=5 SCLE vs 6 DLE keratinocyte ROIs) (**C**) Pathway analysis of the keratinocyte ROI in DLE vs SCLE biopsies as performed in GeoMX software. (**D**) Calpain gene family expression in keratinocytes reveals differences in DLE vs SCLE. (two-way ANOVA with Tukey's posttests significant or trending, as indicated) (**E**) Example CD45+ ROI cartoon. (**F**) Volcano plot of the CD45+ ROI in DLE vs SCLE biopsies. (n=5 SCLE vs 5 DLE CD45+ ROIs) (**G**) Pathway analysis of the CD45+ ROI in DLE vs SCLE biopsies. (**H**) IL-1 Receptor Associated Kinase (IRAK) gene family expression in CD45+ cells reveals differences in DLE vs SCLE. (two-way ANOVA with Tukey's posttests significant or trending, as indicated) (**I**) Example CD3+ ROI cartoon. (**J**) Volcano plot of the CD3+ ROI RNA DSP in DLE vs SCLE biopsies. (n=5 SCLE vs 5 DLE CD3+ ROIs) (**K**) Volcano plot of the CD3+ ROI protein DSP in DLE vs SCLE biopsies. (n= 11 DLE, 6 SCLE and 3 healthy CD3+ ROIs) The immune checkpoint B7-H3 is unique to SCLE and may serve as a novel treatment target. (**L**) Janus kinase (JAK) gene family expression in DLE vs SCLE T cells reveals the treatment target *TYK2* is trending

591 higher in both DLE and SCLE. (n=5 SCLE vs 5 DLE CD3+ ROIs, two-way ANOVA with Tukey's
592 posttests significant or trending, as indicated) Cartoons created with Biorender.com

Fig 4. Examining proximal versus distal keratinocytes, and lesional versus nonlesional blister fluid, reveals pathways that may be required for initiation of chemokine cascades for leukocyte recruitment to form interface dermatitis. (**A**) Schematic of proximal versus distal analysis. Created with Biorender.com. (**B**) Volcano plot of DSP WTA data for proximal versus distal epidermis. (n=9 proximal and 4 distal plus healthy margin keratinocyte ROIs) (**C**) Examination of lesional versus nonlesional blister fluid in 96-plex immunoassay as performed with NPX software. (n= 9 lesional and 9 nonlesional blister fluid samples from 6 patients). (**D**) Box plot of *AIM2* expression in keratinocytes from distal plus healthy margins versus proximal keratinocytes from Q3 normalized counts in the WTA dataset. (n=4 healthy plus distal and 9 proximal ROIs) (**E**) Caspase 8, (**F**) IL-18, (**G**) IFNL1 and (**H**) IFNG were elevated in CLE blisters compared to healthy controls in the 96-plex immunoassay dataset (Normalized Protein eXpression, NPX; panels E, F, H & I: n=9 lesional and 9 nonlesional blister fluid samples from 6 patients, and 5 healthy controls run in the 96-plex immunoassay inflammation panel; panel G: n=7 lesional and 7 nonlesional blister fluid samples from 4 patients, and 3 healthy controls run in the 96-plex immunoassay neuroexploratory panel).

Fig. 5. Combining proteomics and spatial transcriptomics using a cell masking approach allows assignment of biomarkers and chemokine ligand-receptor pairs governing leukocyte recruitment to the skin during CLE. (**A**) We queried the 96-plex immunoassay datasets for all available chemokine measurements. Significant differences were observed in the levels of CCL7, CCL8, CXCL1, CXCL6, CXCL9, CXCL11 and CX3CL1 in the 96-plex immunoassay interstitial skin fluid analysis (two-way ANOVA with Tukey's post tests significant

as indicated; n=5 healthy blisters, and 6 matched CLE lesional and nonlesional blisters). **B**) Heatmaps illustrate the cellular sources of chemokines in the WTA DSP analysis for healthy, SCLE, and DLE samples. (n=41 ROIs assessed from 4 DLE, 3 SCLE, and 2 healthy margin control biopsies for WTA DSP)

Fig. 6. Functional chemotaxis assays reveal that T cells are poised to respond to CXCR3 ligands, whereas CD14+CD16+ myeloid cells are poised to respond to CXCL6 via CXCR1 and CD14+CD16- myeloid cells are poised to respond to CCL8 via CCR2. (**A**) Example flow staining and (**B**) quantification of CXCR3 on CD3+ T cells (n=3 healthy and n=6 CLE, t test ns). (**C**) Chemotaxis assay reveals both healthy and lupus T cells migrate towards CXCL11 (n=4 healthy and 4 lupus donors; one-way ANOVA with significant posttests as indicated). (**D**) Chemotaxis assay reveals higher migration in healthy donor T cells than lupus donor T cells towards CXCL9. (n=4 healthy and 4 lupus donors; one-way ANOVA with significant posttests as

indicated). (**E**) Example flow staining of myeloid cells demonstrating CXCR1 expression in
CD14+, CD14+CD16+ and CD16+ gates. (**F**) Quantification of CXCR1+ cells demonstrates
CD14+CD16+ cells express more CXCR1 than their counterparts. (n=4 healthy and 5 lupus
donors; p=0.0122 and 0.0109 by repeated measures one way ANOVA matched by donor). (**G**)
Chemotaxis assay reveals directed migration of CD14+CD16+ myeloid cells in all but one healthy
donor (n=6 healthy and 1 lupus donor; paired t test p=0.0357). (**H**) Chemotaxis assay reveals
pretreatment of PBMCs from healthy donors with the CXCR1 inhibitor reparixin (REPA) inhibits
migration towards CXCL6. (n=5 healthy donors; one way ANOVA with significant posttests as
indicated). (**I**) Example flow staining of myeloid cells demonstrating CCR2 expression in CD14+,
CD14+CD16+ and CD16+ gates. (**J**) Quantification of CCR2+ myeloid cells demonstrates
CD14+CD16- cells are trending towards more CCR2 expression than their counterparts (n=4
healthy and 5 lupus donors; p=0.0208 by one way ANOVA with significant posttests as indicated).
(**K**) Chemotaxis assay demonstrated that CD14+CD16- cells migrate towards CCL8. (n=6 healthy
and n=3 lupus donors; paired t test p=0.0252). (**H**) Chemotaxis assay reveals pretreatment of
PBMCs from healthy donors with hydroxychloroquine (HCQ) inhibits migration towards CCL8
(n=6 healthy donors; one way ANOVA with significant posttests as indicated).

References:

Garelli CJ, Refat MA, Nanaware PP, *et al.* Current Insights in Cutaneous Lupus Erythematosus Immunopathogenesis. *Front Immunol* 2020; **11**:1353.

Schultz HY, Dutz JP, Furukawa F, *et al.* From pathogenesis, epidemiology, and genetics to definitions, diagnosis, and treatments of cutaneous lupus erythematosus and dermatomyositis: a report from the 3rd International Conference on Cutaneous Lupus Erythematosus (ICCLE) 2013. *J Invest Dermatol* 2015; **135**:7–12.

Reich A, Werth VP, Furukawa F, *et al.* Treatment of cutaneous lupus erythematosus: current practice variations. *Lupus* 2016; **25**:964–72.

Chang AY, Werth VP. Treatment of cutaneous lupus. *Curr Rheumatol Rep* 2011; **13**:300–7.

Stannard JN, Kahlenberg JM. Cutaneous lupus erythematosus: updates on pathogenesis and associations with systemic lupus. *Curr Opin Rheumatol* 2016; **28**:453–9.

Wenzel J, Zahn S, Mikus S, *et al.* The expression pattern of interferon-inducible proteins reflects the characteristic histological distribution of infiltrating immune cells in different cutaneous lupus erythematosus subsets. *Br J Dermatol* 2007; **157**:752–7.

Wenzel J, Henze S, Wörenkämper E, *et al.* Role of the Chemokine Receptor CCR4 and its Ligand Thymus- and Activation-Regulated Chemokine/CCL17 for Lymphocyte Recruitment in Cutaneous Lupus Erythematosus. *J Invest Dermatol* 2005; **124**:1241–8.

Méndez-Flores S, Hernández-Molina G, Azamar-Llamas D, *et al.* Inflammatory chemokine profiles and their correlations with effector CD4 T cell and regulatory cell subpopulations in cutaneous lupus erythematosus. *Cytokine* 2019; **119**:95–112.

Berthier CC, Tsoi LC, Reed TJ, *et al.* Molecular Profiling of Cutaneous Lupus Lesions Identifies Subgroups Distinct from Clinical Phenotypes. *J Clin Med Res* 2019; **8**. doi:10.3390/jcm8081244.

Strassner JP, Rashighi M, Ahmed Refat M, *et al.* Suction blistering the lesional skin of vitiligo patients reveals useful biomarkers of disease activity. *J Am Acad Dermatol* 2017; **76**:847–55.e5.

Komarowska I, Coe D, Wang G, *et al.* Hepatocyte Growth Factor Receptor c-Met Instructs T Cell Cardiotropism and Promotes T Cell Migration to the Heart via Autocrine Chemokine Release. *Immunity* 2015; **42**:1087–99.

Doleckova I, Vidovic T, Jandova L, *et al.* Calpain inhibition protects against UVB-induced degradation of dermal-epidermal junction-associated proteins. *J Invest Dermatol* 2024. doi:10.1016/j.jid.2024.02.020.

Lavazais S, Jargosch M, Dupont S, *et al.* IRAK4 inhibition dampens pathogenic processes driving inflammatory skin diseases. *Sci Transl Med* 2023; **15**:eabj3289.

Pierini R, Juruj C, Perret M, *et al.* AIM2/ASC triggers caspase-8-dependent apoptosis in Francisella-infected caspase-1-deficient macrophages. *Cell Death Differ* 2012; **19**:1709–21.

15   Ratsimandresy RA, Indramohan M, Dorfleutner A, Stehlik C. The AIM2 inflammasome is a
central regulator of intestinal homeostasis through the IL-18/IL-22/STAT3 pathway. *Cell Mol*
Immunol 2016; **14**:127–42.

16   Ishikawa Y, Yoshimoto T, Nakanishi K. Contribution of IL-18-induced innate T cell activation
to airway inflammation with mucus hypersecretion and airway hyperresponsiveness. *Int*
Immunol 2006; **18**:847–55.

17   Struyf S, Proost P, Vandercappellen J, *et al.* Synergistic up-regulation of MCP-2/CCL8
activity is counteracted by chemokine cleavage, limiting its inflammatory and anti-tumoral
effects. *Eur J Immunol* 2009; **39**:843–57.

18   Skopelja-Gardner S, An J, Tai J, *et al.* The early local and systemic Type I interferon
responses to ultraviolet B light exposure are cGAS dependent. *Sci Rep* 2020.URL
https://www.nature.com/articles/s41598-020-64865-w.

19   Ko W-CC, Li L, Young TR, *et al.* Gene expression profiling in skin reveals strong similarities
between subacute and chronic cutaneous lupus that are distinct from lupus nephritis. *J*
Invest Dermatol 2021. doi:10.1016/j.jid.2021.04.030.

20   Mande P, Zirak B, Ko W-C, *et al.* Fas ligand promotes an inducible TLR-dependent model
of cutaneous lupus-like inflammation. *J Clin Invest* 2018; **128**:2966–78.

21   Garelli CJ, Wong NB, Piedra-Mora C, *et al.* Shared inflammatory and skin-specific gene
signatures reveal common drivers of discoid lupus erythematosus in canines, humans and
mice. *Current Research in Immunology* 2021. doi:10.1016/j.crimmu.2021.03.003.

22   Dey-Rao R, Smith JR, Chow S, Sinha AA. Differential gene expression analysis in CCLE
lesions provides new insights regarding the genetics basis of skin vs. systemic disease.
Genomics 2014; **104**:144–55.

23   Jabbari A, Suárez-Fariñas M, Fuentes-Duculan J, *et al.* Dominant Th1 and Minimal Th17
Skewing in Discoid Lupus Revealed by Transcriptomic Comparison with Psoriasis. *J Invest*
Dermatol 2014; **134**:87–95.

24   Wuyts A, Proost P, Lenaerts JP, *et al.* Differential usage of the CXC chemokine receptors 1
and 2 by interleukin-8, granulocyte chemotactic protein-2 and epithelial-cell-derived
neutrophil attractant-78. *Eur J Biochem* 1998; **255**:67–73.

25   Ottaviani C, Nasorri F, Bedini C, *et al.* CD56brightCD16(-) NK cells accumulate in psoriatic
skin in response to CXCL10 and CCL5 and exacerbate skin inflammation. *Eur J Immunol*
2006; **36**:118–28.

26   Ng YY, Tay JCK, Wang S. CXCR1 Expression to Improve Anti-Cancer Efficacy of
Intravenously Injected CAR-NK Cells in Mice with Peritoneal Xenografts. *Mol Ther*
Oncolytics 2020; **16**:75–85.

27   Billi AC, Ma F, Plazyo O, *et al.* Nonlesional lupus skin contributes to inflammatory education
of myeloid cells and primes for cutaneous inflammation. *Sci Transl Med* 2022;
14:eabn2263.

28   Jiang Y, Tsoi LC, Billi AC, *et al.* Cytokinocytes: the diverse contribution of keratinocytes to

immune responses in skin. *JCI Insight* 2020; **5**. doi:10.1172/jci.insight.142067.

29   Flier J, Boorsma DM, van Beek PJ, *et al.* Differential expression of CXCR3 targeting
chemokines CXCL10, CXCL9, and CXCL11 in different types of skin inflammation. *J Pathol*
2001; **194**:398–405.

30   Wenzel J, Wörenkämper E, Freutel S, *et al.* Enhanced type I interferon signalling promotes
Th1-biased inflammation in cutaneous lupus erythematosus. *The Journal of Pathology: A*
Journal of the Pathological Society of Great Britain and Ireland 2005; **205**:435–42.

31   Groom JR, Luster AD. CXCR3 ligands: redundant, collaborative and antagonistic functions.
Immunol Cell Biol 2011; **89**:207–15.

32   Robinson ES, Werth VP. The role of cytokines in the pathogenesis of cutaneous lupus
erythematosus. *Cytokine* 2015; **73**:326–34.

33   Meller S, Winterberg F, Gilliet M, *et al.* Ultraviolet radiation--induced injury, chemokines,
and leukocyte recruitment: an amplification cycle triggering cutaneous lupus
erythematosus. *Arthritis & Rheumatism* 2005; **52**:1504–16.

34   Scholtissek B, Zahn S, Maier J, *et al.* Immunostimulatory Endogenous Nucleic Acids Drive
the Lesional Inflammation in Cutaneous Lupus Erythematosus. *J Invest Dermatol* 2017;
137:1484–92.

35   Dai C-L, Yang H-X, Liu Q-P, *et al.* CXCL6: A potential therapeutic target for inflammation
and cancer. *Clin Exp Med* 2023. doi:10.1007/s10238-023-01152-8.

36   Sun M-Y, Wang S-J, Li X-Q, *et al.* CXCL6 Promotes Renal Interstitial Fibrosis in Diabetic
Nephropathy by Activating JAK/STAT3 Signaling Pathway. *Front Pharmacol* 2019; **10**:224.

37   Wuyts A, Struyf S, Gijsbers K, *et al.* The CXC Chemokine GCP-2/CXCL6 Is Predominantly
Induced in Mesenchymal Cells by Interleukin-1β and Is Down-Regulated by Interferon-γ:
Comparison with Interleukin-8/CXCL8. *Lab Invest* 2003; **83**:23–34.

38   Rodriguez-Pla A, Patel P, Maecker HT, *et al.* IFN priming is necessary but not sufficient to
turn on a migratory dendritic cell program in lupus monocytes. *J Immunol* 2014; **192**:5586–
98.

39   Gill MA, Blanco P, Arce E, *et al.* Blood dendritic cells and DC-poietins in systemic lupus
erythematosus. *Hum Immunol* 2002; **63**:1172–80.

40   Yuan X, Qin X, Wang D, *et al.* Mesenchymal stem cell therapy induces FLT3L and CD1c+
dendritic cells in systemic lupus erythematosus patients. *Nat Commun* 2019; **10**:2498.

41   Robak E, Woźniacka A, Sysa-Jedrzejowska A, *et al.* Serum levels of angiogenic cytokines
in systemic lupus erythematosus and their correlation with disease activity. *Eur Cytokine*
Netw 2001; **12**:445–52.

42   Liu Y, Zheng M, Yin W-H, Zhang B. Relationship of serum levels of HGF and MMP-9 with
disease activity of patients with systemic lupus erythematosus. *Zhejiang Da Xue Xue Bao*
Yi Xue Ban 2004; **33**:340–3, 348.

43   Kuroiwa T, Iwasaki T, Imado T, *et al.* Hepatocyte growth factor prevents lupus nephritis in a

murine lupus model of chronic graft-versus-host disease. *Arthritis Res Ther* 2006; **8**:R123.

44 Mehling A, Loser K, Varga G, *et al.* Overexpression of CD40 ligand in murine epidermis
results in chronic skin inflammation and systemic autoimmunity. *J Exp Med* 2001; **194**:615–
28.

45 Caproni M, Torchia D, Antiga E, *et al.* The CD40/CD40 ligand system in the skin of patients
with subacute cutaneous lupus erythematosus. *J Rheumatol* 2007; **34**:2412–6.

46 AlFadhli S, Ghanem AAM, Nizam R. Genome-wide differential expression reveals
candidate genes involved in the pathogenesis of lupus and lupus nephritis. *Int J Rheum Dis*
2016; **19**:55–64.

47 Yi KH, Chen L. Fine tuning the immune response through B7-H3 and B7-H4. *Immunol Rev*
2009; **229**:145–51.

48 Castellanos JR, Purvis IJ, Labak CM, *et al.* B7-H3 role in the immune landscape of cancer.
Am J Clin Exp Immunol 2017; **6**:66–75.

49 Suh W-K, Gajewska BU, Okada H, *et al.* The B7 family member B7-H3 preferentially down-
regulates T helper type 1-mediated immune responses. *Nat Immunol* 2003; **4**:899–906.

50 Piro G, Carbone C, Agostini A, *et al.* CXCR1/2 dual-inhibitor ladarixin reduces tumour
burden and promotes immunotherapy response in pancreatic cancer. *Br J Cancer* 2022;
128:331–41.

51 Morand E, Pike M, Merrill JT, *et al.* Deucravacitinib, a tyrosine kinase 2 inhibitor, in
systemic lupus erythematosus: A phase II, randomized, double-blind, placebo-controlled
trial. *Arthritis Rheumatol* 2023; **75**:242–52.

52 Bouché N, Al-Saedy MA, Song EJ. Successful treatment of refractory subacute cutaneous
lupus erythematosus with deucravacitinib. *JAAD Case Rep* 2023; **39**:93–5.

53 Teijeira Á, Garasa S, Gato M, *et al.* CXCR1 and CXCR2 Chemokine Receptor Agonists
Produced by Tumors Induce Neutrophil Extracellular Traps that Interfere with Immune
Cytotoxicity. *Immunity* 2020; **52**:856–71.e8.

54 Florey OJ, Johns M, Esho OO, *et al.* Antiendothelial cell antibodies mediate enhanced
leukocyte adhesion to cytokine-activated endothelial cells through a novel mechanism
requiring cooperation between FcγRIIa and CXCR1/2. *Blood* 2007; **109**:3881–9.

55 Edgar R, Domrachev M, Lash AE. Gene Expression Omnibus: NCBI gene expression and
hybridization array data repository. *Nucleic Acids Res* 2002; **30**:207-10.

A Aurora Flow cytometry

#cells per 10,000 live singlets

- HLADR+ — 0.0021, 0.0006, 0.0021
- plasma cells
- pDCs
- CD56+CD16+ NK cells
- CD16+ NK cells
- CD56+ NK cells
- CD4+CD8+ T cells
- CD8+ T cells — 0.0166
- CD4+ T cells
- B cells

- CLE lesional
- CLE nonlesional
- Healthy blister

B Olink Targeted Proteomics

Normalized protein expression (NPX)

- ADAM15 — 0.0060
- CD244
- CD302 — 0.0002, 0.0017
- CD8A — 0.0051, 0.0001
- CD5 — 0.0222, 0.0004

C

CD3+ ROI (T cells) | CD45+CD3- ROI (Other Immune) | Geometric ROI (Keratinocytes) | SMA+ ROI (Endothelium)

D DSP RNA WTA Cell Type Deconvolution of Dermis CD45+ ROIs

0 — 50 — 100

- macrophages
- mast cells
- B cells - naive
- B cells - memory
- plasma cells
- T cells - CD4 naive
- T cells - CD4 memory
- T cells - CD8 naive
- T cells - CD8 memory
- NK cells
- pDCs
- mDCs
- monocytes -classical
- monocytes - nonclassical/intermediate
- neutrophils
- Tregs
- endothelial cells
- fibroblasts

SCLE | DLE | Healthy

E DSP Protein Modules for Cell Type Predictions in Dermis CD45+ ROIs

0 — 5 — 10

Immune Cell Profiling:
- B2M
- CD11c
- CD20
- CD3
- CD4
- CD45
- CD56
- CD68
- CD8
- CTLA4
- Pan-cytokeratin (PanCK)
- Fibronectin
- GZMB
- HLA-DR
- Ki-67
- PD-1
- PD-L1
- SMA

Immune Cell Typing:
- CD14
- CD163
- CD34
- CD45RO
- CD66b
- FAP-alpha
- FOXP3

Housekeeping:
- Rb IgG
- Ms IgG1
- Ms IgG2a
- GAPDH
- S6
- Histone H3

SCLE | DLE | Healthy