Rabphilin-3A negatively regulates neuropeptide release, through its SNAP25 interaction

Adlin Abramian[1*], Rein I. Hoogstraaten[1*], Fiona H. Murphy[1], Kathryn F. mcDaniel[1], Ruud F. Toonen[1], Matthijs Verhage[1,2]

*These authors contributed equally to this work.

[1]Department of Functional Genomics, Faculty of Exact Science, Center for Neurogenomics and Cognitive Research, Vrije Universiteit Amsterdam and Vrije Universiteit Medical Center, Amsterdam 1081 HV, The Netherlands.

[2]Department of Clinical Genetics, Center for Neurogenomics and Cognitive Research, Amsterdam Neuroscience, Vrije Universiteit Medical Center, Amsterdam 1105 AZ, The Netherlands.

To whom correspondence should be addressed: Matthijs Verhage, PhD (matthijs@cncr.vu.nl) or Ruud Toonen, PhD (ruud.toonen@cncr.vu.nl)

Abstract

Neuropeptides and neurotrophins are stored in and released from dense-core vesicles (DCVs). While DCVs and synaptic vesicles (SVs) share fundamental SNARE/SM proteins for exocytosis, a detailed understanding of DCV exocytosis remains elusive. We recently identified the RAB3-RIM1 pathway to be essential for DCV-, but not SV exocytosis, highlighting a significant distinction between the SV- and DCV secretory pathways. Whether RIM1 is the only RAB3 effector that is essential for DCV exocytosis is currently unknown. In this study, we show that rabphilin-3A (RPH3A), a known downstream effector of RAB3A, is a negative regulator of DCV exocytosis. Using live-cell imaging at single vesicle resolution with RPH3A-deficient hippocampal neurons, we show that DCV exocytosis increased 3-fold in the absence of RPH3A. RAB3A-binding deficient RPH3A lost its punctate distribution, but still restored DCV exocytosis to WT levels when re-expressed. SNAP25-binding deficient RPH3A did not rescue DCV exocytosis. In addition, we show that RPH3A did not travel with DCVs, but remained stationary at pre-synapses. RPH3A null neurons also had longer neurites, which was partly restored when ablating all regulated secretion with tetanus neurotoxin. Taken together, these results show that RPH3A negatively regulates DCV exocytosis, potentially also affecting neuron size. Furthermore, RAB3A interaction is required for the synaptic enrichment of RPH3A, but not for limiting DCV exocytosis. Instead, the interaction of RPH3A with SNAP25 is relevant for inhibiting DCV exocytosis.

Introduction

Neuropeptides and neurotrophins are crucial neuronal signaling molecules that play diverse roles in brain development, neurogenesis and synaptic plasticity (Malva et al., 2012; Pang et al., 2004; Park & Poo, 2013; van den Pol, 2012; Zaben & Gray, 2013). These neuromodulators are sorted at the trans-Golgi network (TGN) and packaged into dense core vesicles (DCVs). Similar to neurotransmitter release from synaptic vesicles (SVs), neuropeptide release is calcium and activity-dependent. However, unlike SVs, DCVs can fuse at various locations in the neuron, requiring higher and more sustained stimulation frequencies (Balkowiec & Katz, 2002; Hartmann et al., 2001; Persoon et al., 2018). Interestingly, all SNARE/SM proteins essential for exocytosis are shared between these two secretory pathways (Arora et al., 2017; Farina et al., 2015; Hoogstraaten et al., 2020; Puntman et al., 2021; Südhof, 2013; van de Bospoort et al., 2012). However, while SV secretory pathways are well characterized, a detailed understanding of DCV exocytosis is still emerging. We recently discovered that RAB3, a protein largely dispensable for SV exocytosis (Schlüter et al., 2004, 2006), and its effector RIM1 to be essential for DCV exocytosis (Persoon et al., 2019). This indispensable role of the RAB3-RIM pathway signifies a main difference between SV and DCV exocytosis. Whether RIM is the only RAB3 effector that is essential for DCV exocytosis is currently unknown.

Rabphilin-3A (RPH3A) is a downstream effector of RAB3A that is highly expressed in the brain (Schlüter et al., 1999). RPH3A expression increases throughout development, similar to established synaptic proteins (Baldarelli et al., 2021; Blake et al., 2021; Krupke et al., 2017). RPH3A binds RAB3A and RAB27 and contains two lipid- and calcium-binding C2 domains (Guillén et al., 2013; Tsuboi & Fukuda, 2005). The second C2 domain (C2B) binds SNAP25 in a calcium independent manner (Deák et al., 2006). Similar to RAB3 null mutants, depletion of RPH3A shows no detectable synaptic phenotype in mice under normal physiological conditions (Schlüter et al., 1999), and only a mild phenotype in *C. elegans* (Staunton et al., 2001). However, mutant mice do show increased synaptic recovery after intense stimulation, which is rescued by reintroducing full length (FL) RPH3A but not a mutant that lacks the C2B domain. This suggests that RPH3A negatively regulates SV recycling after depletion of the releasable vesicle pool, and that this function depends on its interaction with SNAP25 (Deák et al., 2006). In C. *elegans*, the absence of RBF-1, the homolog of RPH3A, exacerbates the phenotypes of other SNARE mutants, as observed in the *rbf-1/ric-4* (RPH3A/SNAP25) double mutant, suggesting that RPH3A contributes to SNARE protein function (Staunton et al., 2001).

The role of RPH3A in the context of DCV exocytosis within mammalian neurons remains unclear. Notably, a recent genetic screen in C. *elegans* revealed that *rbf-1* depletion increases the release of fluorescently tagged neuropeptides without affecting the frequency of spontaneous mini-EPSCs. This suggest a potential negative regulatory role for RPH3A in neuropeptide release (Laurent et al., 2018), opposite to RIM1 (Persoon et al., 2019). However, it remains uncertain

whether RPH3A functions as a negative regulator in mammalian neurons, and whether, similar to
synaptic transmission, the interaction with SNAP25 is important.
We set out to determine the role of RPH3A in DCV exocytosis in hippocampal mouse
neurons, and to investigate the relevance of RAB3A and SNAP25 binding. We found that RPH3A
did not travel with DCVs, but remained stationary at synapses. Using RPH3A null mutant mice
(Schlüter et al., 1999) we show that the absence of RPH3A indeed increased DCV exocytosis.
Furthermore, RPH3A null neurons exhibit longer neurites and an increased number of DCVs. This
effect was partly reduced when all regulated secretion was eliminated by tetanus neurotoxin
(TeNT, Hoogstraaten et al., 2020; Shimojo et al., 2015), indicating that the increased neurite length
partially depends on regulated secretion. Finally, expressing a mutant RPH3A unable to bind
RAB3A/RAB27A restored DCV exocytosis, but not when expressing a mutant RPH3A unable to
bind SNAP25. This suggests that limiting DCV exocytosis does not depend on the interaction with
RAB3A, but at least in part on the interaction with SNAP25.
87

Results

RPH3A is enriched in presynaptic structures

RPH3A is reported to localize to both pre- and post-synaptic sites (Stanic et al., 2015), and on
DCVs through its RAB-binding domain (RBD) in PC12 cells (Fukuda et al., 2004). We first
determined whether RPH3A is expressed in excitatory or inhibitory mouse neurons. To test this,
we immunostained for RPH3A in hippocampal VGLUT+ wildtype (WT) neurons and striatal VGAT+
WT neurons at 14 days *in vitro* (DIV14, Figure S1A). We found no difference in RPH3A expression
between hippocampal and striatal neurons (Figure S1B). To localize RPH3A with higher spatial
precision and confirm localization on DCVs, we performed stimulated emission depletion (STED)
microscopy in hippocampal mouse neurons at DIV14. RPH3A staining was predominantly
punctate and overlapped with the presynaptic marker Synapsin1 (Syn1, Figures 1A, B) and, to a
lesser extent, with the post-synaptic marker Homer (Figures 1A, C). RPH3A partly colocalized with
DCV marker chromogranin B (ChgB, Figures 1A, D). Pearson's correlation coefficients were
similar for RPH3A with either Syn1, Homer and ChgB (Figure 1E), however, Mander's overlap
coefficient analysis (Manders et al., 1992) revealed that RPH3A colocalized significantly more with
Syn1 than Homer or ChgB (Figure 1F). In addition, most RPH3A puncta contained Syn1, but not
all Syn1 puncta contained RPH3A (Figure 1G). These data suggest that RPH3A is a synaptic
protein with predominantly presynaptic accumulation.
We next determined which known interactions are important for RPH3A's synaptic
localization by expressing mutant RPH3A constructs fused to mCherry, lacking specific
interactions or a phosphorylation site, in hippocampal WT neurons at DIV1-2 (Figure S1C). FL
RPH3A-mCherry and all mutant constructs were expressed to a similar level in neurites (Figures
S1G, H). We performed STED microscopy at DIV14 and quantified the colocalization of mCherry
with Syn1 and Homer (Figures S1D, E). Similar to endogenous RPH3A, FL WT RPH3A-mCherry

112
Figure 1. RPH3A localizes to the pre-synapses. (**A**) Representative images of WT hippocampal neurons (top) with
zooms (bottom) co-stained for RPH3A (blue) and Syn1 (magenta), Homer (green) or ChgB (yellow). Scale bar, 5
µm (top) and 2 µm (bottom). (**B, C, D**) Line plots show normalized fluorescent intensity across dotted line in example
zooms. (**E**) Pearson's correlation coefficient and (**F**) Mander's overlap coefficient comparing the colocalization of
RPH3A with Syn1, Homer and ChgB. (**G**) Mander's overlap coefficient comparing the colocalization of RPH3A in
either Syn1, Homer or ChgB puncta, and vice versa. RPH3A:Syn1, RPH3A:Homer and RPH3A:ChgB show the
same dataset as in F. Dots represent a field of view. Boxplots represent the median (line), mean (+) and Tukey

showed a punctate distribution (Figure S1D) that colocalized strongly with Syn1, and to a lesser extent with Homer (Figure S1F). Four mutant versions of RPH3A: a C-terminal truncation, a Ca^{2+}-binding deficient mutant (ΔCa^{2+}-binding) and a mutant that does not bind SNAP25 (ΔSNAP25), all retained a similar punctate presynaptic distribution (Figures S1D, E). However, mutant RPH3A unable to bind RAB3A/RAB27A (ΔRAB3A/RAB27A, Fukuda et al., 2004) and a mutant that lacked a CaMKII dependent phosphorylation site (ΔCAMKII-phos site) showed no punctate distribution (Figures S1D-F). These data suggest that the interaction with RAB3A is required for RPH3A's synaptic localization. CaMKII dependent phosphorylation may also be involved in the synaptic localization of RPH3A.

RPH3A does not travel with DCVs

Recent evidence has demonstrated that RAB3A is transported together with DCVs (Persoon et al., 2019). To test if RPH3A travels with DCVs on both axons and dendrites, we co-expressed neuropeptide Y (NPY)-mCherry and FL WT EGFP-RPH3A (Tsuboi & Fukuda, 2005) in DIV14 RPH3A knockout (KO) neurons. In addition, we tested co-transport of NPY fused to pH-sensitive EGFP (NPY-pHluorin), with two RPH3A mutants: a truncated RPH3A lacking both C2 domains but retaining the RAB3A and RAB27A binding sites, and ΔRAB3A/RAB27A mutant (Figure 2A, Fukuda et al., 2004). Live-cell imaging was performed before and after photobleaching a fixed-size area in neurites, without distinguishing between axons and dendrites, to enhance visualization of moving vesicles entering the bleached area (Figures 2B-D). The number of moving versus immobile puncta was determined for each construct prior to photobleaching. Both FL- and truncated RPH3A exhibited low mobility (7-14% moving), although they occasionally showed small movements over short distances (less than 10 µm within 4 minutes). In contrast, 47% of NPY puncta were mobile (Figure 2E), as shown before (Hoogstraaten et al., 2020; Persoon et al., 2019). Only a small fraction of RPH3A (14%) and truncated RPH3A (7%) puncta travelled at velocities characteristic for DCV transport (Bittins et al., 2010; de Wit et al., 2006; Kwinter et al., 2009). These results indicate that the majority of RPH3A organizes in immobile puncta that colocalize with synaptic markers (see above), but do not travel through the axons or dendrites like DCVs do.

We next determined the fraction of RPH3A that travelled together with NPY-labelled vesicles. Few moving NPY vesicles contained RPH3A or truncated RPH3A (NPY:RPH3A = 6%, NPY:Truncated RPH3A = 1%). Conversely, of the already small fraction of moving FL- and truncated RPH3A, only a few travelled together with NPY (RPH3A:NPY = 18%, truncated RPH3A:NPY = 8%; Figure 2F). Truncated RPH3A showed almost no fluorescence recovery after photobleaching compared to NPY (Figure 2G), however ΔRAB3A/RAB27A RPH3A showed a faster recovery compared to FL- and truncated RPH3A (Figure 2H). Overall, these findings

Figure 2. RPH3A does not travel with DCVs. (**A**) Domain structures of FL RPH3A and mutant RPH3A constructs lacking specific interactions: ΔRAB3A/RAB27A mutant RPH3A and truncated RPH3A that lacked its calcium and SNAP25 binding C2A and C2B domain. (**B**) Kymographs of EGFP-RPH3A and NPY-mCherry, (**C**) mCherry-truncated RPH3A and NPY-pHluorin, and (**D**) mCherry-ΔRAB3A/RAB27A RPH3A and NPY-pHluorin before (upper) and after (lower) photobleaching (black bar) to remove most stationary puncta and facilitate analysis. NPY-

pHluorin showed more resistance for bleaching. Merged images show mCherry (pseudo-coloured magenta) and EGFP/pHluorin (green). Scale bar, 20 µm (x-axis) and 20 s (y-axis). (**E**) Moving fraction of NPY, FL RPH3A and truncated RPH3A puncta per kymograph. (**F**) Fraction of co-travel of NPY puncta with either FL- or truncated RPH3A puncta, and co-travel of FL- or truncated RPH3A with NPY puncta. Dots represent a kymograph. (**G**) Fluorescent recovery of the traces shown in Figure 2C after photobleaching truncated RPH3A or NPY-pHluorin. (**H**) Mean fluorescent recovery traces from multiple kymographs after photobleaching FL-, truncated- or ΔRAB3A/RAB27A RPH3A. Lines ± shading represents mean ± SEM. Boxplots represent median (line), mean (+) and Tukey range (whiskers).

suggest that RPH3A does not travel with DCVs and that the stationary organization of RPH3A relies on RAB3A/RAB27A interactions.

RPH3A deficiency increases DCV exocytosis

Since RPH3A appeared to remain mostly stationary at the pre-synapse (Figures 1, 2), we examined the role of RPH3A in neuropeptide release in hippocampal neurons. We recorded DCV fusion events in single hippocampal mouse neurons from RPH3A KO and WT littermates at DIV14-16. We confirmed the loss of RPH3A expression in RPH3A KO neurons with immunocytochemistry (Figures 3A, B). NPY-pHluorin, a validated DCV fusion reporter (Arora et al., 2017; Farina et al., 2015; Persoon et al., 2018; van de Bospoort et al., 2012), was used to quantify single fusion events (Figure 3C). To elicit DCV fusion, neurons were stimulated twice with 8 bursts of 50 action potentials (AP) at 50 Hz, separated by 30 s (Figure 3D) or once with 16 bursts of 50 APs at 50 Hz. The acidity of DCVs quenches NPY-pHluorin, but upon fusion with the plasma membrane, the DCV deacidifies resulting in increased fluorescent NPY-pHluorin intensity (Figure 3C). After stimulation, neurons were briefly perfused with NH_4^+ to de-quench all NPY-pHluorin labelled DCVs (Figures 3C, D) to determine the number of remaining DCVs per cell. We have previously demonstrated that the fluorescent intensity of NPY-pHluorin in confocal imaging directly correlates with the number of fluorescent puncta for endogenous DCV markers using super-resolution imaging (Persoon et al., 2018).

RPH3A KO neurons showed a 3-fold increase in the total number of fusion events compared to WT (Figures 3F, G, S2A, B). To test if re-expressing RPH3A in KO neurons could restore DCV fusion to WT levels, we infected KO neurons with a FL RPH3A construct at DIV0 (Figure 3E). This construct localized to synapses (Figure S1D), similar to endogenous RPH3A expression (Figures 1A-C). Re-expression of RPH3A in KO neurons restored the number of fusion events to WT levels (Figures 3F, G). The released fraction, i.e. the number of fusion events divided by the remaining DCV pool, did not differ between WT and KO neurons (Figure 3H), and a trend towards a larger remaining DCV pool in these KO neurons was observed (Figure S2C).

We have recently shown that different stimulation protocols influence certain fusion dynamics like event duration, but not the total number of fusion events (Baginska et al., 2023). To test this and the robustness of our findings, we used a prolonged stimulation protocol (16 bursts of 50 APs at 50 Hz). The event duration was similar in WT and KO neurons, for both stimulation

 Figure 3. RPH3A deficiency increases DCV exocytosis. (**A**) Typical example of RPH3A WT and KO neurons

immunostained for MAP2 (red) and RPH3A (white). Scale bar, 50 μm. (**B**) RPH3A expression in dendrites of WT
and KO neurons normalized to WT per independent experiment. N numbers of individual experiments and single
neuron observations in brackets: WT: 4(34); KO: 4(33). (**C**) Schematic representation (left) and imaging example
of a WT neurite stretch (right) infected with NPY-pHluorin as optical DCV fusion reporter. NPY-pHluorin is quenched
in the acidic DCV lumen before fusion (baseline) but dequenches upon fusion (stimulation). $NH4^+$ perfusion
dequenches all NPY-pHluorin labelled DCVs (remaining DCV pool). Scale bar, 5 μm. (**D**) Kymograph of a WT
neurite stretch with the stimulation paradigm used to elicit DCV fusion (two bursts of 8 x 50 AP trains at 50 Hz
interspaced by 0.5 s between each train and 30 s between each burst, blue bars) and NH4 perfusion ($NH4^+$) used
to dequench all NPY-pHluorin labelled vesicles. Arrowheads indicate fusion events. (**E**) Domain structure of FL
RPH3A construct. (**F**) Cumulative median histogram of fusion events over time in WT (black), RPH3A KO (red)
and KO neurons infected with FL RPH3A (cyan). Blue bars indicate the stimulation paradigm (two bursts of 8 x 50
AP bursts at 50 Hz). (**G**) Total number of DCV fusion events per condition (two bursts of 8 x 50 AP bursts at 50
Hz). (**H**) Released fraction defined as the number of fusion events normalized to the remaining pool of DCVs. N
numbers of individual experiments and single neuron observations in brackets: WT: 5(51); KO: 5(43); KO + RPH3A:
5(39). (**I**) Cumulative median histogram of events over time in WT (black), RPH3A KO (red) and KO neurons
infected with FL RPH3A (cyan). Blue bars indicate the stimulation paradigm (16 x 50 AP bursts at 50 Hz). (**J**) Total
number of DCV fusion events per condition (16 x 50 AP bursts at 50 Hz). (**K**) Release fraction per cell. N numbers
of individual experiments and single neuron observations in brackets: WT: 4(25); KO: 4(18); KO + RPH3A: 4(16).
Boxplots represent the median (line), mean (+) and Tukey range (whiskers). Each dot represents an individual
neuron. Kruskal-Wallis H test with Dunn's correction: * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$. ns = non-significant,
$P > 0.05$.

paradigms (Figures S2D-F). During prolonged stimulation, we observed an increase in DCV fusion
events in KO neurons compared to WT (Figures 3I, J), similar to the 2 x 8 stimulation protocol. In
addition, the released fraction was significantly increased in KO neurons compared to WT (Figure
3K). We did not observe a difference in remaining pool size between KO and WT in these neurons
(Figure S2G). Overexpression of FL RPH3A in KO neurons again restored the number and
released fraction of fusion events to WT levels (Figures 3I-K). We did not observe an effect on
spontaneous DCV fusion in RPH3A KO neurons (Figure S2H). Together, these findings indicate
that RPH3A is an inhibitor of DCV exocytosis.

The SNAP25-, but not the RAB3A-interaction domain of RPH3A contributes to limiting DCV
exocytosis
To investigate whether the interactions with RAB3A or SNAP25 are relevant to limit DCV
exocytosis, we overexpressed FL WT RPH3A, ΔRAB3A/RAB27A mutant RPH3A (Fukuda et al.,
2004), and ΔSNAP25 mutant RPH3A (Ferrer-Orta et al., 2017) in RPH3A KO neurons (Figure 4A).
Expression of ΔRAB3A/RAB27A restored DCV fusion and the released fraction to WT levels,
similar to FL RPH3A (Figures 4B, C). However, expression of ΔSNAP25 did not fully rescue the
number of fusion events or the released fraction (Figures 4E, F). We did not observe any
difference in the number of remaining DCVs upon ΔRAB3A/RAB27A or ΔSNAP25 expression
(Figures 4D, G). Taken together, these results suggest that RAB3A/RAB27A binding is not

essential for the limiting effect of RPH3A on DCV exocytosis, but that the interaction with SNAP25

appears to contribute to this effect.

Figure 4. RPH3A interaction with SNAP25, but not RAB3A, partly contributes to limiting DCV exocytosis. This

Figure shows the same dataset for WT, KO and FL RPH3A as in Figure 3. (**A**) Domain structures of FL RPH3A

(cyan), ΔRAB3A/RAB27A- (yellow) and ΔSNAP25 mutant RPH3A (purple) with the corresponding mutant sites in

red. (**B**) Total DCV fusion events in WT (black), KO (red) and KO neurons expressing RPH3A (cyan) or

ΔRAB3A/RAB27A mutant RPH3A (yellow). (**C**) Released fraction of the number of fusion events normalized to the

remaining DCV pool per cell. (**D**) Remaining NPY-pHluorin labelled DCV pool estimates derived from NH_4^+

perfusion after stimulation. N numbers of individual experiments and single neuron observations in brackets: WT:

4(27); KO: 4(28); KO + RPH3A: 4(24); KO + ΔRAB3A/RAB27A: 4(28). (**E**) Total number of DCV fusion events in

WT (black), KO (red) and KO neurons expressing RPH3A (cyan) or ΔSNAP25 mutant RPH3A (purple). (**F**) Release

fraction per cell. (**G**) Remaining NPY-pHluorin labelled DCV pool per cell. N numbers of individual experiments and

single neuron observations in brackets: WT: 4(25); KO: 4(18); KO + RPH3A: 4(16); KO + ΔSNAP25: 4(26). Boxplots

show the mean (+), median (line) and Tukey range (whiskers). Each dot represents a single neuron. Kruskal-Wallis

H test with Dunn's correction: * $P < 0.05$, ** P < 0.01. ns = non-significant, $P > 0.05$.

RPH3A deficiency leads to increased neurite length and DCV numbers

Since we observed a trend towards a bigger DCV pool in KO neurons (Figure S2C), and the total

number of DCVs per neuron correlates with dendrite length (Persoon et al., 2018), we examined

the neuronal morphology of RPH3A KO neurons. Single hippocampal neurons (DIV14) were

immunostained for MAP2 to quantify the total dendritic length, and for endogenous DCV cargo

ChgB to determine the number of DCVs (Figure 5A). We have previously shown that this correlates

well with the number of dSTORM ChgB puncta (Persoon et al., 2018). Indeed, neurons lacking

RPH3A had longer dendrites that harbored more DCVs than WT neurons (Figures 5B, C) and KO

neurons contained more DCVs per μm (Figure 5D). The number of DCVs correlated with the total

dendrite length in both genotypes (Figure 5F), as shown previously for WT neurons (Persoon et

al., 2018). Moreover, the intensity of endogenous ChgB was decreased in RPH3A KO neurons

(Figure 5E), suggesting affected vesicle loading or reduced clustering. However, the peak intensity

of fusion events during live recording was unchanged (Figure S2I), indicating that the decrease in

DCV cargo intensity is potentially due to reduced clustering or accumulation. Taken together, these

results suggest that lack of RPH3A leads to longer dendrites with a concomitant increase in the

number of DCVs. RPH3A depletion does not seem to affect the neuropeptide content in DCVs,

instead it might reduce the clustering of DCVs.

Figure 5. Increased neurite length and DCV number upon RPH3A deficiency. (**A**) Typical example of a single WT and RPH3A KO hippocampal neuron (top) with zooms (bottom) stained for MAP2 (green) and the DCV marker ChgB (magenta). Scale bars, 50 µm (top) and 20 µm (bottom). (**B**) Total dendritic length of single hippocampal RPH3A WT or KO neurons normalized to WT per independent experiment. N numbers per condition: WT: 14(112); KO: 14(113). (**C**) Total number of ChgB labelled DCVs per neuron for each group. N numbers per condition: WT: 3(24); KO: 3(25). (**D**) Total ChgB labelled DCVs per µm for each neuron per group. (**E**) Mean intensity of ChgB labelled DCVs per neuron for each group. (**F**) Correlation between ChgB labelled DCVs and dendritic length (mm). Linear regression goodness of fit (r^2) is given for each group. Boxplots represent median (line), mean (+) and Tukey range (whiskers). Each dot represents an individual neuron. Mann-Whitney U or unpaired t test: ** $P < 0.01$, ***$P < 0.001$, **** $P < 0.0001$.

Increased neurite length upon RPH3A deficiency partly depends on regulated secretion

RPH3A deletion resulted in increased DCV exocytosis (Figure 3) and dendrite length (Figure 5B). We reasoned that increased release of neuropeptide- and neurotrophic factors throughout development could contribute to the longer neurites observed. To test this, we inhibited both SV- and DCV exocytosis by cleaving VAMP1, VAMP2, and VAMP3 using TeNT (Hoogstraaten et al., 2020; Humeau et al., 2000), followed by immunostainings for MAP2 and Tau to assess dendritic and axonal length, respectively. Neurons infected with TeNT at DIV1 lacked VAMP2 staining at DIV14, confirming successful cleavage (Figure 6A). TeNT expression in WT neurons had no effect on dendritic (Figure 6B) or axonal length (Figure 6C), as shown before (Harms & Craig, 2005). TeNT expression in KO neurons restored neurite length to WT levels (Figures 6B, C). When comparing KO neurons with and without TeNT, KO neurons with TeNT show a trend towards decreased neurite length, similar to WT. (Figures 6C, B). Re-expression of RPH3A in KO neurons without TeNT restored neurite length to WT levels (Figures 6B, C). To identify the downstream pathway, we overexpressed mutant RPH3A constructs, lacking specific interactions (Figure S1A). No significant differences in dendrite or axon length were observed for any of the mutants compared to WT (Figures S3A-C). These results indicate that regulated secretion is not required for neurite outgrowth, but that the increased neurite length upon RPH3A depletion depends, at least in part, on regulated secretion.

Figure 6. Increased neurite length upon RPH3A deficiency partly depends on regulated secretion. (**A**) Typical example of a single RPH3A KO neurons either infected with TeNT or not, showing successful VAMP2 cleavage with zooms (bottom) stained for MAP2 (green), Tau (blue) and VAMP2 (magenta). Scale bars, 50 μm (top) and 20 μm (bottom). (**B**) Total dendritic and (**C**) axonal length (mm) of WT and KO neurons - / + TeNT, and KO neurons expressing RPH3A. N numbers per condition: WT: 3(19); WT + TeNT: 3(27); KO: 3(28). KO + TeNT: 3(26), KO + RPH3A: 3(28). Boxplots represent median (line), mean (+) and Tukey range (whiskers). Each dot represents an individual neuron. Kruskal-Wallis H test with Dunn's correction: * $P < 0.05$, ** $P < 0.01$, ***$P < 0.001$, **** $P < 0.0001$. ns = non-significant, $P > 0.05$.

Discussion

In this study, we addressed the role of RPH3A in DCV exocytosis in hippocampal neurons. RPH3A predominantly localized to the pre-synapse (Figure 1) and did not travel with DCVs (Figure 2). RPH3A null mutant neurons showed an increase in DCV exocytosis (Figure 3). Expression of a RAB3A/RAB27A-binding deficient, but not a SNAP25-binding deficient RPH3A, in RPH3A KO neurons, restored DCV exocytosis to WT levels. (Figure 4). Finally, RPH3A null neurons had longer

neurites that contained more DCVs (Figure 5). The increase in neurite length may partially depend

on regulated secretion, as TeNT expression showed a strong trend towards reduced neurite length

to WT levels (Figure 6). Taken together, we conclude that RPH3A limits DCV exocytosis, partly

through its interaction with SNAP25.

Presynaptic enrichment and accumulation of RPH3A

We found that RPH3A is enriched at the pre-synapse. This is in line with previous studies showing

synaptic enrichment of RPH3A (Li et al., 1994; Mizoguchi et al., 1994; Stanic et al., 2015, 2017).

RPH3A showed mostly a punctate distribution, except in the pre-synapse, where it tends to

accumulate (Figure 1A). Previous studies showed that inactivation of *synapsin1* and *2* genes

decreased RPH3A levels but increased phosphorylation of the remaining RPH3A (Lonart &

Simsek-Duran, 2006). Hence, synapsins may, directly or indirectly, contribute to presynaptic

RPH3A accumulation, for instance as part of the phase separation of the vesicle cluster. In

addition, phosphorylation reduces RPH3A's affinity for membranes (Foletti et al., 1994) and

therefore preferential presynaptic dephosphorylation may also help to explain the presynaptic

RPH3A accumulation.

RPH3A does not travel with DCVs in hippocampal neurons

We demonstrated that FL and truncated RPH3A are highly stationary at synapses. However, a

RPH3A mutant unable to bind RAB3A/RAB27A was more mobile (faster recovery after

photobleaching, Figure 2), and lost its punctate distribution. This indicates that the synaptic

enrichment of RPH3A depends, at least in part, on RAB3A/RAB27A interactions. This is in line

with previous findings showing the expression levels and localization of RPH3A depend on RAB3

in mammalian neurons (Schlüter et al., 1999). These data do not exclude that RPH3A interacts

with immobile DCVs at synapses, potentially in a docked or primed state at the release sites,

consistent with our conclusion that RPH3A serves as a negative regulator of DCV exocytosis.

Based on these findings, we propose that RAB3A recruits RPH3A to DCV release sites, where it

interacts with (immobile) DCVs and competes with essential DCV proteins, such as

synaptobrevin/VAMP2 and synaptotagmin-1/synaptotagmin-7 (Hoogstraaten et al., 2020; van

Westen et al., 2021), for SNAP25 binding (Figure 7).

RPH3A limits DCV exocytosis in hippocampal neurons

DCV exocytosis was increased by 3-fold upon RPH3A depletion and this effect was rescued by

overexpressing FL RPH3A (Figures 3F, I). The increase in released fraction upon RPH3A

depletion was most significant during intense, prolonged stimulation (16 bursts of 50 action

potentials, Figure 3K). During more distributed stimulation, a similar trend was observed (Figure

3F). Together, these results suggest that RPH3A is a negative regulator of DCV exocytosis

(Figures 7A, B). This is in line with *C. elegans* data, showing that the lack of the nematode homolog

rbf-1 increased DCV exocytosis, but had no effect on spontaneous release of SVs. This suggests
that *rbf-1* limits DCV exocytosis only (Laurent et al., 2018). In addition, expression of FL RPH3A
in PC12 cells reduced high KCl-dependent neuropeptide release (Fukuda et al., 2004). In this
study, we demonstrated that RPH3A also negatively regulated neuropeptide release in mammalian
hippocampal neurons. Our prior estimations in mouse hippocampal neurons indicated that merely
1-6% of the total DCV pool undergo exocytosis upon strong stimulation (Persoon et al., 2018),
implying the existence of an inhibitory release mechanism. To our knowledge, RPH3A is the only
negative regulator of DCV exocytosis in mammalian neurons identified so far.
368
The limiting effect of RPH3A on DCV exocytosis partially depends on SNAP25 binding
We recently discovered that RAB3 and its effector RIM1 are positive regulators of DCV exocytosis
in mammalian hippocampal neurons (Persoon et al., 2019). In contrast, we demonstrated that
RPH3A serves as a negative regulator of DCV exocytosis. Given that RPH3A is a downstream
effector of RAB3, and that RAB3 is necessary for synaptic RPH3A enrichment, we expected that
the interaction with RAB3 plays a role in limiting DCV exocytosis. However, enhanced DCV
exocytosis was rescued upon expression of a RPH3A mutant that was unable to bind
RAB3A/RAB27A, suggesting that the limiting effect of RPH3A on DCV exocytosis is independent
of an interaction with RAB3A (Figure 7C). This indicates that even though RAB3A is important for
the localization of RPH3A, RPH3A can still limit exocytosis when its punctate organization is lost,
suggesting that the interaction with RAB3A is not rate limiting. One other potential mechanism by
which RPH3A could directly limit exocytosis is by limiting SNAP25 availability. Previous research
has demonstrated that RPH3A binding to SNAP25 negatively regulates SV recycling (Deák et al.,
2006). We found that expressing mutant RPH3A that lacks SNAP25 binding in KO neurons does
not fully restore DCV exocytosis to WT levels. This suggests that RPH3A limits DCV exocytosis
by interacting with SNAP25 (Figure 7D). The partial selectivity for the DCV secretory pathway may
be attributed to RPH3A functioning as a downstream effector of RAB3A. RAB3 is essential for
DCV exocytosis (Persoon et al., 2019) but largely redundant for synaptic transmission (Schlüter et
al., 2004, 2006). Based on our findings, we propose that RAB3A plays a role in recruiting RPH3A
to synaptic exocytosis sites, where RPH3A might bind available SNAP25, potentially restricting the
assembly of SNARE complexes and thereby inhibiting DCV exocytosis (Figure 7).
390
Increased regulated secretion in RPH3A KO neurons might lead to longer neurites
RPH3A KO neurons have longer neurites that correlated with the number of DCVs as shown before
(Persoon et al., 2018). In agreement with previous findings (Ahnert-Hilger et al., 1996; Grosse et
al., 1999; Osen-Sand et al., 1996), we find that TeNT expression did not affect neurite length of
WT neurons, but showed a trend towards shorter neurites in RPH3A KO neurons. Based on this,
we speculate that the increased neurite length in RPH3A KO neurons might, at least partially, be
driven by TeNT-dependent regulated secretion, in particular VAMP1, 2 or 3 mediated exocytosis.

Given that neuropeptides and neurotrophic factors are key modulators of neuronal maturation and
outgrowth (Mu et al., 2010), and that RPH3A depletion leads to increased DCV exocytosis, it
stands to reason that we observed longer neurites in RPH3A KO neurons.

The partial rescue by TeNT suggests RPH3A-dependent mechanisms that could explain
the increase in neuron size besides regulated secretion. RPH3A could control neurite length by
regulating the actin cytoskeleton. RPH3A interacts with actin via binding to α-actinin and ß-adducin
(Baldini et al., 2005; Coppola et al., 2001; Kato, Sasaki, et al., 1996). *In vitro* experiments showed
that RPH3A together with α-actinin can bundle actin (Kato, Sasaki, et al., 1996). Regulation of the
actin cytoskeleton has extensively been linked to neurite outgrowth (Meberg & Bamburg, 2000)
making it plausible that lack of RPH3A alters actin regulation, resulting in longer neurites.

Figure 7. Role of RPH3A in DCV exocytosis. (**A**) RPH3A binding to RAB3 through its RAB-binding domain (RBD)
ensures confined presynaptic localization of RPH3A. RPH3A binding to SNAP25 through its C2B domain inhibits
DCV exocytosis, potentially by inhibiting SNAP25-binding of essential DCV proteins synaptobrevin/VAMP2 and/or
synaptotagmin (not depicted). (**B)** In the absence of RPH3A, DCV exocytosis is not limited. (**C**) Upon expression
of a RPH3A mutant that is unable to bind RAB3A/RAB27A, DCV exocytosis is limited. (**D**) When RPH3A is unable
to bind SNAP25, the SNARE assembly is not restricted and therefore DCV exocytosis is not limited, while RPH3A
is still recruited to synapses/release sites via RAB3.

415
416
417
418

Material and methods

Animals

Animals were housed and bred in accordance with Dutch and institutional guidelines. All animal experiments were approved by the animal ethical committee of the VU University / VU University Medical Centre (license number: FGA 11-03 and AVD112002017824). All animals were kept on a C57Bl/6 background. For RPH3A KO (Schlüter et al., 1999) and WT primary hippocampal neuron cultures, RPH3A heterozygous mice mating was timed and P1 pups were used to dissect hippocampi. Pups were genotyped prior to dissection to select RPH3A KO and WT littermates.

Neuron culture

Primary hippocampal neurons were cultured as described before (De Wit et al., 2009; Farina et al., 2015). In short, dissected hippocampi were digested with 0.25% trypsin (Gibco) in Hanks' balanced salt solution (Sigma) with 10 mM HEPES (Life Technologies) for 20 minutes at 37 °C. Hippocampi were washed, triturated, and counted prior to plating. For single hippocampal neurons, 1000-2000 neurons per well were plated on pre-grown micro-islands generated by plating 6000 rat glia on 18 mm glass coverslips coated with agarose and stamped with a solution of 0.1 mg/ml poly-D-lysine (Sigma) and 0.7 mg/ml rat tail collagen (BD Biosciences, Mennerick et al., 1995; Wierda et al., 2007). For continental hippocampal cultures, 20,000 neurons per well were plated on pre-grown glial plates containing 18 mm glass coverslips. All neurons were kept in neurobasal supplemented with 2% B-27, 18 mM HEPES, 0.25% Glutamax and 0.1% Pen-Strep (all Gibco) at 37 °C and 5% CO_2.

Lentiviral vectors and infections

All constructs were cloned into lentiviral vectors containing a synapsin promoter to restrict expression to neurons. Lentiviral particles were produced as described before (Naldini et al., 1996). NPY-mCherry, NPY-pHluorin and TeNT-IRES-mCherry were described before (Emperador Melero et al., 2017; Nagai et al., 2002). For re-expression of RPH3A in single RPH3A KO neurons, FL RPH3A was cloned into a lentiviral vector containing an IRES-NLS-mCherry to verify infection during live cell experiments. For DCV fusion experiments at DIV14-DIV16, neurons were infected with RPH3A-IRES-NLS-mCherry, RAB3A and RAB27A binding deficient RPH3A with two-point mutations (E50A and I54A, Fukuda et al., 2004) and SNAP25 binding deficient RPH3A (K648A, K653A and K660A, Ferrer-Orta et al., 2017) at DIV1-2. The E51A/I54A double mutant of RPH3A was previously validated to lose RAB3A and RAB27A binding activity (Fukuda et al., 2004), and the K651A/K656A/K663A mutant was shown to not bind rat WT SNAP25 (Ferrer-Orta et al., 2017). We determined the corresponding residues to be mutated in a mouse SNAP25-binding deficient RPH3A construct (K648,653,660A). Neurons were infected with NPY-pHluorin at DIV8-9. For neurite length and co-travel experiments, FL RPH3A, truncated RPH3A (1-375), RAB3A/RAB27A binding deficient RPH3A (Fukuda et al., 2004), RPH3A with two point-mutations in the C2B domain

457 (D568N, D574N) and RPH3A phosphorylation deficient (S217A, Foletti et al., 2001; Fykse et al.,
458 1995) were N-terminally tagged with EGFP or mCherry interspaced by a short glycine linker
459 sequence (Tsuboi & Fukuda, 2005). For neurite length experiments at DIV14, all constructs were
460 infected at DIV1-2. For co-travel experiments at DIV10 constructs were infected at DIV4.
461

Immunocytochemistry
Cells were fixed with 3.7% paraformaldehyde (PFA, Merck) in phosphate-buffered saline (PBS;
137 mM NaCl, 2.7 mM KCl, 10 mM Na_2HPO_4, 1.8 mM KH_2PO_4, pH 7.4) for 12 minutes at room
temperature (RT). Cells were immediately immunostained or kept in PBS at 4 °C. For ChgB
immunostainings, cells were permeabilized in 0.5% Triton X-100 (Fisher Chemical) for 10 minutes
and blocked with 5% BSA (Acro Organic) in PBS for 30 minutes at RT. For all other
immunostainings, cells were permeabilized with 0.5% Triton X-100 for 5 minutes and blocked with
2% normal goat serum (Fisher Emergo) in 0.1% Triton X-100 at RT. Cells were incubated with
primary antibodies overnight at 4°C. Primary antibodies used were: polyclonal ChgB (1:500,
SySy), monoclonal RPH3A (1:1000, Transduction Labs), polyclonal MAP2 (1:500, Abcam),
polyclonal Syn1 (1:1000, #P610; 1:500, SySy), VGLUT1 (1:500, Millipore), polyclonal Homer 1
(1:500, SySy), polyclonal Tau (1:1000, SySy), monoclonal VAMP2 (1:1000, SySy) and monoclonal
mCherry (1:1000, Signalway Antibody). Alexa Fluor conjugated secondary antibodies (1:1000,
Invitrogen) were incubated for 1 hour at RT. Abberior secondary antibodies (1:50, Abberior) for
STED imaging were incubated for 2 hours at RT. Coverslips were mounted on Mowiol (Sigma-
Aldrich).
478

STED imaging
STED images were acquired with a Leica SP8 STED 3x microscope with an oil immersion 100x
1.44 numerical aperture objective. Alexa Fluor 488, Abberior STAR 580 and Abberior STAR 635p
were excited with 592 nm and 775 nm using a white light laser. The Abberior STAR 635p was
acquired first in STED mode using 4x line accumulation, followed by Abberior STAR 580, both
were depleted with 775 nm depletion laser (50% of max power). Alexa Fluor 488 was acquired in
STED mode and depleted with 592 nm depletion laser (50% of max power). The Alexa Fluor 405
channel was acquired in confocal mode. The signals were detected using a gated hybrid detector
(HyD) in photon-counting mode (Leica Microsystems). Z-stacks were made with a step size of 150
nm and pixel size of 22.73 × 22.73 nm. Finally, deconvolution was performed with Huygens
Professional software (SVI).
490

491 **Live and fixed imaging**
492 For live cell experiments, neurons were continuously perfused in Tyrode's solution (2 mM CaCl2,
493 2.5 mM KCl, 119 mM NaCl, 2 mM MgCl2, 30 mM glucose, 25 mM HEPES; pH 7.4) at RT. For DCV
494 fusion experiments, imaging was performed at DIV14-16 with a Zeiss AxioObserver.Z1 equipped

with 561 nm and 488 nm lasers, a polychrome V, appropriate filter sets, a 40x oil objective (NA 1.3) and an EMCCD camera (C9100-02; Hamamatsu, pixel size 200 nm). Images were acquired at 2 Hz with AxioVision software (version 4.8, Zeiss). Electrical stimulation was performed with two parallel platinum electrodes placed around the neuron. After 30 seconds of baseline, 2 x 8 (separated by 30 seconds) or 1 x 16 train(s) of 50 APs at 50 Hz (interspaced by 0.5 s) were initiated by a Master-8 (AMPI) and a stimulus generator (A-385, World Precision Instruments) delivering 1 ms pulses of 30 mA. NPY-pHluorin was dequenched 50 or 80 seconds after the last stimulation train by superfusing Tyrode's with 50 mM NH4Cl (replacing 50 mM NaCl). For a detailed protocol of DCV fusion analyses see (Moro et al., 2021).

For co-travel experiments, neurons were infected with NPY-mCherry or NPY-pHluorin and FL- or mutant RPH3A fused to EGFP or mCherry. Imaging was performed at DIV9-10 on a Nikon Ti-E eclipse microscope with a LU4A laser system, appropriate filter sets, 60x oil objective (NA = 1.4) and EMCCD camera (Andor DU-897). Sequential images for both 488 and 561 color channels were acquired for 2 minutes at 2 Hz. After 2 minutes, both axonal and dendritic stretches were photobleached to facilitate analysis. A galvano laser scanner was used to scan a selected area with both lasers at 100% (26.9 μs pixel dwell time). Bleaching was followed by an 8-minute acquisition at 2 Hz in both channels. To visualize NPY-pHluorin, neurons were continuously perfused in Tyrode's containing 25 mM NH$_4$Cl (replacing 25 mM NaCl). NPY-pHluorin showed more resistance for bleaching.

For fixed experiments, confocal images were acquired using a Zeiss LSM 510 confocal laser-scanning microscope (40x objective; NA 1.3) and LSM510 software (version 3.2 Zeiss) or a A1R Nikon confocal microscope with LU4A laser unit (40x objective; NA 1.3) and NIS elements software (version 4.60, Nikon). To determine both dendrite (MAP2) and axon (Tau) length, the whole glial island was visualized by stitching 4 images (604.7 x 604.7 microns).

Analyses

For DCV fusion experiments, ImageJ was used to manually place 3x3 pixel ROIs around each NPY-pHluorin fusion event on both axons and dendrites, excluding the soma. An NPY-pHluorin event was considered a DCV fusion event if it suddenly appeared and if the maximal fluorescence was at least twice the SD above noise. Custom written MATLAB (MathWorks, Inc.) scripts were used to calculate the number and timing of fusion events. The remaining DCV number per neuron was determined as the number of fluorescent puncta during NH$_4^+$ perfusion, corrected for overlapping puncta. The released fraction was calculated by dividing the total number of fusion events by the remaining pool size. For a detailed protocol of DCV fusion analyses see (Moro et al., 2021).

For DCV co-travel experiments, segmented lines were drawn along the neurites and kymographs were created using the KymoResliceWide plugin in ImageJ. Puncta were considered moving if the minimal displacement during the whole 2- or 8-minute acquisition was at least 3/4

microns within a 10 second period. For colocalization experiments, the Pearson's and Manders' correlation coefficients were determined using the JACoP plugin (Bolte & Cordelières, 2006).

For fixed experiments, the number of endogenous ChgB+ puncta were counted using SynD (Schmitz et al., 2011; van de Bospoort et al., 2012) software (version 491). All puncta were divided by the mode of the first quartile of puncta intensity values (an estimate for a single DCV) to estimate the total number of DCVs per neuron. Based on this, the total number of DCVs per neuron was quantified. To determine dendritic and axonal length, a mask was created using MAP2 and Tau immunostaining, respectively. Single-pixel images were obtained from the masks. The distances between the neighboring pixels within the dendrite mask are summed together to obtain the dendrite or axon length (Schmitz et al., 2011).

Statistical analysis

Statistical tests were performed using R or GraphPad Prism. Normal distributions of all data were assessed with Shapiro-Wilk normality tests and Levene's test of homogeneity of variances. Multi-level models were used to account for variance within the animals when variance between animals significantly improved the model (Aarts et al., 2014). To compare two groups, unpaired Student's t-test in the case of normal distributed data or Mann-Whitney U tests for non-parametric data were used. For multiple comparisons the Kruskal-Wallis test was used for non-parametric data followed by Dunn's multiple comparisons test to compare conditions. Data is represented as boxplots (25-75% interquartile range) with the median (line), mean (+) and Tukey whiskers or bar graphs with SEM. N numbers represent number of independent experiments, with the number of individual neurons in brackets. Dots in all graphs indicate single neuron observations, unless stated otherwise.

Acknowledgements

This work is supported by an ERC Advanced Grant (322966) of the European Union (to M.V.) We thank Joke Wortel for animal breeding, Robbert Zalm for cloning and producing viral particles, Desiree Schut and Lisa Laan for astrocyte culture and cell culture assistance, and Ingrid Saarloos for assistance in protein chemistry.

Author contributions

Adlin Abramian*: Validation; formal analysis; investigation; data curation; visualization; writing original draft; writing review and editing. **Rein Hoogstraaten*:** Conceptualization; validation; formal analysis; investigation; data curation; visualization; writing original draft; writing review. **Fiona H. Murphy:** Formal analysis; investigation; writing review. **Kathryn F. McDaniel:** Investigation; formal analysis. **Ruud F. Toonen:** Supervision; conceptualization; writing review. **Matthijs Verhage:** Supervision; conceptualization; writing review; funding acquisition.

*These authors contributed equally to this work.

Data availability statement

All data and constructs used will be made available upon reasonable request.

References

Aarts, E., Verhage, M., Veenvliet, J. V, Dolan, C. V, & Sluis, S. van der. (2014). A solution to dependency: using multilevel analysis to accommodate nested data. *Nature Neuroscience 2014 17:4*, *17*(4), 491–496. https://doi.org/10.1038/nn.3648

Ahnert-Hilger, G., Kutay, U., Chahoud, I., Rapoport, T., & Wiedenmann, B. (1996). Synaptobrevin is essential for secretion but not for the development of synaptic processes. *European Journal of Cell Biology*, *70*(1), 1–11.

Arora, S., Saarloos, I., Kooistra, R., van de Bospoort, R., Verhage, M., & Toonen, R. F. (2017). SNAP-25 gene family members differentially support secretory vesicle fusion. *Journal of Cell Science*, *130*(11), 1877–1889. https://doi.org/10.1242/jcs.201889

Baginska, U., Moro, A., Toonen, R. F., & Verhage, M. (2023). Maximal Fusion Capacity and Efficient Replenishment of the Dense Core Vesicle Pool in Hippocampal Neurons. *The Journal of Neuroscience : The Official Journal of the Society for Neuroscience*, *43*(45), 7616–7625. https://doi.org/10.1523/JNEUROSCI.2251-22.2023

Baldarelli, R. M., Smith, C. M., Finger, J. H., Hayamizu, T. F., McCright, I. J., Xu, J., Shaw, D. R., Beal, J. S., Blodgett, O., Campbell, J., Corbani, L. E., Frost, P. J., Giannatto, S. C., Miers, D. B., Kadin, J. A., Richardson, J. E., & Ringwald, M. (2021). The mouse Gene Expression Database (GXD): 2021 update. *Nucleic Acids Research*, *49*(D1), D924–D931. https://doi.org/10.1093/nar/gkaa914

Baldini, G., Martelli, A. M., Tabellini, G., Horn, C., Machaca, K., Narducci, P., & Baldini, G. (2005). Rabphilin localizes with the cell actin cytoskeleton and stimulates association of granules with F-actin cross-linked by α-actinin. *Journal of Biological Chemistry*, *280*(41), 34974–34984. https://doi.org/10.1074/jbc.M502695200

Balkowiec, A., & Katz, D. M. (2002). Cellular mechanisms regulating activity-dependent release of native brain-derived neurotrophic factor from hippocampal neurons. *Journal of Neuroscience*, *22*(23), 10399–10407. https://doi.org/10.1523/jneurosci.22-23-10399.2002

Bittins, C. M., Eichler, T. W., Hammer, J. A., & Gerdes, H. H. (2010). Dominant-negative myosin Va impairs retrograde but not anterograde axonal transport of large dense core vesicles. *Cellular and Molecular Neurobiology*, *30*(3), 369–379. https://doi.org/10.1007/s10571-009-9459-2

Blake, J. A., Baldarelli, R., Kadin, J. A., Richardson, J. E., Smith, C. L., & Bult, C. J. (2021). Mouse Genome Database (MGD): Knowledgebase for mouse-human comparative biology. *Nucleic Acids Research*, *49*(D1), D981–D987. https://doi.org/10.1093/nar/gkaa1083

Bolte, S., & Cordelières, F. P. (2006). A guided tour into subcellular colocalization analysis in
light microscopy. In *Journal of Microscopy* (Vol. 224, Issue 3, pp. 213–232). Blackwell
Publishing Ltd. https://doi.org/10.1111/j.1365-2818.2006.01706.x
Coppola, T., Hirling, H., Perret-Menoud, V., Gattesco, S., Catsicas, S., Joberty, G., Macara,
L. G., & Regazzi, R. (2001). Rabphilin dissociated from Rab3 promotes endocytosis
through interaction with Rabaptin-5. *Journal of Cell Science*, *114*(9), 1757–1764.
de Wit, J., Toonen, R. F., Verhaagen, J., & Verhage, M. (2006). Vesicular trafficking of
semaphorin 3A is activity-dependent and differs between axons and dendrites. *Traffic*,
7(8), 1060–1077. https://doi.org/10.1111/j.1600-0854.2006.00442.x
de Wit, J., Toonen, R. F., & Verhage, M. (2009). Matrix-dependent local retention of secretory
vesicle cargo in cortical neurons. *Journal of Neuroscience*, *29*(1), 23–37.
https://doi.org/10.1523/JNEUROSCI.3931-08.2009
Deák, F., Shin, O. H., Tang, J., Hanson, P., Ubach, J., Jahn, R., Rizo, J., Kavalali, E. T., &
Südhof, T. C. (2006). Rabphilin regulates SNARE-dependent re-priming of synaptic
vesicles for fusion. *EMBO Journal*, *25*(12), 2856–2866.
https://doi.org/10.1038/sj.emboj.7601165
Emperador Melero, J., Nadadhur, A. G., Schut, D., Weering, J. V., Heine, V. M., Toonen, R.
F., & Verhage, M. (2017). Differential Maturation of the Two Regulated Secretory
Pathways in Human iPSC-Derived Neurons. *Stem Cell Reports*, *8*(3), 659–672.
https://doi.org/10.1016/j.stemcr.2017.01.019
Farina, M., van de Bospoort, R., He, E., Persoon, C. M., van Weering, J. R. T., Broeke, J. H.,
Verhage, M., & Toonen, R. F. (2015). CAPS-1 promotes fusion competence of
stationary dense-core vesicles in presynaptic terminals of mammalian neurons. *ELife*,
2015(4). https://doi.org/10.7554/eLife.05438
Ferrer-Orta, C., Pérez-Sánchez, M. D., Coronado-Parra, T., Silva, C., López-Martínez, D.,
Baltanás-Copado, J., Gómez-Fernández, J. C., Corbalán-García, S., & Verdaguer, N.
(2017). Structural characterization of the Rabphilin- 3A-SNAP25 interaction.
Proceedings of the National Academy of Sciences of the United States of America,
114(27), E5343–E5351. https://doi.org/10.1073/pnas.1702542114
Foletti, D. L., Blitzer, J. T., & Scheller, R. H. (2001). Physiological modulation of rabphilin
phosphorylation. *The Journal of Neuroscience : The Official Journal of the Society for
Neuroscience*, *21*(15), 5473–5483. https://doi.org/10.1523/JNEUROSCI.21-15-
05473.2001
Fukuda, M., Kanno, E., & Yamamoto, A. (2004). Rabphilin and Noc2 Are Recruited to Dense-
core Vesicles through Specific Interaction with Rab27A in PC12 Cells. *Journal of
Biological Chemistry*, *279*(13), 13065–13075. https://doi.org/10.1074/jbc.M306812200

Fykse, E., Li, C., & Sudhof, T. (1995). Phosphorylation of rabphilin-3A by Ca2+/calmodulin-
and cAMP-dependent protein kinases in vitro. *Journal of Neuroscience*, *15*(3), 2385–
2395. https://doi.org/10.1523/JNEUROSCI.15-03-02385.1995

Grosse, G., Grosse, J., Tapp, R., Kuchinke, J., Gorsleben, M., Fetter, I., Höhne-Zell, B.,
Gratzl, M., & Bergmann, M. (1999). SNAP-25 Requirement for Dendritic Growth of
Hippocampal Neurons. *J. Neurosci. Res*, *56*, 539–546.
https://doi.org/10.1002/(SICI)1097-4547(19990601)56:5

Guillén, J., Ferrer-Orta, C., Buxaderas, M., Pérez-Sánchez, D., Guerrero-Valero, M., Luengo-
Gil, G., Pous, J., Guerra, P., Gómez-Fernández, J. C., Verdaguer, N., & Corbalán-
García, S. (2013). Structural insights into the Ca2+ and PI(4,5)P2 binding modes of the
C2 domains of rabphilin 3A and synaptotagmin 1. *Proceedings of the National Academy*
of Sciences of the United States of America, *110*(51), 20503–20508.
https://doi.org/10.1073/pnas.1316179110

Harms, K. J., & Craig, A. M. (2005). Synapse composition and organization following chronic
activity blockade in cultured hippocampal neurons. *Journal of Comparative Neurology*,
490(1), 72–84. https://doi.org/10.1002/CNE.20635

Hartmann, M., Heumann, R., & Lessmann, V. (2001). Synaptic secretion of BDNF after high-
frequency stimulation of glutamatergic synapses. *EMBO Journal*, *20*(21), 5887–5897.
https://doi.org/10.1093/emboj/20.21.5887

Hoogstraaten, R. I., van Keimpema, L., Toonen, R. F., & Verhage, M. (2020). Tetanus
insensitive VAMP2 differentially restores synaptic and dense core vesicle fusion in
tetanus neurotoxin treated neurons. *Scientific Reports*, *10*(1), 10913.
https://doi.org/10.1038/s41598-020-67988-2

Humeau, Y., Doussau, F., Grant, N. J., & Poulain, B. (2000). How botulinum and tetanus
neurotoxins block neurotransmitter release. *Biochimie*, *82*(5), 427–446.
https://doi.org/10.1016/S0300-9084(00)00216-9

Kato, M., Sasakii, T., Ohyai, T., Nakanishi, H., Nishioka, H., Imamura, M., & Takai, Y. (1996).
Physical and functional interaction of rabphilin-3A with α-actinin. *Journal of Biological*
Chemistry, *271*(50), 31775–31778. https://doi.org/10.1074/jbc.271.50.31775

Krupke, D. M., Begley, D. A., Sundberg, J. P., Richardson, J. E., Neuhauser, S. B., & Bult, C.
J. (2017). The mouse tumor biology database: A comprehensive resource for mouse
models of human cancer. *Cancer Research*, *77*(21), e67–e70.
https://doi.org/10.1158/0008-5472.CAN-17-0584

Kwinter, D. M., Lo, K., Mafi, P., & Silverman, M. A. (2009). Dynactin regulates bidirectional
transport of dense-core vesicles in the axon and dendrites of cultured hippocampal
neurons. *Neuroscience*, *162*(4), 1001–1010.
https://doi.org/10.1016/j.neuroscience.2009.05.038

Laurent, P., Ch'ng, Q. L., Jospin, M., Chen, C., Lorenzo, R., & De Bono, M. (2018). Genetic dissection of neuropeptide cell biology at high and low activity in a defined sensory neuron. *Proceedings of the National Academy of Sciences of the United States of America*, *115*(29), E6890–E6899. https://doi.org/10.1073/pnas.1714610115

Li, C., Takei, K., Geppert, M., Daniell, L., Stenius, K., Chapman, E. R., Jahn, R., De Camilli, P., & Südhof, T. C. (1994). Synaptic targeting of rabphilin-3A, a synaptic vesicle Ca2+/phospholipid-binding protein, depends on rab3A/3C. *Neuron*, *13*(4), 885–898. https://doi.org/10.1016/0896-6273(94)90254-2

Lonart, G., & Simsek-Duran, F. (2006). Deletion of synapsins I and II genes alters the size of vesicular pools and rabphilin phosphorylation. *Brain Research*, *1107*(1), 42–51. https://doi.org/10.1016/j.brainres.2006.05.092

Malva, J. O., Xapelli, S., Baptista, S., Valero, J., Agasse, F., Ferreira, R., & Silva, A. P. (2012). Multifaces of neuropeptide Y in the brain - Neuroprotection, neurogenesis and neuroinflammation. In *Neuropeptides* (Vol. 46, Issue 6, pp. 299–308). Churchill Livingstone. https://doi.org/10.1016/j.npep.2012.09.001

Manders, E. M., Stap, J., Brakenhoff, G. J., Van Driel, R., & Aten, J. A. (1992). Dynamics of three-dimensional replication patterns during the S-phase, analysed by double labelling of DNA and confocal microscopy. *Journal of Cell Science*, *103*(3), 857–862. https://doi.org/10.1242/jcs.103.3.857

McMahon, H. T., Ushkaryov, Y. A., Edelmann, L., Link, E., Binz, T., Niemann, H., Jahn, R., & Südhof, T. C. (1993). Cellubrevin is a ubiquitous tetanus-toxin substrate homologous to a putative synaptic vesicle fusion protein. *Nature*, *364*(6435), 346–349. https://doi.org/10.1038/364346a0

Meberg, P. J., & Bamburg, J. R. (2000). Increase in Neurite Outgrowth Mediated by Overexpression of Actin Depolymerizing Factor. *Journal of Neuroscience*, *20*(7), 2459–2469. https://doi.org/10.1523/JNEUROSCI.20-07-02459.2000

Mennerick, S., Que, J., Benz, A., & Zorumski, C. F. (1995). Passive and synaptic properties of hippocampal neurons grown in microcultures and in mass cultures. *Journal of Neurophysiology*, *73*(1), 320–332. https://doi.org/10.1152/jn.1995.73.1.320

Mizoguchi, A., Yano, Y., Hamaguchi, H., Yanagida, H., Ide, C., Zahraoui, A., Shirataki, H., Sasaki, T., & Takai, Y. (1994). Localization of Rabphilin-3A on the Synaptic Vesicle. *Biochemical and Biophysical Research Communications*, *202*(3), 1235–1243. https://doi.org/10.1006/BBRC.1994.2063

Moro, A., Hoogstraaten, R. I., Persoon, C. M., Verhage, M., & Toonen, R. F. (2021). Quantitative analysis of dense-core vesicle fusion in rodent CNS neurons. *STAR Protocols*, *2*(1), 100325. https://doi.org/10.1016/J.XPRO.2021.100325

Mu, Y., Lee, S. W., & Gage, F. H. (2010). Signaling in adult neurogenesis. *Current Opinion in Neurobiology*, *20*(4), 416–423. https://doi.org/10.1016/j.conb.2010.04.010

Nagai, T., Ibata, K., Park, E. S., Kubota, M., Mikoshiba, K., & Miyawaki, A. (2002). A variant of yellow fluorescent protein with fast and efficient maturation for cell-biological applications. *Nature Biotechnology*, *20*(1), 87–90. https://doi.org/10.1038/nbt0102-87

Naldini, L., Blömer, U., Gallay, P., Ory, D., Mulligan, R., Gage, F. H., Verma, I. M., & Trono, D. (1996). In vivo gene delivery and stable transduction of nondividing cells by a lentiviral vector. *Science*, *272*(5259), 263–267. https://doi.org/10.1126/science.272.5259.263

Osen-Sand, A., Staple, J. K., Naldi, E., Schiavo, G., Rossetto, O., Petitpierre, S., Malgaroli, A., Montecucco, C., & Catsicas, S. (1996). Common and distinct fusion proteins in axonal growth and transmitter release. *Journal of Comparative Neurology*, *367*(2), 222–234. https://doi.org/10.1002/(SICI)1096-9861(19960401)367:2<222::AID-CNE5>3.0.CO;2-7

Pang, P. T., Teng, H. K., Zaitsev, E., Woo, N. T., Sakata, K., Zhen, S., Teng, K. K., Yung, W. H., Hempstead, B. L., & Lu, B. (2004). Cleavage of proBDNF by tPA/plasmin is essential for long-term hippocampal plasticity. *Science*, *306*(5695), 487–491. https://doi.org/10.1126/science.1100135

Park, H., & Poo, M. M. (2013). Neurotrophin regulation of neural circuit development and function. In *Nature Reviews Neuroscience* (Vol. 14, Issue 1, pp. 7–23). Nat Rev Neurosci. https://doi.org/10.1038/nrn3379

Persoon, C. M., Hoogstraaten, R. I., Nassal, J. P., van Weering, J. R. T., Kaeser, P. S., Toonen, R. F., & Verhage, M. (2019). The RAB3-RIM Pathway Is Essential for the Release of Neuromodulators. *Neuron*, *104*(6), 1065-1080.e12. https://doi.org/10.1016/j.neuron.2019.09.015

Persoon, C. M., Moro, A., Nassal, J. P., Farina, M., Broeke, J. H., Arora, S., Dominguez, N., Weering, J. R., Toonen, R. F., & Verhage, M. (2018). Pool size estimations for dense-core vesicles in mammalian CNS neurons . *The EMBO Journal*, *37*(20), 1–18. https://doi.org/10.15252/embj.201899672

Puntman, D. C., Arora, S., Farina, M., Toonen, R. F., & Verhage, M. (2021). Munc18-1 Is Essential for Neuropeptide Secretion in Neurons. *Journal of Neuroscience*, *41*(28), 5980–5993. https://doi.org/10.1523/JNEUROSCI.3150-20.2021

Rossi, V., Picco, R., Vacca, M., D'Esposito, M., D'Urso, M., Galli, T., & Filippini, F. (2004). VAMP subfamilies identified by specific R-SNARE motifs. *Biology of the Cell*, *96*(4), 251–256. https://doi.org/10.1016/j.biolcel.2003.12.009

Schiavo, G. G., Benfenati, F., Poulain, B., Rossetto, O., De Laureto, P. P., Dasgupta, B. R., & Montecucco, C. (1992). Tetanus and botulinum-B neurotoxins block neurotransmitter release by proteolytic cleavage of synaptobrevin. *Nature*, *359*(6398), 832–835. https://doi.org/10.1038/359832a0

Schlüter, O. M., Basu, J., Südhof, T. C., & Rosenmund, C. (2006). Rab3 superprimes synaptic
vesicles for release: Implications for short-term synaptic plasticity. *Journal of*
Neuroscience, *26*(4), 1239–1246. https://doi.org/10.1523/JNEUROSCI.3553-05.2006

Schlüter, O. M., Schmitz, F., Jahn, R., Rosenmund, C., & Südhof, T. C. (2004). A complete
genetic analysis of neuronal Rab3 function. *Journal of Neuroscience*, *24*(29), 6629–
6637. https://doi.org/10.1523/JNEUROSCI.1610-04.2004

Schlüter, O. M., Schnell, E., Verhage, M., Tzonopoulos, T., Nicoll, R. A., Janz, R., Malenka,
R. C., Geppert, M., & Südhof, T. C. (1999). Rabphilin knock-out mice reveal that
rabphilin is not required for rab3 function in regulating neurotransmitter release. *The*
Journal of Neuroscience : The Official Journal of the Society for Neuroscience, *19*(14),
5834–5846. https://doi.org/10.1523/JNEUROSCI.19-14-05834.1999

Schmitz, S. K., Hjorth, J. J. J., Joemai, R. M. S., Wijntjes, R., Eijgenraam, S., de Bruijn, P.,
Georgiou, C., de Jong, A. P. H., van Ooyen, A., Verhage, M., Cornelisse, L. N., Toonen,
R. F., & Veldkamp, W. (2011). Automated analysis of neuronal morphology, synapse
number and synaptic recruitment. *Journal of Neuroscience Methods*, *195*(2), 185–193.
https://doi.org/10.1016/j.jneumeth.2010.12.011

Shimojo, M., Courchet, J., Pieraut, S., Torabi-Rander, N., Sando, R., Polleux, F., & Maximov,
A. (2015). SNAREs Controlling Vesicular Release of BDNF and Development of
Callosal Axons. *Cell Reports*, *11*(7), 1054–1066.
https://doi.org/10.1016/j.celrep.2015.04.032

Stanic, J., Carta, M., Eberini, I., Pelucchi, S., Marcello, E., Genazzani, A. A., Racca, C., Mulle,
C., Di Luca, M., & Gardoni, F. (2015). Rabphilin 3A retains NMDA receptors at synaptic
sites through interaction with GluN2A/PSD-95 complex. *Nature Communications*, *6*(1),
10181. https://doi.org/10.1038/ncomms10181

Stanic, J., Mellone, M., Napolitano, F., Racca, C., Zianni, E., Minocci, D., Ghiglieri, V., Thiolat,
M. L., Li, Q., Longhi, A., De Rosa, A., Picconi, B., Bezard, E., Calabresi, P., Di Luca, M.,
Usiello, A., & Gardoni, F. (2017). Rabphilin 3A: A novel target for the treatment of
levodopa-induced dyskinesias. *Neurobiology of Disease*, *108*, 54–64.
https://doi.org/10.1016/J.NBD.2017.08.001

Staunton, J., Ganetzky, B., & Nonet, M. L. (2001). Rabphilin potentiates soluble N-
ethylmaleimide sensitive factor attachment protein receptor function independently of
rab3. *Journal of Neuroscience*, *21*(23), 9255–9264.
https://doi.org/10.1523/jneurosci.21-23-09255.2001

Südhof, T. C. (2013). Neurotransmitter release: The last millisecond in the life of a synaptic
vesicle. In *Neuron* (Vol. 80, Issue 3, pp. 675–690). Cell Press.
https://doi.org/10.1016/j.neuron.2013.10.022

Tsuboi, T., & Fukuda, M. (2005). The C2B domain of rabphilin directly interacts with SNAP-
25 and regulates the docking step of dense core vesicle exocytosis in PC12 cells.

Journal   of   Biological   Chemistry,   *280*(47),   39253–39259.
https://doi.org/10.1074/jbc.M507173200

van de Bospoort, R., Farina, M., Schmitz, S. K., de Jong, A., de Wit, H., Verhage, M., &
Toonen, R. F. (2012). Munc13 controls the location and efficiency of dense-core vesicle
release   in   neurons.   *Journal   of   Cell   Biology*,   *199*(6),   883–891.
https://doi.org/10.1083/jcb.201208024

van den Pol, A. N. (2012). Neuropeptide Transmission in Brain Circuits. *Neuron*, *76*(1), 98–
115. https://doi.org/10.1016/j.neuron.2012.09.014

van Westen, R., Poppinga, J., Arazola, R. D., Toonen, R. F., & Verhage, M. (2021).
Neuromodulator release in neurons requires two functionally redundant calcium
sensors. *Proceedings of the National Academy of Sciences of the United States of*
America, *118*(18). https://doi.org/10.1073/pnas.2012137118

Wierda, K. D. B., Toonen, R. F. G., de Wit, H., Brussaard, A. B., & Verhage, M. (2007).
Interdependence of PKC-Dependent and PKC-Independent Pathways for Presynaptic
Plasticity. *Neuron*, *54*(2), 275–290. https://doi.org/10.1016/j.neuron.2007.04.001

Zaben, M. J., & Gray, W. P. (2013). Neuropeptides and hippocampal neurogenesis. In
Neuropeptides   (Vol.   47,   Issue   6,   pp.   431–438).   Neuropeptides.
https://doi.org/10.1016/j.npep.2013.10.002

 Supplementary figures

Supplementary figure 1. RPH3A localization depends on RAB3A/RAB27A-binding. (**A**) Representative images of WT hippocampal and striatal neurons stained for RPH3A (white), VGLUT/VGAT (not shown) and MAP2 (red). N numbers of individual experiments and single neuron observations in brackets: HC: 2(13); STR: 2(12). Scale bar, 50 μm. (**B**) RPH3A intensity in MAP2+ neurite mask in hippocampal and striatal neurons, normalized to batch. (**C**) Domain structures of FL RPH3A and mutant RPH3A constructs lacking specific interactions: ΔRAB3A/RAB27A; truncated RPH3A that lacked its calcium and SNAP25 binding C2A and C2B domains; RPH3A with two C2B domain mutations to reduce calcium binding affinity (ΔCa^{2+} binding); RPH3A that lacked a CaMKII dependent phosphorylation site (ΔCAMKII-dependent phosphorylation site) and ΔSNAP25. The corresponding mutation sites are indicated in red. (**D, E**) Representative STED images of WT neurons expressing either FL RPH3A, ΔRAB3A/RAB27A, truncated RPH3A, ΔCa^{2+}-binding, ΔCAMKII-dependent phosphorylation site or ΔSNAP25, immunostained for mCherry (pseudo-colored blue), MAP2 (grey), Syn1 (magenta) or Homer (green). Scale bar, 5 μm. (**F**) Mander's overlap coefficient analyses of mCherry with either Syn1 or Homer, in WT neurons expressing FL or mutant RPH3A constructs. (**G**) RPH3A expression in KO neurons expressing FL RPH3A or mutant RPH3A constructs normalized to WT per independent experiment. N numbers per condition: KO + RPH3A: 6(24); KO + ΔRAB3A/RAB27A: 3(12); truncated RPH3A: 6(28); KO + ΔCa2+ binding: 5(23); and KO + ΔCAMKII-dependent phosphorylation: 2(9). (**H**) RPH3A expression in KO neurons expressing FL RPH3A or mutant RPH3A unable to bind SNAP25. N numbers per condition: KO + RPH3A: 1(3); KO + ΔSNAP25: 1(6). Each dot represents a single neuron. Kruskal-Wallis H test with Dunn's correction: ns = non-significant, $P > 0.05$.

Supplementary figure 2. RPH3A depletion increases DCV exocytosis, but does not affect remaining DCV pool size or content. (**A, B**) DCV fusion events per condition for the first and second 8 x 50 APs bursts at 50 Hz. (**C**) Estimate of the remaining NPY-pHluorin labelled DCV pool during 2 x 8 x 50 APs at 50 Hz. (**D, E**) Fusion duration during first and second 2 x 8 burst stimulation protocol. (**F**) Fusion duration during 16 x 50 Hz stimulation protocol. (**G**) Estimate of the remaining NPY-pHluorin labelled DCV pool during 16 x 50 APs at 50 Hz. (**H**) The number of spontaneous DCV fusion events per cell during baseline. (**I**) Mean peak intensity of NPY-pHluorin during live imaging per independent experiment. The black dashed line represents the median. Boxplots show the mean (+), median (line) and Tukey range (whiskers). Each dot represents a single neuron. Kruskal-Wallis H test with Dunn's correction: * $P < 0.05$, ** $P < 0.01$. ns = non-significant, $P > 0.05$.

Supplementary figure 3. Increased neurite length upon RPH3A depletion does not depend on RAB3A/RAB27A-
binding, calcium binding or phosphorylation of RPH3A. (**A**) Total dendritic and (**B**) axonal length of WT, KO and
KO neurons expressing FL or mutant RPH3A (ΔRAB3A/RAB27A, truncated, ΔCa2+ binding or ΔCAMKII-
dependent phosphorylation). N numbers per condition: WT 3(25); KO 3(28); KO + RPH3A: 3(29); KO +
ΔRAB3A/RAB27A: 3(32); truncated RPH3A: 3(29); KO + ΔCa2+ binding: 3(30); and KO + ΔCAMKII-dependent
phosphorylation: 3(30). (**C**) Total dendritic length of WT, KO and KO neurons expressing ΔSNAP25 mutant. N
numbers per condition: WT 3(17); KO 3(21); KO + ΔSNAP25: 3(20). Each dot represents a single neuron. Kruskal-
Wallis H test with Dunn's correction: * $P < 0.05$. ns = non-significant, $P > 0.05$.