Unbiased identification of cell identity in dense mixed neural cultures.

Sarah De Beuckeleer[1], Tim Van De Looverbosch[1], Johanna Van Den Daele[1], Peter Ponsaerts[2],

and Winnok H. De Vos[1,3,4]

[1] Laboratory of Cell Biology and Histology, University of Antwerp

[2] Laboratory of Experimental Hematology, Vaccine and Infectious Disease Institute

(Vaxinfectio), University of Antwerp

[3] Antwerp Centre for Advanced Microscopy, University of Antwerp

[4] µNeuro Research Centre of Excellence, University of Antwerp

e-mail:

sarah.debeuckeleer@uantwerpen.be

tim.vandelooverbosch@uantwerpen.be

johanna.vandendaele@uantwerpen.be

peter.ponsaerts@uantwerpen.be

winnok.devos@uantwerpen.be

Contact information *Corresponding author:

University of Antwerp, Laboratory of Cell Biology and Histology

Universiteitsplein 1, 2610 Antwerp, Belgium

e-mail: winnok.devos@uantwerpen.be

Keywords: morphological phenotyping, multiplexed imaging, iPSC culture validation, neural

differentiation, computational biology

Abstract

Induced pluripotent stem cell (iPSC) technology is revolutionizing cell biology. However, the variability between individual iPSC lines and the lack of efficient technology to comprehensively characterize iPSC-derived cell types hinder its adoption in routine preclinical screening settings. To facilitate the validation of iPSC-derived cell culture composition, we have implemented an imaging assay based on cell painting and convolutional neural networks to recognize cell types in dense and mixed cultures with high fidelity. We have benchmarked our approach using pure and mixed cultures of neuroblastoma and astrocytoma cell lines and attained a classification accuracy above 96%. Through iterative data erosion we found that inputs containing the nuclear region of interest and its close environment, allow achieving equally high classification accuracy as inputs containing the whole cell for semi-confluent cultures and preserved prediction accuracy even in very dense cultures. We then applied this regionally restricted cell profiling approach to evaluate the differentiation status of iPSC-derived neural cultures, by determining the ratio of postmitotic neurons and neural progenitors. We found that the cell-based prediction significantly outperformed an approach in which the time in culture was used as classification criterion (96% *vs.* 86%, resp.). In mixed iPSC-derived neuronal cultures, microglia could be unequivocally discriminated from neurons, regardless of their reactivity state. A tiered strategy, allowed for discriminating microglial cell states as well, albeit with lower accuracy. Thus, morphological single cell profiling provides a means to quantify cell composition in complex mixed neural cultures and holds promise for use in quality control of iPSC-derived cell culture models.

Introduction

Modelling the complexity of the human brain and its (dys)function has proven to be notoriously challenging. This is due to its intricate wiring, the cellular heterogeneity and species-specific differences[1]. To increase the translational value of neuroscientific research, there is a need for physiologically relevant human models. With the advent of human induced pluripotent stem cell (iPSC) technology, it has become possible to generate a wealth of brain-resident cell types including neurons[2], astrocytes[3], microglia[4], oligodendrocytes[5] and endothelial cells[6], allowing the study of complex polygenic pathologies that cannot be modelled in animals, opening avenues to precision pharmacology[7,8]. Furthermore, the ability of iPSC to develop into organoids renders them attractive for studying multi-cellular interactions in a 3D context that is closer to the *in vivo* situation[9]. However, genetic drift, clonal and patient heterogeneity cause variability in reprogramming and differentiation efficiency[10,11]. The differentiation outcome is further strongly influenced by variations in protocol[12]. This can significantly impact experimental outcomes, leading to inconsistent and potentially misleading results and consequently, it hinders the use of iPSC-derived cell systems in systematic drug screening or cell therapy pipelines. This is particularly true for iPSC-derived neural cultures, as their composition, purity and maturity directly affect gene expression and functional activity, which is essential for modelling neurological conditions[13,14]. Thus, from a preclinical perspective, there is the need for a fast and cost-effective quality control (QC) approach to increase experimental reproducibility and cell type specificity[15]. From a clinical perspective in turn, robust QC is required for safety and regulatory compliance (e.g., for cell therapy). This need for improved standardization is underscored by large-scale collaborative efforts such as the International Stem Cell Banking Initiative[16], which focusses on clinical quality attributes and provides recommendations for iPSC validation testing for use as cellular therapeutics, or the CorEuStem network, aiming to harmonize iPSC practices across core facilities in Europe.

Current culture validation methods include (combinations of) sequencing, flow cytometry and immunocytochemistry[15,17]. These methods are often low in throughput, costly and/or destructive. Thence, we set out to develop a method for evaluating the composition of such cultures based on high-content imaging, which is fast, affordable and scalable. The primary goal was to facilitate cell type identification in dense cultures, while ensuring compatibility with subsequent immunocytochemistry or molecular assays for further biological inquiries. To this end, we employed the Cell Painting (CP)[18,19] method, which is based on labelling cells with simple organic dyes and analysing the resulting phenotypes. CP has proven to be a powerful and generic method for predicting the mode-of-action of pharmacological or genetic perturbations, and this sheerly using a cell morphology readout[20–22]. Thus far, CP has primarily been utilized to predict conditions associated with pharmacological treatments or genetic modifications using images as input. For example, CP has allowed predicting patient diversity in lung adenocarcinoma-associated somatic variants[23]. or genetic variation across donors in iPSC cultures [24]. Methods such as PhenoRipper[25] and CP-CHARM[26] use whole-image features for classification, which circumvent the difficulty of cell segmentation allow classification of images with cells of similar phenotype or class [27]. However, they do not consider differences in cell density and disregard the inter-cellular heterogeneity within the field of view. Therefore, we explored the amenability of CP to predict individual cell types in dense and mixed cultures. By combining deep learning for cell segmentation and classification, we established an approach that allows recognizing cell types with high accuracy, even in very dense cultures. We employed the approach to evaluate the composition of iPSC-derived neural cultures as they are often quite dense and composed of heterogenous cell types. Varying the density, composition, and cell state allowed us to benchmark the discriminatory potential of cell-based profiling.

Results

Neural cell lines have a unique morphotextural fingerprint.

Several groups have demonstrated that morphological cell profiling can be used to discriminate pharmacogenomic perturbations based on phenotypic similarity[20–22]. We asked whether a similar approach could be leveraged to unequivocally distinguish individual cell types as well. To this end, we implemented a version of CP based on 4-channel confocal imaging (**Fig. 1A**) and first applied it to monocultures of two neural cell lines from a different lineage, namely astrocyte-derived 1321N1 astrocytoma cells and neural crest-derived SH-SY5Y neuroblastoma cells. First, we explored whether traditional morphotextural feature extraction provided sufficient distinctive power. The features were calculated for each channel in three regions of interest, namely the nucleus, cytoplasm and whole cell. They describe shape, intensity and texture features of each ROI (**Table 5**). Representation of the resulting standardized feature set in UMAP space revealed a clear separation of both cell types along the first UMAP dimension. Clustering of instances was less pronounced after principal component analysis (**Suppl. Fig. 1A**), but UMAP better preserves local and global data structure [28]. Despite some separation of replicates across the second UMAP dimension, the absence of clear replicate clusters showed that the morphological differences between cell types were consistent across biological replicates (**Fig. 1B**). When projecting individual features onto the UMAP space, we found that both texture (*e.g.,* Nucleus channel 3 energy) and shape (*e.g.,* Cellular Area) metrics contributed to the cell type separation (**Fig. 1C**). The contribution of intensity-related features (*e.g.,* Channel 3 Intensity) to cell type separation was less pronounced as they were more correlated with the biological replicate. Thus, we conclude that cell types can be separated across replicates based on a morphotextural fingerprint.

CNN outperforms random forest in cell type classification.

To evaluate whether the morphotextural fingerprint could be used to predict cell type, we

performed Random Forest (RF) classification using the full feature dataset, using different

seeds for splitting up the data into training and validation sets. This resulted in a rather poor

accuracy (F-score: 0,75±0,01), mainly caused by the significant (46%) misclassification of

1321N1 cells (**Fig. 1D**). The imbalance in recall and precision was surprising given the clear

separation of cell populations in UMAP space using the same feature matrix. When inspecting

the main contributions to the RF classifier using the mean decrease in impurity, we found very

similar features as highlighted in UMAP space to add to the discrimination (**Fig. 1C**). Where

the most important features (*e.g.,* cellular area, channel 1 contrast) showed the expected

gradient along the first UMAP direction, lower ranked parameters (e.g., cellular channel 3

mean intensity) had no contribution to UMAP separation (**Fig. 1C**). Reducing noise by

removing redundant features (correlation > 0.95) could not ameliorate RF classification

performance, which may be due to the documented bias in feature selection for node splitting

in high-dimensional data[29]. This result drove us to evaluate a different classification approach

based on a ResNet[30] convolutional neuronal network (CNN). Here, we no longer relied on the

extraction of "hand-crafted" features from segmented cell objects for training the shallow RF

classifier. Instead, we used isotropic image crops of 60µm centred on individual cell centroids

and blanked for their surroundings as input for the CNN (**Fig. 2A**). Using this approach, we

found a significantly higher prediction performance with an F-score of 0,96±0,01, with a much

more balanced recall and precision (**Fig. 2B**). The UMAP space built from the CNN feature

embeddings showed a clear cell type separation (**Suppl. Fig. 1B**). Even with only 100 training

instances per class, the CNN outperformed RF, but optimal performance was attained with

5000 training instances (**Fig. 2C**). A model trained on a combination of three biological

replicates ("Mixed reps.") outperformed those that were only trained on a single replicate

("Single rep.") when predicting instances of an independent unseen replicate (**Fig. 2E**). This

emphasizes the need for including sufficient variation in the training set. Although much more

performant, CNN classification does not allow direct retrieval of intuitive features, which

complicates model interpretation. To gain a visual understanding of image information

contributing most to the classification we therefore resorted to Gradient-weighted Class

Activation Mapping (Grad-CAM)[31]. This revealed that the attention of the CNN was mainly

focused on cell borders (edge information), nuclear and nucleolar signals (**Fig. 2F**). When

scrutinizing CNN misclassifications, we found that these are mainly caused by faulty cell

detection (*e.g.,* oversegmentation of cell ramifications), unhealthy/dead cells, debris or visibly

aberrant cell shapes (**Suppl. Fig. 1C**) - errors not necessarily attributed to the CNN.

To shed light on the contribution of individual markers to the classification, we eroded the

input to single channel images. For all cases, the prediction performance was below or equal

to 85,0% suggesting that no single channel contains all relevant information (**Fig. 3A**).

Combinations of two or three channels could not match the prediction accuracy of the full four-

channel image either. Thus, we concluded that all channels contribute to the successful

classification. As the CNN directly uses image crops as input, the image quality will determine

the classification performance. Therefore, we assessed the impact of resolution and signal to

noise ratio (SNR). We simulated the effect of decreasing spatial resolution through progressive

pixel binning (from 192 pixels (original, pixel size 0.3µm) to 9 pixels (pixel size 6 µm)) (**Fig.**

3B). For each iteration, 3 CNN models were trained and evaluated. A reduction in pixel size

from 0.3µm to 0.6µm did not result in a significantly lower prediction performance, but

decreasing the spatial resolution further caused a progressive decrease in F-score. In a similar

manner, we tested the impact of decreasing SNR (by increasing the level of Gaussian noise) on

classification accuracy. Starting from an original SNR value of 20,05 dB, we found that the F-

score started to decrease when lowering the SNR below 14,27 dB.

We observed that even with low input image quality, the CNN performance still retained approximately 80% accuracy. We attribute this to the predictive power of the nuclear size, which is a dominant discriminating feature between the cells (**Suppl. Fig. 2**). Given the relatively overt phenotypic difference between 1321N1 and SH-SY5Y cells, we asked whether the CNN could also discriminate more subtle phenotypes. To this end, we used 1321N1 cells and two related retinal pigment epithelial cell lines, hTERT-immortalized RPE1 and the spontaneously immortalized variant ARPE (**Fig. 3D**). The CNN discriminated all three cell lines with high accuracy of 95,06±2,51%. Unsupervised UMAP dimensionality reduction using the CNN feature embeddings revealed three clusters of ROIs, with the two RPE lines located closer together (but well separated) in comparison to the morphologically more distinct 1321N1 cell type (**Fig. 3D**). Together this work illustrates that a CNN approach can be used to distinguish diverse cell types within a given image quality window.

Nucleocentric predictions remain accurate regardless of culture density.

Morphological profiling relies on accurate cell detection. This may become difficult in dense cultures such as iPSC-cultures, clustered cells tissues and tissue-mimics. Having established a method to distinguish cell types with high accuracy, we next asked how robust the classification was to increasing cell density. To this end, we acquired images of 1321N1 and SH-SY5Y monocultures, grown at densities ranging from 0 to 100% confluency (**Fig. 4A**). Based on the nuclear count, we binned individual fields into 6 density classes (0-20%, 20-40%, 40-60%, 60-80%, 80-95%, 95-100%) and trained a CNN with equal sampling of cell numbers per density class to avoid bias. No decrease in accuracy was observed until the culture density reached 80% confluency. Only for very high densities (95-100%), we found a significant decrease in the prediction accuracy (F-score: 0,92±0,05) (**Fig. 4B**). We reasoned that under these conditions cell shape would be predominantly determined by neighbouring cells and cell

segmentation performance would decrease (**Suppl. Fig. 1D**). Nuclei are less malleable and even in dense cultures remain largely separated, allowing their robust segmentation and avoiding CNN misclassifications resulting from segmentation errors. Hence, we asked whether using the nuclear ROI as input would improve classification performance at high densities. However, despite the relatively high average F-score of 0,91±0,05 (**Fig. 4C**), the performance was consistently lower than whole cell ROI across the density range and the performance still decreased with full confluency. To understand these results, we inspected the GradCAM output for these predictions, and found that an important part of the attention of the CNN is diverted to the background (**Fig. 4D, Suppl. Fig. 3**). We interpret this result as the CNN using the background as a proxy for nuclear size. To rule out bias by setting the background to zero in nuclear crops, the CNN was also trained on the same crops with randomly speckled background, but despite a shift in attention to the nucleus, similar prediction performance was attained (**Suppl. Fig. 2**). The classification performance was not biased by segmentation errors (**Suppl. Fig. 1E**) but may be influenced by the fact that nuclear area is affected by culture density (**Suppl. Fig. 1F**). In highly dense cultures, the dynamic range of the nuclear size decreases as all nuclei become more compact and the nuclear size range decreases (**Suppl. Fig. 1F**). Thence, we tested an intermediate condition which exploits the more robust nuclear segmentation but also includes part of the (sub-)cellular local surrounding information as input (**Suppl. Fig. 1G**). To identify the optimal patch size, we varied the size of a square box centred around the nuclear centroid from 0.6 to 150 µm (**Fig. 4E**). Within a range of 12-18µm, we found a maximal F-score of 0,96±0,02 (**Fig. 4E**). Further increasing the nucleocentric patch size did not majorly affect the F-score but significantly decreased the precision and variability of recall for CNN predictions (**Fig. 4E**). Hence, for further experiments, a nucleocentric patch size of 18µm was used. GradCAM images revealed that this latter approach led the CNN to focus on perinuclear structures (**Fig. 4D and Suppl. Fig. 3**). Interestingly, when using this

nucleocentric approach, the prediction performance was maintained at almost confluent cell densities in contrast with whole cell approaches (**Fig. 4B**). Thus, we conclude that using a nucleocentric region as input for the CNN could be a valuable strategy for accurate cell type identification in dense cultures.

Cell prediction remains accurate in mixed cell culture conditions.

Although a very high classification accuracy was obtained with nucleocentric CNN predictions, both training and testing were performed with input images drawn from monocultures. As our goal was to allow cell type prediction in complex, heterogeneous cultures, we next switched to a more faithful situation in which we co-cultured both cell types. Ground-truth for these predictions was generated by either performing post-hoc immunofluorescence (IF) with cell-type specific antibodies (CD44 for 1321N1 and TUBB3 for SH-SY5Y cells), or by differential replication labelling (*i.e.,* by incubating the two cell types with EdU and BrdU respectively, prior to mixing them) after dye quenching[32] (**Fig. 5A**). Replication labelling proved significantly more successful for binary ground truth classification (through intensity thresholding) than IF labelling for the cell lines we used (**Fig. 4B**). When training a CNN to recognize the cell types using these markers as ground truth, we found that the prediction accuracy on the left-out dataset of the co-culture (Co2Co) was almost as high as when a model was trained and tested on monocultures (Mono2Mono) (**Fig. 5C**). We then tested whether it was possible to train a classifier based on monocultures only, for cell type prediction of cells in co-culture (Mono2Co). This resulted in an F-score of 0,86±0,01%. This drop in performance may be caused by the effect on cell phenotype exerted by the presence of other cell types in mixed cultures which is not captured in monocultures. As the images from monocultures and co-cultures were obtained from different plates, we suspected inter-replicate variability in culture and staining procedures to contribute in part to the lesser performance.

Therefore, we tested whether we could improve the performance of the CNN by including

monocultures from the same plate as the co-cultures. This finetuning indeed improved the

average performance to 0,88±0,01%, and more importantly, it significantly reduced the

variability (coefficient of variation) of the predictions making it more reproducible (**Fig. 5D**).

Thus, while not yet reaching the same accuracy, it proves that it is possible to establish a model

that recognizes cell types in co-cultures that is solely trained on monocultures.

Cell type profiling can be applied to stage iPSC-derived neuronal cultures.

iPSC-derived neuronal cell cultures suffer from significant inter- and intra-batch variability

and could benefit from an efficient quality control[14,33]. Thence, we applied our nucleocentric

phenotyping pipeline to stage the maturity of a neuronal cell culture based on its cell type

composition. Using a guided protocol[2], two differentiation stages were simulated: a primed

stage, where most cells are assumed to be cycling neural progenitors (NPCs), and a

differentiated stage where most cells are post-mitotic neurons (**Fig. 6A**). The two cell types

were discriminated by post-hoc IF labelling for the cell cycle marker Ki67 (NPC) and the

microtubule marker ß-III-tubulin (TUBB3, neurons) (**Fig. 6B**). Not all cells in the CP image

could be assigned with a ground truth label due to cell detachment upon re-staining or sheer

absence of the tested marker. Since no single monoculture consists of 100% of either cell type,

we applied gates to retain only those cells that show unequivocal staining for either one of both

markers. Based on these gates, ROIs were either classified as neuron (Ki67-/TUBB3+), NPC

(Ki67+/TUBB3-) or undefined (outside of gating boundaries). We assume the latter category

to represent transitioning cells in intermediate stages of neural development, un- or

dedifferentiated iPSCs. This gating strategy resulted in a fractional abundance of neurons vs.

total (neurons + NPC) of 36,4 % in the primed condition and 80,0% in the differentiated

condition (**Fig. 6C**). In a first attempt to classify cells within the guided culture, we trained a

CNN on individual cell inputs using the culture condition (primed or differentiated) as ground truth, which given the heterogeneity, can be considered as a weak label. This resulted in a low F-score of 0,86±0,01%. When we used the cell-level IF ground truth labels instead, we obtained a classification performance of 0,96±0,00 %. For comparison, a shallow learner (RF), showed a significantly lower F-score of 0,87±0,02 (**Fig. 6D**). Applying this cell-based (as opposed to condition-based) CNN to the two culture conditions resulted in predicted fractional abundance of neurons to NPC of 40,5% in images of the primed condition and 74,2% in images of the differentiated condition – aligning well with the manually defined ratios (**Fig. 6E**). Both supervised and unsupervised UMAP dimensionality reduction on the feature embeddings of the cell-based classifier revealed a clear clustering of both phenotypes suggesting that the CNN captures the differences in morphotextural fingerprint between neurons and NPCs well (**Fig. 6F**). We then went on to evaluate the established cell-based classifier to a primed neuronal cell culture undergoing gradual spontaneous differentiation after dual SMAD inhibition[34]. We examined cell cultures at 13, 30, 60 and 90 days *in vitro* (DIV) after the start of the differentiation process and visually confirmed a gradual change in neural maturity. This is evidenced by a gradually smaller cell body and long, thin ramifications (**Fig. 6G**). The cell-based CNN model trained on the guided differentiation dataset tested on the spontaneous cultures, confirmed a shift in the (slower) neuron-to-NPC fractional abundance with increasing time in culture (**Fig. 6H**).

Cell type identification can be applied to mixed iPSC-derived neuronal cultures regardless of the activation state

Next to NPCs and neurons, iPSC-derived neuronal cultures are often studied in conjunction with other relevant cell types that influence neuronal connectivity and homeostasis such as astrocytes and microglia[13]. Therefore, we tested whether the cell-based approach could be

extended to these cell types as well. We generated monocultures of iPSC-derived astrocytes, neurons, and microglia from the same parental iPSC line (**Fig. 7A**) and trained a nucleocentric CNN to identify each cell type. This led to a prediction accuracy of 96,81±0,95% (**Fig. 7A**). Recognizing that these cell types are visually morphologically distinct, we tested the robustness of the model to experimental perturbations in cell state (**Fig. 7B**). To this end, we induced reactivity in the iPSC-derived microglial culture by LPS treatment. A CNN trained on monocultures of neurons, unchallenged or LPS-treated microglia showed high accuracy in differentiating neurons from microglia (98% accurate), although the difference between reactive and resting-state microglia in this tripartite model proved more challenging (71% and 90% accuracy, resp.). However, when implementing a tiered approach, in which after neuron-microglia recognition, a second model was tasked with the classification of reactive vs. resting state microglia without the presence of neurons, an F-score of 0,92±0,00 was obtained. Repeating those experiments in mixed cultures of neurons and microglia (using Tubb3, resp. Iba1 as ground-truth IF markers), yielded an F-score of 0,98±0,01 for the classification of neurons vs. microglia. Again for comparison, CNN outperformed the classical RF approach (0,86±0,03) (**Fig. 7C**). In the presence of LPS, a CNN model yielded a high F-score of 0,97±0,01 (**Fig. 7C**). This implies that the shift in cell state does not significantly affect the CNN's ability to distinguish neurons from microglia. As above, a separate CNN was trained to classify within the microglial subpopulation the LPS-treated (reactive) from the unchallenged (resting-state) microglia. Yet, this phenotype proved more challenging for the CNN to predict, as evidenced by an F-score of 0,80±0,01 (**Fig. 7C**). Based on these results, we conclude that nucleocentric phenotyping can be used to gauge the cell type composition of iPSC-derived cultures. Distinct cell types show sufficient morphological differences to allow highly accurate classification. Different cell states of individual cell types can still be recognized albeit with lower performance.

Discussion

iPSC-derived cell cultures have the potential to improve translatability of biomedical research but represent a challenge due to their variability and multicellular composition. With our work, we have developed a method to identify neural cell types in mixed cultures to aid in their validation and application in routine screening settings. We first benchmarked our approach using neural cell lines and found that a CNN-based approach outperforms shallow learning (RF) based on handcrafted feature extraction, even with a limited number of input images. This aligns well with recent data showing CNN-based feature representations outperform shallow techniques for condition-based drug response prediction[35]. To assess the robustness and sensitivity of deep learning for cell type prediction, we evaluated the CNN performance as a function of input data size and quality. Next, we tested the performance of predictions for dense single and multi-cellular cultures. Finally, we assessed its performance in the presence of different cells states.

Although our benchmarking revealed a minimal image resolution of 0,6 µm/pixel and SNR of approximately 10 dB for optimal CNN performance, lower quality images may still yield acceptable performance, especially with the use of advanced image restoration algorithms[36]. We found that all channels contribute to the overall prediction accuracy but that the whole cell is not required to obtain the highest prediction accuracy. In line with earlier studies that highlight the biomarker potential of the nucleus for patient stratification[37], cell state identification[38] or drug response[20], we found that the nuclear ROI as such already carries highly distinctive information for cell type prediction. However, extension to its direct surroundings proved to be the best and most robust input across a range of cell densities. The nucleocentric approach, which is based on more robust nuclear segmentation, reduces such mistakes whilst still retaining input information from the structures directly surrounding the nucleus. At higher cell density, the whole-cell segmentation becomes more error-prone, while also loosing

morphological information (**Suppl. Fig. 1D**). The nucleocentric approach is more consistent as it relies on a more robust segmentation and does not blank the surrounding region. This way it also buffers for occasional nuclear segmentation errors (*e.g.,* where blebs or parts of the nucleus are left undetected). GradCAM maps on nucleocentric crops highlight specifically those structures surrounding the nucleus (reflecting ER, mitochondria, Golgi) indicating their importance in correct cell classification. We observed a severe imbalance in precision and recall when altering the size of the nucleocentric window. This imbalance was not reflected into an overall worse F-score, since both target classes had an equal number of input instances. However, this imbalance would strongly affect the prediction performance of cases with unknown ground truth were the input numbers are imbalanced. Nucleocentric prediction of other cell types with less pronounced differences within their nucleus and close environment might yield a different optimum. It opens possibilities for applying cell profiling in the future to even more complex 3D cell systems such as tissue or organoid slices, where accurate cell segmentation becomes extremely challenging. Indeed, our results indicate that it might be possible to discriminate individual cells in extremely crowded environments. For specific applications such as classification of healthy vs. tumour tissue, binary classification models can prove useful. It has been shown that a dual-task graph neural network can classify epithelioid and sacromatoid tumor cells with cellular resolution for accurate mesothelioma subtype determination[39]. However, the rich diversity of cell types and intermediates present in tissue may complicate classification and demand a more extensive ground truth based on cyclic immunostaining to further refine the predictions. In the further future, it may even be possible to include volumetric information, but this will require optimization of the sample preparation procedures as well.

One crucial realisation of this work is that cell types can be identified in mixed cultures solely using the input from monocultures. This implies that cells retain their salient characteristics

even in more complex heterogeneous environments. While for now, predictions are still superior when based on training directly on mixed cultures, we found that the prediction accuracy of monoculture trained models can be increased by employing replicate controls. This suggests that it may become possible to apply or adapt existing models without the need for a cell-level ground truth as provided by post-hoc labelling. This technique could potentially be of use for cultures or tissues where no antibody- or pre-labelling is possible (*e.g.,* no unique IF marker is available, non-replicating cells). To increase the accuracy one could resort to intelligent data augmentation[40] or transfer learning[41] strategies. While a ground-truth independent method holds promise for bulk predictions (*e.g.,* for quality control purposes), the use of post-hoc labelling allows building more refined models that can distinguish multiple cell types and/or cell states at once. Especially using cyclic staining or spatial transcriptomics, much richer information content can be gained. With the method presented here, we combine the rich multidimensional information from cyclic IF with morphological information and show that it is possible to predict IF signatures. Multiplexed imaging has previously shown its utility in gaining in-depth information on culture composition and differentiation status in iPSC-derived neural cultures (progenitor, radial glia, astrocytes (immature and mature) and neurons (inhibitory and excitatory)[42]. Similarly, this could be expanded to cell states (apoptosis, DNA damage), albeit with a lesser performance depending on the phenotype[21]. Initial model training requires both CP and multiplexed IF images. After the initial training phase, the IF step can be omitted, and the CNN will predict the phenotype outcomes of the IF with high accuracy on the CP images alone. This significantly reduces experimental costs and time requirements.

Applying the method to iPSC-neuronal cultures revealed its potential to score their differentiation state. Although guided protocols manage to speed up the differentiation process and lead to enhanced culture purity, the neural differentiation process proves to be less than binary, as evidenced by the mixed presence of Ki67+/TUBB3- and Ki67-/TUBB3+ cells. The

spontaneous differentiation protocol illustrated the unpredictable nature and late stage of the

differentiation process. Many groups highlight the difficulty of reproducible neural

differentiation and attribute this to culture conditions, cultivation time and variation in

developmental signalling pathways in the source iPSC material[43,44]. Spontaneous neural

differentiation has previously been shown to require approximately 80 days before mature

neurons arise that can fire action potentials and show neural circuit formation. Although these

differentiation processes display a stereotypical temporal sequence[34], the exact timing and

duration vary. This variation negatively affects the statistical power when testing drug

interventions and thus prohibits the application of iPSC-culture derivatives in routine drug

screening. Current solutions (e.g., immunocytochemistry, flow cytometry, …) are often cost-

ineffective, tedious, and incompatible with longitudinal/multimodal interrogation. CP is a

much more cost-effective solution and ideally suited for this purpose. Routine CP-based could

add confidence to and save costs for the drug discovery pipeline. We have shown that CP can

be leveraged to capture the morphological changes associated with neural differentiation. We

could reliably predict the gradual transition from NPC to a more differentiated state. We realize

that differentiating iPSC cultures are highly heterogenous and are composed of a landscape of

transitioning progenitors in various stages of maturity that our binary models currently fail to

capture. As a result, we filtered out cells with a 'dubious' IF profile (e.g., cells that might be

transitioning or are of a different type) as they would negatively affect the model by introducing

noise. In future iterations, one could envision defining more refined cell (sub-)types in a

population based on richer post-hoc information (e.g., through cyclic immunofluorescence or

spatial single cell transcriptomics). While we emphasize the value of identifying fixed states as

a fast-track to gather the composition of an iPSC-derived culture, it is equally interesting to

focus on the continuum that such cultures represent. To predict the maturation of iPSC-derived

neural progenitors to differentiated neurons in a more continuous manner, would demand

regression-based approaches [45]. Pioneering efforts using live-cell imaging and machine

learning have allowed predicting gradual cell state transitions, for example in the context of

myoblast, adipocyte or osteoblast differentiation [46,47]. Label-free timelapse imaging has also

been shown to aid in the assessment of differentiation without disruption of the sample[46]. Given

that many of the cell painting dyes are compatible with live cells, it is conceivable that our

approach is amenable to further refining the assessment of iPSC differentiation status in real

time.

After testing the CNN performance on heterogenous cultures, we have added an additional

layer of complexity by inducing microglial reactivity in a coculture of neurons and microglia.

We found that we could still predict the cell type regardless of the treatment. The increased

variability within the microglial subpopulation did not impact the CNNs' ability to discriminate

cell types in mixed culture. Furthermore, using the layered approach, the resting-state and

reactive subgroups within the microglial population could also be classified, albeit with a lesser

prediction accuracy. This could be explained by the fact that microglia grown *in vitro* are not

completely homeostatic, even in absence of LPS [48], and may require different cell types to

adopt a more natural state. Having more faithful mixed cultures with cells in their endogenous

states, and an apt tool to recognize both cell type and state, holds promise to enhance the

relevance of preclinical screens, increase the accuracy of drug targets and ultimately lead to

more precise therapeutic strategies[49]. For now, we have tailored models to the individual

datasets, but it is conceivable that a more generalized CNN could be established for multiple

culture types. This would obviously demand a much larger dataset to encompass the variability

encountered in such models (*e.g.,* various starting iPSC lines, various differentiation protocols).

Publicly available datasets (*e.g.,* JUMP-Cell Painting Consortium) can aid in creating an input

database containing a large variability (different iPSC lines, different neural differentiation

protocols, …), which would ultimately lead to a more robust predictor. Our results showing

the prediction accuracy of a guided differentiation model on spontaneously differentiating

cultures indicate that the approach can be transferred to other differentiation protocols as well.

Inclusion of more input images and variability should thus enable developing a generalist

model for other differentiation protocols without the need for ground truth validation and

further CNN training.

In conclusion, we have developed a novel application for unbiased morphological profiling by

extending its use to complex mixed neural cultures using sequential multispectral imaging and

convolutional network-informed cell phenotype classification. We show that the resulting

predictors are robust with respect to biological variation and cell culture density. This method

holds promise for use in quality control of iPSC cultures to allow their routine use in high-

throughput and high-content screening applications.

Methods

Cell culture. Cells were cultured at 37 °C and 5% CO2. 1321N1 and SH-SY5Y cell lines were maintained in DMEM-F12 + Glutamax (Gibco, 10565018) supplemented with 10% Fetal Bovine Serum (Gibco, 10500064). Cell seeding prior to imaging was done in 96-well black multiwell plates with #1.5 glass-like polymer coverslip bottom (Cellvis, P96-1.5P). Only the inner 60 wells were used, while the outer wells were filled with PBS-/- to avoid plate effects. Plates were coated with Matrigel (Corning, 734-1440) After seeding, the imaging plate was centrifuged at 100g for 3min.

iPSCs (Sigma Aldrich, iPSC0028 Epithelial-1) were cultured on Matrigel (Corning, 734-1440) in Essential 8 medium (Gibco, A1517001). Upon cell thawing, Rock inhibitor (Y-27632 dichloride, MedChem, HY-10583) was added in 10μM concentration. Subculturing of iPSCs was performed with ReLeSR (Stemcell Technologies, 05872).

Unguided differentiation[34] of iPS cells to neural progenitor cells (NPCs) was started by subculturing the iPSCs single cell using Tryple Express Enzyme (Life technologies, 12605010) at a density of 10e4 cells/cm² in mTesR1 medium (Stemcell Technologies, 85850) and Rock inhibitor. The following day, differentiation to NPCs was started by dual-SMAD inhibition in neural maintenance medium (1:1 Neurobasal (Life technologies, 21103049):DMEM-F12+Glutamax (Gibco, 10565018), 0.5x Glutamax (Gibco, 35-050-061), 0.5% Mem Non Essential Amino Acids Solution (Gibco, 11140050), 0.5% Sodium Pyruvate (Gibco, 11360070), 50 μM 2-Mercaptoethanol (Gibco, 31350010), 0.025% Human Insulin Solution (Sigma Aldrich, I9278), 0.5X N2 (Gibco, 17502048), B27 (Gibco, 17504044), 50 U/ml Penicillin-Streptomycin (Gibco, 15140122)) supplemented with 1 μM LDN-193189 (Miltenyi, 130-106-540), SB431542 (Tocris, 1614). This dual blockade of SMAD signalling in iPSCs is induces neural differentiation by synergistically causing the loss of pluripotency and push towards neuroectodermal lineage[50]. Daily medium changes were performed for 11 consecutive

days. Following neural induction, neural rosettes were selected by STEMdiff Neural Rosette

Selection Reagent (Stemcell technologies, 05832). Maintenance of neural progenitor cells was

performed in neural maintenance medium with 10 µM bFGF (Peprotech, 100-18C) added after

subculturing. Single cell detachment of NPCs during maintenance was performed using Tryple

Express Enzyme. Cell seeding prior to imaging is done in 96-well black multiwell plates

µCLEAR (Greiner, 655090) coated with Poly-L-ornithine (Sigma-Aldrich, P4957) and laminin

(Sigma-Aldrich, L2020). Only the inner 60 wells were used, while the outer wells were filled

with PBS-/- to avoid plate effects. After seeding, the imaging plate was centrifuged at 100g for

3min.

Guided iPSC differentiation to neurons was performed according to Bell et al. (2019)[2]. The

initial phase of neural induction consisted of 12 days neural induction medium (DMEM-

F12+Glutamax (Gibco, 10565018), 1x N2 (Gibco, 17502048), 1x B27 (Gibco,

17504044),1mg/ml BSA (Sigma-Aldrich, A7979), 0.5% Mem Non-Essential Amino Acids

Solution (Gibco, 11140050)). Of these 12 days, the first 7 were supplemented with cytokines

for dual-SMAD inhibition (1 µM LDN-193189 (Miltenyi, 130-106-540), SB431542 (Tocris,

1614)). Following neural induction, the NPCs were floated in uncoated MW6 culture plates in

NPC medium (DMEM-F12+Glutamax (Gibco, 10565018), 1x N2 (Gibco, 17502048), 1x B27

(Gibco, 17504044), 10 µM bFGF (Peprotech, 100-18C), 10 µM EGF (Peprotech, 100-47)).

After 4 days, NPC clusters of appropriate size were filtered using a cell strainer ((37 µm cell

strainer, Stemcell technologies, 27250) and plated on Matrigel-coated (Corning, 734-1440)

MW6 culture plates. NPCs can now be expanded in NPC medium. Guided differentiation into

forebrain neurons can be induced by switching to neuronal medium (BrainPhys (STEMCELL

Technologies, 05792), 1x N2 (Gibco, 17502048), 1x B27 (Gibco, 17504044), 10 µM BDNF

(PeproTech, AF-450-02), 10 µM GDNF (PeproTech, AF-450-02) for 15 days before fixation.

Differentiation of iPSC to microglia[4] was performed by the formation of embroid bodies (EBs)

with 10e3 iPSCs/well in a 96-well U-bottom plate (Corning, 351177) coated with Anti-

Adherence Rinsing Solution (Stemcell technologies, 07010) in mTeSR medium supplemented

with Rock inhibitor, 50 ng/mL BMP4 (Peprotech, 120-05), 50 ng/mL VEGF (Peprotech, 100-

20), 20 ng/mL SCF (Peprotech, 250-03). 75% medium is changed for 4 consecutive days. After

mesoderm induction, EBs are transferred to a 6-well plate with 20 EBs/well and placed in

macrophage precursor medium (X-vivo15 (Lonza, BE02-060Q), 100 ng/mL M-CSF

(Peprotech, 300-25), 25 ng/mL IL-3 (Peprotech, 213-13), 1x Glutamax, 50 U/ml Penicillin-

Streptomycin, 50 µM 2-Mercaptoethanol). 14 days after macrophage differentiation,

macrophage precursors were harvested using a cell strainer (Stemcell technologies, 27250).

Macrophage precursors were added to the NPC culture in 1:1 neural maintenance

medium:microglia medium ((DMEM-F12+Glutamax, 100 ng/mL M-CSF (Peprotech, 300-25),

100 ng/mL IL-34 (Peprotech, 200-34), 1x Glutamax, 50 U/ml Penicillin-Streptomycin, 50 µM

2-Mercaptoethanol).

Replication labelling. Prior to co-seeding of 1321N1 and SH-SY5Y mixed cultures, individual

cultures were incubated with either 10 µM EdU (Click-iT® EdU Imaging Kit, Life

Technologies, C10340) or 10 µM BrdU (Sigma-Aldrich, B5002) for 24h. This incubation time

exceeded the doubling time, allowing incorporation of the nucleotide analog in all cells. This

labelling period was followed by a 24h washout period in regular cell culture medium. After

washout, the cells were subcultured and plated in coculture. In half of the replicate, SH-SY5Y

cells received BrdU while 1321N1 cells received EdU. For the remainder of wells, the pre-

label switched cell types.

Morphological staining. Morphological staining (Cell Painting) was adapted from Bray et al.

2016[19]. After careful titration, all dye concentrations were adjusted and evaluated for

compatibility with 4-channel laser and filter combinations available on the confocal

microscope (see further). Staining was performed on cell cultures fixed in 4% PFA (roti-histofix 4% paraformaldehyde, Roth, 3105.2) for 15min. Cells were rinsed once with PBS-/- (Life Technologies, 10010015) prior to fixation and 3x 5min post fixation. Staining solutions were prepared fresh before staining in PBS-/- with 0.3% Triton-X-100 (Sigma Aldrich, X100) (**Table 1**). Each staining solution was incubated for 30min on a shaker at RT in the dark. After staining, the cells were washed 1x with PBS -/- and sealed for imaging.

Cyclic staining and immunocytochemistry. Cyclic staining is executed by fluorescence quenching after each sequential imaging round. 1mg/ml in ddH$_2$O LiBH$_4$ solution[32] (Acros Organics, 206810050) was prepared fresh before use. 1.5h incubation of quenching solution was performed before each successive staining series. After incubation, the quenching solution was removed by washing 3x 5min in PBS-/-. Successful fluorescence quenching was microscopically verified before proceeding with immunofluorescence staining (IF) (**Table 2, Suppl. Fig. 4**).

Cells are treated with PAV blocking buffer (Thimerosal 0.5% (Fluka, 71230), NaN3 0.1% (Merck, k 6688), Bovine serum albumin (Sigma-Aldrich, A7284), Normal horse serum, PBS-/-) for 8min. The desired primary antibodies (pAB) are diluted in PAV blocking buffer. pAB incubation was performed 12h (overnight) at 4°C after which the cells were washed 1x 5min in PBS-/- followed by incubation in secondary antibody solution (sAB) in PAV + DAPI for nuclear counterstain. sAB staining was performed for 3h at RT while shaking. Prior to imaging, the cells were washed 2x with PBS-/- and stored in PBS-/- + 0,1% NaN$_3$.

BrdU staining was performed using IF, requiring DNA denaturation before pAB incubation. This was performed by 10 min incubation with 2N HCl at 37°C. HCl was neutralized with 0.1 M sodium borate buffer pH 8.5 for 30min at RT. Cells were washed 3x 5min in PBS-/- before continuing with the general IF protocol. EdU click-it labelling was performed according to the

manufacturer's instructions (Click-iT® EdU Imaging Kit, Life Technologies, C10340) (**Suppl.**

Fig. 5A).

Image acquisition. Images were acquired using a spinning disk confocal microscope (Nikon

CSU-W1 SoRa) with a 20x 0.75 NA objective (Plan APO VC 20x DIC N2) and Kinetix

sCMOS camera (pinhole 50µm; disk speed 4000 rpm; pinhole aperture 10; bit depth 12-bit,

pixel size 0.325µm²). We opted for confocal microscopy instead of widefield to overcome

image quality limitations resulting from highly dense cell clusters. 96-well plates were scanned,

capturing a minimum of 16 images per well spread in a regular pattern (0,8 mm spacing in x

and y) across the surface of the well. If multiple z-slices were acquired (to correct for surface

inclinations in the field of view), a maximum projection was performed before further analysis.

Morphological images were acquired in all 4 channels. (**Table 3**).

Software. Images were captured by the Nikon CSU-W1 confocal system in combination with

the NIS elements software (RRID:SCR_014329). Image visualization was later performed

using Fiji freeware[51,52]. In-depth image analysis, pre-processing and machine learning for cell

classification were performed using Python programming language[53] in combination with

Anaconda[54] (distribution and package managing software) and Visual Studio Code (code

editor). The packages and versions used for data analysis are shown in **Table 4**.

Image pre-processing (Suppl. Fig. 5B). Note: all image and data analysis scripts are available

on GitHub. See further the data availability statement.

Cell/nuclear segmentation. Using a tailor-made script implementing the Cellpose[55] (cell

segmentation) or StarDist[56] (nuclear segmentation) package, images were pre-processed by

normalizing the intensity values of each channel between the 1st and 99th quantile. Individual

channels or channel combinations for segmentation can be selected depending on the desired

outcome mask. Resulting outputs included the segmentation mask alongside the quality control

images. For Stardist implementation, hyperparameter probability was set at 0,6 and overlap at

0,3. For Cellpose segmentation, 4 models (cyto2) were averaged to obtain the final mask.

Segmentation was performed on the composite of all CP channels. An estimation of the cell's

diameter could be included to optimize cell detection. Cell segmentation was performed using

Cellpose and used in all cases where the whole-cell crop was given as input to the CNN or data

from the whole cell was used for feature extraction for RF (Fig. 1-4). Nuclear segmentation

was performed using Stardist and used for nuclear and nucleocentric crops (Fig. 4-7).

Ground truth alignment. Following sequential staining and imaging rounds, multiple images

were captured representing the same cell with different markers. Lifting the plate of the

microscope stage and imaging in sequential rounds after several days results in small linear

translations in the exact location of each image. These linear translations need to be corrected

to align morphological with ground truth image data within the same ROI. All images were

aligned using Fourier-based image cross correlation on the intensity-normalized multichannel

images. The alignment shift between image1 and image2 was determined using scikit-image

phase cross correlation. Image2 was then moved according to the predetermined shift to align

morphological with ground truth images.

Ground truth phenotyping. The true cell phenotype was determined by the fluorescence

intensity of the post-hoc immunostaining with class-specific markers or pre-labelling with

Edu/Brdu. In this latter case, the base analogues were incorporated into each cell line prior to

mixing them, i.e. when they were still growing in monoculture so they could be labelled and

identified after co-seeding and morphological profiling. Each ground truth image was imported

alongside the corresponding cell mask for that image. For each cell label, the fluorescence

intensity was determined and tabulated. The threshold was set manually based on the

fluorescence intensity of the monoculture controls. Ground truth labels were assigned to each

region of interest (ROI).

Feature extraction. Handcrafted features were extracted using the scikit-image package (regionprops and GLCM functions). The definition of each feature extracted from the image is listed in **table 5**. All features were extracted for each channel in the cell painting image and for every region (cell, cytoplasm and nucleus) within the ROI.

Data filtering. Over the entire pipeline, ROIs could be discarded based on 3 conditions: (1) Cell detachment. Due to the harsh sample preparation and repeated washing steps, cells could detach from the imaging substrate thus resulting in lack of ground truth information for those cells. As a result, all ROIs for which there was no DAPI signal detected in the ground truth image, were removed from the dataset as incomplete. (2) Cells for which the ground truth fluorescence intensity was ambiguous. Ground truth labels were determined based on the presence of specific IF staining in either of the phenotype classes. If no class-specific IF staining was detected by thresholding, no true label could be assigned. These ROIs were therefore discarded due to uncertainty. (3) Likelihood of faulty ROI detection. The DAPI signal was used to discard ROIs that do not represent (complete) cells by thresholding on minimal DAPI intensity (mean nuclear DAPI intensity > 500) and minimal nuclear size (nuclear area > 160).

ROI classification. *Train-validation-test split.* Model training requires splitting the available data in 3 subsets: training (60%), validation (10%), and testing (30%). The training dataset was used to train the machine learning models (either RF or CNN). The validation dataset, composing of 10% of the total data, was used to determine the hyperparameters and intermediate testing. The remaining 30% of the dataset was kept apart and used to test the final model when training is completed. For both RF and CNN, the testing dataset was never shown to the model during the training phase, but only used after training to determine the accuracy of predictions to the ground truth. The number of instances for each class was equalized by sampling equal number of instances from each predicted class. To account for variation

between technical replicates, the train-validation-test split was stratified per well. As a result,

no datapoints arising from the same well could appear simultaneously in the training, validation

and testing subset. This data stratification was repeated 3 times with different random seeds

(see Methods – Reproducibility).

Random Forest. For each ROI, a set of manually defined parameters was extracted

corresponding to cell shape (area, convex area, filled area, length minor axis, length major axis,

centroid, eccentricity, equivalent diameter area, ferret diameter max, orientation, perimeter,

solidity), texture (GLCM: contrast, dissimilarity, homogeneity, energy, correlation, ASM) and

intensity (maximal intensity, minimal intensity, mean intensity, standard deviation intensity).

This was done for 3 regions (nucleus, cytoplasm and complete cell), and for all channel

dimensions. Redundant parameters could be removed if above a 0.95 correlation threshold. All

parameters were standardized per ROI, grouped per replicate. The number of trees within the

forest was varied between 10 and 150, reaching maximum accuracy at around 30 trees.

Uniform Manifold Approximation and Projection. Dimensionality reduction using UMAP was

performed using either the same feature matrix as used for RF prediction or the feature

embeddings from the trained CNN classification network. Hyperparameters were set at the

default settings.

Convolutional neural network. A ResNet50 model was trained for image classification. In

contrast to classical machine learning techniques, no handcrafted features were extracted.

Crops were defined based on the segmentation mask for each ROI. The bounding box was

cropped out of the original image with a fixed patch size (60µm for whole cells, 18µm for

nucleus and nucleocentric crops) surrounding the centroid of the segmentation mask. For the

whole cell and nuclear crops, all pixels outside of the segmentation mask were set to zero. This

was not the case for the nucleocentric crops. Each ROI was cropped out of the original

morphological image and associated with metadata corresponding to its ground truth label.

Images alongside their labels were fed to the network. Tensors were normalized per channel. Models are trained on a minimum of 5000 training inputs of each class for 50 epochs (training iterations). Each batch consisted of 100 samples. The training input was augmented by linear transformations (horizontal and vertical flip, random rotation). Each epoch, the current model was tested against a validation dataset (**Suppl. Fig. 5C**). The performance of the model on this validation subset determined whether the model was stored as new best (if the new accuracy exceeded the accuracy of the previous best model) or discarded. The learning rate at the start was set at 0,0001 and automatically reduced with a factor of 0,1 during training when no improvement was seen after 10 epochs. After 50 epochs, the best resulting model was tested on a separate test dataset to determine the accuracy on previously unseen data.

GradCAM. GradCAM analysis was used to visualize the regions used by the CNN for classification. This map is specific to each cell. Images are selected randomly out the full dataset for visualization. To avoid cherry-picking, a set of GradCam maps is reported alongside the random seed used for image selection.

Reproducibility. Each model training was performed 3 independent times (model initializations, repetitions are indicated with 'N' on the figures). This was repeated for 3 different random seeds. Each model received input data arising from a minimum of 16 images per well, at least 15 technical replicates (wells). The optimization experiments (figures 1-4) were performed with cell lines with limited variability. These models were trained on 3 independent experiments where ground truth pre-labelling (Edu/BrdU) was performed at least once on either of the cell lines in coculture. For iPSC-derived cultures, as variability is inherent to these differentiations, 3 biological replicates (independent differentiations) were pooled for model training.

Statistics. All statistical comparisons were made nonparametric using Mann-Whitney U (for two independent sample comparison) or Kruskal-Wallis (for multiple sample comparison) with

pairwise tests using Tukey's honestly significant difference test. We opted for nonparametric testing because the number of models in each group to be compared was < 15. Significance levels are indicated on the figures using ns. (no statistical significance, p-value above 0,05), * (p-value between 0,05 and 5e-4), ** (p-value between 5e-4 and 5e-6) and *** (p-value smaller than 5e-6).

Funding

This work was funded by Fonds Wetenschappelijk Onderzoek Vlaanderen (1SB7423N; 1274822N; G033322N), BOF (FFB210009) and IOF UAntwerpen (FFI210239; FFI230099) and VLAIO (HBC.2023.0155).

Acknowledgements

We thank Marlies Verschuuren and Hugo Steenberghen for their assistance and knowledge regarding the cell lines. By extension, we would like to thank all current and former members of the De Vos lab.

Data availability statement

The authors report that the results of this study are available within the manuscript and supplementary materials. All image analysis scripts are open-source available on GitHub (https://github.com/DeVosLab/Nucleocentric-Profiling)

Conflict of interest

The authors declare no conflict of interest.

Abbreviations

BrdU: Bromodeoxyuridine; CNN: convolutional neural network; CP: cell painting; DAPI: 4',6-

diamidino-2-fenylindool; DIV: days in vitro; EdU: 5-ethynyl-2'-deoxyuridine; ER:

endoplasmic reticulum; Grad-CAM: Gradient-weighted Class Activation Mapping; IF:

immunofluorescence; iPSC: induced pluripotent stem cell; NPC: neural progenitor cell; RF:

random forest; ROI: region of interest; TUBB3: beta-III-tubulin; UMAP: Uniform Manifold

Approximation and Projection

Author contributions

The experiments were conceptualized by SDB, TVDL, JVDD, PP and WDV. Experiments

were executed by SDB and JVDD. TVDL and SDB optimized the data analysis scripts. SDB

analysed the imaging data and performed the data analysis. SDB and WDV prepared the

original draft. PP and WDV supervised the work. All authors took part in reviewing and editing

the manuscript.

References

1.   De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164**, 603–615 (2016).

2.   Bell, S. *et al.* Differentiation of Human Induced Pluripotent Stem Cells (iPSCs) into an Effective Model of Forebrain Neural Progenitor Cells and Mature Neurons. *BIO-PROTOCOL* **9**, (2019).

3.   Neyrinck, K. *et al.* SOX9-induced Generation of Functional Astrocytes Supporting Neuronal Maturation in an All-human System. *Stem Cell Rev and Rep* **17**, 1855–1873 (2021).

4.   Haenseler, W. *et al.* A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports* **8**, 1727–1742 (2017).

5.   Nevin, Z. S. *et al.* Modeling the Mutational and Phenotypic Landscapes of Pelizaeus-Merzbacher Disease with Human iPSC-Derived Oligodendrocytes. *The American Journal of Human Genetics* **100**, 617–634 (2017).

6.   Abutaleb, N. O. & Truskey, G. A. Differentiation and characterization of human iPSC-derived vascular endothelial cells under physiological shear stress. *STAR Protocols* **2**, 100394 (2021).

7.   Paik, D. T., Chandy, M. & Wu, J. C. Patient and Disease–Specific Induced Pluripotent Stem Cells for Discovery of Personalized Cardiovascular Drugs and Therapeutics. *Pharmacol Rev* **72**, 320–342 (2020).

8.   Lopez-Gonzalez, R. *et al.* Poly(GR) in C9ORF72 -Related ALS/FTD Compromises Mitochondrial Function and Increases Oxidative Stress and DNA Damage in iPSC-Derived Motor Neurons. *Neuron* **92**, 383–391 (2016).

9.   Amin, N. D. & Paşca, S. P. Building Models of Brain Disorders with Three-Dimensional Organoids. *Neuron* **100**, 389–405 (2018).

10.   Rouhani, F. *et al.* Genetic Background Drives Transcriptional Variation in Human Induced Pluripotent Stem Cells. *PLoS Genet* **10**, e1004432 (2014).

11.   Salomonis, N. *et al.* Integrated Genomic Analysis of Diverse Induced Pluripotent Stem Cells from the Progenitor Cell Biology Consortium. *Stem Cell Reports* **7**, 110–125 (2016).

12.   Yamamoto, T., Arita, M., Kuroda, H., Suzuki, T. & Kawamata, S. Improving the differentiation potential of pluripotent stem cells by optimizing culture conditions. *Sci Rep* **12**, 14147 (2022).

13.   Kuijlaars, J. *et al.* Sustained synchronized neuronal network activity in a human astrocyte co-culture system. *Sci Rep* **6**, 36529 (2016).

14.   Hernández, D. *et al.* Culture Variabilities of Human iPSC-Derived Cerebral Organoids Are a Major Issue for the Modelling of Phenotypes Observed in Alzheimer's Disease. *Stem Cell Rev and Rep* **18**, 718–731 (2022).

15.   D'Antonio, M. *et al.* High-Throughput and Cost-Effective Characterization of Induced Pluripotent Stem Cells. *Stem Cell Reports* **8**, 1101–1111 (2017).

16.   Sullivan, S. *et al.* Quality Control Guidelines for Clinical-Grade Human Induced Pluripotent Stem Cell Lines. *Regen. Med.* **13**, 859–866 (2018).

17.   Chen, C. X.-Q. *et al. Standardized Quality Control Workflow to Evaluate the Reproducibility and Differentiation Potential of Human iPSCs into Neurons.* http://biorxiv.org/lookup/doi/10.1101/2021.01.13.426620 (2021) doi:10.1101/2021.01.13.426620.

18.   Gustafsdottir, S. M. *et al.* Multiplex Cytological Profiling Assay to Measure Diverse Cellular
States. *PLoS ONE* **8**, e80999 (2013).

19.   Bray, M.-A. *et al.* Cell Painting, a high-content image-based assay for morphological profiling
using multiplexed fluorescent dyes. *Nat Protoc* **11**, 1757–1774 (2016).

20.   Cimini, B. A. *et al.* Optimizing the Cell Painting assay for image-based profiling. *Nat Protoc* **18**,
1981–2013 (2023).

21.   Way, G. P. *et al.* Predicting cell health phenotypes using image-based morphology profiling.
MBoC **32**, 995–1005 (2021).

22.   Schiff, L. *et al.* Integrating deep learning and unbiased automated high-content screening to
identify complex disease signatures in human fibroblasts. *Nat Commun* **13**, 1590 (2022).

23.   Caicedo, J. C. *et al.* Cell Painting predicts impact of lung cancer variants. *MBoC* **33**, ar49
(2022).

24.   Tegtmeyer, M. *et al.* High-dimensional phenotyping to define the genetic basis of cellular
morphology. *Nat Commun* **15**, 347 (2024).

25.   Rajaram, S., Pavie, B., Wu, L. F. & Altschuler, S. J. PhenoRipper: software for rapidly profiling
microscopy images. *Nat Methods* **9**, 635–637 (2012).

26.   Uhlmann, V., Singh, S. & Carpenter, A. E. CP-CHARM: segmentation-free image classification
made accessible. *BMC Bioinformatics* **17**, 51 (2016).

27.   Caicedo, J. C. *et al.* Data-analysis strategies for image-based cell profiling. *Nat Methods* **14**,
849–863 (2017).

28.   McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for
Dimension Reduction. (2018) doi:10.48550/ARXIV.1802.03426.

29.   Nguyen, T.-T., Huang, J. Z. & Nguyen, T. T. Unbiased Feature Selection in Learning Random
Forests for High-Dimensional Data. *The Scientific World Journal* **2015**, 1–18 (2015).

30.   He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition. Preprint
at http://arxiv.org/abs/1512.03385 (2015).

31.   Selvaraju, R. R. *et al.* Grad-CAM: Visual Explanations from Deep Networks via Gradient-Based
Localization. *Int J Comput Vis* **128**, 336–359 (2020).

32.   Radtke, A. J. *et al.* IBEX: an iterative immunolabeling and chemical bleaching method for high-
content imaging of diverse tissues. *Nat Protoc* **17**, 378–401 (2022).

33.   Volpato, V. & Webber, C. Addressing variability in iPSC-derived models of human disease:
guidelines to promote reproducibility. *Disease Models & Mechanisms* **13**, dmm042317 (2020).

34.   Shi, Y., Kirwan, P. & Livesey, F. J. Directed differentiation of human pluripotent stem cells to
cerebral cortex neurons and neural networks. *Nat Protoc* **7**, 1836–1846 (2012).

35.   Dincer, A. B., Celik, S., Hiranuma, N. & Lee, S.-I. *DeepProfile: Deep Learning of Cancer
Molecular Profiles for Precision Medicine.* http://biorxiv.org/lookup/doi/10.1101/278739 (2018)
doi:10.1101/278739.

36.   Weigert, M. *et al.* Content-aware image restoration: pushing the limits of fluorescence
microscopy. *Nat Methods* **15**, 1090–1097 (2018).

37. De Vos, W. H. *et al.* Increased plasticity of the nuclear envelope and hypermobility of telomeres
due to the loss of A–type lamins. *Biochimica et Biophysica Acta (BBA) - General Subjects* **1800**, 448–
458 (2010).

38. Heckenbach, I. *et al.* Nuclear morphology is a deep learning biomarker of cellular senescence.
Nat Aging **2**, 742–755 (2022).

39. Eastwood, M. *et al.* MesoGraph: Automatic profiling of mesothelioma subtypes from histological
images. *Cell Reports Medicine* **4**, 101226 (2023).

40. Mikolajczyk, A. & Grochowski, M. Data augmentation for improving deep learning in image
classification problem. in *2018 International Interdisciplinary PhD Workshop (IIPhDW)* 117–122 (IEEE,
Swinoujście, 2018). doi:10.1109/IIPHDW.2018.8388338.

41. Nguyen, L. D., Lin, D., Lin, Z. & Cao, J. Deep CNNs for microscopic image classification by
exploiting transfer learning and feature concatenation. in *2018 IEEE International Symposium on*
Circuits and Systems (ISCAS) 1–5 (IEEE, Florence, 2018). doi:10.1109/ISCAS.2018.8351550.

42. Tomov, M. L. *et al.* Resolving cell state in iPSC-derived human neural samples with multiplexed
fluorescence imaging. *Commun Biol* **4**, 786 (2021).

43. Strano, A., Tuck, E., Stubbs, V. E. & Livesey, F. J. Variable Outcomes in Neural Differentiation
of Human PSCs Arise from Intrinsic Differences in Developmental Signaling Pathways. *Cell Reports*
31, 107732 (2020).

44. Galiakberova, A. A. *et al.* Different iPSC-derived neural stem cells shows various spectrums of
spontaneous differentiation during long term cultivation. *Front. Mol. Neurosci.* **16**, 1037902 (2023).

45. Du, J. & Xu, Y. Hierarchical deep neural network for multivariate regression. *Pattern*
Recognition **63**, 149–157 (2017).

46. Shakarchy, A. *et al.* Machine learning inference of continuous single-cell state transitions during
myoblast differentiation and fusion. *Mol Syst Biol* (2024) doi:10.1038/s44320-024-00010-3.

47. Mai, M. *et al.* Morphology-based deep learning approach for predicting adipogenic and
osteogenic differentiation of human mesenchymal stem cells (hMSCs). *Front. Cell Dev. Biol.* **11**,
1329840 (2023).

48. Gosselin, D. *et al.* An environment-dependent transcriptional network specifies human
microglia identity. *Science* **356**, eaal3222 (2017).

49. Engle, S. J., Blaha, L. & Kleiman, R. J. Best Practices for Translational Disease Modeling Using
Human iPSC-Derived Neurons. *Neuron* **100**, 783–797 (2018).

50. Chambers, S. M. *et al.* Highly efficient neural conversion of human ES and iPS cells by dual
inhibition of SMAD signaling. *Nat Biotechnol* **27**, 275–280 (2009).

51. Rueden, C. T. *et al.* ImageJ2: ImageJ for the next generation of scientific image data. *BMC*
Bioinformatics **18**, 529 (2017).

52. Schindelin, J., Rueden, C. T., Hiner, M. C. & Eliceiri, K. W. The ImageJ ecosystem: An open
platform for biomedical image analysis. *Molecular Reproduction Devel* **82**, 518–529 (2015).

53. Van Rossum, G., & Drake, F. L. Python 3 Reference Manual. (2009).

54. Anaconda Software Distribution. Anaconda Documentation. Anaconda Inc.

55. Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for
cellular segmentation. *Nat Methods* **18**, 100–106 (2021).

56.   Weigert, M., Schmidt, U., Haase, R., Sugawara, K. & Myers, G. Star-convex Polyhedra for 3D
Object Detection and Segmentation in Microscopy. in *2020 IEEE Winter Conference on Applications of*
Computer   Vision   (WACV)   3655–3662   (IEEE,   Snowmass   Village,   CO,   USA,   2020).
doi:10.1109/WACV45572.2020.9093435.

Tables

Table 1 | Specifications of morphological staining composition.

Staining solution 1							
Dye	**Target**	**Excitation (nm)**	**Emission (nm)**	**Stock concentration**	**End dilution**	**Supplier**	**Catalog #**
DAPI	Nucleus	350	470	2.5 mg/ml	1/500	Sigma Aldrich	D9542
Concanavalin	ER	480	490-540	5 mg/ml	1/300	Life Technologies	C11252
WGA	Golgi	550	560-570	1 mg/ml	1/600	Life Technologies	W32464
Phalloidin	Actin	570	580-610	66μM	1/1000	Life Technologies	A12380
Mitotracker	Mitochondria	640	650-660	1 mM	1/1000	Life Technologies	M22426
Staining solution 2							
Syto14	Nucleoli /RNA	525	540-580	5 mM	1/1000	Life Technologies	S7576

Table 2 | Used antibodies.

Antibody		Host	Supplier	Catalog #
Primary antibodies	Anti-BrdU	Sheep	Abcam	ab1893
	Anti-MAP2	Chicken	Synaptic systems	188006
	Anti-TUBB3	Mouse	BioLegend	801202
	Anti-CD44	Mouse	Millipore	MAB4065
	Anti-GFAP	Goat	Abcam	ab53554
	Anti-Iba1	Rabbit	Wako	019-19741
	Anti-Ki67	Rabbit	Abcam	ab15580
Secondary antibodies	Anti-mouse-Cy3	Donkey	Jackson ImmunoResearch	715-165-150
	Anti-mouse-Cy5	Donkey	Jackson ImmunoResearch	715-175-150
	Anti-sheep-Cy3	Donkey	Jackson ImmunoResearch	713-165-147
	Anti-sheep-Cy5	Donkey	Jackson ImmunoResearch	713-175-147
	Anti-chicken-Cy5	Donkey	Jackson ImmunoResearch	703-175-155
	Anti-goat-Cy3	Donkey	Jackson ImmunoResearch	705-165-147
	Anti-Guinea pig-Cy3	Donkey	Jackson ImmunoResearch	706-165-148
	Anti-Rabbit-Cy5	Donkey	Jackson ImmunoResearch	711-175-152

Table 3 | Specifications of the used laser lines, excitation and emission filters.

Channel dimension	Laser line (nm)	Excitation filter (nm)	Emission filter (nm)
Channel 1	405		B447/60

Channel 2	488	Di01-T405/488/568/647-	B520/35
Channel 3	561	13x15x0.5	B617/73
Channel 4	640		B685/40

Table 4 | Python packages used for image and data analysis alongside the software version.

Package	Version	Reference
pathlib	1.0.1	
numpy	1.21.2	Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., … Oliphant, T. E. (2020). Array programming with NumPy. Nature, 585, 357–362. https://doi.org/10.1038/s41586-020-2649-2
tifffile	2021.7.2	
pytorch	1.13.0	Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., … Chintala, S. (2019). PyTorch: An Imperative Style, High-Performance Deep Learning Library. In Advances in Neural Information Processing Systems 32 (pp. 8024–8035). Curran Associates, Inc.
torchvision	0.14.0	
nd2reader	3.3.0	
matplotlib	3.5.2	Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. Computing in Science & Engineering, 9(3), 90–95.
scikit-image	0.19.3	Van der Walt, S., Sch"onberger, Johannes L, Nunez-Iglesias, J., Boulogne, Franccois, Warner, J. D., Yager, N., … Yu, T. (2014). scikit-image: image processing in Python. PeerJ, 2, e453.

scikit-learn	1.1.3	Pedregosa, F., Varoquaux, Ga"el, Gramfort, A., Michel, V., Thirion, B., Grisel, O., … others. (2011). Scikit-learn: Machine learning in Python. Journal of Machine Learning Research, 12(Oct), 2825–2830.
argparse	1.4.0	
cellpose	2.1.1	Stringer, C., Wang, T., Michaelos, M. et al. Cellpose: a generalist algorithm for cellular segmentation. Nat Methods 18, 100–106 (2021). https://doi.org/10.1038/s41592-020-01018-x
tqdm	4.64.1	
scipy	1.9.3	Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., … SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods, 17, 261–272. https://doi.org/10.1038/s41592-019-0686-2
pandas	1.5.1	McKinney, W., & others. (2010). Data structures for statistical computing in python. In Proceedings of the 9th Python in Science Conference (Vol. 445, pp. 51–56).
seaborn	0.12.1	https://doi.org/10.5281/zenodo.883859
imbalanced-learn	0.10.0	Imbalanced-learn: a python toolbox to tackle the curse of imbalanced datasets in machine learning
StarDist	0.8.5	M. Weigert, U. Schmidt, R. Haase, K. Sugawara and G. Myers, "Star-convex Polyhedra for 3D Object Detection and Segmentation in Microscopy," 2020 IEEE Winter Conference on Applications of Computer Vision (WACV), Snowmass, CO, USA, 2020, pp. 3655-3662, doi: 10.1109/WACV45572.2020.9093435.

Table 5 | Definition of handcrafted features (according to the scikit-image documentation). All of these features were extracted for each fluorescent channel (1-4) and region (nucleus, cytoplasm and whole-cell) in the cell painting images.

Feature name	Description	Category
Intensity max	Maximal pixel intensity inside the ROI	Intensity
Intensity mean	Mean pixel intensity inside the ROI	Intensity
Intensity min	Minimal pixel intensity inside the ROI	Intensity
Intensity Std	Standard deviation of the pixel intensities inside the ROI	Intensity
Area	Area of the ROI	Shape
Area convex	The area of the convex polygon that encloses the ROI	Shape
Area filled	Area of the ROI including all holes inside the ROI	Shape
Axis major length	The length of the major axis of the ellipse that has the same normalized second central moments as the region.	Shape
Axis minor length	The length of the minor axis of the ellipse that has the same normalized second central moments as the region.	Shape
Centroid	Coordinates of the center of the ROI	Shape
Eccentricity	Eccentricity of the ellipse that has the same second-moments as the ROI.	Shape
Equivalent diameter area	The diameter of a circle with the same area as the ROI.	Shape

Extent	The ratio of the Area of the ROI to the Area of the bounding box around the ROI.	Shape
Feret diameter max	The longest distance between points around a ROIs' convex polygon.	Shape
Orientation	The angle between the X-axis and the major axis.	Shape
Perimeter	Length of the contour of the ROI	Shape
Perimeter crofton	Perimeter of the ROI approximated by the Crofton formula in 4 directions.	Shape
Solidity	The ratio of the number of pixels inside the ROI to the number of pixels of the convex polygon.	Shape
Contrast	The difference between the maximum intensity and minimum intensity pixel inside the ROI.	Texture
Dissimilarity	The average difference in pixel intensity between neighbouring pixels.	Texture
Homogeneity	Value for similarity of pixels in the ROI.	Texture
Energy	Value for the local change of pixel intensities in the ROI.	Texture
Correlation	Value for the linear dependency of pixels in the ROI.	Texture
ASM (angular second moment)	Value for the uniformity of pixel values in a ROI.	Texture

Figure 1 | Shallow classification using manually defined features. (A) Image overview of 1321N1 (top) and SH-SY5Y (bottom) cells after CP staining acquired at higher magnification (Plan Apo VC 60xA WI DIC N2, NA = 1,20). Scale bar is 30µm. The channel, wavelength and dye combination used is listed in the table below the figure. This color code and channel numbering is used consistently across all figures. (B) UMAP dimensionality reduction using handcrafted features. Each dot represents a single cell. The color reflects either cell type condition (left) or replicate (right). This shows UMAP clustering is a result of cell type differences and not variability between replicates. (C) Feature importance deducted from the UMAP (feature maps). Each dot represents a single cell. Three exemplary feature maps are highlighted alongside the quantification per cell type. These feature representations help understanding the morphological features that underly the cluster separation in UMAP. (D) Random Forest classification performance on the manually defined feature dataframe with and without exclusion of redundant features. Average confusion matrix (with redundant features) and Mean Decrease in Impurity (reflecting how often this feature is used in decision tree splits across all random forest trees). All features used in the UMAP are used for RF building. Each dot in the violinplot represents the F-score of one classifier (model initialization, N = 30). Classifiers were trained 10x with 3 different random seeds.

Figure 2 | Convolutional neural network classification of monoculture conditions. (A) Schematic of image pre-processing for CNN classification. (B) CNN accuracy and confusion matrix on cell type classification in monocultures. Each dot in the violinplot represents the F-score of one classifier (model initializations, N = 9). (C) Shallow vs. deep learning accuracy for a varying number of input instances (ROIs). Each dot in the violinplot represents the F-score of one classifier (model initialization, N). The ribbon represents the standard deviation on the classifiers. (D) The impact of experimental variability in the training dataset on model performance for shallow vs. deep learning. Classifiers were trained on either 1 (single rep.) or multiple (mixed reps.) replicates (where each replicate consists of N new biological experiment). Mann-Whitney-U-test, p-values resp. 0.7304 and 0.000197. Each violinplot represents the F-score of one classifiers (model initializations). (E) The performance of deep learning classifiers trained in panel D on single replicates (low variability) or mixed replicates (high variability) on unseen images from either the training replicate (cross-validation) or an independent replicate (independent testing) (where each replicate consists of a new biological experiment). Kruskal-Wallis test on single training condition, p-value of 0,026 with post-hoc Tukey test. Mann-Whitney-U-test on mixed training, p-value of 7,47e-6. Each violinplot represents the F-score of one classifier (model initialization, N = 3). (F) Images of example inputs given to the CNN. The composite image contains an overlay of all CP channels (left). The GradCAM image overlays the GradCAM heatmap on top of the composite image, highlighting the most important regions according to the CNN (right). One example is given per cell type.

Figure 3 | Required input quality for cell type classification. (A) Performance of the CNN when only a selection of image channels is given as training input. Each boxplot represents the F-score of one classifier (model initialization, N = 3). Channel numbering is in accordance with fig. 1A. (1 = DAPI; 2 = FITC; 3 = Cy3; 4 = Cy5) (B) Simulation of the effect of increased pixel size (reduced spatial resolution) on the classification performance. Each point is the average of 3 CNN model initializations (N), with bars indicating the standard deviation between models. Red line indicates the average F-score of the original crops. (C) Simulation of the effect of added gaussian noise (reduced signal-to-noise ratio) on the classification performance. Each point is the average of 3 CNN model initializations, with bars indicating the standard deviation between models. Red line indicates the average F-score of the original images. Statistics were performed using a Kruskal-Wallis test with post-hoc Tukey test. (D) CNN performance on 1321N1, RPE1 and ARPE cells. Dots represent different model initializations (N = 3).

Figure 4 | Required regional input for accurate CNN training in cultures of high density. (A) Selected images and insets for increasing culture density with the density categories used for CNN training. (B) Results of 3 CNN models trained using different regional input (the full cell, only nucleus or nucleocentric area) and evaluated on data subsets with increasing density. Each dot represents the F-score of one classifier (model initialization, N = 3) tested on a density subset. Classifiers were trained with the same random seed. Ribbon represents the standard deviation. (C) CNN performance (F-score) of CNN models using different regional inputs (full cell, only nucleus or nucleocentric area). Each boxplot represents 3 model initializations for 3 different random seeds (N = 9). (D) Images of example inputs for both the nuclear and nucleocentric region. The composite image contains an overlay of all CP channels (top). The GradCAM image overlays the GradCAM heatmap on top of the composite image, highlighting the most important regions according to the CNN (bottom). One example is given per cell type. (E) Systematic in- and decrease (default of 18µm used in previous panels) of the patch size surrounding the nuclear centroid used to determine the nucleocentric area. Each dot represents the results of one classifier (model initialization, N = 3). Ribbon represents the standard deviation. The analysis was performed using a mixed culture dataset of 1321N1 and SH-SY5Y cells (Fig. 5).

Figure 5 | Cell type prediction in mixed cultures by post-hoc ground truth identification. (A) Schematic overview of virtual vs. physically mixed cultures and the subsequent CNN models. For virtual mixing, individual cell types arise from distinct cultures, the cell type label was assigned based on the culture condition. For physical mixing, the true phenotype of each crop was determined by intensity thresholding after post-hoc staining. Three model types are defined: Mono2Mono (culture-based label), Co2Co (cell-based label) and Mono2Co (trained with culture-based label and tested on cell-based label). (B) Determining ground-truth phenotype by intensity thresholding on IF (top) or pre-label (bottom). The threshold was determined on the intensity of monocultures. Only the pre-labelled ground-truth resulted in good cell type separation by thresholding for 1321N1 vs. SH-SY5Y cells. (C) Mono2Mono (culture-based ground truth) vs. Co2Co (cell-based ground truth) models for cell type classification. Analysis performed with full cell segmentation. Mann-Whitney-U-test p-value 0,0027. Monoculture-trained models were tested on mixed cultures. Pretrained models were trained on independent biological replicates. These could be finetuned by additional training on monoculture images from the same replicate as the coculture. This was shown to reduce the variation between model iterations (Median performance: Mann-Whitney-U-test, p-value 0,0015; Coefficient of variation: Mann-Whitney-U-test, p-value 3,48e-4). Each dot in the violinplots represents the F-score of one classifier (model initialization, N = 9). Classifiers were trained 3x with 3 different random seeds.

Figure 6 | iPSC-derived differentiation staging using morphology profiling. (A) Schematic overview of guided vs. spontaneous neural differentiation. DIV = days in vitro. Selected timepoints for analysis of the spontaneously differentiation culture were 13, 30, 60 and 90 days from the start of differentiation of iPSCs. (B) Representative images of morphological staining (color code as defined in fig. 1A) and post-hoc IF of primed and differentiated iPSC-derived cultures (guided differentiation). Ground-truth determination is performed using TUBB3 (for mature neurons) and Ki67 (mitotic progenitors). (C) Fraction of neurons vs. NPC cells in the primed vs. differentiated condition as determined by IF staining. Upon guided differentiation, the fraction of neurons increased. (D) left: CNN performance when classifying neurons (Ki67-/TUBB3+) vs. NPC (Ki67+/TUBB3-) cells using either a condition-based or cell-type based ground truth. Each dot in the violinplots represents the F-score of one classifier (model initialization). Classifiers were trained with different random seeds. Mann-Whitney-U-test, p-value 4,04e-4. Right: comparison of CNN vs. RF performance. Mann-Whitney-U-test, p-value 2,78e-4. (E) Fractional abundance of predicted cell phenotypes (NPC vs. neurons) in primed vs. differentiated culture conditions using the cell-based CNN. (F) Unsupervised and supervised UMAP of the cell-based CNN feature embeddings. Plot color coded by cell type. Points represent individual cells. (G) Representative images of spontaneously differentiating neural cultures. Color code as defined in fig. 1A. (H) Prediction of differentiation status using the cell-based CNN model trained on guided differentiated culture.

Figure 7 | iPSC cell type identification using morphology profiling. (A) Representative images of iPSC-derived neurons, astrocytes and microglia in monoculture with morphological staining. Color code as defined in fig. 1A. Prediction accuracy of a CNN trained to classify monocultures of iPSC-derived astrocytes, microglia and neurons with confusion matrix (average of all models). Each dot in the boxplot represents the F-score of one classifier (model initialization, N = 9). Classifiers were trained 3x with 3 different random seeds. (B) Representative images of monocultures of iPSC-derived neurons and microglia treated with LPS or control. Color code as defined in fig. 1A. Prediction accuracy and confusion matrix (average of all models) are given. Each dot in the violinplot represents the F-score of one classifier (model initialization, N = 3). (C) Representative images of a mixed culture of iPSC-derived microglia and neurons. Ground-truth identification was performed using IF. Each dot in the violinplot represents the F-score of one classifier (model initialization). Classifiers were trained 3 different random seeds. Results of the CNN are compared to shallow learning (RF). The same analysis was performed for mixed cultures of neurons and microglia with LPS treatment or control. A layered approach was used where first the neurons were separated from the microglia before classifying treated vs. non-treated microglia. Each dot in the violinplot represents the F-score of one classifier (model initialization, N = 9). Classifiers were trained 3x with 3 different random seeds.

Supplementary figure 1 (A) PCA dimensionality reduction on handcrafted features extracted from monocultures of 1321N1 and SH-SY5Y cells. Each point represents an individual cell. (B) UMAP of feature embeddings of the CNN trained to classify 1321N1 and SH-SY5Y monocultures. Each point represents an individual cell. (C) Examples of misclassified ROIs. (D) Left: Representative images of 1321N1 cells with increasing density alongside their cell and nuclear mask produced using resp. Cellpose and Stardist. Images are numbered from 1-5 with increasing density. Upper right: The number of ROIs detected in comparison to the ground truth (manual segmentation). A ROI was considered undetected when the intersection over union (IoU) was below 0,15. Each bar refers to the image number on the left. The IoU quantifies the overlap between ground truth (manually segmented ROI) and the ROI detected by the segmentation algorithm. It is defined as the area of the overlapping region over the total area. IoU for increasing cell density for cell and nuclear masks is given in the bottom right. Each point represents an individual ROI. Each bar refers to the image number on the left. (E) Examples of segmentation mistakes made by the Stardist segmentation algorithm for nuclear segmentation for different culture densities. (F) Nuclear area in function of density. (G) Definition of cell regions given as training input for nuclear and nucleocentric model training.

Supplementary figure 2 | Influence of the nuclear/cellular size on CNN prediction and association with the input background (space artificially set to zero outside of the segmentation mask). (A) Quantification of average nuclear/cellular size per cell type. (B) GradCAM images for 10 random seeds for crops and CNN models trained with background either set to zero or 'random speckle'. (C) CNN prediction results for models trained on crops with background either set to zero or 'random speckle'. Each dot in the violinplots represents the F-score of one classifier (model initialization, N = 3). (D) Feature map (see UMAP in figure 1 B and C) of 1321N1 and SH-SY5Y cells showing the contribution of nuclear/cellular area to the cell type cluster separation. Each point represents an individual cell.

Density: <20% confluency

Density: 20-40% confluency

Density: 40-60% confluency

Density: 60-80% confluency

Density: 80-95% confluency

Density: >95% confluency

Supplementary figure 3 | GradCAM maps per region and density.

Supplementary figure 4 | Fluorescence quenching over time using LiBH4. (A) Images before and after quenching for all 4 fluorescence channels. (B) Time curve of normalized image-level fluorescence intensity during incubation with 1mg/ml LiBH4.

Supplementary figure 5 | Methods. (A) Pipeline used for morphological profiling in mixed cultures. (B) Overview of the steps within the image analysis pipeline. (C) Evaluation of accuracy, true negative and true positive rate during CNN training (1321N1 vs. SH-SY5Y in monocultures) across all 50 epochs.