Unbiased identification of cell identity in dense mixed neural cultures.

Sarah De Beuckeleer[1], Tim Van De Looverbosch[1], Johanna Van Den Daele[1], Peter Ponsaerts[2],

and Winnok H. De Vos[1,3,4]

[1] Laboratory of Cell Biology and Histology, University of Antwerp

[2] Laboratory of Experimental Hematology, Vaccine and Infectious Disease Institute

(Vaxinfectio), University of Antwerp

[3] Antwerp Centre for Advanced Microscopy, University of Antwerp

[4] μNeuro Research Centre of Excellence, University of Antwerp

e-mail:

sarah.debeuckeleer@uantwerpen.be

tim.vandelooverbosch@uantwerpen.be

johanna.vandendaele@uantwerpen.be

peter.ponsaerts@uantwerpen.be

winnok.devos@uantwerpen.be

Contact information *Corresponding author:

University of Antwerp, Laboratory of Cell Biology and Histology

Universiteitsplein 1, 2610 Antwerp, Belgium

e-mail: winnok.devos@uantwerpen.be

Keywords: morphological phenotyping, multiplexed imaging, iPSC culture validation, neural

differentiation, computational biology

Abstract

Induced pluripotent stem cell (iPSC) technology is revolutionizing cell biology. However, the variability between individual iPSC lines and the lack of efficient technology to comprehensively characterize iPSC-derived cell types hinder its adoption in routine screening settings. To facilitate the validation of iPSC-derived cell culture composition, we have implemented an imaging assay based on cell painting and convolutional neural networks to recognize neural cell types in dense and mixed cultures with high fidelity. We have benchmarked our approach using pure and mixed cultures of neuroblastoma and astrocytoma cell lines and attained a classification accuracy above 96%. Through iterative data erosion we found that inputs containing the nuclear region of interest and its close environment, allow achieving equally high classification accuracy as inputs containing the whole cell for semi-confluent cultures and preserved its accuracy even in very dense cultures. We then applied the nucleocentric cell profiling approach to evaluate the differentiation status of iPSC-derived neural cultures, by determining the ratio of postmitotic neurons and neural progenitors. We found that the cell-based prediction significantly outperformed an approach in which the culture time point was used as classification criterion (96% *vs.* 86%, resp.). Thus, nucleocentric morphological single cell profiling provides a means to quantify cell composition in complex mixed neural cultures and holds promise for use in quality control of iPSC-derived neural cell culture models.

Introduction

Modelling the complexity of the human brain and its (dys)function has proven to be notoriously challenging. This is due to its intricate wiring, the cellular heterogeneity and species-specific differences[1]. To increase the translational value of neuroscientific research, there is a need for physiologically relevant human models. With the advent of human induced pluripotent stem cell (iPSC) technology, it has become possible to generate a wealth of brain-resident cell types including neurons[2], astrocytes[3], microglia[4], oligodendrocytes[5] and endothelial cells[6], allowing the study of complex polygenic pathologies that cannot be modelled in animals, opening avenues to precision pharmacology[7,8]. Furthermore, the ability of iPSC to develop into organoids renders them attractive for studying multi-cellular interactions in a 3D context that is closer to the *in vivo* situation[9]. However, genetic drift and batch-to-batch heterogeneity cause variability in reprogramming and differentiation efficiency[10]. This hinders the use of iPSC-derived cell systems in systematic drug screening or cell therapy pipelines. Quality control in such settings demands a robust and cost-effective approach. Current culture validation methods include sequencing, flow cytometry and immunocytochemistry [10,11]. These methods are often low in throughput, costly and/or destructive. Thence, we set out to develop a method for evaluating the composition of such cultures based on high-content imaging, which is fast and affordable. The primary criteria were to facilitate cell type identification with cellular resolution while preserving spatial information and ensuring compatibility with subsequent immunocytochemistry or molecular assays for further biological inquiries. To this end, we employed the Cell Painting (CP)[12,13] method, which is based on labelling cells with simple organic dyes and analysing the resulting phenotypes. CP has proven to be a powerful and generic method for predicting the mode-of-action of pharmacological or genetic perturbations, and this sheerly using a cell morphology readout[14–16]. Thus far, CP has primarily been utilized to predict conditions associated with pharmacological treatments or genetic modifications

using images as input. An exemplary achievement in this context has been realized by the

JUMP consortium, which has published an open-source CP image dataset comprising over

13,000 distinct conditions[17]. Recognizing the predictive potential of morphological

information, we explored its utility at the single-cell level, shifting focus from the image level

to predict individual cell types. By implementing deep learning for cell classification and

tuning the input space, we established an approach that allows recognizing cell types with high

accuracy, even in very dense cultures. We illustrate its potential by using it to distinguish iPSC-

derived cell types in co-culture and to assess the maturity of differentiating neurons.

Results

Neural cell lines have a unique morphotextural fingerprint.

Several groups have demonstrated that morphological cell profiling can be used to discriminate pharmacogenomic perturbations based on phenotypic similarity[14–16]. We asked whether a similar approach could be leveraged to unequivocally distinguish individual cell types as well. To this end, we implemented a version of CP based on 4-channel confocal imaging (**Fig. 1A**) and applied it to monocultures of two neural cell lines from a different lineage, namely astrocyte-derived 1321N1 astrocytoma cells and neural crest-derived SH-SY5Y neuroblastoma cells. First, we explored whether traditional morphotextural feature extraction from cells and their nuclei provided sufficient distinctive power. Representation of the resulting standardized feature set in UMAP space revealed a clear separation of both cell types without replicate bias (**Fig. 1B**). Clustering of instances was less pronounced after principal component analysis (**Suppl. Fig. 4 A**), but UMAP better preserves local and global data structure [18]. When mapping the individual features to the UMAP space, we found that both texture (*e.g.,* energy, homogeneity) and shape (*e.g.,* nuclear area, cellular area) metrics contributed to the separation (**Fig. 1C**). The contribution of intensity-related features (*e.g.,* mean, max, min, std intensity) to cell type separation was less pronounced as they were more correlated with the biological replicate. Thus, we concluded that cell types can be separated across replicates based on a morphotextural fingerprint.

CNN outperforms random forest in cell type classification.

To evaluate whether the morphotextural fingerprint could be used to predict cell type, we performed Random Forest (RF) classification, using different seeds for splitting up training and validation sets. This resulted in a rather poor prediction accuracy of 71,0±1,0%, mainly caused by the significant (46%) misclassification of 1321N1 cells (**Fig. 1D**). The imbalance in

recall and precision was surprising given the clear separation of cell populations in UMAP space using the same feature matrix. When inspecting the main contributions to the RF classifier, we found very similar features as highlighted in UMAP space to add to the discrimination (**Fig. 1C**). Where the top 2 features (nuclear Cy3 energy and cellular area) showed the expected gradient along the UMAP1 direction, the third parameter (cellular DAPI contrast) had no contribution to UMAP separation. Reducing noise by manually removing redundant and correlating features could not ameliorate RF performance, which may be due to the documented bias in feature selection for node splitting in high-dimensional data[19]. This result drove us to improve the classification accuracy by means of a ResNet[20] convolutional neuronal network (CNN). Here, we no longer relied on the extraction of features from segmented cell objects, but instead used image crops centred around individual cells (bounding box around cell centroid) and blanked for their surroundings as input (**Fig. 2A**). Using this approach, we found a significantly higher prediction accuracy of 96,0±1,8%, with a much more balanced recall and precision (**Fig. 2B**). Even with only 100 training instances per class, CNN outperformed RF, but optimal performance was attained with 5000 training instances (**Fig. 2C**). As expected, a model trained on all three biological replicates performed slightly worse (lower accuracy and higher variability) at predicting a test set of a given replicate than a model trained on that same replicate (**Fig. 2D**). However, models trained on two different replicates outperformed those that were only trained on one when used for predicting instances of a third (unseen) replicate. This emphasizes the need for including sufficient variation in the training set (**Fig. 2E**). Although more performant, CNN classification does not allow direct retrieval of discriminative features, which complicates model interpretation. To gain a visual understanding of image information contributing most to the classification we therefore resorted to Gradient-weighted Class Activation Mapping (Grad-CAM)[21]. This revealed that the cell borders (edge information), nuclear and nucleolar signals were the most distinctive features

(**Fig. 2F**). When scrutinizing CNN misclassifications, we found that these are mainly caused

by faulty cell detection (*e.g.,* oversegmentation of cell ramifications), unhealthy/dead cells,

debris or visibly aberrant cell shapes (**Suppl. Fig. 4B**) - mistakes not necessarily attributed to

the CNN. To shed light on the contribution of individual markers to the classification, we

eroded the input to single channel images. For all cases, the prediction performance was below

or equal to 85,0% suggesting that no single channel contains all relevant information (**Fig. 2**

G). Combinations of two or three channels could not match the prediction accuracy of the full

four-channel image either. Thus, we concluded that all channels contribute at least in part to a

successful classification.

Nucleocentric predictions remain accurate regardless of culture density.

Morphological profiling relies on cell identification for adequate class prediction. This may

become difficult in dense cultures such as iPSC-cultures, clustered cells and tissue (mimics).

Having established a method to distinguish cell types with high accuracy, we next asked how

robust the classification was to increasing cell density. To this end, we acquired images of

1321N1 and SH-SY5Y monocultures, grown at densities ranging from 0 to 100% confluency

(**Fig. 3A**). Based on the nuclear count, we binned individual fields into 6 density classes (0-

20%, 20-40%, 40-60%, 60-80%, 80-95%, 95-100%) and trained a CNN with equal sampling

of cell numbers per density class to avoid bias. No decrease in accuracy was observed until

culture density reached 80% confluency. Only for very high densities (95-100%), we found a

significant decrease in the prediction accuracy (92,0±1,7%) (**Fig. 3B**). We reasoned that under

these conditions cell shape would be predominantly determined by neighbouring cells and cell

segmentation performance would decrease (**Suppl. Fig. 1**). Nuclei are less malleable and even

in dense cultures remain largely separated, allowing their robust segmentation and avoiding

CNN misclassifications resulting from segmentation errors. Hence, we asked whether using

the nuclear ROI as input would improve classification performance at high densities despite containing less cellular information. However, regardless of a relatively high average prediction accuracy of about 92,4±1,0% (**Fig. 3C**), the performance was lower than cell ROI across the density range. Thence, we tested an intermediate condition in which nuclear ROI with their direct surroundings (a square bounding box of 60x60 microns surrounding the nuclear centroid coordinates) were considered as input (**Suppl. Fig. 4C**). This resulted in a classification accuracy of 95,0±0,4% (**Fig. 3C**) comparable to that of models trained on whole-cell inputs for low to moderate cell densities, but also maintained prediction performance at high cell densities (**Fig. 3B**). To understand why this was not the case for nuclear crops, we inspected Grad-CAM output for these predictions, and found that the latter diverted the attention of the CNN to the background (**Fig. 3D**). We interpret this result as the CNN using the background as a proxy for nuclear size. As nuclear area dropped with culture density, the dynamic range decreased, which could explain the increased error rate of the CNN for high densities unrelated to segmentation errors (**Suppl. Fig. 4B**). Thus, we conclude that using the nucleocentric region as input for the CNN is a valuable strategy for accurate cell phenotype identification in dense cultures.

Cell prediction remains accurate in mixed cell culture conditions.

Although a very high classification accuracy was obtained with nucleocentric CNN predictions, both training and testing were performed with input images drawn from monocultures. As our goal was to allow cell type prediction in complex, heterogeneous cultures, we next switched to a more faithful situation in which we co-cultured both cell types. Ground-truth for these predictions was generated by either performing post-hoc immunofluorescence (IF) with cell-type specific antibodies (CD44 for 1321N1 and TUBB3 for SH-SY5Y cells), or by differential replication labelling (EdU or BrdU for either of the cell

types) after dye quenching[22] (**Fig. 4A**). The latter proved significantly more successful for

binary ground truth classification (through intensity thresholding) than the former (**Fig. 4B**).

When training a CNN to recognize the cell types using these post-hoc markers as ground truth,

we found that the prediction accuracy on the left-out dataset of the co-culture (Co2Co) was

almost as high as when a model was trained and tested on monocultures (Mono2Mono) (**Fig.**

4C).

We then tested whether it was possible to use a classifier trained on monocultures for cell type

prediction in co-culture (Mono2Co). This resulted in a prediction accuracy of 85,0±4,6%. We

contributed this drop in accuracy to the effect on cell phenotype exerted by the presence of

other cell types in mixed cultures which is not captured in monocultures. As the images from

monocultures and co-cultures were obtained from different plates, we suspected inter-replicate

variability in culture and staining procedures to contribute in part to this lesser performance.

Therefore, we tested whether we could improve the performance of the CNN by including

monocultures from the same plate as the co-cultures. This finetuning indeed improved the

average performance to 89,0±0,7%, and more importantly, it significantly reduced the

variability (coefficient of variation) of the predictions (**Fig. 4D**). Thus, while not yet reaching

the same accuracy, it proves that it is possible to establish a model that recognizes cell types in

co-cultures that is solely trained on monocultures.

Nucleocentric phenotyping can be applied to stage iPSC-derived neuronal cultures.

iPSC-derived neuronal cell cultures suffer from significant inter- and intra-batch variability

and could benefit from an efficient quality control[23,24]. Thence, we applied our nucleocentric

phenotyping pipeline to stage the maturity of a neuronal cell culture based on its cell type

composition. Using a guided protocol[2], two differentiation stages were simulated: a primed

stage, where most cells are cycling neural progenitors (NPCs), and a differentiated stage where

most cells are post-mitotic neurons (**Fig. 5A**). The two cell types were discriminated by post-hoc IF labelling for the cell cycle marker Ki67 (NPC) and the microtubule marker ß-III-tubulin (TUBB3, neurons). Not all cells in the CP image could be assigned with a ground truth label due to cell detachment upon re-staining or sheer absence of the tested marker. Since no single monoculture consists of 100% of either cell type, we applied gates to retain only those cells that show unequivocal staining for either one of both markers. Based on these gates, ROIs were either classified as neuron (Ki67-/TUBB3+), NPC (Ki67+/TUBB3-) or undefined (outside of gating boundaries). We assume the latter category to represent transitioning cells in intermediate stages of neural development, un- or dedifferentiated iPSCs. (**Fig. 5B**). This gating strategy resulted in a fractional abundance of neurons vs. NPC of 36,4 % in the primed condition and 80,0% in the differentiated condition (**Fig. 5C**). UMAP dimensionality reduction revealed a clear clustering of both phenotypes suggesting that the morphotextural profile captured their distinct phenotypic fingerprint and should enable their separation (**Fig. 5D**).

In a first attempt to classify cells within the guided culture, we trained a CNN on individual cell inputs using the culture condition as a weak label. This resulted in a prediction accuracy of 87,0±0,9%, which is not unexpected given the cellular heterogeneity within each condition. When we used the IF ground truth labels instead, we obtained a classification performance of 96,0±0,5 % (**Fig. 5E**). Applying this cell-based (as opposed to condition-based) CNN to the two culture conditions resulted in predicted fractional abundance of neurons to NPC of 40,5% in images of the primed condition and 74,2% in images of the differentiated condition – aligning well with the manually defined ratios (**Fig. 5F**).

We then went on to evaluate the established cell-based classifier to a primed neuronal cell culture undergoing gradual spontaneous differentiation after dual-SMAD inhibition[25]. We examined cell cultures at 13, 30, 60 and 90 days *in vitro* (DIV) after the start of the differentiation process and visually confirmed a gradual change in neural maturity, as

evidenced by the increasing dendritic outgrowth (**Fig. 5G**). In absence of post-hoc ground truth

labelling, we plotted the cells of the different DIVs onto the UMAP space of the guided

differentiation and quantified their relative abundance in regions encompassing either NPCs or

neurons (**Fig. 5H-I**). This confirmed a shift in the neuron-to-NPC fractional abundance with

an especially strong accumulation of neurons at DIV90. The cell-based CNN model trained on

the guided differentiation dataset recapitulated this shift in its predictions (**Fig. 5J**).

Nucleocentric phenotyping can be applied to characterize mixed iPSC-derived neuronal

cultures

Next to NPCs and neurons, iPSC-derived neuronal cultures are often studied in conjunction

with other relevant cell types that influence neuronal connectivity and homeostasis such as

astrocytes and microglia[26]. Therefore, we tested whether the cell-based approach could be

extended to these cell types as well. We generated iPSC-derived astrocytes, neurons, and

microglia from the same parental iPSC line (**Fig. 6A**) and trained a CNN to identify each cell

type. In monocultures, this led to a prediction accuracy of 96,8±1,0% (**Fig. 6B**) and in mixed

cultures of neurons and microglia (using NeuN, resp. Iba1 as ground-truth IF markers) (**Fig.**

6C) an accuracy of 96,9±0,5% (**Fig. 6D**). Based on these results, we conclude that

nucleocentric phenotyping can be used to gauge the composition of iPSC-derived cultures.

Discussion

iPSC-derived cell cultures can improve translatability of biomedical research but represent a challenge due to their higher variability and multicellular composition. In this research, we have presented a method to efficiently identify neural cell types in mixed cultures to aid in their validation and application in routine screening settings. We first benchmarked our approach using neural cell lines and found that a CNN-based approach outperforms shallow learning (RF) based on handcrafted feature extraction, even with a limited number of input images. This aligns well with recent data showing CNN-based feature representations outperformed shallow techniques for condition-based drug response prediction[27].

We found that all channels contribute to the overall prediction accuracy but that the whole cell is not required to obtain the highest prediction accuracy, especially in dense cultures. In line with earlier studies that highlight the biomarker potential of the nucleus for patient stratification[28], cell state identification[29] or drug response[14], we found that the nuclear ROI as such already carries highly distinctive information for cell type prediction. However, extension to its direct surroundings proved to be the best and most robust input across a range of cell densities. Since errors made by the CNN largely arose from segmentation errors, we hypothesize that such errors will be most pronounced in dense cultures. The nucleocentric approach, which is based on more robust nuclear segmentation, minimizes such mistakes. Furthermore, it opens possibilities for applying cell profiling to even more complex 3D cell systems such as tissue slices or organoids, where cell segmentation becomes extremely challenging.

One important conclusion from this work is that cell types can be identified in mixed cultures solely using the input from monocultures. This implies that cells retain their salient characteristics even in complex heterogeneous environments. While for now, predictions are still superior when based on training on mixed cultures, we found that the prediction accuracy

of monoculture trained models can be increased by employing replicate controls. This suggests

that it may become possible to apply or adapt existing models without the need for a cell-level

ground truth (as provided by post-hoc labelling). This technique could potentially be of use for

cultures or tissues where no antibody- or pre-labelling is possible (*e.g.,* no unique IF marker is

available, non-replicating cells). To increase the accuracy one could resort to intelligent data

augmentation[30] or transfer learning[31] strategies. While a ground-truth independent method

holds promise for bulk predictions (*e.g.,* for quality control purposes), the use of post-hoc

labelling allows building more refined models that can distinguish multiple cell types and/or

cell states at once. Especially using the cyclic staining that we have used[22], much richer

information content can be gained. With the method presented here, we combine the rich

multidimensional information from cyclic IF with morphological information and show that it

is possible to predict these IF outcomes. Multiplexed imaging has previously shown its utility

in gaining in depth information on culture composition and differentiation status in iPSC-

derived neural cultures (progenitor, radial glia, astrocytes (immature and mature) and neurons

(inhibitory and excitatory)[32]. Similarly, this could be expanded to cell states (apoptosis, DNA

damage)[15]. Initial model training requires both CP and multiplexed IF images. After the initial

training phase, the IF step can be omitted, and the CNN will predict the phenotype outcomes

of the IF with high accuracy on the CP images alone. This significantly reduces experimental

costs and time requirements.

Applying the method to iPSC-neuronal cultures revealed its potential to score their

differentiation state. Although guided protocols manage to speed up the differentiation process

and lead to enhanced culture purity, the neural differentiation process proves to be less than

binary, as evidenced by the mixed presence of Ki67+/TUBB3- and Ki67-/TUBB3+ cells. The

spontaneous differentiation protocol illustrated the unpredictable nature and late stage of the

differentiation process. Reports in literature highlight the difficulty of reproducible neural

differentiation and attribute this to culture conditions, cultivation time and variation in

developmental signalling pathways in the source iPSC material[33,34]. Spontaneous neural

differentiation has previously been shown to require approximately 80 days before mature

neurons arise that can fire action potentials and show neural circuit formation. Although these

differentiation processes display a stereotypical temporal sequence[25], the exact timing and

duration might vary. Yet, we could reliably predict the gradual transition to a more

differentiated state using our CNN model. Complementary, the same transition from NPC to

neuronal phenotype could be observed in UMAP space. Although UMAP dimensionality

reduction merely visualizes data structures and is less sensitive as compared to CNN

classification, it aids in visualization and comprehension of the dataset. We realize that

differentiating iPSC cultures are highly heterogenous and are composed of a landscape of

transitioning progenitors in various stages of maturity that our binary models currently fail to

capture. To predict more continuously the maturation of iPSC-derived neural progenitors to

differentiated neurons, it could be of interest to consider deep learning-based regression

approaches [35].

For now, we have tailored models to the individual datasets, but it is conceivable that a more

generalist CNN could be established for multiple culture types. This would obviously demand

a much larger dataset to encompass the variability encountered in such models (*e.g.,* various

starting iPSC lines, various differentiation protocols). Publicly available datasets (*e.g.,* JUMP-

Cell Painting Consortium) can aid in creating an input database containing a large variability

(different iPSC lines, different neural differentiation protocols, …), which would ultimately

lead to a more robust and generalist predictor. Our results showing the prediction accuracy of

a guided differentiation model on spontaneously differentiating cultures indicate that the

approach can be transferred to other differentiation protocols. Inclusion of more input images

and variability should thus enable developing a generalist model for other differentiation

protocols without the need for ground truth validation and further CNN training.

In conclusion, we present a novel application for unbiased morphological profiling by

extending its use to complex mixed neural cultures using sequential multispectral imaging and

convolutional network-informed cell phenotype classification. We show that the resulting

predictors are robust with respect to biological variation and cell culture density. This method

holds promise for use in quality control of iPSC cultures to allow their routine use in high-

throughput and high-content screening applications.

Methods

Cell culture. Cells were cultured at 37 °C and 5% CO2. 1321N1 and SH-SY5Y cell lines were maintained in DMEM-F12 + Glutamax (Gibco, 10565018) supplemented with 10% Fetal Bovine Serum (Gibco, 10500064). Cell seeding prior to imaging was done in 96-well black multiwell plates with #1.5 glass-like polymer coverslip bottom (Cellvis, P96-1.5P). Only the inner 60 wells were used, while the outer wells were filled with PBS-/- to avoid plate effects. Plates were coated with Matrigel (Corning, 734-1440) After seeding, the imaging plate was centrifuged at 100g for 3min.

iPSCs (Sigma Aldrich, iPSC0028 Epithelial-1) were cultured on Matrigel (Corning, 734-1440) in Essential 8 medium (Gibco, A1517001). Upon cell thawing, Rock inhibitor (Y-27632 dichloride, MedChem, HY-10583) was added in 10μM concentration. Subculturing of iPSCs was performed with ReLeSR (Stemcell Technologies, 05872).

Unguided differentiation[25] of iPS cells to neural progenitor cells (NPCs) was started by subculturing the iPSCs single cell using Tryple Express Enzyme (Life technologies, 12605010) at a density of 10e4 cells/cm² in mTesR1 medium (Stemcell Technologies, 85850) and Rock inhibitor. The following day, differentiation to NPCs was started by dual SMAD inhibition in neural maintenance medium (1:1 Neurobasal (Life technologies, 21103049):DMEM-F12+Glutamax (Gibco, 10565018), 0.5x Glutamax (Gibco, 35-050-061), 0.5% Mem Non Essential Amino Acids Solution (Gibco, 11140050), 0.5% Sodium Pyruvate (Gibco, 11360070), 50 μM 2-Mercaptoethanol (Gibco, 31350010), 0.025% Human Insulin Solution (Sigma Aldrich, I9278), 0.5X N2 (Gibco, 17502048), B27 (Gibco, 17504044), 50 U/ml Penicillin-Streptomycin (Gibco, 15140122)) supplemented with 1 μM LDN-193189 (Miltenyi, 130-106-540), SB431542 (Tocris, 1614). Daily medium changes were performed for 11 consecutive days. Following neural induction, neural rosettes were selected by STEMdiff Neural Rosette Selection Reagent (Stemcell technologies, 05832). Maintenance of neural

progenitor cells was performed in neural maintenance medium with 10 µM bFGF (Peprotech, 100-18C) added after subculturing. Single cell detachment of NPCs during maintenance was performed using Tryple Express Enzyme. Cell seeding prior to imaging is done in 96-well black multiwell plates µCLEAR (Greiner, 655090) coated with Poly-L-ornithine (Sigma-Aldrich, P4957) and laminin (Sigma-Aldrich, L2020). Only the inner 60 wells were used, while the outer wells were filled with PBS-/- to avoid plate effects. After seeding, the imaging plate was centrifuged at 100g for 3min.

Guided iPSC differentiation to neurons was performed according to Bell et al. (2019)[2]. The initial phase of neural induction consisted of 12 days neural induction medium (DMEM-F12+Glutamax (Gibco, 10565018), 1x N2 (Gibco, 17502048), 1x B27 (Gibco, 17504044),1mg/ml BSA (Sigma-Aldrich, A7979), 0.5% Mem Non-Essential Amino Acids Solution (Gibco, 11140050)). Of these 12 days, the first 7 were supplemented with cytokines for dual-SMAD inhibition (1 µM LDN-193189 (Miltenyi, 130-106-540), SB431542 (Tocris, 1614)). Following neural induction, the NPCs were floated in uncoated MW6 culture plates in NPC medium (DMEM-F12+Glutamax (Gibco, 10565018), 1x N2 (Gibco, 17502048), 1x B27 (Gibco, 17504044), 10 µM bFGF (Peprotech, 100-18C), 10 µM EGF (Peprotech, 100-47)). After 4 days, NPC clusters of appropriate size were filtered using a cell strainer ((37 µm cell strainer, Stemcell technologies, 27250) and plated on Matrigel-coated (Corning, 734-1440) MW6 culture plates. NPCs can now be expanded in NPC medium. Guided differentiation into forebrain neurons can be induced by switching to neuronal medium (BrainPhys (STEMCELL Technologies, 05792), 1x N2 (Gibco, 17502048), 1x B27 (Gibco, 17504044), 10 µM BDNF (PeproTech, AF-450-02), 10 µM GDNF (PeproTech, AF-450-02) for 15 days before fixation. Differentiation of iPSC to microglia[4] was performed by the formation of embroid bodies (EBs) with 10e3 iPSCs/well in a 96-well U-bottom plate (Corning, 351177) coated with Anti-Adherence Rinsing Solution (Stemcell technologies, 07010) in mTeSR medium supplemented

with Rock inhibitor, 50 ng/mL BMP4 (Peprotech, 120-05), 50 ng/mL VEGF (Peprotech, 100-20), 20 ng/mL SCF (Peprotech, 250-03). 75% medium is changed for 4 consecutive days. After mesoderm induction, EBs are transferred to a 6-well plate with 20 EBs/well and placed in macrophage precursor medium (X-vivo15 (Lonza, BE02-060Q), 100 ng/mL M-CSF (Peprotech, 300-25), 25 ng/mL IL-3 (Peprotech, 213-13), 1x Glutamax, 50 U/ml Penicillin-Streptomycin, 50 μM 2-Mercaptoethanol). 14 days after macrophage differentiation, macrophage precursors were harvested using a cell strainer (Stemcell technologies, 27250). Macrophage precursors were added to the NPC culture in 1:1 neural maintenance medium:microglia medium ((DMEM-F12+Glutamax, 100 ng/mL M-CSF (Peprotech, 300-25), 100 ng/mL IL-34 (Peprotech, 200-34), 1x Glutamax, 50 U/ml Penicillin-Streptomycin, 50 μM 2-Mercaptoethanol).

Replication labelling. Prior to co-seeding of 1321N1 and SH-SY5Y mixed cultures, individual cultures were incubated with either 10 μM EdU (Click-iT® EdU Imaging Kit, Life Technologies, C10340) or 10 μM BrdU (Sigma-Aldrich, B5002) for 24h. This incubation time exceeded the doubling time, allowing incorporation of the nucleotide analog in all cells. This labelling period was followed by a 24h washout period in regular cell culture medium. After washout, the cells were subcultured and plated in coculture. In half of the replicate, SH-SY5Y cells received BrdU while 1321N1 cells received EdU. For the remainder of wells, the pre-label switched cell types.

Morphological staining. Morphological staining (Cell Painting) was adapted from Bray et al. 2016[13]. After careful titration, all dye concentrations were adjusted and evaluated for compatibility with 4-channel laser and filter combinations available on the confocal microscope (see further). Staining was performed on cell cultures fixed in 4% PFA (roti-histofix 4% paraformaldehyde, Roth, 3105.2) for 15min. Cells were rinsed once with PBS-/- (Life Technologies, 10010015) prior to fixation and 3x 5min post fixation. Staining solutions

were prepared fresh before staining in PBS-/- with 0.3% Triton-X-100 (Sigma Aldrich, X100)

(**Table 1**). Each staining solution was incubated for 30min on a shaker at RT in the dark. After

staining, the cells were washed 1x with PBS -/- and sealed for imaging.

Cyclic staining and immunocytochemistry. Cyclic staining is executed by fluorescence

quenching after each sequential imaging round. 1mg/ml in ddH$_2$O LiBH$_4$ solution[22] (Acros

Organics, 206810050) was prepared fresh before use. 1.5h incubation of quenching solution

was performed before each successive staining series. After incubation, the quenching solution

was removed by washing 3x 5min in PBS-/-. Successful fluorescence quenching was

microscopically verified before proceeding with immunofluorescence staining (IF) (**Table 2**).

Cells are treated with PAV blocking buffer (Thimerosal 0.5% (Fluka, 71230), NaN3 0.1%

(Merck, k 6688), Bovine serum albumin (Sigma-Aldrich, A7284), Normal horse serum, PBS-

/-) for 8min. The desired primary antibodies (pAB) are diluted in PAV blocking buffer. pAB

incubation was performed 12h (overnight) at 4°C after which the cells were washed 1x 5min

in PBS-/- followed by incubation in secondary antibody solution (sAB) in PAV + DAPI for

nuclear counterstain. sAB staining was performed for 3h at RT while shaking. Prior to imaging,

the cells were washed 2x with PBS-/- and stored in PBS-/- + 0,1% NaN$_3$.

BrdU staining was performed using IF, requiring DNA denaturation before pAB incubation.

This was performed by 10 min incubation with 2N HCl at 37°C. HCl was neutralized with 0.1

M sodium borate buffer pH 8.5 for 30min at RT. Cells were washed 3x 5min in PBS-/- before

continuing with the general IF protocol. EdU click-it labelling was performed according to the

manufacturer's instructions (Click-iT® EdU Imaging Kit, Life Technologies, C10340).

Image acquisition. Images were acquired using a spinning disk confocal microscope (Nikon

CSU-W1 SoRa) with a 20x 0.75 NA objective (Plan APO VC 20x DIC N2) and Kinetix

sCMOS camera (pinhole 50μm; disk speed 4000 rpm; pinhole aperture 10; bit depth 12-bit,

pixel size 0.325μm²). We opted for confocal microscopy instead of widefield to overcome

image quality limitations resulting from highly dense cell clusters. 96-well plates were scanned,

capturing a minimum of 16 images per well spread in a regular pattern (0,8 mm spacing in x

and y) across the surface of the well. If multiple z-slices were acquired (to correct for surface

inclinations in the field of view), a maximum projection was performed before further analysis.

Morphological images were acquired in all 4 channels. (**Table 3**).

Software. Images were captured by the Nikon CSU-W1 confocal system in combination with

the NIS elements software (RRID:SCR_014329). Image visualization was later performed

using Fiji freeware[36,37]. In-depth image analysis, pre-processing and machine learning for cell

classification were performed using Python programming language[38] in combination with

Anaconda[39] (distribution and package managing software) and Visual Studio Code (code

editor). The packages and versions used for data analysis are shown in **Table 4**.

Image pre-processing. *Cell/nuclear segmentation.* Using a tailor-made script implementing

the Cellpose[40] (cell segmentation) or StarDist[41] (nuclear segmentation) package, images were

pre-processed by normalizing the intensity values of each channel between the 1st and 99th

quantile. Individual channels or channel combinations for segmentation can be selected

depending on the desired outcome mask. Resulting outputs included the segmentation mask

alongside the quality control images. For Stardist implementation, hyperparameter probability

was set at 0,6 and overlap at 0,3. For Cellpose segmentation, 4 models (cyto2) were averaged

to obtain the final mask. Segmentation was performed on the composite of all CP channels. An

estimation of the cell's diameter could be included to optimize cell detection.

Ground truth alignment. Following sequential staining and imaging rounds, multiple images

were captured representing the same cell with different markers. All images were aligned using

Fourier-based image cross correlation on the intensity-normalized multichannel images. The

alignment shift between image1 and image2 was determined using scikit-image phase cross

correlation. Image2 was then moved according to the predetermined shift to align morphological with ground truth images.

Ground truth phenotyping. The true cell phenotype was determined by the fluorescence intensity of the post-hoc immunostaining with class-specific markers. Each ground truth image was imported alongside the corresponding cell mask for that image. For each cell label, the fluorescence intensity was determined and tabulated. The threshold was set manually based on the fluorescence intensity of the monoculture controls. Ground truth labels were assigned to each region of interest (ROI).

Data filtering. Over the entire pipeline, ROIs could be discarded based on 3 conditions: (1) Cell detachment. Due to the harsh sample preparation and repeated washing steps, cells could detach from the imaging substrate thus resulting in lack of ground truth information for those cells. As a result, all ROIs for which there was no DAPI signal detected in the ground truth image, were removed from the dataset as incomplete. (2) Cells for which the ground truth fluorescence intensity was ambiguous. Ground truth labels were determined based on the presence of specific IF staining in either of the phenotype classes. If no class-specific IF staining was detected by thresholding, no true label could be assigned. These ROIs were therefore discarded due to uncertainty. (3) Likelihood of faulty ROI detection. The DAPI signal was used to discard ROIs that do not represent (complete) cells by thresholding on minimal DAPI intensity and minimal nuclear size.

ROI classification. *Train-validation-test split.* Model training requires splitting the available data in 3 subsets: training (60%), validation (10%), and testing (30%). The training dataset was used to train the machine learning models (either RF or CNN). The validation dataset, composing of 10% of the total data, was used to determine the hyperparameters and intermediate testing. The remaining 30% of the dataset was kept apart and used to test the final model when training is completed. For both RF and CNN, the testing dataset was never shown

to the model during the training phase, but only used after training to determine the accuracy

of predictions to the ground truth. The number of instances for each class was equalized by

sampling equal number of instances from each predicted class. To account for variation

between technical replicates, the train-validation-test split was stratified per well. As a result,

no datapoints arising from the same well could appear simultaneously in the training, validation

and testing subset. This data stratification was repeated 3 times with different random seeds

(see Methods – Reproducibility).

Random Forest. For each ROI, a set of manually defined parameters was extracted

corresponding to cell shape (area, convex area, filled area, length minor axis, length major axis,

centroid, eccentricity, equivalent diameter area, ferret diameter max, orientation, perimeter),

texture (GLCM: contrast, dissimilarity, homogeneity, energy, correlation, ASM), intensity

(maximal intensity, minimal intensity, mean intensity, standard deviation intensity) and

distribution (solidity). This was done for 3 regions (nucleus, cytoplasm and complete cell), and

for all channel dimensions. Redundant parameters were removed. All parameters were

standardized per ROI, grouped per replicate. The number of trees within the forest was varied

between 10 and 150, reaching maximum accuracy at around 30 trees.

Uniform Manifold Approximation and Projection. Dimensionality reduction using UMAP was

performed on the same feature matrix as used for RF prediction. Hyperparameters were set at

the default settings.

Convolutional neural network. A ResNet50 model was trained for image classification. In

contrast to classical machine learning techniques, no handcrafted features were extracted. Cell

crops (60µm whole cell – 15µm nucleocentric/nuclear area) were defined based on the

segmentation mask for each ROI. Each ROI was cropped out of the original morphological

image and associated with metadata corresponding to its ground truth label. Images alongside

their labels were fed to the network. Tensors were normalized per channel. Models are trained

on a minimum of 5000 training inputs of each class for 50 epochs (training iterations). Each

batch consisted of 100 samples. The training input was augmented by linear transformations

(horizontal and vertical flip, random rotation). Each epoch, the current model was tested against

a validation dataset. The performance of the model on this validation subset determined

whether the model was stored as new best (if the new accuracy exceeded the accuracy of the

previous best model) or discarded. The learning rate at the start was set at 0,0001 and

automatically reduced with a factor of 0,1 during training when no improvement was seen after

10 epochs. After 50 epochs, the best resulting model was tested on a separate test dataset to

determine the accuracy on previously unseen data.

Reproducibility. Each model training was performed 3 independent times. This was repeated

for 3 different random seeds. Each model received input data arising from a minimum of 16

images per well, at least 15 technical replicates (wells). The optimization experiments (figures

1-4) were performed with cell lines with limited variability. These models were trained on 3

independent experiments where ground truth pre-labelling (Edu/BrdU) was performed at least

once on either of the cell lines in coculture. For iPSC-derived cultures, as variability is inherent

to these differentiations, 3 biological replicates (independent differentiations) were pooled for

model training.

Statistics. All statistical comparisons were made nonparametric using Mann-Whitney U (for

two independent sample comparison) or Kruskal-Wallis (for multiple sample comparison) with

pairwise tests using Tukey's honestly significant difference test. We opted for nonparametric

testing because the number of models in each group to be compared was < 15. Significance

levels are indicated on the figures using ns. (no statistical significance, p-value above 0,05), *

(p-value between 0,05 and 5e-4), ** (p-value between 5e-4 and 5e-6) and *** (p-value smaller

than 5e-6).

Funding

This work was funded by Fonds Wetenschappelijk Onderzoek Vlaanderen (1SB7423N; 1274822N), BOF (FFB210009) and IOF Uantwerpen (FFI210239; FFI230099) and VLAIO (HBC.2023.0155).

Acknowledgements

We thank Marlies Verschuuren and Hugo Steenberghen for their assistance and knowledge regarding the cell lines. By extension, we would like to thank all current and former members of the De Vos lab.

Data availability statement

The authors report that the results of this study are available within the manuscript and supplementary materials. All image analysis scripts are open-source available on GitHub (https://github.com/DeVosLab)

Conflict of interest

The authors declare no conflict of interest.

Abbreviations

BrdU: Bromodeoxyuridine; CNN: convolutional neural network; CP: cell painting; DAPI: 4',6-diamidino-2-fenylindool; DIV: days in vitro; EdU: 5-ethynyl-2'-deoxyuridine; ER: endoplasmic reticulum; Grad-CAM: Gradient-weighted Class Activation Mapping; IF:

553 immunofluorescence; iPSC: induced pluripotent stem cell; NPC: neural progenitor cell; RF:

554 random forest; ROI: region of interest; TUBB3: beta-III-tubulin; UMAP: Uniform Manifold

555 Approximation and Projection

Author contributions

558 The experiments were conceptualized by SDB, TVDL, JVDD, PP and WDV. Experiments

559 were executed by SDB and JVDD. TVDL and SDB optimized the data analysis scripts. SDB

560 analysed the imaging data and performed the data analysis. SDB and WDV prepared the

561 original draft. PP and WDV supervised the work. All authors took part in reviewing and editing

562 the manuscript.

References

1. De Strooper, B. & Karran, E. The Cellular Phase of Alzheimer's Disease. *Cell* **164**, 603–615 (2016).

2. Bell, S. *et al.* Differentiation of Human Induced Pluripotent Stem Cells (iPSCs) into an Effective Model of Forebrain Neural Progenitor Cells and Mature Neurons. *BIO-PROTOCOL* **9**, (2019).

3. Neyrinck, K. *et al.* SOX9-induced Generation of Functional Astrocytes Supporting Neuronal Maturation in an All-human System. *Stem Cell Rev and Rep* **17**, 1855–1873 (2021).

4. Haenseler, W. *et al.* A Highly Efficient Human Pluripotent Stem Cell Microglia Model Displays a Neuronal-Co-culture-Specific Expression Profile and Inflammatory Response. *Stem Cell Reports* **8**, 1727–1742 (2017).

5. Nevin, Z. S. *et al.* Modeling the Mutational and Phenotypic Landscapes of Pelizaeus-Merzbacher Disease with Human iPSC-Derived Oligodendrocytes. *The American Journal of Human Genetics* **100**, 617–634 (2017).

6. Abutaleb, N. O. & Truskey, G. A. Differentiation and characterization of human iPSC-derived vascular endothelial cells under physiological shear stress. *STAR Protocols* **2**, 100394 (2021).

7. Paik, D. T., Chandy, M. & Wu, J. C. Patient and Disease–Specific Induced Pluripotent Stem Cells for Discovery of Personalized Cardiovascular Drugs and Therapeutics. *Pharmacol Rev* **72**, 320–342 (2020).

8. Lopez-Gonzalez, R. *et al.* Poly(GR) in C9ORF72 -Related ALS/FTD Compromises Mitochondrial Function and Increases Oxidative Stress and DNA Damage in iPSC-Derived Motor Neurons. *Neuron* **92**, 383–391 (2016).

9. Amin, N. D. & Paşca, S. P. Building Models of Brain Disorders with Three-Dimensional Organoids. *Neuron* **100**, 389–405 (2018).

10. D'Antonio, M. *et al.* High-Throughput and Cost-Effective Characterization of Induced Pluripotent Stem Cells. *Stem Cell Reports* **8**, 1101–1111 (2017).

11. Chen, C. X.-Q. *et al. Standardized quality control workflow to evaluate the reproducibility and differentiation potential of human iPSCs into neurons.* http://biorxiv.org/lookup/doi/10.1101/2021.01.13.426620 (2021) doi:10.1101/2021.01.13.426620.

12. Gustafsdottir, S. M. *et al.* Multiplex Cytological Profiling Assay to Measure Diverse Cellular States. *PLoS ONE* **8**, e80999 (2013).

13. Bray, M.-A. *et al.* Cell Painting, a high-content image-based assay for morphological profiling using multiplexed fluorescent dyes. *Nat Protoc* **11**, 1757–1774 (2016).

14. Cimini, B. A. *et al.* Optimizing the Cell Painting assay for image-based profiling. *Nat Protoc* **18**, 1981–2013 (2023).

15. Way, G. P. *et al.* Predicting cell health phenotypes using image-based morphology profiling. *MBoC* **32**, 995–1005 (2021).

16. Schiff, L. *et al.* Integrating deep learning and unbiased automated high-content screening to identify complex disease signatures in human fibroblasts. *Nat Commun* **13**, 1590 (2022).

17. Chandrasekaran, S. N. *et al. JUMP Cell Painting dataset: morphological impact of 136,000 chemical and genetic perturbations.* http://biorxiv.org/lookup/doi/10.1101/2023.03.23.534023 (2023) doi:10.1101/2023.03.23.534023.

18. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and Projection for Dimension Reduction. (2018) doi:10.48550/ARXIV.1802.03426.

19.   Nguyen, T.-T., Huang, J. Z. & Nguyen, T. T. Unbiased Feature Selection in Learning Random Forests
for High-Dimensional Data. *The Scientific World Journal* **2015**, 1–18 (2015).

20.   He, K., Zhang, X., Ren, S. & Sun, J. Deep Residual Learning for Image Recognition. Preprint at
http://arxiv.org/abs/1512.03385 (2015).

21.   Selvaraju, R. R. *et al.* Grad-CAM: Visual Explanations from Deep Networks via Gradient-Based
Localization. *Int J Comput Vis* **128**, 336–359 (2020).

22.   Radtke, A. J. *et al.* IBEX: an iterative immunolabeling and chemical bleaching method for high-content
imaging of diverse tissues. *Nat Protoc* **17**, 378–401 (2022).

23.   Hernández, D. *et al.* Culture Variabilities of Human iPSC-Derived Cerebral Organoids Are a Major Issue
for the Modelling of Phenotypes Observed in Alzheimer's Disease. *Stem Cell Rev and Rep* **18**, 718–731 (2022).

24.   Volpato, V. & Webber, C. Addressing variability in iPSC-derived models of human disease: guidelines
to promote reproducibility. *Disease Models & Mechanisms* **13**, dmm042317 (2020).

25.   Shi, Y., Kirwan, P. & Livesey, F. J. Directed differentiation of human pluripotent stem cells to cerebral
cortex neurons and neural networks. *Nat Protoc* **7**, 1836–1846 (2012).

26.   Kuijlaars, J. *et al.* Sustained synchronized neuronal network activity in a human astrocyte co-culture
system. *Sci Rep* **6**, 36529 (2016).

27.   Dincer, A. B., Celik, S., Hiranuma, N. & Lee, S.-I. *DeepProfile: Deep learning of cancer molecular*
profiles for precision medicine. http://biorxiv.org/lookup/doi/10.1101/278739 (2018) doi:10.1101/278739.

28.   De Vos, W. H. *et al.* Increased plasticity of the nuclear envelope and hypermobility of telomeres due to
the loss of A–type lamins. *Biochimica et Biophysica Acta (BBA) - General Subjects* **1800**, 448–458 (2010).

29.   Heckenbach, I. *et al.* Nuclear morphology is a deep learning biomarker of cellular senescence. *Nat Aging*
2, 742–755 (2022).

30.   Mikolajczyk, A. & Grochowski, M. Data augmentation for improving deep learning in image
classification problem. in *2018 International Interdisciplinary PhD Workshop (IIPhDW)* 117–122 (IEEE, 2018).
doi:10.1109/IIPHDW.2018.8388338.

31.   Nguyen, L. D., Lin, D., Lin, Z. & Cao, J. Deep CNNs for microscopic image classification by exploiting
transfer learning and feature concatenation. in *2018 IEEE International Symposium on Circuits and Systems*
(ISCAS) 1–5 (IEEE, 2018). doi:10.1109/ISCAS.2018.8351550.

32.   Tomov, M. L. *et al.* Resolving cell state in iPSC-derived human neural samples with multiplexed
fluorescence imaging. *Commun Biol* **4**, 786 (2021).

33.   Strano, A., Tuck, E., Stubbs, V. E. & Livesey, F. J. Variable Outcomes in Neural Differentiation of
Human PSCs Arise from Intrinsic Differences in Developmental Signaling Pathways. *Cell Reports* **31**, 107732
(2020).

34.   Galiakberova, A. A. *et al.* Different iPSC-derived neural stem cells shows various spectrums of
spontaneous differentiation during long term cultivation. *Front. Mol. Neurosci.* **16**, 1037902 (2023).

35.   Du, J. & Xu, Y. Hierarchical deep neural network for multivariate regression. *Pattern Recognition* **63**,
149–157 (2017).

36.   Rueden, C. T. *et al.* ImageJ2: ImageJ for the next generation of scientific image data. *BMC*
Bioinformatics **18**, 529 (2017).

37.   Schindelin, J., Rueden, C. T., Hiner, M. C. & Eliceiri, K. W. The ImageJ ecosystem: An open platform
for biomedical image analysis. *Molecular Reproduction Devel* **82**, 518–529 (2015).

38.   Van Rossum, G., & Drake, F. L. Python 3 Reference Manual. (2009).

39.   Anaconda Software Distribution. Anaconda Documentation.

40.   Stringer, C., Wang, T., Michaelos, M. & Pachitariu, M. Cellpose: a generalist algorithm for cellular
segmentation. *Nat Methods* **18**, 100–106 (2021).

41.   Weigert, M., Schmidt, U., Haase, R., Sugawara, K. & Myers, G. Star-convex Polyhedra for 3D Object
Detection and Segmentation in Microscopy. in *2020 IEEE Winter Conference on Applications of Computer Vision*
(WACV) 3655–3662 (IEEE, 2020). doi:10.1109/WACV45572.2020.9093435.

  Tables

Table 1 | Specifications of morphological staining composition.

Staining solution 1							
Dye	**Target**	**Excitation (nm)**	**Emission (nm)**	**Stock concentration**	**End dilution**	**Supplier**	**Catalog #**
DAPI	Nucleus	350	470	2.5 mg/ml	1/500	Sigma Aldrich	D9542
Concanavalin	ER	480	490-540	5 mg/ml	1/300	Life Technologies	C11252
WGA	Golgi	550	560-570	1 mg/ml	1/600	Life Technologies	W32464
Phalloidin	Actin	570	580-610	66μM	1/1000	Life Technologies	A12380
Mitotracker	Mitochondria	640	650-660	1 mM	1/1000	Life Technologies	M22426
Staining solution 2							
Syto14	Nucleoli /RNA	525	540-580	5 mM	1/1000	Life Technologies	S7576

Table 2 | Used antibodies.

Antibody		Host	Supplier	Catalog #
Primary antibodies	Anti-BrdU	Sheep	Abcam	ab1893
	Anti-MAP2	Chicken	Synaptic systems	188006
	Anti-TUBB3	Mouse	BioLegend	801202
	Anti-CD44	Mouse	Millipore	MAB4065
	Anti-GFAP	Goat	Abcam	ab53554
	Anti-NeuN	Guinea pig	Millipore	ABN90P
	Anti-Iba1	Rabbit	Wako	019-19741
	Anti-Ki67	Rabbit	Abcam	ab15580
Secondary antibodies	Anti-mouse-Cy3	Donkey	Jackson ImmunoResearch	715-165-150
	Anti-mouse-Cy5	Donkey	Jackson ImmunoResearch	715-175-150
	Anti-sheep-Cy3	Donkey	Jackson ImmunoResearch	713-165-147
	Anti-sheep-Cy5	Donkey	Jackson ImmunoResearch	713-175-147
	Anti-chicken-Cy5	Donkey	Jackson ImmunoResearch	703-175-155
	Anti-goat-Cy3	Donkey	Jackson ImmunoResearch	705-165-147
	Anti-Guinea pig-Cy3	Donkey	Jackson ImmunoResearch	706-165-148
	Anti-Rabbit-Cy5	Donkey	Jackson ImmunoResearch	711-175-152

Table 3 | Specifications of the used laser lines, excitation and emission filters.

Channel dimension	Laser line (nm)	Excitation filter (nm)	Emission filter (nm)

Channel 1	405	Di01-T405/488/568/647-	B447/60
Channel 2	488	13x15x0.5	B520/35
Channel 3	561		B617/73
Channel 4	640		B685/40

Table 4 | Python packages used for image and data analysis alongside the software version.

Package	Version	Reference
pathlib	1.0.1	
numpy	1.21.2	Harris, C. R., Millman, K. J., van der Walt, S. J., Gommers, R., Virtanen, P., Cournapeau, D., … Oliphant, T. E. (2020). Array programming with NumPy. Nature, 585, 357–362. https://doi.org/10.1038/s41586-020-2649-2
tifffile	2021.7.2	
pytorch	1.13.0	Paszke, A., Gross, S., Massa, F., Lerer, A., Bradbury, J., Chanan, G., … Chintala, S. (2019). PyTorch: An Imperative Style, High-Performance Deep Learning Library. In Advances in Neural Information Processing Systems 32 (pp. 8024–8035). Curran Associates, Inc.
torchvision	0.14.0	
nd2reader	3.3.0	
matplotlib	3.5.2	Hunter, J. D. (2007). Matplotlib: A 2D graphics environment. Computing in Science & Engineering, 9(3), 90–95.
scikit-image	0.19.3	Van der Walt, S., Sch"onberger, Johannes L, Nunez-Iglesias, J., Boulogne, Franccois, Warner, J. D., Yager, N., … Yu, T. (2014). scikit-image: image processing in Python. PeerJ, 2, e453.

scikit-learn	1.1.3	Pedregosa, F., Varoquaux, Ga"el, Gramfort, A., Michel, V., Thirion, B., Grisel, O., … others. (2011). Scikit-learn: Machine learning in Python. Journal of Machine Learning Research, 12(Oct), 2825–2830.
argparse	1.4.0	
cellpose	2.1.1	Stringer, C., Wang, T., Michaelos, M. et al. Cellpose: a generalist algorithm for cellular segmentation. Nat Methods 18, 100–106 (2021). https://doi.org/10.1038/s41592-020-01018-x
tqdm	4.64.1	
scipy	1.9.3	Virtanen, P., Gommers, R., Oliphant, T. E., Haberland, M., Reddy, T., Cournapeau, D., … SciPy 1.0 Contributors. (2020). SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods, 17, 261–272. https://doi.org/10.1038/s41592-019-0686-2
pandas	1.5.1	McKinney, W., & others. (2010). Data structures for statistical computing in python. In Proceedings of the 9th Python in Science Conference (Vol. 445, pp. 51–56).
seaborn	0.12.1	https://doi.org/10.5281/zenodo.883859
imbalanced-learn	0.10.0	Imbalanced-learn: a python toolbox to tackle the curse of imbalanced datasets in machine learning
StarDist	0.8.5	M. Weigert, U. Schmidt, R. Haase, K. Sugawara and G. Myers, "Star-convex Polyhedra for 3D Object Detection and Segmentation in Microscopy," 2020 IEEE Winter Conference on Applications of Computer Vision (WACV), Snowmass, CO, USA, 2020, pp. 3655-3662, doi: 10.1109/WACV45572.2020.9093435.

Figure 1 | Shallow classification using manually defined features. (A) Image overview of 1321N1 cells after CP staining acquired at higher magnification (Plan Apo VC 60xA WI DIC N2, NA = 1,20). Scale bar is 30µm. (B) UMAP dimensionality reduction, color coded for either cell type condition (left) or replicate (right). (C) Feature importance deducted from the UMAP (feature maps) and RF (Gini importance). Three exemplary feature maps are highlighted alongside the quantification per cell type. (D) Random Forest classification performance with confusion matrix and Gini importance. All features used in the UMAP are used for RF building.

Figure 2 | Convolutional neural network classification of monoculture conditions. (A) Schematic of image pre-processing for CNN classification. (B) CNN accuracy and confusion matrix on cell type classification in monocultures. (C) Shallow vs. deep learning accuracy for varying training set sizes. Dots represent different training iterations (cross-validation). Train-validation-test dataset selection was performed with the same random seed at 60-10-30 ratios. (D) The impact of variability in the training dataset on model accuracy for shallow vs. deep learning. Mann-Whitney-U-test, p-values resp. 7,47e-6 and 1,96e-4. (E) The performance of deep learning classifiers trained on single replicates (low variability) and mixed replicates (high variability) on unseen images (independent replicates). Kruskal-Wallis test on single training condition, p-value of 0,026 with post-hoc Tukey test. Mann-Whitney-U-test on mixed training, p-value of 7,47e-6. (F) GradCAM map highlighting the regions on which the CNN relies for making its prediction. (G) Performance accuracy of the CNN when only a selection of image channels is given as training input.

Figure 3 | The impact of regional information on CNN performance in relation to culture density. (A) Definition of different culture density categories with corresponding images. (B) The effect of culture density (number of cells within the FOV) on prediction accuracy, for each of the different input regions. Kruskal-Wallis test for each of the regions, statistically significant decrease (p-value 2,07e-5) between the most dense condition with all other density categories for the cell region. (C) CNN accuracy for different input regions. Kruskal-Wallis test, p-value of 4,29e-4 with post-hoc Tukey test. (D) GradCAM intensity plots highlighting the regions on which the CNN relies for making its prediction in nucleocentric (top) and nuclear (bottom) crops.

Figure 4 | Cell type prediction in mixed cultures by post-hoc ground truth identification. (A) Schematic overview of virtual vs. physically mixed cultures and the subsequent CNN models. For virtual mixing, individual cell types arise from distinct cultures, the cell type label was assigned based on the culture condition. For physical mixing, the true phenotype of each crop was determined by intensity thresholding after post-hoc staining. Three model types are defined: Mono2Mono (culture-based label), Co2Co (cell-based label) and Mono2Co (trained with culture-based label and tested on cell-based label). (B) Determining ground-truth phenotype by intensity thresholding on IF (top) or pre-label (bottom). The threshold was determined on the intensity of monocultures. Only the pre-labelled ground-truth resulted in good cell type separation by thresholding for 1321N1 vs. SH-SY5Y cells. (C) Mono2Mono (culture-based ground truth) vs. Co2Co (cell-based ground truth) models for cell type classification. Mann-Whitney-U-test p-value 0,0027. (D) Monoculture-trained models were tested on mixed cultures. Pretrained models were trained on independent biological replicates. These could be finetuned by additional training on monoculture images from the same replicate as the coculture. This was shown to reduce the variation between model iterations (Median performance: Mann-Whitney-U-test, p-value 0,0015; Coefficient of variation: Mann-Whitney-U-test, p-value 3,48e-4)

Figure 5 | iPSC-derived differentiation staging using morphology profiling. (A) Schematic overview of guided vs. spontaneous neural differentiation. (B) Representative images of morphological staining and post-hoc IF of primed and differentiated iPSC-derived cultures (guided differentiation). Ground-truth determination is performed using TUBB3 (for mature neurons) and Ki67 (mitotic progenitors). Scale bar is 200µm. (C) Fraction of neurons vs. NPC cells in the primed vs. differentiated condition as determined by IF staining. Upon guided differentiation, the fraction of neurons increased. (D) Following feature extraction (handcrafted features), supervised UMAP dimension reduction shows separation between the primed vs. differentiated cultures. (E) CNN performance when classifying neurons (Ki67-/TUBB3+) vs. NPC (Ki67+/TUBB3-) cells using either a condition-based or cell-type based ground truth. Mann-Whitney-U-test, p-value 4,04e-4. (F) Fractional abundance of predicted cell phenotypes (NPC vs. neurons) in primed vs. differentiated culture conditions using the cell-based CNN. (G) Representative images of spontaneously differentiating neural cultures. Scale bar is 200µm. (H) Handcrafted features were extracted from the morphological images. These morphological profiles were plotted onto the UMAP feature space created using the guided differentiation (panel D). Plotting the points of spontaneously differentiating cultures onto this space, revealed gradual displacement of the morphological profile from the NPC towards the neuronal cluster with increasing differentiation time. (I) Quantification of the number of points in each cluster of panel H. (J) Prediction of differentiation status using the cell-based CNN model trained on guided differentiated culture.

Figure 6 | iPSC cell type identification using morphology profiling. (A) Representative images of iPSC-derived neurons, astrocytes and microglia with morphological staining. Scale bar is 200μm. (B) Prediction accuracy of a CNN trained to classify monocultures of iPSC-derived astrocytes, microglia and neurons (panel A) with confusion matrix (average of all models). (C) Representative images of a mixed culture of iPSC-derived neurons and microglia. Ground-truth identification was performed using IF. (D) CNN accuracy of a model trained to identify both cell types in mixed culture mounted up to 98%.

Supplementary figure 1 | The influence of cell density on segmentation quantity and quality. (A) Representative images of 1321N1 cells with increasing density alongside their cell and nuclear mask produced using resp. Cellpose and Stardist. (B) the number of ROIs detected in comparison to the ground truth (manual segmentation) (determined as intersection over union (IoU) < 0,15). (C) ROI detection quality (IoU) for increasing cell density for cell and nuclear masks.

Supplementary figure 2 | Fluorescence quenching over time using LiBH4. (A) Images before and after quenching for all 4 fluorescence channels. (B) Time curve of normalized image-level fluorescence intensity during incubation with 1mg/ml LiBH4.

A

Pre-labelling and cell culture
24h 24h 24h

CP
30m 30m

Cyclic IF
1,5h 12h 3h

24h 10µM Edu/BrdU
24h washout
Cell plating 24h before fixation

1x PBS-/- wash, 15min PFA fixation, 3x PBS-/- wash
30min incubation in staining solution 1
30min incubation in staining solution 2
1x PBS-/- wash

1,5h incubation in 1mg/ml LiBH4 , 3x PBS-/- wash
8min PAV+Triton-X-100
12h (ON) pAB incubation in PAV
2x PBS-/- wash
3h sAB incubation in PAV + DAPI
2x PBS-/- wash

B

Extract features → Build UMAP → Train/evaluate RF

Image segmentation → Alignment with ICC → Determine ground truth label

Crop ROIs → Train/evaluate CNN

Only in mixed cultures
Shallow learning
Deep learning

C

Increasing accuracy with number of epochs during CNN training

Accuracy (%)

Number of training epochs

Performance
- Accuracy
- True positive rate
- True negative rate

Supplementary figure 3 | Methods. (A) Pipeline used for morphological profiling in mixed cultures. (B) Overview of the steps within the image analysis pipeline. (C) Evaluation of accuracy, true negative and true positive rate during CNN training (1321N1 vs. SH-SY5Y in monocultures) across all 50 epochs.

Supplementary figure 4 | Additional figures. (A) PCA dimension reduction on handcrafted features extracted from monocultures of 1321N1 and SH-SY5Y cells. In contrast to UMAP dimension reduction, this linear approach does not result in clear cell type clustering. (B) Examples of misclassified ROIs. (C) Definition of cell regions given as training input for nuclear and nucleocentric model training. (D) Decreasing nuclear area in function of culture density. (E) Evaluation of cell phenotype prediction on an image of differentiating neurons (guided differentiation). The predicted phenotype is shown by the color of the square around each ROI. The distinctive phenotype is shown in the insets. (F) Additional GradCAM images for both cell type classes. (G) Predicted neuron-to-NPC ratio in spontaneously differentiating cultures as predicted by the condition-based classifier (Fig. 5E) in contrast to the cell-based classifier performance on the same dataset (Fig. 5H).