1

Acquisition phase-specific contribution of climbing fiber transmission to cerebellum-

dependent motor memory

Jewoo Seo[1,2,†], Seung Ha Kim[1,2,†], Jaegeon Lee[1,2,†], Min Seok Kim[1,2], Yong-Seok Lee [1,2,3],

Sang Jeong Kim[1,2,3*]

Author affiliations

[1]Department of Physiology, Seoul National University College of Medicine, Seoul, Republic

of Korea

[2]Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul,

Republic of Korea

[3]Memory Network Medical Research Center, Neuroscience Research Institute, Wide River

Institute of Immunology, Seoul National University College of Medicine, Seoul, Republic of

Korea

*: Corresponding author

†: Authors contributed equally

Abstract

Climbing fiber (CF) transmission from the inferior olive (IO) triggers complex spikes (Cs) in

Purkinje cells (PCs) driven by a burst of calcium spikes. In the context of motor learning,

especially the compensatory optic response, CF transmission serves as an instructive signal

selectively conveyed to PCs. While the significance of CF input in motor memory formation

is widely acknowledged, a comprehensive understanding of its distinct contribution across

different temporal windows, spanning from the initial learning phase to the retrieval period,

remains incomplete. Therefore, we aimed to investigate the necessity of CF-induced

instructive signals in motor learning by assessing their roles in memory acquisition,

consolidation, and retrieval. We employed optogenetics to selectively inhibit CF transmission

during targeted time windows. Consequently, the absence of CF-induced instructive signals

during motor learning impairs memory acquisition. However, when these signals were

suppressed during the consolidation and retrieval period, there was neither a loss of long-term

memory nor a prevention of memory retrieval. Our results highlight that CF transmission

plays a specialized and critical role primarily in memory acquisition, rather than in

subsequent processes.

Introduction

The cerebellar architecture, characterized by multiple synapses connecting diverse neuron types, enables precise adjustments within intricate neural networks (Boqhyden et al., 2004). Within this framework, the climbing fiber (CF) transmission derived from the inferior olive (IO) projecting to the cerebellar cortex is generally recognized as the primary pathway for relaying instructive signals to correct errors and enhance motor performance, including oculomotor behaviors (Albus, 1971; Frens et al., 2001; Ito, 1972; Marr, 1969).

A prominent example of cerebellum-dependent motor learning is the optokinetic reflex (OKR), a compensatory optic response evoked by an unstable visual field (Cahill & Nathans, 2008; Schweigart et al., 1997). In OKR, adaptation to stabilize retinal images amid mismatched visual input requires the engagement of multiple neural substrates. For instance, complex spikes (Cs) in floccular Purkinje cells (PCs) driven by CF input, emerge in response to unpredicted motor performance errors (Clopath et al., 2014; Eccles et al., 1966; Goossens et al., 2004; Palmer et al., 2010). Rotational visual stimulation induces Cs discharge in floccular PCs, reflecting CF-induced Cs generation in response to error signals generated from errors in oculomotor performance (Frens et al., 2001; Graf et al., 1988). Therefore, this fine-tuning of Cs in PCs is expected to regulate learning by modulating error-correcting adaptations (Yang & Lisberger, 2014).

Previous studies highlight a potential duality in CF involvement in cerebellar motor learning. For example, one study using a modified behavioral paradigm to suppress Cs suggested that simple spikes (Ss) alone may suffice for motor learning (Ke et al., 2009). In contrast, pharmacological lesioning and optogenetic inactivation of the IO impair oculomotor learning (Pham et al., 2020) and prevent eye blink conditioning (Kim et al., 2020; Medina et al., 2002; Welsh & Harvey, 1998). While CF's role in acquisition is well-supported by existing literature, its involvement in downstream memory processes like consolidation and

retrieval remains contentious. This ambiguity stems from methodological limitations in

previous studies, which have lacked the spatial and temporal precision to isolate CF function

at distinct learning stages. Addressing this knowledge gap is essential to clarify whether CF

input is uniformly critical or phase-specific in motor learning and memory.

To address this gap, we investigated the necessity of CF-induced instructive signals in

motor learning by assessing their contributions across memory acquisition, consolidation, and

retrieval phases. Using optogenetics, we selectively inhibited CF signals in the flocculus, a

lobe-like cerebellar structure primarily responsible for ocular reflex, during targeted learning

stages. Our findings provide compelling evidence for a phase-specific role of CF

transmission in motor learning. This indicates that CF input is essential primarily during

acquisition, with minimal involvement in consolidation and retrieval.

Results

Optogenetic inhibition of climbing fiber transmission derived from the inferior olive

To specifically manipulate CF signals in the cerebellar cortex, generally known to originate from the IO, we injected AAV1-CaMKIIα-eNpHR 3.0-EYFP or AAV1-CaMKIIα-EGFP into the IO of wild-type mice (Figure 1A, B). This approach allowed visualization of IO neuron expression and CF terminals in the brainstem nucleus and flocculus. Importantly, the selectively segregated expression in these regions enabled manipulation of CF terminals without disrupting IO somas, which, if damaged, can critically impair motor performance and ocular reflexes due to the IO's role as a prominent center of sensory integration (Pham et al., 2020; Shaikh et al., 2017; Van Der Giessen et al., 2008).

To confirm the effectiveness of optogenetic inhibition of climbing fiber (CF) transmission, we performed whole-cell patch-clamp recordings of Purkinje cells (PCs) in acute slices of cerebellar vermis lobules 4–5. A 593 nm yellow laser was used to activate halorhodopsin (NpHR). PCs were selected based on optical fluorescence expression (Figures 1C and 1E), and a stimulation electrode was positioned in the granule cell layer near the recorded PC. The stimulation intensity was carefully calibrated to evoke minimal CF excitatory postsynaptic currents (EPSCs). Laser illumination (3–5 mW) during stimulation resulted in a significant reduction in EPSC amplitude, indicating robust inhibition of CF transmission. Notably, this inhibition was consistent, with 9 out of 9 cells in the NpHR group exhibiting reliable suppression without failures. Importantly, laser illumination did not affect CF EPSCs in the GFP group (Figure 1D). Under the current-clamp mode, CF stimulation-induced spikelets were also completely abolished in the NpHR group but remained unaffected in the GFP group (Figure 1F). These findings establish that optogenetic inhibition effectively suppresses CF-PC synaptic transmission in vitro.

To extend our findings in vivo, we recorded from PCs in awake, head-fixed mice during

593 nm optogenetic stimulation. AAV1-CaMKIIα-eNpHR 3.0-EYFP or AAV1-CaMKIIα-

EGFP was injected into the IO followed by the creation of a cranial window above the

cerebellar vermis three weeks later (Figure 2A). Due to technical constraints preventing

direct access to the flocculus for in vivo recordings, we targeted cerebellar lobule IV/V. PCs

were identified based on the detection of Cs (Figure 2B).

Optogenetic inhibition selectively suppressed Cs firing rates in the NpHR group without

affecting Ss rates, whereas no changes were observed in the GFP control group (Figures 2C

and 2D). Recordings from the same neurons during both laser-off and laser-on conditions

demonstrated consistent and robust suppression of Cs activity throughout the 40-minute

optostimulation period. Although continuous recordings over several hours were not feasible,

the stability and sustained suppression observed at 40 minutes strongly suggest that the

manipulation would remain effective during the extended durations required for behavioral

experiments. These findings confirm that optogenetic inhibition of NpHR-expressing CFs

selectively suppresses Cs activity in awake mice without altering Ss activity. Furthermore,

they validate the feasibility and specificity of using optogenetic interventions to block CF

transmission both in vitro and in vivo, reinforcing the reliability of this approach for studying

CF contributions to cerebellar function and behavior.

Inhibition of climbing fiber transmission during memory acquisition

An evaluation of CF's direct contribution in a region-specific manner has not been performed.

Thus, after confirming the viability of optogenetic inhibition of CF, we investigated whether

the elimination of CF transmission in the cerebellar flocculus affects cerebellar motor

learning. When images on the retina are unstable due to mismatched visual inputs, CF signals

are conventionally perceived to be responsible for sending error signals to the cerebellar

cortex to improve ocular reflexes (Bloedel & Bracha, 1998; Frens et al., 2001; Zang & De

Schutter, 2019). Using the same mouse model shown in Figure 1, we surgically implanted an

optical fiber in flocculus to suppress CF transmission in vivo during OKR (Figure 3A). Head-

fixed mice were placed in a rotating drum painted with a black striped pattern to provide

visual input, and its rotating motion elicited an unstable retinal image. Hence, in the

beginning, the trace of the mouse's eye movement represented in a sine curve was only

partially overlapping with the trace of the drum's kinetics, but as the mouse underwent more

sessions of OKR learning, the trace of eye movement became almost identical (Figure 3Bi).

Notably, the group with inhibited CF transmission exhibited only slight changes or no

improved trace of eye movements (Figure 3Bii). Overall, the results demonstrated that CF

transmission is required for motor memory acquisition.

This indicates that CF transmission is essential for cerebellar motor learning by presenting

two contrasting learning curves. The control group exhibited an increasing trend in the

learning curve, whereas the NpHR group's learning curve was almost flat, which reflects no

change of gain (Figure 3C). The gain value through the learning sessions was distinctly

increased in the control group, whereas the NpHR group showed no significant change in

gain (Figure 3D). Furthermore, from 0 to 50 min period of learning, the control group

demonstrated almost a doubling in gain, whereas the NpHR group showed little or almost no

change of gain (Figure 3E). These results indicate that CF transmission is required for the

proper acquisition of motor memory.

Inhibition of climbing fiber transmission during memory consolidation and retrieval

phase

When memories are initially acquired, they are generally unstable; hence, it requires some

time to shape a stabilized long-term memory (Attwell et al., 2002). Therefore, memories

undergo multiple processes that allow distinct shifts from short-term to long-term memory,

leading to the construction of adapted behavior (Shutoh et al., 2006). Having confirmed the

active involvement of CF transmission during memory acquisition, we sought to understand

if its role is limited to the acquisition period or extends to memory transfer during the

consolidation phase for long-term storage. We implemented identical surgical and behavioral

procedures as previously conducted. However, instead of optogenetically inhibiting the

transmission during memory acquisition, we suppressed CF activity in two distinct

optogenetic manipulation schemes: one targeting short-term inhibition (30 min) and the other

for long-term inhibition (6 hr) (Figures 4A and 4D). This approach was based on findings that

long-term memories associated with cerebellum-dependent behaviors are vulnerable to

disruption within specific time windows of consolidation, spanning from minutes to hours

(Cooke et al., 2004; Titley et al., 2007). Learning curves confirmed the successful gain

enhancement, validating the integrity of memory acquisition (Figures 4B and 4E). Once the

gain was robustly enhanced, optogenetic inhibition of CF transmission during consolidation

did not yield any change in the maintenance of memory, irrespective of the duration of

manipulation. Consequently, long-term memory was well-established in both NpHR and GFP

groups (Figures 4C and 4F). In addition, we verified whether CF signaling is required during

the retrieval phase (Figure 4G). When mice, fully trained in the optokinetic reflex, displayed

increased gain, disrupting CF signaling did not affect the retrieval of motor memory (Figure

4H). This divergence underscores a specialized role for CF transmission in the early stages of

motor learning, with subsequent phases relying on distinct neural mechanisms. The results

highlight phase-specific contributions of CF-driven error signaling, aligning with the

hypothesis that memory consolidation and retrieval depend on downstream circuits

independent of CF activity.

Discussion

Motor learning is a finely tuned process, yet its underlying cellular mechanisms operate with remarkable precision, adapting selectively to task demands. This adaptability relies on CF activity, originating from the inferior olive, which instructs cerebellar circuits to reduce motor errors and enhance performance. CF-driven Cs in PCs have long been associated with motor learning, but how CF signals vary across learning stages remains less understood. Our study uncovers a distinct role for CF signaling that challenges traditional views: CF input is essential for driving motor adaptation during memory acquisition but surprisingly dispensable during consolidation and retrieval. This phase-specific pattern of CF transmission in cerebellum-dependent learning indicates that its role is not uniform over time, as previously thought, but instead selectively tunes cerebellar circuits to optimize adaptation.

By inhibiting CF transmission optogenetically, we demonstrated that CF activity is crucial for enhancing motor gain in the early phases of memory acquisition, where substantial adjustments and corrections are necessary. Conversely, CF input was found to have a minor influence during memory consolidation and retrieval, indicating that different mechanisms may dominate as motor memories become stable and transition to downstream circuits. This finding refines our understanding of CF function: CFs prompt learning adjustments but do not play a key role in the later phases of memory storage or recall, thereby altering our perspective on the cerebellum's role in memory formation over time.

Motor memories gained through cerebellar-dependent learning are thought to transfer to the vestibular nuclei for long-term retention (Jang et al., 2020; Seo et al., 2024; Shutoh et al., 2006). By employing optogenetic inhibition of CF transmission at various stages of learning, we identified the phase-specific role of CF signaling. CF activity was crucial for motor adaptation, such as gain increase training, during the acquisition phase but was not needed during

consolidation and retrieval. This suggests that the primary function of CF input lies in initial

memory formation rather than in subsequent memory processing.

Our results suggest that CF-dependent and CF-independent mechanisms may operate in a

context-sensitive manner during cerebellar learning, consistent with established insights

derived from the vestibulo-ocular reflex (VOR) and OKR systems. Specifically, CF signaling

appears essential in tasks involving substantial, persistent errors requiring precise error

correction. For instance, in the VOR, the increase in gain is contingent upon CF-driven

complex spikes, which provide adaptive signals in response to large error inputs (Ke et al.,

2009). Conversely, CF-independent processes can facilitate reductions in VOR gain, wherein

smaller or more gradual adjustments are adequate. This selective role of CF signaling reflects

a broader functional adaptation within cerebellar circuits, permitting the engagement of

different pathways according to the demands of motor learning. The findings related to our

OKR results correspond with this notion, emphasizing the involvement of CF-dependent

pathways in tasks characterized by substantial error correction throughout the acquisition

phase. In contrast, CF-independent mechanisms likely enable incremental adjustments of

lesser magnitude, thereby promoting efficient motor adaptation in the absence of CF input.

This delineation of responsibilities between CF-dependent and CF-independent processes

highlights the cerebellum's adaptive plasticity, wherein CF input is selectively activated

based on the magnitude and persistence of motor errors. The cerebellum maintains a balance

between precision and flexibility by dynamically alternating between these mechanisms,

enhancing motor adaptation across various behavioral contexts. This interplay offers a

versatile framework for motor learning in cerebellar-dependent tasks, encompassing VOR,

OKR, and likely other forms of sensorimotor adaptation.

In addition to their robust excitatory drive on PCs, CFs shape cerebellar computation by

recruiting molecular-layer interneurons to impose feedforward inhibition. Optogenetic

activation of molecular-layer interneurons, which silences PCs, markedly suppresses CF-

evoked dendritic calcium spikes, thereby CF transmission may dynamically gate their own

instructive signals, shifting the polarity and magnitude of synaptic plasticity at PF–PC

synapses (Rowan et al., 2018). Beyond the cerebellar cortex, anatomical tracing reveals that

the same IO neurons that innervate the flocculus also send axon collaterals to the vestibular

nuclei (Balaban et al., 1981), suggesting that CF signals can simultaneously modulate cortical

and vestibular outputs. While the behavioral consequences of this direct CF-vestibular

nucleus pathway remain to be defined, it provides a parallel channel through which IO

teaching signals may coordinate cortical plasticity with adaptive vestibular reflexes.

The process of systems consolidation, facilitated by the medial vestibular nuclei (MVN), a

downstream circuitry of floccular PCs, is essential for long-term memory storage (Shutoh et

al., 2006). Given that CF activity modulates floccular PC firing, we hypothesized that CF

signaling might indirectly influence MVN activity during memory consolidation and

retrieval. However, contrary to this expectation, our results revealed no substantial effect of

CF transmission on long-term memory formation or recall, suggesting that CF signaling does

not directly impact MVN function. Instead, previous studies suggest that PC intrinsic

plasticity supports memory consolidation (Seo et al., 2024) , while granule cell activity

contributes to memory retrieval (Wada et al., 2014). These findings reinforce the notion that

CF signaling primarily facilitates the acquisition phase of motor memory, while simple

spiking in PCs may encode timing and frequency information crucial for stabilizing memory

traces in downstream circuits. Ss could either complement or function independently of CF

transmission, especially during phases when error correction demands are minimal.

Mechanistically, suppressing CF signaling likely mimicked key aspects of long-term

depression (LTD) at CF-PC synapses, such as reducing the amplitude of Cs (Hansel &

Linden, 2000). Under normal conditions, CF input promotes parallel fiber-PC (PF-PC) LTD

by driving calcium influx during error correction, such as in response to retinal slip

(Coesmans et al., 2004). CF-LTD modifies Cs waveforms and attenuates CF-evoked

dendritic calcium transients (Weber et al., 2003). In our study, optogenetic CF suppression

likely simulated the functional consequences of CF-LTD, which may have impaired PF-PC

LTD and subsequently hindered gain enhancement during learning (Han et al., 2007).

Our findings underscore a phase-specific role for CF input, demonstrating that CF activity

is critical for motor memory acquisition but less necessary for consolidation and retrieval.

Using optogenetic inhibition, we achieved precise temporal control over CF activity within

distinct learning phases, enabling targeted manipulation of CF signaling. This differs from

prior studies that employed pharmacological or lesion methods, which affect broader

cerebellar pathways and may implicate CF signaling across multiple memory stages. The task

demands inherent to different learning phases may explain these observed differences. Tasks

requiring high-error correction, such as those encountered during the active learning phase,

likely depend on CF-driven signals for precise error correction during acquisition. In contrast,

CF-independent mechanisms may dominate later phases where such adjustments are less

critical. This context-sensitive role highlights the adaptability of CF signaling in tuning

cerebellar circuits to meet the specific demands of each learning phase. Moreover, cerebellar-

dependent memory consolidation exhibits biphasic temporal dynamics: an initial rapid phase

of consolidation followed by prolonged stabilization. Our study specifically focused on the

early consolidation phase, emphasizing cerebellar plasticity mechanisms that are critical for

encoding and stabilizing motor memory (Cooke et al., 2004; Seo et al., 2024; Steinmetz &

Freeman, 2016; Titley et al., 2007). Importantly, changes during this early period

significantly shape memory formation and retention, while alterations beyond this timeframe

have minimal impact on storage or performance. While our findings align with this

framework, extended consolidation periods may involve slower processes, such as systems-

level integration into downstream nuclei or cortical regions, contributing to memory

stabilization or generalization. Future investigations using chronic in-vivo recordings or live

imaging could elucidate these prolonged dynamics and clarify CF transmission's nuanced

contributions across phases of cerebellar-dependent memory consolidation.

In conclusion, inhibiting CF transmission during motor learning impairs gain increases

specifically during acquisition. These findings provide critical insights into CF signaling's

role in cerebellum-dependent learning while underscoring the need for further exploration of

how cerebellar circuits contribute to adaptive motor memory over extended timeframes.

320 **Key resources table**

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Virus	AAV9.CaMKIIα.EGFP	Addgene	RRID: Addgene_50469	Plasmid #50469
Virus	AAV1.CaMKIIα.eNpHR3.0.EYFP	Addgene	RRID: Addgene_26971	Plasmid #26971
Chemical compound, drug	Zoletil	Virvac		
Chemical compound, drug	Rompun	Bayer		
Chemical compound, drug	Dexamethasone	Samyang Pharmaceutical		
Chemical compound, drug	Meloxicam	Boehringer Ingelheim		
Software, algorithm	ZEN	Zeiss	RRID: SCR_013672	
Software, algorithm	Fiji	http://fiji.sc	RRID: SCR_002285	
Software, algorithm	Patchmaster	HEKA	RRID: SCR_000034	
Software, algorithm	Igor Pro	WaveMetrics	RRID: SCR_000325	
Software, algorithm	GraphPad Prism	GraphPad	RRID: SCR_002798	

Animals and Stereotaxic Surgery for Virus Injection

All experimental procedures were approved by the Seoul National University Institutional

Animal Care and Use Committee. First, wild-type mice aged 7–10 weeks were anesthetized

using intraperitoneal injections of a Zoletil/Rompun mixture (30 mg / 10 mg/kg). For

optogenetic expression in CF, the virus was injected into the IO 2–3 weeks before the

behavioral test, as previously described (Kimpo et al., 2014; Roh et al., 2020). Briefly,

bilateral injections were made at the midpoint between the edge of the occipital bone and the

C1 cervical vertebra. The glass pipette was set at a 55° angle from the vertical and 7° from

the midline. After approaching a 3 mm depth, a virus solution containing 100–200 nL of

AAV1-CaMkIIα-eNpHR3.0-EYFP or AAV9.CaMkIIα-EGFP was injected with a Picopump

at 5 nL/s. The pipettes were left in place for 10 min before they were removed to minimize

backflow.

Stereotaxic Surgery for Optic Cannula Implantation

After the mice were fully recovered (a few days to a week) from the first surgery of virus

injection, a second operation of optic cannula implantation (Ø1.25 mm Ceramic Ferrule,

L=3.0 mm) near cerebellar flocculus and head fixation was conducted. Unlike the fiber optic

cannula, head fixation parts were hand-made. Before the surgery, mice were anesthetized by

intraperitoneal injection of Zoletil 50 (Virbac, 15 mg/kg) and xylazine (Rompun, Bayer, 15

mg/kg) mixture. Mice were given 48 hours to recover after surgery.

Confirmation Viral Expression.

After completion of all the experiments, sampling using cardiac perfusion is performed. The

brain was then extracted and coronal sections were made at 30 μm intervals. Images were

acquired and processed using a confocal microscope (Zeiss LSM 7 MP, Carl Zeiss, Jena,

Germany) and Zen software (Zeiss). The location of virus expression was confirmed by

comparison with the brain atlas (Paxinos and Franklin's The Mouse Brain in Stereotaxic

Coordinates 4th Edition by George Paxinos).

Optical suppression of CF terminals in the cerebellar cortex in vivo

To manipulate CF transmission-induced Cs in PCs, AAV1.CaMKIIα.eNpHR3.0.eYFP or

AAV8.CaMKIIa.eGFP was bilaterally injected into the IO. A glass micro-pipette was used to

inject 200 nL per site. To optically suppress the CF terminals and record CF-PC synaptic

response, a small cranial window was made above the cerebellar cortex lobule IV/V a week

after the viral injection. After 2 weeks from the viral injection, mice underwent single-unit

recordings. The recordings were performed as described in Neural data acquisition and

analysis. Cs responses were acquired during 10 seconds of yellow on and off sessions 20

times to test the effect of CF terminal suppression on CF-PC synaptic transmission.

Neural data acquisition and analysis

Single-unit recordings were performed while mice were head-fixed and awake in the

recording rack. A Digital Lynx system (Neuralynx, Bozeman, MT) was used to amplify, band

pass filter (0.1–8000 Hz for simple spikes, 10–200 Hz for complex spikes), and digitize the

electrode recordings. A silicon neural probe (Cambridge NeuroTech) was used to probe and

acquire Cs signals. Cs signals were collected at a 32 kHz sampling rate. Cs waveforms were

manually selected based on visual inspection of the averaged appearance of characteristic

negative Ca^{2+} peaks.

Slice preparation, electrophysiology, and optogenetic validation

An acute brain slice preparation and electrophysiological experiments were carried out as

previously described (Lee et al., 2024). First, mice aged 9–10 weeks with

AAV1.CaMKIIα.eNpHR 3.0.WPRE-EYFP injection into their inferior olivary neuron 2–3

weeks before the preparation were anesthetized by isoflurane and briefly decapitated. Then,

250 μm-thick sagittal slices of the cerebellar vermis were obtained from the mice using a

vibratome (VT1200S, Leica). The ice-cold cutting solution contained 75 mM sucrose, 75 mM

NaCl, 2.5 mM KCl, 7 mM MgCl2, 0.5 mM CaCl2, 1.25 mM NaH2PO4, 26 mM NaHCO3,

and 25 mM glucose with bubbled 95% O2 and 5% CO2. The slices were immediately

transferred into the artificial cerebrospinal fluid (ACSF) containing 125 mM NaCl, 2.5 mM

KCl, 1 mM MgCl2, 2 mM CaCl2, 1.25 mM NaH2PO4, 26 mM NaHCO3, and 10 mM

glucose with bubbled 95% O2 and 5% CO2. They were allowed to recover at 32 °C for 30

min and at room temperature for 1 hour. All recordings were performed within 8 hr of

recovery.

Brain slices were placed in a submerged chamber with perfusion of ACSF for at least 10

min before recording. We used OptoPatcher (HEKA Elektronik) holding recording pipette

(2–3 MΩ) filled with internal solution containing 9 mM KCl, 10 mM KOH, 120 mM K-

gluconate, 3.48 mM MgCl2, 10 mM HEPES, 4 mM NaCl, 4 mM Na2ATP, 0.4 mM Na3GTP,

and 17.5 mM sucrose (pH 7.25). The stimulation electrode was placed on the granule cell

layer near the PC. The stimulation isolator injected a brief current pulse to the stimulation

electrode and was controlled using PatchMaster software (HEKA Elektronik). Stimulation

intensity (6–30 μA) was calibrated to minimally evoke CF EPSC. Regarding the laser on

trial, a 593.5 nm irradiation from a yellow laser was applied 5 s before CF stimulation. The

eEPSC amplitude was averaged over three lasers on and off trials. Under current clamp

mode, complex spikes induced by CF stimulation were recorded using the same validation

protocol. Overall, the laser illumination (3-5 mW) consistently inhibited CF transmission.

Electrophysiological data were acquired using an EPC10 patch-clamp amplifier (HEKA

Elektronik) and PatchMaster software (HEKA Elektronik) with a sampling frequency of 20

kHz, and the signals were filtered at 2 kHz. All electrophysiological recordings were acquired

from the central cerebellar vermis. The amplitude of EPSC was analyzed using Igor Pro

(WaveMetrics).

Optokinetic reflex behavior test

First, two sessions of acclimation were conducted. During each session, the mouse was restrained using a custom-made restrainer for 20 min both with and without light for habituation to the recording environment. Following the procedure reported by (Stahl et al., 2000), a calibration was performed. This allowed for the conversion of the dynamics of pupil-to-eye rotation. Next, three basal oculomotor performances including (OKR), vestibulo-ocular reflex in the dark (dVOR), and vestibulo-ocular reflex in the light (lVOR) were measured. During the recording of eye movements, it was necessary to control pupil dilation that was induced by an absence of light in dark conditions. Thus, physostigmine salicylate solution (Eserine; Sigma Millipore) was given to mice under brief isoflurane anesthesia. The concentration of eserine solution was increased from 0.1%, 0.15%, and 0.2% based on the pupil size. Basal oculomotor performances were described in gains. For the actual learning protocol, we adopted OKR and it consisted of five training sessions (10 min per session), six checkup points, and 24 hr of consolidation period. During the training sessions, the mice were visually stimulated with a sinusoidally rotating drum (±5 degrees). Completely trained mice were placed back into their home cage, which was stored in the dark condition until the last gain checkup points.

Gain Analysis

Gains acquired from basal oculomotor performances and OKR learning were calculated as the ratio of evoked eye movements to the movement of the screen or turn table as visual or vestibular stimuli, respectively. A custom-built LabView (National Instrument) analysis tool was used for all the calculations

Statistical analysis

Graph plotting and statistical analysis were performed using GraphPad Prism (GraphPad

Software Inc, CA, USA). The hypothesis was tested by using paired and unpaired t-tests

between sample pairs using GraphPad Prism. Statistical significance was set at $p < 0.05$.

Asterisks denoted in the graph indicate the statistical significance. * indicates $p < 0.05$, ** <

0.01, *** < 0.001, and **** < 0.0001. The test name and statistical values are presented in

each figure legend.

Acknowledgments

This work was supported by National Research Foundation of Korea (NRF) grants funded by the Korean Government (MSIT) (NRF-2018R1A5A2025964 and 2022M3E5E8017970 to Sang Jeong Kim).

Competing interest

The authors declare that they have no conflicts of interest.

Figure 1. Optogenetic inhibition of climbing fiber transmission derived from the

inferior olive.

A. Schematic diagram showing virus injection for optogenetic manipulation of CF

transmission.

B. Bi. A virus is expressed specifically only in the IO region (IO, inferior olive; py,

pyramidal tract; ml, medial lemniscus). Scale bar, 100 μm. Bii. Virus expression along

CF in the cerebellar flocculus region. (GCL, granule cell layer; PCL, Purkinje cell layer;

ML, molecular layer; FL, flocculus). Scale bar, 100 μm.

C. A scheme of optogenetic suppression using a yellow laser (593 nm, 3–5 mW) while

recording CF EPSCs under voltage clamp mode.

D. Left, representative traces of CF EPSCs of the GFP (top) and NpHR group (bottom) with

(dark line) and without (light line) opto-stimulation. Right, Quantitative analysis of

suppression of CF EPSCs (GFP: n = 10 cells/2 mice, NpHR: n = 9 cells/5 mice, *** $p <$

0.001, Unpaired t-test).

E. A scheme of optogenetic suppression while recording complex spikes under current

clamp mode.

F. Representative traces of complex spike of the GFP (top) and NpHR group (bottom) with

(dark line) and without (light line) opto-stimulation. Quantitative analysis of number of

complex spike spikelets (GFP: n = 8 cells/1 mouse, NpHR: n = 8 cells/5 mice, ** $p =$

0.002; Paired t-test).

Figure 2. Optogenetic inhibition of climbing fiber transmission blocked Cs firing but not Ss firing in vivo

A. Schematic diagram of virus injection into IO and in vivo recording during optogenetic manipulation of CF transmission. Arrow heads indicate NpHR-expressing CF terminals in the molecular layer of the cerebellar cortex. Scale bar, 50 μm

B. Representative electrophysiological recording traces of PC firing from the GFP and NpHR groups, without (OFF) and with (ON) yellow (593 nm) opto-stimulation. The upper plots show 10-second traces without and with opto-stimulation, and the lower plots display a 1-second segment from the respective upper traces. Asterisks denote Cs

C. Comparison of Ss firing rates during optogenetic inhibition. For the GFP group, firing rates with laser off (n = 11 cells from 4 mice) versus on (n = 11 cells from 4 mice) were not significantly different (Mann-Whitney test, $P = 0.8594$). Similarly, for the NpHR

group, no significant difference was observed between OFF (n = 10 cells from 6 mice)

and ON (n = 10 cells from 6 mice) conditions (Mann-Whitney test, p > 0.9999).

D. Comparison of Cs firing rates during optogenetic inhibition. In the GFP group, no

significant difference was observed between OFF (n = 8 cells from 4 mice) and ON (n = 8

cells from 4 mice) conditions (Mann-Whitney test, p = 0.4563). In the NpHR group, Cs

firing rates were significantly reduced during opto-stimulation (OFF, n = 6 cells from 6

mice; ON, n = 6 cells from 6 mice; Mann-Whitney test, ** p = 0.0022). Of note, 6 out of

8 recorded cells were responsive to optogenetic suppression.

Figure 3. Inhibition of climbing fiber transmission during memory acquisition of OKR.

A. Illustration of the optokinetic reflex behavioral test (left) and experimental scheme (right).

For training, visual stimulation was given by constant rotation of the screen, while the

table, where a mouse's head is fixed, remained stationary. For day 1, a mouse was trained

for a total of 50 minutes, and on day 2, an amount of sustained memory was validated via

gain check. Scale bar, 100 μm

B. Representative traces of the screen (black sinusoidal curve) and eye movements (grey

sinusoidal curve) divided into the GFP (i) and NpHR (ii) groups. Top traces (pre) were

acquired before learning, and bottom traces (post) were obtained after 50 min of learning.

The vertical scale bar represents 10 degrees per second and the horizontal scale bar

represents 0.5 seconds.

C. The learning curve from 0 min to 50 min, and +24 hr period. Change of gain is indicated

at each time point (+10 min increment). Yellow boxes indicate opto-stimulation (12.5

mW for 10 min/session, total of 5 sessions). The left graph represents the GFP (n = 8) and

the right graph indicates the NpHR (n = 9) group.

D. Comparison of gain changes between the GFP and NpHR groups (GFP, n = 8 mice;

NpHR, n = 9 mice). Percentages of gain increment from 0 min to 50 min were calculated.

The change of gain of the GFP group was significantly larger than that of the NpHR

group (left; Unpaired t-test, p = 0.0007). The gain of the GFP group was significantly

increased from 0 min to 50 min (right; Paired t-test, **** p < 0.0001), while the NpHR

group showed no significant improvement of gain from 0 min to 50 min (right; Paired t-

test, p = 0.1046).

Figure 4. Inhibition of climbing fiber transmission during memory consolidation and

retrieval phase.

A. Illustration of the experimental scheme. 10 min of optokinetic reflex training was

repeated 5 times, and then opto-stimulation (12.5 mW) was given at 0 min post-learning

for 30 min.

B. The learning curve from 0 min to 50 min, and +24 hr period. Change of gain is indicated

at each time point (+10 min increment). A yellow box indicates opto-stimulation at 0 min

post-learning. The left graph represents the GFP and the right graph indicates the NpHR

group.

C. Comparison of consolidation percentage between the GFP and NpHR groups (GFP group,

n = 7 mice; NpHR group, n = 9 mice). The percentage of sustained memory after 24 hr

was calculated. No difference was observed between the groups (p = 0.3173; Unpaired t-

test).

D. Illustration of the experimental scheme. 10 min of optokinetic reflex training was

repeated 5 times, and then opto-stimulation (12.5 mW) was given at 0 min post-learning

for 6 hr.

E. The learning curve from 0 min to 50 min, and +24 hr period. Change of gain is indicated

at each time point (+10 min increment). A yellow box indicates opto-stimulation at 0 min

post-learning. The left graph represents the GFP group, and the right graph indicates the

NpHR group.

F. Comparison of gain changes between GFP and NpHR groups (GFP group, n = 6 mice;

NpHR group, n = 6 mice). Percentages of gain increment from 0 min to 50 min were

calculated. No difference was observed between the groups (Unpaired t-test, p = 0.6405).

G. Illustration of the experimental scheme. Following proper motor learning on day 1, the

remaining gain value was checked on day 2, after testing 1st gain retrieval without opto-

stimulation, 2^{nd} gain retrieval was tested with opto-stimulation to validate the effect of CF

inhibition on gain retrieval (opto-stimulation was given 3 times for 24 seconds each at

12.5 mW).

H. Gain retrievals before and during opto-stimulation were compared. The left graph

represents the GFP group, and the right shows the NpHR group (GFP group, n = 6 mice;

NpHR group, n = 8 mice). There was no significant variation between the groups (NpHR

group, p = 0.2439; GFP group, p = 0.3180; Paired t-test).

References

Albus, J. S. (1971). A theory of cerebellar function. *Mathematical Biosciences*, *10*(1), 25-61.
https://doi.org/https://doi.org/10.1016/0025-5564(71)90051-4

Attwell, P. J. E., Cooke, S. F., & Yeo, C. H. (2002). Cerebellar function in consolidation of a motor
memory. *Neuron*, *34*(6), 1011-1020. https://doi.org/Doi 10.1016/S0896-6273(02)00719-5

Balaban, C. D., Kawaguchi, Y., & Watanabe, E. (1981). Evidence of a Collateralized Climbing Fiber
Projection from the Inferior Olive to the Flocculus and Vestibular Nuclei in Rabbits.
Neuroscience Letters, *22*(1), 23-29. https://doi.org/Doi 10.1016/0304-3940(81)90279-2

Bloedel, J. R., & Bracha, V. (1998). Current concepts of climbing fiber function. *Anatomical Record*,
253(4), 118-126. <Go to ISI>://WOS:000075784300008

Cahill, H., & Nathans, J. (2008). The Optokinetic Reflex as a Tool for Quantitative Analyses of
Nervous System Function in Mice: Application to Genetic and Drug-Induced Variation. *Plos*
One, *3*(4). https://doi.org/ARTN e2055

10.1371/journal.pone.0002055

Clopath, C., Badura, A., De Zeeuw, C. I., & Brunel, N. (2014). A Cerebellar Learning Model of
Vestibulo-Ocular Reflex Adaptation in Wild-Type and Mutant Mice. *Journal of Neuroscience*,
34(21), 7203-7215. https://doi.org/10.1523/Jneurosci.2791-13.2014

Coesmans, M., Weber, J. T., De Zeeuw, C. I., & Hansel, C. (2004). Bidirectional parallel fiber plasticity
in the cerebellum under climbing fiber control. *Neuron*, *44*(4), 691-700. https://doi.org/DOI
10.1016/j.neuron.2004.10.031

Cooke, S. F., Attwell, P. J. E., & Yeo, C. H. (2004). Temporal properties of cerebellar-dependent
memory consolidation. *Journal of Neuroscience*, *24*(12), 2934-2941.
https://doi.org/10.1523/Jneurosci.5505-03.2004

Eccles, J. C., Llinas, R., & Sasaki, K. (1966). The excitatory synaptic action of climbing fibres on the
Purkinje cells of the cerebellum. *J Physiol*, *182*(2), 268-296.
https://doi.org/10.1113/jphysiol.1966.sp007824

Frens, M. A., Mathoera, A. L., & van der Steen, J. (2001). Floccular complex spike response to
transparent retinal slip. *Neuron*, *30*(3), 795-801. https://doi.org/Doi 10.1016/S0896-
6273(01)00321-X

Goossens, H. H. L. M., Hoebeek, F. E., van Alphen, A. M., van der Steen, J., Stahl, J. S., De Zeeuw,
C. I., & Frens, M. A. (2004). Simple spike and complex spike activity of floccular Purkinje cells
during the optokinetic reflex in mice lacking cerebellar long-term depression (vol 19, pg 687,
2004). *European Journal of Neuroscience*, *19*(6), 1673-1673. https://doi.org/10.1111/j.1460-
9568.2004.03351.x

Graf, W., Simpson, J. I., & Leonard, C. S. (1988). Spatial-Organization of Visual Messages of the
Rabbits Cerebellar Flocculus .2. Complex and Simple Spike Responses of Purkinje-Cells.
Journal of Neurophysiology, *60*(6), 2091-2121. https://doi.org/DOI 10.1152/jn.1988.60.6.2091

Han, V. Z., Zhang, Y. P., Bell, C. C., & Hansel, C. (2007). Synaptic plasticity and calcium signaling in

Purkinje cells of the central cerebellar lobes of mormyrid fish. *Journal of Neuroscience*, *27*(49),
13499-13512. https://doi.org/10.1523/Jneurosci.2613-07.2007

Hansel, C., & Linden, D. J. (2000). Long-term depression of the cerebellar climbing fiber-Purkinje
neuron synapse. *Neuron*, *26*(2), 473-482. https://doi.org/Doi 10.1016/S0896-6273(00)81179-4

Ito, M. (1972). Neural design of the cerebellar motor control system. *Brain Res*, *40*(1), 81-84.
https://doi.org/10.1016/0006-8993(72)90110-2

Jang, D. C., Shim, H. G., & Kim, S. J. (2020). Intrinsic Plasticity of Cerebellar Purkinje Cells
Contributes to Motor Memory Consolidation. *Journal of Neuroscience*, *40*(21), 4145-4157.
https://doi.org/10.1523/Jneurosci.1651-19.2020

Ke, M. C., Guo, C. C., & Raymond, J. L. (2009). Elimination of climbing fiber instructive signals during
motor learning. *Nature Neuroscience*, *12*(9), 1171-U1123. https://doi.org/10.1038/nn.2366

Kim, O. A., Ohmae, S., & Medina, J. F. (2020). A cerebello-olivary signal for negative prediction error
is sufficient to cause extinction of associative motor learning. *Nature Neuroscience*, *23*(12),
1550-U1517. https://doi.org/10.1038/s41593-020-00732-1

Kimpo, R. R., Rinaldi, J. M., Kim, C. K., Payne, H. L., & Raymond, J. L. (2014). Gating of neural error
signals during motor learning. *Elife*, *3*. https://doi.org/ARTN e02076

10.7554/eLife.02076

Lee, J., Kim, S. H., Jang, D. C., Jang, M., Bak, M. S., Shim, H. G., Lee, Y. S., & Kim, S. J. (2024).
Intrinsic plasticity of Purkinje cell serves homeostatic regulation of fear memory. *Mol*
Psychiatry, *29*(2), 247-256. https://doi.org/10.1038/s41380-023-02320-8

Marr, D. (1969). A Theory of Cerebellar Cortex. *Journal of Physiology-London*, *202*(2), 437-&.
https://doi.org/DOI 10.1113/jphysiol.1969.sp008820

Medina, J. F., Nores, W. L., & Mauk, M. D. (2002). Inhibition of climbing fibres is a signal for the
extinction of conditioned eyelid responses. *Nature*, *416*(6878), 330-333. https://doi.org/DOI
10.1038/416330a

Palmer, L. M., Clark, B. A., Grundemann, J., Roth, A., Stuart, G. J., & Hausser, M. (2010). Initiation of
simple and complex spikes in cerebellar Purkinje cells. *Journal of Physiology-London*,
588(10), 1709-1717. https://doi.org/10.1113/jphysiol.2010.188300

Pham, N. C., Kim, Y. G., Kim, S. J., & Kim, C. H. (2020). Differential effects of inferior olive lesion on
vestibulo-ocular and optokinetic motor learning. *Neuroreport*, *31*(1), 9-16.
https://doi.org/10.1097/Wnr.0000000000001333

Roh, S. E., Kim, S. H., Ryu, C., Kim, C. E., Kim, Y. G., Worley, P. F., Kim, S. K., & Kim, S. J. (2020).
Direct translation of climbing fiber burst-mediated sensory coding into post-synaptic Purkinje
cell dendritic calcium. *Elife*, *9*. https://doi.org/ARTN e61593

10.7554/eLife.61593

Rowan, M. J. M., Bonnan, A., Zhang, K., Amat, S. B., Kikuchi, C., Taniguchi, H., Augustine, G. J., &
Christie, J. M. (2018). Graded Control of Climbing-Fiber-Mediated Plasticity and Learning by
Inhibition in the Cerebellum. *Neuron*, *99*(5), 999-+.

https://doi.org/10.1016/j.neuron.2018.07.024

Schweigart, G., Mergner, T., Evdokimidis, I., Morand, S., & Becker, W. (1997). Gaze stabilization by
optokinetic reflex (OKR) and vestibulo-ocular reflex (VOR) during active head rotation in man.
Vision Research, 37(12), 1643-1652. https://doi.org/Doi 10.1016/S0042-6989(96)00315-X

Seo, J., Kim, Y. G., Lee, Y.-S., & Kim, S. J. (2024). Cerebellar systems consolidation driven by the
temporal dynamics of Purkinje cell excitability. *bioRxiv*, 2024.2007.2024.604888.
https://doi.org/10.1101/2024.07.24.604888

Shaikh, A. G., Wong, A. L., Optican, L. M., & Zee, D. S. (2017). Impaired Motor Learning in a Disorder
of the Inferior Olive: Is the Cerebellum Confused? *Cerebellum, 16*(1), 158-167.
https://doi.org/10.1007/s12311-016-0785-x

Shutoh, F., Ohki, M., Kitazawa, H., Itohara, S., & Nagao, S. (2006). Memory trace of motor learning
shifts transsynaptically from cerebellar cortex to nuclei for consolidation. *Neuroscience,*
139(2), 767-777. https://doi.org/10.1016/j.neuroscience.2005.12.035

Stahl, J. S., van Alphen, A. M., & De Zeeuw, C. I. (2000). A comparison of video and magnetic search
coil recordings of mouse eye movements. *J Neurosci Methods, 99*(1-2), 101-110.
https://doi.org/10.1016/s0165-0270(00)00218-1

Steinmetz, A. B., & Freeman, J. H. (2016). Cannabinoid modulation of memory consolidation within
the cerebellum. *Neurobiol Learn Mem, 136*, 228-235.
https://doi.org/10.1016/j.nlm.2016.11.002

Titley, H. K., Heskin-Sweezie, R., Chung, J. Y. J., Kassardjian, C. D., Razik, F., & Broussard, D. M.
(2007). Rapid consolidation of motor memory in the vestibuloocular reflex. *Journal of*
Neurophysiology, 98(6), 3809-3812. https://doi.org/10.1152/jn.01056.2007

Van Der Giessen, R. S., Koekkoek, S. K., van Dorp, S., De Gruijl, J. R., Cupido, A., Khosrovani, S.,
Dortland, B., Wellershaus, K., Degen, J., Deuchars, J., Fuchs, E. C., Monyer, H., Willecke, K.,
De Jeu, M. T. G., & De Zeeuw, C. I. (2008). Role of olivary electrical coupling in cerebellar
motor learning. *Neuron, 58*(4), 599-612. https://doi.org/10.1016/j.neuron.2008.03.016

Wada, N., Funabiki, K., & Nakanishi, S. (2014). Role of granule-cell transmission in memory trace of
cerebellum-dependent optokinetic motor learning. *Proceedings of the National Academy of*
Sciences of the United States of America, 111(14), 5373-5378.
https://doi.org/10.1073/pnas.1402546111

Weber, J. T., De Zeeuw, C. I., Linden, D. J., & Hansel, C. (2003). Long-term depression of climbing
fiber-evoked calcium transients in Purkinje cell dendrites. *Proceedings of the National*
Academy of Sciences of the United States of America, 100(5), 2878-2883.
https://doi.org/10.1073/pnas.0536420100

Welsh, J. P., & Harvey, J. A. (1998). Acute inactivation of the inferior olive blocks associative learning.
European Journal of Neuroscience, 10(11), 3321-3332. https://doi.org/DOI 10.1046/j.1460-
9568.1998.00400.x

Yang, Y., & Lisberger, S. G. (2014). Purkinje-cell plasticity and cerebellar motor learning are graded
by complex-spike duration. *Nature, 510*(7506), 529-+. https://doi.org/10.1038/nature13282

Zang, Y. L., & De Schutter, E. (2019). Climbing Fibers Provide Graded Error Signals in Cerebellar

654   Learning. *Frontiers in Systems Neuroscience*, *13*. https://doi.org/ARTN 46

655   10.3389/fnsys.2019.00046

656