1 **The value of livestock abortion surveillance in Tanzania: identifying disease priorities**

2 **and informing interventions**

3 Lankester F.[1,2,*], Kibona T.[2], Allan K.J.[3], de Glanville W.A.[3], Buza J.J.[4], Katzer F.[5], Halliday

4 J.E.B.[3], Mmbaga BT.[6], Wheelhouse N.[7], Innes E.A.[5], Thomas K.M.[8], Nyasebwa O.M.[9], Swai

5 E.S.[9], Claxton J.R.[3], and Cleaveland S.[3]

6
7 **Affiliations**

8
[1.] Paul G. Allen School for Global Health, Washington State University, Pullman,
Washington, USA
[2.] Global Animal Health Tanzania, Arusha, Tanzania
[3.] School of Biodiversity, One Health, and Veterinary Medicine, College of Medical,
Veterinary and Life Sciences, University of Glasgow, Glasgow, UK
[4.] Nelson Mandela African Institution of Science and Technology, Arusha, Tanzania
[5.] Moredun Research Institute, Pentlands Science Park, Edinburgh, UK
[6.] Kilimanjaro Clinical Research Institute, Moshi, Tanzania
[7.] School of Applied Sciences, Edinburgh Napier University, Edinburgh, UK
[8.] Centre for International Health, University of Otago, Dunedin, New Zealand
[9.] Ministry of Livestock and Fisheries, Dodoma, Tanzania

20
21 *Corresponding author: F. Lankester – felix.lankester@wsu.edu
22

Abstract

Background: Lack of reliable data on the aetiology of livestock diseases, especially in Africa, is a major factor constraining the design of effective livestock health interventions to improve livelihoods, food security and public health.

Rationale: Livestock abortion is an important disease syndrome that affects productivity and livestock economies, and poses risks to public health. Worldwide, several pathogens are associated with livestock abortions but across Africa livestock disease surveillance data rarely include information from abortion events, little is known about the aetiology and impacts of livestock abortions, and data are not available to inform prioritisation of disease interventions.

Methods: This paper describes outcomes from a surveillance platform that was set up in 15 wards in northern Tanzania spanning pastoral, agropastoral and smallholder agro-ecological systems to investigate the causes and impacts of livestock abortion. Abortion cases were reported by farmers to Ministry of Livestock and Fisheries livestock field officers and on to investigation teams. The abortion was eligible for inclusion if the research field team or livestock field officer could attend within 72 hours of the event occurring. If so, a field investigation was carried out to collect diagnostic samples from aborted materials and aborting dams and questionnaire data on herd/flock management. Laboratory diagnostic tests were carried out for a range of abortigenic pathogens, and laboratory and questionnaire data analysed to investigate a) determinants of attribution; b) patterns of investigated events, including species and breed affected, history of previous abortion and recent stressful events, and the seasonality of cases; c) determinants of reporting, investigation and attribution; (d) cases in which zoonotic pathogens were detection.

Results: Over a two-year period (2017-2019), 215 abortion events in cattle (n=71), sheep (n=44) and goats (n=100) were investigated from 13 of the 15 wards. The number of investigated cases varied widely across wards, with three of the 15 field officers (20%) reporting 70% of investigated cases. Aetiological attribution, achieved for 19.5% of cases through PCR-based diagnostics, was significantly affected by delays in field investigation. Histopathology proved less useful than PCR diagnostics due to rapid deterioration of field samples. Vaginal swabs from aborting dams provided a practical and sensitive source of diagnostic material for pathogen detection.

Discussion / Conclusion: Livestock abortion surveillance, even at a small scale and when
capturing only a small proportion of events, can generate valuable information on causes of
zoonotic disease outbreaks, livestock reproductive losses and can identify important
pathogens that are not easily captured through other forms of livestock disease surveillance.
The study demonstrated the feasibility of establishing an effective reporting and investigation
system that could be implemented across a range of settings, including remote rural areas,
with engagement of community-based field officers, and establishment of practical sample
collection protocols and molecular diagnostic platforms all being important elements of
effectiveness.

64
65

Introduction

Livestock reproductive losses, including spontaneous abortion, are a major concern for the livestock industry worldwide, resulting in significant economic loss and posing a threat to public health. The impacts of livestock abortion on the world's poorest livestock-keepers, who are heavily dependent on livestock for food security and livelihoods (Food and Agriculture Organization 2011), is likely to be substantial. For these families, the loss of a livestock foetus and a subsequent decline in milk production reduces the availability of a high-quality food source (milk) that can be essential for childhood growth and cognitive development (Neumann et al. 2003). In addition, livestock reproductive losses reduce income (from sales of meat and milk), and results in a loss of livestock assets that are a critical store of wealth, collateral or a safety net in times of need (The World Bank 2021). Recent analyses have shown that because of the need for increased spending on animal management, livestock abortions can also have indirect negative impacts on household expenditure and education (Ahmed et al. 2019). Livestock abortion can also pose a direct threat to human health because many abortigenic agents are also zoonotic (Givens and Marley 2008; Thomas et al. 2022).

Surveillance is defined as the real-time (or near real-time) collection, analysis, interpretation, and dissemination of health-related data to enable the early identification of the impact (or absence of impact) of potential health threats, which require effective action (Hulth 2014). Effective livestock health surveillance provides critical data for evidence-based approaches to livestock disease control and management but this requires reliable, high-quality, and timely data that can be drawn from multiple sources (George et al. 2021). Over the past decade, increasing attention has been given to animal health syndromic surveillance (Dórea and Vial 2016), which relies on detection of health indicators, such as livestock abortion, that are discernible before a confirmed diagnosis is made. However, systematic reviews of the literature that collectively cover a period from 2000 to 2016 indicate that syndromic surveillance programmes have mostly been implemented in Europe, North America or Australasia with only a single pilot project identified in Africa (Dórea and Vial 2016; Dorea, Sanchez, and Revie 2011).

In public health, event-based surveillance has also been gaining attention for the early detection of unusual events that might signal acute and emerging human health risks (World

Health Organization 2008; Africa Centres for Disease Control and Prevention 2018). This
involves the collection, analysis and interpretation of information through both formal and
informal channels to rapidly identify unusual or unexpected health events, such as disease
outbreaks, emerging infectious diseases, or other public health threats. Although the WHO
guidelines for event-based surveillance (World Health Organization 2008) make explicit
reference to the capture of information around unusual disease events in animals, the
development of integrated early-warning systems involving animal disease events is still very
limited. Livestock surveillance has a clear potential for identifying and preventing outbreaks
of zoonotic and emerging diseases; a substantial proportion (39.4%) of livestock pathogens
infect humans (Cleaveland et al. 2001), and ungulates harbour more zoonotic pathogens than
any other taxonomic group (Woolhouse and Gowtage-Sequeria 2005). With several
abortigenic agents also causing emerging livestock diseases, for example those caused by
Schmallenberg virus, bluetongue virus and porcine reproductive and respiratory syndrome
virus, livestock abortion may represent a particularly important syndromic target for zoonotic
and emerging disease surveillance. However, there is limited information available on current
practices, effectiveness, and challenges of livestock abortion surveillance, particularly in low-
and middle-income countries.

Livestock abortion has many causes, including infectious, nutritional, physical and genetic
factors and the diversity of agents makes abortion diagnosis complex and challenging (Wolf-
Jäckel et al. 2020). Challenges relate to timely collection of diagnostic samples, sample
availability and deterioration, biases in detectability of agents, as well as complexities around
establishment of a causal aetiology, particularly for pathogens such as *Coxiella burnetii* and
Escherichia coli, which are commonly present in livestock samples (Thomas et al. 2022; Wolf-
Jäckel et al. 2020). As a result, aetiological attribution of livestock abortion rarely exceeds
35%, even in well-resourced industrialised livestock systems (Cabell 2007; Wolf-Jäckel et al.
2020) and data are particularly sparse in African livestock systems.

A recent prospective study of the aetiologies of livestock abortion, carried out in northern
Tanzania (Thomas et al. 2022), investigated 215 cases of livestock abortion of which an
attribution was made in 41 cases (19%). The infectious causes of abortion were identified as
Rift Valley fever virus (RVFV) in 14 cases (6.6%), followed by *Neospora caninum* in ten (4.7%),

pestiviruses in nine (4.2%), *Coxiella burnetii* in six (2.8%), and *Brucella* sp., and *Toxoplasma*

gondii each in one case (0.5%). Our study draws on the operational data generated from the

Thomas et al. (2022) aetiological study to examine characteristics of reporting and

investigation of cases of livestock abortion. While this study was not designed as a systematic

or comprehensive evaluation of a surveillance system, we present data on several key

attributes (ECDC 2014) of the platform, including simplicity, data quality, representativeness,

timeliness and usefulness. We discuss our findings in relation to the feasibility, practicality,

and value of establishing a livestock abortion surveillance framework to support evidence-

based interventions to improve livestock development, livelihoods, and human health in

Africa.

Methods

Logic model

A logic model was created to provide a conceptual framework that described the logical links

between the main activities, outputs, and outcomes that were expected from the programme

(Figure 1). The model depicts the overarching assumption that building better livestock

abortion surveillance systems and strong community partnerships will lead to data-driven

prevention actions and changes in knowledge, attitudes, behaviours, or practices that could

improve livestock productivity, livelihoods, and health.

Figure 1. A logic model illustrating the conceptual links between the inputs, activities, outputs, and short to long term impacts expected from effective livestock health surveillance with a particular focus on abortion.

Abortion surveillance platform

The abortion surveillance platform was set up in northern Tanzania through a collaboration between the Ministry of Livestock and Fisheries, local government authorities, and the research team. The study was undertaken from October 2017 through September 2019 in 15 wards of five districts of Arusha, Kilimanjaro, and Manyara Regions in northern Tanzania (Figure 2). These study wards were selected from randomly selected wards included in earlier cross-sectional exposure studies (Bodenham et al. 2021). Seventeen wards were selected at random and two additional wards were selected purposively because of strong existing relationships with the livestock-keeping community (Thomas et al. 2022). These 15 wards comprised five wards that were expected to be predominantly pastoral, three were expected to be predominantly agropastoral and seven expected to predominantly smallholder, with categories assigned by the research team following discussion with local experts (typically the district level veterinary officer) (Bodenham et al. 2021). Following recruitment into the study,

169 livestock field officers (LFOs) associated with each target ward received training on the causes
170 and safe investigation of livestock abortion. Livestock field officers are government
171 employees that are equivalent to para-veterinarians in other settings. They were requested
172 to ask livestock owners to report any incidents of livestock abortion, stillbirths, and peri-natal
173 death (hereafter referred to as abortion cases).

174
175

Figure 2: Map of the study area in northern Tanzania showing selected pastoral, agropastoral
and smallholder wards in Kilimanjaro, Arusha and Manyara Regions. The number of
investigated cases per ward are shown and the study base (Moshi) is shown.

Investigation of cases

Cases were investigated if, following a report from the LFO, the event could be followed up
within 72 hours of the abortion occurring. Full details of sample collection are provided in
Thomas et al. (2022). Briefly, where available, blood, milk and vaginal swabs were collected
from the aborting dam and tissue and swab samples collected from the foetus and placental
membranes. Information was collected about the abortion event and a household
questionnaire conducted to collect information on livestock demographics, livestock abortion
history, the aborting dam (age, breed), household livestock parturition practices and
household socio-economic data (Supplementary Material 1). Questionnaire data were used

to investigate underlying patterns of abortion, risks associated with abortion cases, and

operational aspects of the surveillance platform. These data were not collected from abortion

cases that were not investigated by the research team.

LFOs were instructed to provide farmers with advice as to locally appropriate preventive

measures that could be taken to reduce transmission or contamination risks associated with

abortion cases, which included safe removal of abortion tissues from livestock-occupied areas

(e.g. burning, burying or covering the tissues in thorny branches) (Supplementary Materials

2). Diagnostic results were reported back to LFOs and livestock owners within 10 days of the

investigation and, where pathogens were detected, more specific advice provided as to

management strategies that could minimise further transmission to livestock and people.

Event data were collected using a paper-based Cardiff Teleform system (Cardiff Inc., Vista,

Ca., USA) into an Access database (Microsoft Corp, Va., USA). Household questionnaire data

were collected using handheld digital devices programmed with the Open Data Kit™ survey

tool. Data were imported into R (R Core Team 2023) for cleaning, coding and analysis. The

survey instruments were pre-tested in wards that were not targets for this study. Geographic

co-ordinates from a central point within the household were collected with a handheld GPS

(Garmin eTrex™).

Sample analysis

Laboratory diagnostic analyses have previously been described in detail (Thomas et al. 2022).

Briefly, samples were tested for: (a) *Brucella* spp., *Coxiella burnetii*, *Chlamydia* spp., and

Leptospira spp. using quantitative polymerase chain reaction (qPCR) assays: (b) *Neospora* spp.

and *Toxoplasma gondii* using conventional nested PCR assays; (c) Bluetongue virus (BTV),

bovine viral diarrhoea virus / border disease virus (BVDV/BDV), and Rift Valley fever virus

(RVFV) using quantitative reverse transcriptase PCR (RT-qPCR); and (d) additional tests for *C.*

burnetii and *Neospora* spp. using immunohistochemistry. Additionally, serum was tested for

antibodies to *Brucella* spp., bovine herpesvirus (BHV-1), BVDV, *C. burnetii, Leptospira hardjo*,

and RVFV.

Summarising the investigated cases and description of the livestock study population

The number of abortion cases was recorded for each agro-ecological zone (pastoral, agro-pastoral and small holder) and study ward, together with the number of households that had an abortion event, the composition of the livestock herds (cattle, goats and sheep) kept at each household, and the number of previous abortion cases in each household that had an event.

The sensitivity of the platform was examined by dividing the number of investigated abortion cases by the *expected* number of abortions for the livestock population of the study wards over the study period. The expected number of abortions was estimated by multiplying (a) the number of abortion cases per head of livestock reported over a 12 month period obtained through a previous randomized cross-sectional study (described in de Glanville et al. 2022) by (b) the number of livestock in the ward reported by surveys conducted by LFOs from the Tanzania Ministry of Livestock and Fisheries covering a period from 2011 to 2016 (E. Swai, *unpublished data*). This figure was multiplied by two to account for the 24-month duration of the study. Where data were not available for a specific ward, figures were estimated from the average of other wards in the same division (Rau) or by current estimates provided by the LFO (Machame Mashariki) (Supplementary Materials 3).

Determinants of investigation

The determinants of event investigation were analysed to determine: first, the number of cases that each LFO investigated; second, the distribution of time taken between the report and the subsequent investigation by the research team; and third, using a Pearson correlation analysis, the relationship between the time taken between a report and its subsequent investigation and the distance (km) of the ward from the research team headquarters (in the town of Moshi) (Figure 2).

Sample collection data

Data were recorded and summarized for: (a) the types of samples collected for each event; (b) the availability of placental and foetal materials and, when not available, the reasons why these materials were not available; and c) the relationship between sample types and pathogen detection and pathogen attribution.

Observed patterns in investigated abortions

The distribution of investigated abortion cases was examined in relation to a) livestock species and breed in which they occurred, b) history of previous abortion cases in the aborting dams, c) a history of recent stressful events affecting the pregnant dam, and d) seasonality of the cases.

a) Breeds were recorded as indigenous (local) and non-indigenous (cross-bred and exotic). We used the breed distribution in the investigated herds to estimate the expected number of abortions if breed type did not affect the likelihood of abortion. Using an exact binomial test, we compared the expected with the observed number of cases.

b) Data on the occurrence of previous abortion cases and recent stressful incidents were obtained from the questionnaire data collected at the time of the investigation.

c) Stressful incidents were recorded as open-ended questions but were prompted by asking about events such as being chased by predators, drought, unusual handling or recent change in diet or grazing.

d) Seasonality was investigated in relation to typical 'dry' and 'wet' seasons and in relation to monthly rainfall in the Arusha region during the project period. Mean monthly rainfall was calculated from daily rainfall data, which was obtained across the Arusha Region (Ashouri et al. 2015; Sorooshian et al. 2019) and plotted alongside the temporal pattern of reported cases. The relationship between mean monthly rainfall and the number of reported abortions was investigated using linear regression.

All processing and analysis of data was carried out in R.

Determinants of attribution

World Organization for Animal Health (WOAH – previously OIE) guidelines and case definitions, in conjunction with recommendations from specialist/reference laboratories and

peer-reviewed literature, were used to inform diagnosis and attribution of infectious causes

of abortion (described in detail in Thomas et al. 2022). Binomial logistic regression analysis

was carried out with attribution (yes or no) as the dependent variable. Potential independent

explanatory variables, including the delay between the event and its investigation, species,

and livestock management system, were selected for inclusion in the model through

univariable analysis (with all variables with *p* value < 0.25 selected for inclusion). Potential

independent explanatory variables were retained or de-selected through a stepwise

approach using Akaike Information Criterion (AIC) as an indicator for model efficiency to

achieve a final model.

Exposure to zoonotic pathogens

The percentage of cases in which zoonotic pathogens (*Brucella* spp., *C. burnetii*, *T. gondii* and

RVFV) were detected, and the percentage of these cases in which someone assisted with the

delivery, was calculated. The age and sex of the persons assisting with delivery was also

collected.

Ethical Approval

Ethics approval for this research was granted by Kilimanjaro Christian Medical Centre (KCMC)

Ethics Committee (No, 535 and No. 832); National Institute of Medical Research (NIMR),

Tanzania (NIMR/HQ/R8a/Vol IX/1522 & NIMR/HQ/R.8a/Vol IX.2028); Research Ethics

Coordination Committee and the Institutional Review Board for Clinical Investigations of Duke

University Health System in the United States (Pr00037356); University of Otago Ethics

Committee (H15/069 & H17/069); University of Glasgow College of Medical Veterinary and

Life Sciences Ethics Committee (200140152 & 200170006).

Results

Descriptive Statistics

Between October 2017 and September 2019, 215 abortion cases were reported from 150

households in 13 of the 15 target wards (Fig. 2). The distribution of investigated cases in

relation to agro-ecological system and herd/flock composition is shown in Table 1. Of the 150

households investigated, most (n=115) had only one event investigated. Of the remaining

households, 21 had two cases investigated, eight had three, two had four, three had five and

one household had 11 cases investigated. Herd level summary statistics have been provided
in Supplementary Material 4.
319
The sensitivity of the platform (the percentage of expected abortion cases that were
investigated) ranged from 0%-12.4% for cattle, 0-1.2% for goats and 0-0.3% for sheep. A
higher percentage of expected abortions in cattle were reported in smallholder wards (2.7%)
than in other wards (0% for agropastoral and 0.31% for pastoral wards), with particularly high
reporting in one smallholder ward, Machame Mashariki, close to Moshi town, where 12.4%
of expected cattle abortions were investigated.
326
327 Over the 12 months preceding a reported abortion event, abortions, peri-natal mortalities, or
328 stillbirths occurred in 52 of 184 (28.3%) cattle herds, 98 of 168 (58.3%) goat herds and 93 of
329 140 (66.4%) sheep herds that had at least one adult female. Of herds that had at least one
330 abortion case, the mean (median, range) number of cases of abortion per herd in the past 12
331 months was 3.7 (3, 1 – 12) for cattle, 9.0 (6, 1 – 70) for goats and 6.7 (4, 1 – 67) for sheep.
332
Determinants of investigation
The number of cases reported to the research team by each LFO varied considerably (median
= 5, range 0 – 84). Of the 215 cases, 70% were reported by three (20%) of the LFOs, with one
reporting 84 cases (39.1%). Two LFOs did not report a single case (Figure 3a). The range in
the interval between the report and the subsequent investigation by the research team was
zero to four days with a median of one day (Figure 4).
339
340 From the Pearson correlation analysis, no significant association was found between the
341 interval between reporting and investigation and the direct distance between the centroid of
342 the ward in which the LFO worked and the research team headquarters (R^2 = -0.07, t = -0.32,
343 p = 0.75) (Figure 3b).
344
345
346
347

Event	Category	Number (%)
Number of abortion cases	All species	215
	Cattle	71 (33%)
	Goats	100 (46.5%)
	Sheep	44 (20.5%)
Number of abortion cases in each agricultural ecological zone	Pastoral	144 (67.0%)
	Agro-pastoral	1 (0.5%)
	Peri-urban	70 (32.5%)
Number of households that had an abortion event	Households	150
Number of households in each agricultural ecological zone	Pastoral	84 (56.0%)
	Agro-pastoral	1 (0.7%)
	Peri-urban	65 (43.3%)
Composition of the 150 herds	Cattle, goats and sheep	77 (51.3%)
	Cattle and goats	17 (11.3%)
	Goats and sheep	7 (4.7%)
	Cattle and sheep	1 (0.7%)
	Cattle only	40 (26.6%)
	Goats only	8 (5.3%)

348 **Table 1:** The number (and percentage) of abortion cases by species and agro-ecological zone

349 and the composition of the livestock herds (and percentage) in investigated households

356   **Figure 3:** a) The number and percentage of abortion cases reported by each LFO. b) The

357   relationship between the number of reports per LFO and the distance to the research based

358   in the town of Moshi.

361   **Figure 4:** The number of days between abortion and the investigation. No cases were

362   investigated more than after four days after the abortion.

Sample collection data

Out of the 215 cases, placental and foetal tissues were collected in 116 (24.0%) and 141 (34.1%), respectively. The reasons given for failure to collect placental and foetal tissues included: a) the tissues not being seen by owners (for example if the animal aborted while away grazing); b) the tissues being burned by the owner; and c) the tissues being consumed by dogs or other animals. Vaginal and milk samples from aborting dams were collected in 213 (99.1%) and 167 (77.7%) cases, respectively.

Observed patterns in investigated abortions

a) **Pattern of abortions in species and breeds**

In cattle, reported abortions occurred significantly more often than expected in non-indigenous cross-bred animals (expected proportion = 0.11, actual proportion = 0.52, 95% CI: 0.42 – 1.00, $p < 0.001$) and non-indigenous exotic animals (expected proportion = 0.01, actual proportion = 0.25, 95% CI: 0.16 – 1.00, $p < 0.001$). In goats, reported abortions occurred significantly more often than expected in non-indigenous cross-bred animals (expected proportion = 0.02, actual proportion = 0.18, 95% CI: 0.11 – 1.00, $p < 0.001$) and more often than expected in non-indigenous exotic animals, although this difference was not significant (expected proportion = 0.01, actual proportion = 0.03, 95% CI: 0.01 – 1.00, $p = 0.053$) (Figure 5 and Supplementary Materials 5). There was no significant difference in the distribution of abortions in different breeds of sheep.

Figure 5. The difference between the expected and actual proportion of abortion cases in each species and breed. Value of 0 = the expected number of cases occurred, > 0 more than expected, < 0 less than expected (LOC= indigenous (local), XB= non-indigenous cross-bred, EX= non-indigenous exotic breed).

b) **History of previous abortion cases in the aborting dams**

Of the cattle, goat and sheep dams that were investigated in this study (and that had had previous pregnancies), 33.3% (n=12), 29.8% (n=17) and 16.7% (n=5), respectively, were reported by the owner to have experienced a previous abortion event. Of these cattle and goats, 41.7% (n=5) and 47.1% (n=8), respectively, had suffered multiple previous abortions. One cow had experienced four previous abortion cases and one goat had experienced seven.

c) **History of recent stressful events**

Of dams that aborted, 16 of 71 cattle (22.5%), 37 of 98 goats (37.7%) and 15 of 43 sheep (34.9%) were reported to have experienced recent stress. Dams for which an aetiological attribution was not made were no more or less likely to have experienced a stressful event than dams that aborted for which an attribution was made. Regarding recent illnesses over the previous four weeks (diagnosed by the farmer), cattle were reported to have suffered

from a range of conditions including anaplasmosis, diarrhoea, lumpy skin disease, and

trypanosomiasis, whilst goats were reported to have suffered predominantly from respiratory

disease.

d) **Seasonality of cases**

Cases of abortion were reported in every month of the 24-month study period, and although

these fluctuated over time, with more cases reported during the drier periods (Figure 6), there

was no significant effect of mean monthly rainfall on the number of cases (correlation

coefficient = -0.005. t = -0.25, p = 0.8).

Figure 6. The number of abortion cases investigated per month (blue columns) shown against

mean rainfall recorded in the Arusha region over each month of the study period (red line).

Figure 7: The type of samples in which pathogens were detected in the 41 abortion cases for which an attribution was made using PCR are shown. Each row represents one abortion event. Red - sample type returned a positive result; blue - sample type returned a negative result; grey - sample type was not collected.

Determinants of Attribution

As described in Thomas et al. (2022), the number of cases for which an abortigenic agent was attributed was 42 out of 215 (19.5%). Of these, an attribution was made using PCR in 41 cases. One event in a cow met the case definition for two pathogens (both BHV-1 through seroconversion and *Neospora* through PCR) and in this event both pathogens were attributed. The attribution of the single case that was not determined using PCR (BHV-1) was made by serology alone. The sample types that were collected in each of these 41 cases for which an attribution was made using PCR, and whether the samples returned a positive or negative result, are shown in Figure 7. The time period between the abortion incident and the investigation (*Delay*) had a negative impact on attribution ($z = -2.1$, $p = 0.03$), with each daily increase in the delay corresponding to a decrease in the odds of an attribution being made

438   by 46.1% (i.e. 1 - 0.539) (Table 2 and Figure 8). Additionally, when the abortion occurred in

439   goats (*Goat*) an attribution was significantly less likely than when it occurred in cattle. Finally,

440   the odds of achieving an attribution were not affected by the availability of the placenta

441   (*Placenta present*), foetus (*Foetus present*) or milk (*Milk collected*) for sampling.

	Odds Ratio	2.5%	97.5%	z value	*p*
(Intercept)	0.353	0.053	2.361	-1.074	0.283
Delay	0.539	0.306	0.95	-2.139	0.032
Goat	0.195	0.058	0.653	-2.651	0.008
Sheep	0.615	0.142	2.664	-0.65	0.516
Foetus present	0.936	0.345	2.544	-0.129	0.897
Placental present	1.779	0.621	5.094	1.074	0.283
Milk collected	2.405	0.639	9.055	1.297	0.195

444   **Table 2**: Output of final regression model investigating determinants of attribution

Figure 8. Predicted probability of attribution being made as a function of increasing delay

between abortion and case investigation, as determined by the regression model output. The

blue line indicates the regression line with the 95% confidence interval shaded blue.

Exposure to zoonotic pathogens

Zoonotic pathogens (*Brucella* spp., *Coxiella burnetii*, *Toxoplasma gondii*. and RVFV) were

detected in 61 of the 77 (79.2%) abortion cases where a pathogen was detected. Respondents

reported that someone had assisted with the delivery in 13 (21.3%) of these cases, similar to

the proportion of assisted deliveries across all abortion cases (40 out of 215, 23.5%). Of those

assisting with delivery, the median age was 37, the youngest was seven, the eldest 84, and

20% were female.

Discussion

This study demonstrated the feasibility of establishing a surveillance platform for reporting

and diagnosing cases of livestock abortion and that these investigations, even at small scale,

have generated important information on livestock diseases, and their investigation, in

Tanzania. This included information that, to our knowledge, has not been captured or

reported through existing surveillance systems.

Key findings were that: i) livestock abortion can be a target for syndromic and/or event-based

surveillance, with abortion incidents sufficiently distinctive and noteworthy to be reported by

farmers to livestock field officers for investigation; ii) pragmatic and robust protocols for

sample collection and laboratory diagnosis can be established, even in resource-limited

settings, to generate important etiological and epidemiological data; iii) the likelihood of

obtaining an etiological diagnosis depended on the timeliness of reporting and sample

collection; iv) there was wide variation in reporting and investigation of cases by different

LFOs and this did not appear to be associated with the distance between the ward and the

investigation center; v) in many herds prior livestock abortions were reported to have

occurred in the 12 months preceding the investigated abortion event; vi) abortions and repeat

abortions reported disproportionately more frequently in non-indigenous breeds (cross-bred

and exotic) than local breeds; and vii) abortion cases were reported and investigated more

frequently during drier periods, although there was no evidence of a relationship with

monthly rainfall.

This study suggests that livestock abortion cases are sufficiently distinctive and observable to

be reported by farmers to LFOs, would have widespread acceptability as a target for

syndromic or event-based surveillance and would have considerable utility for farmers as well

as for the livestock and public health sectors. The platform demonstrated the potential for

generating high-quality aetiological data on previously relatively unrecognised livestock

disease problems in Tanzania, such as *Neospora caninum* and novel pestiviruses, as well as

data on endemic zoonoses, such as Q fever, brucellosis and toxoplasmosis (Thomas et al.

2022). Reports of unusual livestock abortion cases also provided an early warning of acute

human health risks, as shown in this study area by the detection of an outbreak of RVF in

livestock in peri-urban smallholder cattle (de Glanville et al. 2022) and which was

subsequently linked with human RVF cases and deaths (Madut et al., n.d.).

An attempt was made to gauge the sensitivity of the surveillance platform by estimating the

percentage of expected abortions that the investigated cases represented. This was carried

out using data on ward-level livestock numbers drawn from Ministry of Livestock and

Fisheries records. However, given the dynamic nature of livestock populations in the study

area and shifts in ward configurations, these estimates should be considered approximate

figures only. Overall the sensitivity of this platform was low, with the highest ward level

sensitivity (12.4%) below a mean sensitivity of 34% reported for bovine abortion surveillance

systems in France (Bronner et al. 2015). Nonetheless, given that this was the first time that

an abortion surveillance platform of this kind had been introduced, and the logistic and

communication challenges inherent to this type of low income setting, our study gives

confidence of the feasibility and potential utility of this approach. Despite the limitations of

the data, our conclusion that there is wide variation in the sensitivity of the surveillance

platform across wards is likely to be robust. This finding, together with the ability of the

platform to investigate only a very small proportion of expected abortion cases overall, has

implications for the representativeness of data generated, which needs to be considered

when interpreting the results reported here.

Contrary to expectations, the location and distance of the ward and the type of farming

system was not associated with the probability of cases being investigated. Relatively few

cases were reported and investigated in agropastoral systems but both high and low rates of

reporting occurred in smallholder and pastoral systems. This suggests that LFOs and farmers

in some wards were much more engaged and motivated than in others. To date, much

investment in animal and human surveillance has been directed towards strengthening
laboratory diagnostic capacity. While this is clearly essential, support also needs to be given
to engaging, training, and motivating farmers and front-line animal health workers to
effectively report both routine and unusual animal disease events. Understanding why and
how animal keepers and LFOs are incentivised to report and investigate livestock health
events will be essential for improving the reach and sensitivity of future surveillance
platforms. The infrastructure for LFO engagement across Tanzania provides a valuable
platform for timely and cost-effective reporting of abortions and other disease events but
seems to be underutilised in existing surveillance systems.

The effectiveness of the platform in allowing an aetiological diagnosis to be reached was
significantly impacted by the timeliness of investigation. The impact of delays may be
explained by degradation of diagnostic samples, which also affected the utility of
histopathological diagnosis (Thomas et al. 2022). Delays were caused by several factors,
including the time taken by the investigation team to reach households due to remoteness of
some areas and multiple cases being reported in different areas on the same day. These
findings underscore the importance of being able to respond rapidly to cases and, where
access is difficult, of providing locally suitable means of transport.

A reactive surveillance platform that is managed locally, for example by ward-based LFOs, will
require the establishment of effective and safe protocols for collection and transport of
samples for laboratory diagnosis that can be carried out without the need for long-distance
travel of more highly trained investigators. Protocols for collection of vaginal swabs from
aborting dams, which proved effective for both pathogen detection and attribution, may be
of value. In this study, vaginal swabs performed better than placental samples and foetal
samples when detecting abortigenic pathogens (Figure 7). They also require less handling of
potentially infectious tissues and are more reliably accessible than placental and foetal
tissues, which are often consumed by scavengers or disposed of by the farmer. Indeed,
collection of vaginal swabs from the dam was possible in almost all cases investigated in this
study. Thus, while logistic, financial, and capacity constraints for comprehensive sampling and
investigation of livestock abortions are likely to exist across Africa, these need not preclude
the establishment of simple and robust protocols that can yield valuable surveillance data.

Attribution

From this platform, an attribution was reached in 19.5% of cases (Thomas et al. 2022), which

was not far below the typical range of 25-45% achieved in industrialized farming systems in

high-income settings (Campero et al. 2003; Amouei et al. 2019; Derdour et al. 2017;

Anderson, Barr, and Hoffman 1990). During this study, samples were tested for only ten

abortigenic pathogens and it is likely that rates of detection and attribution would be higher

with inclusion of tests for other known abortigenic agents (such as *Listeria* spp.,

Campylobacter spp., *Salmonella* spp. and fungal pathogens), with more locally relevant

aetiological data informing the suite of pathogens to be tested, and with metagenomic

approaches.

As elsewhere in the world, non-infectious conditions, including nutritional, metabolic, and

toxic conditions (Alemayehu et al. 2021; Mee et al. 2023; Woodburn et al. 2021), are also

likely to cause livestock abortions in Tanzania. Our study suggests a possible role for recent

stressful events, including episodes of drought and attacks by wild predators, which were

reported from approximately a third of unattributed abortion cases. However, these events

are a common occurrence in our study population, so attribution is likely to be challenging.

We do not have an immediate explanation as to why the likelihood of attribution was lower

for goats than for sheep or cattle. Given this is just one study, it could just be a random finding.

Another possible explanation is that, in comparison to sheep and cattle, fewer data are

available worldwide on causes of abortions in goats and very little is known about the

aetiology of goat abortions in Africa, so the pathogens included in our diagnostic panel may

not have been as relevant for goats as for sheep and cattle. This lack of knowledge is

highlighted by the unusual finding of *Neospora caninum* as a cause of abortion in a goat

detected through this platform (Thomas et al. 2022).

The challenges of attribution, discussed also in Thomas et al. (2022), should not detract from

the value of reporting and investigating livestock abortion cases. Recent studies have

indicated that the economic costs of livestock abortion in Tanzania and impacts on food

security are much more substantial than previously recognized (Semango et al. under review)

and data collected on the number of cases and species/breed affected will be of considerable

value in highlighting the importance of this syndrome in the context of livestock productivity,

household livelihoods and food security.

Patterns of livestock abortion in northern Tanzania

Data from the household questionnaire indicated that livestock abortion was a common

occurrence across the study area with almost a third of cattle herds and two-thirds of small

ruminant herds having experienced a previous case of abortion in the preceding twelve

months. Our results also suggest a possible link between abortion and low rainfall, although

further work will be needed to examine this relationship in detail.

This study indicated that investigated cases occurred more often than expected in non-

indigenous breeds (cross-bred and exotic cattle and cross-bred goats) than in local breeds.

This is consistent with studies reporting a four-times higher risk of abortion in cross-bred

cattle in Ethiopia in comparison to local breeds (Deresa, Tulu, and Deressa 2020) and studies

in Nigeria (Yakubu, Awuje, and Omeje 2015) and India (Khan, Manoj, and Pramod 2016)

reporting breed effects in relation to cattle abortion. Our results may simply reflect a higher

likelihood of reporting of abortion in animals that are of higher value and more likely to be

zero-grazed or tethered while grazing (and hence abortions would be more apparent) and for

which an existing relationship with animal health services is more likely.

More than a fifth of the dams that aborted were reported to have experienced a previous

abortion, and several had experienced multiple abortion losses, with non-indigenous breeds

more likely than indigenous breeds to have had a previous abortion. Chronic repeat abortions

can arise from infection with pathogens such as *Neospora caninum* and Pestiviruses which

were detected as common causes of abortion in this study. Repeat aborting dams could be

identified as candidates for slaughter with accruing potential benefits for herd fertility and

productivity, as well as reduced risks of potential zoonotic transmission. However, any advice

around removal of these animals from a herd may result in sale rather than slaughter, with

the risk of further spread of the pathogen.

The risk of exposure to zoonotic pathogens through abortion cases was underscored by two

findings: first that, in nearly 80% of cases in which a pathogen was detected, the pathogen

was zoonotic and second, that in nearly a quarter of all cases, someone had assisted with the

aborted delivery, very likely without any form of personal protective equipment. Of these

people who assisted, 20% were female, of which most were of reproductive age and therefore

at particular risk following infection with pathogens such as Toxoplasma gondii.

In conclusion, this study has demonstrated that livestock abortion surveillance, even at a

relatively small scale, can capture valuable aetiological and epidemiological information on

important livestock pathogens, including pathogens that are zoonotic and those with

epidemic potential. The study demonstrated the utility, acceptability, and feasibility of

livestock abortion as a target for both syndromic and event-based surveillance and showed

that an effective reporting and investigation system could be operationalized across a range

of settings in Tanzania, including remote rural areas. Key elements of effectiveness were high

levels of engagement of community-based field officers and the establishment of practical

and robust field sample collection and molecular diagnostic protocols, with prompt response

to reporting of cases, and prompt feedback of results.

Acknowledgements

This study was supported by the Supporting Evidence Based Interventions project, University

of Edinburgh (grant number R83537). We are grateful for the support of the Ministry of

Livestock and Fisheries, the District Veterinary Officers, the Livestock Field Officers and the

communities that contributed to this work. We are also grateful to the research team who

implemented the project. This work was also supported by the BBSRC, the Department for

International Development (DFID), the Economic & Social Research Council (ESRC), the

Medical Research Council (MRC), the Natural Environment Research Council (NERC) and the

Defence Science & Technology Laboratory, under the Zoonoses and Emerging Livestock

Systems (ZELS) programme (BB/L018926/1).

638
639

References:

Africa Centres for Disease Control and Prevention. 2018. "Africa CDC - Event Based Surveillance Framework." http://www.africacdc.org/resources/strategic-framework/Strategic%0AFramework/Africa%0ACDC%0AEBS%0AFramework%0A-%0AEN.pdf/download.

Ahmed, Haseeb, Jonathan Yoder, William A. de Glanville, Alicia Davis, Tito J. Kibona, Blandina T. Mmbaga, Felix Lankester, Emmanuel S. Swai, and Sarah Cleaveland. 2019. "Economic Burden of Livestock Disease and Drought in Northern Tanzania." *Journal of Development and Agricultural Economics* 11 (6): 140–51. https://doi.org/10.5897/jdae2018.1028.

Alemayehu, Gezahegn, Gezahegne Mamo, Biruk Alemu, Hiwot Desta, Biniam Tadesse, Teferi Benti, Adane Bahiru, Muhabaw Yimana, and Barbara Wieland. 2021. "Causes and Flock Level Risk Factors of Sheep and Goat Abortion in Three Agroecology Zones in Ethiopia." *Frontiers in Veterinary Science* 8 (January): 1–16. https://doi.org/10.3389/fvets.2021.615310.

Amouei, Afsaneh, Mehdi Sharif, Shahabeddin Sarvi, Ramin Bagheri, Sargis A Aghayan, Mohammad Bagher Hashemi-soteh, Mohammad Sarafrazi, and Ahmad Daryani. 2019. "Aetiology of Livestock Fetal Mortality in Mazandaran Province, Iran." *PeerJ* 6: 1–14. https://doi.org/10.7717/peerj.5920.

Anderson, Mark L., Bradd C. Barr, and Rick L. Hoffman. 1990. "A Survey of Causes of Bovine Abortion Occurring in the San Joaquin Valley, California." *Journal of Veterinary Diagnostic Investigation* 2 (4): 283–87. https://doi.org/10.1177/104063879000200405.

Ashouri, Hamed, Kuo Lin Hsu, Soroosh Sorooshian, Dan K. Braithwaite, Kenneth R. Knapp, L. Dewayne Cecil, Brian R. Nelson, and Olivier P. Prat. 2015. "PERSIANN-CDR: Daily Precipitation Climate Data Record from Multisatellite Observations for Hydrological and Climate Studies." *Bulletin of the American Meteorological Society* 96 (1): 69–83. https://doi.org/10.1175/BAMS-D-13-00068.1.

Bodenham, Rebecca F., Stella Mazeri, Sarah Cleaveland, John A. Crump, Folorunso O. Fasina, William A. de Glanville, Daniel T. Haydon, et al. 2021. "Latent Class Evaluation of the Performance of Serological Tests for Exposure to Brucella Spp. In Cattle, Sheep, and Goats in Tanzania." *PLoS Neglected Tropical Diseases* 15 (8): 1–23. https://doi.org/10.1371/journal.pntd.0009630.

Bronner, Anne, Emilie Gay, Nicolas Fortané, Mathilde Palussière, Pascal Hendrikx, Viviane Hénaux, and Didier Calavas. 2015. "Quantitative and Qualitative Assessment of the Bovine Abortion Surveillance System in France." *Preventive Veterinary Medicine* 120 (1): 62–69. https://doi.org/10.1016/J.PREVETMED.2015.02.019.

Cabell, Emma. 2007. "Bovine Abortion: Aetiology and Investigations." *In Practice* 29 (8): 455–63. https://doi.org/10.1136/inpract.29.8.455.

Campero, C. M., D. P. Moore, A. C. Odeón, A. L. Cipolla, and E. Odriozola. 2003. "Aetiology of Bovine Abortion in Argentina." *Veterinary Research Communications* 27 (5): 359–69. https://doi.org/10.1023/A:1024754003432.

Cleaveland, S., L. Kusiluka, J. Ole Kuwai, C. Bell, and R. Kazwala. 2001. "Assessing the Impact of Malignant Catarrhal Fever in Ngorongoro District, Tanzania." *Report to the DFID Animal Health Programme, Centre for Tropical Veterinary Medicine, University of Edinburgh*.

Derdour, Salima-yamina, Fella Hafsi, Naouelle Azzag, Safia Tennah, Abdelouahab Laamari,

and Farida Ghalmi. 2017. "Prevalence of the Main Infectious Causes of Abortion in
Dairy Cattle in Algeria." *Journal of Veterinary Research* 61: 337–43.
https://doi.org/10.1515/jvetres-2017-0044.

Deresa, Benti, Dereje Tulu, and Feyissa Begna Deressa. 2020. "Epidemiological Investigation
of Cattle Abortion and Its Association with Brucellosis in Jimma Zone, Ethiopia."
Veterinary Medicine: Research and Reports Volume 11: 87–98.
https://doi.org/10.2147/vmrr.s266350.

Dorea, Fernanda C., Javier Sanchez, and Crawford W. Revie. 2011. "Veterinary Syndromic
Surveillance: Current Initiatives and Potential for Development." *Preventive Veterinary*
Medicine 101 (1–2): 1–23. https://doi.org/10.1016/j.prevetmed.2011.05.004.

Dórea, Fernanda C, and Flavie Vial. 2016. "Animal Health Syndromic Surveillance: A
Systematic Literature Review of the Progress in the Last 5 Years (2011–2016)."
Veterinary Medicine: Research and Reports Volume 7 (November): 157–70.
https://doi.org/10.2147/vmrr.s90182.

ECDC. 2014. "European Centre for Disease Prevention and Control. Data Quality Monitoring
and Surveillance System Evaluation - A Handbook of Methods and Applications." *ECDC*
Technical Document. Stockholm.
http://ecdc.europa.eu/en/publications/_layouts/forms/Publication_DispForm.aspx?Lis
t=4f55ad51-4aed-4d32-b960-
af70113dbb90&ID=1171%250Ahttps://ecdc.europa.eu/en/publications-data/data-
quality-monitoring-and-surveillance-system-evaluation-handbook-methods-an.

Food and Agriculture Organization. 2011. "World Livestock 2011 - Livestock in Food
Security." Rome.

George, Janeth, Barbara Häsler, Erick Komba, Calvin Sindato, Mark Rweyemamu, and James
Mlangwa. 2021. "Towards an Integrated Animal Health Surveillance System in
Tanzania: Making Better Use of Existing and Potential Data Sources for Early Warning
Surveillance." *BMC Veterinary Research* 17 (1). https://doi.org/10.1186/s12917-021-
02789-x.

Givens, D. M., and M. S.D. Marley. 2008. "Infectious Causes of Embryonic and Fetal
Mortality." *Theriogenology* 70 (3): 270–85.
https://doi.org/10.1016/j.theriogenology.2008.04.018.

Glanville, William A. de, Kathryn J. Allan, James M. Nyarobi, Kate M. Thomas, Felix
Lankester, Tito J. Kibona, John R. Claxton, et al. 2022. "An Outbreak of Rift Valley Fever
among Peri-Urban Dairy Cattle in Northern Tanzania." *Transactions of the Royal Society*
of Tropical Medicine and Hygiene, 1–9. https://doi.org/10.1093/trstmh/trac076.

Hulth, A. 2014. "First European Guidelines on Syndromic Surveillance in Human and Animal
Health Published." *Eurosurveillance* 19 (41): 9806.
https://doi.org/10.2807/ese.19.41.20927-en.

Khan, M. H., K. Manoj, and S. Pramod. 2016. "Reproductive Disorders in Dairy Cattle under
Semi-Intensive System of Rearing in North-Eastern India." *Veterinary World* 9 (5): 512–
18. https://doi.org/10.14202/vetworld.2016.512-518.

Madut, DB, MP Rubach, K Allan, KM Thomas, WA de Glanville, JEB Halliday, C Costales, et al.
n.d. "Epidemiologic and Genomic Characterization of an Outbreak of Rift Valley Fever
among Humans and Dairy Cattle in Northern Tanzania." *Under Review*.

Mee, J. F., C. Hayes, T. Stefaniak, and P. Jawor. 2023. "Review: Bovine Foetal Mortality - Risk
Factors, Causes, Immune Responses and Immuno-Prophylaxis." *Animal : An*
International Journal of Animal Bioscience 17: 100774.

https://doi.org/10.1016/j.animal.2023.100774.

Neumann, Charlotte G., Nimrod O. Bwibo, Suzanne P. Murphy, Marian Sigman, Shannon
Whaley, Lindsay H. Allen, Donald Guthrie, Robert E. Weiss, and Montague W.
Demment. 2003. "Animal Source Foods Improve Dietary Quality, Micronutrient Status,
Growth and Cognitive Function in Kenyan School Children: Background, Study Design
and Baseline Findings." *Journal of Nutrition* 133 (11 SUPPL. 2): 3941S-3949S.
https://doi.org/10.1093/jn/133.11.3941s.

R Core Team. 2023. "R: A Language and Environment for Statistical Computing." Vienna: R
Foundation for Statistical Computing. http://www.r-project.org/.

Semango, G., J. Yoder, T.K. Kibona, J.R. Claxton, J. Buza, S. S. Johnson, S. Cleaveland, and F.
Lankester. 2023. "Economic Burden of Livestock Abortions in Northern Tanzania."
Journal of Agricultural and Applied Economics.

Sorooshian, Soroosh, Kuolin Hsu, Dan Braithwaite, Hamed Ashouri, and NOAA CDR Program.
2019. "NOAA Climate Data Record (CDR) of Precipitation Estimation from Remotely
Sensed Information Using Artificial Neural Networks (PERSIANN-CDR), Version 1
Revision 1." NOAA National Centers for Environmental Information. 2019.
https://doi.org/10.7289/V51V5BWQ.

The World Bank. 2021. "Moving towards Sustainability: The Livestock Sector and the World
Bank." World Bank Understanding Poverty. 2021.
https://www.worldbank.org/en/topic/agriculture/brief/moving-towards-sustainability-
the-livestock-sector-and-the-world-bank.

Thomas, Kate M, Tito Kibona, John R Claxton, William A de Glanville, Felix Lankester, Joram J
Buza, Ryan Carter, et al. 2022. "Prospective Cohort Study Reveals Unexpected
Aetiologies of Livestock Abortion in Northern Tanzania." *Scientific Reports* 12: 1–50.
https://doi.org/10.1038/s41598-022-15517-8.

Wolf-Jäckel, Godelind Alma, Mette Sif Hansen, Gitte Larsen, Elisabeth Holm, Jørgen Steen
Agerholm, and Tim Kåre Jensen. 2020. "Diagnostic Studies of Abortion in Danish Cattle
2015-2017." *Acta Veterinaria Scandinavica* 62 (1). https://doi.org/10.1186/s13028-
019-0499-4.

Woodburn, Daniel B., Johan Steyl, Elizabeth C. Du Plessis, Rick D. Last, Bjorn Reininghaus,
and Emily P. Mitchell. 2021. "Pathological Findings in African Buffaloes (Syncerus
Caffer) in South Africa." *Journal of the South African Veterinary Association* 92: 1–11.
https://doi.org/10.4102/JSAVA.V92I0.2117.

Woolhouse, Mark E.J., and Sonya Gowtage-Sequeria. 2005. "Host Range and Emerging and
Reemerging Pathogens Emerging Infectious Diseases • Www." *Emerging Infectious
Diseases* 11 (12): 1842–47. https://doi.org/10.3201/eid1112.050997.

World Health Organization. 2008. "A Guide to Establishing Event-Based Surveillance." *World
Health Organiszation*, 21.

Yakubu, A., A. D. Awuje, and J. N. Omeje. 2015. "Comparison of Multivariate Logistic
Regression and Classification Tree to Assess Factors Influencing Prevalence of Abortion
in Nigerian Cattle Breeds." *Journal of Animal and Plant Sciences* 25 (6): 1520–26.

775