Diverse bacteriohemerythrin genes of *Methylomonas denitrificans* FJG1 provide insight into

the survival and activity of methanotrophs in low oxygen ecosystems

Running Title: Bacteriohemerythrin genes in methanotrophs

Cerrise Weiblen[1], K. Dimitri Kits[1], Manuel Kleiner[2], Dominic Sauvageau[3], Lisa Y. Stein[1#]

[1]Department of Biological Sciences, University of Alberta, Edmonton, AB, Canada

[2]Department of Plant and Microbial Biology, North Carolina State University, Raleigh, NC, USA

[3]Department of Chemical and Materials Engineering, University of Alberta, Edmonton, AB,

Canada

Correspondence: lisa.stein@ualberta.ca

Abstract word count: 240

Text word count: 4860

Abstract

Proteobacterial methanotrophs are dependent on the oxidation of methane for ATP

production and assimilation of carbon into biomass. Interestingly, some types of

gammaproteobacterial methanotrophs thrive in oxygen-depleted zones of lakes and other aquatic

ecosystems despite their reliance on oxygen to support methane oxidation. The model

gammaproteobacterial methanotroph, *Methylomonas denitrificans* FJG1, oxidizes methane

coupled to nitrate reduction under hypoxia and highly upregulates its expression of

bacteriohemerythrin (Bhr). Bhr is a homolog of eukaryotic hemerythrin, which is a protein

associated with oxygen binding. Ten *bhr* homologs were identified in the genome of *M.*

denitrificans FJG1, requiring phylogenetic and gene expression analyses to pinpoint which

homolog is likely responsible for delivering oxygen to support methane oxidation to under low

oxygen conditions. This study examined the prevalence and phylogeny of the 10 *bhr* homologs

from *M. denitrificans* FJG1 in the genomes of other methanotrophs and across the Bacteria. The

homolog denoted "*bhr*-00" was specific to methanotroph genomes, was highly expressed in *M.*

denitrificans FJG1 under hypoxia, and its predicted structure was nearly identical to a purified

oxygen-scavenging hemerythrin protein from *Methylococcus capsulatus* Bath. Other *bhr*

homologs upregulated from denitrifying cultures of *M. denitrificans* FJG1 included those with

gene neighborhoods related to oxygen sensing, denitrification and chemotaxis. Together, this

study uncovered potential multifunctional roles of bacteriohemerythrin genes of *M. denitrificans*

FJG1 under low oxygen conditions and identified the *bhr* homolog that most likely enables and

supports oxygen delivery to methane monooxygenase enzymes in anoxic ecosystems.

Importance

Aerobic gammaproteobacterial methanotrophs can survive and grow in anoxic lakes, but

mechanisms that provide them with oxygen to support methane oxidation remain

uncharacterized. *Methylomonas denitrificans* FJG1 encodes 10 copies of bacteriohemerthyrin

(*bhr*), of which 7 are expressed at the mRNA level under low oxygen conditions. Comparing the

10 *bhr* homologs from *M. denitrificans* FJG1 with those from other methanotrophs and bacterial

genomes shows that two are specific to methanotrophs. Gene neighbourhoods surrounding

conserved *bhr* genes in methanotrophs suggest a range of potential functions including oxygen

respiration, oxygen sensing, chemotaxis, and nitrate reduction. The results from this study

illuminate a previously undescribed diversity of structures and potential functions of *bhr*

homologs in *M. denitrificans* FJG1 and related methanotrophic bacteria. The results pinpoint a

methanotroph-specific homolog, *bhr*-00, that is likely responsible for oxygen binding and

delivery to methane monooxygenase enzymes to promote methane oxidation in low oxygen

ecosystems.

Introduction

Hemerythrin (Hr), and its bacterial homolog bacteriohemerythrin (Bhr), is a

multifunctional and structurally diverse protein family linked to multiple metabolic functions in

eukaryotes and prokaryotes, including reactive oxygen and nitrogen species detoxification, metal

detoxification, iron storage, transmembrane signaling, oxygen sensing, chemotaxis, biofilm

formation, cellular respiration, and oxygen binding and transport (1, 2). Previous research has

shown that genetically similar Hr and Bhr enzymes perform an array of different functions,

placing Hr and Bhr proteins at the crossroads of multiple pathways, even within a single

organism. For instance, Baert et al. (3) identified Hr from the leech *Theromyzon tessulatum*

which was classified as an iron storage molecule named "ovohemerythrin." In bacterial systems,

Kendall et al. (4) investigated the functions of three Bhr homologs in *Campylobacter jejuni*: one

had a protective role in preventing molecules with iron-sulfur clusters (Fe-S) from incurring

oxygen damage and the other two were related to flagellar biosynthesis and expression of sigma-

factor 28. In the bacterium *Neanthes diversicolour*, a Bhr was found associated with cadmium

detoxification functions (5). Research from Xiong et al. (6) showed that the oxygen-sensitive

bacterium *Desulfovibrio vulgaris* used a Bhr-like protein as an oxygen sensor with a role in

chemotaxis. French et al. (7) showed that Bhr in *Vibrio cholerae* was part of the cyclic-di-GMP

signaling system with involvement in biofilm formation.

Several reports of Bhr proteins in gammaproteobacterial methanotrophs have emerged in

recent years. These aerobic bacteria use methane as a sole energy and carbon source. A previous

survey of *bhr* homologs in methanotroph genomes showed that 80% of those analyzed possessed

at least one, and some up to three, *bhr* gene copies (8). In the model strain *Methylococcus*

capsulatus Bath, *in vitro* studies showed that purified Bhr could supply oxygen to membrane

fractions enriched with particulate methane monooxygenase (pMMO) enzymes, and a crystal

structure of the Bhr protein was resolved (9-11). In *Methylotuvimicrobium alcaliphilum* 20Z,

Nariya and Kalyuzhnaya (12) showed that overexpression of Bhr supported aerobic respiration

but did not enhance methane oxidation capacity under hypoxia or anoxia. A *bhr* knock-out strain

of *Methylomonas* sp. LW13 showed decreased growth in a hydrogel with a methane-oxygen

counter gradient, but showed no growth impairment in planktonic cultures (13). These studies

indicate a relationship between Bhr and oxygen binding, delivery, and/or respiration in

methanotrophic bacteria, but none showed the range of potential Bhr activities or their direct

roles in supporting methanotrophic metabolism under low oxygen.

Aerobic methanotrophs can use alternate terminal electron acceptors including nitrate

(14), nitrite (15), and iron oxides (16). They can also link methane oxidation to fermentation

pathways in the absence of oxygen (17). However, there is no known substitute for oxygen that

can facilitate the turnover of methane by pMMO enzymes. The present study examines the

phylogeny, gene neighborhoods, expression levels, and predicted structures of the 10 *bhr*

homologs identified in the genome of *M. denitrificans* FJG1, a denitrifying

gammaproteobacterial methanotroph (14), as it transitions from oxic to hypoxic growth

conditions. The results highlight a wide diversity of *bhr* contexts in *M. denitrificans* FJG1 and

other methanotroph genomes and pinpoints the *bhr* homolog that likely enables

gammaproteobacterial methanotrophs to access oxygen to thrive in low oxygen environments.

Results

Phylogeny of *bhr* genes identified in *Methylomonas denitrificans* FJG1

    M. denitrificans FJG1 encodes 10 *bhr* homologs annotated as "hemerythrin family,"

"hemerythrin domain-containing protein," or "bacteriohemerythrin" in its genome (Table 1). The

genes, numbered *bhr-00* through *bhr-99*, were designated based on their proximity to the origin

of replication in the *M. denitrificans* FJG1 genome sequence (Supp. Fig. S1). Eight of the 10 *bhr*

genes are approximately 400 bp, typical of the oxygen-carrying Hr and Bhr proteins originally

discovered in eukaryotes (2), whereas *bhr-66* is approximately 1,000 bp and *bhr-55* is

approximately 4,000 bp (Table 1). BLASTn searches were performed with each of the 10 *bhr*

genes to identify homologs across methanotrophic and non-methanotrophic bacterial genomes.

Candidate genes from these BLASTn searches were aligned to a concatenated sequence of the 10

bhr homologs (separated by 100 bp spacers to prevent mismatching) from *M. denitrificans* FJG1.

and a phylogenetic tree was generated (Fig. 1). The resulting unrooted phylogram of positive

matches revealed that *bhr-66* is the most common *bhr*-containing gene found among a diverse

assemblage of Bacteria (Fig. 1). In contrast, *bhr-00* and *bhr-99* appear nearly exclusive to

methanotrophs in the Methylococcales order. Closer examination of complete *bhr* genes from

methanotroph genomes revealed that *bhr*-00 and *bhr*-66 homologs were the most common, but

only *M. denitrificans* FJG1 encoded *bhr*-11,-33,-44 and -55 among the proteobacterial

methanotrophs (Fig. 1; Supp. Data, Sheet 4).

    As *bhr*-66 was the most common homolog found in bacterial genomes, we examined its

detailed phylogeny with organisms for which full genomes were available (Fig. 2; Supp. Data,

Sheet 4). A cladogram using full-length *bhr*-66 alignments showed its polyphyletic distribution,

several instances of horizontal gene transfer, and a range of annotations (e.g. histidine kinase,

response regulator, hypothetical protein) that suggest a relationship to the Bhr class of oxygen-

sensing and signal transducing types (18). Expectedly, *bhr*-66 was monophyletic within the

methanotrophic bacteria. We compared the phylogeny of the methanotroph *bhr*-66 genes with

their 16S rRNA genes and found that they were highly congruent, suggesting an ancestral vertical

transmission (Fig. 3). Similar congruence was found for the methanotroph-specific *bhr*-00 gene.

When comparing phylogenies between *bhr*-00 and *bhr*-99, their high congruence and sequence

similarity to each other suggests that they are paralogs (Fig. 3). Similarly, two pairs: *bhr*-33/-44,

and *bhr*-77/-88, share high sequence similarity to one another, suggesting that each of the two

pairs may also be paralogs (Supplemental data sheet 3). The *bhr*-77/-88 genes are encoded

contiguously, perhaps due to a recent gene duplication event, and thus share the same gene

neighbourhood. As very few genomes of methanotrophs or other bacteria were found to encode

bhr-11, -22, -33, -44, -77 and -88, we could not perform a detailed phylogenetic analysis.

Conservation and predicted functions of *bhr* gene neighbourhoods in methanotrophic

bacteria

136   To explore the possible functionalities of commonly shared *bhr* genes among

methanotrophic bacteria, we examined the surrounding gene neighbourhoods for *bhr*-00, -66 and

-99 across available complete methanotroph genomes (Fig. 4). The *bhr*-00 gene neighbourhood

is highly conserved in the *Methylomonas* and *Methylotuvimicrobium* genera with flanking gene

clusters for biotin synthesis (*bio*BFHCD), an enzyme co-factor, and cytochrome *c* oxidase genes

(*cox*/*cta*) for oxygen respiration. The *bhr*-66 gene neighbourhood is less well conserved, with

genes related to signal transduction (i.e. response regulator) and ATP-binding domains flanking

bhr-66 in most *Methylomonas* genomes, and genes for the electron carrying cytochrome P-460

flanking *bhr*-66 in *Methylmicrobium* and *Methylotuvimicrobium* genomes (Fig. 4). Although *bhr*-

99 appears to be a paralog of *bhr*-00, its gene neighbourhood is distinct and conserved in four of

the *Methylomonas* genomes (Fig. 4) which show the AlkB DNA-repair gene encoded on the

same DNA strand as *bhr*-99. An operon for the Na$^+$-translocating NADH:quinone oxidoreductase

(NQR) complex, which is involved in creating sodium motive force for transporter and flagellar

function (19), is encoded on the opposing strand. The upstream region of *bhr*-99 is not well

conserved among the analyzed methanotroph genomes (Fig. 4).

Expression of *bhr* homologs and their neighboring genes in *M. denitrificans* FJG1

transitioning from high to low oxygen

154   Expression of the *bhr* homologs and their surrounding gene neighbourhoods under oxic

(24 h) and hypoxic (48 and 72 h) conditions were determined for cells grown in ammonium

(AMS) or nitrate (NMS) mineral salts media (14). Among the *bhr* genes, only *bhr*-00 showed a

strong (>2 norm log2 ratio) increase in mRNA abundance under hypoxia in both NMS and AMS

media (Fig. 5, Table 2, Supp. Fig. 3). This transcript was also, by orders of magnitude, the most

highly expressed among the *bhr* homologs (Fig. 5, Table 2). Even though previous research

showed that *M. denitrificans* FJG1 reduces nitrate only when grown in NMS (14), the AMS-

grown bacteria also expressed *bhr*-00 mRNA to the same magnitude under hypoxia (Fig. 5, Table

2), indicating that oxygen deprivation and not N-source was the signal affecting its expression.

Other *bhr* homologs (*bhr*-11, -55, -66, -77, -88) showed significantly increased mRNA

expression (P<0.05) under hypoxia (ratio of 24:48 h time points) when *M. denitrificans* FJG1 was

grown in NMS, but not in AMS, media suggesting a potential linkage to denitrification functions

(Table 2). Proteomic analysis was performed only for NMS-grown cells to compare expression

responses at the protein and mRNA levels. We detected both Bhr-00 and Bhr-55 in the proteome

(accession numbers GI:998900537, GI:1011416966 respectively) (Fig. 5, Supp. Fig. 3, Supp.

Data sheet 6). Bhr-00 increased by around 6-fold from the oxic to the hypoxic condition, making

up >1% of the total proteome. In contrast, Bhr-55 was of much lower abundance (<0.001%) and

it decreased from the oxic to hypoxic condition, contrary to expression of its transcript.

While genes for cytochrome *c* oxidase in the gene neighborhood of *bhr*-00 showed

decreased expression at the mRNA level under hypoxia, the protein abundances remained relatively

constant for this gene cluster (Fig. 5). In the same gene neighborhood, the *bioB* biotin synthase gene

showed decreased mRNA, but increased protein abundance with hypoxia. Genes encoding the

enzymes for nitrate reduction within the gene neighbourhood of *bhr*-11 showed increased mRNA

levels under hypoxia for only NMS-grown cells, which is consistent with the onset of

denitrification activity (14) (Fig. 5). Only the NarI protein in the nitrate reductase gene cluster

was detected in the proteome, but its expression was elevated under hypoxia. The gene

neighbourhood of *bhr*-66 showed increased expression at the mRNA and protein levels for the

response regulator histidine kinase and polyphosphate kinase as NMS-grown cells transitioned

from oxic to hypoxic conditions, suggesting potential involvement in oxygen sensing under

denitrifying conditions (Fig. 5). Furthermore, the gene neighbourhood surrounding *bhr*-77/-88

includes multiple genes for chemotaxis that showed increased expression at both the mRNA and

protein levels in NMS-grown cells as they transitioned from oxic to hypoxic conditions, in the

same range as the nitrate reduction genes (Fig. 5, Supp. Table 1). The results suggest that

chemotaxis and oxygen sensing may be facilitated by *bhr*-domain proteins in *M. denitrificans*,

which is consistent with Bhr functions characterized in other bacteria (7, 18, 20).

Diversity of Bhr protein structures from *M. denitrificans* FJG1

Because the 10 *bhr* genes of *M. denitrificans* FJG1 vary in size and genomic context, we

explored their predicted protein structural diversity using AlphaFold2 (Fig. 6A). Each predicted

structure showed unique features, although the Bhr-33 and -44 structures were similar to one

another. When the predicted structure of Bhr-00 was overlaid with the experimentally derived

crystal structure of the Bhr-Bath protein, which was demonstrated as an oxygen binding and

delivery system to pMMO enzymes in cell-free membrane systems (11), they were congruent

even though their amino acid identities overlapped by only 57.58% (Fig. 6B, Supp. Fig. S2,

Supp. Movie).

The hybrid structure of the Bhr-66 protein of *M. denitrificans* FJG1explains the

inconsistency in its gene annotation and phylogeny. This predicted structure has a distinct

bacteriohemerythrin of four alpha-helices arranged in a chimney shape at the N-terminus, which

was the segment likely identified as a bacteriohemerythrin by nBLAST, appended by a linear C-

terminal extension containing two structures with distinctive folds reminiscent of PAS domains

of s-box histidine kinases and response regulators. The overall structure suggests that Bhr-66 acts

as part of an oxygen-sensing two-component regulatory system that has been well described

previously in other bacteria (1, 21, 22). This may also explain the proximity of genes grouped as

response regulators in the neighbourhoods of several methanotroph *bhr*-66 genes (Fig. 4).

Similarly, the predicted Bhr-55 structure appears to have a bacteriohemerythrin segment at its N-

terminus with a long trailing section at the C-terminus, which includes up to seven folded PAS-

like domains (Fig. 6). Interestingly, no sequences related to *bhr*-55 were identified in other

genomes, suggesting that this protein may be unique to *M. denitrificans* FJG1.

Discussion

Methylomonas denitrificans FJG1 consumes methane and reduces nitrate to survive

hypoxia (14). Close examination of the *M. denitrificans* FJG1 genome revealed 10 homologs of

bhr genes, suggesting multiple potential functions associated with oxygen sensing and binding in

this bacterium. By providing a detailed examination of the phylogeny, gene neighbourhood,

expression, and predicted protein structures of the 10 *bhr* genes of *M. denitrificans* FJG1, we

provide compelling evidence that Bhr-00, a bacteriohemerythrin that appears nearly exclusive to

gammaproteobacterial methanotrophs, is likely the oxygen-binding protein responsible for

supporting methane oxidation in low oxygen environments. This assumption is partially based on

a close match of the predicted structure of Bhr-00 to Bhr-Bath, which was shown to bind and

deliver oxygen to pMMO enzymes *in vitro* (11). Elevated expression of *bhr*-00 at both the

transcript and protein levels in *M. denitrificans* FJG1 under hypoxic conditions further supports

this hypothesis. Also in support, a study of *Methylomonas* sp. LW13 with a deleted *bhr*-00 gene

was unable to grow in methane-oxygen counter gradient, but was able to grow normally in

planktonic culture with 50% methane:50% air (13). Aside from Bhr-00, this study identified *bhr*

homologs co-localized with denitrification (*bhr*-11), oxygen sensing (*bhr*-66), and chemotaxis

(*bhr*-77, -88) associated genes. These co-localized gene clusters showed increased mRNA

expression under hypoxic conditions with NMS, suggesting multiple roles for *bhr* domain-

containing proteins in *M. denitrificans* FJG1 for its survival and respiration under low oxygen,

denitrifying, conditions.

The hemerythrin originally discovered in marine worms, Hr, is encoded by a ~400 bp

gene for a monomeric protein that assembles into an octameric oxygen transport protein, causing

the pink hue of the worms' blood (2). The Bhr-Bath protein purified from *M. capsulatus* Bath

was crystallized as a monomer of similar size and shape to Hr and was shown to enhance oxygen

delivery from the cytoplasm to pMMO-enriched membrane fractions (9, 10). Purified Hr and

Bhr-Bath proteins turn an opaque reddish-purple or bright pink colour when bound to oxygen

(10, 23). This is the same hue as *M. denitrificans* FJG1 cells when grown to stationary phase in

liquid culture or on plates, further suggesting that increased expression of Bhr could be

responsible for oxygen binding and thus the change in their coloration beyond that of orange-

hued carotenoid proteins (24).

Aside from the identification and proposed role of Bhr-00, this study revealed several

diverse *bhr* gene homologs encoded by gammaproteobacterial methanotrophs. Based on our

findings, relevant potential functions of *bhr*-containing proteins in methanotrophs could include

oxygen binding and delivery to support methane oxidation and respiration, oxygen sensing,

nitrate reduction, and chemotaxis. Gammaproteobacterial methanotrophs often in the

Methylomonas genus are found in hypoxic to anoxic ecosystems, particularly in freshwater lakes

(25). The discovery and hypoxic regulation of the methanotroph-specific *bhr*-00 gene suggests a

mechanism to promote the binding and delivery of oxygen to support methane oxidation.

Moreover, the apparent restriction of *bhr*-00 to only a few related genera (*Methylomonas*,

Methylotuvimicrobium, *Methylomicrobium*) could explain their enrichment in these ecosystems

(25). From an environmental perspective, methanotrophs play a key role in biogeochemical

cycling of methane and nitrogen (26). Methane saturation of nutrient-impacted lakes and

reservoirs may further enrich for methanotrophs that rely on *bhr* to survive and thrive as these

ecosystems transition into hypoxia. The emergence of methanotroph phylotypes similar to *M.*

denitrificans FJG1 should be carefully monitored in nutrient impacted freshwater ecosystems as

they are bellwethers for hypoxia, methane cycling, and N_2O production (25, 27).

Materials and Methods

Phylogenetic analyses. The genome sequence of *M. denitrificans* FJGI (GenBank

accession CP014476A) was downloaded and analyzed using Geneious Prime 2019 (v1.2.3). The

FJG1 genome was found to contain 10 coding sequences (CDS) annotated as hemerythrin,

bacteriohemerythrin, hemerythrin family, or hemerythrin domain-containing proteins in the

GenBank and RefSeq annotation databases (Table 1). To facilitate further analyses, the 10

sequences were designated *bhr*-00 through *bhr*-99 in the order of location from the genome

origin of replication for the purposes of this study.

Unrooted phylogrammatic overview of all *M. denitrificans* FJG1 *bhr* BLASTn hits in

prokaryotes. Phylogenetic analyses of the 10 *bhr* sequences from *M. denitrificans* FJG1 were

performed using Geneious Prime 2019 (v1.2.3). The BLASTn plug-in was used with default

settings. The resulting dataset was filtered by excluding eukaryotic organisms and prokaryotic

organisms containing fewer than two *bhr* BLASTn hits.) Because the bhr-66 sequence is so

widespread among prokaryotes of multiple lineages, we chose to focus on a smaller dataset

including organisms that contain not just the ubiquitous *bhr*-66 but multiple hemerythrin

sequences that are more differentiated.

The resulting BLASTn hits were collected into a database for analysis and categorized by

aligning them to a concatenated sequence consisting of the 10 FJG1 genes. The reference

sequence was created by extracting the 10 *bhr* genes from the FJG1 genome and concatenating

them in order, separated by spacers of 100 base pairs. The concatenated reference sequence

eliminates duplicate matches and ensures that BLASTn hits are correlated to the FJG1 numbered

sequence with which they share the greatest sequence similarity (Supp. Data, Sheet 3). After

aligning all of the selected prokaryotic BLASTn hits with the reference sequence, an unrooted

phylogram was generated showing the inclusive overview of a broad range of *bhr* genes among

prokaryotic organisms (Geneious Prime 1.2.3, Global alignment with free end gaps, Cost matrix

51% similarity (5.0/-3.0), Tamura-Nei, Neighbor-Joining (NJ), No Outgroup, Gap open penalty

12, Gap extension penalty 3, Automatically determine direction). The accompanying data file of

sequence similarity presented in a distance matrix is available as Supp. Data, Sheet 3.

The BLASTn hits were used to locate prokaryotic organisms containing putative *bhr*

sequences with available complete genomes. These *bhr*-containing prokaryote genomes were

downloaded from GenBank and assembled into a local genome database. A circular alignment of

methanotroph genomes containing *bhr* sequences was generated using BLAST Ring Image

Generator (BRIGv0.95 default settings, Upper threshold 70%, Lower threshold 50%) to show the

location of their *bhr* sequences in relation to each other (Accession numbers CP014476,

CP033381, CP023669, CP002738, CP014360, CP022129, CP035467, CP024202, CP046565,

FO082060, NC_002977.6, AP017928). The circular alignment is available as Supp. Fig. S1.

Working in the locally assembled *bhr*-containing genome database, the *bhr* BLAST hit

sequences were used to locate full length CDSs in all collected genomes. These full sequences

including start and stop codons were extracted and compiled into a comprehensive *bhr* CDS

database to facilitate more accurate and complete phylogenetic analyses. Sequences were first

aligned using the Geneious Alignment tool (Global Alignment with free end gaps, Cost Matrix:

51%, default settings). Based on these alignments, three rooted cladograms were generated using

Geneious Prime 2019, one for each of the *M. denitrificans* FJG1 genes. CDSs with four or more

matching organisms (total n ≥ 5) (Geneious Tree Builder v1.2.3, Genetic Distance Model:

Tamura-Nei, Tree Build Method: Neighbour-joining, Resampling Method: 1,000 bootstrap,

Support Threshold: 50%, Image Scale: Substitutions per site). Less than four genomes could be

located encoding *bhr*-11, -22, -33, -44, -77 and/or -88, and therefore we could not perform a

detailed analysis of these genes across bacterial or other methanotroph species.

Cladograms describing *bhr* sequences *bhr*-00 and *bhr*-99 from methanotrophs were

rooted with the prokaryotic outlier *Chitinomonas* sp R3-44 (CP041730); however, in order to

include all prokaryotes for the widely distributed *bhr*-66 sequence, the cladogram was rooted

using hemerythrin sequences from a eukaryote outgroup, *Themiste pyroides* (KY007473.1,

KY007474.1, KY007475.1, KY007476.1, KY007477.1, KY007478.1, KY007479.1). For

phylogenetic comparison, we also generated 16S rRNA trees using the same methods.

Comparative images were created using Geneious Prime 2019 and colour was added to highlight

branch differences.

Identification and comparison of *bhr* gene neighbourhoods. Three of the 10 *bhr*

sequences (*bhr*-00, *bhr*-66, and *bhr*-99) were selected for gene neighbourhood analysis, as at

least five methanotroph genomes contained homologous copies to those found in *M. denitrificans*

FJG1. Less than four genomes encoding *bhr*-11, -22, -33, -44, -77 and/or -88, could be located

therefore we could not perform a meaningful comparative analysis of these gene neighbourhoods.

Approximately 16,000 bp surrounding each *bhr* sequence was extracted from each annotated

genome. In cases where multiple genomes were found for a species, the most recent annotated

genome assembly was used to leverage the most up-to-date annotations available. In cases where

full genomes were found, but lacked gene annotations, they were annotated in-house using

Geneious Prime 2019 (Default settings, Similarity > 50%), drawing from the in-house database

of all annotated methylotroph genomes downloaded for this project (> 2,000 genomes).

Comparative genomics based on these neighbourhood extractions was used to investigate each

bhr sequence. Annotations were compared manually to overcome inconsistencies in labeling

methodology. Colour schemes were used to identify sequences sharing similar or related

functional categories, while maintaining the annotation labeling of the original authors. An

accompanying comparison of the *bhr*-66 gene

neighbourhood in non-methanotroph genomes is available as Supp. Fig. S4.

RNAseq analysis. The RNAseq experiments were performed as described previously (14).

Briefly, *M. denitrificans* FJG1 was cultured in NMS or AMS medium at 30% methane to 70% air to

allow for natural oxygen depletion over time. The cultures achieved hypoxia and entry into

stationary phase by 48 h (Supp. Fig. S5). Replicate cultures (n=5) were destructively sampled at 24

and 48 h time points to determine differential mRNA expression as a function of oxygen depletion.

Cells were collected and mRNA was purified, converted to cDNA, and processed for transcriptome

analysis using Ilumina Hi-Seq 2000 sequencing technology as described (13). Geneious Prime 2019

(v1.2.3) was employed with the DESeq2 plug-in (default model settings "Parametric") to re-calculate

expression values and differential expression (14). PCA plots revealed zero outliers with < 7% total

within-treatment variance. TPM values for transcripts at 24, 48 and 72 h time points for *bhr*

homologs and members of their gene neighborhoods were calculated using DESeq2 in Geneious

Prime 2019 (v1.2.3), and the ratio of differential expression of *bhr* homologs between 24 and 48 h

were determined using R.

Protein extraction and peptide preparation. For each of the three treatments we

prepared tryptic digests from six biological replicates following the filter-aided sample

preparation (FASP) protocol described by Wisniewski et al. (28). SDT-lysis buffer (4% (w/v)

SDS, 100 mM Tris-HCl pH 7.6, 0.1 M DTT) was added in a 1:10 sample/buffer ratio to the

sample pellets. Samples were heated for lysis to 95° C for 10 minutes followed by pelleting of

debris for 5 min at 21,000 x g. 30 µl of the cleared lysate were mixed with 200 µl of UA solution

(8 M urea in 0.1 M Tris/HCl pH 8.5) in a 10 kDa MWCO 500 µl centrifugal filter unit (VWR

International) and centrifuged at 14,000 x g for 40 min. 200 µl of UA solution were added again

and centrifugal filter spun at 14,000 x g for 40 min. 100 µl of IAA solution (0.05 M

iodoacetamide in UA solution) were added to the filter and incubated at 22° C for 20 min. The

IAA solution was removed by centrifugation and the filter was washed three times by adding 100

µl of UA solution and then centrifuging. The buffer on the filter was then changed to ABC (50

mM Ammonium Bicarbonate), by washing the filter three times with 100 µl of ABC. 1 µg of MS

grade trypsin (Thermo Scientific Pierce, Rockford, IL, USA) in 40 µl of ABC were added to the

filter and filters incubated overnight in a wet chamber at 37° C. The next day, peptides were

eluted by centrifugation at 14,000 x g for 20 min, followed by addition of 50 µl of 0.5 M NaCl

and again centrifugation. Peptides were desalted using C18 spin columns (Thermo Scientific

Pierce, Rockford, IL, USA) according to the manufacturer's instructions. Approximate peptide

concentrations were determined using the Pierce Micro BCA assay (Thermo Scientific Pierce,

Rockford, IL, USA) following the manufacturer's instructions.

Proteomics 1D-LC-MS/MS. Samples were analyzed by 1D-LC-MS/MS using a block-

randomized design as outlined by Oberg and Vitek (29). Two blank runs were done between

samples to reduce carry over. For each run 800 ng of peptide were loaded onto a 2 cm, 75 µm ID

C18 Acclaim® PepMap 100 pre-column (Thermo Fisher Scientific) using an EASY-nLC 1000

Liquid Chromatograph (Thermo Fisher Scientific) set up in 2-column mode. The pre-column was

connected to a 50 cm x 75 µm analytical EASY-Spray column packed with PepMap RSLC C18,

2µm material (Thermo Fisher Scientific), which was heated to 35° C via the integrated heating

module. The analytical column was connected via an Easy-Spray source to a Q Exactive Plus

hybrid quadrupole-Orbitrap mass spectrometer (Thermo Fisher Scientific). Peptides were

separated on the analytical column at a flow rate of 225 nl/min using a 260 min gradient going

from buffer A (0.2% formic acid, 5% acetonitrile) to 20% buffer B (0.2% formic acid in

acetonitrile) in 200 min, then from 20 to 35% B in 40 min and ending with 20 min at 100% B.

Eluting peptides were ionized with electrospray ionization (ESI) and analyzed in the Q Exactive

Plus. Full scans were acquired in the Orbitrap at 70,000 resolution. MS/MS scans of the 15 most

abundant precursor ions were acquired in the Orbitrap at 17,500 resolution. The mass (m/z)

445.12003 was used as lock mass. Ions with charge state +1 were excluded from MS/MS

analysis. Dynamic exclusion was set to 30 sec. Roughly 160,000 MS/MS spectra were acquired

per sample run.

Protein identification, quantification and statistics. For protein identification a

database was created using all protein-coding gene sequences from the complete genome of

Methylomonas denitrificans strain FJG1 (Genbank Identifier CP014476.1). The database was

submitted to the PRIDE repository (see below). For protein identification MS/MS spectra were

searched against the database using the Sequest HT node in Proteome Discoverer version

2.0.0.802 (Thermo Fisher Scientific) with the following parameters: Trypsin (Full), max. 2

missed cleavages, 10 ppm precursor mass tolerance, 0.1 Da fragment mass tolerance and max. 3

equal dynamic modifications per peptide. The following three dynamic modifications were

considered: oxidation on M (+15.995 Da), carbamidomethyl on C (+57.021 Da) and acetyl on

protein N-terminus (+42.011 Da). False discovery rates (FDRs) for peptide spectral matches

(PSMs) were calculated and filtered using the Percolator Node in Proteome Discoverer (30).

Percolator was run with the following settings: Maximum Delta Cn 0.05, a strict target FDR of

0.01, a relaxed target FDR of 0.05 and validation based on q-value. Search results for all 18

samples were combined into a multiconsensus report using the FidoCT node in Proteome

Discoverer to restrict the protein-level FDR to below 5% (FidoCT q-value <0.05)(31). Based on

these filtering criteria a total of 1833 proteins were identified in all samples together.

400   For protein quantification normalized spectral abundance factors (NSAFs) were

calculated based on number of PSMs per protein using the method described by Florens et al.

(32) and multiplied by 10,000. The NSAFx10,000 gives the relative abundance of a protein in a

sample as a fraction of 10,000. The table with NSAFx10,000 values was loaded into the Perseus

software (version 1.5.1.6, http://www.perseus-framework.org/doku.php) (33) an annotation row

was added to group replicates by treatment and proteins that did not have abundance values >0

for all replicates in at least one treatment group were removed (1831 proteins remained).

NSAFx10,000 values were log2 transformed. Missing values produced by log2(0) were replaced

by sampling from a normal distribution assuming that the missing values are on the lower end of

abundance (normal distribution parameters in Perseus: width 0.3, down shift 1.8, do separately

for each column). A t-test with permutation-based FDR calculation to account for the multiple

hypothesis testing problem was used to detect proteins that differed significantly in their

abundance level between two treatments. The following parameters were used for the test:

groupings were not preserved for randomizations, both sides, 250 randomizations, FDR of 1%

and s0 of 0.

415   **Protein Structure Predictions**. Predicted protein structures corresponding to the *bhr*

gene loci for *M. denitrificans* FJG1 were downloaded from AlphaFold2 (alphafold.ebi.ac.uk) (34,

35). Nine structures were predicted that corresponded to Bhr-00 (A0A140E3I1), Bhr-11

(A0A140E4Z5), Bhr-22 (A0A126T104), Bhr-33 (A0A126T1L7), Bhr-44 (A0A126T494), Bhr-

66 (A0A126T8C7), Bhr-77 (A0A126T8U2), Bhr-88 (A0A126T8Y8), and Bhr-99

(A0A140E6J4). The predicted structure of Bhr-66 was further investigated by extracting the

hemerythrin section at the N-terminus and separating it from the C-terminal section of the gene

into two unique nucleotide subsequences. The *bhr*-like subsequence (~400 bp) and the C-terminal

tail subsequence (~1,200 bp) were used for BLASTn searches in the GenBank nucleotide

database (Geneious Prime 2019, v1.2.3 with BLAST plugin, default settings, filtered by Grade >

40%) and the annotations from matching sequences were examined for each individual

subsection of the atypical *bhr*-66 gene, revealing a hemerythrin domain at the N-terminus and a

chain of multiple PAS sensing domains at the C-terminal end. No corresponding structure for

Bhr-55 was available in the AlphaFold database, thus an alternative protein structure prediction

tool was used (https://robetta.bakerlab.org/). The FASTA sequence of the *bhr*-55 gene was

uploaded to the online tool and prediction was performed using RoseTTAFold, with default

settings on the translated protein. The resulting visualization was coloured according to Error

Estimate in the Robetta interface.

The predicted Bhr-00 (A0A140E3I1) structure and the Bhr-Bath structure (PDB ID:

4XPX) were downloaded from the RCSB Protein Data Bank (36) and imported into the MolStar

Viewer (37) to compare their structures. Protein structures were superposed using a single Lys

residue in each amino acid sequence (Lys 68, Bhr-Bath and Lys 70, Bhr-00). The Mol*

animation is available as Supp. Movie. The AA alignment of FJG1 Bhr-00 and Bhr-Bath is

available as Supp. Fig. S2.

Acknowledgments: We thank Marc Strous (Univ. Calgary) for access to the proteomics

equipment. This work was funded by the Canada First Research Excellence Fund Future Energy

Systems to DS and LYS, an NSERC Discovery grant to LYS, and a Mitacs Accelerate grant to

DS. The purchase of the proteomics equipment was supported by a grant of the Canadian

Foundation for Innovation to Marc Strous. MK was supported by a NSERC Banting Postdoctoral

Fellowship. The funders had no role in study design, data collection and interpretation, or the

decision to submit the work for publication.

Conflict of interest: the authors have no conflicts of interest to declare.

Author Contributions: C.W., D.S. and L.Y.S. participated in the planning of the study. C.W.

performed the phylogenetic analyses, comparative genomics, and drafted the manuscript. K.D.K.

performed the transcriptome and proteome experiments, analyzed data, and edited the

manuscript. M.K. provided the proteomic data, statistical analysis, and methodology description.

D.S. and L.Y.S. provided financial support, critical analysis, writing and editing of the

manuscript. All authors have read and approved the manuscript for publication.

Originality statement: None of the work in this manuscript has been published or

submitted for publication elsewhere.

Data Availability Statement

Transcriptome data can be found in GenBank (accession number SRX696231). The genomic

datasets can be found by provided accession numbers in the article. Additional details on data

used in this study are in the Supp. Data file. The mass spectrometry proteomics data and the

protein sequence database have been deposited to the ProteomeXchange Consortium (38) via the

PRIDE partner repository with the dataset identifier PXD004041 [Reviewer access: log in at

http://www.ebi.ac.uk/pride/archive/ with Username: reviewer61804@ebi.ac.uk and Password:

B6gFiqkh].

References

1.   Alvarez-Carreño C, Alva V, Becerra A, Lazcano A. 2018. Structure, function and evolution of
the hemerythrin-like domain superfamily. Protein Sci 27:848-860.
2.   Costa-Paiva EM, Whelan NV, Waits DS, Santos SR, Schrago CG, Halanych KM. 2017. Discovery
and evolution of novel hemerythrin genes in annelid worms. BMC Evol Biol 17.
3.   Baert JL, Britel M, Sautiere P, Malecha J. 1992. Ovohemerythrin, a major 14-kDa yolk protein
distinct from vitellogenin in leech. Eur J Biochem 209:563-569.
4.   Kendall JJ, Barrero-Tobon AM, Hendrixson DR, Kelly DJ. 2014. Hemerythrins in the
microaerophilic bacterium *Campylobacter jejuni* help protect key iron-sulphur cluster
enzymes from oxidative damage. Environ Microbiol 16:1105-1121.
5.   Demuynck S, Li KW, Vanderschors R, Dhainautcourtois N. 1993. Amino acid sequence of the
small cadmium-binding protein (MP II) from *Nereis diversicolour* (annelida, polychaeta) -
Evidence for a myohemerythrin structure. Eur J Biochem 217:151-156.
6.   Xiong JJ, Kurtz DM, Ai JY, Sanders-Loehr J. 2000. A hemerythrin-like domain in a bacterial
chemotaxis protein. Biochemistry 39:5117-5125.
7.   French CE, Bell JML, Ward FB. 2008. Diversity and distribution of hemerythrin-like proteins
in prokaryotes. FEMS Microbiol Lett 279:131-145.

8.   Rahalkar MC, Bahulikar RA. 2018. Hemerythrins are widespread and conserved for
methanotrophic guilds. Gene Rep 11:250-254.
9.   Chen KHC, Wu HH, Ke SF, Rao YT, Tu CM, Chen YP, Kuei KH, Chen YS, Wang VCC, Kao WC,
Chan SI. 2012. Bacteriohemerythrin bolsters the activity of the particulate methane
monooxygenase (pMMO) in *Methylococcus capsulatus* (Bath). J Inorg Biochem 111:10-17.
10.   Kao WC, Wang VCC, Huang YC, Yu SSF, Chang TC, Chan SI. 2008. Isolation, purification and
characterization of hemerythrin from *Methylococcus capsulatus* (Bath). J Inorg Biochem
102:1607-1614.
11.   Chen KHC, Chuankhayan P, Wu HH, Chen CJ, Fukuda M, Yu SSF, Chan SI. 2015. The
bacteriohemerythrin from *Methylococcus capsulatus* (Bath): Crystal structures reveal that
Leu114 regulates a water tunnel. J Inorg Biochem 150:81-89.
12.   Nariya S, Kalyuzhnaya MG. 2020. Hemerythrins enhance aerobic respiration in
Methylomicrobium alcaliphilum 20Z [R] a methane-consuming bacterium. FEMS Microbiol Lett
367.
13.   Beals DG, Puri AW. 2024. Linking methanotroph phenotypes to genotypes using a simple
spatially resolved model ecosystem. ISME J 18.
14.   Kits KD, Klotz MG, Stein LY. 2015. Methane oxidation coupled to nitrate reduction under
hypoxia by the Gammaproteobacterium *Methylomonas denitrificans*, sp. nov. type strain
FJG1. Environ Microbiol 17:3219-3232.
15.   Kits KD, Campbell DJ, Rosana AR, Stein LY. 2015. Diverse electron sources support
denitrification under hypoxia in the obligate methanotroph *Methylomicrobium album* strain
BG8. Front Microbiol 6.
16.   Li B, Tao Y, Mao ZD, Gu QJ, Han YX, Hu BL, Wang HW, Lai AX, Xing P, Wu QL. 2023. Iron
oxides act as an alternative electron acceptor for aerobic methanotrophs in anoxic lake
sediments. Water Res 234.
17.   Kalyuzhnaya MG, Yang S, Rozova ON, Smalley NE, Clubb J, Lamb A, Gowda GAN, Raftery D, Fu
Y, Bringel F, Vuilleumier S, Beck DAC, Trotsenko YA, Khmelenina VN, Lidstrom ME. 2013.
Highly efficient methane biocatalysis revealed in a methanotrophic bacterium. Nature Comm
4:7.
18.   Kitanishi K. 2022. Bacterial hemerythrin domain-containing oxygen and redox sensors:
Versatile roles for oxygen and redox signaling. Front Molec Biosci 9.
19.   Barquera B. 2014. The sodium pumping NADH:quinone oxidoreductase (Na+-NQR), a
unique redox-driven ion pump. J Bioenerg Biomembr 46:289-298.
20.   Alvarez-Carreño C, Becerra A, Lazcano A. 2016. Molecular evolution of the oxygen-binding
hemerythrin domain. Plos One 11.
21.   Taylor BL, Zhulin IB. 1999. PAS domains: Internal sensors of oxygen, redox potential, and
light. Microbiol Molec Biol Rev 63:479-+.
22.   Yamada S, Sugimoto H, Kobayashi M, Ohno A, Nakamura H, Shiro Y. 2009. Structure of PAS-
linked histidine kinase and the response regulator complex. Structure 17:1333-1344.
23.   Stenkamp RE, Sieker LC, Jensen LH, McCallum JD, Sandersloehr J. 1985. Active site structures
of deoxyhemerythrin and oxyhemerythrin. Proc Natl Acad Sci 82:713-716.
24.   Oshkin IY, Tikhonova EN, Suleimanov RZ, Ashikhmin AA, Ivanova AA, Pimenov NV, Dedysh
SN. 2023. All kinds of sunny colors synthesized from methane: genome-encoded carotenoid
production by *Methylomonas* species. Microorganisms 11.
25.   Reis PCJ, Tsuji JM, Weiblen C, Schiff SL, Scott M, Stein LY, Neufeld JD. 2024. Enigmatic
persistence of aerobic methanotrophs in oxygen-limiting freshwater habitats. ISME J 18.
26.   Stein LY. 2020. The long-term relationship between microbial metabolism and greenhouse
gases. Trends Microbiol 28:500-511.
27.   Schorn S, Graf JS, Littmann S, Hach PF, Lavik G, Speth DR, Schubert CJ, Kuypers MMM,
Milucka J. 2024. Persistent activity of aerobic methane-oxidizing bacteria in anoxic lake

536   waters due to metabolic versatility. Nat Commun 15.

537   28.   Wisniewski JR, Zougman A, Nagaraj N, Mann M. 2009. Universal sample preparation method
538   for proteome analysis. Nat Meth 6:359-U60.

539   29.   Oberg AL, Vitek O. 2009. Statistical design of quantitative mass spectrometry-based
540   proteomic experiments. J Proteome Res 8:2144-2156.

541   30.   Spivak M, Weston J, Bottou L, Käll L, Noble WS. 2009. Improvements to the percolator
542   algorithm for peptide identification from shotgun proteomics data sets. J Proteome Res
543   8:3737-3745.

544   31.   Serang O, MacCoss MJ, Noble WS. 2010. Efficient marginalization to compute protein
545   posterior probabilities from shotgun mass spectrometry data. J Proteome Res 9:5346-5357.

546   32.   Florens L, Carozza MJ, Swanson SK, Fournier M, Coleman MK, Workman JL, Washburn MP.
547   2006. Analyzing chromatin remodeling complexes using shotgun proteomics and
548   normalized spectral abundance factors. Methods 40:303-311.

549   33.   Cox J, Mann M. 2012. 1D and 2D annotation enrichment: a statistical method integrating
550   quantitative proteomics with complementary high-throughput data. BMC Bioinformatics 13.

551   34.   Jumper J, Evans R, Pritzel A, Green T, Figurnov M, Ronneberger O, Tunyasuvunakool K, Bates
552   R, Žídek A, Potapenko A, Bridgland A, Meyer C, Kohl SAA, Ballard AJ, Cowie A, Romera-
553   Paredes B, Nikolov S, Jain R, Adler J, Back T, Petersen S, Reiman D, Clancy E, Zielinski M,
554   Steinegger M, Pacholska M, Berghammer T, Bodenstein S, Silver D, Vinyals O, Senior AW,
555   Kavukcuoglu K, Kohli P, Hassabis D. 2021. Highly accurate protein structure prediction with
556   AlphaFold. Nature 596:583-589.

557   35.   Varadi M, Bertoni D, Magana P, Paramval U, Pidruchna I, Radhakrishnan M, Tsenkov M, Nair
558   S, Mirdita M, Yeo J, Kovalevskiy O, Tunyasuvunakool K, Laydon A, Zídek A, Tomlinson H,
559   Hariharan D, Abrahamson J, Green T, Jumper J, Birney E, Steinegger M, Hassabis D, Velankar
560   S. 2023. AlphaFold Protein Structure Database in 2024: providing structure coverage for
561   over 214 million protein sequences. Nuc Acid Res 52:D368-D375.

562   36.   Burley SK, Berman HM, Bhikadiya C, Bi CX, Chen L, Di Costanzo L, Christie C, Dalenberg K,
563   Duarte JM, Dutta S, Feng ZK, Ghosh S, Goodsell DS, Green RK, Guranovic V, Guzenko D,
564   Hudson BP, Kalro T, Liang YH, Lowe R, Namkoong H, Peisach E, Periskova I, Prlic A, Randle C,
565   Rose A, Rose P, Sala R, Sekharan M, Shao CH, Tan LH, Tao YP, Valasatava Y, Voigt M,
566   Westbrook J, Woo J, Yang HW, Young J, Zhuravleva M, Zardecki C. 2019. RCSB Protein Data
567   Bank: biological macromolecular structures enabling research and education in fundamental
568   biology, biomedicine, biotechnology and energy. Nuc Acid Res 47:D464-D474.

569   37.   Sehnal D, Bittrich S, Deshpande M, Svobodová R, Berka K, Bazgier V, Velankar S, Burley SK,
570   Koča J, Rose AS. 2021. Mol* Viewer: modern web app for 3D visualization and analysis of
571   large biomolecular structures. Nucleic Acids Res 49:W431-W437.

572   38.   Vizcaíno JA, Deutsch EW, Wang R, Csordas A, Reisinger F, Ríos D, Dianes JA, Sun Z, Farrah T,
573   Bandeira N, Binz PA, Xenarios I, Eisenacher M, Mayer G, Gatto L, Campos A, Chalkley RJ,
574   Kraus HJ, Albar JP, Martinez-Bartolomé S, Apweiler R, Omenn GS, Martens L, Jones AR,
575   Hermjakob H. 2014. ProteomeXchange provides globally coordinated proteomics data
576   submission and dissemination. Nat Biotechnol 32:223-226.

578 **Table 1**. Hemerythrin genes identified in the *Methylomonas denitrificans* FJG1 genome (CP014476)

Name	Locus_tag/ GI_number	Min	Max	Length	Direction	Annotation	Protein_id	Protein	GO_function
bhr-00	JT25_RS00380 GI:998900537	85767	85372	396	R	hemerythrin family protein	WP_036277397.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-11	JT25_RS03015	663113	663445	333	F	hemerythrin domain-containing protein	WP_132325259.1	hemerythrin domain-containing protein	-
bhr-22	JT25_RS04600	989403	989813	411	F	bacteriohemerythrin	WP_036273626.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-33	JT25_RS05590	1204483	1204097	387	R	hemerythrin family protein	WP_036274438.1	hemerythrin family protein	-
bhr-44	JT25_RS10375	2251509	2251123	387	R	hemerythrin family protein	WP_036280274.1	hemerythrin family protein	-
bhr-55	JT25_RS11125 GI:1011416966	2420834	2416305	4530	R	bacteriohemerythrin	WP_062328608.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-66	JT25_RS17815	3859160	3860944	1785	F	bacteriohemerythrin	WP_062329310.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-77	JT25_RS18710	4064071	4063667	405	R	bacteriohemerythrin	WP_036274237.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-88	JT25_RS18715	4064508	4064095	414	R	hemerythrin family protein	WP_036274239.1	bacteriohemerythrin	GO:0046872 - metal ion binding
bhr-99	JT25_RS21450	4674029	4673625	405	R	hemerythrin family protein	WP_052142344.1	hemerythrin family protein	-

580   **Table 2.** Differential expression of *bhr* gene homologues from *M. denitrificans* FJG1 based on

581   RNAseq data collected after 24 and 48 h growth in NMS or AMS media with methane.

	Nitrate Treatment (NMS)				
	Raw Expression (TPM)		Differential Expression 24h vs 48h		
	24h	48h	log2 ratio	adjusted p-value	normalized log2 ratio
bhr-00	4,657.50	13,150.82	1.35	0.000	3.33
bhr-11	38.22	62.58	0.47	0.034	2.45
bhr-22	93.16	61.85	-0.65	0.000	1.33
bhr-33	29.11	41.03	0.50	0.051	2.48
bhr-44	83.43	72.20	-0.28	0.104	1.70
bhr-55	30.82	46.69	0.41	0.000	2.39
bhr-66	48.44	63.83	0.28	0.002	2.26
bhr-77	107.16	176.26	0.62	0.000	2.60
bhr-88	50.66	80.73	0.59	0.000	2.57
bhr-99	170.36	172.87	-0.05	0.732	1.93
rpoB	418.02	113.34	-1.98	0.000	-

583
584
585
586

	Ammonium Treatment (AMS)				
	Raw Expression (TPM)		Differential Expression 24h vs 48h		
	24h	48h	log2 ratio	adjusted p-value	normalized log2 ratio
bhr-00	7,706.16	28,626.86	1.91	0.000	3.21
bhr-11	75.47	23.55	-1.14	0.000	0.16
bhr-22	122.98	57.33	-0.57	0.000	0.73
bhr-33	22.58	39.44	0.50	0.004	1.80
bhr-44	74.32	65.09	-0.45	0.000	0.85
bhr-55	32.34	36.47	0.25	0.000	1.55
bhr-66	45.44	53.34	0.17	0.013	1.47
bhr-77	146.80	119.38	0.07	0.470	1.37
bhr-88	64.90	73.10	0.22	0.071	1.52
bhr-99	193.38	181.93	-0.15	0.036	1.15
rpoB	421.34	162.99	-1.30	0.000	-

Figure Legends

Figure 1. Unroooted phylogram of prokaryotic genes related to the 10 *bhr* genes from *M.*

denitrificans FJG1. Homologs were identified from the GenBank database using BLASTn

(Geneious Prime 2019 v1.2.3). Sequences from *M. denitrificans* FJG1 are highlighted in pink and

genes from other proteobacterial methanotrophs are highlighted in blue. Non-methanotrophs are

colored grey.

Figure 2. Cladogram created with full-length CDSs matching the *bhr*-66 gene of *M. denitrificans*

FJG1 (rooted with eukaryotic outgroup *Themiste pyroides* (KY007473 – KY007479; Geneious

Prime 2019 v1.2.3). Green: *Deltaproteobacteria*, blue: *Betaproteobacteria*, red:

Gammaproteobacteria, purple: *Actinobacteria*. Genes from methanotrophic organisms are

highlighted with a yellow bar.

Figure 3. Cladograms of phylogenetic pairings between *bhr*-00 and *bhr*-66 with 16S rRNA genes

in methanotrophs and of *bhr*-00 with *bhr*-99 genes (Geneious Prime 2019 v1.2.3). Yellow

highlights indicate multiple branch differences, green highlights indicate one branch difference,

and blue highlights indicate consistent branch topology.

Figure 4. Gene neighborhoods and colour-coded gene annotations surrounding *bhr*-00, *bhr*-66,

and *bhr*-99 (Geneious Prime 2019 v1.2.3). Key genes were colour-coded to discriminate

functional categories as shown above each gene neighborhood.

Figure 5. Gene neighborhoods of *bhr*-00, -11, -66, and -77/-88 with gene expression and protein

abundance data. Gene neighborhoods are shown on the left in their operon structures and are

colour-coded by functional group as per the legend. Up arrows indicate genes encoded on the

forward DNA strand, and down arrows indicate genes encoded on the reverse DNA strand.

Columnar data displayed to the right represents mRNA and protein expression abundance for *M*.

denitrificans FJG1 cells transitioning from oxic (24 h) through hypoxic (48 h) to anoxic (72 h)

conditions in rows from left to right as noted in the column headings. mRNA expression (TPM)

in NMS: low (yellow) to high (red) and AMS: low (yellow) to high (green). Protein expression

(NSAF) for NMS: low (pink) to high (purple). Rows are aligned with the center point of

corresponding gene icons in the neighborhoods to the left. CDS lengths are proportional and gaps

are preserved. (Neighborhood genes colourized in Geneious Prime 2019 v1.2.3. Data columns

prepared in LibreOffice Calc. Components organized and legend added in GIMP 2.10.)

Figure 6. AlphaFold2 Predicted Protein Structures visualized in Mol* Viewer. **A.** Bhr protein

structures predicted from *M. denitrificans* FJG1 genes from the AlphaFold2 database. Br-55 was

not found using AlphaFold2 and was instead created using Robetta, as described in Methods. **B.**

Structural comparison of predicted AlphaFold2 structure for Bhr-00 (ID: A0A140E3I1) with

experimentally determined Bhr-Bath structure (Chen et al. 2015, PDB ID: 4XPX). Diiron

cofactor is shown in orange. Animated superposition in .mp4 format is available as a

Supplemental Movie.

Supplemental Figure Legends

Supp. Fig. S1. Circular alignment of methanotroph genomes created with Blast Ring Image

Generator (BRIG). The location of *bhr* genes in *M. denitrificans* FJG1 and other methanotroph

genomes are designated by colored bars.

Supp. Fig. S2. Alignment of Bhr-00 protein sequence from *Methylomonas denitrificans* FJG1

with Bhr-Bath protein sequence from *Methylococcus capsulatus* Bath.

Supp. Fig. S3. Gene neighborhoods of bhr-22, -33, -44, -55 with corresponding gene expression

and protein abundance data. Gene neighborhoods are shown on the left in their operon structures

and are colour-coded by functional group as per the legend. Up arrows indicate genes encoded on

the forward DNA strand, and down arrows indicate genes encoded on the reverse DNA strand.

Columnar data displayed to the right represents mRNA and protein expression abundance for M.

denitrificans FJG1 cells transitioning from oxic (24 h) through hypoxic (48 h) to anoxic (72 h)

conditions in rows from left to right as noted in the column headings. mRNA expression (TPM)

in NMS: low (yellow) to high (red) and AMS: low (yellow) to high (green). Protein expression

(NSAF) for NMS: low (pink) to high (purple). Rows are aligned with the center point of

corresponding gene icons in the neighborhoods to the left. CDS lengths are proportional and gaps

are preserved. (Neighborhood genes colourized in Geneious Prime 2019 v1.2.3. Data columns

prepared in LibreOffice Calc. Components organized and legend added in GIMP 2.10.)

Supp. Fig. S4. Gene neighborhoods of *bhr*-66 of non-methanotroph genomes.

Supp. Fig. S5. Growth curve of *M. denitrificans* FJG1 on NMS or AMS media. (A) Growth over time. (B) Oxygen consumption over time. Reprinted with permission from Environmental Microbiology (14).

Supp. Table 1. Differential expression of neighboring genes upstream and downstream of the *bhr*-00, -11, -66, and -77/-88 homologs from *M. denitrificans* FJG1 based on RNAseq data at 24 h (oxic) versus 48 h (hypoxic) growth in NMS and AMS media.

Supp. Data. Spreadsheet in .xls format containing metadata: Sheet 1 - Metadata; Sheet 2- list of methanotroph genes used to create rooted phylograms; Sheet 3- distance matrix for BLAST overview tree (Fig. 1); Sheet 4- percent identity of FJG1 hemerythrins found in other methanotrophs; Sheet 5- summary of previous research regarding prokaryotic hemerythrins; Sheet 6 - analyzed proteome data.

Supp. Movie. MP4 showing animated overlay of AlphaFold structure for Bhr-00 and crystal structure of Bhr-Bath proteins.

bhr-00 neighbourhood

Gene Locus	Gene Product	NMS TPM 24 hr	48 hr	72 hr	NMS NSAF×10K 24 hr	48 hr	72 hr	AMS TPM 24 hr	48 hr	72 hr
JT25_RS00360	SURF1 family protein									
JT25_RS00355	twin-transmembrane helix small protein									
JT25_RS00365	cytochrome c oxidase subunit 3									
JT25_RS00365	cytochrome c oxidase assembly protein									
JT25_RS00370	ctaD, cytochrome c oxidase subunit I									
JT25_RS00380	coxB, cytochrome c oxidase subunit II									
JT25_RS00385	homerylohm family protein									
JT25_RS00390	bioD, debiobiotin synthase									
JT25_RS00390	bioC, malonyl-ACP O-methyltransferase BioC									
JT25_RS00395	bioH, pimeloyl-ACP methyl ester esterase BioH									
JT25_RS00400	bioF, 8-amino-7-oxononanoate synthase									
JT25_RS00405	bioB, biotin synthase BioB									

bhr-66 neighbourhood

Gene Locus	Gene Product
JT25_RS17790	EAL domain-containing protein
JT25_RS17795	AEC family transporter
JT25_RS17800	competence/damage-inducible protein A
JT25_RS17805	ABC transporter ATP binding protein
JT25_RS17810	hypothetical protein
JT25_RS17815	bacteriohemerythrin
JT25_RS17820	response regulator, histidine kinase
JT25_RS17825	polyphosphate kinase
JT25_RS17830	hypothetical protein
JT25_RS17835	hypothetical protein
JT25_RS17840	hypothetical protein

bhr-11 neighbourhood

Gene Locus	Gene Product
JT25_RS02985	nitrate reductase subunit alpha
JT25_RS02990	narH, nitrate reductase subunit beta
JT25_RS02995	narJ, nitrate reductase molybdenum cofactor
JT25_RS03000	narI, respiratory nitrate reductase subunit gamma
JT25_RS03005	multicopper oxidase domain-containing protein
JT25_RS03010	hypothetical protein
JT25_RS03015	hemerythrin domain-containing protein
JT25_RS03020	HPP family protein
JT25_RS03025	NAD(P)/FAD-dependent oxidoreductase
JT25_RS03030	PhzF family phenazine biosynthesis protein
JT25_RS03035	nirD, nitrite reductase small subunit NirD
JT25_RS03040	nirB, nitrite reductase large subunit
JT25_RS03045	formate/nitrite transporter family protein

bhr-77, bhr-88 neighbourhood

Gene Locus	Gene Product
JT25_RS16680	hypothetical protein
JT25_RS16685	response regulator
JT25_RS16690	STAS domain-containing protein
JT25_RS16695	chemotaxis protein CheA
JT25_RS16700	HAMP domain-containing protein
JT25_RS16705	purine-binding chemotaxis protein CheW
JT25_RS16710	bacteriohemerythrin
JT25_RS16715	hemerythrin family protein
JT25_RS16720	methyl-accepting chemotaxis protein
JT25_RS16725	formate–tetrahydrofolate ligase
JT25_RS16730	ribose-5-phosphate isomerase RpiA
JT25_RS16735	threonine ammonia-lyase, biosynthetic
JT25_RS16740	fucosamine kinase family protein
JT25_RS16745	glutamine–tRNA ligase/YqeY domain fusion

A.

Bhr-00   Bhr-11   Bhr-22   Bhr-33   Bhr-44

Bhr-66   Bhr-77   Bhr-88   Bhr-99

AlphaFold2

Robetta

Bhr-55

B.

Bhr-Bath   Bhr-00

Peptide Legend

Bhr-00 superposed with Bhr-Bath