A developmental brain-wide screen identifies retrosplenial cortex as a key player in the emergence of persistent memory

Abbreviated Title: Circuit mechanisms underlying infantile amnesia

Authors: Benita Jin[1,2], Michael W. Gongwer[1,3,4], Lilit Ohanian[1], Lucinda Holden-Wingate[1], Bryan Le[1], Alfonso Darmawan[1], Yuka Nakayama[1], Sophia A. Rueda Mora[1] and Laura A. DeNardo[1,5]*

1. Department of Physiology, University of California Los Angeles, Los Angeles, CA 90095

2. Molecular Cellular and Integrative Physiology Interdepartmental Program, University of California Los Angeles, Los Angeles, CA 90095

3. Neuroscience Interdepartmental Program, University of California Los Angeles, Los Angeles, CA 90095

4. Medical Scientist Training Program, University of California Los Angeles, Los Angeles, CA 90095

5. Department of Neurobiology, University of California Los Angeles, Los Angeles, CA 90095

*Address correspondence to: ldenardo@ucla.edu

Number of pages: 33

Numbers of figures: 6 main, 6 supplemental

Number of tables: 2

Keywords: infantile amnesia, memory, retrosplenial cortex, neural circuits, brain clearing, light sheet fluorescence microscopy

Summary: 150 words

Introduction: 571 words

Discussion: 1642 words

Summary

Memories formed early in life are short-lived while those formed later persist. Recent work revealed that infant memories are stored in a latent state. But why they fail to be retrieved is poorly understood. Here we investigated brain-wide circuit mechanisms underlying infantile amnesia in mice. We performed a screen that combined activity-dependent neuronal tagging at different postnatal ages, tissue clearing and light sheet microscopy. We observed striking developmental transitions in the organization of fear memory networks and changes in the activity and functional connectivity of the retrosplenial cortex (RSP) that aligned with the emergence of persistent memory. 7 days after learning, chemogenetic reactivation of tagged RSP ensembles enhanced memory in adults but not in infants. But after 33 days, reactivating infant-tagged RSP ensembles recovered forgotten memories. These studies show that RSP ensembles store latent infant memories, reveal the time course of RSP functional maturation, and suggest that immature RSP functional networks contribute to infantile amnesia.

Introduction

Infantile amnesia describes a process wherein episodic memories formed early in life are rapidly forgotten[1–3]. In contrast, memories formed later can last for decades. In humans, memories become persistent around age five[4]. Infantile amnesia is not a uniquely human phenomenon[5,6]. In rodents, memories become persistent during the fourth postnatal week[7–11]. But even though early formed memories cannot be consciously recalled, they can influence later experiences by promoting the maturation of learning systems and altering the mechanisms of later memory encoding[12–14]. Their continuing influence suggests that early-formed memories leave an enduring trace in the brain.

Several studies showed that infant memories are not erased but rather inaccessible. Forgotten memories leave a molecular trace in the amygdala[15]. Providing strong reminders of the original memory[13,16,17] or optogenetically reactivating infant memory ensembles in the hippocampus[11] can recover lost infant memories. Importantly, early adversity such as neglect or abuse can accelerate the development of emotional memory systems, ending the period of infantile amnesia prematurely[18,19]. Early adversity is a major risk factor for developing mental health disorders, suggesting that infantile amnesia may actually protect against the emergence of problematic behaviors[7,20]. Despite their importance for brain and behavioral development, how infant memories are stored and why they remain inaccessible remains poorly understood.

In adults, initial memory encoding in the hippocampus is followed by a period of consolidation

during which episodic memories come to be stored in distributed networks across the brain[21].

Memory consolidation and retrieval involves coordinated activity of numerous regions including

the prefrontal cortex, temporal cortical association areas, medial and anterior thalamic nuclei,

and retrosplenial cortex (RSP)[21–23]. Long range connectivity amongst these regions continually

refines throughout postnatal development until early adulthood[24–26]. It is therefore possible that

developmental remodeling of the long-term memory networks render infant memories

inaccessible. But little is known about the organization of memory networks in the developing

brain and how they change with age. To date, mechanistic studies of infantile amnesia have

focused on the hippocampus, showing that infantile amnesia is associated with naturally

elevated hippocampal neurogenesis[9] and immature hippocampal synapses[13]. As such, we have

only limited understanding of which aspects of brain circuit development are associated with the

emergence of persistent memories.

Here we set out to investigate why infant memories become inaccessible. We considered two

potential models. In the first, the infant brain may use different circuits to store and retrieve

memories that are no longer used in adulthood. In the second, the same circuits are involved

but memories are stored in a different way. We designed a developmental screen to distinguish

between these possibilities. By performing contextual fear conditioning, whole-brain activity-

dependent neuronal tagging[27–29] at different postnatal ages, tissue clearing, light sheet

fluorescence microscopy, and high throughput computational analyses[30], we identified key brain

regions whose activity and functional connectivity change significantly during memory retrieval

across development. Our analyses indicated that while similar regions are involved in memory

retrieval over development, functional connectivity amongst those regions changes significantly

between infancy and adulthood. RSP emerged as a key region whose functional maturation

closely aligns with changes in memory persistence. Then, in targeted functional studies, we

discovered that 1) that memory ensembles in RSP are more stable in adults compared to

juveniles and infants, 2) that chronically reactivating RSP memory for a week after learning

ensembles enhances memory in adults but not in infants, and 3) that after 33 days, reactivation

of infant-tagged RSP ensembles can recover infant memories.

Results

A brain-wide screen to identify neural signatures of infantile amnesia

Using an established protocol for modeling infantile amnesia[8,9,11], we performed contextual fear

conditioning (CFC) in mice when they were infants (postnatal day (P) 17), juveniles (P25), or

adults (P60). In separate cohorts of mice, we tested the strength of their fear memory 1 day (d)

or 7d after CFC by placing mice in the original training context and measuring freezing behavior

(Figure 1A). While mice of all ages exhibited robust freezing during CFC and the 1d memory

retrieval test, mice trained as infants froze significantly less than adults and juveniles during a

7d memory retrieval test (Figure 1B).

In our developmental screen, we combined activity-dependent tagging, tissue clearing and light

sheet fluorescence microscopy to screen for developmental changes in brain-wide cellular

activation patterns during memory retrieval (Figure 1C). We labeled activated neuronal

populations using Targeted Recombination in Active Populations (TRAP2: $Fos^{icre-2A-ERT2}$)[28,29], in

which the Fos promoter drives expression of tamoxifen-inducible Cre recombinase. We crossed

TRAP2 to the Ai14 Cre reporter line[31] so all TRAPed neurons could be labeled with a red

fluorophore. We performed CFC in infant, juvenile or adult *TRAP2;Ai14* mice. One day later, we

TRAPed fear memory-activated neurons by pairing re-exposure to the training context with a

tamoxifen injection. We chose to TRAP 1d instead of 7d memory retrieval because infant mice

still freeze 1d after CFC. This allowed us to relate neuronal activation patterns to freezing

behavior while comparing the neural signatures underlying retrieval of infant memories that

would fade vs. juvenile and adult memories that would persist. A week after TRAPing we

perfused mice, immunostained for tdTomato and rendered brains transparent using the Adipo-

Clear protocol[32]. We then imaged TRAPed cells in intact brain hemispheres using light sheet

fluorescence microscopy, automatically quantified the number of TRAPed cells per brain region,

and compared the resulting patterns of neuronal activation across age groups (Figure 1C).

First, we investigated whether brain regions were differentially activated during memory retrieval

over development (P17–P60). To do this, we calculated the number of TRAPed cells per brain

region as a fraction of the total TRAPed cells counted per brain. Cortical association areas,

regions in the hippocampal formation (HPF), and dorsal polymodal thalamic nuclei (DORpm)

had high fractional counts in adults compared to infants and juveniles whereas nuclei in the

hypothalamus and midbrain had higher fractional counts in infants compared to juveniles and

adults (Figure 1D,E). Cortical association areas, including the anterior cingulate area (ACA) and

RSP, HPF, and DORpm play key roles in memory consolidation[21,33–37]. These findings suggest

that maturation of regions involved in long-term memory in adults contributes to developmental

transitions in memory persistence.

Developmental changes in short-term fear memory network organization

To better understand which brain regions contribute to the behavioral expression of fear

memories across development, we conducted a partial least-squares (PLS) analysis[38] of

TRAPed cells across brain regions. These analyses identify latent variables that maximally

differentiate high versus low freezing levels during 1d memory retrieval. Bootstrap ratios (or

saliences) from the PLS analysis were used to determine the extent that individual regional

activation (based on numbers of TRAPed cells) contributed to the ability to distinguish freezing

levels among mice within each age group. We identified regions contributing to significant

contrasts by thresholding salience values at 3, which corresponds to P=0.01 (Figure 1F–H). In

all age groups, distinctions between freezing levels were primarily driven by TRAPed ensembles

in ACA, RSP, lateral septum (LSX), entorhinal cortex (ENT), and the lateral zone of the

hypothalamus (LH). This suggests that over development, similar regions are involved in the

behavioral expression of conditioned fear during 1d fear memory retrieval.

Memory retrieval and consolidation involves coordinated activity across brain regions.

Interregional functional connectivity strength is often estimated by assessing co-variance in

activity across brain regions[38,39]. In many brain regions, long-range connectivity continually

refines until early adulthood[24–26]. We therefore hypothesized that these structural changes would

produce distinct patterns of functional connectivity underlying 1d fear memory retrieval in

different stages of development. To examine this, we computed interregional correlations based

on the number of TRAPed cells and compared across the age groups. Based on our findings

that cortical association areas and DORpm are more highly activated during memory retrieval in

adults, we focused our covariance analysis on these and connected regions.

This analysis revealed an overall decrease of interregional correlation strength in juvenile mice

compared to infants and adults as well as age-dependent switches in correlation values among

several regions (Figure 2A). A Fisher r-to-z transformation test revealed several statistically

significant changes in interregional correlations across development (Figure S1), especially

between RSP, parahippocampal regions (PHR) and DORpm. Correlated activity between some

sets of regions changed between infant and juvenile stages and then stabilized. In other sets of

regions, correlated activity changed continuously from infant to adult stages, suggesting that

their circuits develop along a more protracted timeline.

We summarized some of the most notable changes by calculating the mean Pearson correlation

coefficient (r) for several combinations of regions (Figure 2B). In RSP, correlations with the

hippocampus and striatum were strongly negative in infants but increased in the juvenile stage

and then leveled off. RSP correlations with BLA parahippocampal regions were positive in

infants and adults but exhibited a transient decrease in juveniles. On the other hand,

correlations between RSP and DORpm decreased steadily from infancy to adulthood. We also

examined changes in correlations between DORpm regions and several cortical areas (PL, ILA,

ACA and RSP). In lateral (LAT), medial (MED), intralaminar (ILM) and midline (MTN) thalamic

nuclei, correlations with cortical areas switched from positive to negative from infancy to

adulthood. In anterior thalamic nuclei (ATN), correlations remained positive but decreased from

infant to juvenile stages and then plateaued (Figure 2B).

We next generated network graphs in which the nodes and connections represent brain regions

and correlations that survived thresholding using p<0.1 as a cutoff. In the network plots, node

size is proportional to degree, or the number of connections for that node (Figure 2C). Our

network analysis revealed that fear memory networks became sparser with age. In infants, most

nodes had high degrees of connectivity with all other nodes (66 edges). In juveniles, the

network was sparser and particular sets of regions were interconnected. For instance, subplate

areas including the BLA were interconnected with each other and with the hippocampal

formation (8/40 edges, or 20% of total connections within the network). On the other hand,

cortical areas were mostly connected with other cortical areas and not with the rest of the

regions. In adults, connectivity between cortical areas, hippocampal formation and BLA were

most prominent (29/50 edges, or 58% of the total connections in the network).

We calculated several measures of network tendencies to quantify changes in the organization

of fear memory networks across development. Network density, the ratio of the number of

actual links to possible links, decreased steadily from infant to adult stages. Analysis of network

heterogeneity revealed that networks become more clustered with age (Figure 2C). We also

calculated degree centrality, or the number of edges per node. Nodes in the infant network had

significantly more connections than in juvenile networks, but adults trended back slightly higher

(Figure 2D). We next computed the average shortest path length, or the number of steps along

the shortest paths connecting all pairs of nodes, for each age group. Nodes in adult networks

had significantly higher average shortest path lengths compared to both infants and juveniles

(Figure 2D). These results revealed significant developmental reorganization of memory

networks, with a shift from dense connectivity in infants to sparser more structured networks in

adults.

Overall, our developmental screen provided convergent evidence that a network of regions

known to play a key role in memory consolidation and long-term storage in adults –including

medial prefrontal cortex (mPFC), RSP, DORpm and PHR– is also involved in memory retrieval

during postnatal development. However, functional connectivity between these regions

reorganizes significantly between infancy and adulthood. Notably, RSP was identified in all

analyses in our screen (Figure 2E). RSP subregions were less activated in the infant brain

(Figure 1E), RSP had high salience values in our PLS analysis (Figure 1F-H), RSP was a key

node in memory networks at all ages (Figure 2C), and RSP underwent some of the most

pronounced changes in functional connectivity over development (Figure 2A,B; Figure S1).

While RSP has established roles in memory consolidation and long-term memory retrieval in

adults[34,40–43], its role in memory is poorly understood in the developing brain. Based on the

convergent findings in our screen, we chose to further examine the role of RSP in memory

retrieval and consolidation across development.

Reactivation of TRAPed RSP neurons during memory retrieval increases with age

Memory persistence has been ascribed to the reinstatement of patterns of neuronal ensemble

activity that were established during initial learning[44,45]. In hippocampus, mPFC and RSP,

neuronal ensembles that express Fos during learning or recent memory retrieval overlap with

Fos+ ensembles evoked during later recall[29,46]. The extent of ensemble reactivation predicts

retrieval success[5,26]. While RSP is known to play key roles in memory consolidation and remote

memory retrieval[41,47], how ensemble stability in RSP may change across development remains

unknown.

To determine to what extent ensembles of RSP neurons are reactivated during memory retrieval

across time, we used a combination of TRAP2 and immunostaining for Fos, a marker of recently

activated neurons (Figure 3A). Infant, juvenile or adult mice were TRAPed during 1d retrieval,

permanently labeling activated neurons with tdTomato. Then, following a 7d memory retrieval

test in the same mice, brains were collected for Fos immunostaining (Figure 3B–D). In RSP, we

observed no differences in the numbers of TRAPed or Fos+ cells (Figure 3E$_{1,2}$). However,

compared to infants and juveniles, adult mice had significantly higher levels of reactivated

neurons in RSP (Figure 3E$_3$). To explore the specificity of these findings, we examined

ensemble overlap in ACA, a neighboring cortical area that also has an established role in

memory consolidation and long-term storage[21,22]. In ACA, all age groups had similar numbers of

TRAPed, Fos+ and reactivated cells (Figure 3F$_{1-3}$). These results suggest that in the adult brain

RSP memory ensembles are more likely to be reactivated by salient contextual cues.

In the cortex, neurons residing in different layers (L) have distinct projection targets. L2/3

neurons are primarily intratelencephalic, projecting to other cortical areas, striatum and

subplate. L5 and L6 contain a mixture of intratelencephalic neurons, pyramidal tract neurons

that project to thalamus, midbrain, and spinal cord and corticothalamic neurons[48]. To gain

insight into whether distinct classes of projection neurons may be activated during memory

retrieval in different developmental stages, we calculated the number of TRAPed, Fos+ and

double-positive (reactivated) neurons in different cortical layers (Figure S2). In RSP, mice of all

ages had the most TRAPed and Fos+ neurons in L2/3 and the fewest in L6 (Figure S2A$_{1,2}$). We

observed significantly more reactivated cells in L2/3 for adults compared to juveniles and a

trend-level effect for adults vs. infants (Figure S2A$_3$). In ACA, juveniles had significantly more

TRAPed and Fos+ cells in L2/3 (Figure S2B$_{1,2}$) but we did not observe any layer- or age-

dependent differences in the distribution of reactivated neurons in ACA (Figure S2B$_3$). These

findings suggest that in RSP, L2/3 neurons play a prominent role in memory persistence and

that their likelihood of being reactivated by contextual cues increases with age.

Reactivation of TRAPed RSP ensembles shortly after learning enhances memory in an

age-dependent manner

Off-line activity within memory networks is thought to underlie memory consolidation[21]. A recent

study revealed that reactivation of tagged RSP memory ensembles can accelerate memory

consolidation[40]. We therefore hypothesized that offline reactivation of RSP memory ensembles

could recover lost infant memories. To test this, we injected into RSP an AAV expressing the

excitatory cre-dependent DREADD (designer receptor exclusively activated by designer drugs)

hM3Dq[49] or a control AAV that encoded a cre-dependent fluorophore. Two weeks later – when

mice were infants, juveniles, or adults – we performed contextual fear conditioning. The

following day, we TRAPed 1d memory-activated RSP neurons, thereby driving expression of

hM3Dq in those cells. We then performed off-line reactivation of the TRAPed ensembles by

administering daily the hM3Dq ligand clozapine-N-oxide (CNO, 1 mg/kg/day, intraperitoneally).

We administered a final dose of CNO (2 mg/kg) 30 min prior to a test retrieval session on day 7

(Figure 4A,B). Histology confirmed that hM3Dq-mCherry was expressed in a similar location

within RSP and to a similar extent across age groups (Figure S3) and that hM3Dq expressing

mice had significantly higher levels of activity in RSP (as assayed with Fos immunostaining)

compared to controls (Figure S4).

Chronic chemogenetic activation of TRAPed RSP ensembles shortly after learning affected fear

memory retrieval in an age-dependent manner. In infant mice, both control and hM3Dq groups

froze minimally during 7d retrieval, indicating that enhancing activity in RSP memory ensembles

in infant mice cannot prevent infantile amnesia (Figure 4C). In juveniles, control mice froze

slightly but significantly less during 7d vs. 1d retrieval, suggesting some fear extinction may

have occurred. On the other hand, juvenile hM3Dq-expressing mice froze similarly during the 1d

and 7d retrieval tests and significantly more than controls during 7d retrieval (Figure 4D). This

suggests that in juveniles, chronic reactivation of TRAPed RSP ensembles mildly enhances

memory. In line with previous findings[40], RSP ensemble reactivation in adult mice significantly

increased conditioned freezing during 7d retrieval compared to both the 1d retrieval session and

to 7d retrieval in control mice (Figure 4E). To further compare age-specific behaviors, we

computed for each hM3Dq-expressing mouse the change in freezing levels from 1d to 7d

retrieval (Figure 4F). All infants froze less, juveniles had no net change in freezing, and adults

froze more by the 7d test. Together, these data suggest that mature functions of RSP emerge

between juvenile and adult stages, and that memories are consolidated or retrieved via a

distinct mechanism in the infant brain.

To investigate the circuit-level differences underlying the distinct, age-specific effects of

reactivating RSP ensembles, we sacrificed the mice 60 min after the 7d retrieval test for Fos

immunostaining. We sectioned the brains, aligned individual slices to the Allen Brain Atlas and

then quantified the number of Fos+ neurons across a variety of brain regions. We focused on

brain regions that receive direct anatomical projects from RSP or have known functional

connectivity with RSP[25,35,41,50–53]. Compared to adults, infants had significantly fewer Fos+

neurons in the dorsal caudate putamen (CPd), several anterior thalamic nuclei (AM, AV, LD and

LP), and the anterior pretectal nucleus (APN), a midbrain motor nucleus (Figure 4G,H),

suggesting that immature functional connectivity between RSP and these regions may lead to

retrieval failure in infants.

Reactivation of infant-TRAPed RSP ensembles weeks after learning can recover lost memories

We next investigated whether infant-TRAPed RSP ensembles could promote memory retrieval after development had progressed. To test this, we injected into RSP an AAV encoding cre-dependent hM3Dq or a fluorophore control. Two weeks later, when mice were P17, we TRAPed RSP ensembles during contextual fear conditioning. 33d later, we delivered CNO before a memory retrieval test (Figure 5A,B). As before, histology confirmed that hM3Dq-mCherry was targeted to RSP and that hM3Dq expressing mice had significantly higher levels of activity in RSP (as assayed with Fos immunostaining) compared to controls (Figure S5).

In this experiment, hM3Dq-expressing mice froze significantly more than controls. We observed this effect in the conditioning context, but not in an altered context that had a distinct floor, wall decorations and scent (Figure 5C). Together these results reveal that infant RSP ensembles store latent memories and that once RSP cells and circuits develop, reactivating those ensembles can recover the stored infant memory. However, because activating TRAPed RSC cells only drove freezing in the conditioning context, our findings also indicate that the function of infant-TRAPed RSP ensembles are gated by the presence of salient contextual cues. Reactivating infant-TRAPed neurons in ACA or LSX did not enhance freezing 33d later (Figure S6), suggesting the effects we observed are specific to RSP.

To gain insight into which circuits may allow for recovery of lost infant memories, we then performed a week of chronic reactivation as described above in order to compare Fos downstream of infant-TRAPed RSP neurons to Fos downstream of adult-TRAPed RSP ensembles (Figure 4). We performed offline reactivation of TRAPed RSC neurons, performed another memory test on day 40 (Figure 5D), and then stained for Fos so we could compare to the findings described in Figure 4. We found that after allowing development to progress for 33d, reactivation of infant-TRAPed RSP ensembles largely recapitulated the Fos activation patterns we observed in adults (Figure 5E). There were two exceptions in the regions analyzed – AD and AM, suggesting that enhancement of activity in other regions (CPd, APN, LD, LP) may be responsible for the recovery of conditioned freezing.

Discussion

Even though very early experiences cannot be explicitly recalled, they can profoundly impact

the rest of our life[5,7,54]. Recent work shed light on this paradox, showing that infant memories

are not erased but rather inaccessible[11,13,15–17]. But why infant memories become inaccessible

was poorly understood. We hypothesized that one of two potential models may be at play. In the

first, distinct circuits store memories early in development and are no longer used in adulthood.

In the second, the same circuits are involved but function in different ways over development.

To investigate this, we designed a brain-wide screen to identify developmental changes in

memory retrieval circuits. Evidence from our developmental screen supported the second

model. We found that while similar brain regions encode memory throughout development,

changes in their behavioral functions and network organization align with the emergence of

persistent memory.

Our focused analyses revealed that RSP functional maturation aligns with the emergence of

persistent memories. RSP had more activated cells during memory retrieval in adults compared

to juveniles and infants. RSP functional connectivity with association cortices, hippocampus,

and amygdala - all key regions for memory encoding, consolidation and retrieval - changed

significantly across development. We defined the time course of RSP functional maturation,

showing that RSP memory ensembles were more likely to be reactivated by salient contextual

cues in adults compared to juveniles and infants (Figure 6). Along the same lines, reactivating

TRAPed RSP ensembles at 7 days enhanced memory in adults but not in infant mice and had

an intermediate effect in juveniles. But when we TRAPed RSP ensembles in infancy and then

allowed development to progress, reactivating TRAPed RSP ensembles after 30d successfully

recovered lost infant memories. This effect was only present in the conditioning context and not

in a modified context with distinct sensory cues. These findings indicate that in infancy,

memories are consolidated or retrieved through distinct mechanisms compared to adults due to

an immature RSP. Furthermore, our data indicate that while learning-activated RSP ensembles

encode latent infant memories over time, salient sensory cues do not activate those original

cells, so retrieval does not occur.

Our work extends models of infantile amnesia to include a key role for the retrosplenial cortex.

Current models center on hippocampal maturation. In infants, learning promotes hippocampal

synaptic maturation via signaling pathways that are involved in critical periods in other brain

regions, suggesting that the infant hippocampus is in a critical period for learning systems.[13]

These studies led the authors to propose the retrieval hypothesis, in which an immature

hippocampus stores information in a latent form such that it fails to be retrieved without

extensive reminders. In this model, learning promotes hippocampal maturation and drives

transitions through sequential critical periods. In each critical period, the hippocampus becomes

better at storing information such that it can be actively retrieved later[14]. Other studies showed

that in the infant brain, naturally high levels of hippocampal neurogenesis promote forgetting but

infant memories are stored in a dormant state in hippocampal ensembles[9,11]. This led the

authors to propose a (non-mutually exclusive) neurogenic hypothesis of infantile amnesia, in

which neurogenesis-induced circuit remodeling underlies infantile amnesia[5,55].

Our findings build upon this work, suggesting that the immature RSP contributes to the retrieval

failure of infant memories. Our data suggest that in infancy, an immature RSP can store

memories but cannot retrieve them due to immature functional connectivity with

parahippocampal and thalamic regions. Previous studies showed that offline reactivation of

tagged RSP ensembles accelerates memory consolidation in adults[40]. In our developmental

studies of RSP functions, we activated TRAPed neurons offline and immediately before memory

retrieval. So, we cannot distinguish the role of RSP in memory consolidation vs. retrieval. Still,

our results suggest that while activating TRAPed RSP enhances memory in adults, these

functions are absent in infants and only begin to emerge in juveniles. We also found that infant-

tagged RSP ensembles are less likely to be reactivated by salient contextual cues as compared

to adults. Furthermore, chemogenetically reactivating infant-TRAPed RSP neurons 33d later

enhanced freezing, but only in the conditioning context. This suggests that neither conditioned

sensory cues nor chemogenetic activation of RSP ensembles are alone sufficient to evoke

freezing behavior, but that they can do so in combination. In infants, immature RSP functional

connectivity and weak activation of RSP ensembles by conditioned stimuli may contribute to the

inability to retrieve infant memories. During development, these processes may be linked or

may be independently at play.

It remains to be determined how an immature hippocampus may influence the ability of the

infant RSP to encode, consolidate and retrieve memories. RSP serves as an input structure to

the hippocampus, providing important spatial cues. RSP also serves as a target of hippocampal

output during systems consolidation[56]. In adults, transfer of information from the hippocampus to

RSP is critical for the formation of long-term memories in the RSP[57–59] and for subsequent

memory retrieval[56,60,61]. In infants, the immature hippocampus could lead to weaker memory

representations in RSP as compared to adults. But we show that directly stimulating infant-

TRAPed RSP ensembles can recover the infant memory, suggesting that some post-learning

hippocampal-RSP information transfer has taken place. Further, our analyses indicate that in

infants, weaker functional connectivity between RSP and anterior thalamic nuclei and midbrain

motor areas may underlie retrieval failure. Future studies can examine how RSP connections

with the hippocampus, anterior thalamic nuclei and midbrain influence developmental transitions

in memory persistence.

The neural mechanisms of memory retrieval are well studied in adults but remain poorly

understood in the developing brain. Our studies provide novel insights into the organization of

fear memory networks in the developing brain and point to RSP as a key player in the

maturation of memory. Our whole brain analyses revealed that during recent memory retrieval

(1d after learning), similar sets of regions underlie conditioned freezing (Figure 2). These

included RSP, ACA, motor and sensory cortices, lateral hypothalamus, and midbrain. RSP was

a key node in recent (1d) fear memory networks at each age examined, but these networks had

distinct organization (Figure 3). Juvenile networks were sparser than in infants and adults. In

infants, most network nodes were interconnected, while adult networks had more structure, with

notable connections between RSP, frontal association areas, parahippocampal regions and

BLA. Even though juveniles and adults had similar freezing levels across time, distinct circuit

architectures underlie fear memory retrieval in these two age groups. This suggests that juvenile

and adult fear memories likely have unique mechanisms of consolidation, storage and retrieval.

In adults, in both rodents and humans, RSP plays a key role in learning[42,62,63], consolidation[64,65]

and recent and remote memory retrieval[61,66–68], but how and when RSP functionally matures

was largely unknown. We showed that infant RSP ensembles active during fear conditioning

encoded long-term memories but were not reactivated by contextual cues. We also found that

offline reactivation of RSP ensembles can promote memory retrieval beginning in juvenile

stages and these functions are further enhanced in adults. Together with the whole brain

analyses of TRAPed cells, these studies indicate that RSP is involved in learning and recent

memory retrieval beginning in infancy and plays an increasing role in memory consolidation and

long-term memory retrieval between juvenile and adult stages.

Our studies provide novel insights into the RSP circuit mechanisms underlying memory retrieval

and consolidation in adults and align maturation of RSP functional connectivity with its changing

role in memory. RSP is densely interconnected with sensory cortices, with limbic regions that

are important for memory encoding, and with anterior thalamic nuclei and cortical regions

important for long-term memory storage and retrieval[21,57,69–79]. In developing rats, bidirectional

connections linking RSP to the hippocampal formation reach adult-like densities by P12[25,26]. In

humans, resting state functional connectivity between RSP and the prefrontal cortex increases

between adolescence and early adulthood while RSP-hippocampal functional connectivity

matures earlier[80]. But little is known about how and when RSP circuits underlying memory

retrieval functionally mature.

We show that activating RSP in the conditioning context led to age-specific patterns of Fos

induction in downstream regions. Infants had fewer Fos+ cells in anterior thalamic nuclei, dorsal

striatum and midbrain motor nuclei. Because we did not observe differences in the layer

distribution of TRAPed cells (Figure S2), it is unlikely that the distinct patterns of Fos arose from

TRAPing different classes of projection neurons at each age. Instead, immature connectivity

with striatum, anterior thalamic nuclei and midbrain areas may influence the ability to

consolidate or retrieve memories, leading to retrieval failure in infants. Reactivating infant

TRAPed RSP neurons one month after training was sufficient to induce freezing behavior, but

only in the conditioning context, indicating that RSP ensemble function is gated by salient

sensory cues. Memory retrieval is thought to involve brain-wide reinstantiation of patterns of

activity established during learning, but in the cortex, synaptic connections remodel until early

adulthood. Consistent with previous work that suggests hippocampal circuit remodeling

promotes forgetting[55], our studies indicate that during development, synaptic remodeling within

infant RSP memory-encoding ensembles may diminish the likelihood of instantiating original

patterns of ensemble activity. This could break the links between salient sensory stimuli and

conditioned behaviors, leading to retrieval failure.

Early life adversity like chronic stress is a major risk factor for developing mental health

disorders. Early adversity can lead to premature emergence of persistent memories, suggesting

that accelerated development of memory systems may be linked to poor mental health

outcomes[7,18–20]. In aging models of neurodegeneration, in which subjects display a similar level

of forgetting as in infantile amnesia, memory impairment corresponds to the degeneration of

memory circuits that include the RSP[81,82]. By illustrating how memory retrieval networks change

across development, our work builds a foundation for understanding how adverse rearing

conditions may influence neurodevelopmental trajectories, leading to problematic behaviors in

adulthood. Further, revealing the organization and function of memory networks in animals with

infantile amnesia may provide mechanistic insight into memory disorders associated with

forgetting[83].

Acknowledgments

We thank Carlos Portera-Cailliau and Scott Wilke for feedback on the manuscript. This

publication was supported by a NARSAD Young Investigator Award from BBRF (L.A.D.),

Whitehall Foundation Grant (L.A.D.), Klingenstein-Simons Fellowship in Neuroscience (L.A.D.),

R01MH127214 (L.A.D), T32 MH073526 (B.J.), Jessamine Hilliard Neurobiology Graduate

Student Grant Program (B.J) and UCLA Dissertation Year Fellowship (B.J.). Confocal imaging

was performed in the UCLA Broad Center for Regenerative Medicine and Stem Cell Research

microscopy core. Figure 6 was produced using BioRender.com.

Author Contributions

450   Conception, B.J. and L.A.D.; Methodology, B.J., M.W.G., L.O., L.H.W., B.W., A.D., Y.N.

451   S.A.R.M., and L.A.D.; Investigation, B.J., ; Writing – Original Draft, B.J. and L.A.D.; Writing –

452   Review & Editing, B.J., M.W.G., L.A.D.; Funding Acquisition, B.J. and L.A.D.

Declaration of Interests

454   The authors declare no competing financial interests.

Figure Titles & Legends

Figure 1. A developmental screen to identify neural signatures of infantile amnesia.

(A) Infant, juvenile, and adult mice underwent contextual fear conditioning and then memory

testing 1 or 7 days later.

(B) Percent time freezing in infant, juvenile and adult wildtype mice during memory retrieval

tests occurring 1 or 7 days after CFC ($F_{age}(2,45)=15.89$, P<0.0001, $F_{day}(1,45)=3.56$, P=0.07,

$F_{int}(2,45)=4.21$, P=0.02, Infant: N=7 1d, N=10 7d; Juvenile: N=14 1d, N=8 7d; Adult: N=7 1d,

N=6 7d, two-way ANOVA with Tukey post-hoc test).

(C_1) Experimental design for developmental screen.

(C_2) Freezing behavior in TRAP mice used in screen (F(2,22)=1.87, P=0.18, One way ANOVA;

N=7 infant, N=9 juvenile, N=9 adult).

(D) Representative 100-µm thick optical z-stacks showing TRAPed cells from 1-day retrieval.

(E) TRAPed cell counts by age (Sensory cortex: $F_{region}(2.03,44.7)=267.30$, $P<0.0001$;

$F_{age}(2,22)=0.02$, $P=0.97$; $F_{int}(10,110)=0.65$, $P=0.77$; Association cortex:

$F_{region}(3.45,75.93)=132.7$, $P<0.0001$; $F_{age}(2,22)=3.82$, $P=0.38$; $F_{int}(24,264)=2.14$, $P=0.002$;

Sensory Thalamus: $F_{region}(1.18,26.04)=80.39$, $P<0.0001$; $F_{age}(2,22)=0.69$, $P=0.51$;

$F_{int}(10,110)=1.05$, $P=0.41$; Polymodal thalamus: $F_{region}(2.22,48.84)=28.77$, $P<0.0001$;

$F_{age}(2,22)=7.60$, $P=0.003$; $F_{int}(12,132)=1.59$, $P=0.10$; Hypothalamus/Midbrain:

$F_{region}(3.38,74.38)=141.0$, $P<0.0001$; $F_{age}(2,22)=3.10$, $P=0.07$; $F_{int}(10,110)=2.61$, $P=0.007$;

Subplate: $F_{region}(2.96,65.07)=149.7$, $P<0.0001$; $F_{age}(2,22)=1.93$, $P=0.17$; $F_{int}(12,132)=1.42$,

$P=0.17$; Hippocampal Formation: $F_{region}(1.94,42.58)=43.78$, $P<0.0001$; $F_{age}(2,22)=2.16$, $P=0.14$;

$F_{int}(24,264)=0.82$, $P=0.72$. Infant N=7, Juvenile N=9, Adult N=9; Two-way repeated measure

ANOVAs with Tukey post-hoc correction test).

(F-H) Task PLS analysis of TRAPed cell densities in (F) infants, (G) juveniles and (H) adults.

Dashed lines indicate threshold of 3. Color labels on x-axis correspond to major brain

subdivisions according to the Allen Brain Atlas.

See Tables S1 and S2 for abbreviations and statistics. Graphs show mean $\pm$ SEM. *$P< 0.05$,

**$P<0.01$, ***$P<0.001$. Scale bars, 100 μm.

Figure 2. Developmental changes in fear memory networks.

(A) Pearson correlation matrices showing interregional correlations based on numbers of

TRAPed cells at 1-day memory retrieval across ages. Colors reflect Pearson correlation

coefficients (scale, right) and labels within squares correspond to p values from pairwise

comparisons.

(B) Mean pearson r values for left: RSP ($F_{region}(4,48) = 12.47$, $P<0.0001$; $F_{age}(2,48) = 1.04$,

$P=0.37$; $F_{interaction}(8,48) = 5.60$, $P<0.0001$; HPF: N=3; STR: N=4; BLA: N=3; PHR: N=6; TH: N=5

brain regions) and right: between dorsal cortex (PL, ILA, ACA, RSP) and dorsal polymodal

thalamic nuclei (DORpm, $F_{region}(4, 45) = 5.82$, $P=0.0007$; $F_{age}(2, 45) = 26.48$, $P<0.0001$;

$F_{interaction}(8, 45) = 1.36$, $P=0.24$; N=4 cortical brain regions). Two-way ANOVAs with Tukey post-

hoc correction test.

(C) Network graphs generated for Pearson's correlations $p<0.1$ Regions are grouped and

colored according to major brain subdivision as defined by the Allen Brain Atlas. The node size

is proportional to the number of connections (degree) and the weights of the lines are

proportional to the correlation strength.

(D) Average degree (left, $F_{(2,75)}=5.21$, P=0.008, One-way ANOVA) and shortest path length

(right, $F_{(2,75)}=18.34$, P<0.0001, One-way ANOVA) for each network node for each age (Infant:

N=24; Juvenile: N=28; Adult: N=26 nodes).

(E) Summary of convergent findings from analyses presented in Figures 1 and 2.

*P< 0.05, **P<0.01, ***P<0.001, ****P<0.0001. Graphs show mean $\pm$ SEM. See also Figure S1

for additional network analyses.

Figure 3. Selectively increased reactivation of memory-encoding ensembles in RSP in

adults.

(A) Illustration of stable vs. dynamic ensembles in a TRAP/Fos overlay assay.

(B) Experiment design for activity-dependent tagging of ensembles between 1- and 7-day

retrieval.

(C) Illustration of analysis sites for RSP and ACA.

(D) Representative examples of TRAPed (magenta) and Fos+ cells (green) in the anterior

cingulate (top) and retrosplenial cortices (anterior, middle; posterior, bottom).

(E_1) TRAPed cells per ROI for ACA ($F_{(2,16)}=0.34$, P=0.72, one-way ANOVA with Tukey post-

hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(E_2) Fos+ cells per ROI for ACA each region and age group ($F_{(2,16)}=2.11$, P=0.15, one-way

ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(E_3) Double-labeled neurons per ROI for ACA ($F_{(2,16)}=0.39$, P=0.68, one-way ANOVA with

Tukey post-hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(F_1) TRAPed cells per ROI for RSP ($F_{(2,11)}=0.09$, P=0.92, one-way ANOVA with Tukey post-

hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

(F_2) Fos+ cells per ROI for RSP each region and age group ($F_{(2,11)}=0.70$, P=0.52, one-way

ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

(F_3) Double-labeled neurons per ROI for RSP ($F_{(2,11)}=7.1$, P=0.01, one-way ANOVA with

Tukey post-hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

Graphs show mean $\pm$ SEM. *P <0.05. See also Figure S2 for layer analysis.

Figure 4. Chemogenetic reactivation of TRAPed RSC neurons increased freezing during

short-term memory retrieval in an age-dependent manner.

(A) Experimental design.

(B) Representative coronal section showing AAV injection site in RSP. Scale bar, 125 μm.

(C) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (brown) groups for infants ($F_{virus}(1,7)=1.29$, P=0.29, $F_{day}=(1,7)=24.18$, P=0.002; $F_{int}(1,7)=0.47$, P=0.51, Cont: N=4, hM3Dq: N=5, two-way ANOVA with Holm-Sidak post hoc test).

(D) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (magenta) groups for juveniles ($F_{virus}(1,28)=1.81$, P=0.19, $F_{day}=(1,7)=3.33$, P=0.07; $F_{int}(1,7)=4.34$, P=0.04, Cont: N=16, hM3Dq: N=14, two-way ANOVA with Holm-Sidak post hoc test).

(E) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (teal) groups for adults ($F_{virus}(1,28)=1.81$, P=0.19, $F_{day}=(1,7)=3.33$, P=0.07; $F_{int}(1,7)=4.34$, P=0.04, Cont: N=8, hM3Dq: N=10, two-way ANOVA with Holm-Sidak post hoc test).

(F) Differences in freezing levels between 7-day and 1-day retrieval sessions for hM3Dq mice in each age group (F(2,26)=12.92, P=0.001, Infant: N=5, Juvenile: N=14, Adult: N=10, one-way ANOVA with Tukey post hoc test).

(G) Representative images showing Fos immunoreactive cells in dorsal caudate putamen (CPd), anteromedial thalamus (AM), laterodorsal thalamus (LD), and anterior pretectal nucleus (APN). Scale bars, 100 μm.

(H) Quantification of Fos+ cells per ROI in 19 brain regions ($F_{region}(18,160)=30.03$, P<0.05; $F_{age}(2,9)=3.69$, P>0.05; $F_{int}(36,160)=5.78$, P<0.05, Mixed Effects Model; Infant: N=4; Juvenile: N=4; Adult: N=3-4).

Graphs show mean $\pm$ SEM. * P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. See also Figure S3 and S4.

Figure 5. Chemogenetic reactivation of learning encoding ensembles in RSC 33d after training recovers infant memories

(A) Experimental design.

(B) Representative coronal section showing AAV injection site in RSP. Scale bar, 125 μm.

(C) Percent freezing levels in contexts A and B during the 33-day retrieval test in control (gray) and hM3Dq (brown) mice ($F_{virus}(1,14)=9.06$, P=0.009; $F_{context}(1,14)=7.41$, P=0.01; $F_{int}(1,14)=5.23$, P=0.03, Cont: N=8, hM3Dq: N=8, two-way ANOVA with Holm-Sidak posthoc test).

(D) Percent freezing levels in contexts A and B during the 40-day retrieval test in control (gray) and hM3Dq (brown) mice ($F_{virus}(1,14)=9.81$, P=0.007; $F_{context}(1,14)=12.3$, P=0.003; $F_{int}(1,14)=4.65$, P=0.07, Cont: N=8, hM3Dq: N=8, two-way ANOVA with Holm-Sidak posthoc test).

565   (E) Comparison of regional Fos levels in P17-TRAP+40d vs. Adult TRAP from Figure 4

566   (F_{region}(18,121)=18.97, P<0.0001; F_{age}(1,7)=3.05, P=0.12; F_{int}(18,121)=3.5, P<0.0001, Inf: N=5,

567   Adult: N=4, two-way mixed effects repeated measures analysis with Sidak posthoc test).

568   Graphs show mean $\pm$ SEM. *P<0.05. See also Figures S5 and S6.

STAR Methods

Resource availability

Further information and request for resources and reagents should be directed to and will be

fulfilled by the lead contact, Laura A. DeNardo (ldenardo@ucla.edu)

Materials availability

This study did not generate new unique reagents.

Data and code availability

Whole brain TRAP data are presented in Table 2 and previously published code for

DeepCOUNT can be found in the DeNardo lab GitHub database

(https://github.com/DeNardoLab). Any additional information required to reanalyze the data

reported in this paper is available from the lead contact upon request.

Experimental model and study participant details

Animals

Female and male C57B16/J mice (JAX Stock No. 000664) or TRAP2;Ai14 mice (JAX Stock

Nos. 030323 and 007914) were group housed (2–5 per cage) and kept on a 12 hour light cycle

(lights on 7am-7pm). All animal procedures followed animal care guidelines approved by the

University of California, Los Angeles Chancellor's Animal Research Committee.

Method Details

Surgery

Animals underwent surgery at P3, P9 or P46. Depending on the experiment, mice were

anesthetized through hypothermia (P3 mice), or induced in 5% isoflurane in oxygen until loss of

righting reflex (P9 or older) and transferred to a stereotaxic apparatus with rubber ear bars. P9+

mice were maintained under 2% isoflurane in oxygen, and warmed with a circulating water

heating pad throughout surgery and eye gel was applied to the animal's eyes. The mouse's

head was shaved and prepped with three scrubs of alternating betadine and then 70% ethanol.

Following a small skin incision, a dental drill was used to drill through the skulls above the

injection targets. A syringe pump (Kopf, 693A) with a Hamilton syringe was used for injections

delivered at a rate 100nL/min. The syringe was left in the brain for 1 or 7 minutes (min) following

injection for P3 and P9+ mice respectively. For DREADD studies, a small volume of AAV8-

hSyn-DIO-hM3D(Gq)-mCherry virus (Addgene 44361, final titer 2 x 10^{12} pp per mL) was

injected bilaterally into RSC (AP: -2.7, ML: $\pm$ 0.3, DV: 1.0). At P3 and P9, this coordinate was

scaled using Neurostar StereoDrive software (Neurostar) which corrects for head size,

orientation, and tilt. To compensate for age-dependent changes in cortical volume, we made

proportional adjustments to the volume of injected virus based on previous developmental

studies[84]: 0.15 µL for P3, 0.2 µL for P9, and 0.4 µL for P46 mice. Control mice received

injections of age-matched volumes of AAV1-pCAG-FLEX-eGFP-WPRE (Addgene 51502, final

titer 1 x 10^{13} pp per mL) into RSC. For pain management mice received 5 mg/kg carprofen

diluted in 0.9% saline subcutaneously. Mice received one carprofen injection 30 min prior to

surgery and daily injections for two days following surgery. Following confirmatory histology,

samples with mistargeted injection sites were excluded from analysis.

4-Hydroxytamoxifen Preparation

4-hydroxytamoxifen (4-OHT; Sigma, Cat# H6278) was dissolved at 20 mg/mL in ethanol by

shaking at 37°C for 15 min and was then aliquoted and stored at –20°C for up to several weeks.

Before use, 4-OHT was redissolved in ethanol by shaking at 37°C for 15 min, a 1:4 mixture of

castor oil:sunflower seed oil (Sigma, Cat #s 259853 and S5007) was added to give a final

concentration of 10 mg/mL 4-OHT, and the ethanol was evaporated by vacuum under

centrifugation. The final 10 mg/mL 4-OHT solutions were always used on the day they were

prepared. All injections were delivered intraperitoneally (i.p.).

Behavioral assays

Mice were placed in an operant chamber with a shock floor. They received 5 1 second, 0.5 mA

foot shocks separated by 1 min. On retrieval days, mice were placed in the conditioning

chamber for 5 min. For TRAP experiments, mice were injected with 4-OHT solution immediately

following the retrieval session. Point-tracking of behavioral videos was performed in

DeepLabCut[85]. Locomotion and freezing behavior was analyzed using BehaviorDEPOT[86].

Brain Clearing

Two weeks after TRAPing, mouse brains were collected and processed based on the published Adipo-Clear protocol[32] with slight modifications. Mice were perfused intracardially with 20mL of PBS (Gibco) followed by 4% paraformaldehyde (PFA, Electron Microscopy Sciences) on ice. Brains were hemisected approximately 1 mm past midline and postfixed overnight in 4% PFA at 4°C. The following day, samples were dehydrated with a gradient of methanol (MeOH, Fisher Scientific):B1n buffer (1:1,000 Triton X-100, 2% w/v glycine, 1:10,000 NaOH 10N, 0.02% sodium azide) for 1 hour for each step (20%, 40%, 60%, 80%) on a nutator (VWR). Samples were then washed with 100% MeOH 2x for 1 hour each and then incubated in a 2:1 dicholoromethane (DCM):MeOH solution overnight. The following day, two washes of 1 hour in 100% DCM were performed followed by three washes of 100% MeOH for 30 min, 45 min then 1 hour. Samples were bleached for 4 hours in 5:1 H2O2/MeOH buffer. A cascade of MeOH/B1n washes (80%, 60%, 40%, 20%) for 30 min each rehydrated the samples followed by a 1 hour wash in B1n buffer. 5%DMSO/0.3M Glycine/PTxWH permeabilized tissue for one hour and then again for 2 hour with fresh solution. Samples were washed with PTxwH for 30 min and then incubated in fresh PTxwH overnight. The following day two more PTxwH washes lasted 1 hour then 2 hours. For TRAP2 experiments, samples were incubated in primary RFP antibody (Rockland 600-401-379) at 1:300 in PTxwH shaking at 37°C for 11 days, washed in PTxwH 2x1 hour and then 2x2 hours, then for two days with at least one PTxwH change per day while shaken at 37°C. Samples were then incubated in secondary antibody AlexaFluor 647, catalog #A-31573, 1:300, ThermoFisher Scientific) for 8 days shaken at 37°C. Samples were washed in PTxwH 2x1 hour, then 2x2 hour, then 2 days with at least one PTxwH change per day while shaken at 37°C. Samples were then washed in 1x PBS twice 1x1 hour, 2x2 hours and then overnight. To dehydrate samples, a gradient of washes in MeOH:H2O (20%, 40%, 60% and 80%) were conducted for 30 min each, followed by 3x100% MeOH for 30 min, 1 hour, then 1.5 hours. Samples were incubated overnight in 2:1 DCM:MeOH on a nutator. The next day, samples were washed in 100% DCM 2x1 hour each. Samples were then cleared in 100% dibenzyl ether (DBE). DBE was changed after 4 hours. Samples were stored in DBE in a dark place at room temperature. Imaging took place at least 24 hours after clearing. Samples with poor antibody penetration following tissue clearing were excluded from analysis.

Whole Brain Imaging

Brain samples were imaged on a light-sheet microscope (Ultramicroscope II, LaVision Biotec)

equipped with a sCMOS camera (Andor Neo) and a 2x/0.5 NA objective lens (MVPLAPO 2x)

equipped with a 6 mm working distance dipping cap. Image stacks were acquired at 0.8x optical

zoom using Imspector Microscope v285 controller software. For TRAP2;Ai14 brains, we imaged

using 488-nm (laser power 20%) and 640-nm (laser power 50%) lasers. The samples were

scanned with a step-size of 3 µm using the continuous light-sheet scanning method with the

included contrast adaptive algorithm for the 640-nm channel (20 acquisitions per plane), and

without horizontal scanning for the 488-nm channel.

DeepCOUNT Analysis

Image Registration & Cell Segmentation

Whole-brain TRAP counts were obtained using DeepCOUNT as previously described[30]. Briefly,

whole-brain image stacks from the 640nm channel from each brain were segmented in

TrailMap[87] using the TRAP2-Ai14 trained model. Images were registered to the template brain

using elastix and transformix using the 488nm autofluorescence channel. MATLAB was used to

identify 3D maxima of the transformed probability map. Connected component analysis was

used to reduce any maxima that consisted of multiple pixels into a single pixel per cell.

Cell Quantification

Regional TRAPed cell density was quantified in MATLAB (Mathworks) by counting the number

of labeled pixels (i.e. cells) in each brain region, then dividing this pixel count by the total

number of pixels in a region. This regional pixel count was then divided by the total number of

detected cells across the brain to normalize for differences in tamoxifen-induced recombination.

Regions were defined by a collapsed version of the LSFM atlas in which maximum granularity

was balanced with the need to account for slight differences in registration which would lead to

inaccurate quantification of small brain regions. This atlas was cropped on the anterior and

posterior ends to match the amount of tissue visible in our data. Fiber tracts, ventricular

systems, cerebellum, and olfactory bulb were excluded from analysis. For comparison of

regional TRAPed counts between ages, Two-way repeated measures ANOVA with post-hoc

multiple comparison correction was performed to identify statistically significant differences

between brain regions. All statistical comparisons were performed in GraphPad Prism v10.

Functional Network Construction

In MATLAB, we calculated simple linear correlations between all pairs of brain regions based on

the normalized, per area TRAPed cell counts. The nodes and connections in the networks

represent brain regions and correlations that survived thresholding using $P<0.1$ as a cutoff.

While potentially interesting, we did not consider negative correlations in the current analysis.

We used Cytoscape software to visualize and analyze networks. Degree was calculated by

counting all the above-threshold connections for a given node. Node size is proportional to

degree and lines represent above-threshold correlations.

Brain Slice Histology and Immunostaining

Mice were transcardially perfused with phosphate-buffered saline (PBS) followed by 4%

paraformaldehyde (PFA) in PBS. Brains were dissected, post-fixed in 4% PFA for 12–24 hours

and placed in 30% sucrose for 24–48 hours. They were then embedded in Optimum Cutting

Temperature (OCT, Tissue Tek) and stored at -80°C until sectioning. 60 µm floating sections

were collected into PBS. Sections were washed 3x10 min in PBS and then blocked in 0.3%

PBST containing 10% normal donkey serum (catalog #17-000-121, Jackson Immunoresearch)

for 2 hours. Sections were then incubated in a carrier solution (3% donkey serum in 0.3%

PBST) containing the primary antibodies for two nights at 4°C. Two days later, sections were

washed 3x5 min in PBS and then stained in a carrier solution (5% donkey serum in 0.3% PBST)

with secondary antibody for 2 h at room temperature. The following primary antibodies were

used: polyclonal chicken anti-GFP (catalog #1020, 1:2000, Aves), monoclonal rabbit anti-RFP

(catalog #600-401-379, 1:2000, Rockland), monoclonal rabbit anti-Fos (catalog #226-008,

1:2000, Synaptic Systems) , and mCherry monoclonal anti-rat (catalog #M11217, 1:500,

ThermoFisher). The following secondaries were used: Cy2 polyclonal donkey anti-chicken

(catalog #703-225-155, 1:1000, Jackson Immunoresearch), Cy3 polyclonal donkey anti-rabbit

(catalog #711-005-152, 1:1000, Jackson Immunoresearch), Alexa647 polyclonal donkey anti-

rabbit (catalog #711-605-152, 1:1000, Jackson Immunoresearch), and Cy3 polyclonal donkey

anti-rat (catalog #712-165-153, 1:1000, Jackson Immunoresearch). Sections were then washed

5 min with PBS, 15 min with PBS+DAPI (catalog #D1306, 1:4000, Thermofisher), and then 5

min with PBS. Sections were mounted on glass slides using FluoroMount-G (catalog #00-4958-

02, ThermoFisher) and then imaged at 10x with a Zeiss 700 confocal microscope or at 10x on a

Leica DM6 B scanning microscope.

DREADD Studies

To activate neurons, we used a cre-dependent virus expressing the excitatory DREADD (pAAV-

hSyn-DIO-hM3D(Gq)-mCherry) or a fluorophore control (pAAV-hSyn-DIO-mCherry) in TRAP2

mice two weeks prior to fear-conditioning. Viral expression was driven exogenously by

intraperitoneally delivered 4-hydroxytamoxifen, which induced fos-dependent expression of cre-

recombinase in the TRAP2 mice, allowing specific tagging and long-term labeling of memory

encoding ensembles for weeks later. To determine the effects of chronic activation of TRAPed

RSC ensembles during development (i.e. comparing freezing levels between retrieval days 1

and 7), 4-hydroxytamoxifen was intraperitoneally injected immediately after 1-day retrieval to tag

ensembles involved in retrieval and consolidation (Figure 4) or immediately after contextual fear

conditioning (Figure 5).

30 min prior to their first retrieval session, mice were intraperitoneally injected with CNO

dissolved in 0.9% saline (1mg/kg, i.p., Fisher Scientific). After the first retrieval session, mice

were returned to their home cage. A subset of mice were also tested in an alternate context 3

hours after the first testing session, and then returned to their home cage. Following the first

retrieval test, mice were injected daily with CNO (2 mg/kg, i.p.) for 5 days and then returned to

the homecage. 30 min prior to the last retrieval day, CNO (1 mg/kg, i.p.) was injected prior to

testing in a similar manner as the first retrieval day. Mice were perfused 1 hour after the last

retrieval session and their brains collected and processed for Fos immunostaining.

Fos Analysis in 2-D sections

Alignment

Following Fos immunostaining and imaging as described above, sections were aligned to the

Allen Brain Atlas[88] using DeepSlice[89], a deep-learning based algorithm for automatically

calculating the position and slicing angle of each section in relation to the atlas. The DAPI

channel was used for alignment. Alignments were imported to the ABBA plugin for FIJI[90,91]

(https://biop.github.io/ijp-imagetoatlas/) which was used to manually check the alignment for

each section. ROIs from ABBA were then exported to FIJI, and ROIs for regions to be analyzed

were manually checked for tissue damage and image quality.

Fos+ Cell Detection

Images of the Fos channel were processed using a custom deep learning pipeline. This pipeline

has largely the same architecture as our existing 3D pipeline, DeepCOUNT[30], but uses a 2D U-

Net instead of a 3D U-Net. A network for segmenting Fos$^+$ cells was trained and validated using

manually annotated 100x100 pixel images. Following model training, each section was

segmented for Fos$^+$ cells. Cells were quantified using 2D maxima detection on the segmented

images in FIJI. The number of detected cells in each region was divided by the area of that

region to generate Fos density counts.

Quantification and Statistical Analysis

Results were plotted and tested for statistical significance with MATLAB, R and GraphPad

Prism v10. Statistical tests are described in the figure legends. Multiple comparisons with single

variables were analyzed with one-way ANOVA with post-hoc Tukey test (comparing the mean of

each column with that of every other column). For multiple comparisons with more than one

variable, two-way ANOVA with post hoc Tukey or Sidak test was used. The relationships

between TRAPed cell counts and freezing levels were analyzed using a partial least-squares

analysis implemented in R. Resampling statistics were used to determine the reliability of brain

region saliences. The reliability of brain region saliences was determined using bootstrap ratios

in which subjects retained their freezing levels but were resampled with replacement 1000

times. Original saliences were divided by the bootstrap-derived standard deviations to generate

bootstrap ratios (saliences). Salience above 3 was considered reliable. For the correlations

shown in Figure 2, all possible pairwise correlations between the TRAP signal in the 29 regions

were determined by computing Pearson correlation coefficients. Each complete set of

correlations was displayed as color-coded correlation matrices using MATLAB. The Fisher Z-

Transformation (to transform the sampling distribution of Pearson's R) was performed in

MATLAB. No statistical methods were used to predetermine sample size. Sample sizes were

determined based on similar published studies. In all figures, graphs show group means

superimposed on individual data points from each animal.

Supplemental Information Titles & Legends

Figure S1. Developmental changes in fear memory networks. Related to Figure 1.

Correlation matrices showing the differences in functional connectivity (Fisher Z-transformation)

between ages. Fisher correlation coefficients (scale, right; brown and cobalt representing

significant differences that are higher in older or younger age groups respectively) and labels

within squares correspond to p values of pairwise comparisons of Fisher r-to-z-transformation of

the Pearson's correlation coefficient matrices corresponding to respective age groups. *P< 0.05,

**P<0.01, ***P<0.001 by one-way ANOVA test.

Figure S2. Layer-specific distribution of TRAPed and Fos+ cells in ACA and RSP. Related

to Figure 3.

(A_1) TRAPed cell distributions across layers in RSP ($F_{layer}(2,33)=45.06$, $P<0.0001$;

$F_{age}(2,33)=0.00005$, $P=0.99$; $F_{int}(4,33)=0.96$, $P=0.43$; Infant: N=4; Juvenile: N=5, Adult: N=5).:

(A_2) Fos+ cell distribution across layers in RSP: $F_{layer}(2,33)=93.86$, $P<0.0001$; $F_{age}(2,33)=0.0003$,

$P=0.99$; $F_{int}(4,33)=0.23$, $P=0.91$; Infant: N=4; Juvenile: N=5, Adult: N=5).

(A_3) Reactivated cell distribution across layers in RSP ($F_{layer}(2,33)=8.55$, $P=0.001$;

$F_{age}(2,33)=7.97$, $P=0.002$; $F_{int}(4,33)=0.53$, $P=0.71$; Infant: N=4; Juvenile: N=5, Adult: N=5).

(B_1) TRAPed cell distributions across layers in ACA ($F_{layer}(2,48)=23.82$, $P<0.0001$;

$F_{age}(2,48)=0.0005$, $P=99$; $F_{int}(4,48)=4.25$, $P=0.005$; Infant: N=4; Juvenile: N=8, Adult: N=7).

(B_2) Fos+ cell distribution across layers in ACA ($F_{layer}(2,48)=20.99$, $P<0.0001$;

$F_{age}(2,48)=0.0012$, $P=0.99$; $F_{int}(4,48)=5.34$, $P=0.0012$; Infant: N=4; Juvenile: N=8, Adult: N=7).

(B_3) Reactivated cell distribution across layers in ACA ($F_{layer}(2,48)=0.51$, $P=0.60$;

$F_{age}(2,48)=1.11$, $P=0.33$; $F_{int}(4,48)=0.14$, $P=0.96$; Infant: N=4; Juvenile: N=8, Adult: N=7).

Figure S3. Mapping viral targeting in RSP. Related to Figure 4.

(A) Coronal sections showing location of RSP hM3D-mCherry expression in infant-TRAPed

mice

(B) Coronal sections showing location of RSP hM3D-mCherry expression in juvenile-TRAPed

mice

(C) Coronal sections showing location of RSP hM3D-mCherry expression in adult-TRAPed mice

Figure S4. RSP Fos levels in hM3D-expressing vs. control mice. Related to Figure 4.

(A_1) Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice.

(A_2). Quantification of Fos+ cell density in RSP for infant-TRAPed mice (Cont: N=4, hM3Dq:

N=5, P=0.003, two-tailed t-test).

(B_1). Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice.

in juvenile-TRAPed mice (Cont: N=6, hM3Dq: N=6, P=0.0001, two-tailed t-test).

(B_2). Quantification of Fos+ cell density in RSP for juvenile-TRAPed mice.

(C_1). Example images RSP showing AAV-mCherry and Fos+ RSP cells adult-TRAPed mice.

(C_2). Quantification of Fos+ cell density in RSP for adult -TRAPed mice (Cont: N=6, hM3Dq:

N=6, P<0.0001, two-tailed t-test).

Scale bars, 125 μm. Graphs show mean ± SEM.

Figure S5. Mapping viral targeting in RSP. Related to Figure 5.

(A) Coronal sections showing location of RSP hM3D-mCherry expression in infant-TRAPed

mice.

(B) Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice.

Scale bars, 125μm.

(C) Quantification of Fos+ cell density in RSP for infant-TRAPed mice (Cont: N=6, hM3Dq: N=6,

P<0.0001, two-tailed t-test). Graphs show mean ± SEM.

Figure S6. RSP DREADD validation and controls for Figure 5.

(A) Representative coronal section showing AAV-hM3Dq injection sites in LSX and ACA.

(B) Percent freezing levels in contexts A and B during the 33d retrieval test in LSX- and ACA-

targeted mice (LSX: P=0.54, N=3, ACA: P=0.56, N=3, paired t-test). Graphs show mean ± SEM.

References

1.   Bauer, P. J. & Larkina, M. Childhood amnesia in the making: Different distributions of

autobiographical memories in children and adults. *J. Exp. Psychol. Gen.* **143**, 597–611

(2014).

2.   Bauer, P. J., Tasdemir-Ozdes, A. & Larkina, M. Adults' reports of their earliest memories:

Consistency in events, ages, and narrative characteristics over time. *Conscious. Cogn.* **27**,

76–88 (2014).

3.   Campbell, B. A. & Campbell, E. H. Retention and extinction of learned fear in infant and

adult rats. *J. Comp. Physiol. Psychol.* **55**, 1–8 (1962).

4.   Lloyd, M. E. & Newcombe, N. S. Implicit memory in childhood: Reassessing developmental

invariance. in *The development of memory in infancy and childhood, 2nd ed.* 93–113

(Psychology Press, 2009).

5.   Ramsaran, A. I., Schlichting, M. L. & Frankland, P. W. The ontogeny of memory persistence

and specificity. *Dev. Cogn. Neurosci.* (2018) doi:10.1016/j.dcn.2018.09.002.

6.   Campbell, B. A. & Spear, N. E. *ONTOGENY OF MEMORY 1. Psychological Review* vol. 79

215–236 (1972).

7.   Callaghan, B. *et al.* Using a Developmental Ecology Framework to Align Fear Neurobiology

Across Species. *Annu. Rev. Clin. Psychol.* (2019) doi:10.1146/annurev-clinpsy-050718-

095727.

8. Akers, K. G., Arruda-Carvalho, M., Josselyn, S. A. & Frankland, P. W. Ontogeny of
contextual fear memory formation, specificity, and persistence in mice. *Learn. Mem.* **19**,
598–604 (2012).

9. Akers, K. G. *et al.* Hippocampal Neurogenesis Regulates Forgetting During Adulthood and
Infancy. *Science* **598**, 598–603 (2014).

10. Guskjolen, A., Josselyn, S. A. & Frankland, P. W. Age-dependent changes in spatial
memory retention and flexibility in mice. *Ontog. Learn. Mem.* **143**, 59–66 (2017).

11. Guskjolen, A. *et al.* Recovery of "Lost" Infant Memories in Mice. *Curr. Biol.* 1–8 (2018)
doi:10.1016/j.cub.2018.05.059.

12. Li, S., Kim, J. H. & Richardson, R. Updating memories: Changing the involvement of the
prelimbic cortex in the expression of an infant fear memory. *Neuroscience* **222**, 316–325
(2012).

13. Travaglia, A., Bisaz, R., Sweet, E. S., Blitzer, R. D. & Alberini, C. M. Infantile amnesia
reflects a developmental critical period for hippocampal learning. *Nat. Neurosci.* **19**, 1225–
1233 (2016).

14. Alberini, C. M. & Travaglia, A. Infantile Amnesia: A Critical Period of Learning to Learn and
Remember. *J. Neurosci.* **37**, 5783–5795 (2017).

15. Kim, J. H., Li, S., Hamlin, A. S., McNally, G. P. & Richardson, R. Phosphorylation of
mitogen-activated protein kinase in the medial prefrontal cortex and the amygdala following
memory retrieval or forgetting in developing rats. *Neurobiol. Learn. Mem.* **97**, 59–68 (2012).

16. Spear, N. & Parsons, P. Analysis of a reactivation treatment: Ontogenetic determinants of
alleviated forgetting. in *Processes of Animal Memory* vol. 1 135–165 (Lawrence Erlbaum
Associates, 1976).

17. Campbell, B. A. & Jaynes, J. Reinstatement. *Psychol. Rev.* **73**, 478–480 (1966).

18. Callaghan, B. L. & Richardson, R. Maternal Separation Results in Early Emergence of
Adult-Like Fear and Extinction Learning in Infant Rats. *Behav. Neurosci.* (2011)
doi:10.1037/a0022008.

19. Bath, K. G., Manzano-Nieves, G. & Goodwill, H. Early life stress accelerates behavioral and
neural maturation of the hippocampus in male mice. *Horm. Behav.* (2016)
doi:10.1016/j.yhbeh.2016.04.010.

20. Callaghan, B. L. & Tottenham, N. The Stress Acceleration Hypothesis: Effects of early-life
adversity on emotion circuits and behavior. *Curr. Opin. Behav. Sci.* **7**, 76–81 (2016).

21. Frankland, P. W. & Bontempi, B. The organization of recent and remote memories. *Nat.*
Rev. Neurosci. **6**, 119–30 (2005).

22. Frankland, P. W. & Bontempi, B. Fast track to the medial prefrontal cortex. *Pnas* **103**, 509–
510 (2006).

23. Tonegawa, S., Morrissey, M. D. & Kitamura, T. The role of engram cells in the systems
consolidation of memory. *Nat. Rev. Neurosci.* **19**, 485–498 (2018).

24. Klune, C. B., Jin, B. & DeNardo, L. A. Linking mPFC circuit maturation to the developmental
regulation of emotional memory and cognitive flexibility. *eLife* **10**, 1–33 (2021).

25. Sugar, J. & Witter, M. P. Postnatal development of retrosplenial projections to the
parahippocampal region of the rat. *eLife* **5**, e13925 (2016).

26. Haugland, K. G., Sugar, J. & Witter, M. P. Development and topographical organization of
projections from the hippocampus and parahippocampus to the retrosplenial cortex. *Eur. J.*
Neurosci. **50**, 1799–1819 (2019).

27. DeNardo, L. & Luo, L. Genetic strategies to access activated neurons. *Curr Opin Neurobiol*
45, 121–129 (2017).

28. Allen, W. E. *et al.* Thirst-associated preoptic neurons encode an aversive motivational drive.
Science **357**, 1149–1155 (2017).

29. DeNardo, L. A. *et al.* Temporal evolution of cortical ensembles promoting remote memory
retrieval. *Nat. Neurosci.* **22**, 460–469 (2019).

30. Michael W. Gongwer *et al.* Brain-wide projections and differential encoding of prefrontal
neuronal classes underlying learned and innate threat avoidance. *J. Neurosci.* JN-RM-0697-
23 (2023) doi:10.1523/JNEUROSCI.0697-23.2023.

31. Madisen, L. *et al.* A robust and high-throughput Cre reporting and characterization system
for the whole mouse brain. *Nat. Neurosci.* **13**, 133–140 (2010).

32. Chi, J. *et al.* Three-Dimensional Adipose Tissue Imaging Reveals Regional Variation in
Beige Fat Biogenesis and PRDM16-Dependent Sympathetic Neurite Density. *Cell Metab.*
(2018) doi:10.1016/j.cmet.2017.12.011.

33. Do Monte, F., Quirk, G., Li, B. & Penzo, M. Retrieving fear memories, as time goes by….
Mol. Psychiatry **21**, 1027–1036 (2016).

34. Kwapis, J. L., Jarome, T. J., Lee, J. L. & Helmstetter, F. J. The retrosplenial cortex is
involved in the formation of memory for context and trace fear conditioning. *Neurobiol.*
Learn. Mem. **123**, 110–116 (2015).

35. Corcoran, K. A., Frick, B. J., Radulovic, J. & Kay, L. M. Analysis of coherent activity between
retrosplenial cortex, hippocampus, thalamus, and anterior cingulate cortex during retrieval of
recent and remote context fear memory. *Neurobiol. Learn. Mem.* **127**, 93–101 (2016).

36. Marchand, A., Faugère, A., Coutureau, E. & Wolff, M. A role for anterior thalamic nuclei in

contextual fear memory. *Brain Struct. Funct.* **219**, 1575–1586 (2014).

37. Vetere, G. *et al.* An inhibitory hippocampal–thalamic pathway modulates remote memory
retrieval. *Nat. Neurosci.* **24**, 685–693 (2021).

38. Wheeler, A. L. *et al.* Identification of a Functional Connectome for Long-Term Fear Memory
in Mice. *PLoS Comput. Biol.* **9**, e1002853 (2013).

39. Vetere, G. *et al.* Chemogenetic Interrogation of a Brain-wide Fear Memory Network in Mice.
Neuron **94**, 363–374 (2017).

40. De Sousa, A. F. *et al.* Optogenetic reactivation of memory ensembles in the retrosplenial
cortex induces systems consolidation. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 8576–8581
(2019).

41. Trask, S., Ferrara, N. C., Jasnow, A. M. & Kwapis, J. L. Contributions of the rodent
cingulate-retrosplenial cortical axis to associative learning and memory: A proposed circuit
for persistent memory maintenance. *Neurosci. Biobehav. Rev.* **130**, 178–184 (2021).

42. Keene, C. S. & Bucci, D. J. Contributions of the retrosplenial and posterior parietal cortices
to cue-specific and contextual fear conditioning. *Behav. Neurosci.* **122**, 89–97 (2008).

43. Frankland, P. W., Josselyn, S. A. & Köhler, S. The neurobiological foundation of memory
retrieval. *Nat. Neurosci.* **22**, 1576–1585 (2019).

44. Tonegawa, S., Liu, X., Ramirez, S. & Redondo, R. Memory Engram Cells Have Come of
Age. *Neuron* **87**, 918–931 (2015).

45. Josselyn, S. A. & Tonegawa, S. Memory engrams: Recalling the past and imagining the
future. *Science* **367**, (2020).

46. Kitamura, T. *et al.* Engrams and circuits crucial for systems consolidation of a memory.
Science **356**, 73–78 (2017).

47. Todd, T. P., Mehlman, M. L., Keene, C. S., DeAngeli, N. E. & Bucci, D. J. Retrosplenial
cortex is required for the retrieval of remote memory for auditory cues. *Learn. Mem. Cold*
Spring Harb. N **23**, 278–88 (2016).

48. Harris, K. D. & Shepherd, G. M. G. The neocortical circuit: Themes and variations. *Nat.*
Neurosci. **18**, 170–181 (2015).

49. Armbruster, B. N., Li, X., Pausch, M. H., Herlitze, S. & Roth, B. L. Evolving the lock to fit the
key to create a family of G protein-coupled receptors potently activated by an inert ligand.
Proc. Natl. Acad. Sci. **104**, 5163–5168 (2007).

50. Sugar, J., Witter, M., van Strien, N. & Cappaert, N. The Retrosplenial Cortex: Intrinsic
Connectivity and Connections with the (Para)Hippocampal Region in the Rat. An Interactive
Connectome. *Front. Neuroinformatics* **5**, (2011).

51. van Groen, T. & Michael Wyss, J. Connections of the retrosplenial granular a cortex in the
rat. *J. Comp. Neurol.* **300**, 593–606 (1990).

52. van Groen, T. & Wyss, J. M. Connections of the retrosplenial dysgranular cortex in the rat.
J. Comp. Neurol. **315**, 200–216 (1992).

53. Vann, S. D., Aggleton, J. P. & Maguire, E. A. What does the retrosplenial cortex do? *Nat.*
Rev. Neurosci. **10**, 792–802 (2009).

54. Li, S., Callaghan, B. L. & Richardson, R. Infantile amnesia: Forgotten but not gone. *Learn.*
Mem. **21**, 135–139 (2014).

55. Axel Guskjolen *et al.* Neurogenesis-mediated circuit remodeling reduces engram
reinstatement and promotes forgetting. *bioRxiv* 2023.10.10.561722 (2023)
doi:10.1101/2023.10.10.561722.

56. Miller, A. M. P., Vedder, L. C., Law, L. M. & Smith, D. M. Cues, context, and long-term
memory: the role of the retrosplenial cortex in spatial cognition. *Front. Hum. Neurosci.* **8**,
(2014).

57. Maviel, T. Sites of Neocortical Reorganization Critical for Remote Spatial Memory. *Science*
305, 96–99 (2004).

58. Thomas Wolbers & Christian Büchel. Dissociable Retrosplenial and Hippocampal
Contributions to Successful Formation of Survey Representations. *J. Neurosci.* **25**, 3333
(2005).

59. Staresina, B. P., Alink, A., Kriegeskorte, N. & Henson, R. N. Awake reactivation predicts
memory in humans. *Proc. Natl. Acad. Sci.* **110**, 21159–21164 (2013).

60. Opalka, A. N. & Wang, D. V. Hippocampal efferents to retrosplenial cortex and lateral
septum are required for memory acquisition. *Learn. Mem.* **27**, 310–318 (2020).

61. Katche, C. *et al.* On the role of retrosplenial cortex in long-lasting memory storage.
Hippocampus **23**, 295–302 (2013).

62. Keene, C. S. & Bucci, D. J. Neurotoxic lesions of retrosplenial cortex disrupt signaled and
unsignaled contextual fear conditioning. *Behav. Neurosci.* **122**, 1070–1077 (2008).

63. Robinson, S., Keene, C. S., Iaccarino, H. F., Duan, D. & Bucci, D. J. Involvement of
retrosplenial cortex in forming associations between multiple sensory stimuli. *Behav.*
Neurosci. **125**, 578–587 (2011).

64. Freeman, J. H. & Gabriel, M. Changes of Cingulothalamic Topographic Excitation Patterns
and Avoidance Response Incubation over Time Following Initial Discriminative Conditioning
in Rabbits. *Neurobiol. Learn. Mem.* **72**, 259–272 (1999).

65. Bussey, T. J., Muir, J. L., Everitt, B. J. & Robbins, T. W. Dissociable effects of anterior and

995 posterior cingulate cortex lesions on the acquisition of a conditional visual discrimination:

996 Facilitation of early learning vs. impairment of late learning. *Behav. Brain Res.* **82**, 45–56

997 (1996).

998 66. Corcoran, K. A. *et al.* NMDA Receptors in Retrosplenial Cortex Are Necessary for Retrieval

999 of Recent and Remote Context Fear Memory. *J. Neurosci.* **31**, 11655–11659 (2011).

1000 67. VALENSTEIN, E. *et al.* RETROSPLENIAL AMNESIA. *Brain* **110**, 1631–1646 (1987).

1001 68. Gainotti, G., Almonti, S., Di Betta, A. M. & Silveri, M. C. Retrograde amnesia in a patient

1002 with retrosplenial tumour. *Neurocase* **4**, 519–526 (1998).

1003 69. Cowansage, K. K. *et al.* Direct Reactivation of a Coherent Neocortical Memory of Context.

1004 *Neuron* **84**, 432–441 (2014).

1005 70. Aggleton, J. P. EPS Mid-Career Award 2006: Understanding anterograde amnesia:

1006 Disconnections and hidden lesions. *Q. J. Exp. Psychol.* **61**, 1441–1471 (2008).

1007 71. Insausti, R., Amaral, D. G. & Cowan, W. M. The entorhinal cortex of the monkey: II. Cortical

1008 afferents. *J. Comp. Neurol.* **264**, 356–395 (1987).

1009 72. van Groen, T. & Michael Wyss, J. Connections of the retrosplenial granular a cortex in the

1010 rat. *J. Comp. Neurol.* **300**, 593–606 (1990).

1011 73. van Groen, T. & Wyss, J. M. Connections of the retrosplenial dysgranular cortex in the rat.

1012 *J. Comp. Neurol.* **315**, 200–216 (1992).

1013 74. Van Groen, T. & Wyss, J. M. Connections of the retrosplenial granular b cortex in the rat. *J.*

1014 *Comp. Neurol.* **463**, 249–263 (2003).

1015 75. WA Suzuki & DG Amaral. Topographic organization of the reciprocal connections between

1016 the monkey entorhinal cortex and the perirhinal and parahippocampal cortices. *J. Neurosci.*

1017 **14**, 1856 (1994).

1018 76. Burwell, R. D. & Amaral, D. G. Cortical afferents of the perirhinal, postrhinal, and entorhinal

1019 cortices of the rat. *J. Comp. Neurol.* **398**, 179–205 (1998).

1020 77. Morris, R., Petrides, M. & Pandya, D. N. Architecture and connections of retrosplenial area

1021 30 in the rhesus monkey (macaca mulatta). *Eur. J. Neurosci.* **11**, 2506–2518 (1999).

1022 78. Lavenex, P., Suzuki, W. A. & Amaral, D. G. Perirhinal and parahippocampal cortices of the

1023 macaque monkey: Projections to the neocortex. *J. Comp. Neurol.* **447**, 394–420 (2002).

1024 79. Kobayashi, Y. & Amaral, D. G. Macaque monkey retrosplenial cortex: II. Cortical afferents.

1025 *J. Comp. Neurol.* **466**, 48–79 (2003).

1026 80. Choi, H. *et al.* Maturation of metabolic connectivity of the adolescent rat brain. *eLife* **4**,

1027 e11571 (2015).

1028 81. Yasuno, F. *et al.* Resting-state synchrony between the retrosplenial cortex and anterior

medial cortical structures relates to memory complaints in subjective cognitive impairment.

Neurobiol. Aging **36**, 2145–2152 (2015).

82. Düzel, E., Schütze, H., Yonelinas, A. P. & Heinze, H. Functional phenotyping of successful

aging in long-term memory: Preserved performance in the absence of neural compensation.

Hippocampus **21**, 803–814 (2011).

83. Callaghan, B. L., Li, S. & Richardson, R. The elusive engram: what can infantile amnesia tell

us about memory? *Trends Neurosci.* **37**, 47–53 (2014).

84. Arruda-Carvalho, M., Wu, W.-C., Cummings, K. A. & Clem, R. L. Optogenetic Examination

of Prefrontal-Amygdala Synaptic Development. *J. Neurosci.* **37**, 2976–2985 (2017).

85. Mathis, A. *et al.* DeepLabCut: markerless pose estimation of user-defined body parts with

deep learning. *Nat. Neurosci.* **21**, (2018).

86. Gabriel, C. J. *et al.* Behavior DEPOT is a simple, flexible tool for automated behavioral

detection based on marker less pose tracking. *eLife* **11**, 1–33 (2022).

87. Friedmann, D. *et al.* Mapping mesoscale axonal projections in the mouse brain using a 3D

convolutional network. *Proc. Natl. Acad. Sci.* (2020) doi:10.1073/pnas.1918465117.

88. Wang, Q. *et al.* The Allen Mouse Brain Common Coordinate Framework: A 3D Reference

Atlas. *Cell* **181**, 936-953.e20 (2020).

89. Carey, H. *et al.* DeepSlice: rapid fully automatic registration of mouse brain imaging to a

volumetric atlas. *Nat. Commun.* **14**, 5884 (2023).

90. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat. Methods*

9, 676–682 (2012).

91. Chiaruttini, N. *et al.* An Open-Source Whole Slide Image Registration Workflow at Cellular

Precision Using Fiji, QuPath and Elastix. *Front. Comput. Sci.* **3**, (2022).

1052

1053

Figure 1. A developmental screen to identify neural signatures of infantile amnesia.

(A) Infant, juvenile, and adult mice underwent contextual fear conditioning and then memory testing 1 or 7 days later.

(B) Percent time freezing in infant, juvenile and adult wildtype mice during memory retrieval tests occurring 1 or 7 days after CFC ($F_{age}(2,45)=15.89$, P<0.0001, $F_{day}(1,45)=3.56$, P=0.07, $F_{int}(2,45)=4.21$, P=0.02, Infant: N=7 1d, N=10 7d; Juvenile: N=14 1d, N=8 7d; Adult: N=7 1d, N=6 7d, two-way ANOVA with Tukey post-hoc test).

(C_1) Experimental design for developmental screen.

(C_2) Freezing behavior in TRAP mice used in screen (F(2,22)=1.87, P=0.18, One way ANOVA; N=7 infant, N=9 juvenile, N=9 adult).

(D) Representative 100-µm thick optical z-stacks showing TRAPed cells from 1-day retrieval.

(E) TRAPed cell counts by age (Sensory cortex: $F_{region}(2.03,44.7)=267.30$, P<0.0001; $F_{age}(2,22)=0.02$, P=0.97; $F_{int}(10,110)=0.65$, P=0.77; Association cortex: $F_{region}(3.45,75.93)=132.7$, P<0.0001; $F_{age}(2,22)=3.82$, P=0.38; $F_{int}(24,264)=2.14$, P=0.002; Sensory Thalamus: $F_{region}(1.18,26.04)=80.39$, P<0.0001; $F_{age}(2,22)=0.69$, P=0.51; $F_{int}(10,110)=1.05$, P=0.41; Polymodal thalamus: $F_{region}(2.22,48.84)=28.77$, P<0.0001; $F_{age}(2,22)=7.60$, P=0.003; $F_{int}(12,132)=1.59$, P=0.10; Hypothalamus/Midbrain: $F_{region}(3.38,74.38)=141.0$, P<0.0001; $F_{age}(2,22)=3.10$, P=0.07; $F_{int}(10,110)=2.61$, P=0.007; Subplate: $F_{region}(2.96,65.07)=149.7$, P<0.0001; $F_{age}(2,22)=1.93$, P=0.17; $F_{int}(12,132)=1.42$, P=0.17; Hippocampal Formation: $F_{region}(1.94,42.58)=43.78$, P<0.0001; $F_{age}(2,22)=2.16$, P=0.14; $F_{int}(24,264)=0.82$, P=0.72. Infant N=7, Juvenile N=9, Adult N=9; Two-way repeated measure ANOVAs with Tukey post-hoc correction test).

(F-H) Task PLS analysis of TRAPed cell densities in (F) infants, (G) juveniles and (H) adults. Dashed lines indicate threshold of 3. Color labels on x-axis correspond to major brain subdivisions according to the Allen Brain Atlas.

See Tables S1 and S2 for abbreviations and statistics. Graphs show mean ± SEM. *P< 0.05, **P<0.01, ***P<0.001. Scale bars, 100 µm.

Figure 2. Developmental changes in fear memory networks.

(A) Pearson correlation matrices showing interregional correlations based on numbers of TRAPed cells at 1-day memory retrieval across ages. Colors reflect Pearson correlation coefficients (scale, right) and labels within squares correspond to p values from pairwise comparisons.

(B) Mean pearson r values for left: RSP ($F_{region}(4,48)$ = 12.47, P<0.0001; $F_{age}(2,48)$ = 1.04, P=0.37; $F_{interaction}(8,48)$ = 5.60, P<0.0001; HPF: N=3; STR: N=4; BLA: N=3; PHR: N=6; TH: N=5 brain regions) and right: between dorsal cortex (PL, ILA, ACA, RSP) and dorsal polymodal thalamic nuclei (DORpm, $F_{region}(4, 45)$ = 5.82, P=0.0007; $F_{age}(2, 45)$ = 26.48, P<0.0001; $F_{interaction}(8, 45)$ = 1.36, P=0.24; N=4 cortical brain regions). Two-way ANOVAs with Tukey post-

hoc correction test.

(C) Network graphs generated for Pearson's correlations p<0.1 Regions are grouped and colored according to major brain subdivision as defined by the Allen Brain Atlas. The node size is proportional to the number of connections (degree) and the weights of the lines are proportional to the correlation strength.

(D) Average degree (left, $F_{(2,75)}=5.21$, P=0.008, One-way ANOVA) and shortest path length (right, $F_{(2,75)}=18.34$, P<0.0001, One-way ANOVA) for each network node for each age (Infant: N=24; Juvenile: N=28; Adult: N=26 nodes).

(E) Summary of convergent findings from analyses presented in Figures 1 and 2.

*P< 0.05, **P<0.01, ***P<0.001, ****P<0.0001. Graphs show mean $\pm$ SEM. See also Figure S1 for additional network analyses.

Figure S1. Developmental changes in fear memory networks. Related to Figure 1.
Correlation matrices showing the differences in functional connectivity (Fisher Z-transformation) between ages. Fisher correlation coefficients (scale, right; brown and cobalt representing significant differences that are higher in older or younger age groups respectively) and labels within squares correspond to p values of pairwise comparisons of Fisher r-to-z-transformation of the Pearson's correlation coefficient matrices corresponding to respective age groups. *P< 0.05, **P<0.01, ***P<0.001 by one-way ANOVA test.

Figure 3. Selectively increased reactivation of memory-encoding ensembles in RSP in adults.

(A) Illustration of stable vs. dynamic ensembles in a TRAP/Fos overlay assay.

(B) Experiment design for activity-dependent tagging of ensembles between 1- and 7-day retrieval.

(C) Illustration of analysis sites for RSP and ACA.

(D) Representative examples of TRAPed (magenta) and Fos+ cells (green) in the anterior cingulate (top) and retrosplenial cortices (anterior, middle; posterior, bottom).

(E_1) TRAPed cells per ROI for ACA ($F_{(2,16)}=0.34$, P=0.72, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(E_2) Fos+ cells per ROI for ACA each region and age group ($F_{(2,16)}=2.11$, P=0.15, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(E_3) Double-labeled neurons per ROI for ACA ($F_{(2,16)}=0.39$, P=0.68, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=8, Adult: N=7).

(F_1) TRAPed cells per ROI for RSP ($F_{(2,11)}=0.09$, P=0.92, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

(F_2) Fos+ cells per ROI for RSP each region and age group ($F_{(2,11)}=0.70$, P=0.52, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

(F_3) Double-labeled neurons per ROI for RSP ($F_{(2,11)}=7.1$, P=0.01, one-way ANOVA with Tukey post-hoc correction; Infant: N=4, Juvenile: N=5, Adult: N=5).

Graphs show mean $\pm$ SEM. *P <0.05. See also Figure S2 for layer analysis.

Figure S2. Layer-specific distribution of TRAPed and Fos+ cells in ACA and RSP. Related to Figure 3.

(A_1) TRAPed cell distributions across layers in RSP ($F_{layer}(2,33)=45.06$, $P<0.0001$; $F_{age}(2,33)=0.00005$, $P=0.99$; $F_{int}(4,33)=0.96$, $P=0.43$; Infant: N=4; Juvenile: N=5, Adult: N=5).:

(A_2) Fos+ cell distribution across layers in RSP: $F_{layer}(2,33)=93.86$, $P<0.0001$; $F_{age}(2,33)=0.0003$, $P=0.99$; $F_{int}(4,33)=0.23$, $P=0.91$; Infant: N=4; Juvenile: N=5, Adult: N=5).

(A_3) Reactivated cell distribution across layers in RSP ($F_{layer}(2,33)=8.55$, $P=0.001$; $F_{age}(2,33)=7.97$, $P=0.002$; $F_{int}(4,33)=0.53$, $P=0.71$; Infant: N=4; Juvenile: N=5, Adult: N=5).

(B_1) TRAPed cell distributions across layers in ACA ($F_{layer}(2,48)=23.82$, $P<0.0001$; $F_{age}(2,48)=0.0005$, $P=99$; $F_{int}(4,48)=4.25$, $P=0.005$; Infant: N=4; Juvenile: N=8, Adult: N=7).

(B_2) Fos+ cell distribution across layers in ACA ($F_{layer}(2,48)=20.99$, $P<0.0001$; $F_{age}(2,48)=0.0012$, $P=0.99$; $F_{int}(4,48)=5.34$, $P=0.0012$; Infant: N=4; Juvenile: N=8, Adult: N=7).

(B_3) Reactivated cell distribution across layers in ACA ($F_{layer}(2,48)=0.51$, $P=0.60$; $F_{age}(2,48)=1.11$, $P=0.33$; $F_{int}(4,48)=0.14$, $P=0.96$; Infant: N=4; Juvenile: N=8, Adult: N=7).

Figure 4. Chemogenetic reactivation of TRAPed RSC neurons increased freezing during short-term memory retrieval in an age-dependent manner.

(A) Experimental design.

(B) Representative coronal section showing AAV injection site in RSP. Scale bar, 125 μm.

(C) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (brown) groups for infants ($F_{virus}(1,7)=1.29$, $P=0.29$, $F_{day}=(1,7)=24.18$, $P=0.002$; $F_{int}(1,7)=0.47$, $P=0.51$, Cont: N=4, hM3Dq: N=5, two-way ANOVA with Holm-Sidak post hoc test).

(D) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (magenta) groups for juveniles ($F_{virus}(1,28)=1.81$, $P=0.19$, $F_{day}=(1,7)=3.33$, $P=0.07$; $F_{int}(1,7)=4.34$, $P=0.04$, Cont: N=16, hM3Dq: N=14, two-way ANOVA with Holm-Sidak post hoc test).

(E) Percent freezing levels at 1- and 7-day retrieval in control (grey) and hM3Dq (teal) groups for adults ($F_{virus}(1,28)=1.81$, P=0.19, $F_{day}=(1,7)=3.33$, P=0.07; $F_{int}(1,7)=4.34$, P=0.04, Cont: N=8, hM3Dq: N=10, two-way ANOVA with Holm-Sidak post hoc test).

(F) Differences in freezing levels between 7-day and 1-day retrieval sessions for hM3Dq mice in each age group (F(2,26)=12.92, P=0.001, Infant: N=5, Juvenile: N=14, Adult: N=10, one-way ANOVA with Tukey post hoc test).

(G) Representative images showing Fos immunoreactive cells in dorsal caudate putamen (CPd), anteromedial thalamus (AM), laterodorsal thalamus (LD), and anterior pretectal nucleus (APN). Scale bars, 100 µm.

(H) Quantification of Fos+ cells per ROI in 19 brain regions ($F_{region}(18,160)=30.03$, P<0.05; $F_{age}(2,9)=3.69$, P>0.05; $F_{int}(36,160)=5.78$, P<0.05, Mixed Effects Model; Infant: N=4; Juvenile: N=4; Adult: N=3-4).

Graphs show mean $\pm$ SEM. * P<0.05, **P<0.01, ***P<0.001, ****P<0.0001. See also Figure S3 and S4.

Figure S3. Mapping viral targeting in RSP. Related to Figure 4.
(A) Coronal sections showing location of RSP hM3D-mCherry expression in infant-TRAPed mice
(B) Coronal sections showing location of RSP hM3D-mCherry expression in juvenile-TRAPed mice
(C) Coronal sections showing location of RSP hM3D-mCherry expression in adult-TRAPed mice

Figure S4. RSP Fos levels in hM3D-expressing vs. control mice. Related to Figure 4.
(A_1) Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice.
(A_2). Quantification of Fos+ cell density in RSP for infant-TRAPed mice (Cont: N=4, hM3Dq: N=5, P=0.003, two-tailed t-test).
(B_1). Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice. in juvenile-TRAPed mice (Cont: N=6, hM3Dq: N=6, P=0.0001, two-tailed t-test).
(B_2). Quantification of Fos+ cell density in RSP for juvenile-TRAPed mice.
(C_1). Example images RSP showing AAV-mCherry and Fos+ RSP cells adult-TRAPed mice.
(C_2). Quantification of Fos+ cell density in RSP for adult -TRAPed mice (Cont: N=6, hM3Dq: N=6, P<0.0001, two-tailed t-test).
Scale bars, 125 μm. Graphs show mean ± SEM.

Figure 5. Chemogenetic reactivation of learning encoding ensembles in RSC 33d after training recovers infant memories

(A) Experimental design.

(B) Representative coronal section showing AAV injection site in RSP. Scale bar, 125 μm.

(C) Percent freezing levels in contexts A and B during the 33-day retrieval test in control (gray) and hM3Dq (brown) mice ($F_{virus}(1,14)=9.06$, P=0.009; $F_{context}(1,14)=7.41$, P=0.01; $F_{int}(1,14)=5.23$, P=0.03, Cont: N=8, hM3Dq: N=8, two-way ANOVA with Holm-Sidak posthoc test).

(D) Percent freezing levels in contexts A and B during the 40-day retrieval test in control (gray) and hM3Dq (brown) mice ($F_{virus}(1,14)=9.81$, P=0.007; $F_{context}(1,14)=12.3$, P=0.003; $F_{int}(1,14)=4.65$, P=0.07, Cont: N=8, hM3Dq: N=8, two-way ANOVA with Holm-Sidak posthoc test).

(E) Comparison of regional Fos levels in P17-TRAP+40d vs. Adult TRAP from Figure 4 ($F_{region}(18,121)=18.97$, P<0.0001; $F_{age}(1,7)=3.05$, P=0.12; $F_{int}(18,121)=3.5$, P<0.0001, Inf: N=5, Adult: N=4, two-way mixed effects repeated measures analysis with Sidak posthoc test). Graphs show mean ± SEM. *P<0.05. See also Figures S5 and S6.

Figure S5. Mapping viral targeting in RSP. Related to Figure 5.
(A) Coronal sections showing location of RSP hM3D-mCherry expression in infant-TRAPed mice.
(B) Example images RSP showing AAV-mCherry and Fos+ RSP cells infant-TRAPed mice. Scale bars, 125μm.
(C) Quantification of Fos+ cell density in RSP for infant-TRAPed mice (Cont: N=6, hM3Dq: N=6, P<0.0001, two-tailed t-test). Graphs show mean ± SEM.

Figure S6. RSP DREADD validation and controls for Figure 5.
(A) Representative coronal section showing AAV-hM3Dq injection sites in LSX and ACA.
(B) Percent freezing levels in contexts A and B during the 33d retrieval test in LSX- and ACA-targeted mice (LSX: P=0.54, N=3, ACA: P=0.56, N=3, paired t-test). Graphs show mean ± SEM.

Figure 6. Working model

A progressive maturation of RSP ensemble stability and functional connectivity during memory retrieval underlies the emergence of persistent memories across development.