Early survival in Atlantic salmon is associated with parental genotypes at loci linked
to timing of maturation

Tutku Aykanat[1]*, Darryl McLennan[2], Neil B. Metcalfe[2], Jenni M. Prokkola[1,3]

[1]Organismal and Evolutionary Biology Research Programme, Faculty of Biological and Environmental Sciences, University of Helsinki,
Helsinki, Finland

[2]School of Biodiversity, One Health and Veterinary Medicine, University of Glasgow, Glasgow, United Kingdom

[3]Natural Resources Institute Finland (LUKE), Oulu, Finland

*Corresponding author, tutku.aykanat@helsinki.fi

Short running title: Parental maturation genotypes influence offspring survival

Author contributions

14 Conceptualization: TA, DM, NBM, JMP; Formal analysis: TA; Funding acquisition: TA, NBM, JMP;

15 Lab and Fieldwork: TA, DM, NBM, JMP; Writing: TA, DM, NBM, JMP

Abstract

Large effects loci often contain genes with critical developmental functions with potentially broad effects
across life-stages. However, the life-stage-specific fitness consequences are rarely explored. In Atlantic salmon,
variation in two large-effect loci, *six6* and *vgll3*, is linked to age at maturity, and several physiological and
behavioural traits in early life. By genotyping the progeny of wild Atlantic salmon that were planted into natural
streams with nutrient manipulations, we tested if genetic variation in these loci is associated with survival in
early life. We found that higher early life survival was linked to the genotype associated with late maturation in
the *vgll3*, but with early maturation in the *six6* locus. These effects were significant in high-nutrient, but not in
in low-nutrient streams. The differences in early survival were not explained by additive genetic effects in the
offspring generation, but by maternal genotypes in the *six6* locus, and by both parents' genotypes in the *vgll3*
locus. Our results suggest that indirect genetic effects by large-effect loci can be significant determinants of

offspring fitness. This study demonstrates an intriguing case of how large-effect loci can exhibit complex fitness

associations across life stages in the wild and indicates that predicting evolutionary dynamics is difficult.

Keywords: age at maturity, indirect genetic effects, parental fitness effect, *vgll3*, *six6*, Atlantic salmon

Introduction

Adaptation in the wild is a complex process, composed of interactions between many mechanisms, such as

context-dependency of fitness across time and space. Understanding the genetic architecture of fitness has

been one of the key aims of evolutionary quantitative genetics. Classical theory suggests that mutations with

large adaptive effects may hinder fine scale adaptation (Orr 2000, 2005). Yet, several quantitative trait loci

(QTLs) have been identified to be associated with fitness in natural systems (Bradshaw and Schemske 2003;

Colosimo et al. 2005; Johnston et al. 2013; Tishkoff et al. 2007), indicating that adaptive genetic variation via

large effect loci can persist. Orr (2000), on the other hand, suggests that genetic correlations between fitness

components should be selected against. However, Yeaman and Whitlock (2011) predict that under certain

migration-selection scenarios, adaptive loci will be selected for to cluster tightly within a genomic region. As

such, adaptive genomic regions with large phenotypic effects may contain multiple loci associated with

different fitness components, resulting in high genetic correlations among them. How such genetic architecture

is tied to physiological processes remains poorly understood. For example, a genomic region with a large

phenotypic effect may be centrally located within cellular and developmental networks, and the genetic

variation may in turn be linked to broad changes in body plan and/or physiology (Chan et al. 2010; Colosimo et

al. 2005; Jimenez-Gomez et al. 2010; Snoek et al. 2012; Weith et al. 2023). Such broad expression of

phenotypes is likely to alter fitness via different pathways, perhaps across different tissues or life stages. Yet,

little is known of the fitness landscapes of large effect loci across life stages in the wild. Determining the

direction of genetic correlations between fitness components and their potential costs across different

environmental conditions is crucial to a better understanding of how genetic variation is maintained in

populations, and of evolutionary responses to environmental change.

The Atlantic salmon has been emerging as a promising wild model to explore the multitude of fitness effects

associated with large effect loci. Wild salmon populations are adapted to their local environments, so that even

those in close proximity and lacking barriers to migration can exhibit genetic and phenotypic differences

(Taylor 1991; Garcia de Leaniz et al. 2007). At the genetic level, two genomic regions, so-called *vgll3* and *six6*

(named after the most predominant genes within the loci, *vestigial-like family member 3* and *SIX homeobox 6*,

respectively) exhibit substantial signatures of adaptation within and among populations (Barson et al. 2015;

Pritchard et al. 2018). These genomic regions harbour genes encoding transcription (co)factors, and are

expressed as early as embryonic development (Halperin et al. 2013; Kurko et al. 2020; Lee et al. 2012;

Moustakas-Verho et al. 2020). Genetic variation in the *vgll3* underpins key life-history traits in Atlantic salmon,

such as the age at maturity in wild individuals (Ayllon et al. 2015; Barson et al. 2015) and in captivity (Asheim et

al. 2023), the probability of males becoming precociously mature prior to sea migration (Debes et al. 2021),

and iteroparity (Aykanat et al. 2019). *Vgll3* is also associated with physiological and behavioural traits in

juvenile fish, such as lipid profiles (House et al. 2021), aggression (Bangura et al. 2022), and (sex-specific) smolt

migration (Niemela et al. 2022), suggesting that this single QTL can have broad effects across multiple life

stages. In addition, the genetic variation in these loci is associated with metabolic phenotypes, whereby

juvenile salmon with the *vgll3* early genotype exhibit a higher maximum metabolic rate and aerobic scope, with

the former trait also affected by *six6* loci via epistasis (Prokkola et al. 2022). These genomic regions also show

physiological epistasis via the maximal activities of mitochondrial and anaerobic metabolic enzymes in the

heart and intestine (Prokkola et al. 2024). Variation in *six6*, in turn, has been associated with stomach content

volume during ocean feeding (Aykanat et al. 2020). Similar to *vgll3,* the allele frequencies of the *six6* region are

highly correlated with the average age at maturity in wild populations (Barson et al. 2015), and explain

variation in age-at-maturity of a domesticated aquaculture strain of Atlantic salmon (Sinclair-Waters et al.

2020), as well as in Pacific salmonids (Waters et al. 2021). Taken together, these studies suggest that genetic

variation in these two loci may be linked to differences in resource acquisition and allocation in juvenile

salmon, with cascading effects on life-history strategies.

The early life stages of Atlantic salmon are characterized by high rates of mortality, in large part due to intense

density-dependent resource competition (Einum et al. 2006; Grossman and Simon 2019). The availability of

resources such as food and territorial refuge may strengthen or weaken this competition. The likelihood of an

individual surviving to maturation may depend on what resources are available and how capable that individual

is at competing for such resources. Considering the associations of *vgll3* and *six6* with resource acquisition and

allocation, we predict that these loci interact with variation in stream nutrient levels to predict survival rates of

juvenile salmon. However, directional predictions on how freshwater conditions (i.e., nutrient availability)

should lead to adaptive differences in the maturation age of Atlantic salmon (i.e., the orientation and strength

of selection) are challenging to formulate. For example, favourable growth opportunities and increased

freshwater growth could induce early maturation (Salminen 1997), but also reduce the age when salmon

juveniles migrate to sea, which imposes an opposite effect on maturation timing (Marschall et al. 1998). Such

physiological complexities, coupled with context-dependent adaptive responses, make it difficult to formulate

a priori predictions on the gene-phenotype relationship in relation to the adaptive significance of maturation

timing (Barrett et al. 2009; Reznick 2016).

In this study we test whether the *vgll3* and *six6* loci are linked to early life survival in Atlantic salmon. We used

data from a field experiment in which fertilized eggs from the same wild-origin families were divided among 10

experimental streams, all of which had become increasingly oligotrophic in recent years as a result of land

management practices and anthropogenic barriers to the migration of anadromous fish (e.g., Bernthal et al.

2022). Five of these streams were given an experimental supplement to restore nutrient that imitates

conditions of natural spawning levels prior to the river became inaccessible to migrating salmon (Auer et al.

2018). A previous study using the same field system showed that family-level juvenile survival was influenced

by this manipulation of stream nutrient availability in addition to family-level maximum metabolic rate and egg

mass (Auer et al. 2018). In this current study, we test for associations between juvenile survival and genetic

variation in the two life-history loci, *vgll3* and *six6*. More specifically, we first tested whether differences in

survival are due to family-wise genetic effects, i.e., associated with family-level allele frequencies of *vgll3* and

six6, and second, we tested for direct additive effects of the loci, and parental effects, as occurs when the

outcome on offspring survival depends on which parent passed on the allele.

Materials and Methods

The full field protocol is described by Auer et al. (2018). In brief, wild adult Atlantic salmon were trapped on

their upstream spawning migration as part of fishery management operations in the River Conon catchment in

northern Scotland in late autumn 2015 and used to generate 30 full-sib crosses (different parental fish were

used for each full-family) by using *in vitro* fertilization. Genotype data were lost for one family, and so 29

families were used in all analyses in the present study. All parental fish were selected to be fish that had spent

one winter at sea (the most common marine life history phenotype in this catchment, subsequently confirmed

by scale reading) to avoid major differences in egg (and hence initial offspring) size, which could affect among-

family differences in early life survival. Egg mass was estimated for all families using a subset of eggs. Parents

were weighed (to nearest 10 g) and measured for fork length (to nearest 5 mm), and a fin clip was taken for

DNA analyses, i.e., for the parental assignment of each of the recaptured offspring (see below). Condition

factor was calculated using a modified version of Fulton's condition factor, $CF = 100 \times Weight/(Length)^m$, where

m is the slope coefficient of the linear regression between log(weight) (response variable) and log(length), as

the covariate (Bolger and Connolly 1989). This formula corrects for the residual relationship between length (or

weight) and condition factor, hence condition factor can be modelled independently from the length of

individuals.

Eggs were initially reared as family-specific groups and were overwintered in a local hatchery until the eyed

stage of development, when a subsample of eggs from each of the full-sib families were transferred to the

aquarium facilities at the University of Glasgow to be reared under standardized conditions. Hatchery mortality

of eggs and subsequently of fry in the aquarium was negligible (<5%). Meanwhile, another subsample of 100

eggs from each of the same families were planted at the same density in each of 10 streams (in the same River

Conon catchment in which the parent fish had been caught) in February/early March 2016, thus equalling

3,000 eggs per stream. There was no natural spawning of Atlantic salmon in these streams due to the presence

of downstream dams that prevented the upstream migration of spawning fish. All ten streams were

oligotrophic at the start of the experiment. Half of these streams were then experimentally supplemented with

nutrient pellets at the time of egg planting, thereby simulating the decay of parental carcasses (referred to as

the high nutrient streams, Auer et al. 2018), whereas nutrient levels in the other five streams were unaltered

(referred to as the low nutrient streams). Previous ecological analyses of data from this experiment have

shown that these nutrient additions led to significant increases in macroinvertebrate densities, and

consequently faster growth of the young salmon (details in Auer et al. 2018; McLennan et al. 2019).

The maximum metabolic rate (MMR) and standard metabolic rate (SMR) of 10 individuals from each family

were measured using continuous flow-through respirometry in late June 2016, using the offspring that had

been reared under common garden conditions (Auer et al. 2018) following protocols from (Auer et al. 2015).

Standardized triple-pass electrofishing surveys were conducted in each of the experimental streams in July

2016, when the fish were approximately three months post-first-feeding (Auer et al. 2018), to calculate-family

level rates of growth and survival. A small sample of adipose fin tissue was taken from the recaptured

individuals to allow parental (and thus family) assignment (Auer et al. 2018). In the present study, we used

these same DNA samples of the parental fish and of the offspring that survived in the streams to genotype

individuals at the *vgll3* and *six6* loci, using kompetitive allele specific PCR (KASP) method, as described in

Prokkola et al. (2022). Throughout the paper, we refer to the alleles associated with early and late maturation

as E and L, respectively. Consequently, EE and LL will be used for fish that are homozygous for E and L alleles

and EL for heterozygotes, respectively, for both loci (so that the possible genotypes are $vgll3^{EE}$, $vgll3^{LL}$, $vgll3^{EL}$,

and $six6^{EE}$, $six6^{EL}$, $six6^{LL}$). Out of 1246 juveniles that were collected and genotyped, the genotyping success was

96.3% and 97.5% for *six6* and *vgll3* genotypes, respectively. Three individuals were scored with improbable genotypes (one for *six6*, and two for *vgll3* loci) which did not conform to the Mendelian expectations based on parental genotypes. The genotype of one of these individuals, which could not be inferred, was excluded, while the genotypes of two individuals were corrected to the genotype value that was invariable within the full-sib family. Across the 58 parents that were used to generate the 29 families, the gene frequencies were unbalanced towards the early genotypes (0.85 and 0.63 for *vgll3*^{E} and *six6*^{E}, respectively, see Supplementary Table 1), likely due to selecting parents that matured after one year at sea, and of the bias towards fish maturing after one sea winter in this population. In particular, for the *vgll3* locus, there were no *vgll3*^{LL} individuals, and only one family (out of 29) that could generate *vgll3*^{LL} offspring (i.e., family 9 in Supplementary Table 1).

Family-wise early life survival in relation to life-history genotype

We performed a generalized linear mixed model to quantify survival among full-sib families within the high and low nutrient streams in relation to their genotype. We measured local survival as the number of individuals recaptured from the 100 eggs per family that were planted in each stream. While emigration is possible, juvenile salmon are predominantly sedentary in their first year of life, with limited movement and a high rate of mortality (Saunders and Gee 1964), suggesting that the local recapture rate can be mostly attributable to survival rather than emigration. Models were run using the *glmmTMB* package in R version 4.2.2 (Brooks et al. 2022; R Core Team 2021) and fitted using a negative binomial error structure with quadratic parameterization of mean in relation to variance (the *negbin2* family structure in the *glmmTMB* function).

The full model structure is as follows:

$$Y = \mu + SMR_{family} + MMR_{family} + \text{egg mass}_{mother} + vgll3 + six6 + vgll3:six6 + \varepsilon_{family} + \varepsilon_{stream} + \varepsilon_{error}$$

Whereby Y is the number of surviving individuals per family and stream, μ is the intercept, and *vgll3* and *six6* are the family-wise expected genotype frequency in these genomic regions, and *vgll3:six6* indicating the interaction term between *vgll3* and *six6*. In addition, the model included SMR_{family}, MMR_{family}, and egg mass$_{mother}$ which are family-wise mean values for maximum metabolic rate (MMR) and standard metabolic rate (SMR), and mean egg mass per mother, respectively. These physiological indices were added as covariates, since Auer et al. (2018) showed that family-wise local survival was positively related to family-wise MMR and egg mass in the low nutrient streams, and marginally positively related to SMR in the high nutrient streams.

Family-wise expected genotype frequencies were modelled continuously, and calculated based on parental genotypes and segregation of alleles according to the Mendelian inheritance principles. For example, in a

family where the parents' genotypes are EE (0) and EL (0.5), the family-wise expected genotype frequency is

0.25, with 50% probability of offspring to be EE and 50% probability to be EL (see Supplementary Table 1 for

parental genotype information). Family and stream were included in the model as random terms to account for

non-independence among individuals of a given family or in each stream.

We performed a semi-automated model evaluation, using the *dredge* function in the *MuMin* package (version

1.47.1, Barton 2022) in R, by which we evaluated all possible fixed effect structures reduced from the full

model, and ranked their fit according to the AICc value (a version of Akaike information criterion that corrects

for small sample size), in which models within ΔAICc < 2 to the best model considered equally parsimonious. In

a few instances, where convergence have not been archived, models were re-run using *glmer.nb* in *lme4*

package (version 1.1.27.1, Bates et al. 2015), which provides nearly identical parameter estimates and AICc

values. The model fits were checked using the *DHARMa* package (version 0.4.6, Hartig 2022). Marginal means

for genotype effects and their confidence intervals were calculated using the *emmeans* package (version, 1.8.5,

Lenth 2023), whereby the values of continuous covariates were fixed at the mean value, and the value of the

non-focal locus' genotype was fixed at 0.5 (i.e., the EL genotype). We performed separate analyses for low- and

high- nutrient streams to avoid overly complex model structures, since those two environments have been

shown to have different associations with physiological indices (Auer et al. 2018).

Partitioning additive and parental genetic effects

Given that the genotype effects for both *vgll3* and *six6* were significant in high nutrient streams (see Results

below), we then employed individual level analyses to decompose the genetic variation to its additive (direct

genetic effects) and parental genetic effects (indirect genetic effects), and further, to assess the contributions

of each parent's effects (i.e., maternal vs paternal, or both). Teasing out additive vs. parental genetic effects

requires individual level survival data from the offspring, but we obviously did not have the genotype data for

the missing (dead or emigrated) progeny. Therefore, the genotype of the missing progeny was inferred in

accordance with Mendelian segregation of the parental genotypes, and after accounting for the known

genotypes of the recaptured (surviving) individuals. As an example, suppose that for a family with parental

vgll3^{EE} and *vgll3*^{EL} genotypes, nine offspring out of initial 100 implanted eggs survived in a particular stream,

with seven of them being *vgll3*^{EE} and two of them being *vgll3*^{EL} genotype. In this case, and in accordance to

Mendelian segregation laws, we can estimate that N=43 and N=48 offspring are missing (dead or emigrated)

for the *vgll3*^{EE} and *vgll3*^{EL} genotype groups, respectively. A small subset of recaptured individuals with no

genotype data (3.7% and 2.5% of the recaptured individuals for *six6* and *vgll3*, respectively) were randomly

imputed a genotype based on Mendelian segregation of parental genotypes. (As an example, in a family with

parental $vgll3^{EE}$ and $vgll3^{EL}$ genotypes, offspring with missing genotypes would be randomly assigned $vgll3^{EE}$ or

$vgll3^{EL}$ genotypes with an equal (50%) probability). The dataset contained 29000 data points (100 offspring per

29 family and 10 streams), of which N=1246 consisted of recaptured individuals. The effect of sampling

variation, due to both imputing missing genotypes of the recaptured individuals and missing individuals, was

accounted for and quantified by resampling the missing genotypes 1000 times, which then re-analysed as

described below.

We then built and compared the fit of binomial models with different parametrization of parental effects. All

these models included family-wise SMR as a covariate (since this was significant in the high nutrient streams in

the family-wise model – see Results), and the additive effects of the genotypes, but they differed in the

parametrization of the parental effects, whereby each model contained a different combination of parental

effects for $six6$ and $vgll3$ genotypes. As such, the general model structure could be written as follows:

$Y = \mu + SMR_{family} + vgll3_{add.} + six6_{add.} + vgll3_{parent/mother/father} + six6_{parent/mother/father} + \varepsilon_{family} + \varepsilon_{stream} + \varepsilon_{error}$

whereby, Y is a Boolean response variable for local survival (with missing and recaptured individuals scored as 0

and 1, respectively), SMR_{family} is the family-wise mean SMR value, $vgll3_{add.}$ and $six6_{add.}$ are the additive genetic

effects, and $vgll3_{parent/mother/father}$ and $six6_{parent/mother/father}$ are different combinations of parental effects

associated with $six6$ and $vgll3$ loci. In total nine different parental genotype combinations were tested in nine

models, whereby the parental covariates included in the models are as follows: (1) $vgll3_{parent} + six6_{parent}$, (2)

$vgll3_{parent} + six6_{mother}$, (3) $vgll3_{parent} + six6_{father}$, (4) $vgll3_{mother} + six6_{parent}$, (5) $vgll3_{mother} + six6_{mother}$, (6) $vgll3_{mother} +$

$six6_{father}$, (7) $vgll3_{father} + six6_{parent}$, (8) $vgll3_{father} + six6_{mother}$, (9) $vgll3_{father} + six6_{father}$. Family and stream were

included in the models as random terms. The parental genotype value is the average genotype value of mother

and father genotype, e.g., $vgll3_{parent} = (vgll3_{mother} + vgll3_{father}) / 2$, and $six6_{parent} = (six6_{mother} + six6_{father}) / 2$, which

quantifies an overall combined parental genotype effect.

Genotypes were modelled additively (i.e., EE, EL, and LL alleles were coded as 0, 0.5, and 1, respectively) with

no dominance effects for either genotype. The $vgll3$ locus is known to have a sex-dependent dominance effect

on sea-age at maturity, hence testing dominance in this locus would have provided further insights on the

genetic architecture. However, the lack of $vgll3^{LL}$ genotypes in the parental fish and suited cross designs, we

had one family with $vgll3^{LL}$ offspring (Supplementary Table 1), making predicting dominance effect unreliable.

The fit of the models containing different combinations of parental genotypes were then compared according

to the AICc criteria, in which models within ΔAICc < 2 to the best model were considered equally parsimonious.

To account for uncertainty associated with sampling variation of genotype distributions, we also performed

random sampling of genotypes, in which genotypes of missing (dead or emigrated) individuals were randomly

drawn from expected genotype probabilities, in accordance with Mendelian segregation of the parental

genotypes. In this case, we simulated the genotypes of 100 offspring per family/stream (i.e., the initial number

of individuals released per family to each stream) and assigned genotypes for missing individuals after

excluding the genotypes of recaptured (surviving) individuals from the simulated set. Similar to above, a small

subset of recaptured (survived) individuals with no genotype data (3.7% and 2.5% of the recaptured individuals

for *six6* and *vgll3*, respectively) were also randomly assigned a genotype based on Mendelian segregation of

parental genotypes in each replicate, to prevent any sampling biases. The random data sampling described

here was repeated 1000 times.

Results

The observed local survival per family was dependent on genotype in high but not low nutrient streams (see

also the observed family-wise data in Supplementary Figure 1). In the low nutrient streams all parsimonious

models (models within ΔAICc = 2 of the best model) contained egg mass and MMR as covariates, a result that is

consistent with Auer et al. (2018). While the *vgll3* effect was present in the best model, it was insignificant, and

the parameter was not present in all parsimonious models (Figure 1, Supplementary Table 2). In the high

nutrient streams, all parsimonious models contained *six6* and SMR as covariates. In the best model, a higher

SMR (p = 0.018) and $six6^E$ allele (p = 0.025) were significantly and positively associated with local survival, while

there was a marginal negative association with the $vgll3^E$ allele (p = 0.047). Allelic substitution from $vgll3^E$ to

$vgll3^L$ increased the odds of survival 1.67 times (SE = 0.432, p = 0.047), whereby the marginal survival

probability associated with $vgll3^{EE}$ was 3.16% (95% CI 2.46 -4.05), compared to 5.28% (95% CI 3.47 − 8.02) for

$vgll3^{EL}$ (Figure 1). Allelic substitution effect from $six6^E$ to $six6^L$ was in opposite direction, decreasing the odds of

survival 1.49 times (SE = 0.118, p = 0.025), whereby the marginal survival probability associated with $six6^{EE}$ was

7.85% (95% CI 4.94 -12.46), compared to 5.28% (95% CI 3.47 − 8.02) for $six6^{EL}$ (Figure 1). Models were

substantially more parsimonious when both genotype and physiological indices (SMR, MMR and egg mass)

were included as covariates compared to the models that included only genotypes. However, family-wise SMR,

MMR and egg mass were not significantly associated with either genomic region (Supplementary Table 3),

suggesting an independent effect of genotypes and physiological traits on local survival rates. Similarly, an

epistasis between *vgll3* and *six6* loci was not supported (Supplementary Table 3).

For the high nutrient streams, we then used an individual-based, binomial model to partition the significant family-wise genotype effects into additive effects and different combinations of parental genetic effects. A model with only additive genetic terms had substantially lower fit compared to models that additionally parametrized parental genotypes as covariates (Figure 2). Among different combinations of parental genetic effects, the best model included parental *vgll3* and maternal *six6* genetic effects (Figure 2). This model was superior to the second-best model with an AICc differential 2.08 (SE=0.25, Figure 2). An allelic substitution from *vgll3*^{E} to *vgll3*^{L} in parental genotype increased the odds of survival by 1.98 times (SE = 0.60, *p* = 0.023), whereby the marginal survival probability associated with parental genotype *vgll3*^{EE} was 2.44% (95% CI 1.71 -3.46), compared to 4.72% (95% CI 3.09 – 7.15) for *vgll3*^{EL} (Figure 3, Table 1). Likewise, an allelic substitution from *six6*^{E} to *six6*^{L} associated with maternal genotype decreased the odds of survival by 1.58 times (SE = 0.10, *p* = 0.003), whereby the marginal survival probability associated with *six6*^{EE} maternal genotype was 7.25% (95% CI 4.69 - 11.07), compared to 4.72 % (95% CI 3.09 – 7.15) for *six6*^{EL} (Figure 3). The additive (direct) *E* to *L* allelic substitutions was insignificant both for *vgll3* (*p* = 0.51) and *six6* loci (*p* = 0.36, Figure 3, Table 1). The individual-based analyses were robust to sampling variation associated with imputing the missing genotypes of missing individuals, or recaptured individuals that were not successfully genotyped, since similar effect sizes were obtained across 1000 simulated datasets (Supplementary Figure 2). The results were also robust to some families being invariant in *vgll3* and *six6* genotypes, which may result in lower power for quantifying the additive genetic variance compared to the parental genetic effects. As such, when the analyses included only families which harboured genetic variation in these loci (21 and 14 families for *vgll3* and *six6*, respectively, see Supplementary Table 3), the effect sizes and associated *p-values* were similar to those obtained from the full dataset, and parental genetic effect sizes were substantially higher than the additive genetic effects (Supplementary Table 4).

In salmonids, the maternal genotype might be more important to offspring survival than the paternal genotype, since mothers also determine the egg size and composition that influences offspring viability (Fleming 1996). To test the if maternal body size drives the association between local survival and the maternal genetic effects (in *six6* locus), maternal body size traits (i.e., length, weight, and condition factor) were added separately to the most parsimonious models explaining the maternal genotype effects on offspring local survival in the high nutrient streams. None of the maternal size traits had a significant effect on survival and did not improve the model fit, with or without the *six6* maternal effects present in the models (Supplementary Table 5). This suggests that maternal size is not driving the association between offspring survival and maternal *six6* genotypes, although it should be noted that variation in maternal size within the sample was constrained as a result of only using females that had spent a single winter at sea. On the other hand, a maternal *six6*

302 genotype had strong associations with her size and condition (Supplementary Table 6), with the late genotype

303 being associated with larger size and higher body condition. This indicates that a genetic correlation exists

304 between parental size and early offspring survival via *six6*, but the association between *six6* and offspring

305 survival could be mediated by a different process than female body size.

Discussion

Understanding the genetic architecture of adaptation is crucial for predicting how populations respond to

changing environmental conditions at contemporary time scales. Major effect loci may alter phenotypes at

early life stages (i.e., during development), leading to phenotypic and fitness changes operated via multiple

pathways (Colosimo et al. 2005). This study has demonstrated that the loci that are associated with maturation

timing in Atlantic salmon can also affect early life survival, but in a manner that interacts with nutrient levels

and the availability of food in the streams that they inhabit. Intriguingly, this effect did not operate via additive

genetic effects associated with offspring own genotype, but via a maternal effect in *six6* loci, and via parental

effects in *vgll3*. The results suggest that the offspring of parents carrying the *vgll3* late-maturation genotype

and of mothers carrying the *six6* early-maturation genotype will benefit from higher survival, regardless of the

inherited alleles. Such indirect genetic effects add further complexity to evolutionary dynamics compared to

simple Mendelian inheritance (Bonduriansky and Day 2009; Wolf et al. 1998), but have been little studied in

wild populations.

Salmon are well known for exhibiting plasticity in life history traits, particularly in response to environmental

conditions that influence growth rates (Mobley et al. 2021; Sloat et al. 2014). Age at maturity is an important

life history trait that can influence fitness and that has a genetic as well as an environmental component, likely

maintained in the population via balancing selection (Barson et al. 2015). Our results suggest that the viability

selection acting on *vgll3* and *six6* loci in early juvenile stages of salmon will likely alter the age-structure, hence

the dynamics of the evolutionary response in the population. However, the response has a plastic component

and operates in environments with higher nutrient levels. As such, the rate of evolutionary response in age at

maturity may be attributed, in part, to ecological processes operating very early in life. However, the effects of

the freshwater environment on variation in age at maturity are poorly understood, limiting the scope for *a*

priori predictions of the effect of these life-history loci on early survival (see also the Introduction). Similarly,

while our results indicate that a genetic correlation exists between early- and late-life fitness components, it

does not reveal the possible physiological basis or how the observed correlation between fitness components

contributes to the complexity of adaptation (Orr 2000). For example, it is unclear if the direction of selection

associated with early- and late-life fitness components is aligned (i.e., synergistic pleiotropy) or inflicts a cost to

the life-history adaptations (i.e., constraining selection via trade-offs, Orr 2000; Rennison and Peichel 2022;

Thorholludottir et al. 2023). A recent study showed that a rapid evolutionary response towards earlier

maturation (i.e., younger age structure) in the *vgll3* locus was partly driven by the abundance of prey species

once the salmon reach the ocean (Czorlich et al. 2022). However, our results in the high nutrient streams

suggest that selection in earlier life stages favours the opposite *vgll3* allele (i.e., late). One might then predict a

larger evolutionary cost to age-at-maturity evolution at the *vgll3* loci, i.e., via a slower evolutionary response to

a changing environment (and a larger genetic load to the population, Orr 2000). However, we found that the

genetic effects on survival was dependent on nutrient levels, and therefore the potential for correlated

selection with age at maturity is context dependent, suggesting that the cost of any correlation between fitness

components might be population/condition dependent. Finally, the survival effects associated with *six6* and

vgll3 loci are in opposite directions to each other with respect to their effect on maturation timing: while the

$vgll3^L$ allele (the allele linked to later age at maturity) appears to be associated with higher offspring survival, it

is the early maturation allele ($six6^E$) that positively influences offspring survival in the *six6* region, which

necessitates that variation in both loci is accounted for when formulating predictions. Put together, our results

highlight the complexity related to predicting the evolutionary dynamics of maturation age in Atlantic salmon,

and potentially in other taxa where indirect genetic effects have been found (e.g., Bonduriansky and Day 2009;

Tomar and Teperino 2020, Wolf et al. 1998). The observed genetic correlation, however, suggests that a

physiological, mechanistic link exists between genetic polymorphisms in the life-history loci and early survival.

The indirect genetic effect on viability selection via parental genotypes might have theoretical implications for

evolutionary processes (e.g., Bonduriansky and Day 2009; Rausher 1992; Räsänen and Kruuk 2007). In the

classical sense, parental effects are a component of phenotypic plasticity, in which the parental environment

influence the offspring phenotype, through developmental reprogramming (Kappeler and Meaney 2010). As

such, indirect genetic effects via parental genetic variation help to maintain genetic diversity i.e., by reducing the

narrow sense heritability by increasing environmental variance, and provide a means for transmission of acquired

traits between generations (Bonduriansky and Day 2009; Wolf et al. 1998). However, in our context, the indirect

genetic effects of parents are also correlated with the offspring's future life-history and thereby with the fitness of

the offspring via the direct effect of genotypes on age at maturity (Wolf et al. 1998). Therefore, selection on

offspring survival would alter the phenotypic variation of life-history traits, resulting in deviating from the optimal

trait variation maintained under balancing selection (Turelli and Barton 2004), though this prediction is based on

a relatively limited range of genotypes (since the $vgll3^{LL}$ genotypes was completely absent in the dataset) and

phenotypes (since all parents were 1SW fish). Theoretical predictions accounting for such a complexity have not

been explored explicitly (e.g., Oomen et al. 2020).

Our results highlight that the effect of viability selection on these genomic regions is dependent on the

environment, as it appears to only operate under conditions of high nutrient availability. Both loci are linked to

a number of different physiological and behavioural traits that mediate energy homeostasis (Mobley et al.

2021). However, we did not find a direct link between these genomic regions and SMR or MMR (measured at

the family level), thereby suggesting that metabolic rates are not underlying mechanisms for mediating the

effects of *vgll3*/*six6* loci on early survival. Earlier studies have shown a broad range of effects on physiological

and behavioural traits linked to genetic variation in *vgll3* locus, such as MMR (Prokkola et al. 2022), lipid

profiles (House et al. 2021), and aggressive behaviour (Bangura et al. 2022), and that the *six6* genomic region

can be broadly expressed during embryonic developmental (Moustakas-Verho et al. 2020). The effect of these

on fitness is likely to be context-dependent, supporting our conclusions, although the exact mechanisms are

yet to be elucidated. In our study, the genotype dependent selection operating under a more favourable

environment (high nutrient) may seem counterintuitive. However, the higher relative fitness (of different

genotypes) in a less stressful environment is not only theoretically possible, but an earlier meta-analysis

indicated no overall differences in relative fitness under more stressful conditions (Arbuthnott and Whitlock,

2018). Furthermore, there were no differences in absolute fitness between low- and high-nutrient conditions,

despite substantial differences in juvenile growth rates and different metabolic processes shaping the fitness

landscape (Auer et al. 2018), and different impacts of habitat features on the rate of senescence in the fish

(McLennan et al. 2021). These suggests a complex landscape is likely underlying the fitness dynamics across

environment.

The higher local survival of juvenile fish with the early maturation genotype at the *six6* locus was explained

solely by the maternal genotype. Indirect genetic effects associated with the maternal environment (i.e.,

maternal effects) are commonly observed in salmonids and in other organisms (e.g., Aykanat et al. 2012; Heath

et al. 1999; Mousseau and Fox 1998). However, because salmonids lack parental provisioning, these effects are

mostly limited to egg size and/or composition, which in turn can be influenced by the size of the mother

(Munoz et al. 2014). A significant trait-genotype association (i.e., between $six6_{mother}$ and offspring survival /

mother size) despite that the mother's size was constrained (due to using only females that spent a single

winter at sea), suggests a robust inference can be drawn to test for a potential effect of an intermediate

phenotype (egg mass) mediating the relation. Our results showed that in the high nutrient streams, family-level

egg mass was neither correlated with survival (Supplementary Table 3), nor with variation in the $six6_{mother}$

(Supplementary Table 2), thereby ruling out a genetic effect of egg mass on survival via $six6_{mother}$ (within the

constrained size range of mothers). Similarly, we found that maternal size did not explain the correlation

between $six6_{mother}$ and the survival of her offspring, despite the observed correlation between maternal size

and $six6_{mother}$ (Supplementary Table 6).

A plausible mechanism explaining our results is that that variation in *six6* alters early life survival via changes in

egg composition. In oviparous species, the composition of eggs, such as hormone levels, albumin and fat

content, may mediate the phenotype and behaviour of the offspring, and subsequently their fitness (Thorn and

Morbey 2019; Widowski et al. 2022; Williams 1994). In Chinook salmon, carotenoid composition in the eggs has

been shown to be associated with early life survival (Tyndale et al. 2008). Therefore, despite the apparent

absence of a link between the *six6* region and family-level egg mass, we cannot rule out potential effects on

offspring survival via changes in egg composition. Future studies could test such a hypothesis in a common

garden cross design, by quantifying potential markers associated with egg composition, and their effect on the

survival of their full sibs. Alternatively, but less parsimoniously, epigenetic mechanisms may be associated with

a mothers' *six6* genotype, whereby the maternal *six6* locus controls the epigenetic signature of the offspring

genome and thus influences early life developmental pathways (see, Lawson et al. 2013; Venney et al. 2021;

Wolf and Wade 2009). *Six6* and other candidate genes located in the same genomic region (i.e., *six1* and *six4*)

are known transcription factors, and *six6* expression has been documented in Atlantic salmon in early

development (Moustakas-Verho et al. 2020). It is therefore plausible that a mother's *six6* genotype may play a

role in the regulation of her offspring's genome.

The effect of a parent's *vgll3* genotype on offspring early life survival was not solely limited to the maternal

contribution. In salmonids, and generally in fishes, the lack of parental provisioning rules out social

environment such as behavioural variation as a potential basis of indirect genetic effects (e.g., Bailey et al.

2018, Bonduriansky and Day 2009). A plausible mechanism that is consistent with the observed results is that

the genomes inherited from parents are already modified via epigenetic mechanisms, such as via DNA

methylation (Bonduriansky and Day 2009; Kappeler and Meaney 2010; Lawson et al. 2013). Epigenetic

regulation is known to alter phenotypic variation in natural populations, but the underlying genetic

mechanisms and evolutionary predictions are still unclear (Husby 2022). In salmonids, epigenetic effects via

DNA methylation have been documented, for example, between hatchery and wild origin fish associated with

both parents of origin, with implications for evolutionary genetics and conservation efforts (Koch et al. 2022; Le

Luyer et al. 2017; Smith and Kornfield 2002). Our results are consistent with a hypothesis whereby variation in

epigenetic signals (associated with genetic variation in the *vgll3* locus) could result in differences in gene

expression during early development, subsequently altering offspring survival in the natural settings. Such a

hypothesis may be tested by designing cross-breeding experiments, where offspring epigenetic signals, such as

genome wide methylation patterns, could be examined among crosses from parents with different *vgll3*

genotypes, and in relation to offspring phenotype and survival (i.e., Bossdorf et al. 2008). Alternatively, histone

modification is another plausible mechanism that might generate parental genotype-specific modification of

chromatin structure, and subsequent survival differences between families (Koch et al. 2022).

Even though we have shown that the same genomic regions can account for variance in different fitness

components, it is unclear whether covariation between fitness components is causally linked to a single genetic

mutation (true pleiotropy), or takes place as a result of physically linked loci acting on different fitness

components. While molecular evidence suggests that both *vgll3* and *six6* regulate multiple functions across

multiple life stages (Mobley et al. 2021), implying pleiotropy, genetic linkage within the genomic regions cannot

be ruled out. There is accumulating evidence to suggest that co-adapted gene complexes are common in wild

species (Li et al. 2022; Matschiner et al. 2022; Thompson and Jiggins 2014). Moreover, theoretical models have

predicted the evolution of genetic architecture with fewer, larger, and more tightly linked divergent alleles in

species that exhibits marked local adaptation between migratory populations (Yeaman and Whitlock 2011). In

keeping with this model, Atlantic salmon exhibit local adaptation across closely situated populations with both

six6 and *vgll3* genomic regions harbouring substantial signals of divergence, even between adjacent

populations (e.g., Barson et al. 2015; Pritchard et al. 2018). Therefore, it is plausible that the genetic

correlations of fitness components via the *six6* and *vgll3* genomic regions might be explained by independent

loci with tight linkage within these genomic regions. Such genetic architecture may be instrumental in

maintaining the genetic variation and local adaptation of populations amidst high gene flow, although explicit

mathematical models and empirical tests of this hypothesis is still lacking.

Our study re-purposed a dataset from Auer et al. (2018) in order to test the effect of life history genomic

regions on early life survival. While such approaches provide a substantial savings in resources, they come with

unavoidable drawbacks. In particular, we could not explicitly test for dominance effects in the *vgll3* locus,

which exhibits a (sex-dependent) dominance architecture associated with the age-at-maturity genotype

(Barson et al. 2015). This is because the genotypes in the parental fish were either *vgll3*[EE] or *vgll3*[EL], and only

one family (out of 29) had a parental genotype combination that would generate *vgll3*[LL] offspring (i.e., both

parents heterozygous for *vgll3*), making inferences about dominance confounded with the additive effect, and

thus unreliable. For *six6*, we did not have a priori belief that dominance should explain the trait value, and it

was not considered in the main framework. However, modelling dominance in the *six6* locus did not improve

the model fit ($AICc_{dominance} - AICc_{additive} = 2.00$), and the *six6* effect was still insignificant, as it is when modelled additively.

In this study, we provide a unique example from a natural setting of how indirect genetic effects from large-effect loci influencing adult age at maturity and adaptive divergence shape the early-life survival of offspring. Indirect genetic effects are common across organisms and can alter phenotypic variation and evolutionary trajectories, but most empirical evidence is based on transmission of behavioural traits in higher organisms, or has explored typical maternal effects (Wolf et al. 1998; Bonduriansky and Day 2009). The complex genetic architecture of large effect loci that comprise indirect genetic effects between parents and offspring revealed by this study highlights the challenges associated with predicting evolutionary dynamics in a changing world. Little is known of the fitness landscapes of large effect loci across life stages in the wild, and the generality of our results across different Atlantic salmon populations and taxa needs further work. Our study also calls for a better understanding of the mechanisms, ecological role and evolutionary consequences of indirect genetic effects.

Data accessibility

The compiled datasets and R-codes to reproduce the results are available from the dryad repository
https://doi.org/10.5061/dryad.866t1g1xk.

Acknowledgements

We thank the Cromarty Firth Fishery Trust (in particular the late Simon McKelvey), Sonya Auer and our field team for assistance in fieldwork, and Leena Laaksonen and the MES Lab in the University of Helsinki for help with the KASP assays. Funding came from the European Research Council (Advanced Grants 322784 and 834653 to NBM), and from the Research Council of Finland (353388 and 325964 to TA, 348965 to JMP).

References

Arbuthnott, D., & Whitlock, M. C. (2018). Environmental stress does not increase the mean strength of selection. *Journal of Evolutionary Biology, 31*(10), 1599-1606. doi:10.1111/jeb.13351

484
485 Asheim, E. R., Debes, P. V., House, A., Liljestrom, P., Niemela, P. T., Siren, J. P., Erkinaro, J., & Primmer, C. R.
486   (2023). Atlantic salmon (*Salmo salar*) age at maturity is strongly affected by temperature, population
487   and age-at-maturity genotype. *Conservation Physiology, 11*(1), coac086. doi:10.1093/conphys/coac086
488
489 Auer, S. K., Anderson, G. J., McKelvey, S., Bassar, R. D., McLennan, D., Armstrong, J. D., Nislow, K. H., Downie, H.
490   K., McKelvey, L., Morgan, T. A. J., Salin, K., Orrell, D. L., Gauthey, A., Reid, T. C., & Metcalfe, N. B. (2018).
491   Nutrients from salmon parents alter selection pressures on their offspring. *Ecology Letters, 21*(2), 287-
492   295. doi:10.1111/ele.12894
493
494 Auer, S. K., Salin, K., Rudolf, A. M., Anderson, G. J., Metcalfe, N. B., & Hopkins, W. (2015). The optimal
495   combination of standard metabolic rate and aerobic scope for somatic growth depends on food
496   availability. *Funct. Ecol.* 29:479-486. doi:10.1111/1365-2435.12396
497
498 Aykanat, T., Bryden, C. A., & Heath, D. D. (2012). Sex-biased genetic component distribution among
499   populations: additive genetic and maternal contributions to phenotypic differences among populations
500   of Chinook salmon. *Journal of Evolutionary Biology, 25*(4), 682-690. doi:10.1111/j.1420-
501   9101.2012.02462.x
502
503 Aykanat, T., Ozerov, M., Vaha, J. P., Orell, P., Niemela, E., Erkinaro, J., & Primmer, C. R. (2019). Co-inheritance of
504   sea age at maturity and iteroparity in the Atlantic salmon *vgll3* genomic region. *Journal of Evolutionary
505   Biology, 32*(4), 343-355. doi:10.1111/jeb.13418
506
507 Aykanat, T., Rasmussen, M., Ozerov, M., Niemela, E., Paulin, L., Vaha, J. P., Hindar, K., Wennevik, V., Pedersen,
508   T., Svenning, M. A., & Primmer, C. R. (2020). Life-history genomic regions explain differences in Atlantic
509   salmon marine diet specialization. *Journal of Animal Ecology, 89*(11), 2677-2691. doi:10.1111/1365-
510   2656.13324
511
512 Ayllon, F., Kjaerner-Semb, E., Furmanek, T., Wennevik, V., Solberg, M. F., Dahle, G., Taranger, G. L., Glover, K.
513   A., Almen, M. S., Rubin, C. J., Edvardsen, R. B., & Wargelius, A. (2015). The *vgll3* locus controls age at
514   maturity in wild and domesticated Atlantic salmon (*Salmo salar L.*) males. *PLoS Genetics, 11*(11),
515   e1005628. doi:10.1371/journal.pgen.1005628
516
517 Bailey, N. W., Marie-Orleach, L., Moore, A. J., & Simmons, L. (2018). Indirect genetic effects in behavioral
518   ecology: does behavior play a special role in evolution? *Behavioral Ecology, 29*(1), 1-11.
519   doi:10.1093/beheco/arx127
520
521 Bangura, P. B., Tiira, K., Niemela, P. T., Erkinaro, J., Liljestrom, P., Toikkanen, A., & Primmer, C. R. (2022). Linking
522   *vgll3* genotype and aggressive behaviour in juvenile Atlantic salmon (*Salmo salar*). *Journal of Fish
523   Biology, 100*(5), 1264-1271. doi:10.1111/jfb.15040
524
525 Barrett, R. D., Rogers, S. M., & Schluter, D. (2009). Environment specific pleiotropy facilitates divergence at the
526   *Ectodysplasin* locus in threespine stickleback. *Evolution, 63*(11), 2831-2837. doi:10.1111/j.1558-
527   5646.2009.00762.x
528
Barson, N. J., Aykanat, T., Hindar, K., Baranski, M., Bolstad, G. H., Fiske, P., Jacq, C., Jensen, A. J., Johnston, S. E.,
Karlsson, S., Kent, M., Moen, T., Niemela, E., Nome, T., Naesje, T. F., Orell, P., Romakkaniemi, A.,
Saegrov, H., Urdal, K., Erkinaro, J., Lien, S., & Primmer, C. R. (2015). Sex-dependent dominance at a

single locus maintains variation in age at maturity in salmon. *Nature, 528*(7582), 405-408.
doi:10.1038/nature16062

Bartoń K (2022). MuMIn: Multi-Model Inference. R package version 1.47.1.

Bates, D., Mächler, M., Bolker, B., & Walker, S. (2015). Fitting linear mixed-effects models using lme4. *Journal
of Statistical Software, 67*(1). doi:10.18637/jss.v067.i01

Bernthal, F. R., Armstrong, J. D., Nislow, K. H., & Metcalfe, N. B. (2022). Nutrient limitation in Atlantic salmon
rivers and streams: Causes, consequences, and management strategies. *Aquatic Conservation: Marine
and Freshwater Ecosystems, 32*(6), 1073-1091. doi:10.1002/aqc.3811

Bolger, T., & Connolly, P. L. (2006). The selection of suitable indices for the measurement and analysis of fish
condition. *Journal of Fish Biology, 34*(2), 171-182. doi:10.1111/j.1095-8649.1989.tb03300.x

Bonduriansky, R., & Day, T. (2009). Nongenetic inheritance and its evolutionary implications. *Annual Review of
Ecology Evolution and Systematics, 40*, 103-125. doi:10.1146/annurev.ecolsys.39.110707.173441

Bossdorf, O., Richards, C. L., & Pigliucci, M. (2008). Epigenetics for ecologists. *Ecology Letters, 11*(2), 106-115.
doi:10.1111/j.1461-0248.2007.01130.x

Bradshaw, H. D., & Schemske, D. W. (2003). Allele substitution at a flower colour locus produces a pollinator
shift in monkeyflowers. *Nature, 426*(6963), 176-178. doi:10.1038/nature02106

Brooks, M. E., Kristensen, van Benthem, K. J., Magnusson, A., Berg, C. W., Nielsen, A., Skaug, H.J., Maechler, M.,
Bolker, B. M., (2017). glmmTMB balances speed and flexibility among packages for zero-inflated
generalized linear mixed modeling. *The R Journal*, 9(2), 378-400.

Chan, Y. F., Marks, M. E., Jones, F. C., Villarreal, G., Jr., Shapiro, M. D., Brady, S. D., Southwick, A. M., Absher, D.
M., Grimwood, J., Schmutz, J., Myers, R. M., Petrov, D., Jonsson, B., Schluter, D., Bell, M. A., & Kingsley,
D. M. (2010). Adaptive evolution of pelvic reduction in sticklebacks by recurrent deletion of a *Pitx1*
enhancer. *Science, 327*(5963), 302-305. doi:10.1126/science.1182213

Colosimo, P. F., Hosemann, K. E., Balabhadra, S., Villarreal, G., Jr., Dickson, M., Grimwood, J., Schmutz, J.,
Myers, R. M., Schluter, D., & Kingsley, D. M. (2005). Widespread parallel evolution in sticklebacks by
repeated fixation of *Ectodysplasin* alleles. *Science, 307*(5717), 1928-1933. doi:10.1126/science.1107239

Czorlich, Y., Aykanat, T., Erkinaro, J., Orell, P., & Primmer, C. R. (2022). Rapid evolution in salmon life history
induced by direct and indirect effects of fishing. *Science, 376*(6591), 420-423.
doi:10.1126/science.abg5980

Debes, P. V., Piavchenko, N., Ruokolainen, A., Ovaskainen, O., Moustakas-Verho, J. E., Parre, N., Aykanat, T.,
Erkinaro, J., & Primmer, C. R. (2021). Polygenic and major-locus contributions to sexual maturation
timing in Atlantic salmon. *Molecular Ecology, 30*(18), 4505-4519. doi:10.1111/mec.16062

Einum, S., Sundt-Hansen, L., & H. Nislow, K. (2006). The partitioning of density-dependent dispersal, growth
and survival throughout ontogeny in a highly fecund organism. *Oikos, 113*(3), 489-496.
doi:10.1111/j.2006.0030-1299.14806.x

Fleming, I. A. (1996). Reproductive strategies of Atlantic salmon: ecology and evolution. *Reviews in Fish Biology*
and Fisheries, 6(4), 379-416. doi:10.1007/bf00164323

Garcia de Leaniz, C., Fleming, I. A., Einum, S., Verspoor, E., Jordan, W. C., Consuegra, S., Aubin-Horth, N., Lajus,
D., Letcher, B. H., Youngson, A. F., Webb, J. H., Vollestad, L. A., Villanueva, B., Ferguson, A., & Quinn, T.
P. (2007). A critical review of adaptive genetic variation in Atlantic salmon: implications for
conservation. *Biological Reviews of the Cambridge Philosophical Society, 82*(2), 173-211.
doi:10.1111/j.1469-185X.2006.00004.x

Grossman, G. D., & Simon, T. N. (2019). Density-dependent effects on salmonid populations: A review. *Ecology*
of Freshwater Fish, 29(3), 400-418. doi:10.1111/eff.12523

Halperin, D. S., Pan, C., Lusis, A. J., & Tontonoz, P. (2013). Vestigial-like 3 is an inhibitor of adipocyte
differentiation. *Journal of Lipid Research, 54*(2), 473-481. doi:10.1194/jlr.M032755

Hartig F (2022). DHARMa: residual diagnostics for hierarchical (multi-Level / mixed) regression models. R
package version 0.4.6.

Heath, D. D., Fox, C. W., & Heath, J. W. (1999). Maternal effects on offspring size: variation through early
development of chinook salmon. *Evolution, 53*, 1605-1611. doi: 10.1111/j.1558-5646.1999.tb05424.x

House, A. H., Debes, P. V., Kurko, J., Erkinaro, J., Kakela, R., & Primmer, C. R. (2021). Sex-specific lipid profiles in
the muscle of Atlantic salmon juveniles. *Comparative Biochemistry and Physiology Part D: Genomics*
and Proteomics, 38, 100810. doi:10.1016/j.cbd.2021.100810

Husby, A. (2022). Wild epigenetics: insights from epigenetic studies on natural populations. *Proceedings of the*
Royal Society B: Biological Sciences, 289(1968), 20211633. doi:10.1098/rspb.2021.1633

Jimenez-Gomez, J. M., Wallace, A. D., & Maloof, J. N. (2010). Network analysis identifies *ELF3* as a QTL for the
shade avoidance response in Arabidopsis. *PLoS Genetics, 6*(9), e1001100.
doi:10.1371/journal.pgen.1001100

Johnston, S. E., Gratten, J., Berenos, C., Pilkington, J. G., Clutton-Brock, T. H., Pemberton, J. M., & Slate, J.
(2013). Life history trade-offs at a single locus maintain sexually selected genetic variation. *Nature,*
502(7469), 93-95. doi:10.1038/nature12489

Kappeler, L., & Meaney, M. J. (2010). Epigenetics and parental effects. *Bioessays, 32*(9), 818-827.
doi:10.1002/bies.201000015

Koch, I. J., Nuetzel, H. M., & Narum, S. R. (2022). Epigenetic effects associated with salmonid supplementation
and domestication. *Environmental Biology of Fishes, 106*(5), 1093-1111. doi:10.1007/s10641-022-
01278-w

Kurko, J., Debes, P. V., House, A. H., Aykanat, T., Erkinaro, J., & Primmer, C. R. (2020). Transcription profiles of
age-at-maturity-associated genes suggest cell fate commitment regulation as a key factor in the
Atlantic salmon maturation process. *G3: Genes, Genomes, Genetics, 10*(1), 235-246.
doi:10.1534/g3.119.400882

Lawson, H. A., Cheverud, J. M., & Wolf, J. B. (2013). Genomic imprinting and parent-of-origin effects on
complex traits. *Nature Reviews: Genetics, 14*(9), 609-617. doi:10.1038/nrg3543

Le Luyer, J., Laporte, M., Beacham, T. D., Kaukinen, K. H., Withler, R. E., Leong, J. S., Rondeau, E. B., Koop, B. F.,
& Bernatchez, L. (2017). Parallel epigenetic modifications induced by hatchery rearing in a Pacific
salmon. *Proceedings of the National Academy of Sciences of the United States of America, 114*(49),
12964-12969. doi:10.1073/pnas.1711229114

Lee, B., Rizzoti, K., Kwon, D. S., Kim, S. Y., Oh, S., Epstein, D. J., Son, Y., Yoon, J., Baek, K., & Jeong, Y. (2012).
Direct transcriptional regulation of *Six6* is controlled by *SoxB1* binding to a remote forebrain enhancer.
Developmental Biology, 366(2), 393-403. doi:10.1016/j.ydbio.2012.04.023

Lenth R (2023). emmeans: Estimated marginal means, aka least-squares means. R package version 1.8.5.

Li, Q., Lindtke, D., Rodriguez-Ramirez, C., Kakioka, R., Takahashi, H., Toyoda, A., Kitano, J., Ehrlich, R. L., Chang
Mell, J., & Yeaman, S. (2022). Local adaptation and the evolution of genome architecture in threespine
stickleback. *Genome Biology and Evolution, 14*(6). doi:10.1093/gbe/evac075

Marschall, E. A., Quinn, T. P., Roff, D. A., Hutchings, J. A., Metcalfe, N. B., Bakke, T. A., . . . Poff, N. L. (1998). A
framework for understanding Atlantic salmon (*Salmo salar*) life history. *Canadian Journal of Fisheries
and Aquatic Sciences, 55*(S1), 48-58. doi:10.1139/d98-007

Matschiner, M., Barth, J. M. I., Torresen, O. K., Star, B., Baalsrud, H. T., Brieuc, M. S. O., Pampoulie, C.,
Bradbury, I., Jakobsen, K. S., & Jentoft, S. (2022). Supergene origin and maintenance in Atlantic cod.
Nature Ecology and Evolution, 6(4), 469-481. doi:10.1038/s41559-022-01661-x

McLennan, D., Auer, S. K., Anderson, G. J., Reid, T. C., Bassar, R. D., Stewart, D. C., Cauwelier, E., Sampayo, J.,
McKelvey, S., Nislow, K. H., Armstrong, J. D., Metcalfe, N. B., & Rohr, J. (2019). Simulating nutrient
release from parental carcasses increases the growth, biomass and genetic diversity of juvenile Atlantic
salmon. *Journal of Applied Ecology, 56*(8), 1937-1947. doi:10.1111/1365-2664.13429

McLennan, D., Auer, S. K., McKelvey, S., McKelvey, L., Anderson, G., Boner, W., Duprez, J. S., & Metcalfe, N. B.
(2022). Habitat restoration weakens negative environmental effects on telomere dynamics. *Molecular
Ecology, 31*(23), 6100-6113. doi:10.1111/mec.15980

Mobley, K. B., Aykanat, T., Czorlich, Y., House, A., Kurko, J., Miettinen, A., Moustakas-Verho, J., Salgado, A.,
Sinclair-Waters, M., Verta, J.-P., & Primmer, C. R. (2021). Maturation in Atlantic salmon (*Salmo salar*,
Salmonidae): a synthesis of ecological, genetic, and molecular processes. *Reviews in Fish Biology and
Fisheries, 31*(3), 523-571. doi:10.1007/s11160-021-09656-w

Mousseau, T. A., & Fox, C. (1998). *Maternal effects as adaptations*: Oxford University Press.

Moustakas-Verho, J. E., Kurko, J., House, A. H., Erkinaro, J., Debes, P., & Primmer, C. R. (2020). Developmental
expression patterns of six6: a gene linked with spawning ecotypes in Atlantic salmon. *Gene Expression
Patterns, 38*, 119149. doi:10.1016/j.gep.2020.119149

Munoz, N. J., Anttila, K., Chen, Z., Heath, J. W., Farrell, A. P., & Neff, B. D. (2014). Indirect genetic effects underlie oxygen-limited thermal tolerance within a coastal population of chinook salmon. *Proceedings of the Royal Society B: Biological Sciences, 281*(1789), 20141082. doi:10.1098/rspb.2014.1082

Niemela, P. T., Klemme, I., Karvonen, A., Hyvarinen, P., Debes, P. V., Erkinaro, J., Sinclair-Waters, M., Pritchard, V. L., Harkonen, L. S., & Primmer, C. R. (2022). Life-history genotype explains variation in migration activity in Atlantic salmon (*Salmo salar*). *Proceedings of the Royal Society B: Biological Sciences, 289*(1978), 20220851. doi:10.1098/rspb.2022.0851

Oomen, R. A., Kuparinen, A., & Hutchings, J. A. (2020). Consequences of single-locus and tightly linked genomic architectures for evolutionary responses to environmental change. *Journal of Heredity, 111*(4), 319-332. doi:10.1093/jhered/esaa020

Orr, H. A. (2000). Adaptation and the cost of complexity. *Evolution, 54*(1), 13-20. doi:10.1111/j.0014-3820.2000.tb00002.x

Orr, H. A. (2005). The genetic theory of adaptation: a brief history. *Nature Reviews: Genetics, 6*(2), 119-127. doi:10.1038/nrg1523

Pritchard, V. L., Makinen, H., Vaha, J. P., Erkinaro, J., Orell, P., & Primmer, C. R. (2018). Genomic signatures of fine-scale local selection in Atlantic salmon suggest involvement of sexual maturation, energy homeostasis and immune defence-related genes. *Molecular Ecology, 27*(11), 2560-2575. doi:10.1111/mec.14705

Prokkola, J. M., Asheim, E. R., Morozov, S., Bangura, P., Erkinaro, J., Ruokolainen, A., Primmer, C. R., & Aykanat, T. (2022). Genetic coupling of life-history and aerobic performance in Atlantic salmon. *Proceedings of the Royal Society B: Biological Sciences, 289*(1967), 20212500. doi:10.1098/rspb.2021.2500

Prokkola, J. M., Chew, K. K., Anttila, K., Maamela, K. S., Yildiz, A., Asheim, E. R., Primmer, C. R., & Aykanat, T. (2024). Tissue-specific metabolic enzyme levels covary with whole-animal metabolic rates and life-history loci via epistatic effects. *Philosophical Transactions of the Royal Society B: Biological Sciences, 379*(1896), 20220482. doi:10.1098/rstb.2022.0482

R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL https://www.R-project.org/.

Räsänen, K., & Kruuk, L. E. B. (2007). Maternal effects and evolution at ecological time-scales. *Functional Ecology, 21*(3), 408-421. doi:10.1111/j.1365-2435.2007.01246.x

Rausher, M. D. (1992). The measurement of selection on quantitative traits: biases due to environmental covariances between traits and fitness. *Evolution, 46*(3), 616-626. doi:10.2307/2409632

Rennison, D. J., & Peichel, C. L. (2022). Pleiotropy facilitates parallel adaptation in sticklebacks. *Molecular Ecology, 31*(5), 1476-1486. doi:10.1111/mec.16335

Reznick, D. (2016). Hard and soft selection revisited: how evolution by natural selection works in the real world. *Journal of Heredity, 107*(1), 3-14. doi:10.1093/jhered/esv076

723 Salminen, M. (1997). Relationships between smolt size, postsmolt growth and sea age at maturity in Atlantic
724   salmon ranched in the Baltic Sea. *Journal of Applied Ichthyology, 13*(3), 121-130. doi:10.1111/j.1439-
725   0426.1997.tb00111.x
726
727 Saunders, R. L., & Gee, J. H. (1964). Movements of young Atlantic salmon in a small stream. *Journal of the
728   Fisheries Research Board of Canada, 21*(1), 27-36. doi:10.1139/f64-004
729
730 Sinclair-Waters, M., Odegard, J., Korsvoll, S. A., Moen, T., Lien, S., Primmer, C. R., & Barson, N. J. (2020). Beyond
731   large-effect loci: large-scale GWAS reveals a mixed large-effect and polygenic architecture for age at
732   maturity of Atlantic salmon. *Genetics Selection Evolution, 52*(1), 9. doi:10.1186/s12711-020-0529-8
733
734 Smith, P. F. and I. Kornfield. 2002. Phylogeography of Lake Malawi cichlids of the genus Pseudotropheus:
735   significance of allopatric colour variation. *Proc. Roy. Soc. B-Biol. Sci.* 269:2495-2502.
736
737 Snoek, L. B., I. R. Terpstra, R. Dekter, G. Van den Ackerveken, and A. J. Peeters. 2012. Genetical Genomics
738   Reveals Large Scale Genotype-By-Environment Interactions in *Arabidopsis thaliana*. *Front. Genet.* 3:317
739
740 Sloat, M. R., Fraser, D. J., Dunham, J. B., Falke, J. A., Jordan, C. E., McMillan, J. R., & Ohms, H. A. (2014).
741   Ecological and evolutionary patterns of freshwater maturation in Pacific and Atlantic salmonines.
742   *Reviews in Fish Biology and Fisheries, 24*(3), 689-707. doi:10.1007/s11160-014-9344-z
743
744 Taylor, E. B. (1991). A review of local adaptation in Salmonidae, with particular reference to Pacific and Atlantic
745   salmon. *Aquaculture, 98*(1-3), 185-207. doi:10.1016/0044-8486(91)90383-i
746   JOCT 15
747
748 Thompson, M. J., & Jiggins, C. D. (2014). Supergenes and their role in evolution. *Heredity (Edinb), 113*(1), 1-8.
749   doi:10.1038/hdy.2014.20
750
751 Thorholludottir, D. A. V., Nolte, V., & Schlotterer, C. (2023). Temperature-driven gene expression evolution in
752   natural and laboratory populations highlights the crucial role of correlated fitness effects for polygenic
753   adaptation. *Evolution, 77*(9), 2081-2089. doi:10.1093/evolut/qpad132
754
755 Thorn, M. W., & Morbey, Y. E. (2019). Survival under the gravel: the effects of egg size and parental identity on
756   the in situ hatching success of chinook salmon. *Transactions of the American Fisheries Society, 148*(5),
757   991-1003. doi:10.1002/tafs.10192
758
759 Tishkoff, S. A., Reed, F. A., Ranciaro, A., Voight, B. F., Babbitt, C. C., Silverman, J. S., Powell, K., Mortensen, H.
760   M., Hirbo, J. B., Osman, M., Ibrahim, M., Omar, S. A., Lema, G., Nyambo, T. B., Ghori, J., Bumpstead, S.,
761   Pritchard, J. K., Wray, G. A., & Deloukas, P. (2007). Convergent adaptation of human lactase persistence
762   in Africa and Europe. *Nature Genetics, 39*(1), 31-40. doi:10.1038/ng1946
763
764 Tomar, A., & Teperino, R. (2020). Genetic control of non-genetic inheritance in mammals: state-of-the-art and
765   perspectives. *Mammalian Genome, 31*(5-6), 146-156. doi:10.1007/s00335-020-09841-5
766
767 Turelli, M., & Barton, N. H. (2004). Polygenic variation maintained by balancing selection: pleiotropy, sex-
768   dependent allelic effects and G × E interactions. *Genetics, 166*(2), 1053-1079.
769   doi:10.1093/genetics/166.2.1053
770

Tyndale, S. T., Letcher, R. J., Heath, J. W., & Heath, D. D. (2008). Why are salmon eggs red? Egg carotenoids and
early life survival of Chinook salmon (*Oncorhynchus tshawytscha*). *Evolutionary Ecology Research, 10*,
1187-1199.

Venney, C. J., Wellband, K. W., & Heath, D. D. (2021). Rearing environment affects the genetic architecture and
plasticity of DNA methylation in Chinook salmon. *Heredity (Edinb), 126*(1), 38-49. doi:10.1038/s41437-
020-0346-4

Waters, C. D., Clemento, A., Aykanat, T., Garza, J. C., Naish, K. A., Narum, S., & Primmer, C. R. (2021).
Heterogeneous genetic basis of age at maturity in salmonid fishes. *Molecular Ecology, 30*(6), 1435-
1456. doi:10.1111/mec.15822

Weith, M., Grossbach, J., Clement-Ziza, M., Gillet, L., Rodriguez-Lopez, M., Marguerat, S., Workman, C. T.,
Picotti, P., Bahler, J., Aebersold, R., & Beyer, A. (2023). Genetic effects on molecular network states
explain complex traits. *Molecular Systems Biology, 19*(8), e11493. doi:10.15252/msb.202211493

Widowski, T. M., Cooley, L., Hendriksen, S., & Peixoto, M. (2022). Maternal age and maternal environment
affect egg composition, yolk testosterone, offspring growth and behaviour in laying hens. *Scientific
Reports, 12*(1), 1828. doi:10.1038/s41598-022-05491-6

Williams, T. D. (1994). Intraspecific variation in egg size and egg composition in birds: effects on offspring
fitness. *Biological Reviews of the Cambridge Philosophical Society, 69*(1), 35-59. doi:10.1111/j.1469-
185x.1994.tb01485.x

Wolf, J. B., Brodie Iii, E. D., Cheverud, J. M., Moore, A. J., & Wade, M. J. (1998). Evolutionary consequences of
indirect genetic effects. *Trends in Ecology & Evolution, 13*(2), 64-69. doi:10.1016/s0169-
5347(97)01233-0

Wolf, J. B., & Wade, M. J. (2009). What are maternal effects (and what are they not)? *Philosophical
Transactions of the Royal Society B-Biological Sciences, 364*, 1107-1115. doi:10.1098/rstb.2008.0238

Yeaman, S., & Whitlock, M. C. (2011). The genetic architecture of adaptation under migration-selection
balance. *Evolution, 65*(7), 1897-1911. doi:10.1111/j.1558-5646.2011.01269.x

Table

Table 1: Estimated coefficients (fixed effects) and variance (random effects) for individual based binomial
model with local survival as the response variable. The coefficients are from the best-fitted model across
different parametrization of the parental genetic effects in the high nutrient environment.

Fixed effects	Estimate	Std. Error	*z* value	*p* value
(Intercept)	-5.959	1.156	-5.154	<0.0001
Family SMR	16.142	6.945	2.324	0.0201
vgll3 additive	-0.141	0.213	-0.662	0.5080
six6 additive	0.145	0.159	0.911	0.3625
vgll3 parental	1.368	0.602	2.274	0.0230
six6 maternal	-0.914	0.305	-3.002	0.0027
Random effects	Variance			
Family origin	0.1543			
Stream	0.0047			

Figures

Figure 1: Marginal means for probability of survival of offspring in relation to family-wise *vgll3* and *six6*
genotypes for (a) low, and (b) high nutrient streams. Data are plotted for the homozygote early (EE) and
heterozygote (EL) genotypes (no homozygote-LL families were present for either locus). The estimated values
are from the most parsimonious models including both *vgll3* and *six6* effects, which is the best fitting model for
the high nutrient streams and the fourth best model for the low nutrient streams (see Materials and Methods
and Supplementary Table 2 for details). Thin and thick grey bars indicate 50% and 95% confidence intervals,
respectively.

Figure 2: The likelihood differences among individual-based models with different parameterization of the
parental genetic effects explaining survival in the high nutrient environment. ΔAICc = 0 indicates the best fitted
model. All models contained SMR and additive *vgll3* and *six6* genetic effects as covariates. Y-axis indicates the
ΔAICc to the best model, and the dashed line indicates ΔAIC = 2. Error bars represents one standard error of
ΔAICc across 1000 models using 1000 data that simulated the genotypes of dead/emigrated individuals (see
Materials and Methods for details).

Figure 3: Genetic effects on marginal offspring survival probability for the best-fitted parental genetic structure model in the high nutrient streams. Thin and thick grey bars indicate 50% and 95% confidence intervals, respectively. Data are plotted for the homozygote early (EE) and heterozygote (EL) genotypes (no homozygous-LL families were present for either locus).