Title

Glioblastoma cells imitate neuronal excitability in humans

Authors

Tong, T[1,2], Ozsvar, A[2,3], Eschen, JT[1,2], Elbæk, KJ[1,2], Cortnum, SOS[1], Sindby, AKR[1], Kristensen, BW[4,5], Meier, K[1,2,6], Mikic, N[2,7], Sørensen, JCH[1,2], Ting, JT[8], Capogna, M[3], Hou, WH[3*], Korshøj, AR[1,2*]

Affiliations

[1]Department of Neurosurgery, Aarhus University Hospital, Palle Juul-Jensens Boulevard 165, 8200 Aarhus N, Denmark

[2]Department of Clinical Medicine, Aarhus University, Palle Juul-Jensens Boulevard 99, 8200 Aarhus N, Denmark

[3]Department of Biomedicine, Aarhus University, Høegh-Guldbergs Gade 10, 8000 Aarhus C, Denmark.

[4]Department of Pathology, The Bartholin Institute, Rigshospitalet, Copenhagen University Hospital, Blegdamsvej 9, 2100 Copenhagen Ø, Denmark

[5]Department of Clinical Medicine and Biotech Research and Innovation Center (BRIC), University of Copenhagen, Ole Maaløes vej 5, 2200 Copenhagen N, Denmark

[6]Department of Anesthesiology, Aarhus University Hospital, Palle Juul-Jensens Boulevard 165J, 8200 Aarhus N, Denmark

[7]Department of Neurosurgery, Aalborg University Hospital, Syd, Hobrovej 18-22, 9000 Aalborg, Denmark

[8]Allen Institute for Brain Science, 615 Westlake Ave N, Seattle, WA, 98109, USA.

*Corresponding authors:

Anders Rosendal Korshøj, Aarhus University Hospital, Department of Neurosurgery, Palle Juul-Jensens Boulevard J165, 8200, Aarhus N, Denmark. Email: andekors@rm.dk. Telephone: +45 78454400

Wen-Hsien Hou, Aarhus University, Department of Biomedicine, Høegh-Guldbergs Gade 10, 8000 Aarhus C, Denmark. Email: wen-hsien.hou@biomed.au.dk. Telephone: +45 52784084.

Summary

Glioblastomas (GBM) are renowned for their pronounced intratumoral heterogeneity, characterized by a diverse array of plastic cell types, which poses a significant challenge to effective targeting and treatment [1]. Recent research has documented the presence of neuronal-progenitor-like transcriptomic cell states of GBM [2, 3], notably in the leading edge of the tumor, where synaptic input from adjacent neurons drives disease proliferation [4]. However, conflicting observations regarding GBM cell excitability, ranging from non-excitable [5] to neuron-like excitability [6], add complexity to our comprehension of the pathophysiological diversity of GBM cells. Here we established a novel experimental workflow enabling comprehensive and selective investigation of the electrophysiological characteristics of cancer cells and neurons within cancer-infiltrated organotypic tissue specimens from GBM patients, using viral genetic labelling to target cellular subtypes. We observed that GBM cells exhibit distinct electrophysiological features in humans, characterized by hyperexcitability and neuron-like action potential generation. Our research provides direct evidence of excitability and a comprehensive description of the electrophysiological characteristics of GBM cells in the cancer-infiltrated cortex of humans, contributing to a deeper understanding of the cellular biology of GBM. These insights have broader implications for understanding cell-cell interactions in malignant tumors and could inform targeted therapies across diverse cancer types, offering a new lens for tackling tumor heterogeneity.

Main

Glioblastoma (GBM) is known as a highly aggressive and incurable primary brain cancer [7]. These tumors display significant resistance to comprehensive treatment including radiotherapy, chemotherapy, and surgical removal, leaving GBM patients facing a dismal prognosis.[8-10] GBMs are notably diverse in their cellular markup [1-3, 11]. In the dense tumor core, astrocytic (AS-like) and mesenchymal (MES-like) cell states form a resilient, self-repairing network, interconnected by multiple tumor microtubes (TMs) [4, 12-14] and Connexin 43 gap junctions [12, 14, 15]. In contrast, unconnected neural progenitor (NPC-like) states, resembling immature neurons in their migration dynamics, are prevalent at the tumor's infiltrative edge, moving outward from the core to colonize the surrounding brain [4]. These edge cells receive synaptic input from surrounding neurons onto one or two neurite-like TMs stimulating invasion and growth [4, 12-14, 16, 17].

Despite these intriguing features, the electrophysiological properties of GBM cells remain elusive. Specifically, it is unclear if GBM cells at the tumor edge have excitable membranes potentially enabling active integration into the surrounding tumor network and adjacent brain tissue. Preliminary analyses based on animal models and GBM cell lines have indicated a non-excitable nature of GBM cells [5, 18, 19], akin to astrocytes [20]. Conversely, one study on acute human GBM-infiltrated brain slices reported action potentials (APs) in presumed tumor cells with abnormal bipolar and multipolar morphology expressing typical glial markers, such as glial fibrillary acidic protein (GFAP) [21]. By similar notion, a few studies in animal models have reported that some low-grade oligodendroglioma and astrocytoma cells generate aberrant APs [6, 21-23]. Recently, one of these finding was further supported by the identification of a hyperexcitable neuron-like tumor cell entity at the leading edge of IDH-mutated low-grade gliomas in a PDX model, firing single aberrant APs, although these findings could not be reproduced in IDH-wildtype GBM [23].

These contradictory observations highlight the need to elucidate the intrinsic excitability of GBM cells and their physiological role in the interactions with the surrounding brain microenvironment. One of the challenges is to tackle the difficulty of identifying cellular subtypes within the leading edge of the GBM-infiltrated brain regions, as these can be elusive under acute conditions. In addition, the current understanding of the physiological attributes of GBM cells derives mainly from animal models and cell lines, limiting generalizability to humans and highlighting the need for human-based model systems.

To overcome this, we utilized a cutting-edge approach based on stable human organotypic slice cultures in combination with viral transduction tools to label and investigate cells of astrocytic and neuronal lineage selectively. This approach enabled us to identify and characterize the electrophysiological and morphological properties of both GBM cells and neurons in the cancer-infiltrated leading edge of living organotypic human samples to further corroborate their functional importance.

Results

We investigated cancer-infiltrated samples of neocortical brain tissue obtained from GBM patients during fluorescence-guided neurosurgical resection (Extended Data Fig. 1). The

samples were processed with standardized protocols [24-26], rapidly transported to the

laboratory, and cut into 300-μm slices (Fig. 1a). Eligible slices contained areas with structurally

preserved six-layered neocortex as well as regions of dense tumor infiltration (Fig. 1b). Using

whole-cell patch-clamp recording, we immediately investigated the basic electrophysiological

properties of GBM cells at the leading tumor edge and neurons residing in neocortex with no

visible tumor infiltration (Fig. 1b). The leading tumor edge was defined as the area less

infiltrated by the tumor, located immediately superficial to the dense tumor core and deep to

the preserved neocortex (Fig. 1b). Neurons were identified directly under bright field

microscopy at 40X amplification based on morphology and preserved six-layer neocortical

structure, while GBM cells were identified based on morphology and tumor infiltration density.

To expand our sampling capability, and to facilitate accurate preferential cell targeting, we

adopted an *ex vivo* slice culture paradigm [24-26] to selectively label putative GBM cells

expressing GFAP and cells of presumed neuronal lineage expressing human Synapsin-1

(hSyn1) [27], using eGFP-tagged enhancer-based Adeno-associated virus (AAV). GFAP is

extensively expressed in GBM cells and not in neurons, while hSyn1 is a marker of neuronal

identity [27]. We then recorded from GFAP- and hSyn1-labelled cells with soma location in

the leading tumor edge and hSyn1-labelled cells in the structurally preserved cortex (Fig. 1b,

middle). Cultures were preserved for recording for up to 14 days (Fig. 1a, right).

The electrophysiological properties were examined by a standardized protocol testing active-

and passive membrane properties [28] (Fig.1c). Last, we confirmed the morphological and

molecular identity of the recorded cells in the leading edge by using *post hoc*

immunohistochemistry and 3D morphology morphological reconstruction (Fig. 1d).

Infiltrating GBM cells display aberrant action potentials

In acute brain slices, neurons located outside the GBM-infiltrated area displayed typical evoked

APs, serving as a baseline for the electrophysiological behavior of GBM cells (Fig. 2a).

Intriguingly, using whole-cell patch clamp recordings on acute GBM-infiltrated neocortical

slices, we found that morphologically defined GBM cells at the leading tumor edge exhibited

aberrant action potentials (aAPs) when depolarizing the membrane with a positive current

injection (Fig. 2b; Extended Data Fig 2a, b). The identity of the patched and Alexa594/biocytin

labeled GBM cells was validated based on morphology and immunostaining with

colocalization with Olig2, which is commonly expressed in GBM cells at the leading tumor

edge [29]. This marker was extensively expressed in all studied cases based on bulk NGS

diagnostics from biopsies obtained at neighboring sites in the tumor. Olig2$^+$ cells were

differentiated from oligodendrocytes based on morphology and tumor-infiltration density

(Extended Data Fig. 2b, c, Extended Data Table 1).

To better characterize the active membrane properties of GBM cells and neurons, we applied

AAVs to the slice cultures (Extended Data Fig. 3). This allowed for selective fluorescence

labeling, identification, and targeted whole-cell patch clamping of hGFAP-expressing putative

GBM cells (AAV5-hGFAP-hHEbl/E-EGFP-bGHp(A)) and hSyn1-expressing neurons

(AAV1-hSyn1-EGFP-WPRE-hGHp(A)) (Extended Data Fig. 2c, Extended Data Table 1).

Based on their soma location, the labeled cells were classified into three categories: 1) hSyn1-

eGFP$^+$ cells residing outside the GBM-infiltrated area in structurally preserved neocortex,

named hSyn1$^+$ OUT (Fig. 2c), 2) hsyn1-eGFP$^+$ cells within the GBM-infiltrated leading tumor

edge, named hSyn1$^+$ IN (Fig. 2f), and 3) GFAP-eGFP$^+$ cells (Fig. 2i) within the leading tumor

edge, named GFAP$^+$ IN cells, were considered to be GBM cells given the extensive tumor

infiltration and abnormal cellular morphology. APs and aAPs were further defined and

categorized through the normality test of the electrophysiological parameters, followed by a

hierarchical cluster analysis (Fig. 3, Extended Data Fig. 4). We observed that 84.4% of hSyn1$^+$

OUT cells residing outside the GBM-infiltrated area generated normal APs (Fig. 2c-e) [24, 26].

By contrast, only 6.9% of hSyn1$^+$ IN generated APs, while 55.2% exhibited aAP properties

characterized by reduced amplitude and larger half-width (Fig.2f-h). Intriguingly, 56% of

hGFAP$^+$ IN cells exhibited active membrane properties upon depolarization, similar to the

aAPs observed in GBM cells in the acute slices (Fig. 2i-k). The GBM identity of the patched

and Alexa594/biocytin labeled GFAP$^+$ IN cells was further supported by the abnormal

morphology (Extended Data Fig. The GBM identity of the patched and Alexa594/biocytin

labeled GFAP$^+$ IN cells was further supported by the abnormal morphology (Extended Data

Fig. 2b-c), akin to recently identified neuron-like tumor cells in low-grade glioma [23], and by

their colocalization with the GBM marker Olig2 (Extended Data Fig 2b-c). We also

reconstructed the virally labelled hSyn1+ IN cells which tend to have elongated processes and

less primary branching from soma compared to GFAP$^+$ IN cells (Extended Data Fig. 5).

Collectively, these results revealed neuron-like excitability of GBM cells and the abnormal

morphology at the leading tumor edge in the GBM-infiltrated cortex.

We further evaluated the constancy of active membrane properties over culture time for hsyn$^+$

OUT, hsyn1$^+$ IN, and GFAP$^+$ IN (Extended Data Fig. 6). hSyn$^+$ OUT and hsyn1$^+$ IN cells

showed weak or no correlation in each parameter, while GFAP$^+$ IN cells exhibited moderate

correlations, indicating dynamic changes in their voltage-dependent channels, possibly due to rapid GBM cell proliferation during culture.

GBM aAPs differ from neuronal APs

We further analyzed the properties of the observed APs and aAPs by detecting both AP and aAP with an initiation threshold of 20 mV/ms (Fig. 3a). We assessed the active membrane properties such as amplitude, half-width, and rise/decay slopes through a long-pulse depolarization step (Fig. 3c). APs and aAPs were further quantitatively defined and categorized through the normality test of the electrophysiological parameters, followed by a hierarchical cluster analysis (Extended Data Fig. 4). We found that GFAP$^+$ IN cells required a more depolarized threshold to initiate an aAP compared to hSyn1$^+$ OUT cells and hSyn1$^+$ IN cells (Fig. 3d). The aAP generated by GFAP$^+$ IN and hSyn1$^+$ IN cells had lower amplitude, wider half-width, and slower rise and decay slopes compared to hSyn1$^+$ OUT (Fig. 3e-h), while there was no significant difference between GFAP$^+$ IN and hSyn1$^+$ IN cells. Our results thus indicated that the GBM cells displayed physiological features comparable to hSyn1+ IN cells in this area.

Na$_V$ and K$_V$ channels in GBM aAPs

To test if the ion channels required for GFAP+ IN aAP generation were similar to those of neurons, we first applied tetraethylammonium (TEA, 30 mM) to block voltage-gated potassium channels in general. This caused the amplitude of the aAP to decrease notably compared to the baseline (Fig. 4a-c). Additionally, we observed an increase in aAP half-width (Fig. 4d) along with slower rise and decay slopes (Fig. 4e, f). Subsequent application of tetrodotoxin (TTX, 1 µM), a voltage-gated sodium channel blocker, eliminated the aAP (Fig. 5g). These findings elucidate the cellular components involved in GFAP+ IN aAPs and further confirm that the electrical excitability of GBM cells imitates that of neurons.

Passive membrane properties differ between GFAP$^+$ IN and hSyn1$^+$ OUT/IN cells

The resting membrane potential (RMP) of GFAP$^+$ IN and hSyn1$^+$ IN cells in the infiltrated areas was significantly depolarized compared to that of hSyn1$^+$ OUT cells (Fig. 5a). In addition, GFAP$^+$ IN and hSyn1$^+$ IN cells exhibited higher input resistance (Fig. 5b) compared to hSyn1+ OUT cells (Fig. 5b). The RMP and R$_{in}$ of all groups had no significant correlation with days *in vitro*, indicating stable passive membrane properties over the culture period. (Extended Data

Fig. 6). These unique characteristics of GFAP[+] IN cells, relatively depolarized RMP and high

R_{in}, mimic those of NLTs in low-grade gliomas and set them apart from astrocytes, which

typically have a hyperpolarized membrane, lower membrane resistance, and are unable to

generate aAP/APs [20], further supporting the pathological identity of these cells. hSyn1[+] IN

cells exhibited a relatively depolarized resting membrane potential, higher input resistance

compared to hSyn1[+] OUT cells.

Discussion

GBM is an incurable brain cancer characterized by significant cellular heterogeneity and

complex interactions with the local neuronal micro-environment. As the electrophysiological

features of GBM cells remain elusive, their role in neuron-cancer interactions is not fully

understood. Using a human-based acute slice configuration, our study provides novel and

robust evidence of neuron-like electrophysiological features of GBM cells at the leading cancer

invasion zone, contradicting recent findings from organoid and xenograft animal models [5, 18,

19]. These results align with recent findings of neuronal transcriptomic signatures in GBM

cells in this region [4, 23, 29], including NPC-like cell states [4]. In support of our findings, a

recent study has documented the presence of a neuron-like tumor cellular subtype in a

xenograft model of IDH-mutated glioma, using morphology, electrophysiology, and patch-seq

investigation [23]. Although similar findings could not be documented for GBM in the same

study, the electrophysiological and morphological features of the neuron-like low-grade tumor

cells were highly similar to our observations, and exhibited high input resistance,

hyperexcitability, glia-like morphology, and fired single aberrant APs of similar configuration.

The authors further observed that these neuron-like cells had co-occuring transcriptomic

annotations of tumor states and neurons at different levels of differentiation, raising the

question whether neoplastic features are induced in healthy cell states during tumor invasion,

or alternatively that tumor cell states potentially acquire neuronal features at the leading edge

to promote interaction with the surrounding microenvironment [23].

We further introduce a novel *in vitro* organotypic slice culture model designed to facilitate

long-term investigation of human cancer-infiltrated brain tissue using cutting edge viral genetic

labelling to target investigation of specific cellular subtypes. With this approach we found that

the majority of GFAP expressing GBM cells and hSyn1 expressing putative neurons at the

leading tumor edge, both displayed aberrant APs with several similarities [26, 30, 31]. In our

study, we found that 59% of GFAP$^+$/Olig2$^+$ GBM cells at the leading tumor edge were electrically active, displaying hyperexcitability and aAPs, while exhibiting distinct morphology compared to cortical neurons. Moreover, the aAP generation required voltage-gated Na$^+$ and K$^+$ channels. These active membrane properties have not been observed in previous animal studies documenting astrocyte-like passive electrical properties GBM cells residing in the tumor core [23]. The presence of aAPs in the GBM cells may indicate unique and specialized functions influencing the network excitability within the tumor-infiltrated region, perhaps leading to different releasing machinery, invasion ability, proliferation potential [32-34], and local network integration [6, 15], as proposed for neuron-like tumor cells in low-grade glioma [23]. Furthermore, AP generation could be linked to the generation of spontaneous high-frequency pathological discharge, such as epilepsy-like activity commonly observed in GBM patients [35]. Understanding the functional consequence of aAPs in GBM cells may provide a new understanding of the systemic tumor influence of nearby neurons within the infiltrated areas.

Intriguingly, we found a striking similarity in electrophysiological properties between GFAP$^+$/Olig2$^+$ GBM cells and hSyn$^+$/NeuN$^+$ cells located at the leading tumor edge. Specifically, hSyn1$^+$/NeuN$^+$ cells and GFAP$^+$/Olig2$^+$ cells exhibited comparably depolarized resting membrane potentials, higher membrane resistance, and a preference to generate aAPs, relative to neurons in the preserved cortex. As hSyn and NeuN expression indicates neuronal lineage, hSyn1$^+$/NeuN$^+$ cells may represent neurons with an altered and aberrant electrophysiological phenotype due to GBM infiltration. The findings of hyperexcitability within the GBM-infiltrated region, may possibly be influenced by factors like paracrine glutamate secretion from GBM cells [34, 36], loss of GABAergic interneurons [37], glioma-induced neuronal synaptic remodeling using GPC3 as a driver [38], and alterations of potassium handling through the interaction between IGSF3 and Kir4.1 channels [18, 35]. Alternatively, hSyn1$^+$/NeuN$^+$ cells may represent an NPC-like cellular state of GBM, which is commonly known to be enriched at the leading tumor edge [4, 23].

Collectively, our findings support emerging evidence of a complex electrical interplay between neurons and GBM cells at the tumor invasion zone [5, 19]. It is known that neuronal activity can promote the progression of glioma cells through paracrine signaling [34, 36] and direct electrochemical synapses input [4, 5, 19]. Conversely, gliomas can influence neurons by secreting non-synaptic glutamate [32, 33] and synaptogenic factors [38, 39], that drive aberrant

synaptogenesis and enhance hyperexcitability [37]. Our findings provide evidence for one missing link in this loop by which GBM cells are capable of playing an active role in the complex electrical network of the cancer-infiltrated brain. Further exploration using a diverse array of targeted viral tools is needed to unravel the role and importance of specific cell-types in this network, including GABAergic and glutamatergic subtypes, both in the tumor core and invasion zone. To this end multi-modal investigations including cellular electrophysiology and morphology in combination with single-cell, patch-seq, and spatial omics, may provide important and detailed insights towards a comprehensive cellular taxonomy of GBM. Specifically, signature morpho-electric properties of individual cell-types, akin to the recent Brain Cell Census Initiative for healthy cell-type, may reveal crucial mechanisms involved in GBM growth, invasion, resilience, and malignant self-repair, with the prospect of developing targeted therapies.

References

1. Eisenbarth, D. and Y.A. Wang, *Glioblastoma heterogeneity at single cell resolution.* Oncogene, 2023. **42**(27): p. 2155-2165.
2. Wang, Q., et al., *Tumor Evolution of Glioma-Intrinsic Gene Expression Subtypes Associates with Immunological Changes in the Microenvironment.* Cancer Cell, 2017. **32**(1): p. 42-56.e6.
3. Neftel, C., et al., *An Integrative Model of Cellular States, Plasticity, and Genetics for Glioblastoma.* Cell, 2019. **178**(4): p. 835-849.e21.
4. Venkataramani, V., et al., *Glioblastoma hijacks neuronal mechanisms for brain invasion.* Cell, 2022. **185**(16): p. 2899-2917.e31.
5. Venkataramani, V., et al., *Glutamatergic synaptic input to glioma cells drives brain tumour progression.* Nature, 2019. **573**(7775): p. 532-538.
6. Labrakakis, C., et al., *Action potential-generating cells in human glioblastomas.* J Neuropathol Exp Neurol, 1997. **56**(3): p. 243-54.
7. Louis, D.N., et al., *The 2021 WHO Classification of Tumors of the Central Nervous System: a summary.* Neuro Oncol, 2021. **23**(8): p. 1231-1251.
8. Wen, P.Y., et al., *Glioblastoma in adults: a Society for Neuro-Oncology (SNO) and European Society of Neuro-Oncology (EANO) consensus review on current management and future directions.* Neuro Oncol, 2020. **22**(8): p. 1073-1113.
9. Wick, W., et al., *Treatment of glioblastoma in adults.* Therapeutic Advances in Neurological Disorders, 2018. **11**: p. 1756286418790452.
10. Ostrom, Q.T., et al., *CBTRUS Statistical Report: Primary Brain and Other Central Nervous System Tumors Diagnosed in the United States in 2015–2019.* Neuro-Oncology, 2022. **24**(Supplement_5): p. v1-v95.
11. Verhaak, R.G., et al., *Integrated genomic analysis identifies clinically relevant subtypes of glioblastoma characterized by abnormalities in PDGFRA, IDH1, EGFR, and NF1.* Cancer Cell, 2010. **17**(1): p. 98-110.

307   12.   Winkler, F., et al., *Cancer neuroscience: State of the field, emerging directions.* Cell,
308   2023. **186**(8): p. 1689-1707.

309   13.   Osswald, M., et al., *Brain tumour cells interconnect to a functional and resistant*
310   *network.* Nature, 2015. **528**(7580): p. 93-98.

311   14.   Heuer, S. and F. Winkler, *Glioblastoma revisited: from neuronal-like invasion to*
312   *pacemaking.* Trends Cancer, 2023. **9**(11): p. 887-896.

313   15.   Hausmann, D., et al., *Autonomous rhythmic activity in glioma networks drives brain*
314   *tumour growth.* Nature, 2023. **613**(7942): p. 179-186.

315   16.   Jung, E., et al., *Tweety-Homolog 1 Drives Brain Colonization of Gliomas.* The Journal
316   of Neuroscience, 2017. **37**(29): p. 6837-6850.

317   17.   Gritsenko, P.G., et al., *p120-catenin-dependent collective brain infiltration by glioma*
318   *cell networks.* Nature Cell Biology, 2020. **22**(1): p. 97-107.

319   18.   Madadi, A., et al., *Correlation between Kir4.1 expression and barium-sensitive*
320   *currents in rat and human glioma cell lines.* Neuroscience Letters, 2021. **741**: p. 135481.

321   19.   Venkatesh, H.S., et al., *Electrical and synaptic integration of glioma into neural*
322   *circuits.* Nature, 2019. **573**(7775): p. 539-545.

323   20.   McNeill, J., et al., *Ion Channels and Electrophysiological Properties of Astrocytes:*
324   *Implications for Emergent Stimulation Technologies.* Frontiers in Cellular
325   Neuroscience, 2021. **15**.

326   21.   Bordey, A. and H. Sontheimer, *Electrophysiological properties of human astrocytic*
327   *tumor cells In situ: enigma of spiking glial cells.* J Neurophysiol, 1998. **79**(5): p. 2782-
328   93.

329   22.   Patt, S., et al., *Neuron-like physiological properties of cells from human*
330   *oligodendroglial tumors.* Neuroscience, 1996. **71**(2): p. 601-11.

331   23.   Curry, R.N., et al., *Identification of functional immune and neuronal tumour cells in*
332   *glioma.* bioRxiv, 2022: p. 2022.11.05.515316.

333   24.   Ting, J.T., et al., *A robust ex vivo experimental platform for molecular-genetic*
334   *dissection of adult human neocortical cell types and circuits.* Scientific Reports, 2018.
335   **8**(1): p. 8407.

336   25.   Louth, E.L., et al., *Dopaminergic Neuromodulation of Spike Timing Dependent*
337   *Plasticity in Mature Adult Rodent and Human Cortical Neurons.* Frontiers in Cellular
338   Neuroscience, 2021. **15**.

339   26.   Lee, B.R., et al., *Signature morphoelectric properties of diverse GABAergic*
340   *interneurons in the human neocortex.* Science, 2023. **382**(6667): p. eadf6484.

341   27.   Finneran, D.J., et al., *Toward Development of Neuron Specific Transduction After*
342   *Systemic Delivery of Viral Vectors.* Front Neurol, 2021. **12**: p. 685802.

343   28.   Berg, J., et al., *Human neocortical expansion involves glutamatergic neuron*
344   *diversification.* Nature, 2021. **598**(7879): p. 151-158.

345   29.   De Silva, M.I., B.W. Stringer, and C. Bardy, *Neuronal and tumourigenic boundaries of*
346   *glioblastoma plasticity.* Trends Cancer, 2023. **9**(3): p. 223-236.

347   30.   Maroso, M., *A quest into the human brain.* Science, 2023. **382**(6667): p. 166-167.

348   31.   Siletti, K., et al., *Transcriptomic diversity of cell types across the adult human brain.*
349   Science, 2023. **382**(6667): p. eadd7046.

350   32.   Campbell, S.L., S.C. Buckingham, and H. Sontheimer, *Human glioma cells induce*
351   *hyperexcitability in cortical networks.* Epilepsia, 2012. **53**(8): p. 1360-70.

352   33.   Buckingham, S.C., et al., *Glutamate release by primary brain tumors induces epileptic*
353   *activity.* Nat Med, 2011. **17**(10): p. 1269-74.

354   34.   Venkatesh, H.S., et al., *Neuronal Activity Promotes Glioma Growth through*
355   *Neuroligin-3 Secretion.* Cell, 2015. **161**(4): p. 803-16.

35. Curry, R.N., et al., *Glioma epileptiform activity and progression are driven by IGSF3-*
mediated potassium dysregulation. Neuron, 2023.
36. Venkatesh, H.S., et al., *Targeting neuronal activity-regulated neuroligin-3 dependency*
in high-grade glioma. Nature, 2017. **549**(7673): p. 533-537.
37. Campbell, S.L., et al., *GABAergic disinhibition and impaired KCC2 cotransporter*
activity underlie tumor-associated epilepsy. Glia, 2015. **63**(1): p. 23-36.
38. Yu, K., et al., *PIK3CA variants selectively initiate brain hyperactivity during*
gliomagenesis. Nature, 2020. **578**(7793): p. 166-171.
39. Krishna, S., et al., *Glioblastoma remodelling of human neural circuits decreases*
survival. Nature, 2023.
40. *Declaration of Helsinki – Ethical Principles for Medical Research Involving Human*
Subjects. 2008.
41. Ahrens, L.C., et al., *Effect of 5-Aminolevulinic Acid and Sodium Fluorescein on the*
Extent of Resection in High-Grade Gliomas and Brain Metastasis. Cancers (Basel),
2022. **14**(3).
42. Hansen, R.W., et al., *Comparison of 5-aminolevulinic acid and sodium fluorescein for*
intraoperative tumor visualization in patients with high-grade gliomas: a single-center
retrospective study. J Neurosurg, 2019: p. 1-8.
43. Hou, W.H., et al., *Wiring Specificity and Synaptic Diversity in the Mouse Lateral*
Central Amygdala. J Neurosci, 2016. **36**(16): p. 4549-63.
44. Myatt, D.R., et al., *Neuromantic - from semi-manual to semi-automatic reconstruction*
of neuron morphology. Front Neuroinform, 2012. **6**: p. 4.

Figure legends

Fig.1 Workflow for utilizing patient-derived GBM-infiltrated cortical tissue. a, Schematic outline of the processing of a human GBM-infiltrated neocortical specimen using acute and cultured paradigms. **b**, Cell targeting via brightfield microscopy and an adeno-associated virus-assisted gene transduction technique to selectively express fluorescence proteins in hSyn1-eGFP[+] cells (neurons) or GFAP-eGFP[+] cells (GBM cells). Scale bar under bright field is 20µm. In the 10× magnification image, the scale bar corresponds to 100µm. In the 40× magnification image, the scale bar represents 20µm. **c**, Electrophysiological characterization using fluorescence-guided whole-cell patch clamp recordings. **d**, Immunohistochemistry co-localization of Olig2 onto a Alexa594/biocytin loaded patched GFAP-eGFP[+] cells indicating GBM identity.

Fig. 2 GBM cells exhibit aAPs. a, Representative whole-cell recordings of neurons in acute brain slices, exhibiting APs. **b**, Representative whole-cell recordings of GBM cells in acute brain slices, displaying aAP activity. **c**, representative bright field image of a hSyn1[+] OUT cell expressing virally transduced hSyn1-eGFP and loaded with Alexa594 during whole-cell patch clamp recording. Scale bar: 20µm. **d**, Representative whole-cell recordings of hSyn1[+] OUT cells from cultured brain slices, with the majority of hSyn1[+] OUT displayed characteristic APs, while a subset of hSyn1[+] OUT exhibits aAP. **e**, Among the 28 hSyn1[+] OUT cells recorded from cultured slices, 22 displayed normal AP (78.5%), 4 exhibited aAP (14.3%), and 2 neurons (7.1%) show none despite membrane depolarization. **f**, Representative bright field image depicting a hSyn1[+] IN expressing eGFP fluorescent protein within the GBM-infiltrated cortical area. The hSyn1[+] IN cell was loaded with Alexa594 during whole-cell patch clamp recording. Scale bar: 20µm. **g**, Representative hSyn1[+] IN recording, displaying both AP and aAP. **h**, In the hSyn1[+] IN group, 16/29 (55.2%) cells exhibited aAP, 2/29 (6.9%) displayed APs, and 11/29 (37.9%) showed no activation. **i**, The illustrated image is a patched- GFAP[+] IN cell expressing eGFP within the cancer-infiltrated cortical area. The GFAP[+] IN cells were loaded with Alexa594 during whole-cell patch clamp recording. Scale bar: 20µm. **j**, Illustration showing the recorded aAP observed in GFAP[+] IN cells. **k**, Out of the 50 recorded GFAP[+] IN cells, 28 exhibited aAP (56%), and 22 showed no detectable response (44%).

Fig. 3. Electrophysiological characteristics of aAP/APs in GBM cells and Neurons. a-b, depictions of an AP **(a)** and aAP **(b)** during a 3ms-depolarization. Parameters measured were threshold, amplitude, half-width, and rise/decay slopes. **c.** Five parameters were compared between neurons (n = 8 cells) and GBM cells (n = 8 cells) from acute slices. Neurons showed a more depolarized threshold potential (-33.40 ± 4.14 mV, **** $P < 0.001$), a higher amplitude (67.28 ± 9.075 mV, *** $P < 0.005$), a shorter half-width (1.55 ± 0.18 ms, * $P < 0.05$), a faster rise slope (204.5 ± 49.46 mV/ms, ** $P < 0.01$), and a faster decay slope (54.37 ± 6.43 mV/ms, **** $P < 0.001$) than the GBM cells (threshold: -4.67 ± 2.22 mV, amplitude: 16.00 ± 3.23 mV, half-width: 3.02 ± 0.46 ms, rise slope: 26.51 ± 3.83 mV/ms, decay slope: 15.66 ± 2.02 mV/ms). **d**, from a holding potential of –60 mV, GFAP[+] IN cells showed a significantly more depolarized threshold potential (-14.24 ± 1.31 mV, n = 28 cells) than hSyn1[+] OUT cells (-31.60 ± 1.21 mV, n = 26 cells **** $P < 0.001$) with no significant difference between GFAP[+] IN cells and hSyn1[+] IN cells (-18.25 ± 1.82mV, n = 18 cells). hSyn1[+] IN cells demanded more

depolarization to generate APs/aAPs compared to hSyn1$^+$ OUT cells (**** $P < 0.001$). **e,** GFAP$^+$ IN cells (32.91 ± 2.67 mV) and hSyn1$^+$ IN cells (38.61 ± 5.1 mV) showed significantly lower amplitudes than hSyn1$^+$ OUT cells (75.84 ± 2.77 mV) (**** $P < 0.001$) with no significant difference between GFAP$^+$ IN cells and hSyn1$^+$ IN cells. **f,** GFAP$^+$ IN cells (5.21± 0.70 ms, *** $P < 0.005$) and hSyn1$^+$ IN cells (4.92 ± 1.90 ms, ** $P < 0.01$) exhibited a broader half-width compared to hSyn1$^+$ OUT cells (1.56 ± 0.14 ms) with no significant difference between GFAP$^+$ IN cells and hSyn1$^+$ IN cells. **g,** GFAP$^+$ IN cells (46.24 ± 5.72 mV/ms, **** $P < 0.001$) and hSyn1$^+$ IN cells (82.81 ± 20.17 mV/ms, **** $P < 0.001$) exhibited slower rise slopes than hSyn1$^+$ OUT cells (247.3 ± 24.88 mV/ms) with no significant difference between GFAP$^+$ IN cells and hSyn1$^+$ IN cells. **h,** GFAP$^+$ IN cells (20.18 ± 1.38 mV/ms, **** $P < 0.001$) and hSyn1$^+$ IN cells (33.83 ± 4.10 mV/ms, **** $P < 0.001$) had slower decay slopes than hSyn1$^+$ OUT cells (70.93 ± 5.45 mV/ms), without no significant difference between GFAP$^+$ IN cells and hSyn1$^+$ IN cells. **i,** Hierarchical cluster analysis of all recorded excitable cells with electrophysiological parameters for classification in Extended Data Fig 4. The x-axis of the dendrogram represents the individual cells. Distinct cell types are marked below the dendrogram.

Fig. 4. Contribution of Na$_V$ and K$_V$ Channels in aAP Generation. a, Illustrative aAPs induced by a 1-second square pulse depolarization of GFAP+ IN cells. **b,** Illustrations presenting changes in aAP morphology upon application of the K$_V$ blocker TEA, highlighting alterations in comparison to the control condition. **c,** comparison of amplitude parameter for 14 GFAP IN cells under three conditions: Control (35.13 ± 3.53 mV), TEA (25.32 ± 3.34 mV), and TEA+TTX (0 mV), exhibiting a reduction in aAP amplitude in the presence of TEA ($P < 0.05$) and a complete absence in the combined presence of TEA and TTX ($P < 0.001$). TTX is a Na$_V$ channel blocker. **d,** comparison of half-width parameter under three conditions: Control (3.39 ± 0.43 ms), TEA (11.93 ± 2.64 ms), and TEA+TTX (0 ms), exhibiting a significant wider in the presence of TEA ($P < 0.05$) and a complete absence in the combined presence of TEA and TTX ($P < 0.001$). **e,** comparison of rise slope parameter under three conditions: Control (46.97 ± 7.44 mV/ms), TEA (26.80 ± 4.70 mV/ms), and TEA+TTX (0 mV/ms), exhibiting a significant reduction in the presence of TEA ($P < 0.01$) and a complete absence in the combined presence of TEA and TTX ($P < 0.001$). **f,** comparison of decay slope parameter under three conditions: Control (20.77 ± 2.14 mV/ms), TEA (13.34 ± 0.94 mV/ms), and TEA+TTX (0 mV/ms), exhibiting a significant reduction in the presence of TEA ($P < 0.01$) and a complete absence in the combined presence of TEA and TTX ($P < 0.001$). Comparative analysis among paired data from three groups was performed using a Two-way ANOVA followed by Tukey's multiple comparison test. **g,** Absence of aAP induction upon depolarization after the combined application of both TEA and TTX.

Fig. 5. GBM cells exhibit different intrinsic membrane properties compared to neurons. **a,** hSyn1$^+$ OUT cells displayed a resting membrane potential (RMP) of -57.68 ± 1.76 mV (n = 31 cells) compared to -31.95 ± 2.18 mV (n = 55 cells) in GFAP$^+$ IN cells and -37.66 ± 4.32 mV (n = 25 cells) in hSyn1$^+$ IN cells, thus GFAP$^+$ IN cells and hSyn1$^+$ IN were more depolarized compared to hSyn1$^+$ OUT cells (One-way ANOVA, Tukey's multiple comparison test, **** $P < 0.001$). No significant difference was observed between hSyn1$^+$ IN and GFAP$^+$ IN cells. **b,** hSyn1$^+$ IN (1.53 ± 0.25 GΩ, n = 28 cells) and GFAP$^+$ IN cells (1.27 ± 0.19 GΩ, n

464 = 62 cells) displayed a significantly higher input resistance (R_{in}) compared to hSyn1$^+$ OUT
465 (0.51 ± 0.13 GΩ, n = 33 cells) (One-way ANOVA, Tukey's multiple comparison test, ** $P <$
466 0.01, * $P < 0.05$).
467

Methods

Glioblastoma tissue acquisition

All procedures with human tissue and data were approved by and conducted in accordance
with the Scientific Ethics Committee for the Region of Midtjylland Denmark (official name in
Danish: De Videnskabsetiske Komitéer for Region Midtjylland) (journal number: 1-10-72-82-
17) and conducted in accordance with the ethical principles of the World Medical Association
Declaration of Helsinki [40]. Surgical specimens were collected at the neurosurgical
department at Aarhus University Hospital (Denmark) through collaboration with local
surgeons. Prior to participating in the study, all patients gave informed consent for the donation
of tissue. The patients included in the study were undergoing surgery for GBM, and the
obtained specimens were surgically excised cancer-infiltrated cortical tissue that required
removal as part of the surgical treatment. A total of nine patients were included in the study
(Extended Data Table 1), and the diagnoses were confirmed based on pathology examination
and standardized WHO criteria [7].
Preoperatively, the patients were administered the fluorescent compounds 5-Aminolevulinic
acid hydrochloride (5-ALA) (20 mg/kg bodyweight orally administered) and sodium
fluorescein (200 mg IV) [41, 42], which allowed the neurosurgeon to visualize the tumor and
the areas infiltrated by the tumor. The surgical specimens, ranging between 1-2 cm^3 in volume,
were resected from the superficial cortical regions and mpassed the contrast-enhancing rim of
the tumor. The procedure did not impose any constraints on the patients or their surgical
treatment. Immediately after resection, the excised specimens were placed in a sterile container
filled with slicing artificial cerebrospinal fluid (ACSF) as quickly as possible. The slicing
ACSF comprised (in mM):75 Sucrose, 84 NaCl, 2.5 KCl, 1 NaH_2PO_4, 25 $NaHCO_3$, 25 D-
glucose, 0.5 $CaCl_2 \cdot 2H_2O$ and 4 $MgCl_2 \cdot 6H_2O$. The pH of the solution was adjusted to 7.3–7.4
using concentrated hydrochloric acid, and its osmolality was confirmed to be between 295–
305 mOsm kg^{-1}. Before collection, the solution was pre-chilled and thoroughly bubbled with
carbogen (95% O_2, 5% CO_2) gas. During transportation, the tissue was continuously kept in a
carbogenated slicing ACSF and maintained at a constant temperature. The duration of tissue
transport from the operating room to the laboratory site ranged from 15 to 30 minutes.
497

Tissue processing

The human *ex vivo* brain slice culture method was developed based on previously established procedures [24-26]. The surgical specimens were carefully examined to determine the optimal approach for blocking and mounting them onto the vibrating microtome platform. To preserve the integrity of dendrites and axons in the GBM-infiltrated region, the slicing angle was maintained perpendicular to the pial surface. Human GBM-infiltrated brain specimens were cut in sterile-filtered and chilled slicing solution that was saturated with carbogen gas to maintain tissue oxygenation throughout the processing steps. Slices of 300 μm thickness were cut using a Leica 1200S vibratome (Leica Microsystems, Denmark) and subsequently transferred to a carbogenated (95% O_2/5% CO_2) and tempered (34 °C) slicing ACSF to recover for 30 min. All the brain slices were transferred to sterile-filtered, room temperature, and low-Calcium ACSF of the composition (in mM): 130 NaCl, 3.5 KCl, 1 NaH_2PO_4, 24 $NaHCO_3$, 10 D-glucose, 1 $CaCl_2 \cdot 2H_2O$ and 3 $MgCl_2 \cdot 6H_2O$ for the remainder of the day until used for patch clamp recordings. Before utilization, the low-Calcium ACSF was equilibrated with 95% O_2, 5% CO_2, and its pH was adjusted to 7.3–7.4 using NaOH. Osmolality was verified to be between 295–305 mOsm kg^{-1}. The slices were then divided into two groups for further experiments. The first group was used directly for acute whole-cell patch-clamp. The remaining brain slices were cultured and infected with fast viral vectors. Slices for interface culture were placed onto membrane inserts (Millipore) in 6 well plates with 800μL per well of slice culture media of the composition: 8.4 g/L MEM Eagle medium, 20% heat-inactivated horse serum, 30 mM HEPES, 13 mM D-glucose, 15 mM $NaHCO_3$, 1 mM ascorbic acid, 2 mM $MgSO_4 \cdot 7H_2O$, 1 mM $CaCl_2.4H_2O$, 0.5 mM GlutaMAX-I and 1 mg/L insulin, 25 U/mL Penicillin/Streptomycin. The slice culture medium was carefully adjusted to pH 7.2–7.3 by adding 1 M Tris base and osmolality of 295–305 mOsmoles kg^{-1} by addition of pure H_2O, and sterile-filtered. Culture plates were placed in a humidified 5% CO_2 incubator at 35°C, and the slice culture medium was replaced every one to two days until endpoint analysis. One to three hours after brain slices were plated on cell culture inserts, brain slices were transduced by direct application of concentrated AAV viral particles over the slice surface for fluorescent labelling.

AAV-viral transduction

The adeno-associated virus (AAV) vectors used in this study were acquired from the University of Zurich Viral Vector Facility. We used an optimized hGFAP-AAV vector virus (ssAAV5-hGFAP-hHEbl/E-EGFP-bGHp(A)) to label astrocyte or glioblastoma cells with fluorescent eGFP. For labeling neurons in human brain slices, we utilized hsyn-AAV vector virus

(ssAAV1-hSyn-EGFP-WPRE-hGHp(A)) containing the eGFP marker. The functional virus

titers ranged between $1–3 \times 10^{13}$ units per mL. To transduce human slice cultures with AAV

amplicons, we directly applied the concentrated viral stock solutions to the slice surface using

a fine micropipette, at least one hour after the initial plating to allow the slice to equilibrate.

The virus was carefully spread over the entire slice surface, ensuring that the surface tension

of the viral solution was maintained without allowing it to spread onto the surrounding

membrane insert, to the best possible extent.

Patch clamp recording

Human GBM-infiltrated brain slices were individually transferred to the recording chamber

mounted on a SliceScope microscope (Scientfica, United Kingdom). The slices were perfused

with carbogenated recording ACSF at a temperature of 32–35 °C, which composed of 130 mM

NaCl, 3.5 mM KCl, 1 mM NaH_2PO_4, 24 mM $NaHCO_3$, 10 mM glucose, 1.3 mM $CaCl_2 \cdot 4H_2O$

and 0.9 mM $MgSO_4 \cdot 7H_2O$. A SciCam Pro camera (Scientifica, United Kingdom) was used for

visualization and image capture. For whole-cell recordings, borosilicate glass pipettes (Harvard

glass capillaries, GC120F-10) were pulled on a horizontal pipette puller (DMZ universal

electrode puller, Zeitz, Germany) to a resistance of 5–7 MΩ. The pipettes were filled with an

internal solution comprising 126 mM K-gluconate, 10 mM HEPES, 4 mM Mg-ATP, 0.3 mM

Na_2-GTP, 4 mM KCl and 10 mM phosphocreatine. The pH was adjusted to 7.3–7.4 with

concentrated KOH and the osmolality was adjusted to 290 mOsmoles/Kg using sucrose. To

assist with visualization of cell filling, the internal solution was supplemented with 8 mM

biocytin and 50 μM Alexa 594 dye. The pipette was mounted on a Multiclamp 700B amplifier

headstage (Molecular Devices, United States) fixed to a micromanipulator (PatchStar,

Scientifica). Electrophysiology signals were recorded using a Digidata 1550B low-noise data

acquisition system (Molecular Devices, United States). The data were low-pass filtered at 2

kHz and digitized at 20 kHz. The theoretical liquid junction potential was calculated at −11

mV but was not corrected. Once a whole-cell recording was established, pipette capacitance

was neutralized, and bridge balance was applied. After forming a stable seal and breaking in,

the resting membrane potential was recorded within the first minute. The holding potential in

the voltage clamp mode was set at -60 mV. For inclusion in the analysis, cells were required to

have a > 1 GΩ seal recorded before break-in and a series n initial access resistance < 20 MΩ.

After the patch-clamp experiment, the X-, Y-, and Z-positions of all patched cells, as well as

the margin of slices, were recorded in a coordinate file. This coordinate data was subsequently

utilized to discern the identity of the patched cells in the immunohistochemistry experiment.

Electrophysiological data analysis

Electrophysiological data were analyzed using Clampfit 10.7 software (Molecular Devices, United States). The resting membrane potential was determined by averaging the voltage values recorded for 1 minute under current clamp mode (I = 0). Input resistance (R_{in}) was calculated from the linear portion of the current−voltage relationship, generated in response to a series of 1-second current injections (ranging from -100 to +50 pA in 10 or 20 pA steps). The maximum and steady-state voltage deflections were used to determine the maximum and steady-state R_{in}, respectively.

For the differentiation of APs and aAPs, individual potentials were elicited by applying 3 ms direct current injections with increasing amplitudes, which was selected for analysis of AP/aAPkinetics. The initiation of APs/aAPs was defined as the voltage at the time corresponding to when the first derivative of the voltage response first exceeded 20 mV/ms [43]. Five distinct parameters were computed for each AP/aAP: Threshold potential (mV), amplitude (mV), half-width, maximum rise slope (mV/ms), and maximum decay slope (mV/ms) (Supplementary Fig. 3a). The amplitude of APs/aAP was measured from the threshold to the peak voltage. The half-width of AP/aAP was determined as the width at half-maximal amplitude. To determine the rise and decay slopes of APs/aAPs, measure the duration for which the membrane potential ascends from 10% to 90% of the peak amplitude and descends from 90% to 10% of the peak amplitude. Subsequently, divide the amplitude by the corresponding time intervals.

To ascertain the optimal parameter combinations that could effectively classify AP and aAP signatures, a hierarchical clustering analysis was performed by using squared Euclidean distances [43] . Five AP parameters were tested by the D'Agostino–Pearson normality test. Parameters with a non-Gaussian distribution were used for subsequent unsupervised clustering. Before clustering, we first normalized variables into the range (0, 1) by performing min–max normalization. Hierarchical clustering arranges data items into a tree-like dendrogram in which tree leaves represent data items, and the length, or the computed Euclidean distance, of tree edges between leaves represents the dissimilarity between data items. The hierarchical clustering analysis was performed with the Free Statistics Software (Office for Research Development and Education, version 1.1.23-r7, http://www.wessa.net/).

The electrophysiological data obtained from acute and cultured brain slice experiments, evaluating features over time post-slicing, were analyzed using linear regression and Pearson's correlation. The sample size, p-values, and R^2 values were reported.

To confirm the components of aAPs, Tetraethylammonium chloride (TEA, Sigma-Aldrich,

30mM) was first added to the superfusing ACSF, followed by the subsequent addition of

tetrodotoxin citrate (TTX, TOCRIS, 1 µM) to the ACSF. The amplitude, half-width, and rise

and decay slopes of AP/aAP from each cell were compared between the control, TEA, and

TTX+TEA conditions.

Immunocytochemistry

All the virus transduced slices or patched slices were fixed in 4% paraformaldehyde in

phosphate buffered saline (PBS) at 4 °C overnight and then transferred to 0.1 M PBS. The fixed

slices were then transferred to 0.1 M PBS and washed thoroughly three times to remove any

remaining fixative. Next, the slices were incubated in 0.1 M PBS containing 0.5% Triton-X-

100 (0.5% PBST) for 30 minutes at room temperature. To block nonspecific binding sites, the

slices were incubated in 0.5% PBST containing 10% bovine serum albumin (BSA) (Sigma-

Aldrich, A4503) for 5-6 hours at room temperature. Following the blocking step, the slices

were incubated at 4 °C for 48 hours in 0.5% PBST containing 5% BSA and one or more primary

antibodies. After the 48-hour incubation with primary antibodies, the slices were washed three

times for 10 minutes each with 0.1 M PBS. Subsequently, they were incubated for 2 hours at

4 °C with the appropriate secondary antibodies. After incubation with appropriate secondary

antibodies, slices were washed for 10 min three times with PBS, followed by DAPI nuclear

staining for 15-20 min at room temperature. After washing the slices for 10 min three times

with PBS with. Slices were treated with 0.1% Sudan Black B (Sigma-Aldrich, Saint Louis,

MO) in 70% ethanol to quench lipofuscin autofluorescence. The slices were then dried on glass

microscope slides and mounted with anti-fade solution (Sigma-Aldrich, F4680). Slides were

stored at room temperature in the dark prior to imaging. Confocal z-stack images were acquired

on a spinning disk confocal microscope (Olympus IX83) equipped with 504 nm, 488 nm, 594

nm, and 647 nm excitation laser lines. Images were acquired using a ×20 objective with a

numerical aperture (NA) of 0.9.

Morphology reconstruction

The morphology of the cells was reconstructed from a stack of 116–196 images per cell for

hSyn+ IN cells using a 488 nm excitation laser for eGFP fluorescence signal. GFAP+ IN cells

were reconstructed from a stack of 24–43 images per cell using a 594 nm excitation laser for

the Alexa-594 loading dye. Image stacks belonging to one cell were imported into the

Neuromantic 1.7.5 software [44] for 3D reconstruction.

Antibodies

The primary antibodies: Mouse anti-NeuN (1:500 dilution, Chemicon, MAB377); Rabbit anti-

GFAP (1:500 dilution, Abcam, ab68428). Goat anti-Olig2 (1:500 dilution, Biotchne, AF2418).

The secondary antibodies: CY^{TM}5 (Donkey anti Mouse, 1:400, Jacksonimmuno, AB2340820);

CY^{TM}5 (Donkey anti Rabbit, 1:400 dilution, Jacksonimmuno, AB2340607); Alexa 647

(Donkey anti Goat, 1:400 dilution, Invitrogen, A-21447); Alexa Fluor™ 594 (Donkey anti

Rabbit 1:400 dilution, Invitrogen, A-21207); Alexa Fluor™ 594 (Goat anti-Mouse, 1:400,

Jacksonimmuno, AB2338871); Alexa Fluor™ 594 (conjugated streptavidin binding for

biocytin, 1:400, Thermofisher, S11227).

Statistical Analysis

Sample sizes and statistical tests are provided in the figure legends. Statistical analysis was

performed using GraphPad Prism 8 (GraphPad Software, United States). All data are presented

as mean ± SEM. When more than two groups were compared, a One-way ANOVA was used,

followed by Tukey's multiple comparison test. Comparative analysis among paired data from

three groups was performed using a Two-way ANOVA followed by Tukey's multiple

comparison test. To evaluate correlations for each electrophysiological parameter, using linear

regression and Pearson's correlation is applied to the data. R^2 and P values are calculated for

each fitting to indicate correlation strength and significance. The independent t-test was used

to assess differences among the five parameters of APs and aAPs. Data are presented as the

mean ± SEM. Levels of statistical significance are indicated as follows: *p<0.05, **p<0.01,

***p<0.001, and ****p<0.0001.

Acknowledgements

We thank K. Widzsiz for technical assistance and L.M. Fitting for logistical and technical

assistance; H. Thygesen, A. Kobberø, and A.S. Møller-Andersen for administrative assistance;

the OR staff at the Department of Neurosurgery, Aarhus University Hospital, for logistical

assistance; E.M.R. Jensen for assisting patient enrolment.

Author contributions

A.R.K. and M.C. conceived and conceptualized the project. T.T., A.O., W-H.H., MC, and
A.R.K designed the experiments. T.T., A.O, and W-H.H. conducted and analyzed the
electrophysiology experiments. J.T.E. contributed to immunofluorescence and morphology
experiments and manuscript editing. S.O.S.C, A.K.R.S., and A.R.K. contributed with patient
inclusion and performed surgical procedures. K.M. provided data structure and management
of patient data and N.M. contributed with entry of data and meta data. B.W.K. supervised
immunostaining and histopathology experiments and contributed to manuscript editing. T.T.,
W-H.H., K.J.E., and A.R.K. wrote the manuscript draft and all authors contributed critical
editing.   A.R.K. contributed funding and was responsible for project administration and
management. A.R.K. and W-H.H. supervised all aspects of the work.

Competing interests declarations
None.

Fig. 1

Fig. 2

Acute brain slices

Cultured brain slices

Fig. 4

Fig. 5

a

b