The Cancer Testis Antigen Testis Specific Serine Kinase 6 (TSSK6) is abnormally expressed in colorectal cancer and is essential for oncogenic behaviors

Magdalena Delgado[1], Steve Stippec[1], Kathleen McGlynn[1], Melanie H. Cobb[1], Angelique W. Whitehurst[1]*

Department of Pharmacology, UT Southwestern Medical Center, Dallas, TX 75390, USA.

*Correspondence: Angelique.Whitehurst@UTSouthwestern.edu

Running Title: TSSK6 is an oncogenic kinase

Keywords: TSSK6, focal adhesion, colorectal cancer, tensin, paxillin, Cancer testis antigen

The authors declare no potential conflicts of interest.

Abstract

Cancer testis antigens (CTAs) are a collection of proteins whose expression is normally restricted to the testis, but abnormally activated in a wide variety of tumors. The CTA, Testis specific serine kinase 6 (TSSK6), is essential for male fertility in mice. Functional relevance of TSSK6 to cancer, if any, has not previously been investigated. Here we find that TSSK6 is frequently anomalously expressed in colorectal cancer and patients with elevated TSSK6 expression have reduced regression free survival. Depletion of TSSK6 from colorectal cancer cells attenuates anchorage independent growth, invasion and growth in vivo. Conversely, overexpression of TSSK6 promotes neoplastic behaviors in vitro and enhances in vivo tumor growth. Notably, ectopic expression of TSSK6 in non-transformed human colonic epithelial cells is sufficient to confer anchorage independence and enhance invasion. In somatic cells, TSSK6 co-localizes with and enhances the formation of paxillin and tensin positive foci at the cell periphery, suggesting a function in focal adhesion formation. Importantly, TSSK6 kinase activity is essential to induce these tumorigenic behaviors. Our findings establish TSSK6 as a previously unknown oncogenic kinase. The requirement of TSSK6 for tumor cell growth presents it as an intervention target for therapy, which could exhibit an exceptionally broad therapeutic window.

Introduction

Cancer testis antigens (CTAs) are a set of > 200 proteins whose expression is normally confined to the testis or placenta, but frequently anomalously activated in cancer(1, 2). As the testis is an immune-privileged site, expression of these proteins in somatic cells can generate immunogenic antigens recognized by T-cells(1). Indeed T-cell mediated therapies targeting CTAs have exhibited clinical efficacy(3, 4). In addition to immunogenic properties, recent studies suggest CTAs can be functionally integrated into the tumor cell regulatory environment. Specifically, individual CTAs have been implicated in degrading tumor suppressors, enhancing oxidative phosphorylation, regulating transcriptional networks, influencing genomic integrity, and altering immune recognition(5-14). Thus, an emerging hypothesis is that CTAs are not innocent bystanders during tumorigenesis, but are intimately engaged in promoting neoplastic behaviors.

The CTA family includes Testis Specific Serine Kinase 6 (TSSK6)(15). TSSK6 is a member of a 5-protein kinase family found on the CAMK branch of the kinome (16). Deletion of TSSK6 in mice leads to male, but not female, infertility(17). As expected given TSSK6's testes-restricted expression pattern, these mice exhibit no defects in development or adult tissues and female mice are unaffected(17). In sperm, TSSK6 is expressed post-meiotically and localizes to the nucleus in spermatids and the posterior head in spermatocytes. TSSK6-null sperm exhibit defective histone-to-protamine transition (required for post-meiotic chromatin remodeling), as well as a decrease in actin polymerization(18, 19). The results are striking morphological defects in sperm including nuclear deformities, a hairpin shape with heads facing backwards, or completely detached heads(19).

TSSK6 was initially classified as a cancer-testes antigen based on its restricted mRNA expression to the testes and detection in lung cancer, sarcoma and lymphoma(15).

A recent study has indicated that TSSK6 mRNA expression correlates with T-cell diversity(20). However, TSSK6 protein expression in cancer and any functional relevance to tumorigenesis have not been investigated. If TSSK6 is active when aberrantly expressed in somatic cells it could theoretically induce a tumor-specific signaling pathway that may have dramatic consequences on cellular behaviors.

Here we demonstrate that TSSK6 is frequently abnormally expressed in colorectal cancer, where it correlates with reduced regression free survival. We find that TSSK6 is necessary and sufficient for tumor cell viability, anchorage independence, invasion and growth in vivo. Notably, expression of TSSK6 in non-transformed cells is sufficient to confer anchorage independence, suggesting TSSK6 is a bona-fide oncogenic kinase. Importantly, TSSK6 exhibits kinase activity when ectopically expressed and this activity is crucial for inducing oncogenic behaviors. Mechanistically, we find that TSSK6 co-localizes with cytoskeletal regulatory proteins at focal adhesions and enhances paxillin and tensin-positive foci formation. Our findings establish TSSK6 as a novel oncogenic protein, which when abnormally expressed in somatic cells alters signaling networks to promote proliferation, invasion and tumor growth in vivo.

Results

TSSK6 is frequently abnormally expressed in colorectal cancer. TSSK6 mRNA expression is normally restricted to the testes, but its expression in tumors has not been investigated (Supplemental Figure 1A). Querying the PanCancer Atlas indicated that TSSK6 expression is elevated in ~6% of cancers (z-score > 2.0) and expressed in a wide range of tumors types (Supplemental Figure 1B)(21). We then asked whether TSSK6 mRNA expression correlated with patient outcomes in breast, lung, ovarian, epigastric, pancreatic and colon cancer (Supplemental Figure 1C). Here, we observed a correlation between elevated TSSK6 and reduced regression free survival time exclusively in colon cancer

(P=0.0092; log rank test; Hazard Ratio (HR), 1.35; 95% confidence interval (CI), 1.08-1.69. (Supplemental Figure 1C and Figure 1A). To determine the extent of TSSK6 protein accumulation in human colorectal cancer tissue, we developed an immunohistochemical staining protocol in human testis (Figure 1B). Analysis of 20 IHC specimens revealed that ~65% of CRC tumor cores stained positive for TSSK6 protein with a range of expression levels (Figure 1B). We examined endogenous expression of TSSK6 in human tumor derived CRC cell lines, HCT116 and RKO as compared to immortalized mouse spermatagonial cell lines (GC-1, GC-2). In the spermatagonial-derived cell lines, TSSK6 accumulated at two different sizes. We observed that HCT116, but not RKO cells, expressed TSSK6 corresponding to the heavier band in spematagonia (Figure 1C). This band was depleted upon transfection of HCT116 with siRNA's targeting the TSSK6 coding sequence (Figure 1D). We expanded the panel to five CRC cell lines and found that HCT116, HT-29 and LOVO CRC cell lines express endogenous TSSK6, while it is not detectable in RKO and DLD-1 cell lines (Figure 1E).

TSSK6 is essential for CRC oncogenicity. We next directly tested whether TSSK6 is necessary for CRC tumorigenic behaviors. In viability assays, we found that TSSK6 depletion reduced survival of HCT116 cells, but had little impact on HT-29 and LOVO cells. We next asked whether TSSK6 was required for anchorage independent growth, a hallmark of transformation. Here, we observed a decrease in growth in HCT116 as expected. In addition, HT-29 and LOVO cells exhibited a reduction in soft agar growth following TSSK6 depletion; no effect was observed in TSSK6-negative cell lines (Figure 2B). To control for off-target effects of siRNA, we also verified that a second, independent siRNA pool recapitulated a defect in cell growth (Figure 2C and Figure 1D). We next measured TSSK6's contribution to invasion using a transwell assay. We used LOVO cells, which, unlike HCT116, do not exhibit a loss of viability following TSSK6 depletion in 2-D culture (Figure 2A). LOVO cells were

depleted of TSSK6 for 48 hours and then measured for invasion over a 48-hour period. In this setting, depletion of TSSK6 was sufficient to dramatically reduce invasive capacity of CRC cells (Figure 2D).

To determine if ectopic TSSK6 expression can enhance oncogenic phenotypes, we stably expressed TSSK6 in DLD-1 cells, CRC cells that do not express detectable endogenous TSSK6 (Figure 1E & 2E). The DLD-1 TSSK6-V5 cell line did not exhibit a significant increase in cell division in 2-dimenionsal culture as compared to control (Figure 2E). However, we did observe an increase in soft agar growth and invasion in TSSK6-V5 expressing DLD-1 cells (Figure 2F & 2G). Together this analysis indicates that TSSK6 is both necessary and sufficient to confer tumorigenic phenotypes in colorectal cancer cells.

TSSK6 exhibits bonafide oncogene activity dependent on its kinase activity. We next asked whether ectopic expression of TSSK6 in a non-transformed setting can confer oncogenic behaviors. Here, we used normal human colonic epithelial cells that are immortalized with hTERT and CDK4. These cells also express oncogenic K-RAS (V12) and lack p53 (HCEC)(22, 23). Stable expression of TSSK6-V5 led to the accumulation of TSSK6 protein at both the 35 and 30 kDa molecular weights in this setting (Figure 3A). We did not observe an appreciable increase in cell viability following TSSK6-V5 expression in 2-dimensional culture (Figure 3B). HCEC cells do not form appreciable colonies in soft agar, indicating they are not capable of anchorage independent growth. However, HCEC cells expressing TSSK6-V5 exhibited robust anchorage independent growth in (Figure 3C), suggesting that TSSK6 can induce transformation in pre-malignant cells.

TSSK6 is reported to phosphorylate histones during the histone to protamine transition in developing sperm(16). In cells, TSSK6 requires HSP90 for kinase activity. In addition, TSSK6 associates with TSSK6 Activating Co-Chaperone (TSACC, SIP), a tetratricopeptide repeat domain containing protein. TSSK6 recruits TSACC to HSP90 and the formation of this

complex leads to maximal TSSK6 activity in 293 cells(21). While HSP90 is widely expressed, TSACC is exclusively expressed in elongated spermatids and not detected in any other normal tissues(21). Thus, we asked whether TSSK6 is an active kinase when ectopically expressed in HCEC cells, outside of its native environment. We performed kinase assays on immunoprecipitated (IP) TSSK6-V5 from HCEC cells using a generic myelin basic protein substrate. TSSK6 immunoprecipitates exhibited ~70 fold activity kinase activity over background (Figure 3D). Mutation of the catalytic lysine (K41M) or the activation loop threonine (T170A), both dramatically reduced kinase activity (Figure 3D) (21). We subsequently asked whether TSSK6 kinase activity is required for its oncogenic activity. Indeed, HCEC cells expressing wild-type, but not K41M or T170A, TSSK6 exhibited enhanced soft agar growth (Figure 3E). In invasion assays, expression of wild-type TSSK6 but not the activation loop mutant, T170A, dramatically increased invasive capacity of HCEC cells (Figure 3F). Together, these findings indicate that when abnormally expressed in pre-malignant cells, TSSK6 is sufficient to confer and enhance transformation phenotypes.

TSSK6 is associated with and promotes focal adhesion formation.

TSSK6 is reported to localize to the nucleus and head and neck region of sperm where it regulates histone phosphorylation and actin polymerization, respectively(17-19). Thus, we asked where TSSK6 localizes when it is inappropriately expressed in cancer cells. In HCT116 cells, we observed endogenous TSSK6 localization throughout the cell, with an enrichment at the nuclear and cellular membrane (Figure 4A). In HCEC cells, ectopically expressed TSSK6-V5 also localized throughout the cell and at the cell membrane (Figure 4B). We noted focal enrichment of TSSK6-V5 at the cell membrane. These foci stained positive for paxillin, focal adhesion protein (Figure 4B). Moreover, the number of paxillin positive foci were significantly increased in HCEC-TSSK6-V5 cells as compared to HCEC-HcRed cells (Figure 4C). We observed a similar trend for localization and enrichment with

tensin, another focal adhesion protein (Figure 4D&E). This finding suggests that TSSK6 promotes focal adhesion formation, potentially through local interactions with focal adhesion proteins and regulates cell-ECM attachments that may influence motility.

TSSK6 is necessary and sufficient for tumor growth in vivo. Given the robust in vitro effects observed upon TSSK6 expression, we next examined the impact of TSSK6 on tumor growth in vivo. We established xenograft tumors expressing shCTRL and shTSSK6 in HCT116 cells. We measured tumor growth regularly over 30 days and observed a significant decrease in tumor growth in the mice injected with shTSSK6 cells (Figure 5A). Conversely, we established tumors using DLD-1 cells stably expressing TSSK6-wild type or the TSSK6-T170A mutant. Here, we observed enhanced tumor growth upon of TSSK6-V5 expressing cells as compared to control cells. Notably, TSSK6-170A cells exhibited a nearly identical growth rate to that of control cells, indicating that TSSK6 must be fully active to enhance in vivo growth(Figure 5B).

Discussion

Abnormal activation of kinases is a frequent feature of cancers. Chronic activation of kinases including PIK3CA, EGFR, ALK and RAF are well established to drive tumor progression, survival and metastases. In turn, these kinases present targetable vulnerabilities that when inhibited attenuate tumor growth(24). Our findings here present TSSK6 as a previously unrecognized oncogenic kinase. TSSK6 expression is normally restricted to the testes, where it is required for male fertility. Our findings suggest that TSSK6 is frequently aberrantly expressed in colorectal tumors, where it enhances tumorigenic behaviors including anchorage independence, invasion and growth in vivo. Based on our findings, we hypothesize that these behaviors are due to TSSK6-induced changes in cell-ECM adhesion properties.

The strong phenotypes induced by TSSK6 expression present a number of implications for future studies. First, it is unknown how TSSK6's expression and kinase activity are induced when it is abnormally expressed in cancer cells. With respect to expression, CTA promoters are typically highly methylated in somatic cells and DNA methylation can be sufficient to induce CTA expression(25). Whether DNA methylation of TSSK6 is a mechanism for regulating its expression as well as which transcription factors directly activate expression upon demethylation remain unknown. Similarly, mechanisms regulating TSSK6 kinase activity are poorly understood, even in the testes. Studies to-date indicate that HSP70/90 is essential for TSSK6 activation(17, 26). Notably, TSSK6 is inhibited following exposure to the HSP inhibitor 17-allylamino-17-demethoxygeldanamycin (17-AAG)(26). Thus, TSSK6-positive tumors may exhibit enhanced sensitivity to HSP90 inhibitors, which are currently under clinical evaluation in a variety of tumor types (27). TSSK6 activity in sperm is also modulated by the sperm-specific chaperone, TSACC(26). mRNA expression of TSACC is also detectable in tumors and in a subset that overlaps with TSSK6 in some tumor types (not shown), suggesting that both proteins may be working in concert in a subset of tumors. Finally, mechanisms that negatively regulate TSSK6 activity in sperm or somatic cells, including phosphatases or degradation pathways have not been explored. If sperm-specific mechanism exists, but are absent in tumors, TSSK6 activity could be relatively unopposed when it is abnormally expressed in somatic cells.

Second, it will be essential to identify TSSK6 substrates and downstream signaling events in cancer cells. A pool of TSSK6 appears to localize to and increase the formation of focal adhesions. Thus, TSSK6 may be regulating cell adhesion through local signaling events at these sites. Studies in sperm and flies, suggest that TSSK6 can regulate cytoskeletal proteins including actin and microtubules, respectively(19, 28). How TSSK6 signaling influences the cytoskeleton and impact cells shape, adhesion and motility are important next

questions. The localization of TSSK6 in the cytoplasm, nucleus and at the nuclear periphery also suggest that it shuttles between the cytoplasm and nucleus. TSSK6's function, if any, in these compartments may also be contributing to its impact on growth, survival and motility of tumor cells. The localization of TSSK6 throughout the cell along with the substantial behavioral changes upon it expression, suggest it could be inducing a multitude of signaling changes that may permit tumor cell adaptation to diverse environments.

Third, our study implicates TSSK6 as a target for anti-cancer therapy. Previous studies have correlated TSSK6 expression with CD8+ T-cell infiltration in multiple tumor types(20). This finding suggests that TSSK6 could generate antigenic peptides in tumors. A number of peptides have been identified with algorithmic tools, but these have not yet been tested for antigenicity(20). Our studies extend the therapeutic potential of TSSK6 and implicate it as a direct interventional target. While TSSK6-specific inhibitors have not been reported, inhibitors have been reported for TSSK2, suggesting this family of kinases is therapeutically tractable (29). If inhibitable with high specificity, TSSK6 could represent an anti-tumor target with an extraordinarily broad therapeutic window.

Experimental Procedures

Cell Lines and Culture conditions

GC-1spg and GC-2spd(ts) cells were obtained from the American Type Culture Collection (ATCC). HCT116 cells were obtained from Cyrus Vaziri (UNC). RKO, HT29, DLD-1 and LOVO cell lines were obtained from Jerry Shay (UT Southwestern). Colorectal cancer cells lines were maintained in high glucose DMEM supplemented with 10% FBS (Cat# F0926, Sigma-Aldrich) 1% antibiotic/antimycotic (Cat# 400-101, GeminiBio) and 2.5 mM HEPES pH7.4 at 37°C in a humidified 5% CO_2 atmosphere. HCEC cells (immortalized with hTERT and CDK4, expressing RASV12 and lacking p53 (HCEC1CTRP)) were developed by Jerry Shay (UT Southwestern) (22, 23). HCECs were maintained in high glucose DMEM

supplemented with 5 nM sodium selenite (Cat# S5261, Sigma-Aldrich), 2 ug/mL apo-transferrin (Cat# T1147, Sigma-Aldrich), 20 ng/mL epidermal growth factor (Cat# 236-EG, R&D), 10 ug/mL insulin (Cat# 12585-014, Gibco) , 1 ug/mL hydrocortisone (Cat# H0888, Sigma-Alrich), 2% FBS and 1% antibiotic/antimycotic at 37°C in a humidified 5% CO_2 atmosphere. HEK293T cells were obtained from ATCC and cultured in high glucose DMEM supplemented with 10% FBS, 1% anitbiotic/antimycotic and 2.5 mM HEPES pH7.4 at 37°C in a humidified 5% CO_2 atmosphere. All cells were periodically evaluated for mycoplasma contamination by a mycoplasma PCR detection kit (Cat# G238, ABM). Authenticity of cell lines was evaluated by detection of ten genetic loci using the GenePrint® 10 System (Promega) and cross referencing to ATCC or internal genetic profiles.

Antibodies

Antibodies used: TSSK6 (sc-514076, Santa Cruz) (1:100 IHC, 1:500 IB, 1:200 IF), Vinculin (sc-73614, Santa Cruz) (1:2000), Beta-Tubulin (2128, CST) (1:5000), Cofilin (3312, CST) (1:1000), V5-tag (13202, CST) (1:5000 IB, 1:500 IF), V5-tag (46-0705, Invitrogen)(1.12 ug antibody/500 ug protein IP, 1:2000 IB, 1:100 IF), Paxillin (sc-365379, Santa Cruz) (1:275), Tensin (11990, CST) (1:100).

Transfections

For siRNA transfections, cells were trypsinized and seeded in Opti-MEM containing Lipofectamine™ RNAiMAX (Thermo Fisher Scientific, Waltham, MA) complexed with siRNAs at a final concentration of 50 nM for all cells except LOVOs, which were transfected with a final concentration of 100 nM. siRNAs for were purchased from Sigma as follows: non-targeting controls (VC30002), targeting TSSK6 (pool #1: SASI_Hs01_00190487, SASI_Hs01_00190489) or (pool #2WD07275781/82, WD07275783/84).

Plasmids

cDNAs encoding human *TSSK6* were obtained in pDONR223 (DNASU) and cloned into pLX302 or pLX304 (Addgene Plasmid #25896, #25890) using the Gateway Cloning system (Thermo Fisher Scientific). For control transfections, Hc-Red-V5 in pLX302/pLX304 was used. TSSK6 kinase mutations were made using the QuikChange XL Site-Directed Mutagenesis Kit (200516, Agilent). pLX302-TSSK6-V5 was used as a template. Primers for the K41M mutation were as follows: FWD 5'-CCGTGGCCATCATGGTGGTGGACCG-3' and REV 5'-CCGTGGCCATCATGGTGGTGGACCG. Primers for the T170A mutation were as follows: FWD 5'-GACCTGAGCACCGCCTACTGCGGCTC-3' and REV 5'-GAGCCGCAGTAGGCGGTGCTCAGGTC-3'. psPAX2 and pMD2.G lentiviral constructs were purchased from Addgene (Plasmid #12260, #12259). For shRNA experiments, pLKO.1 vectors from TRC expressing TSSK6-targeted shRNAs (TRCN0000199044; sequence: 5'-CGTCGTGCTCTACGTCATGGT-3', TRCN0000356035 5'- TCGTGCTCTACGTCATGGTCA-3') were used as shRNA pool. Nontargeting shRNA in pLKO.1 (shSCR) was used as a control (Addgene Plasmid #17920).

Immunohistochemistry Analysis

Non-malignant testis, lung and colorectal adenocarcinoma tissues were obtained with informed consent from the UTSW Tissue Management Shared Resource (TMSR) in compliance with the UTSW Internal Review Board committee. TSSK6 IHC was optimized and performed by the TMSR according to internal protocols using the Dako Autostainer Link 48 system. Briefly, the slides were baked for 20 minutes at 60°C, then deparaffinized and hydrated before the antigen retrieval step. Heat-induced antigen retrieval was performed at low pH for 20 minutes in a Dako PT Link. The tissue was incubated with a peroxidase block and then an antibody incubated (TSSK6, 1:100, sc-514076, Santa Cruz Biotechnology) for 35 minutes. The staining was visualized using the Keyence BZ-X710 fluorescence microscope.

Staining scores of TSSK6 ranging from 0-3 were set based on the positive scores quantitated by using IHC Profiler of Image J: score 0 (<30), score 1 (30-45), score 2 (45-60), and score 3 (>60)(30).

Lentiviral Transduction

Stable cell lines were generated through lentiviral-mediated transduction. HEK293T cells were co-transfected with the target gene vectors and lentiviral packaging plasmids. 48 hours later, virus-conditioned medium was harvested, passed through 0.45 μm pore-size filters, and then used to infect target cells in the presence of Sequabrene™ (S2667, MilliporeSigma) for at least six hours. After infection, medium was replaced, and cells were allowed to recover. DLD-1 cells were infected with pLX304-HcRed-V5 or pLX304-TSSK6-V5 and selected with blasticidin for 3 days or were infected with pLX302-HcRed, pLX302-TSSK6-V5, pLX302-TSSK6-T170A-V5 and selected with puromycin for 3 days. HCEC1CTRP cells were infected with pLX302-HcRed, pLX302-TSSK6-V5, pLX302-TSSK6-T170A, or pLX302-TSSK6-K41M and selected with puromycin for three days. HCT116 cells were infected with pLKO.1 shSCR or shTSSK6 for 120 hours used immediately for xenograft mouse experiments. Immunoblotting for TSSK6 and/or V5 was used to check for overexpression or depletion.

Xenograft Assays

All animal experiments were conducted with IACUC approval. 6-10 week old male NOD.cg-PRKDC[SCID]Il2rg[tm1Wjl]/SzJ (NSG) (Rolf Brekken, UT Southwestern) mice were subcutaneously injected in the flank with 1 million cells (HCT116) or 800,000 cells (DLD-1) in 100 μl PBS. Once tumors were visible, tumor volume was measured by caliper in two dimensions, and volumes were estimated using the equation $V = \text{length} \times \text{width}^2/2$. Caliper measurements were made twice a week. At the end of the experiment, tumors were harvested and weighed for final tumor mass.

Immunofluorescence Assays

Cells plated on glass coverslips were washed twice with PBS, fixed with 3.7% formaldehyde for 10 minutes at room temperature. Cells were then washed twice with PBS and incubated in 0.25% Triton-100 for 10 minutes prior to washing three times with 1X PBS. Next coverslips were blocked for at least 30 minutes at room temperature or overnight at 4°C in blocking buffer (1% BSA in 0.001%TBST mixed with PBS). Primary antibodies were diluted with blocking buffer and incubated with samples for 1 hour at room temperature. Next, samples were washed three times in blocking buffer for 5 minutes each, followed by incubation with Alexa Fluor-conjugated secondary antibodies (Thermo Fisher Scientific) in blocking buffer at a dilution of 1:1000 for one hour at room temperature. Lastly, samples were washed three times for five minutes with 0.5% BSA in water and 1 time with water, followed by mounting onto glass slides using ProLong™ Gold Antifade reagent with DAPI. Images were acquired by confocal microscope Zeiss LSM700. Staining was quantitated using Zen 3.8 software (Zeiss).

Cell Lysis and Immunoblotting (IB)

Samples were lysed in preheated (100°C for five minutes) 2X Laemmli sample buffer with Beta-mercaptoethanol and boiled for six minutes. Samples were resolved using SDS-PAGE (% for TSSK6 gels noted in legends) , and transferred to an Immobilon PVDF membrane (Millipore), blocked in 5% non-fat dry milk followed by incubation in the respective primary antibodies overnight. Following incubation, membranes were washed three times with Tris Buffered Saline (20 mM Tris, 150 mM NaCl, 0.1% Tween-20) (TBST), and incubated for one hour with horseradish peroxidase-coupled secondary antibodies (Jackson Immunoresearch). Subsequently, membranes were washed three times with TBST and then developed using SuperSignal™ West Pico PLUS chemiluminescence substrate (Thermo Fisher scientific, 45-000-875). Immunoblots were scanned using the EPSON perfection v700 photo scanner. For Figure 3D and Figure 5B, after incubation with primary antibodies, membranes were incubated for one hour with fluorophore-coupled secondary antibodies (Licor). Subsequently,

membranes were washed three times with TBST and then imaged using Licor Odyssey XF Imaging System.

Soft agar assays

Cells were suspended into 0.366% bacto-agar and then plated onto solidified 0.5% bacto-agar. Cells were seeded at a density of 500 cells (HCT116 and HCEC1CTRPs), 2000 cells (LOVO, HT29) or5000 cells (DLD-1 and RKO) per 12-well plate. LOVO, HT29 and HCT116 cells were transfected with siRNA for 48 hours and then collected for plating into soft agar. After 2-3 weeks, colonies were stained for 1 hour with 0.01% crystal violet in 20% methanol. Background stain was removed by washing 3x in water for 30 minutes and 1x overnight. Images were captured with a Leica S9D dissecting microscope, and quantitated by ImageJ software.

Invasion Assay

Corning BioCoat Growth Factor Reduced Matrigel Invasion Chambers with 8.0 uM PET membranes (Cat# 354483) were used for invasion assays according to the manufacture's guidelines for use. Briefly, inserts were allowed to come to room temperature for 30 minutes in a sterile hood. Next inserts were allowed to rehydrate in media at 37°C in a humidified 5% CO_2 atmosphere. After two hours, cell suspensions were prepared at 50,000 cells in 500 uL of 0.1% FBS DMEM per well insert. Media was carefully removed from the invasion inserts and placed into 24-well plates with 10% FBS DMEM. Cell suspensions were immediately added to the wells. Plated inserts were incubated for 24 hours (DLD-1) or 48 hours (HCEC, LOVO). For experiments in which siRNA were used, cells were transfected for 48 h and then collected for plating into transwell inserts.

Cell Titer Glo Assays (Viability)

Cell-Titer Glo (Promega) (CTG) was performed by manufacture's protocol but modified to use 15 µl of CTG for 150 µl of cells in media. HCT116, HT29, LOVO cells were reverse

transfected with siRNA as indicated and plated at 1000 cells per well and allowed to grow for 96 hours. DLD-1 cells were plated at 4000 cells per well and allowed to grow for 96 hours. HCECs were plated at 1000 cells per well and allowed to grow for 96 h. Luminescence was read using a CLARIOstar Plus plate reader (BMG Labtech).

IP-Kinase assays

Cells were lysed for 30 minutes on ice in non-denaturing lysis buffer (50 mM HEPES pH 7.4, 50 mM NaCl, 0.1 % Triton X-100, 100 mM NaF, 30 mM sodium pyrophosphate, 50 mM β-glycerophosphate 1 mM EGTA, 10% glycerol, 1 mM Na_3VO_4, 0.4 ug/mL pepstatin, 0.4 ug/mL leupeptin, 4 ug/mL TAME, 4 ug/mL tos-lys-chloromethylketone, 4 ug/mL Na-benzoyl-L-arginine methyl ester carbonate, 4 ug/mL soybean trypsin inhibitor). Lysates were clarified at 12,000 x g for 10 minutes and pre-cleared with Protein A/G agarose beads for one hour at 4˚C. Five percent of each lysate was set aside as input material and the remainder was immunoprecipitated with 1.12 ug of antibody for one hour at 4˚C. Protein A/G agarose beads were washed three times with lysis buffer and 35 ul of beads incubated with lysate for one hour. Beads were washed three times with 1 mL of 1M NaCl, 20 mM Tris pH 7.4 and one time with minimal kinase reaction buffer (10 mM HEPES pH 8.0, 10 mM MgCl2). The buffer was removed and beads were resuspended in 30 uL kinase reaction buffer with 10 mM HEPES pH 8.0, 10 mM MgCl2, 1 mM benzamidine, 1 mM DTT, 50 µm ATP (1 cpm/fmol) with 2 uL 5mg/mL MBP at 30 °C for 30 minutes. Reactions were stopped by adding 10 µl of GenScript 4X LDS sample buffer, followed by boiling for 3 minutes. Samples were run on GenScript 4-20% Sure Page gels. Gels were stained with coomassie blue and bands corresponding to MBP were excised and raw counts measured using a liquid scintillator. Kinase activation was calculated by subtracting blank MBP raw counts per millions from sample counts per million. All samples were then normalized to control immunoprecipitates.

Normal and tumor expression analysis

The Gtex Portal was used to analyze TSSK6 mRNA expression across normal tissues on 12/2023. The Genotype-Tissue Expression (GTEx) Project was supported by the Common Fund of the Office of the Director of the National Institutes of Health, and by NCI, NHGRI, NHLBI, NIDA, NIMH, and NINDS. cBIO Portal was used to analyze expression in tumors in the TCGA PanCancer Atlas Study (10,528 cases) using a z-score cutoff of 2 (21).

Survival Analysis

Survival analysis was performed using the KM Plotter program(31). Affymetrix 224409_s_t was used for survival analysis for TSSK6. Populations were split based on median expression and regression and overall survival was measured. Analysis was completed in December of 2023.

Statistical Analysis

Graphpad Prism (Graphpad Software) was used to perform statistical analyses. Normality of data was determined by the Shapiro-Wilk normality test. Statistical differences in all cases were determined by Mann-Whitney (not normal distribution) or t-test (normal distribution) as specified in the figure legends. : ns= not significant, $P > 0.05$; $*P < 0.05$; $**P < 0.01$; $***P < 0.001$; $****P < 0.0001$. Outliers were detected by ROUT analysis.

Author contributions

A.W.W., M.D. conceptualization; A.W.W., M.D., methodology; A.W.W., K.M., S. P., M.D., investigation; A.W.W. and M.D. writing—original draft, review & editing; A.W.W., supervision; A.W.W. funding acquisition.

Funding

Funding was provided by the National Institutes of Health to A.W.W. (R01CA251172 and R01CA196905, R03CA250021). M.G. was supported by the Cancer Prevention Research Institute of Texas (CPRIT) Training Grant (RP210041) and a T32 NCI Training Grant (CA124334-15). This work was also supported by The Welch Foundation to A.W.W (I-2087-

20210327). Support for use of core services was made possible through NCI Cancer Center Support Grant (5P30CA142543) to the Harold C. Simmons Cancer Center. The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health.

Conflict of Interest

The authors declare that they have no conflict of interest

Acknowledgements

The authors would like to thank Jerry Shay for the HCEC cells, and Michael Reese and Melanie Cobb for helpful discussions.

References

1. Simpson, A. J., Caballero, O. L., Jungbluth, A., Chen, Y. T., andOld, L. J. (2005) Cancer/testis antigens, gametogenesis and cancer Nat Rev Cancer **5**, 615-625 10.1038/nrc1669
2. Almeida, L. G., Sakabe, N. J., deOliveira, A. R., Silva, M. C., Mundstein, A. S., Cohen, T. *et al.* (2009) CTdatabase: a knowledge-base of high-throughput and curated data on cancer-testis antigens Nucleic Acids Res **37**, D816-819 10.1093/nar/gkn673
3. D'Angelo, S. P., Melchiori, L., Merchant, M. S., Bernstein, D., Glod, J., Kaplan, R. *et al.* (2018) Antitumor Activity Associated with Prolonged Persistence of Adoptively Transferred NY-ESO-1 (c259)T Cells in Synovial Sarcoma Cancer Discov **8**, 944-957 10.1158/2159-8290.CD-17-1417
4. Hunder, N. N., Wallen, H., Cao, J., Hendricks, D. W., Reilly, J. Z., Rodmyre, R. *et al.* (2008) Treatment of metastatic melanoma with autologous CD4+ T cells against NY-ESO-1 N Engl J Med **358**, 2698-2703 10.1056/NEJMoa0800251
5. Yang, X., andPotts, P. R. (2020) CSAG2 is a cancer-specific activator of SIRT1 EMBO Rep **21**, e50912 10.15252/embr.202050912
6. Florke Gee, R. R., Chen, H., Lee, A. K., Daly, C. A., Wilander, B. A., Fon Tacer, K. *et al.* (2020) Emerging roles of the MAGE protein family in stress response pathways J Biol Chem **295**, 16121-16155 10.1074/jbc.REV120.008029
7. Herrera, L. R., Johnson, R. A., McGlynn, K., Gibbs, Z. A., Davis, A. J., andWhitehurst, A. W. (2023) The cancer testes antigen, HORMAD1, limits genomic instability in cancer cells by protecting stalled replication forks J Biol Chem **299**, 105348 10.1016/j.jbc.2023.105348
8. Maine, E. A., Westcott, J. M., Prechtl, A. M., Dang, T. T., Whitehurst, A. W., andPearson, G. W. (2016) The cancer-testis antigens SPANX-A/C/D and CTAG2 promote breast cancer invasion Oncotarget **7**, 14708-14726 10.18632/oncotarget.7408
9. Maxfield, K. E., Taus, P. J., Corcoran, K., Wooten, J., Macion, J., Zhou, Y. *et al.* (2015) Comprehensive functional characterization of cancer-testis antigens defines obligate participation in multiple hallmarks of cancer Nat Commun **6**, 8840 10.1038/ncomms9840
10. Cheng, C. C., Wooten, J., Gibbs, Z. A., McGlynn, K., Mishra, P., andWhitehurst, A. W. (2020) Sperm-specific COX6B2 enhances oxidative phosphorylation, proliferation, and survival in human lung adenocarcinoma Elife **9**, 10.7554/eLife.58108
11. Gibbs, Z. A., Reza, L. C., Cheng, C. C., Westcott, J. M., McGlynn, K., andWhitehurst, A. W. (2020) The testis protein ZNF165 is a SMAD3 cofactor that coordinates oncogenic TGFbeta signaling in triple-negative breast cancer Elife **9**, 10.7554/eLife.57679

12. Whitehurst, A. W., Xie, Y., Purinton, S. C., Cappell, K. M., Swanik, J. T., Larson, B. *et al.*
     (2010) Tumor antigen acrosin binding protein normalizes mitotic spindle function to promote
     cancer cell proliferation Cancer Res **70**, 7652-7661 10.1158/0008-5472.CAN-10-0840
13. Sandhu, S., Sou, I. F., Hunter, J. E., Salmon, L., Wilson, C. L., Perkins, N. D. *et al.* (2021)
     Centrosome dysfunction associated with somatic expression of the synaptonemal complex
     protein TEX12 Commun Biol **4**, 1371 10.1038/s42003-021-02887-4
14. Dervovic, D., Malik, A. A., Chen, E. L. Y., Narimatsu, M., Adler, N., Afiuni-Zadeh, S. *et al.*
     (2023) In vivo CRISPR screens reveal Serpinb9 and Adam2 as regulators of immune therapy
     response in lung cancer Nat Commun **14**, 3150 10.1038/s41467-023-38841-7
15. Hofmann, O., Caballero, O. L., Stevenson, B. J., Chen, Y. T., Cohen, T., Chua, R. *et al.* (2008)
     Genome-wide analysis of cancer/testis gene expression Proc Natl Acad Sci U S A **105**, 20422-
     20427 10.1073/pnas.0810777105
16. Salicioni, A. M., Gervasi, M. G., Sosnik, J., Tourzani, D. A., Nayyab, S., Caraballo, D. A. *et al.*
     (2020) Testis-specific serine kinase protein family in male fertility and as targets for non-
     hormonal male contraceptiondagger Biol Reprod **103**, 264-274 10.1093/biolre/ioaa064
17. Spiridonov, N. A., Wong, L., Zerfas, P. M., Starost, M. F., Pack, S. D., Paweletz, C. P. *et al.*
     (2005) Identification and characterization of SSTK, a serine/threonine protein kinase essential
     for male fertility Mol Cell Biol **25**, 4250-4261 10.1128/MCB.25.10.4250-4261.2005
18. Jha, K. N., Tripurani, S. K., andJohnson, G. R. (2017) TSSK6 is required for gammaH2AX
     formation and the histone-to-protamine transition during spermiogenesis J Cell Sci **130**, 1835-
     1844 10.1242/jcs.202721
19. Sosnik, J., Miranda, P. V., Spiridonov, N. A., Yoon, S. Y., Fissore, R. A., Johnson, G. R. *et al.*
     (2009) Tssk6 is required for Izumo relocalization and gamete fusion in the mouse J Cell Sci
     122, 2741-2749 10.1242/jcs.047225
20. Li, B., Li, T., Pignon, J. C., Wang, B., Wang, J., Shukla, S. A. *et al.* (2016) Landscape of
     tumor-infiltrating T cell repertoire of human cancers Nat Genet **48**, 725-732 10.1038/ng.3581
21. Cerami, E., Gao, J., Dogrusoz, U., Gross, B. E., Sumer, S. O., Aksoy, B. A. *et al.* (2012) The
     cBio cancer genomics portal: an open platform for exploring multidimensional cancer
     genomics data Cancer Discov **2**, 401-404 10.1158/2159-8290.CD-12-0095
22. Roig, A. I., Eskiocak, U., Hight, S. K., Kim, S. B., Delgado, O., Souza, R. F. *et al.* (2010)
     Immortalized epithelial cells derived from human colon biopsies express stem cell markers and
     differentiate in vitro Gastroenterology **138**, 1012-1021 e1011-1015
     10.1053/j.gastro.2009.11.052
23. Zhang, L., Theodoropoulos, P. C., Eskiocak, U., Wang, W., Moon, Y. A., Posner, B. *et al.*
     (2016) Selective targeting of mutant adenomatous polyposis coli (APC) in colorectal cancer
     Sci Transl Med **8**, 361ra140 10.1126/scitranslmed.aaf8127
24. Cicenas, J., Zalyte, E., Bairoch, A., andGaudet, P. (2018) Kinases and Cancer Cancers
     (Basel) **10**, 10.3390/cancers10030063
25. Van Tongelen, A., Loriot, A., andDe Smet, C. (2017) Oncogenic roles of DNA hypomethylation
     through the activation of cancer-germline genes Cancer Lett **396**, 130-137
     10.1016/j.canlet.2017.03.029
26. Jha, K. N., Wong, L., Zerfas, P. M., De Silva, R. S., Fan, Y. X., Spiridonov, N. A. *et al.* (2010)
     Identification of a novel HSP70-binding cochaperone critical to HSP90-mediated activation of
     small serine/threonine kinase J Biol Chem **285**, 35180-35187 10.1074/jbc.M110.134767
27. Li, Z. N., andLuo, Y. (2023) HSP90 inhibitors and cancer: Prospects for use in targeted
     therapies (Review) Oncol Rep **49**, 10.3892/or.2022.8443
28. Zhang, X., Peng, J., Wu, M., Sun, A., Wu, X., Zheng, J. *et al.* (2023) Broad phosphorylation
     mediated by testis-specific serine/threonine kinases contributes to spermiogenesis and male
     fertility Nat Commun **14**, 2629 10.1038/s41467-023-38357-0
29. Hawkinson, J. E., Sinville, R., Mudaliar, D., Shetty, J., Ward, T., Herr, J. C. *et al.* (2017) Potent
     Pyrimidine and Pyrrolopyrimidine Inhibitors of Testis-Specific Serine/Threonine Kinase 2
     (TSSK2) ChemMedChem **12**, 1857-1865 10.1002/cmdc.201700503

30.   Varghese, F., Bukhari, A. B., Malhotra, R., andDe, A. (2014) IHC Profiler: an open source plugin for the quantitative evaluation and automated scoring of immunohistochemistry images of human tissue samples PLoS One **9**, e96801 10.1371/journal.pone.0096801

31.   Kovacs, S. A., Fekete, J. T., andGyorffy, B. (2023) Predictive biomarkers of immunotherapy response with pharmacological applications in solid tumors Acta Pharmacol Sin **44**, 1879-1889 10.1038/s41401-023-01079-6

Figure 1. TSSK6 is frequently ectopically expressed in colon cancer (**A**) Kaplan Meier curve for regression free survival based on median expression of TSSK6 in colorectal cancer. **(B)** Representative images of IHC staining and associated intensity score (inset). Scale bars = 100 m. (C) Indicated cells were lysed, resolved on 14% SDS-Page and immunoblotted with indicated antibodies. (D) Indicated cells were transfected with indicated siRNAs for 96 hours, lysed and resolved on a 12% SDS-PAGE page and blotted with indicated antibodies. (E) As in (C).

Figure 2. TSSK6 is necessary and sufficient for colorectal cancer oncogenic behaviors **(A)** Indicated cells were transfected for 96 hours followed by measurement of ATP using Cell Titer Glo. Bars represent mean ± s.d. (n=6). Statistical analysis by Mann-Whitney t-test. (B) Top: Indicated cell lines were transfected for 48 hours then plated into soft agar. 14-21 days later, samples were stained and quantitated. Below: Quantitation of colonies above. Mean is plotted ±s.d. (n≥6). Statistical analysis by Student's t-test. (C) As in B for second siRNA pool targeting TSSK6 in HCT116 cells. Bars represent mean. Statistical analysis by Mann-Whitney t-test. (D) LOVO cells were transfected with indicated siRNA for 48 hours and plated into invasion chambers. 48 hours later, cells were stained and quantitated. Bars represent mean ± s.d. (n=9). Statistical analysis by student's t-test. (E) Left: Immunoblot of indicated cell lines with indicated antibodies. Samples resolved on 10% SDS-PAGE Right: 4000 cells were seeded for 96 hours followed by Cell Titer Glo assay. Bars represent mean ± s.d. (n=9). (F) Indicated DLD-1 cells were plated into soft agar for 14 days, stained and quantitated. Left: Representative images with arrow heads indicating colonies. Right: Graph represents mean colonies ± s.d. (n=9). Statistical analysis by Mann Whitney t-test. (G) 50,000 DLD1 cells were plated into invasion chambers for 24 hours and subsequently stained and quantitated. Left: Example image. Right: Graphs represent mean± s.d. (n=9). Statistical analysis by student's t-test.

Figure 3: TSSK6 kinase activity is required for transforming activity
 (A) Immunoblot from indicated cell lines with indicated antibodies. Samples were run on a 14% SDS-PAGE. (B) Indicated cell lines were plated for 96 hours and ATP quantitated with Cell Titer Glo. Bars represent mean ±s.d. (n=6). (C) Indicated HCEC cells were grown in soft agar for 21 days then stained and counted. Left: Representative images are shown. Right: Quantitation of colonies. Mean is plotted ±s.d. (n=6). Statistical analysis performed by student's t-test. (D) Left: TSSK6 was immunoprecipitated with a V5 antibody from HCEC cells and IP-Kinase assays performed. Bars represent mean ±s.d. (n=3). Right: Representative blot of parallel lysates resolved on 14% SDS-Page and immunoblotted with indicated antibodies. (E) Indicated HCEC cells were plated into soft agar for 21 days then stained and counted. Left: Representative images. Right: Quantitation of colonies under indicated conditions. Mean is plotted ±s.d. (n=12). Statistical analysis by student's t-test. Below: Representative immunoblot of indicated cell lines with indicated antibodies. (F) Indicated HCEC cell lines were plated into invasion chambers for 48 hours then stained and counted. Representative images are displayed. Invading cells were quantitated and graph represents mean ±s.d. (n=6). Statistical analysis by Mann Whitney t-test.

Figure 4. TSSK6 localizes to and enhances focal adhesion formation.
(A) Confocal images of HCT116 cells transfected with indicated siRNAs for 72 hours, fixed and immunostained with indicated antibodies. Scale bar is 20 M. (B and C) Left: Confocal images of HCEC cells plated onto glass coverslips for one day prior to staining with indicated

antibodies. Arrowheads indicate areas of overlap. Scale bar is 20 M. Hatched box is magnified in lower images with scale bar representing 10 M. Right: Paxillin and Tensin positive foci were manually quantitated in both the HCEC-Ctrl and HCEC-TSSK6-V5 cells. Mean is indicated with S.D. P-value calculated by Mann-Whitney test.

Figure 5. TSSK6 is required for tumor growth in vivo. (A) 1x 10^6 HCT116 cells stably expressing indicated constructs were xenografted into the flank of NSG mice Left: Tumor volume measurements were taken by caliper on indicated days. Each data point represent the mean (n=9) ± S.D. Right: Mass of excised tumors. Bars represent the mean (n≥7) ± S.D P-value calculated by Mann Whitney t-test. Upon excision of tumors, two of the largest shCTRL tumors excreted internal contents, thus final masses were not obtainable. Right: Immunoblot of lysates prior to injection resolved on 10% SDS-PAGE and immunoblotted with indicated antibodies (B) 800,000 DLD-1 cells transfected with indicated constructs were injected into the flank of a NSG mouse. Left: Tumor volume measurements were taken by caliper on indicated days. Each data point represent the mean (n=4) ± s.d. Statistical test performed by Mann Whitney. Right: Mass of excised tumors. Bars represent the mean. Statistical analysis performed by student's t-test (n=4).

Delgado et al
Figure 1

A.

B. **Testis (10X)**   **Colon Adenocarcinoma (20X)**
Negative   Low

Medium   High

Staining Intensity	#Cases
Negative	7
Low	3
Medium	5
High	5

C.

D.

E.

Delgado et al
Figure 2

A.

B.

TSSK6 Positive

C.

D.

E.

DLD-1

F.

G.

A.

B.

C.

D.

E.