Methane and ethane production rates by methyl-coenzyme M reductase in cell extracts from different methanogens

V. P. Thinh Nguyen[†], Sebastian Keller & Silvan Scheller*

Department of Bioproducts and Biosystems, School of Chemical Engineering, Aalto University, FI-02150, Espoo, Finland

*Corresponding author: silvan.scheller@aalto.fi

[†]Current address: Canadian Glycomics Network, E5-33 Gunning/Lemieux Chemistry Centre, T6G 2E2, Edmonton, Alberta, Canada

ORCID

Silvan Scheller 0000-0002-0667-9224

V. P. Thinh Nguyen 0000-0002-2499-2900

Sebastian Keller 0000-0001-7477-1201

Abstract

Methyl-Coenzyme M reductase (Mcr) plays an important role in the regulation of the global carbon cycle. The enzyme catalyzes the reversible conversion of methyl-coenzyme M and coenzyme B to methane and the corresponding heterodisulfide CoM-S-S-CoB, which constitutes the first step of the anaerobic oxidation of methane. Mcr homologs of varies methanogenic archaea also catalyze the anaerobic oxidation of extended alkanes. Fully active, purified enzyme is exclusively achievable *via* strictly anaerobic purification from *Methanothermobacter marburgensis.* This microbe expresses two isoenzymes and most studies were performed with isoenzyme I that exhibits a limited substrate-promiscuity (c.a. 0.5%) towards the homologous substrate ethyl-coenzyme M. Here, cell-free lysates from the different species such as *Methanosarcina mazei*, *Methanococcus maripaludis*, *Methanothermococcus okinawansis,* and *Mt. marburgensis* were screened for Mcr activity and substrate promiscuity. An assay relying on titanium (III) citrate and cobalamin to regenerate coenzyme B and coenzyme M was used to test the ability of different cell extracts for the catalytic activity towards the C2-substrate ethyl-coenzyme M. Cell extracts from *M. mazei* showed a ratio of ethane- to methane- production of ca. 8% at 37 °C, and about 14% at 49 °C. The level of substrate-promiscuity towards ethyl-coenzyme M for *M. marburgensis* cell extracts under our assay conditions were much higher than previously reported for purified isoenzyme I, indicating that isoenzyme II is much more promiscuous for ethane formation than isoenzyme I. Our experiments demonstrate that Mcr activity can be quickly and conveniently studied *via* cell-free lysates, and that substrate promiscuity towards ethane formation is generally larger than anticipated.

Keywords

Enzyme catalysis, methanogenic archaea, methyl-coenzyme M reductase, biological methane formation, C1-metabolism, substrate promiscuity, enzyme assays, response surface methodology

Introduction

Methyl-Coenzyme M reductase (Mcr) catalyzes the reversible last step of methanogenesis, where it catalyzes the conversion of methyl-Coenzyme M (CoM) and the thiol coenzyme B (CoB-SH) to methane and the corresponding heterodisulfide coenzyme B–coenzyme M (CoB-S-S-CoM) in all methanogenic archaea [1] (**Figure 1**). Mcr also catalyzes the first step of anaerobic oxidation of methane, and even more extended alkanes in alkanotrophic archaea [2–4]. This archaeal reductase plays an important role in the global carbon cycle by catalyzing the last step in archaeal methane formation with an estimated global flux of c.a. 1 Gt per year of methane [5]. Investigating the anaerobic oxidation of ethane to corresponding ethyl-CoM mediated by *Methanothermobacter marburgensis's* Mcr has shown the promiscuous conversion of ethyl-CoM to the corresponding ethane [6]. Although an early study has reported an accumulation of ethane from ethyl-CoM carried out by cell-free extracts from *Methanosarcina barkeri* under hydrogen supply [7], the Mcr enzyme activity for alkyl-CoM of *M. barkeri* and other methanogens, besides *Mt. marburgensis*, remains limited.

As mentioned previously, most of the kinetic information was collected from *Mt. marburgensis* Mcr's studies [8,9], whereas insights into the substrate specificity and enzyme activity of other methanogens remain limited up-to-date. It is reported that Mcr isolated from *Mt. marburgensis* possesses two different isoenzymes. The Mcr activity of cell lysates harvested in the late log phase is mainly attributed to the type I isozyme [10]. This later constitutes 0.14% of the Mcr remaining in the harvested cell [9,11], while the isoenzyme II was exclusively expressed in the early log phase. The specific enzyme activity of isoenzyme II (11.7 U/mg) was reported to be significantly higher than that of isoenzyme I (4.5 u/mg) [11]. It is noteworthy that the specific activity of Mcr also varies dependently on the amount of the MCR-red1 form residing in the harvested cell. Specific activities for the methyl-CoM reduction of purified *Mt. marburgensis* Mcr were shown to be elevated over 20 U/mg when Ti(III) citrate and cobalamin were added for the reactivation of the NI(I) state of cofactor F_{430} [11,12], with maximal 100 U/mg upon fully activation of the enzyme [11].

Methyl-CoM reductase requires the nickel-porphinoid [13] cofactor F_{430} in its Ni(I) oxidation state to be active [11,12]. In the *in vitro* assay, Mcr owes the efficient reduction of CoM-S-S-CoB heterodisulfide to the presence of Co(I) or Co(II) of Ti(III)citrate-reduced cyanocobalamin. The hypothesis is this reduction system can be implemented to other Mcr homologues from other methanogenic and alkanotrophic species to maintain or reactivate the enzyme activity for methyl-CoM and even for more extended alkyl-CoM derivatives (e.g., ethyl-CoM). However, the purification of active Mcr from other methanogens appears to be challenging due to the high O_2-sensitivity, as no such successful procedure has been communicated in the literature. This inactivation of Mcr is prompted by the presence of oxidized form of Ni-porphinoid to (II) and (III) oxidation states [13]. Nevertheless, Mcr-Ni(III) can be further

reactivated by using strong electron-donors such as Ti(III) citrate or DTT under either H_2 or

CO atmosphere. A lack of critical post-translational modifications furthermore impactfully

contribute to the inactivity of Mcr [14,15]. In spite of the aforementioned limitations, Mcr-Ni(III)

can be further reactivated by using strong electron-donors such as Ti(III) citrate or DTT under

either H_2 or CO atmosphere. Taken together, the functional heterologous expression of Mcr

remains challenging [15] hampering the study of this of interest enzyme. Moreover, the

utilization of the Ti(III) citrate/cobalamin reduction system to rescue the Mcr activity is

exclusively limited to *Mt. marburgensis* purified Mcr [6,9,11].

Cell-free lysate can be used instead of purified Mcr [7]. This approach will allow one to

address the limit of arduous purification or reactivation processes of Mcr. Employing cell-free

extract is thereby an excellent strategy for rapid assay of Mcr enzyme activity for alkyl-CoM to

form corresponding alkanes. In this context, we aim to provide herein the optimal conditions

of Mcr enzyme activity for ethane formation from corresponding ethyl-CoM basing on the study

conducted with cell-free lysate isolated from *M. acetivorans* (strain WWM73) grown on

methanol (MeOH), as well as to investigate the substrate promiscuity of Mcr from different

methanogen cell-free extracts.

Results

In this study, we have adapted the previously published enzyme activity assay of Mcr

purified from *M. marburgensis* [9] to cell-free lysates isolated from the other cultivable

methanogens such as *Methanosarcina* and *Methan*ococcaceae. We aimed to compare the

enzyme activities towards C1- and C2-alkyl donor (methyl- and ethyl-CoM, respectively). The

promiscuity of cell-free lysates, defined by the ratio between ethane and methane formation

from respective ethyl- and methyl-CoM, were also studied. The optimization of the Mcr assay

was conducted using cell-free lysates from *M. acetivorans* (strain WWM73) regarding its fast

growth pace on methanol (MeOH) [16]. The culture was supplemented with 100 mM MeOH

and 2.5% Na_2S as sulfur source under strict anaerobic conditions. Once the desired growth

phase reached, the cell-free lysate was then prepared from the active culture and will be used

to catalyze the conversion of methyl- and ethyl-CoM to corresponding methane and ethane,

respectively. The standard Mcr assay contained 0.5 mM alkyl-CoM, 0.25 mM CoB, 0.2 mM

vitamin B_{12}, 20 mM of Ti(III) citrate, and 100 µg of cell-free lysate at pH 7 and 25 °C, while the

enzyme activity was determined by methane and ethane level detection via GC-FID.

An experimental design approach employing response surface methodology was

performed (see **Supporting Information)** to provide further insights concerning the effects of

predefined parameters onto enzyme activity for ethyl-CoM alongside with the ratio between

enzyme activity for ethyl-CoM and that for methyl-CoM (henceforth referred as the

promiscuity). The outcomes of this experimental design afforded optimal operating conditions

which would be next implemented to assay enzyme activity for ethyl-CoM and the promiscuity
of different methanogens. We would expect to identify the most promiscuous methanogen-
Mcr among tested methanogens.

Determination of parameters having effects onto Mcr activities

It appeared that selected parameters such as the growth phase at harvesting, the
incubation temperature and the B_{12} concentration involving in the CoB-S-S-CoM
heterodisulfide reduction have significant effects on Mcr enzyme activity for methyl-CoM. In
order to study these effects, we devised a response surface design involving 3 factorial
variables (OD_{600} at harvesting, incubation temperature and vitamin B_{12} concentration) that
included, not only the enzyme activity for methyl-CoM as a response variable, but also
comprised the enzyme activity for ethyl-CoM, and the promiscuity (ratio between ethane and
methane formed from ethyl-CoM and methyl-CoM, respectively) with regards to the main goal
of this study is to assay the ethane formation ability of studying methanogens. Thus, the
optimal operating conditions were implemented to explore the methane and ethane formation
ability, as well as the promiscuity of cell-free lysates from different methanogens.

Growth phase

Cell lysates of different growth phases, with OD600 at harvesting were categorized as early
(low OD, value between 0.1-0.3), middle (moderate OD, value between 0.5-0.7) and late
growth phase (high OD, value between 0.9-1.2). Mcr assays employing these cell lysates were
conducted to determine the effects of growth phase on methanogenic activities. Methyl-CoM
was used as an alkyl donor, and methane was quantified by GC-FID using methane as an
external standard curve. Our results indicated that *M. acetivorans* Mcr was most active at mid
growth phase (OD600 value between 0.5-0.8), whereas harvesting at early growth phase or
late growing time resulted in lower enzyme activity as depicted in **Figure 2**.

Incubation temperature and vitamin B_{12} concentration

We investigated the effects of incubation temperature during the assay onto the enzyme
activity by conducting Mcr assay at different temperatures (25 and 37 °C) employing different
cell-free lysates prepared at different growth phases (low and high OD levels). Our results
presented in **Figure 3** shows that low OD culture exhibited higher enzyme activity at 37 °C.
On the other hand, the high OD cell-free lysate did not show any improvement in enzyme
activity for methyl-CoM at higher temperature.

Ti(III) citrate is used along with the addition of cobalamin to reduce the generated CoB-S-
S-CoM heterodisulfide complex in the Mcr activity assay [11,12]. The concentration of Ti(III)
must be in excess, whereas the cobalamin concentration was reported to be between 0.2-2
mM to ensure the highest Mcr activity [12]. Taken together, we hypothesized that cobalamin
concentration might have an indirect effect on Mcr assay. Therefore, the concentration of

vitamin B_{12}–the source of cobalamin in our assay–was also included as a factorial variable for

our design of experiment. We investigated the concentration range of vitamin B_{12} between 0.2

to 1.1 mM, as higher concentration might not be directly beneficial to our assay. Our results

depicted in **Figure 3** showed that low OD and low cobalamin concentration had significant

positive effect onto enzyme activity for methyl-CoM, whereas increasing cobalamin

concentration only had a slight declination of the enzyme activity for high OD cell-free lysate.

These results confirmed our aforementioned hypothesis concerning the indirect effect of

cobalamin onto Mcr enzyme activity.

Based on our previous results, we proceeded with the optimization of the Mcr enzyme

activity with a non-C1 substrate, ethyl-CoM, in parallel in order to determine the enzyme

activity of Mcr in cell-free lysates towards more extended substrate (C2 in this study), as well

as the promiscuity of Mcr towards these substrates. For the latter purpose, we determined the

ratio between ethane and methane formed by separately providing methyl-CoM and ethyl-

CoM as alkyl donor was therefore determined. Response surface methodology is a powerful

statistical technique that has been successfully employed to optimize the methanogenic

activity of methanogens [17]. Thereby, we believed that employing this statistical approach

would afford an efficient and highly reproducible procedure that enables alkane formation from

corresponding alkyl-CoM. The outcomes of this optimization process would allow us to

investigate the enzyme activity for ethyl-CoM and promiscuity of cell-free lysate from *M.*

acetivorans. The model fitting and further optimization of enzyme activity are presented in **the**

Supporting Information. Confirmatory experiments were conducted in duplicate according

to the operating conditions afford *via* the prediction model to confirm the estimating values.

The outcomes were reported in **Table 1**, with the enzyme activity for methyl-CoM appeared to

be higher than the predicted value. Nevertheless, the value of this latter remained satisfactory

in the context of our study. Thus, we proceeded to the implementation of our optimized assay

conditions onto well-known cultivable methanogens. The results of this process are described

in the following section.

Implementation of optimal Mcr assay to investigate the enzyme activity and promiscuity of

different cell-free lysate harvested from available methanogens

Enzyme activity towards ethyl-CoM and the promiscuity of cell-free lysates from different

cultivable methanogens such as *M. mazei*, M. *M. maripaludis*, *Mt. okinawansis* and *Mt.*

marburgensis were studied under previously optimized operating conditions (OD_{600} at

harvesting at moderate/low, 49 °C and 1.1 mM B_{12}). In addition, experiments under

conventional assay conditions (moderate OD600, incubation temperature 37 °C and vitamin

B_{12} concentration 0.2 mM) were also conducted, as we were concerned that the high

incubation temperature and vitamin B_{12} concentration might significantly affect onto the Mcr

activity of non-thermal methanogens, originally living in cold conditions [17]. For *M. mazei,*

low protein concentration in lysate prepared from cell harvested at OD600 between 0.4-0.6

occurred. As a result, the cell was instead harvested at OD600 value between 1.4-1.6 to

afford the sufficient amount of protein for the Mcr assay. In contrast to other methanogens,

Mt. okinawansis cell-free lysates appeared inactive under both conventional and optimal

conditions. The results are reported in **Figure 4**.

Dependence of substrate promiscuity of *Methanothermobacter marburgensis* on different

growth conditions

We investigated the high promiscuity and ethane formation from ethyl-CoM of *Mt.*

marburgensis by conducting the Mcr assays with different cell-free lysates obtained *via*

different growth methods: batch growth in serum vials with frequently supplement of H_2/CO_2,

and biofermentation in bioreactor with a continuous flow of H_2/CO_2 [6,9]. OD_{600} value at

harvesting of the former growth method was reported to be between 0.5-0.7, whereas the

latter one was harvested during the plateau growth phase (OD_{600} between 5-6). In both

cases, the cell suspension appeared to be slightly fluorescent green due to enrichment of

flavin-based cofactor F_{420}. Upon lysis, the cell-free extracts were retrieved and stored at - 80

ºC under H_2 until further uses.

Upon obtaining the cell-free lysates, we conducted assays of Mcr activity using either

methyl- or ethyl-CoM alone, or mixing methyl- and ethyl-CoM at different ratios to study Mcr

activities of lysates. The conventional assay conditions (0.2 mM vitamin B_{12} concentration,

20 mM Ti(III) citrate) and high incubation temperature (65 °C) were implemented to compare

the enzyme activity for methyl- & ethyl-CoM. Our results in **Figure 5** indicated that cell-free

lysate of *M. marburgensis* grown in serum vial and low-OD600 value at harvesting exhibited

better enzyme activity for ethyl-CoM, with the enzyme activity for methyl-CoM is c.a. 19

times higher than that for ethyl-CoM. In contrast, the lysate growth in bioreactor showed

lower activity for ethyl-CoM, with the enzyme activity ratio between for methyl-CoM and for

ethyl-CoM was determined to be c.a. 230 times, which is in agreement with our previously

reported data [6].

We observed that the presence of ethyl-CoM in high concentration drastically inhibited

the conversion of methyl-CoM to corresponding methane in cell-free lysates of *M.*

marburgensis grown in bioreactor while varying the concentration of methyl- and ethyl-CoM

in the alkyl-CoM mixture (**Figure 5**). In contrast, *M. marburgensis* cell-free lysate harvested

at early growth phase showed higher tolerance towards the presence of ethyl-CoM in the

alkyl-CoM mixture. Ethane formation was observed in both cases where only ethyl-CoM was

supplemented.

In order to rationalize our observation, Mcr composition was analyzed via SDS-PAGE
following the protocol described by Rospert et al. [18]. Thus, lysates of *Mt. marburgensis*,
collected either at a low (vial), or high OD (bioreactor) were step-wise enriched with
ammonium sulfate to saturated level for a partial purification of the Mcr isoenzymes (**Figure**
6). The bioreactor sample from a late growth phase showed basically three main bands
corresponding well to Mcr isoform I [10,18]. Both Mcr isoforms can be distinguished basing
the size difference of the gamma subunit *via* SDS-PAGE analysis. The respective sizes of
small γ subunits are 38 kDa for isozyme I and 33 kDa for isozyme II [10]. In the case of the
vial sample that originated from an earlier growth phase, Mcr isoform I appeared to dominate
the total Mcr fraction. However, a faint band occurred at the corresponding size of the Mcr II
gamma subunit (33 kDa), which was absent in the bioreactor sample. We also observed a
ratio of ca. 3:1 for Mcr I to Mcr II based on the intensity of the two gamma subunit bands in
the vial sample.

Discussion
Optimal Methyl-Coenzyme M reductase assay enhancing enzyme activity and the
promiscuity for ethyl-CoM
In this study, we have optimized the operating conditions for the Mcr methane formation
assay, that so far exclusively employs purified Mcr from *M. marburgensis,* onto other
methanogens such as *M. acetivorans*, *M. mazei*, *M. maripaludis,* and *M. okinawansis*. The
optimal conditions of 3 factorial variables such as OD_{600} at harvesting, incubation temperature,
and B_{12} concentration were afforded *via* an experimental design using the response surface
methodology with Box-Behnken design. Our prediction model suggested that increasing the
incubation temperature at high B_{12} concentration efficiently enhance the formation of ethane
from ethyl-CoM along with the promiscuity towards more extended alkane from corresponding
alkyl-CoM. Experiments conducted with cell-free lysate of *M. acetivorans* harvested at
low/moderate OD_{600} (OD_{600} value at harvesting 0.3-0.5) at high B_{12} concentration (1.1 mM)
and temperature (49 °C) afforded higher enzyme activity for methyl- and ethyl-CoM. These
outcomes suggested the thermostability of Mcr from *M. acetivorans*. This can involve the
different composition of the whole Mcr complex between these species [19]. Nevertheless, the
improvement of the conversion of methyl- and ethyl-CoM to corresponding alkane prompt by
increasing incubation temperature and B_{12}-concentration deserves further investigation, with
high regards to limited respective information relating to Mcr thermostability and enzyme
reactivation mechanism.
Our results depicted in **Figure 5** showed that cell-free lysate from different methanogens
exhibiting the activity for conversion of methyl- and ethyl-CoM to corresponding alkanes
under both conventional conditions. Despite the high OD600 value at harvesting, the cell-

free lysate from *M. mazei* also showed the highest Mcr activity for both provided alkyl-CoM substrates as well as the promiscuity. These values even exceeded that of thermo-methanogens such as *Mt. marburgensis* and *Mt. okinawansis*. In contrast, cell-free lysate *M. maripaludis* was determined to be the less efficient material to carry out Mcr activity for both methyl- and ethyl-CoM. Overall, the Mcr activity appeared to improve at increasing vitamin B_{12} concentration and incubation temperature. Regardless the low activity, this is the very first time Mcr activity of two different cultivable members of *Methanococcacaea* (*M. maripaludis* and *Mt. okinawansis*) has been reported according to our best knowledge.

We have successfully implemented the optimal Mcr conditions for ethane formation onto different methanogenic archaea and have identified *M. mazei* to be the most promiscuous Mcr for ethane formation as well as the promiscuity for ethyl-CoM as substrate.

Cell-free lysate as materials to assay Mcr activity towards non-C_1 alkyl-Coenzyme M

As previously mentioned, data relating to Mcr enzyme activity is mainly reported by studying pure Mcr isoenzyme I isolated from *Mt. marburgensis* [6,9,15]. In this study, we have demonstrated that cell-free lysate can also be used to quantify, in a preliminary manner, the Mcr activity for methyl-CoM and ethyl-CoM. Optimized Mcr assay employing Ti(III) citrate/B_{12} heterodisulfide reducing system furthermore resulted in our findings about the enhancement of activity and promiscuity with increasing assay temperature and B_{12}-concentration for extended alkyl-CoM of multiple methanogen Mcr such as *M. mazei*, *M. acetivorans*, *Mt. marburgensis* (**Figure 6**). In contrast, Mcr from *Methanococcacaea* appears to be thermos- and B_{12}-sensitive. Further investigation is needed to fully understand the behavior of these latter.

The cell-free lysate approach employed in this study enables a quick analytical method to assay the promiscuity of Mcr towards non-C1 alkyl-CoM. Our results furthermore confirmed that this assay is applicable on a broad range of cultivable methanogens. We also showed that *Methanosarcina* is a potential role model for overexpression of yet-to-be alkyl-Coenzyme M reductase, with high regards to its promiscuity towards non-C1 substrates. Moreover, our method emphasized the H_2 availability- and/or the abundance of Mcr isoenzyme-dependence of substrate promiscuity of *Mt. marburgensis*.

Optimal Mcr assay allowing further understanding onto Mcr activity of *Methanothermobacter marburgensis*

Our results showed that cell-extracts of *Mt. marburgensis* grown under non-hydrogen limitation, in which Mcr-isoenzyme II is stronger expressed, exhibited an ethane-formation rate of ca. 7.7 %. This later ethane formation level showed that the well-studied enzyme variant Mcr-I from *M. marburgensis* has a low substrate promiscuity towards C2, and substrate promiscuity is generally higher than anticipated. Mixtures of methyl- and ethyl-

coenzyme M showed a substantial inhibitory effect of ethyl-coenzyme M for methane formation, in particular for Mcr-I, which is in line with the previous finding that ethyl-coenzyme M is readily processed by Mcr, but ethane formation is blocked in this enzyme variant.

We have attempted without success the conversion of propyl-CoM employing different promiscuous cell-free lysates such as *M. mazei*, *M. acetivorans*, and *Mt. marburgensis* under different assay conditions (data not shown). Nevertheless, multiple genetic modification tools for methanogens have been developed [20,21] along with expression of uncultured archaea Mcr [22] which may allow better conversion of longer alkyl-CoM (C3 and above) in the near future. To conclude, employing our cell-free lysate Mcr assay approach to investigate this enzyme family is a rapid and efficient assay to test the enzyme activity of these recombinant enzymes in order to provide a first milestone for further engineering of artificial cultivable alkanogens.

Materials and Methods

Chemical and Materials

Methyl-CoM, ethyl-CoM, and coenzyme B were synthesized by following the reported protocol [19]. Gas formation was recorded on an Agilent 6890 GC-FID system (Agilent Technologies, CA, USA) and analyzed using OpenLab CDS software (Agile Technologies, CA, USA). OD600 value and Bradford assay were recorded on BioPhotometer Plus 6132 spectrophotometers (Eppendorf, Germany) at 595 nm wavelength. All chemicals were purchased from Merck. Titanium (III) citrate were prepared following the description elsewhere [23]. MilliQ water was produced by Milli-Q Direct (Merck Millipore, MA, USA). Universal Hood II-Gel Doc system (Bio-Rad) was used to display the SDS-PAGE gel.

Strains, cultivation, and cell-free lysate preparation

M. barkeri (strain Fusaro), *M. acetivorans* (strain WWM73) [16], and *M. mazei* (strain Gö I) [21], *M. maripaludis* (strain JJ:Δupt) [24], *Mt. okinawansis* [25] and *M. marburgensis* [9], were used for this study. For *M. acetivorans,* cell pellets were obtained at different OD600 values. For the other strains, cell pellets were obtained at OD600 value between 0.3-0.5.

For *M. acetivorans*, cultures were first grown statically in High-Salt (HS) medium supplemented with 150 mM MeOH as substrate and 1 mM Na_2S as sulfur source at 37 °C for 1 day [26]. Typical OD_{600} reaches 0.18-0.20. For low OD_{600} experiments, cultures were then harvested by centrifugation, resuspended in HS medium to yield the cell pellet. For moderate and high OD_{600} experiments, cultures were grown at 25 °C (for moderate OD_{600}) or 37 °C (for high OD_{600}) until OD_{600} reaching the desired value range. The cell pellet was then obtained by centrifugation.

For *M. barkeri,* the desired OD_{600} value (0.3-0.5) can be reached within 3-4 days by
statically cultivating at 37 °C in HS medium supplemented with H_2 and 1% of H_2S as growth
substrate and sulfur source, respectively. For *M. marburgensis*, the cell pellet was obtained
using reported cultivation method [27] under H_2/CO_2 with H_2S as growth substrate and sulfur
source, respectively. For *M. mazei*, the HS medium was used as growth liquid supplemented
with 125 mM MeOH as growth substrate and 1% H_2S as sulfur source [28]. For *M. maripaludis*
and Mt. okinawansis, the liquid McFc medium was used for cultivation [29].

Cell-free lysate was prepared by sonication (Bandelin Sonopuls, Germany) at 30%
amplitude, 30 s pulse and 30 s rest time for 10 min. The cell-lysate fraction was then retrieved
by centrifugation and the protein concentration was determined by Bradford assay [30]. Cell-
free lysates were then aliquoted in 10 mL serum vials, of which the headspace was filled with
100% H_2 and then stored at -80 °C until further uses.

Enzyme activity assay employing cell-free lysate

The enzyme assay was performed following previously published protocol [31]. In brief, the
assay solution was prepared in 10 mL serum vial containing 0.5 mM alkyl-CoM, 0.25 mM CoB,
0.2-1.1 mM of vitamin B_{12}, 20 mM of Ti(III) citrate, and 100 µg of cell-free lysate (gradually
warmed at RT for 30 min. prior to the assay). Tris-HCl (50 mM, pH 7) was then added to make
up to 2 mL total volume. The solution was then mixed by pipetting and then closed with a
rubber stopper. The head space was flushed with 100% H_2 and then immediately transferred
into heat block at different temperature (25, 37, and 49 °C) for incubation. The gas formation
was monitored by injecting 100 µL of the head space into the GC-FID.

For the assay employing mixture of methyl- and ethyl-CoM, the previously described
protocol was employed with slight modification. The assay solution was prepared in 10 mL
serum vial containing methyl- and ethyl-CoM at different concentration were used as following;
for only methyl-CoM–5mM of methyl-CoM; for 9 equivalence (Eq) methyl-CoM:Eq ethyl-CoM–
4.5 mM of methyl-CoM and 0.5 mM of ethyl-CoM; for 1 Eq methyl-CoM:1 Eq ethyl-CoM–2.5
mM of methyl-CoM and 2.5 mM of ethyl-CoM; for 1 Eq methyl-CoM:9 Eq ethyl-CoM–0.5 mM
of methyl CoM and 4.5 mM of ethyl-CoM; for only ethyl-CoM: 5 mM of ethyl-CoM. 0.25 mM
CoB, 0.2 mM of vitamin B_{12}, 20 mM of Ti(III) citrate, and 100 µg of cell-free lysate (gradually
warmed at RT for 30 min. prior to the assay) were then sequentially added. Tris-HCl (50 mM,
pH 7) was then added to make up to 2 mL total volume. The solution was then mixed by
pipetting and then closed with a rubber stopper. The head space was flushed with 100% H_2
and then immediately transferred into heat block at 65 °C) for incubation. The gas formation
was monitored by injecting 100 µL of the head space into the GC-FID.

SDS-PAGE analysis of cell-free lysates of *Mt. marburgensis* harvested under different growth conditions

A fractionated ammonium sulfate precipitation of the *Mt. marburgensis* lysates was performed as described elsewhere [18]. The cell extracts were first well mixed with ammonium sulfate reaching a 70 % salt concentration and kept on ice for 2 h. After centrifugation (14,000 rpm, 5 min, 4 °C) the supernatants were further enriched with ammonium sulfate until the salt concentration reached about 100 %. After a second centrifugation step the pellets were resuspended in 10 µl sterile water. The protein concentration was estimated by the NanoDropTM Lite Spectrophotometer (Thermo Fisher). The samples were subsequently mixed with loading dye and incubated for 5 min at 95 °C.

For SDS PAGE a 10 % polyacrylamide gel was loaded with 2 µg protein/lane for both samples. Precision Plus Protein Dual Color Standard (#1610374, Bio-Rad) was applied as a size marker. The electrophoresis was performed at 120 V for 80 min before the gel was processed by Coomassie staining.

Statistical analysis

The results were expressed as the mean ± standard deviation. Replication was conducted in duplication. The significance of each coefficient was determined by using the Student *t-test* and p-value. The ANOVA analysis was conducted at 95% significance level. The analysis software used for this study was Minitab® 21 (version 3.1) (Minitab Inc, State College, Palo Alto, CA, USA.).

Acknowledgement

The authors would like to acknowledge Norman Adlung (VTT Finland) for guidance in the early development of the project. The authors would like to acknowledge the Academy of Finland (#13326020) and NovoNordisk Foundation (NNF310279) for fundings.

Conflict of interest

The authors declare no conflict of interest.

Author contribution

T.N. and S.S: conceptualization and data curation, T.N. & S.K: conducting experiments, T.N., S.K., & S.S. drafting the manuscript, S.S.: supervising, project administration and funding acquisition. All authors commented and approved the final version of the manuscript.

References

[1] Wang Y, Wegener G, Ruff SE, Wang F. Methyl/alkyl-coenzyme M reductase-based anaerobic alkane oxidation in archaea. Environ Microbiol 2021;23:530–41. https://doi.org/10.1111/1462-2920.15057.

[2] Wegener G, Laso-Pérez R, Orphan VJ, Boetius A. Anaerobic Degradation of Alkanes by Marine Archaea 2022. https://doi.org/10.1146/annurev-micro-111021.

[3] Chen SC, Musat N, Lechtenfeld OJ, Paschke H, Schmidt M, Said N, et al. Anaerobic oxidation of ethane by archaea from a marine hydrocarbon seep. Nature 2019;568:108–11. https://doi.org/10.1038/s41586-019-1063-0.

[4] Chen SC, Musat N, Lechtenfeld OJ, Paschke H, Schmidt M, Said N, et al. Anaerobic oxidation of ethane by archaea from a marine hydrocarbon seep. Nature 2019;568:108–11. https://doi.org/10.1038/s41586-019-1063-0.

[5] Thauer R. Biochemistry of methanogenesis : a tribute to Marjory Stephenson. vol. 144. 1998.

[6] Scheller S, Goenrich M, Thauer RK, Jaun B. Methyl-Coenzyme M Reductase from Methanogenic Archaea: Isotope Effects on Label Exchange and Ethane Formation with the Homologous Substrate Ethyl-Coenzyme M. J Am Chem Soc 2013;135:14985–95. https://doi.org/10.1021/ja4064876.

[7] Ahn Y, Krzycki JA, Floss HG. Steric course of the reduction of ethyl coenzyme M to ethane catalyzed by methyl coenzyme M reductase from Methanosarcina barkeri. J Am Chem Soc 1991;113:4700–1. https://doi.org/10.1021/ja00012a059.

[8] Wongnate T, Ragsdale SW. The Reaction Mechanism of Methyl-Coenzyme M Reductase. Journal of Biological Chemistry 2015;290:9322–34. https://doi.org/10.1074/jbc.M115.636761.

[9] Scheller S, Goenrich M, Thauer RK, Jaun B. Methyl-coenzyme M reductase from methanogenic archaea: Isotope effects on the formation and anaerobic oxidation of methane. J Am Chem Soc 2013;135:14975–84. https://doi.org/10.1021/ja406485z.

[10] Bonacker LG, Baudner S, Morschel E, Bocher R, Thauer RK. Properties of the two isoenzymes of methyl-coenzyme M reductase in Methanobacterium thermoautotrophicum. Eur J Biochem 1993;217:587–95. https://doi.org/10.1111/j.1432-1033.1993.tb18281.x.

[11] Goubeaud M, Schreiner G, Thauer RK. Purified Methyl-Coenzyme-M Reductase is Activated when the Enzyme-Bound Coenzyme F430 is Reduced to the Nickel(I) Oxidation State by Titanium(III) Citrate. Eur J Biochem 1997;243:110–4. https://doi.org/10.1111/j.1432-1033.1997.00110.x.

[12] Ellermann J, Rospert S, Thauer RK, Bokranz M, Klein A, Voges M, et al. Methyl-coenzyme-M reductase from Methanobacterium thermoautotrophicum (strain Marburg). Purity, activity and novel inhibitors. Eur J Biochem 1989;184:63–8. https://doi.org/10.1111/j.1432-1033.1989.tb14990.x.

[13] Lemaire ON, Wagner T. A Structural View of Alkyl-Coenzyme M Reductases,
the First Step of Alkane Anaerobic Oxidation Catalyzed by Archaea.
Biochemistry 2022;61:805–21. https://doi.org/10.1021/acs.biochem.2c00135.

[14] Prakash D, Wu Y, Suh SJ, Duin EC. Elucidating the process of activation of
methyl-coenzyme M reductase. J Bacteriol 2014;196:2491–8.
https://doi.org/10.1128/JB.01658-14.

[15] Thauer RK. Methyl (Alkyl)-Coenzyme M Reductases: Nickel F-430-Containing
Enzymes Involved in Anaerobic Methane Formation and in Anaerobic Oxidation
of Methane or of Short Chain Alkanes. Biochemistry 2019;58:5198–220.
https://doi.org/10.1021/acs.biochem.9b00164.

[16] Guss AM, Rother M, Zhang JK, Kulkkarni G, Metcalf WW. New methods for
tightly regulated gene expression and highly efficient chromosomal integration
of cloned genes for *Methanosarcina* species. Archaea 2008;2:193–203.
https://doi.org/10.1155/2008/534081.

[17] Jiménez J, Guardia-Puebla Y, Romero-Romero O, Cisneros-Ortiz ME, Guerra
G, Morgan-Sagastume JM, et al. Methanogenic activity optimization using the
response surface methodology, during the anaerobic co-digestion of agriculture
and industrial wastes. Microbial community diversity. Biomass Bioenergy
2014;71:84–97. https://doi.org/10.1016/j.biombioe.2014.10.023.

[18] Rospert S, Böcher R, Albracht SPJ, Thauer RK. Methyl-coenzyme M reductase
preparations with high specific activity from H_2-preincubated cells of
Methanobacterium thermoautotrophicum. FEBS Lett 1991;291:371–5.
https://doi.org/10.1016/0014-5793(91)81323-Z.

[19] Chadwick GL, Joiner AMN, Ramesh S, Mitchell DA, Nayak DD. McrD binds
asymmetrically to methyl-coenzyme M reductase improving active-site
accessibility during assembly. Proceedings of the National Academy of
Sciences 2023;120. https://doi.org/10.1073/pnas.2302815120.

[20] Zhu P, Bao J, Scheller S. CRISPR/Cas12a toolbox for genomic manipulation in
Methanosarcina acetivorans. BioRxiv 2023:526428.
https://doi.org/10.1101/2023.01.31.526428.

[21] Bao J, de Dios Mateos E, Scheller S. Efficient CRISPR/Cas12a-Based
Genome-Editing Toolbox for Metabolic Engineering in *Methanococcus*
maripaludis. ACS Synth Biol 2022;11:2496–503.
https://doi.org/10.1021/acssynbio.2c00137.

[22] Shao N, Fan Y, Chou CW, Yavari S, Williams R v., Amster IJ, et al. Expression
of divergent methyl/alkyl coenzyme M reductases from uncultured archaea.
Commun Biol 2022;5. https://doi.org/10.1038/s42003-022-04057-6.

[23] Scheller S. Methyl-coenzyme M reductase Mechanistic studies with ^2H and ^{13}C
labels. ETH Zurich, 2011. https://doi.org/10.3929/ethz-a-006570975.

[24] Walters AD, Smith SE, Chong JPJ. Shuttle Vector System for *Methanococcus*
maripaludis with Improved Transformation Efficiency. Appl Environ Microbiol
2011;77:2549–51. https://doi.org/10.1128/AEM.02919-10.

[25] Takai K, Inoue A, Horikoshi K. Methanothermococcus okinawensis sp. nov., a
thermophilic, methane-producing archaeon isolated from a Western Pacific

481   deep-sea hydrothermal vent system. Int J Syst Evol Microbiol 2002;52:1089–
482   95. https://doi.org/10.1099/00207713-52-4-1089.

483 [26]   Sowers KR, Boone JE, Gunsalus RP. Disaggregation of Methanosarcina spp.
484   and Growth as Single Cells at Elevated Osmolarity. vol. 59. 1993.

485 [27]   Mahlert F, Grabarse W, Kahnt J, Thauer RK, Duin EC. The nickel enzyme
486   methyl-coenzyme M reductase from methanogenic archaea: in vitro
487   interconversions among the EPR detectable MCR-red1 and MCR-red2 states.
488   JBIC Journal of Biological Inorganic Chemistry 2002;7:101–12.
489   https://doi.org/10.1007/s007750100270.

490 [28]   Thomsen J, Weidenbach K, Metcalf WW, Schmitz RA. Genetic Methods and
491   Construction of Chromosomal Mutations in Methanogenic Archaea, 2022, p.
492   105–17. https://doi.org/10.1007/978-1-0716-2445-6_6.

493 [29]   Long F, Wang L, Lupa B, Whitman WB. A Flexible System for Cultivation of
494   *Methanococcus* and Other Formate-Utilizing Methanogens. Archaea
495   2017;2017:1–12. https://doi.org/10.1155/2017/7046026.

496 [30]   Bradford MM. A rapid and sensitive method for the quantitation of microgram
497   quantities of protein utilizing the principle of protein-dye binding. Anal Biochem
498   1976;72:248–54. https://doi.org/10.1016/0003-2697(76)90527-3.

499 [31]   Scheller S, Goenrich M, Mayr S, Thauer RK, Jaun B. Intermediates in the
500   Catalytic Cycle of Methyl Coenzyme M Reductase: Isotope Exchange is
501   Consistent with Formation of a σ-Alkane-Nickel Complex. Angewandte Chemie
502   International Edition 2010;49:8112–5. https://doi.org/10.1002/anie.201003214.

Figures

Figure 1. Anaerobic oxidation of alkane pathway occurred in archaea. The
discontinued box highlights the conversion of alkyl-Coenzyme M to corresponding
alkane catalyzed by Mcr including the heterodisulfide reducing system involving the
utilization of Ti(III) citrate and Co(II)/Co(I).

Figure 2. Influence of the growth phase onto Mcr methane formation activity from

methyl-CoM and Coenzyme B in cell-free lysates prepared from M. acetivorans

supplemented with MeOH.

Figure 3. Comparison of M. acetivorans cell-free lysates enzyme activity under (A) different incubation temperatures (25 and 37 °C) of different harvested at different growth phase (low and high OD$_{600}$); and (B) different vitamin B$_{12}$ concentrations (0.2 mM, 1.1 mM) at different growth phase (low and high OD$_{600}$. Assays were conducted in duplicate under conventional assay conditions (100 µg of protein in cell lysate, 0.2 mM vitamin B12, 0.2 mM B$_{12}$, 37 °C).

A)

B)

Figure 4. Promiscuity and enzyme activity for ethyl-CoM and promiscuity of cell-free
lysates from different methanogens at different conditions: **(A)** conventional
conditions: 37 °C, 0.2 mM B12, 20 mM Ti (III) citrate, and OD600 at harvesting
between 0.3-0.5; **(B)** optimal conditions: 49 °C, 1.1 mM B12, 20 mM Ti(III) citrate,
and OD600 at harvesting between 0.3-0.5. *For M. mazei, due to low protein
concentration in cell lysate, cell was harvested at OD600 between 1.4-1.6.

A)

B)

Figure 5. Comparison of enzyme activity for methyl- and ethyl-Coenzyme at different mixture concentrations employing different cell-free lysate from Mt. marburgensis grown either **(A)** in bioreactor with a continuous flow of growth substrate or **(B)** in serum vial with a periodic fill of growth substrate under high pressure. The experiments were carried out under conventional conditions: 37 °C, 0.2 mM B12, 20 mM Ti (III) citrate, and OD600 at harvesting between 0.3-0.5.

Figure 6. SDS-PAGE of ammonium sulfate-precipitated Mt. marburgensis Mcr

extracts collected from two different growth conditions. The bioreactor and the vial

sample samples were harvested at high- and low-optical density, respectively. The

sizes of the subunits of both Mcr isoforms are: 65 kDa (alpha), 49 kDa (beta), 38 kDa

(gamma of Mcr I), and 33 kDa (gamma of Mcr II).

Table 1. Enzyme activity for methyl- and ethyl-CoM, and the promiscuity of cell-free

lysate under optimal conditions generated by the prediction models.

	Activity methyl-CoM (mU/mg)	Activity ethyl-CoM (mU/mg)	Promiscuity (%)
Prediction value	8.4	1.5	10.9
Replicate 1	16.3	1.37	8.5
Replicate 2	16.7	1.43	8.9
Mean ± S.D.	16.5 ± 0.21	1.43 ± 0.06	8.7 ± 0.2