Modularity of the segmentation clock and morphogenesis

James E. Hammond[1]**, Ruth Baker**[2]**, Berta Verd**[1*]

*For correspondence:

berta.verdfernandez@biology.ox.ac.uk (FMS)

[1]Biology Department, University of Oxford, Oxford, United Kingdom;
[2]Mathematical Institute, University of Oxford, Oxford, United Kingdom

Abstract

Vertebrates have evolved great diversity in the number of segments dividing trunk body features such as the musculature, nervous system, and skeleton, however the developmental origin for this evolvability is poorly understood. The number of segments is determined by the number of somites, transient blocks of mesodermal tissue, that form in the early embryo. These form rhythmically at a rate thought to be determined by the dynamics of a molecular oscillator known as the segmentation clock, and the morphogenesis of the pre-somitic mesoderm (PSM) is thought to control the total length of time over which somites are formed. Here we explore whether the clock and PSM morphogenesis exhibit developmental modularity, as independent evolution of these two processes may explain the high degree of evolvability observed in vertebrate segment number. Using a computational model of the clock and PSM development parameterised for Zebrafish (*Danio rerio*), we find that the clock is broadly robust to variation in morphogenetic processes such as cell ingression, motility, compaction, and cell division. We show that this robustness is in part determined by the length of the PSM and the strength of phase coupling in the clock. As previous studies report no changes to morphogenesis upon perturbing the clock, we conclude based on our study that the clock and morphogenesis of the PSM exhibit developmental modularity.

Introduction

Vertebrates exhibit great diversity in the number of segments that divide the skeleton, musculature, and nervous system of the trunk body (*Richardson et al.* (*1998*)). However, the developmental basis for this evolvability remains poorly understood. Segmentation

of the trunk body is established by transient paired blocks of mesodermal tissue, known as somites, that form periodically on either side of the developing embryo as it elongates posteriorly, in a process known as somitogenesis (*Morin-Kensicki et al.* (*2002*); *Gomez and Pourquié* (*2009*); *Oates et al.* (*2012*)). The total number of segments corresponds to the number of somites formed in the embryo, which is thought to be an emergent property of the morphogenesis of the pre-somitic mesoderm (PSM) and the dynamics of a molecular oscillator known as the segmentation clock (*Morin-Kensicki et al.* (*2002*); *Cooke and Zeeman* (*1976*); *Gomez et al.* (*2008*); *Gomez and Pourquié* (*2009*); *Harima et al.* (*2013*); *Schröter and Oates* (*2010*)). Here, we hypothesise that the evolvability of vertebrate segment number may be underpinned by developmental modularity of the segmentation clock and morphogenesis of the PSM (*Raff* (*1996*)), and that their potential to evolve independently from one another may help to explain the diversity observed in vertebrate segment number.

The segmentation clock is a complex gene regulatory network thought to be driven by cell-autonomous oscillations of transcription factors in the Hes/Her family (*Palmeirim et al.* (*1997*); *Lewis* (*2003*); *Harima et al.* (*2013*); *Schröter et al.* (*2012*); *Webb et al.* (*2016*)), however the degree to which oscillations are truly cell-autonomous is unclear and may vary across species (*Venzin and Oates* (*2020*)). Noisy oscillations are synchronised across cells by delta-notch signalling (*Horikawa et al.* (*2006*); *Lewis* (*2003*); *Venzin and Oates* (*2020*)), creating synchronous travelling waves of gene expression that pulse from the PSM posterior to the anterior. In the anterior PSM somite boundaries are patterned by the interaction of clock oscillations with a posterior-anterior decreasing gradient of FGF signalling that is thought to act as a 'wavefront', reading-out the phase of the clock to create a spatially periodic pattern of somite boundaries (*Cooke and Zeeman* (*1976*); *Soroldoni et al.* (*2014*); *Simsek and Özbudak* (*2018*); *Simsek et al.* (*2023*)). The clock synchronises the differentiation of PSM cells as they adopt somite fates at the anterior of the PSM (*Cooke and Zeeman* (*1976*)), and so controls both the tempo at which somites are formed, and the anterior-posterior polarity of the somites (*Simsek et al.* (*2023*)). Thus, clock synchrony is important for the accuracy of somitogenesis and the frequency of clock oscillations determines the frequency of somite formation, which together with the total duration of the process will be important in determining the total number of somites formed.

Concurrent with somitogenesis, the PSM undergoes elongation. Comparative studies have shown that a diverse range of mechanisms are responsible for the elongation of the PSM (*Gomez et al.* (*2008*); *Bénazéraf et al.* (*2010*); *Steventon et al.* (*2016*); *Mongera*

et al. (*2018*); *Thomson et al.* (*2021*); *Michaut et al.* (*2022*)). Importantly, the elongation dynamics of this tissue are thought to control the total duration of somitogenesis which terminates once the PSM becomes sufficiently short in the AP direction (*Gomez et al.* (*2008*); *Gomez and Pourquié* (*2009*); *Steventon et al.* (*2016*)). Therefore, the total number of segments formed in the developing vertebrate is thought to be the product of both the dynamics of the clock (which controls the rate of somite formation), and the morphogenesis of the PSM, via its control on the total duration of somitogenesis.

Developmental modularity is a property of two or more developmental processes where their uncoupling in space or time permits their evolution independent of one another (*Raff* (*1996*)). This property is thought to give rise to increased phenotypic diversity by enhancing the evolvability of the system (*Raff* (*1996*)). In species examined thus far, such as the Corn Snake *Pantherophis guttatus*, it appears that the evolution of segment number is driven by changes in both the dynamics of the clock and the elongation of the PSM (*Gomez et al.* (*2008*)), and indeed it is possible that independent evolution of these two processes is responsible for the high degree of diversity observed in vertebrate segment number. However, it is not obvious whether the two processes are truly modular, and if not, to which degree. Perturbing the clock's periodicity and function does not appear to affect elongation of the body axis (*Schröter and Oates* (*2010*); *Forero et al.* (*2018*)), so morphogenesis of the PSM is likely to be robust to changes in the segmentation clock through evolution; but many of the cellular- and tissue-level processes which drive PSM elongation could exert an effect on the dynamics of the clock.

For instance, cell rearrangements thought to drive elongation of the PSM in zebrafish (*Danio rerio*) (*Lawton et al.* (*2013*); *Mongera et al.* (*2018*)), and in chicken (*Gallus gallus*) (*Bénazéraf et al.* (*2010*); *Michaut et al.* (*2022*)), promote synchronisation of the segmentation clock (*Uriu et al.* (*2010*, 2013); *Uriu and Morelli* (*2014*); *Uriu et al.* (*2017*)). Additionally arrest of transcription during chromatin condensation is known to cause PSM cells to lag their clock expression relative to neighbours after cell division (*Horikawa et al.* (*2006*); *Delaune et al.* (*2012*)), causing defects in clock synchrony (*Murray et al.* (*2013*)) and suggesting clock synchrony may depend on the degree of proliferative growth in the PSM. The ingression of PSM progenitor cells from dorso-posterior and lateral donor tissues (*Kanki and Ho* (*1997*); *Steventon et al.* (*2016*)) can also create clock asynchrony, as incoming progenitor cells do not appear to show clock gene expression (*Mara et al.* (*2007*)) and thus may be asynchronous with their neighbours once ingressed into the tissue. Finally, tissue convergence movements associated with the elongation of the PSM (*Thomson et al.* (*2021*)) have been proposed to negate the effect of random cell mixing

and be deleterious for clock synchronisation (*Uriu and Morelli* (*2014*)). It is thus non-trivial to determine whether the duration and rate of somitogenesis are modular and in order to do so, one must examine the effect of varying morphogenesis on the dynamics of the clock.

To do this, we use a computational approach to simulate clock dynamics and cell movements within the PSM, and study how the clock responds to changes in morphogenesis of the PSM. We use this approach as it is much quicker and more flexible than experimentally manipulating morphogenesis *in vivo*. As we are constrained by computational complexity and cannot simulate all possible means by which the PSM can elongate (this would require simulation of the growth and dynamics of cells in surrounding tissues which exert forces on the PSM (*Xiong et al.* (*2020*))), we limit our study to simulating regimes of cell movement and growth within the PSM that are known to be associated with, or causal in, PSM growth and elongation across model vertebrates.

For this we use a previously established 3D model of cell movements and clock dynamics within the zebrafish PSM (*Uriu et al.* (*2021*)), and adapt this model to simulate various regimes of cell movement and growth. Briefly (a more detailed description of the model can be found in the methods), the model assumes the PSM is in an inertial frame of reference, with elongation being encoded by the advection of cells towards the anterior PSM (denoted $x = x_a$). Besides advection, cells are subject to random cell mixing, cell-cell repulsion, and a boundary force confining the cells within a hollow horseshoe-shaped domain. This model uses a phase-oscillator model to describe segmentation clock dynamics and each cell is assigned a clock phase which is frozen when the cell exits the PSM at the anterior, crossing the wavefront and patterning a segment. To replenish cells lost at the anterior PSM, new cells are added with random phase and position to keep the tissue at a constant density.

This model is appealing for several reasons, the first being that the model is three-dimensional and so can quantitatively recapture the rates of cell mixing that we observe within the PSM (*Uriu et al.* (*2017*, 2021)). The second is that the segmentation clock is described by an abstract phase-oscillator, broadly agnostic to the molecular details of the segmentation clock which are known to vary between vertebrates (*Krol et al.* (*2011*)). This allows us to make predictions about how the clock responds to changes in morphogenesis in any species which satisfy the basic assumptions of the model while remaining indifferent to the species-specific molecular details (see methods for details). Finally the model makes extensive use of experimentally-derived parameters for cell movements

and tissue dimension, but also more importantly for the phase-oscillator model of the clock (*Uriu et al.* (*2021*)). This allows us to make quantitative predictions of how the zebrafish segmentation clock would react to changes in PSM morphogenesis.

Using this setup we test the clock's response to changes in the position of cell ingression into the PSM, changes in the anterior-posterior profile of random cell motility, changes in the length and density of the PSM, and the presence of mitosis. We find that the clock's synchrony and frequency are robust to these changes except when mitosis is introduced, where we find that synchrony is negatively impacted except where mitosis is confined to the posterior PSM. We find that this robustness is underpinned by the tissue length and cell density in the model, as well as the rate of cell mixing and strength of clock phase coupling. Together, these results suggest that segmentation clock dynamics and PSM morphogenesis are modular components of somitogenesis that can evolve broadly independently from one another, conferring evolvability to this process and helping explain the diversity in vertebral counts across the vertebrates.

Results

Varying the position of cell ingression has minimal effect on clock dynamics

The position of cell ingression into the PSM can vary over the course of development. During somitogenesis new cells enter the PSM posteriorly via ingression from dorsal tissues (*Kanki and Ho* (*1997*); *Banavar et al.* (*2021*)). In the early stages of zebrafish somitogenesis, there is also a contribution of cells from lateral tissues (*Steventon et al.* (*2016*)). As ingressing cells do not appear to express segmentation clock genes (*Mara et al.* (*2007*)), the position at which cells ingress into the PSM can create challenges for clock patterning, as only in the 'off' phase of the clock will ingressing cells be synchronous with their neighbours. Therefore continuous ingression of PSM progenitor cells is likely to create local asynchrony of oscillations where it occurs, and so varying the position of cell ingression is likely to have an effect on anterior clock dynamics.

To test this hypothesis, we alter how cells are added to the tissue in the model. To model the ingression of cells from dorsal tissues into the posterior tailbud (*Banavar et al.* (*2021*); *Kanki and Ho* (*1997*)), we define a density-dependent cell addition process where cells are added onto the dorsal surface of the posterior PSM domain if the total average density in this domain falls below ρ_0, with a constant phase $\theta = 0$. To maintain density in the rest of the PSM, we found it necessary to add cells in the two cylindrical PSM do-

mains as well. To avoid bias we spawn cells at random positions within the domain if the average density falls below ρ_0. These cells are initialised with a random phase drawn from the uniform distribution $[0, 2\pi)$. To model the early stages of Zebrafish somitogenesis, where concurrent with dorso-posterior ingression of cells the PSM also experiences a contribution of cells from lateral tissues, we modify the above simulation so that cell density in the lateral cylindrical domains is maintained by addition of cells with constant phase ($\theta = 0$) on the ventral surface of these two lateral domains. These two ingression scenarios are termed 'Dorso-Posterior' ('DP') and 'Dorso-Posterior + Lateral-Ventral' ('DP+LV') ingression (figure 1A), respectively, and for brevity we refer to them as 'DP' and 'DP+LV' in the rest of the text. We compare the effect of these scenarios of cell ingression with 'Random' ingression (figure 1A), where cell density is maintained by addition of cells at random positions within the tissue with random phase ($\theta \in [0, 2\pi)$).

Figure 1. The effect of cell ingression position on clock frequency and synchrony. A Diagram highlighting the position of cell addition across the three ingression scenarios tested here. Magenta shading shows where cells are added onto the tissue surface. The pale magenta and blue planes respectively correspond to the anterior limit of the PSM, $x = x_a$, and the anterior limit of cell addition, $x = x_d$. In the 'Random' condition cells are added at random positions within the PSM posterior to the plane $x = x_d$, and accordingly no magenta surface is shown. In the dorso-posterior case ('DP') cells are added in this manner in the two lateral cylinders to maintain density, and no magenta surface is shown there. In the dorso-posterior + lateral-ventral ('DP+LV') case cells are only added on the tissue surface at the positions shown by magneta shading. **B** Oscillation synchrony (r) at the PSM anterior after 1000 mins, for the three tested scenarios of cell ingression. N=100 simulations. **C** Mean frequency of oscillations for cells at the PSM anterior after 1000 mins of simulation, for the three tested scenarios of cell ingression. N=100 simulations. **D** Kymographs of synchrony along the x-axis on the left-hand side of the PSM for single simulations from each of the three scenarios of cell ingression tested. Black arrowheads highlight strongly asynchronous populations of cells being transported to the tissue anterior by advection.

We test each of these scenarios by initialising each simulation with a synchronous initial condition for the clock and simulating cell movements and clock dynamics in the presence of each cell ingression scenario for 1000 mins (see methods for further justification). After 1000 mins we measure the synchrony (r) and mean frequency ($\frac{d\theta}{dt}$) of a cylindrical region of tissue at the left-hand PSM anterior, one cell diameter (d_c) in length (see methods). We restrict our measurement to the PSM anterior as it is here that the clock patterns somites (*Simsek and Özbudak* (*2018*); *Simsek et al.* (*2023*)), and thus dynamics elsewhere in the tissue are unlikely to present a phenotype in the animal. Using this method, we find that DP ingression has no effect on synchrony at the PSM anterior after 1000 mins when compared to Random ingression, but that DP+LV ingression creates a minor decrease in anterior synchrony (figure 1B). However, in neither scenario is the anterior frequency changed when compared to Random ingression (figure 1C). Kymographs of synchrony along the anterior-posterior axis reveal that cell addition onto the tissue surface creates asynchronous populations of cells that travel towards the tissue anterior with cell advection (figure 1 D, black arrowheads). These are more frequent than in the DP case, likely explaining the lower anterior synchrony observed in the DP+LV simulations.

The clock is robust to changes in mode of cell ingression regardless of the cell motility profile

As cell mixing promotes clock synchrony (*Uriu et al.* (*2010*, 2013); *Uriu and Morelli* (*2014*)), and an anterior-posterior profile of increasing cell mixing is present in the PSM of zebrafish and chicken (*Kanki and Ho* (*1997*); *Bénazéraf et al.* (*2010*); *Lawton et al.* (*2013*); *Mongera et al.* (*2018*); *Michaut et al.* (*2022*)) (and presumably most vertebrate taxa), we speculated that the shape of this gradient may confer robustness to clock dynamics against cell ingression. For instance, a steep profile (e.g. as reported from the early somitogenesis stages of zebrafish (*Mongera et al.* (*2018*))) might confer robustness to lateral ingression as mixing is higher along a greater length of the PSM than a more graded profile (e.g. as reported from chicken embryos (*Bénazéraf et al.* (*2010*); *Michaut et al.* (*2022*))). Therefore the shape of the cell motility profile in the PSM may constrain the evolution of cell ingression position, and vice versa.

Figure 2. Effect of cell motility profile on clock frequency and synchrony. A Overview of how intrinsic cell motion is encoded in the model. Each cell is given a random direction vector $v_0(\mathbf{x}_i)\mathbf{n}_i(t)$ (black arrows) whose magnitude $v_0(x)$ increases towards the PSM posterior. **B** Magnitude of intrinsic cell motion $v_0(x)$ for the parameters used by Uriu *et al.* (*Uriu et al.* (*2021*)) (blue), and how the shape of the function can change when increasing (yellow) or decreasing (green) the inflexion point and curve steepness parameters, X_v and h respectively. **C** Clock synchrony r (**top**) and difference from expected mean frequency $\Delta\frac{d\theta}{dt}$ (**bottom**) for different $v_0(x)$ specified by combinations of X_v and h. The corresponding pixels display the synchrony or frequency at the PSM anterior after 1000 mins of simulation using the specified motility profile, averaged across $N = 100$ simulations. Parameter ranges used are $X_v \in \{0, 0.1, 0.2, \dots, 1\}$ and $h \in \{0, 0.5, 1, \dots, 5\}$.

To test this, we systematically enumerated the steepness h and inflexion point X_v of the function $v_0(x)$ (figure 2B), which controls the speed of random cell movement in the PSM (see methods, figure 2A). For each parameter pair we performed $N = 100$ simulations and recorded the median anterior synchrony and median anterior mean frequency (see methods) after 1000 mins. The results for each ingression scenario are shown in figure 2C. We find no trend in median anterior synchrony and frequency across different parameter pairs for Random ingression, DP ingression, or DP+LV ingression (figure 2C). Similar to before, results for DP ingression are indistinguishable from Random. For DP+LV ingression, anterior synchrony is lower across all parameter sets than DP or Random ingression (figure 2C) but the effect is still very minor in this case. We therefore predict that at least with respect to effects on the zebrafish segmentation clock, that the evolution of the motility profile and the position of cell ingression are not constrained by one another.

Length and density of the PSM confer robustness to the clock

The PSM is a transient tissue that shrinks in length over time. While this feature is conserved across vertebrate species, the absolute length and rate of shrinkage is known to vary between taxa (*Gomez et al.* (*2008*); *Steventon et al.* (*2016*); *Thomson et al.* (*2021*)). We postulate that PSM length is an important factor in determining the clock's response to the different morphogenetic modes; for example, intuitively cells will have more time to synchronise oscillations with their neighbours before reaching the PSM anterior in a longer PSM, or if cell ingression is spread out along a longer tissue the degree of asynchrony imparted by ingression will be lessened. Similarly, we predict that the density of the tissue confers robustness to clock dynamics as increasing the average number of neighbours for each cell may act to 'correct' against clock noise. Overall tissue density has been observed to increase over time in zebrafish (*Thomson et al.* (*2021*)), and has been observed to vary in space in chicken, with density decreasing towards the PSM posterior (*Bénazéraf et al.* (*2010*); *Michaut et al.* (*2022*)).

We thus sought to explore whether length and density confer robustness to changing morphogenesis. We test this by co-varying the position of cell ingression (figure 1A) with either tissue density or length. When varying the average tissue density ρ_0 and studying model dynamics after 1000 mins we see that increasing cell density marginally increases anterior synchrony (figure 3B), and decreases the variability in frequency at the PSM anterior (figure 3C). However the increase in density does not appear to be sufficient to 'correct' the decrease in synchrony in the DP+LV case (Fig.3B). Increasing the length of the tissue does however appear to 'correct' this decrease, after 1000 mins of simulation

(figure 3E). Here the PSM length L is increased but the anterior limit of cell division x_d is held constant at $x_d = 100 \ \mu m$. Anterior synchrony after 1000 mins positively correlates with L and decreasing L reveals a decrease in anterior synchrony for DP ingression (figure 3E). No obvious trend for mean anterior frequency is observed (figure 3F). Note that the results for the DP and DP+LV case are exactly equal for $L = 185 \ \mu m$, as for $x_d = 100 \ \mu m$ cell addition is confined to the dorsal-posterior domain and the two ingression methods are numerically equivalent in the model (see methods). While much of this recovery in synchrony may be due to cell addition occuring along a longer domain (and in so doing 'diluting' the effect of cell ingression), we observe similar results when cell addition can only occur along a fixed length of the tissue and the total length of the tissue varies (figure 3 supp. 1). Here, the anterior limit of cell addition x_d is fixed by $x_d = X_c - 100$, and we observe a recovery in synchrony across all three cell ingression scenarios as the total length of the tissue increases (figure 3 supp. 1).

Figure 3. Effect of tissue density and length. A Stills from exemplar simulations illustrating changes in tissue density. **B** Anterior synchrony after 1000 mins for changing tissue density ρ_0 and varying position of cell ingression. N=100 simulations. **C** Anterior mean frequency after 1000 mins for changing tissue density ρ_0 and varying position of cell ingression. N=100 simulations. **D** Stills from exemplar simulations illustrating changes in tissue length. **E** Anterior synchrony after 1000 mins for changing tissue length L and varying position of cell ingression. N=100 simulations. **F** Anterior mean frequency after 1000 mins for changing tissue length L and varying position of cell ingression. N=100 simulations.

However in zebrafish density and length are known to co-vary after the 16-somite stage, with the PSM increasing in density as it shrinks in length (*Thomson et al.* (*2021*)). It is not clear whether the increase in density in the PSM is sufficient to compensate for the shrinking in PSM length towards the end of somitogenesis. Here we test this by shrinking the PSM radius r and length L, while increasing its density ρ_0, over time, using published rates and initial conditions for these values (figure 4A-B, methods) (*Thomson et al.* (*2021*)).

We also decrease the cell diameter d_c as has been observed in zebrafish (**Thomson et al.** (**2021**)), as decreasing cell radius may counteract the increase in tissue density by reducing the average number of neighbours for each cell in the tissue (defined as those cells whose nuclei are within a cell's diameter of each cells).

As these changes are dynamic, we depart from previous methodology for measuring dynamics and plot the anterior synchrony and frequency over time for N=100 simulations. For small PSM length and width the model generates voids in the tissue where cells do not exit the toroid domain at a sufficient rate to maintain tissue homogeneity, which we deem to be biologically inaccurate. Therefore, to exclude such cases we only plot synchrony and frequency up until the time at which one of the N=100 replicate simulations encounters such a gap. Thus it is important to note that the simulations presented here do not cover the whole timespan of zebrafish somitogenesis, and we cannot examine dynamics towards the end of the process.

The anterior synchrony and mean anterior frequency over time for N = 100 simulations are shown in figures 4C and 4D respectively. They are compared with simulations that do not shrink the PSM and retain the pre-16-somite values for L, r, d_c, and ρ_0. We find that towards the end of these simulations the clock experiences more noise and fluctuations in synchrony and frequency (figure 4C-D) than the non-shrinking equivalent simulations, however for the majority of the duration of simulation the dynamics of the compacting tissue are broadly comparable with those of the non-compacting tissue, but with synchrony minorly decreasing and frequency becoming noisier towards the end of the simulation (figure 4C-D). We treat the effects observed towards the end of the simulation with caution in light of the irregularities in tissue structure mentioned above. In addition, other model parameters that are fixed throughout the simulation, such as advection velocity or intrinsic frequency, are thought to change towards the end of somitogenesis (**Schröter et al.** (**2008**); **Uriu et al.** (**2021**)), and therefore for the latter stages of somitogenesis the results of our simulations may be inaccurate. However, data for the final stages of zebrafish somitogenesis is extremely limited and we cannot make estimates for model parameters at this stage. Therefore we conclude that for much of somitogenesis in zebrafish the clock is not majorly impacted by the compaction-extension of the tissue, but that there may be effects at the terminal stages of somitogenesis undetected by this model.

Figure 4. Effect of compaction-extension on clock synchrony and frequency. A Snapshots of an exemplar simulation showing how the PSM shrinks in length and diameter as time progresses. **B** Functions for PSM length L, radius r, density ρ, and cell diameter d_c, derived from (**_Thomson et al._** (**_2021_**)) (yellow), and the constant functions (blue) with which the effect of these functions is compared. **C** Anterior synchrony and **D** mean anterior frequency over time, for N=100 simulations. The solid line indicates the median and the inter-quartile range is given by a shaded band either side of this line. Blue shows simulations where the tissue does not undergo compaction-extension after t_{shrink}, and yellow shows the results for simulations where after t_{shrink} the tissue undergoes compaction-extension according to the functions shown in **B**. Results are plotted until the time at which at least one of the replicate simulations encounters a gap in the tissue at the tissue anterior (see methods).

Beyond dynamic changes in density, we also attempted to encode a spatial gradient of decreasing cell density towards the posterior, as reported from chicken (**_Bénazéraf et al._** (**_2010_**); **_Michaut et al._** (**_2022_**)). However, we could not generate a gradient that quantita-

tively recaptures that reported in (*Bénazéraf et al.* (*2010*)) (data not shown). Based on our results for changing average cell density ρ_0 it is highly probable that this gradient would affect clock dynamics. We suggest that another model, such as one encoding cell-cell adhesion, may be a more powerful tool for exploring the effect of this gradient, and dynamics at the terminal stages of somitogenesis.

Clock arrest during cell division creates asynchrony

During M phase of the cell cycle segmentation clock gene expression in zebrafish is known to arrest (*Horikawa et al.* (*2006*)), causing a lag in nascent daughter cells relative to their neighbours (*Delaune et al.* (*2012*)). If cell divisions are asynchronous, as appears to be the case in zebrafish (*Kanki and Ho* (*1997*); *Steventon et al.* (*2016*)), then such lags can create asynchrony of oscillations in the tissue (*Murray et al.* (*2013*)).

As previous models of mitosis and the segmentation clock have been 2D and may not quantitatively recapture the cell mixing and geometry present in zebrafish (*Murray et al.* (*2013*, 2019)), we sought to investigate the effect of mitosis in the presence of the more accurate 3D geometry and cell mixing in the present model. To do so, we simulated cell division by assigning each cell a cell cycle phase τ that increases at a constant rate of $1\ \text{min}^{-1}$, and spawning a new cell adjacent to that cell once that its $\tau \geq T_G + T_M$, where T_G denotes the cumulative time to complete G1, S, and G2 phases of the cell cycle, and T_M denotes the time spent in M phase. Once a new cell is spawned, the τ for both cells is reset to 0. As daughter cells share the same clock phase after division *in vivo* (*Delaune et al.* (*2012*)), the new spawned cell shares the same clock phase θ as its sibling. To simulate the arrest of the clock during M phase, if $\tau \in [T_G, T_G + T_M)$ oscillations cease, i.e. $\frac{d\theta_i}{dt} = 0$. For further details, we refer the reader to the methods section below.

To avoid measuring trivial changes in frequency and synchrony due to arrest of the clock, we measure synchrony and frequency of the clock only in cells where $\tau < TG$. Therefore our results represent a lower bound on the effect of mitosis on the clock. Using the estimated values of T_G and T_M for Zebrafish (see methods for calculation), we find that the presence of cell division creates asynchrony as reported previously (*Murray et al.* (*2013*)), and has no identifiable effect on frequency (figure 5B-C). Fixing $T_G + T_M = 187.5$ mins and varying T_M shows a T_M-dependent trend where synchrony decreases with increasing T_M (figure 5E). After $T_M = 15$ mins the median value for synchrony recovers but there is increased noise (figure 5E). No noticeable trend is observed for frequency (figure supp 1A).

In previous simulations (figure 5), cell division is permitted to occur right up to the PSM anterior where segments are pre-patterned. However if division is confined to the posterior (**Bouldin et al.** (**2014**)) it may be possible that cells have time to re-synchronise before reaching the wavefront. To test whether this is possible within the timescales of zebrafish cell movement, division, and clock dynamics, we define a point in space $x = x_a + x_{div}$ anterior to which (i.e. for $x < x_a + x_{div}$) we halt cell cycle progression by fixing $\tau = 0$ mins in all cells. Simulating for $x_{div} = 25\mu m, 175\mu m, 325\mu m$ we see that while for $x_{div} = 25\mu m$ we observe defects in synchrony, for the more posterior values synchrony is rescued (figure 5D). This suggests that there exists some point along the PSM posterior to which cell division can occur without affecting anterior dynamics of the clock. To try and resolve the value of this point and how it depends on the length of M-phase T_M, we systematically enumerate values of x_{div} from the tissue anterior to the tissue posterior, and T_M across the values tested in Fig.5E. We find that for $T_M < 15$ mins division can occur up to the tissue anterior without a major defect in synchrony (figure supp 2A). For $T_M > 15$ mins division can also occur close to the tissue anterior, though not so close as for $T_M < 15$ mins (figure supp 2A). Only for $T_M = 15$ mins is the anterior limit of cell division significantly displaced posteriorly (figure supp 2A). This result is perhaps intuitive, as the parameters in the model set the clock period to $\frac{2\pi}{\omega_0} = 30$ mins, and a clock arrest lasting $T_M = 15$ mins would be sufficient to move a cell into antiphase relative to its neighbours. Indeed we see that mitosis is most deleterious to synchrony when M-phase lasts approximately half the length of a clock cycle, however for long clock cycles this trend becomes nonlinear and the most deleterious values of T_M lie between the length of one or two clock cycles (figure 5 supp.2B).

Figure 5. Effect of cell division on clock frequency and synchrony. A Diagram showing how during cell division, clock phase θ arrests, causing a cell to fall out of phase with its neighbour. On the right hand side a still from a simulation for T_M = 15 mins is shown, illustrating how this creates asynchrony of oscillations. The effect of cell division on anterior synchrony and frequency is shown in figures **B** and **C**, respectively, for $T_M = 15$ mins after 1000 mins. $N = 100$. **D** The effect on anterior synchrony when division is restricted to only occur posterior to $x = x_{div}$, for $T_M = 15$ mins. $N = 100$. **E** The effect on anterior synchrony after 1000 mins when T_M is varied. In each case, the total length of the cell cycle is maintained at a constant length, i.e. $T_M + T_G = 187.5$ mins. $N = 100$. To rule out trivial changes in synchrony and frequency, in all analysis here we restrict measurement to non-dividing cells, i.e. cells such that $\tau \in [0, TG)$.

Zebrafish clock coupling and cell mixing confers robustness to changes in cell ingression

Across many of the morphogenetic scenarios tested here, we have observed a high degree of robustness in clock dynamics with respect to changing cell movements and pro-

cesses. As shown above, some of this robustness can be attributed to the choice of tissue length L in simulations, and choices of the parameters v_a and v_p governing the advection velocity of cells. However we would also expect that the rate of cell mixing (**Uriu et al.** (**2010**, 2013, 2017)), and the strength of clock coupling confer robustness too. Within the present model, the global rate of cell mixing is controlled by the parameter v_s and the clock coupling strength is denoted by κ. Notably, the values taken for these parameters have been experimentally measured in zebrafish (**Riedel-Kruse et al.** (**2007**); **Herrgen et al.** (**2010**); **Uriu et al.** (**2017**, 2021)). Therefore we were interested in where these parameters lie relative to the space of parameter pairs that generate clock dynamics robust to changes in morphogenesis - do these results reflect something unique about the zebrafish parameters or can many (κ, v_s) pairs exhibit robustness?

Figure 6. Anterior synchrony after 1000 mins, for varying maximum magnitude of intrinsic cell motion v_s and clock phase coupling strength κ, for three different scenarios of cell ingression. Each pixel corresponds to the median value of anterior synchrony across N = 100 simulations. A black + marks the experimental values for Zebrafish, $v_s = 1 \ \mu m \cdot min^{-1}$, $\kappa = 0.07 \ min^{-1}$, that are used elsewhere in this paper. All other parameters are held constant at their normal values (see table 1).

To determine this, we take changing cell ingression as an example of varying morphogenesis and for each ingression scenario, systematically enumerate the parameters v_s and κ, studying the anterior synchrony after 1000 mins with N=100 replicates per parameter pair. This reveals a threshold curve of (κ, v_s) pairs below which the clock tends to asynchrony (figure 6). Comparing across the three scenarios of cell ingression we see that the position of this curve changes with the cell ingression position (figure 6), indicating that the space of (κ, v_s) pairs that achieve clock synchrony is constrained by the cell ingression mode present within the tissue. Notably the experimentally-derived parame-

ter pair ($\kappa = 0.07$ min^{-1}, $v_s = 1$ μm · min^{-1}) lies very close to the threshold for the DP+LV case of cell ingression in the κ direction, suggesting that much of the robustness we have observed in response to changes in cell ingression is sensitive to the choice of κ. However in each case our experimental choices for κ and v_s lie above the threshold, implying that at least part of the robustness we observed with respect to cell ingression is due to the experimental values for zebrafish being sufficiently strong to confer this robustness.

Discussion

Here we have tested a broad range of PSM cellular behaviours and mechanisms that are associated with the elongation of the developing PSM across vertebrate species. We predict that the dynamics of the zebrafish segmentation clock, specifically synchrony and frequency, are generally robust to changes in the local processes of PSM morphogenesis such as cell ingression, motility, and, under certain conditions, division. However clock dynamics show a dependence on global processes of morphogenesis such as elongation rate (equivalent to cell advection velocity) and tissue length.

Our results suggest that a major determinant of this robustness is the rate of clock coupling κ, which in zebrafish appears to be sufficiently strong to confer robustness to changes in cell ingression (figure 6). While synchrony of the clock shows a dependence on cell mixing rate (v_s) and tissue length (L), the dependence on length is weak compared to its dependence on coupling strength (supplementary figure 1), suggesting that this robustness may be more dependent on properties of the segmentation clock than those of the tissue. Nevertheless, there is a dependence of clock synchrony on these global processes and further work is needed to elucidate the relationship between the local processes of morphogenesis driving axial elongation, their effect on tissue length and elongation rate, and the effect on clock dynamics. It is possible that the rate of axial elongation generated by each process, and the corresponding changes in PSM length, may differ, and the robustness of the clock to each process may be less than we present here. This is something we cannot test with the present model formulation, which cannot generate elongation or deform the tissue geometry, and we suggest that the development of more sophisticated and computationally complex models capable of simulating those behaviours will be beneficial in this area.

It is also worth noting that while in the model formulation κ represents the strength of delta-notch signalling between cells, the experimental value for this parameter is measured at a tissue level, incorporating coupling effects from other pathways e.g. cell mixing (*Riedel-Kruse et al.* (*2007*)). It is therefore possible that our choice of κ represents an over-

estimate of the strength of delta-notch signalling and when we combine this with cell mixing in the model, our results present an over-estimate of robustness. However, the precise value of r at which a somite is correctly patterned is unknown, and indeed there exist processes independent of the segmentation clock that can act to correct against inaccuracies in somite length (*Naganathan et al.* (*2022*)), so it is plausible that somitogenesis may be more tolerant of clock asynchrony than we suppose here and that somites are correctly patterned for $r < 1$. As our results appear less sensitive to cell mixing (v_s) than κ (figure 6), we suggest that the contribution of cell mixing to an experimental estimate of κ may be minor. Therefore, combined with a tolerance of somitogenesis for values of $r < 1$, we suggest that our findings are robust to any overestimates of κ incorporating cell mixing.

Our analysis is restricted to the modes of PSM morphogenesis that are known to occur in vertebrates, and we cannot rule out the possibility that the robustness we observe merely reflects evolution of mechanisms of PSM elongation under selective pressure to minimise the effect on the clock. Nevertheless, we argue that there exists a qualitatively distinct set of processes that underpin PSM elongation that the clock may be robust to. These processes are qualitatively different and employed by different species to differing degrees, suggesting that evolution alters these processes in parallel to achieve diversity in PSM elongation dynamics. We suggest that this can be achieved due to the robustness of clock dynamics conferred by cell-cell coupling, cell mixing, elongation rate, and PSM length.

We suggest that as clock dynamics appear to be robust to changes in morphogenesis, the clock and morphogenesis of the PSM comprise two developmental modules within zebrafish, and this is true more generally whenever the phase coupling of a clock is sufficiently strong or the PSM sufficiently long. We have no reason to expect that these properties are uniquely satisfied by zebrafish development and indeed this modularity may have facilitated the evolution of diversity in segment number in vertebrates. Further study of the morphogenesis of the PSM in other vertebrate species, and quantification of the dynamics of their segmentation clocks (*Riedel-Kruse et al.* (*2007*)), will reveal whether this modularity is indeed a general property of vertebrate axial development. We note that the evolution of modularity is an active area of research (*Wagner and Altenberg* (*1996*); *Wagner et al.* (*2007*)), and suggest that studying the evolution of delta-notch signalling in vertebrates, and the evolutionary origin of the pre-somitic mesoderm and somitogenesis, may be an illuminating model system for this field. We predict that the modularity of the clock and morphogenesis of the PSM permits a synergy between

these two processes that heightens the evolvability of segment number in vertebrates, and that this may be responsible for the high diversity we observe in this trait (***Richardson et al.*** (***1998***)).

Methods

Computational Model of the Segmentation Clock and PSM

To simulate the movements of cells within the PSM and the dynamics of the segmentation clock, we used the model of ***Uriu et al.*** (***2021***), who sought to explain defective somite patterning after loss of Notch-Delta signalling (***Uriu et al.*** (***2021***)). Here an abstract phase oscillator kuramoto model is used to describe intracellular clock oscillations and cell-cell phase coupling. Cells are confined within a 3-D tissue domain whose geometry approximates that of the PSM.

This model is suitable for our purposes because it provides a three-dimensional model of cell movements within the PSM, allowing us to alter morphogenesis in order to test hypotheses. Additionally as the segmentation clock here is described as an abstract phase oscillator, the model has broad applicability to different vertebrate species which are known to differ in the structure and regulation of their segmentation clock gene regulatory networks (***Krol et al.*** (***2011***)). Finally, the validity of the model has been shown in several ways from comparison with data for zebrafish (***Webb et al.*** (***2016***); ***Liao et al.*** (***2016***); ***Uriu et al.*** (***2021***)), and uses parameters experimentally inferred from Zebrafish embryos (***Riedel-Kruse et al.*** (***2007***); ***Soroldoni et al.*** (***2014***); ***Uriu et al.*** (***2021***)). This gives us an accurate read-out of the effect of changing morphogenesis on the segmentation clock of a vertebrate.

The model is described elsewhere (***Uriu et al.*** (***2021***)), but for completeness we include an account of it here. For clarity, vector variables are shown in bold, and scalar variables in normal font.

Tissue Geometry and frame of reference

The model assumes that the posterior tip of the tailbud is held in an inertial frame of reference, and models the posterior-ward elongation of the PSM by movement of cells towards the tissue anterior.

Cells are confined within a horseshoe-shaped domain resembling the shape of the PSM. This domain is comprised of two cylindrical sub-domains and one half-toroid subdo-

main, each representing the lateral sides of the PSM and tailbud respectively. A schematic can be seen in figure 7. All domains are open-ended, and cells are free to move between them.

Figure 7. Top: schematic of the PSM in the xy plane. The PSM is comprised of two cylinders, centred at $y = r$ and $y = 2R + r$ respectively, with radius r. The 'Tailbud' is represented as a half-torus domain centred at $\mathbf{x} = (X_c, Y_c, Z_c)^T$, with internal radius r and radius from torus centre R. Cross-sections of the PSM and Tailbud are highlighted in magenta, and illustrated on the **bottom**, showing how a point within the tissue $\mathbf{x}_i$ is assigned the polar coordinates r_i and q_i.

The centre of the toroid domain is denoted by the coordinate vector $(X_c, Y_c, Z_c)^T$, and the anterior of the PSM is demarcated by the x-coordinate $x = x_a$ (see figure 7). The major radius of the torus is denoted by R and the minor radius (and the radius of the two cylindrical domains) is denoted by r (see figure 7). The resulting length of the PSM in the x direction, L, is therefore given by $L = r + R + X_c - x_a$.

Cell Movement

To approximate cell mixing and the advection of cells out of the PSM towards the anterior (in the chosen tailbud-inertial frame of reference), cell movement in the model is described by an equation of motion that incorporates cell advection, cell-cell repulsion, random cell motion, and a boundary force confining cells within the tissue domain (Equation 1).

Specifically, each cell is assigned a position vector $\mathbf{x}_i \in \mathbb{R}^3$, where $\mathbf{x}_i = (x_i, y_i, z_i)^T$. This position $\mathbf{x}_i$ corresponds to the cell centre. Cells are assumed to be spheres with diameter d_c. The equation of motion for cell i is

$$\frac{d\mathbf{x}_i}{dt} = \mathbf{v}_d(x_i) + v_0(x_i)\mathbf{n}_i(t) + \sum_{j=1, j \neq i}^{N} \mathbf{F}(\mathbf{x}_i, \mathbf{x}_j) + \mathbf{F}_b(\mathbf{x}_i) \tag{1}$$

where $\mathbf{v}_d(x_i)$ represents the advection of the cell towards the PSM anterior, $v_0(x_i)\mathbf{n}_i(t)$ the intrinsic random motion of the cell, $\mathbf{F}(\mathbf{x}_i, \mathbf{x}_j)$ the repulsion force between cell i and an adjacent cell j, and $\mathbf{F}_b(\mathbf{x}_i)$ the boundary force confining the cell within the tissue.

The advection velocity $\mathbf{v}_d(x_i)$ is given by

$$\mathbf{v}_d(x_i) = (-v_d(x_i), 0, 0)^T \tag{2}$$

where the function $v_d(x_i)$ describes the local strain rate along the AP axis:

$$v_d(x_i) = \begin{cases} -\frac{v_a - v_p(1-x_q)}{x_q} \cdot \frac{x_i - x_a}{L} + v_a & \frac{x_i - x_a}{L} \leq x_q \\ \\ -\frac{v_p(x_i - x_a)}{L} + v_p & \frac{x_i - x_a}{L} > x_q \end{cases} \tag{3}$$

The model assumes that the random cell motility observed within the PSM is intrinsic, i.e. cells would exhibit random motility in isolation from the tissue. This is modelled using the random evolution of the direction vector $\mathbf{n}_i(t)$, which evolves according a random walk on the surface of the unit sphere specified by the following differential equation:

$$\frac{d\mathbf{n}_i}{dt} = \sqrt{2D}\xi_{\phi_i}(t)\mathbf{m}_x + \sqrt{2D}\xi_{\varphi_i}(t)\mathbf{m}_y - 2D\mathbf{n}_i(t) \tag{4}$$

where the orthogonal vectors (for $\mathbf{e}_z = (0, 0, 1)^T$)

$$\mathbf{m}_x = \frac{\mathbf{n}_i(t) \times \mathbf{e}_z}{|\mathbf{n}_i(t) \times \mathbf{e}_z|}, \quad \mathbf{m}_y = \frac{\mathbf{m}_x \times \mathbf{n}_i(t)}{|\mathbf{m}_x \times \mathbf{n}_i(t)|} \tag{5}$$

together define a plane tangent to the unit sphere, and $\xi_{\phi_i}(t)$, $\xi_{\varphi_i}(t)$ are white gaussian noise terms that satisfy, for any t, t', $\langle\xi_{\phi_i}(t)\rangle = \langle\xi_{\varphi_i}(t)\rangle = \langle\xi_{\phi_i}(t)\xi_{\varphi_j}(t)\rangle = 0$, and $\langle\xi_{\phi_i}(t)\xi_{\phi_j}(t)\rangle = \langle\xi_{\varphi_i}(t)\xi_{\varphi_j}(t)\rangle = \delta_{ij}\delta(t - t')$. A derivation for equation 4 can be found in *Uriu et al.* (*2021*).

To model the increase in random cell motion towards the PSM posterior, the the magnitude of intrinsic random cell movement, $v_0(x)$, increases towards the posterior according to equation 6:

$$v_0(x_i) = \frac{v_s}{1 + (\frac{1 - \frac{x_i - x_a}{L}}{X_v})^h}$$

(6)

The parameters v_s, X_v, and h determine the maximum magnitude of intrinsic cell motion, profile inflexion point, and steepness of the profile, respectively. By varying these parameters one can create a variety of motility profiles with different shapes and amplitudes (see figure 2B). Here the x-position of cell i is normalised according to the length of the PSM L, and so the profile of cell motility scales with the length of the PSM. This is a valid assumption as the length of the PSM is thought to be partially specified by FGF signalling (*Simsek and Özbudak* (*2018*)), and the random motility of PSM cells is also under the control of FGF (*Bénazéraf et al.* (*2010*)).

Cell-cell repulsion is encoded by repulsion of two cells if their centres are within a cell's diameter d_c of one another, according to the cell-cell repulsion force $\mathbf{F}(\mathbf{x}_j - \mathbf{x}_i)$:

$$\mathbf{F}(\mathbf{x}_j - \mathbf{x}_i) = F(\mathbf{x}_j - \mathbf{x}_i)\frac{\mathbf{x}_j - \mathbf{x}_i}{|\mathbf{x}_j - \mathbf{x}_i|}$$

(7)

where the magnitude of the cell-cell repulsion force $F(\mathbf{x}_j - \mathbf{x}_i)$ is given by

$$F(\mathbf{x}_j - \mathbf{x}_i) = \begin{cases} \mu(\frac{|\mathbf{x}_j - \mathbf{x}_i|}{d_c} - 1) & |\mathbf{x}_j - \mathbf{x}_i| \leq d_c \\ 0 & |\mathbf{x}_j - \mathbf{x}_i| > d_c \end{cases}$$

(8)

and the parameter μ controls the strength of cell-cell repulsion throughout the tissue. For computational simplicity, cell-cell adhesion is not encoded within this model.

The form of the boundary force $\mathbf{F}_b(\mathbf{x}_i)$ varies depending on whether the cell i is in one of the lateral PSM cylindrical domains or in the half-toroid tailbud (figure 7). If cell i lies within the cylinders, one can use the cylindrical coordinates for cell i, r_i and q_i (see figure 7) to naturally define the boundary force:

$$\mathbf{F}_b(\mathbf{x}_i) = \begin{pmatrix} 0 \\ -\mu_b e^{-\frac{\delta y}{r_b}} \cos q_i \\ -\mu_b e^{-\frac{\delta z}{r_b}} \sin q_i \end{pmatrix}$$

(9)

where

$$\delta y = |(r - r_i)\cos q_i|$$

(10)

$$\delta z = |(r - r_i)\sin q_i|$$

(11)

The magnitude of the boundary force is denoted by μ_b and the parameter r_b determines the length-scale of the boundary force. In the half-toroid tailbud the position $\mathbf{x}_i$ of each cell i can be described according to the coordinates r_i, p_i, and q_i:

$$\mathbf{x}_i = \begin{pmatrix} X_c + R\cos p_i + r_i \cos p_i \cos q_i \\ Y_c + R\sin p_i + r_i \sin p_i \cos q_i \\ Z_c + r_i \sin q_i \end{pmatrix} \tag{12}$$

One can then naturally define

$$\delta x = |(r - r_i)\cos p_i \cos q_i| \tag{13}$$

$$\delta y = |(r - r_i)\sin p_i \cos q_i| \tag{14}$$

$$\delta z = |(r - r_i)\sin q_i| \tag{15}$$

describing the distance of cell i from the tissue surface in the x, y, and z directions respectively. The boundary force then becomes

$$\mathbf{F}_b(\mathbf{x}_i) = \begin{pmatrix} -\mu_b e^{-\frac{\delta x}{r_b}}\cos p_i \cos q_i \\ -\mu_b e^{-\frac{\delta y}{r_b}}\sin p_i \cos q_i \\ -\mu_b e^{-\frac{\delta z}{r_b}}\sin q_i \end{pmatrix} \tag{16}$$

Note that in both definitions of $\mathbf{F}_b(\mathbf{x}_i)$, the tissue domains are open-ended, i.e. cells are free to move between subdomains, and out of the PSM at the anterior.

For further details we urge the reader to consult *Uriu et al.* (*2021*).

Cell Addition

As cells are removed from the tissue by PSM, new cells must be added to replenish the PSM. Loss of cells is monitored by comparing the total density of cells in each of the subdomains with a target density, ρ_0. If the density in any of these subdomains falls below ρ_0, a new cell is added to that domain. If the density of more than one of the domains falls below ρ_0 then a cell is only added to the subdomain with the lowest density. The position and phase of the nascent cell both vary across different simulations discussed here. Below is a description of how these quantities vary.

In simulations where cell addition is 'random', cells are added with a random position within each domain, with a random phase, i.e. if a cell is added within one of the cylindrical domains, it is given a random $x_i \in [x_a + x_d, X_c]$, a random $r_i \in [0, r]$, and $q_i \in [0, 2\pi)$. Alternatively, if the cell is added within the half-toroid domain, it is given a

random $p_i \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, and random r_i and q_i, defined as above. The cell is also assigned a random phase $\theta_i \in [0, 2\pi)$. This method of cell addition is unbiased in terms of its effect on the model but a biologically implausible scenario of cell ingression, so we use it here as a negative control.

In simulations where we seek to more accurately model cell ingression, cells are added on the cell surface, i.e. $r_i = r$. Additionally, the surface domain on which cells can be added is restricted. For the half-toroid domain, we restrict cell addition to the dorsal quarter of the domain, i.e. $q_i \in [\frac{\pi}{4}, \frac{3\pi}{4}]$, $p_i \in [-\frac{\pi}{2}, \frac{\pi}{2}]$. For the cylindrical domains we restrict addition to the bottom quarter of the surface, i.e. $q_i \in [\frac{5\pi}{4}, \frac{7\pi}{4}]$. To more accurately model conditions of cell ingression combinations of these three modes of cell addition are employed (figure 1).

In newly added cells the direction vector **n** is initialised with a random vector of unit length. For simulations where cell division is present, cells are added with a random cell cycle phase $\tau \in 0, T_G + T_M$.

Coupled Oscillator Model of the Segmentation Clock

Here only the phase of each cell's segmentation clock cycle is considered, and oscillation amplitudes are neglected. This simplification is appropriate when oscillator coupling is weak (***Kuramoto*** (***1984***)). Here oscillators have a spatially-dependent intrinsic frequency $\omega(x_i)$, and couple to adjacent cells (defined as those within d_c, a cell's diameter) with a magnitude controlled by the scalar value κ:

$$\frac{d\theta_i}{dt} = \omega(x_i) + \frac{\kappa}{N_i(t)} \sum_{|\mathbf{x}_j - \mathbf{x}_i| \leq d_c} \sin(\theta_j - \theta_i) + \sqrt{2D_\theta}\xi_{\theta_i}(t) \tag{17}$$

Oscillations are noisy, with phase noise intensity D_θ. $\xi_{\theta_i}(t)$ is a gaussian noise term with $\langle \xi_{\theta_i}(t) \rangle = 0$, $\langle \xi_{\theta_i}(t)\xi_{\theta_j}(t') \rangle = \delta_{ij}\delta(t - t')$ for all t, t'. $N_i(t)$ is the number of cells within a radius of length d_c of cell i at time t, i.e. $N_i(t) = |\{j \mid |\mathbf{x}_j(t) - \mathbf{x}_i(t)| \leq d_c\}|$.

The intrinsic frequency $\omega(x_i)$ decreases toward the PSM anterior, creating a frequency gradient and travelling phase waves that move toward the anterior (***Soroldoni et al.*** (***2014***)). $\omega(x_i)$ is defined as:

$$\omega(x_i) = \omega_0\left(\sigma + (1 - \sigma) \cdot \frac{1 - e^{-\frac{k(x - x_a)}{L}}}{1 - e^{-k}}\right) \tag{18}$$

where ω_0 corresponds to the frequency of cells at the posterior of the tissue, σ the fold-change in frequency at the anterior compared to the posterior, and k controls the

shape of the gradient.

To simulate the effect of the wavefront, where the clock phase is interpreted by cells to pre-pattern somite boundaries (**Cooke and Zeeman** (**1976**)), beyond $x = x_a$ oscillations arrest, forming stable patterns of phase (figure 1A), i.e. $\frac{d\theta_i}{dt} = 0$ if $x_i < x_a$.

Cell division and segmentation clock arrest

During M phase of the cell cycle, the segmentation clock arrests (**Horikawa et al.** (**2006**); **Delaune et al.** (**2012**)). To simulate this, we assign each cell i a cell cycle phase $\tau \in [0, T_G + T_M)$, where T_G and T_M are the duration of G1-G2 and M phases in mins respectively, and evolve τ according to

$$\tau = (t - t_{addition}) \mod (T_G + T_M) \tag{19}$$

where $t_{addition}$ denotes the time at which the cell was added.

To simulate the arrest of segmentation clock gene expression during M phase of the clock, we evolve θ_i according to

$$\frac{d\theta_i}{dt} = H(T_G - \tau_i) \cdot \left(\omega(x_i) + \frac{\kappa}{N_i(t)} \sum_{|\mathbf{x}_j - x_i| \leq d_c} \sin(\theta_j - \theta_i) + \sqrt{2D_\theta} \xi_{\theta_i}(t) \right) \tag{20}$$

where $H(x)$ is the Heaviside step function:

$$H(x) = \begin{cases} 1 & x > 0 \\ 0 & x \leq 0 \end{cases} \tag{21}$$

If $\tau_i \geq T_G + T_M$ then we consider the cell cycle complete and a new cell k is spawned at a random position of length d_{new} away from cell i, i.e.

$$\mathbf{x}_k = \mathbf{x}_i + d_{new} \begin{pmatrix} \cos\varphi \sin\phi \\ \sin\varphi \sin\phi \\ \cos\phi \end{pmatrix}$$

for uniformly distributed random $\phi \in [0, \pi]$, $\varphi \in [0, 2\pi)$. Other variables are inherited from cell i, i.e. $n_k = n_i$ and $\theta_k = \theta_i$. The choice of d_{new} is somewhat arbitrary, however as this distance decreases the magnitude of the cell-cell repulsion force (equation 8) grows very large, and one must choose a d_{new} sufficiently small to be biologically accurate while large enough to avoid implausible cell velocities due to a possibly very large cell-cell repulsion force. We find that $d_{new} = d_c/10$ is a suitable compromise between these two

scenarios.

We are not aware of any direct measurements of T_G within the Zebrafish PSM, and so have calculated a value for this parameter based on some assumptions. M phase within the PSM lasts approximately 15 mins, and approximately 8% of cells in the PSM at a single time point are in M phase (***Horikawa et al.*** (***2006***); ***Kanki and Ho*** (***1997***)). Cells in M phase appear to be distributed homogenoulys in space (***Kanki and Ho*** (***1997***)), and therefore cell cycles in the PSM may be independent, in which case the total duration $T_G + T_M$ is approximately equal to $15/0.08 = 187.5$ mins, and $T_G = 172.5$ mins. Unless otherwise stated, we fix $T_G + T_M = 187.5$.

Modelling compaction-extension

To model the compaction-extension of the PSM (***Thomson et al.*** (***2021***)), we shrink the PSM radius r, and length L over time, concurrent with an increase in density ρ_0 and a step-wise decrease in cell diameter d_c (see figure 4). This is derived from the observations of ***Thomson et al.*** (***2021***) who observed changes in the PSM height, width, length, density, and cell diameter, during the latter stages of zebrafish somitogenesis. We sought to model this process using the rates of change in these parameters described by ***Thomson et al.*** (***2021***), however as their published rates are in units of change per somite stage, and the rate of somitogenesis is nonlinear in zebrafish (***Schröter et al.*** (***2008***)), we needed some interpolation of somite number over time in order to describe the according change in rate over time.

To this, we extracted the data for somite number over time from ***Thomson et al.*** (***2021***) using the web app WebPlotDigitizer (***Rohatgi*** (***2022***)), and fitted a function of the form

$$s(t) = 6 + \frac{t}{24.7} - ae^{b(t-300)} \tag{22}$$

for unknown parameters a, b, using the Julia package LsqFit.jl. We obtained values of $a = 0.5001$ and $b = 0.0049$ using this process. Using this interpolation of somite number over time, we then could change the tissue radius, length, cell diameter, and density, using the values published in ***Thomson et al.*** (***2021***). As this only occurs in the latter stages of somitogenesis (measured only from 16ss onwards), we hold these variables constant before the time at which 16 somites are formed in zebrafish.

The equations describing change in tissue length (L), wavefront position (x_a), and anterior limit of cell addition therefore become:

$$L(t) = \begin{cases} L_0 & s(t) < 16 \\ L_0 - m_a(s(t) - 16) & s(t) \geq 16 \end{cases} \tag{23}$$

$$x_a(t) = \begin{cases} x_{a0} & s(t) < 16 \\ x_{a0} + m_a(s(t) - 16) & s(t) \geq 16 \end{cases} \tag{24}$$

$$x_d(t) = x_a(t) + 100 \tag{25}$$

where m_a denotes the rate at which the PSM shrinks in length, and L_0 and x_{a0} the initial values for L and x_a before shrinking, respectively. The equation describing the change in radius r is

$$r(t) = \begin{cases} r_0 & s(t) < 16 \\ r_0 - m_r(s(t) - 16) & s(t) \geq 16 \end{cases} \tag{26}$$

where m_r denotes the rate at which the PSM shrinks in radius, and r_0 the initial value for r before shrinking. In their work *Thomson et al.* (*2021*) reported that the PSM shrinks in height more rapidly at the PSM posterior than at the anterior, however the rate of shrinkage of the PSM in width (along the medio-lateral axis) was the same across the anterior and posterior halves of the PSM (*Thomson et al.* (*2021*)). As our model assumes the tissue is as wide as it is tall, we take only the rate of shrinkage in the medio-lateral axis in our value for m_r, and calculate m_r by taking the mean of the two (similar) rates published by *Thomson et al.* (*2021*) for the anterior and posterior halves of the PSM.

The functions describing change in tissue density ρ and cell diameter d_c are similarly

$$\rho(t) = \begin{cases} \rho_{t0} & s(t) < 16 \\ \rho_{t0} + m_\rho(s(t) - 16) & s(t) \geq 16 \end{cases} \tag{27}$$

and

$$d_c(t) = \begin{cases} d_{c0} & s(t) < 16 \\ d_{c0} - m_{d_c}(s(t) - 16) & s(t) \geq 16 \end{cases} \tag{28}$$

where m_ρ and m_{d_c} represent the rates of change in density and cell diameter respectively, and ρ_{t0} and d_{c0} the initial values for density and cell diameter respectively, before shrinking. In their work *Thomson et al.* (*2021*) only report the cell diameter for 16ss and 26ss zebrafish embryos, but we choose to shrink the diameter of cells at the rate they

describe between these two timepoints. Our results are insensitive to choice of a function for $d_c(t)$ where cells continue shrinking in diameter at the same rate past 26ss (see figure 4 supp. 1).

Initial Conditions

To initialise the simulation, N cells are generated with random positions

$$\mathbf{x}_i = \begin{pmatrix} x_i \\ r + r_i \cos q_i \\ Z_c + r_i \sin q_i \end{pmatrix}$$

and a further N with positions

$$\mathbf{x}_i = \begin{pmatrix} x_i \\ r + 2R + r_i \cos q_i \\ Z_c + r_i \sin q_i \end{pmatrix}$$

and then a further N_{TB} with positions

$$\mathbf{x}_i = \begin{pmatrix} X_c + R \cos p_i + r_i \cos p_i \cos q_i \\ Y_c + R \sin p_i + r_i \sin p_i \cos q_i \\ Z_c + r_i \sin q_i \end{pmatrix}$$

for $x_i \in [x_a, X_c]$, $p_i \in [-\frac{\pi}{2}, \frac{\pi}{2}]$, $q_i \in [0, 2\pi]$, and $r_i \in [0, r]$, where $N = \lfloor \rho_0 \pi r^2 X_c \rfloor$ and $N_{TB} = \lfloor \rho_0 \pi^2 r^2 R \rfloor$. As this generates more cells near the tissue mid-line than at the periphery, in order to homogenise local density the tissue is 'relaxed' for 10 mins, whereby $\mathbf{x}_i$ evolves according to

$$\frac{d\mathbf{x}_i}{dt} = v_0(x_i)\mathbf{n}_i(t) + \sum_{j=1, j\neq i}^{N} \mathbf{F}(\mathbf{x}_i, \mathbf{x}_j) + \mathbf{F}_b(\mathbf{x}_i) \tag{29}$$

and $\mathbf{n}_i(t)$ evolves according to equation 4. This causes cells to be evenly distributed throughout the tissue.

When cells are initially 'spawned' prior to relaxation, $n_i(t)$ is assigned for each cell at random, i.e. $\mathbf{n}_i(0) = (\cos \varphi_i \sin \phi_i, \sin \varphi_i \sin \phi_i, \cos \phi_i)^T$ for random $\varphi_i \in [0, 2\pi)$, $\phi_i \in [0, \pi]$. The segmentation clock phase θ_i is also assigned as a constant value $\theta_i(0) = \frac{3\pi}{2}$.

Parameter Values

Parameter values, and their source, are shown in Table 1.

Symbol	Value	Units	Symbol	Value	Units
X_c	300	μm	L	$X_c - x_a + R + r$	μm
Y_c	85	μm	x_a	0	μm
Z_c	25	μm	x_d	100	μm
R	60	μm	x_q	0.7	-
r	25	μm	d_c	11	μm
v_a	1.67	μm·min^{-1}	D	0.026	min^{-1}
v_p	3	μm·min^{-1}	μ	8.71	μm·min^{-1}
X_v	0.4	-	μ_b	20	μm·min^{-1}
h	3	-	r_b	1	μm
v_s	1	μm·min^{-1}	ρ_0	0.0015	μm^{-3}
L_0	325	μm	T_G	187.5 - 15	min
m_a	14.1	μm ·somite^{-1}	T_M	15	min
r_0	27.6	μm	κ	0.07	min^{-1}
m_r	1.1625	μm ·somite^{-1}	ω_0	0.2094	min^{-1}
d_{c0}	9.2	μm	σ	0.66	-
$m_{d_{c0}}$	0.2	μm ·somite^{-1}	k	3.07	-
ρ_{t0}	0.002123	μm^{-3}	$D\theta$	0.0013	min^{-1}
m_ρ	0.000121	μm ·somite^{-1}			

Table 1. Parameter values used in our work, unless otherwise stated.

Implementation

Equations 17 and 1 are solved from the initial conditions outlined above across a 1000 min time-span using an forward euler scheme in Julia v1.8.2, using a time step dt = 0.01 mins. Code for simulation and analysis can be found at
https://github.com/jewh/ModularityPSMClock.

Analysis and Metrics

Here we measure clock dynamics using two metrics. The first is the synchrony of the segmentation clock across cells, denoted r, calculated by

$$re^{i\psi} = \frac{1}{N} \sum_{j=1}^{N} e^{i\theta_j} \tag{30}$$

which is sometimes referred to as the Kuramoto Phase Order parameter. When $r \approx 0$, oscillations are asynchronous, and when $r \approx 1$, oscillations are synchronous.

The second metric is the average (mean) frequency of cells in a given domain, where the frequency $\frac{d\theta}{dt}$ is simply the value of equation 17. In cases where we are interested in changes in frequency upon varying model parameters, we find it convenient to define

$$\Delta\frac{d\theta}{dt} = \frac{d\theta}{dt} - \frac{1}{d_c}\int_{x_a}^{x_a+d_c}\omega(x)dx \tag{31}$$

in order to highlight changes in frequency.

Figure 8. Top Trace of synchrony r for a d_c-wide domain of cells at the left-hand anterior edge of the PSM over 1000 mins. Data drawn from a single simulation with random cell addition, using parameters as per *Uriu et al.* (*2021*). **Bottom** Trace of mean frequency $\frac{d\theta}{dt}$ for a d_c-wide domain of cells at the left-hand anterior edge of the PSM over 1000 mins. Data drawn from the same simulation as above. Plotted with a black dashed line is the average intrinsic frequency $\omega(x)$ across the domain, calculated using the formula shown.

In order to analyse the output of a simulation in terms of one value, for many analyses we calculate r, $\frac{d\theta}{dt}$, or $\Delta\omega$, for the cells within one cell's diameter of the anterior (i.e. $x_a \le x_i \le x_a+d_c$). We do this because at the wavefront oscillations arrest and pre-pattern somites, and therefore the frequency and synchrony of oscillations immediately adjacent to the PSM anterior will determine the somite length and precision of pre-patterning. We thus regard the values of r, $\frac{d\theta}{dt}$, and $\Delta\omega$ as proxies for phenotype exhibited by the simulation. As discussed above, unless stated otherwise we measure clock dynamics after 1000 mins, as we find this sufficient to ensure that the dynamics of the clock have reached a steady state that we deem would be reached in vivo (see figure 8).

Acknowledgments

We thank Koichiro Uriu for his help in providing code and first establishing the model. We thank Usha Kadiyala for her comments regarding estimates for T_G. We thank Ben Steventon for his feedback on the project. The authors would like to acknowledge the use of the University of Oxford Advanced Research Computing (ARC) facility in carrying out this work. http://dx.doi.org/10.5281/zenodo.22558

References

Banavar SP, Carn EK, Rowghanian P, Stooke-Vaughan G, Kim S, Campàs O. Mechanical control of tissue shape and morphogenetic flows during vertebrate body axis elongation. Scientific reports. 2021; 11(1):1–14. doi: 10.1038/s41598-021-87672-3.

Bouldin CM, Snelson CD, Farr GH, Kimelman D. Restricted expression of cdc25a in the tailbud is essential for formation of the zebrafish posterior body. Genes Development. 2014; 28:384–395. doi: 10.1101/gad.233577.113.

Bénazéraf B, Francois P, Baker RE, Denans N, Little CD, Pourquié O. A random cell motility gradient downstream of FGF controls elongation of an amniote embryo. Nature. 2010; 466:248–52. doi: 10.1038/nature09151.

Cooke J, Zeeman EC. A clock and wavefront model for control of the number of repeated structures during animal morphogenesis. Journal of Theoretical Biology. 1976; 58(2):455–476. doi: 10.1016/s0022-5193(76)80131-2.

Delaune EA, François P, Shih NP, Amacher SL. Single-cell-resolution imaging of the impact of Notch signaling and mitosis on segmentation clock dynamics. Developmental cell. 2012; 23(5):995–1005. doi: 10.1016/j.devcel.2012.09.009.

Forero LL, Narayanan R, Huitem LF, VanBergen M, Apschner A, Peterson-Maduro J, Logister I, Valentin G, Morelli LG, Oates AC, Schulte-Merker S. Segmentation of the zebrafish axial skeleton relies on notochord sheath cells and not on the segmentation clock. eLife. 2018; 7. doi: 10.7554/eLife.33843.

Gomez C, Özbudak EM, Wunderlich J, Baumann D, Lewis J, Pourquié O. Control of segment number in vertebrate embryos. Nature. 2008; 454(7202):335–339. doi: 10.1038/nature07020.

Gomez C, Pourquié O. Developmental control of segment numbers in vertebrates. Journal of Experimental Zoology Part B: Molecular and Developmental Evolution. 2009; 312(6):533–544. doi: 10.1002/jez.b.21305.

Harima Y, Takashima Y, Ueda Y, Ohtsuka T, Kageyama R. Accelerating the Tempo of the Segmentation Clock by Reducing the Number of Introns in the Hes7 Gene. Cell Reports. 2013; 3(1):1–7. doi: 10.1016/j.celrep.2012.11.012.

Herrgen L, Ares S, Morelli LG, Schroter C, Julicher F, Oates AC. Intercellular Coupling Regulates the Period of the Segmentation Clock. Current Biology. 2010; 20. doi: 10.1016/j.cub.2010.06.034.

Horikawa K, Ishimatsu K, Yoshimoto E, Kondo S, Takeda H. Noise-resistant and synchronized oscillation of the segmentation clock. Nature. 2006; 441:719–723. doi: 10.1038/nature04861.

Kanki JP, Ho RK. The development of the posterior body in zebrafish. Development. 1997 02; 124(4):881–893. doi: 10.1242/dev.124.4.881.

Krol AJ, Roellig D, Dequéant ML, Tassy O, Glynn E, Hattem G, Mushegian A, Oates AC, Pourquié O. Evolutionary plasticity of segmentation clock networks. Development. 2011; 138(13):2783–2792. doi: 10.1242/dev.063834.

Kuramoto Y. Chemical turbulence. In: *Chemical oscillations, waves, and turbulence* Springer; 1984.p. 111–140.

Lawton AK, Nandi A, Stulberg MJ, Dray N, Sneddon MW, Pontius W, Emonet T, Holley SA. Regulated tissue fluidity steers zebrafish body elongation. Development. 2013; 140(3):573–582. doi: 10.1242/dev.090381.

Lewis J. Autoinhibition with Transcriptional Delay: A Simple Mechanism for the Zebrafish Somitogenesis Oscillator. Current Biology. 2003; 13(16):1398–1408. doi: 10.1016/s0960-9822(03)00534-7.

Liao BK, Jörg DJ, Oates AC. Faster embryonic segmentation through elevated Delta-Notch signalling. Nature Communications. 2016; 7(1):1–12. doi: 10.1038/ncomms11861.

Mara A, Schroeder J, Chalouni C, Holley SA. Priming, initiation and synchronization of the segmentation clock by deltaD and deltaC. Nature cell biology. 2007; 9(5):523–530. doi: 10.1038/ncb1578.

Michaut A, Mongera A, Gupta A, Serra M, Rigoni P, Lee JG, Duarte F, Hall AR, Mahadevan L, Guevorkian K, Pourquié O. Activity-driven extracellular volume expansion drives vertebrate axis elongation. bioRxiv. 2022; doi: 10.1101/2022.06.27.497799.

Mongera A, Rowghanian P, Gustafson HJ, Shelton E, Kealhofer DA, Carn EK, Serwane F, Lucio AA, Giammona J, Campàs O. A fluid-to-solid jamming transition underlies vertebrate body axis elongation. Nature. 2018; 561(7723):401–405. doi: 10.1038/s41586-018-0479-2.

Morin-Kensicki EM, Melancon E, Eisen JS. Segmental relationship between somites and vertebral column in zebrafish. Development. 2002; 129. doi: 10.1242/dev.129.16.3851.

Murray PJ, Maini PK, Baker RE. Modelling Oscillator synchronisation during vertebrate axis segmentation. In: *Pattern Formation in Morphogenesis* Springer; 2013.p. 95–105. doi: 10.1007/978-3-642-20164-6_9.

Murray PJ, Carrieri FA, Dale JK. Cell cycle regulation of oscillations yields coupling of growth and form in a computational model of the presomitic mesoderm. Journal of Theoretical Biology. 2019; 481. doi: 10.1016/j.jtbi.2019.05.006.

Naganathan SR, Popović M, Oates AC. Left–right symmetry of zebrafish embryos requires somite surface tension. Nature. 2022; 605(7910):516–521. doi: 10.1038/s41586-022-04646-9.

Oates AC, Morelli LG, Ares S. Patterning embryos with oscillations: structure, function and dynamics of the vertebrate segmentation clock. Development. 2012 02; 139(4):625–639. doi: 10.1242/dev.063735.

Palmeirim I, Henrique D, Ish-Horowicz D, Pourquié O. Avian hairy Gene Expression Identifies a Molecular Clock Linked to Vertebrate Segmentation and Somitogenesis. Cell. 1997; 91:639–648. doi: 10.1016/s0092-8674(00)80451-1.

Raff RA. The Shape of Life – Genes, Development and Evolution of Animal Forms. University of Chicago Press; 1996.

Richardson MK, Allen SP, Wright GM, Raynaud A, Hanken J. Somite number and vertebrate evolution. Development. 1998; 125(2):151–160. doi: 10.1242/dev.125.2.151.

Riedel-Kruse IH, Müller C, Oates AC. Synchrony Dynamics During Initiation, Failure, and Rescue of the Segmentation Clock. Science. 2007; 317(5846):1911–1915. doi: 10.1126/science.1142538.

Rohatgi A, Webplotdigitizer: Version 4.6; 2022. https://automeris.io/WebPlotDigitizer.

Schröter C, Herrgen L, Cardona A, Brouhard GJ, Feldman B, Oates AC. Dynamics of zebrafish somitogenesis. Developmental dynamics: an official publication of the American Association of Anatomists. 2008; 237(3):545–553. doi: 10.1002/dvdy.21458.

Schröter C, Ares S, Morelli LG, Isakova A, Hens K, Soroldoni D, Gajewski M, Jülicher F, Maerkl SJ, Deplancke B, Oates AC. Topology and Dynamics of the Zebrafish Segmentation Clock Core Circuit. PLOS Biology. 2012 07; 10(7):1–20. doi: 10.1371/journal.pbio.1001364.

Schröter C, Oates AC. Segment Number and Axial Identity in a Segmentation Clock Period Mutant. Current Biology. 2010; 20(14):1254–1258. doi: 10.1016/j.cub.2010.05.071.

Simsek MF, Chandel AS, Saparov D, Zinani OQ, Clason N, Özbudak EM. Periodic inhibition of Erk activity drives sequential somite segmentation. Nature. 2023; 613(7942):153–159. doi: 10.1038/s41586-022-05527-x.

Simsek MF, Özbudak EM. Spatial fold change of FGF signaling encodes positional information for segmental determination in zebrafish. Cell reports. 2018; 24(1):66–78. doi: 10.1016/j.celrep.2018.06.023.

Soroldoni D, Jörg DJ, Morelli LG, Richmond DL, Schindelin J, Jülicher F, Oates AC. A Doppler effect in embryonic pattern formation. Science. 2014; 345(6193):222–225. doi: 10.1126/science.1253089.

Steventon B, Duarte F, Lagadec R, Mazan S, Nicolas JF, Hirsinger E. Species-specific contribution of volumetric growth and tissue convergence to posterior body elongation in vertebrates. Development. 2016; 143(10):1732–1741. doi: 10.1242/dev.126375.

Thomson L, Muresan L, Steventon B. The zebrafish presomitic mesoderm elongates through compaction-extension. Cells & development. 2021; 168:203748. doi: 10.1016/j.cdev.2021.203748.

Uriu K, Ares S, Oates AC, Morelli LG. Dynamics of mobile coupled phase oscillators. Physical Review E. 2013; 87(3):032911. doi: 10.1103/PhysRevE.87.032911.

Uriu K, Bhavna R, Oates AC, Morelli LG. A framework for quantification and physical modeling of cell mixing applied to oscillator synchronization in vertebrate somitogenesis. Biology open. 2017; 6(8):1235–1244. doi: 10.1242/bio.025148.

Uriu K, Liao BK, Oates AC, Morelli LG. From local resynchronization to global pattern recovery in the zebrafish segmentation clock. eLife. 2021 feb; 10:e61358. doi: 10.7554/eLife.61358.

Uriu K, Morelli LG. Collective cell movement promotes synchronization of coupled genetic oscillators. Biophysical journal. 2014; 107(2):514–526. doi: 10.1016/j.bpj.2014.06.011.

Uriu K, Morishita Y, Iwasa Y. Random cell movement promotes synchronization of the segmentation clock. Proceedings of the National Academy of Sciences. 2010; 107(11):4979–4984. doi: 10.1073/pnas.0907122107.

Venzin OF, Oates AC. What are you synching about? Emerging complexity of Notch signaling in the segmentation clock. Developmental biology. 2020; 460(1):40–54. doi: 10.1016/j.ydbio.2019.06.024.

Wagner G, Pavlicev M, Cheverud J. The road to modularity. Nature Reviews Genetics. 2007; 8. doi: 10.1038/nrg2267.

Wagner GP, Altenberg L. Perspective: Complex Adaptations and the Evolution of Evolvability. Evolution. 1996; 50. doi: 10.1111/j.1558-5646.1996.tb02339.x.

Webb AB, Lengyel IM, Jörg DJ, Valentin G, Jülicher F, Morelli LG, Oates AC. Persistence, period and precision of autonomous cellular oscillators from the zebrafish segmentation clock. Elife. 2016; 5:e08438. doi: 10.7554/eLife.08438.

Xiong F, Ma W, Bénazéraf B, Mahadevan L, Pourquié O. Mechanical coupling coordinates the co-elongation of axial and paraxial tissues in avian embryos. Developmental Cell. 2020; 55(3):354–366. doi: 10.1016/j.devcel.2020.08.007.

Figure 3—figure supplement 1. Effect of co-varying coupling strength κ, tissue length L, and position of cell ingression on the median anterior synchrony of N=100 simulations.

Figure 3—figure supplement 2. Top Anterior synchrony after 1000 mins for changing tissue length L and varying position of cell ingression, where x_d is fixed at $x_d = X_c - 100$. N=100 simulations. **Bottom** Anterior mean frequency after 1000 mins for changing tissue length L and varying position of cell ingression where $x_[d]$ is fixed at $x_d = X_c - 100$. N=100 simulations.

$$s(t) = 6 + \frac{t}{24.7} - 0.5001\, e^{0.0049(t-300)}$$

Figure 4—figure supplement 1. Fitted function interpolating the data for zebrafish somitogenesis from *Schröter et al.* (*2008*) (black), against the raw data (magenta).

Figure 4—figure supplement 2. Comparing the choice of a 'stepwise' function for decreasing cell diameter d_c against a function where d_c decreases continuously until the simulation's end. **A** The stepwise (blue) and continuous (yellow) functions compared here. The intercept and gradient for these functions are derived from data from *Thomson et al.* (*2021*). **B** Anterior synchrony (r) over time for stepwise shrinking cells (blue) and continuously shrinking cells (yellow). Dark line shows the median of N=100 simulations, and the shaded area either side of this line shows the inter-quartile range (IQR). Results are plotted until the time at which at least one of the replicate simulations encounters a gap in the tissue at the tissue anterior (see methods). **C** Mean anterior frequency ($\frac{d\theta}{dt}$) over time for stepwise shrinking cells (blue) and continuously shrinking cells (yellow). Dark line shows the median of N=100 simulations, and the shaded area either side of this line shows the IQR. Results are plotted until the time at which at least one of the replicate simulations encounters a gap in the tissue at the tissue anterior.

Figure 4—figure supplement 3. Effect of starting tissue density d_0 on anterior synchrony and frequency over time, for a compacting tissue. **A** Functions showing tissue density $\rho(t)$ for different intercept values d_0. **B** Anterior synchrony (r) over time. Dark line shows the median of N=100 simulations, and the shaded area either side of this line shows the inter-quartile range (IQR). Results are plotted until the time at which at least one of the replicate simulations encounters a gap in the tissue at the tissue anterior (see methods). **C** Mean anterior frequency ($\frac{d\theta}{l}dt$) over time. Dark line shows the median of N=100 simulations, and the shaded area either side of this line shows the IQR. Results are plotted until the time at which at least one of the replicate simulations encounters a gap in the tissue at the tissue anterior.

Figure 5—figure supplement 1. The effect of cell division on frequency. **A** Effect on median frequency when varying x_{div} after 1000 mins for N=100 simulations. **B** Effect on median frequency when varying T_M after 1000 mins for N=100 simulations.

Figure 5—figure supplement 2. A Dependence of anterior synchrony on x_{div} and T_M. The median anterior synchrony after 1000 mins for N=100 simulations is plotted. **B** Dependence of anterior synchrony on T_M and intrinsic clock frequency ω_0. The median anterior synchrony after 1000 mins for N=100 simulations is plotted.