High-Throughput DNA melt measurements enable improved models of DNA folding thermodynamics

Yuxi Ke[1], Eesha Sharma[2,3], Hannah K. Wayment-Steele[4], Winston R. Becker[5], Anthony Ho[2], Emil Marklund[2,6], and William J. Greenleaf[2,7*]

[1]Department of Bioengineering, Stanford University, Stanford, CA 94305, USA

[2]Department of Genetics, Stanford University School of Medicine, Stanford, CA 94305, USA

[3]Lamar Health, Palo Alto, CA 94305, USA

[4]Department of Chemistry, Stanford University, Stanford, CA 94305, USA

[5]Program in Biophysics, Stanford University, Stanford, CA 94305, USA

[6]Science for Life Laboratory, Department of Biochemistry and Biophysics, Stockholm University, 17165, Solna, Sweden

[7]Department of Applied Physics, Stanford University, Stanford, CA 94305, USA

*Correspondence: wjg@stanford.edu (W.J.G.)

Abstract

DNA folding thermodynamics are central to many biological processes and biotechnological applications involving base-pairing. Current methods for predicting stability from DNA sequence use nearest-neighbor models that struggle to accurately capture the diverse sequence-dependency of elements other than Watson-Crick base pairs, likely due to insufficient experimental data. We introduce a massively parallel method, Array Melt, that uses fluorescence-based quenching signals to measure equilibrium stability of millions of DNA hairpins simultaneously on a repurposed Illumina sequencing flow cell. By leveraging this dataset of 27,732 sequences with two-state melting behavior, we derived a refined NUPACK-compatible nearest-neighbor model, a richer parameterization nearest-neighbor model that exhibits higher accuracy, and a graph neural network (GNN) model that identifies relevant interactions within DNA beyond nearest neighbors. All models provide improved accuracy in predicting DNA folding thermodynamics, providing improvements relevant for *in silico* design of qPCR primers, oligo hybridization probes, and DNA origami.

Introduction

The thermodynamics of DNA secondary structure formation is fundamental to understanding diverse biological processes, such as DNA replication[1] and repair[2], and is widely used in biotechnological applications, including the design of PCR primers[3], the engineering of in-situ hybridization (ISH) probes[4], the design of guide sequences for genome editing tools[5], and for DNA nanotechnology[6,7]. Numerous algorithms have been developed to predict DNA secondary structure thermodynamics, many of which rely on nearest-neighbor models[8–10].

Briefly, nearest-neighbor models assume that the total folding energy of DNA can be calculated by summing up the energies of each pair of neighboring base pairs. With this assumption, the "folded state" can be defined as the minimum

free energy (MFE) base pairing configuration, identified using dynamic programming[11]. Additionally, by using

partition functions, it is possible to calculate probabilities of the DNA in alternative secondary structures based on

these computed free energies. However, nearest-neighbor models often struggle to accurately capture the sheer

diversity and complexity of DNA secondary structure motifs, including hairpin loops, mismatches, and bulges. One

reason for this limitation is the lack of experimental data spanning those loop features upon which these models are

built, leading to inaccuracies in prediction. For example, the most widely used parameter set from *SantaLucia et al.*

2004 used data from 108 sequences to derive 12 parameters for Watson-Crick (WC) base pairs, and 174 sequences to

derive 44 parameters for internal single mismatches[8].

This data bottleneck has been due to the laborious nature of optical melting[12] and differential scanning calorimetry[13]

experiments, traditionally considered the gold standards of DNA secondary structure dynamics. Several groups have

attempted to overcome this bottleneck, usually with fluorescent melting in bulk solutions in well plates[14–17], reaching

a throughput on the scale of thousands. Still, due to the extremely large combinatorial DNA sequence space, these

works are limited to a specific type of DNA structural element in a fixed sequence scaffold.

To address this data generation bottleneck, we developed a method for systematic, accurate, high-throughput

measurements of nucleic acid secondary-structure motif thermal stability, enabling large-scale, quantitative

measurements of the thermodynamics of nucleic acid secondary structure. We demonstrate that nearest-neighbor

thermodynamic parameters inferred from these data are nearly identical to those observed from bulk thermal melting

experiments for WC base pairs but are improved for loop sequence motifs. The improved nearest-neighbor model

parameter sets are capable of accurately predicting DNA duplex folding energy on orthogonal unseen data sets.

Furthermore, by deploying advanced computational methods including deep learning to model DNA thermodynamics,

we develop a state-of-the-art thermodynamic model for DNA hairpin folding with accuracy comparable to

measurement uncertainties.

Results

Array melt: A high-throughput technique for quantifying the melting behavior of nucleic acids.

We developed the Array Melt technique, a method that allows for the rapid and comprehensive quantification of DNA

folding thermodynamics in high throughput. This technique is based on an Illumina MiSeq chip repurposed for high-

throughput measurements[18,19]. To assay the melting behavior of a nucleic acid construct, we engineered a common

region for annealing a 3'-fluorophore-labeled oligonucleotide to the 5'-end of a hairpin to be investigated. Similarly,

we designed another region for annealing a 5'-quencher-labeled oligonucleotide to the 3'-end of the hairpin (**Fig. 1a**).

After determining the position and sequence of the variant on the MiSeq chip through sequencing, we annealed Cy3-

labeled and Black Hole Quencher (BHQ)-labeled oligonucleotides to these respective binding sites. These annealed

oligos have predicted melting temperatures of 74°C, much higher than the highest temperature in our experiments

(**Table S1**). At room temperature, this fluorophore-quencher pair results in low levels of fluorescence in each cluster,

as the bulk of our hairpin library is folded at room temperature. As the sequence-variable hairpin regions are exposed

to increasing temperatures between 20 °C to 60 °C, the distance between Cy3 and BHQ increases, leading to brighter

fluorescence signals (**Fig. 1a-b**). Using this measurement scheme, we synthesized a comprehensive library of 42,177

unique hairpin sequences for measurement (**Fig. 1c**). These sequences integrated diverse structural elements into a

hairpin scaffold, including Watson-Crick pairing (WC), mismatches, bulges, hairpin loops of various lengths, and

other control sequences, including super-stable stem, polynucleotide repeats, and single-strand fluorescence controls

(see **Fig. 1c** and Methods).

We performed thorough quality control on the dataset by first requiring variants to accurately fit a two-state model

with melting behavior observable within our temperature measurement range (see Methods for our filtering strategy).

Total variants that pass these stringent thresholds include 6,393,050 individual melt curves (including technical

replicates) spanning 27,732 sequence variants for subsequent analysis with a standard two-state melt model.

Sensitivity and precision of the Array Melt method

To calibrate the amount of fluorescence observed as a function of the distance between Cy3 and BHQ, we evaluated

the fluorescence signals of a series of repeat sequence variants. These variants were designed to systematically increase

the distance between the fluorophore and quencher by incorporating increasing numbers of mono-, di- or tri-nucleotide

repeats. The fluorescent signals, aggregated over repeats of the same length, showed a nearly linear response to

distance variations up to approximately 8 nucleotides (**Fig. 1d**). Our observed length-fluorescence curve closely aligns

with a theoretical static quenching curve (Methods). We normalized the fluorescent signals using unfolded and folded

controls designed within the library (**Fig. S1b**). Three example melt curves in **Fig. 1f** illustrated increasing observed

melting temperatures and increasing GC content of the variable region. Technical replicate measurements from Array

Melt data showed high correlation, with $R > 0.94$ (**Fig. S1c**). Melt curves were also highly correlated with anneal

curves ($R = 0.964$), confirming that the DNA molecules were measured at equilibrium (**Fig. 1e**). Upon analyzing our

measurement precision (**Methods**), we found that nearly 90% of the tested variants displayed ΔG_{37} uncertainties

within 0.3 kcal/mol, or 0.5 k_BT (**Fig. 1h**). Overall, these data demonstrate that the equilibrium, aggregate fluorescence

signals obtained from the Array Melt method provide highly accurate melting information and can serve as a

"molecular ruler" capable of resolving single nucleotide distance variations.

FIGURE 1

Figure 1. High-Throughput thermodynamic measurement of DNA secondary structures using a fluorophore–quencher system.

a. Schematic representation of DNA molecules in a folded (quenched) and an unfolded (fluorescent) state.

b. Images of fluorescent DNA clusters immobilized on the sequencing chip surface. Top: Image with only fluorophore-conjugated oligo (Cy3), and with both fluorophore- and quencher-conjugated oligo at increasing temperatures. Bottom: Schematics of DNA molecules in each image. All images are normalized to super-stable stem and repeat control variants for temperature-dependent effects on fluorescence and quenching during the experiment, as shown in **Fig. S1d**.

c. Design of library variants between the constant-sequence binding sites for fluorophore- and quencher-conjugated oligonucleotides. Red represents scaffold nucleotides constant within each type, and blue represents the variable nucleotides systematically permuted ('N'). Numbers under each class indicate the number of unique sequences in each class.

d. Fluorescence measurement of control constructs where the single-strand distance between the fluorophore and the quencher increases in single nucleotide steps. The orange line shows the theoretical fit.

e. Correlation of ΔG_{37} derived from library variants under increasing temperatures (melt curve, x-axis) and decreasing temperatures (annealing curve, y-axis)

f. Representative examples of melt curves for constructs that vary in GC content.

g. Pearson correlation coefficient (R) of ΔG_{37} for replicates of three melt and one annealing curve experiment.

h. Standard error of ΔG_{37} for various construct classes as a function of ΔG_{37}.

UV melting validation

To further validate our method, we compared the results from the Array Melt method to those obtained using conventional UV melting methods on a cloud lab platform. We selected random representative variants from each category within the Array Melt library and sent the synthesized oligonucleotides to Emerald Cloud Lab (ECL) for analysis. Using the variant information and UV melting experiment parameters as input, scripts written in Symbolic Lab Language (SLL) automatically generated experiments as ECL protocol objects. These protocols were then queued for remote execution in a physical laboratory. Raw melting curve data from ECL were automatically analyzed by custom Python code, and samples that failed to pass quality control were resubmitted to ECL for re-measurement (**Fig. 2a**, **Fig. S2a**, **Methods**). A comparison of the melting profiles of random library variants showed good agreement between the two methods (**Fig. 2c-e,** R=0.85 for ΔG_{37}, or R^2= 0.62 after the linear adjustment described later). The melting temperature was found to be independent of DNA oligo concentration (**Fig. S2b**), suggesting intramolecular folding. Notably, we observed a systematic offset between the UV measurements and our Array Melt results, which could be due to the presence of the hairpins on a flow-cell surface, or the additional sequence elements included to facilitate our quenching-based readout. To account for these potential systematic differences, we applied a single offset to the melting points observed in our dataset and used the corrected results for subsequent analysis (**Methods**; on average, a correction of 9.35°C in Tm or approximately 0.7 kcal/mol in ΔG_{37} was applied). Compared to standard nearest-neighbor model predictions from NUPACK software, the measured values showed moderate improvement for ΔG and Tm, and considerable improvement for ΔH (**Fig. 2e, Fig. S2c**). This UV melting validation demonstrated the ability of the Array Melt method to accurately measure the folding energies of DNA and generalize to in-solution systems.

FIGURE 2

Figure 2. Cloud lab UV melting validation of the method

    a.   Schematic representation of UV melting data collection and analysis using the Emerald Cloud Lab (ECL) platform. Magenta arrows indicate semi-automatic steps, and teal arrows indicate automatic steps.

    b.   Example UV melting curve of the variant WC2318. Pink dots represent measured data points of the cooling curve; the green solid line represents the fitted line; the green dotted line represents the fitted melting temperature.

    c.   Mean absolute error (MAE) of ΔG, ΔH and Tm between Array Melt data and UV melting data before and after systematic error correction. Gray bars represent errors prior to correction; colored bars represent errors after applying correction with a single offset parameter.

    d.   Direct comparison of measured ΔG, ΔH and Tm between Array Melt and UV melting data. Each datapoint represents one hairpin variant. The gray dots represent measurements prior to correction; the colored dots represent measurements after applying the correction.

    e.   Pearson's correlation coefficient (R) between UV melting data and either Array Melt data or nearest-neighbor model predictions. The colored bars represent the correlation between Array Melt and UV melting data; the black bars represent that between nearest-neighbor model predictions and UV melting data.

Quantifying nearest-neighbor model performance on non-WC hairpins

After validation of our experimental system, we aimed to examine the performance of the standard nearest-neighbor model on hairpins whose stems are not perfect Watson-Crick pairs. When predicting folding free energies associated with different single mismatch types (**Fig. 3a**), the nearest-neighbor model achieved an R^2 of 0.61 on the average free energy for each of the types (implying it can explain 61% of the observed experimental variance), or $R^2 = 0.31$ on all individual variants. However, this model was unable to consistently predict significant variances in folding energies that arise from the same mismatch type with differing flanking sequences ($R^2 = -0.61$ for T>C mismatches, as shown on the right side panel, $R^2 = 0.09$ for A>G, $R^2 = -0.48$ for C>T, and $R^2 = -1.66$ for G>A) (**Fig. 3b, Fig. S3a**). Indeed, for three out of the four mismatches we analyzed, the nearest-neighbor model introduced extra variance (i.e. negative R^2 value) compared to the class-average prediction, implying that the variance in the nearest-neighbor model's prediction residual was greater than the actual variance present in the data (**Fig. 3c, Fig. S3b**). While the nearest-neighbor model succeeded in capturing variance for some variant groups with the same mismatch type and nearest-neighbor base pairs, it is unable to capture others (**Fig. S3c**). This limitation of the nearest-neighbor model led us to conclude that there are significant variances in folding energy that are not accounted for by the nearest-neighbor model.

Influence of sequences on energy across different secondary structures

We hypothesized that some of the observed variance in melting behavior not explained by the nearest-neighbor model may be due to the influence of sequences adjacent to mismatches. To assess this possibility, we calculated the impact of each nucleotide position on folding energy. We first determined a nucleotide importance score at each nucleotide position in different sequence groups with the same target secondary structures by calculating the variance in either ΔG_{37} or $\Delta\Delta G_{37}$ values for the four different identities of the nucleotide at the target position (see **Methods**). While it is possible that variants might adopt alternative secondary structures yet maintain two-state behavior, our analysis still provides valuable insights into how sequences affect folding energy. For single mismatches, we used $\Delta\Delta G_{37}$ values

relative to Watson-Crick pairing "parent" variants to regress out the energy expected from Watson-Crick stacks; for

other secondary structures, we directly used ΔG_{37} values. As anticipated, nucleotides along the hairpin stem displayed

consistent nucleotide importance scores for Watson-Crick pairing variants, and the two mismatching nucleotides have

the highest contribution to this folding energy for single mismatch variants, both in the Array Melt data and the nearest-

neighbor model (**Fig. 3d, Fig. S3d**). Extending our analysis to hairpin loops and bulges, we observed that specific

sequences within these structures critically influence their thermodynamic energy (**Fig. 3e-f**). In triloops and

tetraloops, loop center nucleotides had a more pronounced influence on folding energy in the Array Melt data than

nearest-neighbor model predictions. We also noted a slight asymmetry between the 5' and 3' ends of the loop,

regardless of loop size (**Fig. 3e, Fig. S3e**). In bulges, flanking Watson-Crick pairs had greater influence than the bulge

sequence itself, with double bulges showing more sequence dependence than singles (**Fig. 3f, Fig. S3f**).

These observations prompted us to examine the energetic interactions between bases by calculating the additional

proportion of variance explained in a linear model that considered base interactions (i.e. allowing free parameters for

pairwise combinations of bases in the two probed positions) compared to one that treats each position independently

(i.e. only allowing free parameters for each individual position). Notably, nucleotides opposite the mismatch showed

interactions unaccounted for in the nearest-neighbor model (**Fig. 3g, Fig. S3g**). This interaction score decays with

distance along the stem (**Fig. S3h**). Our quantification showed that individual nucleotides, nearest neighbors, and

longer-range interactions contribute 74%, 14%, and 12% respectively in this specific case of single mismatches (**Fig.**

3h). In hairpin loops, the interactions between the loop ends and the loop center nucleotides were stronger than

expected in tetraloops, but not in triloops (**Fig. 3i, Fig. S3i-j**).

Subsequently, we fit linear regression models to Watson-Crick and single-mismatch variant Array Melt data, varying

the length of stack features from one (base pairs only), to two (nearest neighbors) and three (next nearest neighbors).

Although prediction errors were similar for nearest- and next-nearest-neighbor models in Watson-Crick variants, next-

nearest-neighbor models outperformed the nearest-neighbor ones in single-mismatch variants (**Fig.4a**).

FIGURE 3

Figure 3. Decomposition of sources of variances in measured energies

a. Schematic representation of mismatch types and ΔΔG calculation for single mismatches.

b. Comparison of single mismatch ΔΔG between nearest-neighbor model prediction and Array Melt measurements. Each variant is colored by the identity of the mismatch (left), and the T>C mismatches are zoomed in on the right.

c. Unexplained expected biological variance for each group of mismatches and combined. Variance explained or introduced by the nearest-neighbor model are shown in green and red, respectively.

d. Nucleotide importance scores for each individual nucleotide in mismatch constructs, grouped by secondary structure. The importance score is calculated as the sum of squares introduced by varying the given nucleotide position.

e. Nucleotide importance scores for tetraloops and triloops.

f. Nucleotide importance scores for bulges.

g. Pairwise interaction scores for each pair of nucleotide locations for Watson-Crick and single mismatch variants. A yellow star indicates the anchor point in each subplot, and the color of each position indicates the strength of the interaction with the anchor. The interaction score is calculated as the R-squared of a linear model given all combinations of the pair as features, subtracted by that of the two individual nucleotides.

h. Percentage contribution of single nucleotides, nearest neighbors, and longer-range interactions to the total variance in $\Delta\Delta G_{37}$ for single mismatches.

i. Pairwise interaction scores for tetraloops and triloops.

Revised parameter set for NUPACK integration

We aimed to develop a refined set of parameters to integrate our findings with the NUPACK software[7,20], thereby enabling more accurate DNA thermodynamic predictions that are easily accessible. We divided Array Melt data into training, validation, and test sets, respectively with 24979, 1315, and 1438 variants, stratified by major variant types as shown in **Fig. 1c**. Initially, we fitted compact linear regression models to the Array Melt data, focusing on structural elements prevalent in our variant library. By allowing the 10 Watson-Crick pairing stack parameters to vary freely (totaling 188 free parameters), we reproduced the established nearest-neighbor parameters set[8] (**Fig. 4b**). Subsequently, we fixed the values of the 10 Watson-Crick stack parameters and fitted a compact 178-parameter nearest-neighbor model. We compared the single mismatch parameters of our new parameter set with the original *dna04* set and observed similar overall trends. Our new parameters also captured context-dependent variations of GT mismatches and highlighted the instability of CC mismatches, which are shown to be the most thermodynamically unstable DNA single mismatch[21] (**Fig. 4c, Fig. S4c**). Next, we obtained parameters for tetraloops and triloops by aggregating thermodynamic measurements of relevant Array Melt variants and converting the aggregated ΔΔH and $\Delta\Delta G_{37}$ values to ΔH and ΔG_{37} parameters (**Methods**). We combined these corrected hairpin loop parameters with the compact 178-parameter nearest-neighbor model. This combined model, thereafter referred to as the NUPACK-compatible model, is easily accessible by pointing NUPACK to our parameter set file (**Fig. S4a**; parameter file available in supplementary material). Note that when using NUPACK to predict free energy, we assumed a two-state model in alignment with the Array Melt dataset, which was filtered based on the two-state criterion. Additionally, the ensemble model yielded inferior results compared to the two-state model, both with the original *dna04* and with our new parameter sets (**Fig. S4b**).

We evaluated the models on held-out Array Melt data and two orthogonal DNA duplex melting datasets from the

literature. To assess model performance, we used both the raw mean absolute error (MAE) and the adjusted MAE,

which is more relevant for assessing $\Delta\Delta G$ errors (**Supplementary Notes**). Testing our NUPACK-compatible model

on the held-out Array Melt test data resulted in an adjusted MAE of 0.59 kcal/mol for ΔG_{37} (**Fig. 4d-e**). This new

parameter set also improved predictions for DNA hairpins designed with random sequences measured with UV

melting (**Fig. S4d**). We further asked whether the new parameter set, trained on DNA hairpin data, extends to unseen

DNA duplexes. Using NUPACK to predict unseen duplex melting data, we observed a similar performance for

Watson-Crick strands with both the original *dna04* and the new parameter set. However, the new parameter set showed

improved performance on more complex variants. For a UV melting data set consisting of 384 Watson-Crick pairing

DNA duplexes[22], split evenly into validation and test sets, our parameter set performed comparably to *dna04* (**Fig.**

S4e). On a dataset from *Oliveira et al.* containing DNA duplex melting temperatures[15] with varying numbers of

mismatches (also split evenly between validation and test sets), both parameter sets had a similar adjusted MAE of

around 0.6 °C for Watson-Crick duplexes on the test set; while for single mismatches, the new parameter set achieved

a much lower adjusted MAE of 1.15 °C ($R^2 = 0.68$) compared to 1.66 °C ($R^2 = 0.25$) with *dna04*. The improvement

was even greater for double and triple mismatches (**Fig. 4f-g**). In conclusion, this updated parameter set, while simple,

enhances the prediction accuracy of NUPACK for DNA secondary structures, particularly those with mismatches and

hairpin loops. This improvement extends to orthogonal DNA duplex data that the model was not trained on.

Enhanced linear model with rich parameterization

Building on this NUPACK-compatible model, we next developed an enhanced linear model that incorporates a richer

parameter set ("richer parameter model"), allowing still more accurate and comprehensive prediction of DNA

thermodynamics on unseen validation data. Because this model has more parameters than are implemented in the

NUPACK, it is not immediately compatible with the software. We fit this model with 1338 free parameters,

incorporating still more context information (including interactions between the two stacks flanking a mismatch, bulge

sequence identity, and full hairpin loop sequence for triloops and tetraloops, see **Methods**), to the Array Melt training

data. This rich parameter nearest-neighbor model yields an even better performance both on the Array Melt test data

(adjusted $MAE = 0.34\ kcal/mol$, $R^2 = 0.81$) and the external duplex datasets (**Fig. 4d-g, Fig. S4e**). Notably, when

comparing the prediction error on Array Melt validation dataset as we titrate the percentage of training data used to

fit the model, the 178 parameter linear regression model used to build the NUPACK-compatible model plateaus with

just a small fraction of training data used, while the 1338 parameter model requires more measurements to plateau,

but still does so after being fit on less than half of the training data set (**Fig. 4h**). Given the relatively overdetermined

nature of even this rich parameter nearest-neighbor model, this behavior is expected, and clearly demonstrates that

increasing data size allows for more variance to be captured by a more expressive model.

FIGURE 4

Figure 4. Enhanced linear regression models for DNA folding energetics

a.   Adjusted mean absolute error (adjusted MAE) of linear regression models with features based on single base
pair, nearest neighbor, or next nearest neighbor stacks, applied to Array Melt test set data.

b.   Comparison of nearest-neighbor parameters for Watson-Crick pairs fitted from Array Melt data and those
reported in the literature[8].

c.   Visualization of new and the original *dna04* free energy parameters for single mismatches. "Context" and
"mismatch" definitions are consistent with those in **Figure 3a**.

d-e. Performance comparison of the original (*dna04*), NUPACK-compatible, and richer parameter nearest-
neighbor models on held-out Array Melt data. All calculations were performed at 1M Na$^+$ concentration. NN: nearest-
neighbor.

f-g. Performance comparison of the original (*dna04*), NUPACK-compatible, and richer parameter nearest-
neighbor models on an orthogonal dataset of DNA duplexes with varying numbers of mismatches (*Oliveira et al.*[15]).

h. Adjusted MAE on Array Melt data plotted as a function of the percentage of training data used to fit the linear
regression model.

Graph neural network (GNN) model for DNA energy prediction

Because our rich linear model likely had insufficient power to capture the biological variance in our data, we developed

a graph neural network model[23,24] specifically designed for predicting DNA energy. In this model, a DNA molecule

is represented as a graph where nodes represent nucleotides and edges represent chemical bonds. Nucleotides (A, T,

C, or G) and chemical bonds (either 5' to 3' backbone, 3' to 5' backbone, or hydrogen bond) are one-hot encoded as

node and edge features, respectively. Taking these graph representations as inputs, the GNN iteratively refines node

embeddings across graph convolutional layers. Essentially, nodes exchange information through different edge types,

allowing, over multiple layers, for the embedding of one node to encapsulate information from the entire graph.

Investigation of various types of graph convolutional layers revealed that the graph transformer layers[25] outperformed

other architectures, such as graph attention layers[26] (**Fig. S5a**). After graph convolution, each node possesses a learned

embedded feature, reflecting not only the original node features, but also the edges and overarching graph structures.

To embed this graph into a fixed-length vector representation, we employed a global pooling layer employing the

Set2Set algorithm[27], which used a Long Short-Term Memory (LSTM) processing block. This block systematically

ingests node embeddings while updating its internal state, resulting in a vector representation of the entire graph that

is order-invariant to the input node features. This processed summary vector is then fed into a small, fully connected

neural network, which regresses for ΔH and Tm values (**Fig. 5a**).

We trained this end-to-end model on Array Melt data. A portion of the Array Melt data was held out as validation and

test set. We first explored the effect of the number of graph convolutional layers on prediction accuracy (**Fig. 5b, Fig.**

S5b). Performance converged at around 4 graph convolutional layers, and did not improve beyond 4 layers, consistent

with our observations in **Fig. S3h** that pairwise nucleotide interactions span around a distance of 4 nt. When evaluating

model performance over an increasing percentage of training data used as in **Fig. 4h**, we observed a slower plateauing

of the curve, indicating that the model can make use of this large-scale data set (**Fig. 5c**). The trained GNN achieved

prediction error of 0.18 kcal/mol in dG_{37} ($R = 0.97$, $R^2 = 0.94$), compared to 0.34 kcal/mol of a baseline model that

simply looks up the k=8 closest variants in the training set and gives a weighted average of the lookup values as

prediction (**Fig. 5d-e, Fig. S5c-d**), and the GNN predictions are consistently high quality for different types of

sequence variants (**Fig. S5e**). Interestingly, although the GNN was trained entirely on graph representations of DNA hairpins without knowing the nature of backbones and base pairing, it is able to extend to the annealing and melting properties of DNA duplexes (**Fig. 5f**). While the GNN performance is slightly worse than our rich parameter nearest-neighbor model (**Fig. 4f-g, Fig. S4e**) on DNA duplexes, this level of performance suggests that the neural network has grasped the "gist" of the physical system of DNA folding thermodynamics and can extrapolate to non-hairping annealing contexts. Compared to the slow decrease of prediction error on the Array Melt data, prediction error rapidly decreased within around the first 20 epochs and then plateaued for the two DNA duplex datasets (**Fig. S5f**), likely because of the relatively low complexity of those datasets.

To better understand how the GNN makes predictions, we analyzed the intermediate activation from the output of the Set2Set aggregation layer. The GNN projects each DNA molecular graph into a 250-dimensional embedding vector. Many of the top principal components of the embeddings correlate strongly with predicted thermodynamics parameters ($R = 0.8$ between PC1 and ΔH, $R = -0.77$ between PC5 and Tm, and $R = 0.79$ between PC1 and ΔG_{37}), but none with other qualities of the sequence variants including DNA length or GC content (**Fig. 5g-h**), underscoring the GNN's focus on thermodynamics. When examining embeddings from different datasets (Array Melt, literature UV, and Oliveira et al.), each dataset's distinct design structure mapped to unique areas in a two-dimensional UMAP visualization (**Fig. 5i**). This observation indicates that the embeddings capture more than just predicted thermodynamics like melting temperatures (**Fig. S5g**), and that the learned embedding space encompasses sequence and structure details beyond mere parameter prediction.

Finally, we compared prediction errors across various models. The graph neural network is the top performer on Array Melt test data, while the 1778-parameter richer parameterization nearest-neighbor model performs best when applied to duplex data. Precision of the GNN is noteworthy, nearing the uncertainty range of bootstrapped experimental measurements (mean of uncertainty is 0.14 kcal/mol in ΔG_{37}), highlighting its effectiveness in capturing complex relationships in DNA sequences and structures (**Fig. 5j, Table 1**).

FIGURE 5

Figure 5. Graph neural network model for DNA folding thermodynamics
a.   Schematic representation of the graph neural network (GNN) architecture.
b.   Mean absolute error (MAE) of melting temperature on Array Melt data as a function of the number of graph
convolution layers in the GNN.
c.   MAE of ΔG_{37} on Array Melt data as a function of the percentage of training data used to fit the GNN.
d.   Pearson's R correlation coefficient between GNN predictions and held-out Array Melt data, literature UV
melting data, or Oliveira et al. duplex melting data.
e.   Scatter plot comparing GNN predictions with held-out Array Melt data. Each dot represents a single sequence
variant.
f.   Scatter plot comparing GNN predictions with literature UV melting data or Oliveira et al. duplex melting
data.
g.   Pearson's R between principal components of the learned variant embeddings and measured thermodynamic
parameters or variant properties.
h.   Comparison between the most correlated principal components and ΔH or Tm.
i.   UMAP visualization of learned variant embeddings after aggregation, colored by dataset (Array Melt,
literature UV melting, or Oliveira et al. duplex melting data).
j.   Benchmark of all models on held-out Array Melt data, compared to the measurement error of Array Melt.
k.   Benchmark of all models on the orthogonal Oliveira et al. dataset, plotted by the number of mismatches.
Colors correspond to models as in **Fig. 5j**.

		Model	Original NUPACK	NUPACK-compatible	Rich parameterization	Closest lookup	Graph neural network	Measurement
Array	ΔH	Adjusted MAE	8.59	6.27	4.97	5.27	3.13	3.09
		MAE	14.33	10.79	5.03	4.03	3.13	3.09
		R^2	0.12	0.48	0.64	0.77	0.86	NA
	Tm	Adjusted MAE	7.94	6.52	5.09	3.72	1.80	1.28
		MAE	8.38	6.53	5.65	3.72	1.80	1.28
		R^2	-0.06	0.26	0.20	0.78	0.94	NA
	ΔG_{37}	Adjusted MAE	0.87	0.59	0.34	0.34	0.19	0.14
		MAE	1.07	0.67	0.35	0.34	0.18	0.14
		R^2	-0.15	0.44	0.81	0.81	0.94	NA
Oliveira et.al	Tm (all)	Adjusted MAE	2.06	1.77	2.76	5.12	5.30	NA
		MAE	5.43	4.70	10.20	9.01	5.56	NA
		R	0.81	0.86	0.66	0.35	0.78	NA
		R^2	0.65	0.73	0.42	-1.06	0.49	NA
	R^2 of Tm	No mismatch	0.87	0.86	0.92	0.07	0.78	NA
		Single	0.25	0.68	0.75	-0.14	0.56	NA
		Double	0.13	0.26	0.21	-0.23	-0.12	NA
		Triple	0.03	0.24	0.27	-8.33	-1.21	NA

353 **Table 1. Model benchmarks.**
354 MAE: mean absolute error. Units of adjusted MAE and MAE are kcal/mol for ΔH and ΔG_{37} and °C for Tm.
355 Numbers in the "Measurement" column are means of bootstrapped uncertainties from measurement.

Discussion

In this work, we developed an accurate, scalable fluorescence-based method for measuring the melting of DNA hairpins (Array Melt) and used it to generate the largest-to-date DNA folding thermodynamics dataset. The Array Melt method, with its high throughput and precise measurement capabilities, allows for a more detailed exploration of the thermodynamics of diverse DNA structures, including those with mismatches, bulges, and various loop motifs. The scale of this data set revealed the significant role played by sequences adjacent to mismatches and within loops to the thermodynamic stability of hairpins. This observation implies that traditional models that parameterize only the immediate nearest neighbors must be underpowered to capture the variance observed in these molecular variants. To this end, we developed three quantitative folding energy models of increasing complexity, including revised parameterization of the linear regression model compatible with NUPACK, a richer nearest-neighbor model especially accurate for duplex annealing of mismatched molecules, and graph neural network (GNN) model that accells at predicting the melting behaviors of hairpins. To our knowledge, this GNN model is the first deep learning model applied to DNA folding. These results support the notion that while traditional nearest-neighbor models are accurate in predicting Watson-Crick stack behaviors, they fall short in accurately modeling the thermodynamics of more complex DNA structures, such as hairpins and interior loops. We anticipate that future library designs will be more oriented towards machine learning, enabling a more comprehensive and less biased sampling of the DNA sequencing space.

Overall, by providing a more accurate model for DNA folding thermodynamics, we pave the way for improved design of biotechnological tools and applications. For example, our models will provide a more accurate starting point for *in silico* design in areas such as qPCR primer design, DNA oligo hybridization probe design, and DNA origami. However, the study's focus on two-state folding presents a notable limitation, although our comparison between the two-state and ensemble methods (**Fig. S4b**) revealed greater accuracy of the former in predicting Array Melt measured thermodynamic parameters. The inferior performance of the ensemble method may be attributed to the inference methods of parameters. Additionally, the experimental design of the datasets used to derive nearest-neighbor models may also play a role. Therefore, interpreting nearest-neighbor model predictions and dynamic programming algorithm outputs as "true" energies associated with monolithic secondary structures warrants careful consideration. In this study, the two-state folding energies are better interpreted as an aggregated average across multiple folding states, rather than that of a single folded microstate within this ensemble. We anticipate that future analysis incorporating experimental design, data preprocessing, and parameter inference[28] techniques that consider ensemble folding could result in further improvements in prediction accuracy. This is particularly relevant when predicting probabilities of microstates of dynamic molecules is important.

This work also indicates that cloud labs can provide researchers with practical access to specialized instruments and support, addressing traditional challenges in equipment access and maintenance. These platforms also facilitate streamlined data collection and metadata tracking, improving open data and experiment reproducibility. This study

represents an early instance of using commercial cloud labs in research setting[29], suggesting potential broader
academic use.

Finally, while our measurements were carried out at a single monovalent salt concentration, large-scale measurements
exploring the stability of DNA structure to the concentration and type of monovalent or divalent cation would be
relatively straightforward. Furthermore, the potential application of our method to RNA structures presents an exciting
avenue for future research. Given the diverse and critical biological functions of RNA, which are heavily dependent
on its secondary structure, applying our high-throughput and precise thermodynamic measurement technique could
unveil new dimensions in RNA biology and facilitate the development of RNA-based therapeutics.

Acknowledgement

We thank Dr. Rhiju Das (Stanford University) for comments on the manuscript; Dr. Mark Schnitzer (Stanford
University) for supporting the project; Dr. Niles Pierce and Dr. Mark Fornace (Caltech) for answering NUPACK-
related questions. We thank Emerald Cloud Lab (ECL) for the UV melting experiments, with special attention to
Malav Desai and Ben Kline for tutorials and technical support of Symbolic Lab Language (SLL) programming and
Brian Frezza for funding.

This work was supported in part by NIH grants R01GM111990, P50HG007735, R01HG009909, P01GM066275,
UM1HG009436 and R01GM121487 to W.J.G. W.J.G. acknowledges support as a Chan Zuckerberg Investigator.
E.M. acknowledges support from the Swedish Research Council (grant 2020-06459), the Foundation Blanceflor, and
the Science for Life Laboratory (SciLifeLab).

Author contributions

409   Conceptualization, W.J.G., Y.K., E.S., and H.K.W.; Methodology, Y.K, E.S.; Software, Y.K. and H.K.W.; Validation,
410   Y.K; Formal Analysis, Y.K.; Investigation, Y.K, E.S., H.K.W., W.R.B., and A.H.; Data Curation, Y.K.; Writing -
411   Original Draft, Y.K. and W.J.G.; Writing - Review & Editing, Y.K., E.S., H.K.W., E.M., and W.J.G.; Visualization,
412   Y.K.; Supervision, E.M. and W.J.G.; Funding Acquisition, W.J.G.

Declaration of Interest

414   Stanford filed a patent application on aspects of this work with H.K.W., E.S., R.D., W.J.G., A.H., W.B., and Y.K. as
415   inventors (WO2023028618). W.J.G. is a consultant and equity holder for 10x Genomics, Guardant Health, Quantapore
416   and Ultima Genomics, and cofounder of Protillion Biosciences. The other authors declare no competing interests.

Methods

Library assembly and sequencing

Designed library variants were synthesized into DNA by Twist Biosciences (South San Francisco, CA). Sequence
map of the library is provided in supplementary information as *array_melt_library.gb*, where the "filler sequence" is

a 5' partial segment of AACAACAACAACATACTAACAACAACATAACAAATCAAAA, its length and the variable

region adding up to 40 bp, such that all synthesized fragments have the same length despite different variable region

length. Reverse complement of the sequence map was ordered as a synthesized oligo pool, amplified using internal

primers to enrich for full-length library variants. The PCR reaction consisted of 1 in 8 dilution of the synthesized oligo

pool (final concentration 10 nM, but concentration of full-length library variants could be much lower), 200 nM of

each primer (T7A1library, D-TruSeqR2 Table S1), 1x Phire Hot Start II PCR Master Mix (ThermoFisher Scientific

F125L). The reaction proceeded for 9 cycles of 98˚C for 10 seconds, 56˚C for 30 seconds, and 72˚C for 30 seconds.

Reaction mixtures were purified using QIAquick PCR Purification Kit (Qiagen 28104) to remove primers and

proteins, then eluted into 20 uL elution buffer.

After initial amplification, the library was amplified with primers to bring in sequences compatible with Illumina

sequencing. This five-piece assembly PCR included two outside primers and two adapter sequences. The PCR reaction

consisted of 1 μl of the previous reaction, 137 nM of outside primers (*short_C* and *short_D*; Table S1), 3.84 nM of

the adapter sequences (*C-i7pr-bc-T7A1* and *D_TruSeqR2*; Table S1), 1x Phire Hot Start II PCR Master Mix

(ThermoFisher Scientific F125L). The reaction proceeded for 14 cycles of 98˚C for 10 seconds, 56˚C for 30 seconds,

and 72˚C for 30 seconds. Reactions were purified using the QIAquick PCR Purification Kit and quantified with a

Qubit Fluorometer (ThermoFisher Scientific).

Sequencing was run with MiSeq Reagent Kit v3, 150 Cycles (MS-102-3001). The amplified library was diluted and

pooled to 1.16 nM final concentration in the pool (27% sample), together with a final concentration of 2.8 nM PhiX

control v3 (70% sample, Illumina), 0.04 nM fiducial marker (1% sample), 0.04 nM Cy3 no RNA control (1% sample),

and 0.04 nM single-strand fluorescence control sequences (1% sample, not included in analysis), adding up to a total

sample concentration of 4 nM. Paired-end sequencing was done with a custom read 1 primer (*stall_R1_primer*) and a

standard read 2 primer to cover the variable region in both read 1 and read 2 directions.

Imaging station setup

An imaging station was used to image the Miseq chip at increasing temperatures. This station was built from a

combination of custom-designed parts from a disassembled Illumina genome analyzer IIx[18,30]. Two channels were

employed: the "red" channel used the 660 nm laser and 664 nm long pass filter (Semrock BLP01-664R-25), and the

"green" channel used the 532 nm laser and 590 nm band pass filter (Semrock FF01-590/104-25). All images were

taken with 600 ms exposure times at 150 mW fiber input laser power. Focusing at each temperature was achieved by

sequentially adjusting the z-position and re-imaging the four corners of the flow cell; the adjusted z-positions were then

fit to a plane. We chose Cy3 as the fluorophore because the "green" channel is relatively more stable in our instrument

setup.

Post-sequencing, the chip was washed with Cleavage buffer (100 mM Tris-HCl, 125 mM NaCl, 0.05% Tween20, 100

mM TCEP, pH 7.4) to remove residual fluorescence from the reversible terminators used in the sequencing reaction

at 60˚C for 5 minutes. Any strands of DNA not covalently attached to the surface of the chip was removed by washing

in 100% formamide at 55˚C. The resulting single-stranded DNA fragments were incubated with 500 nM of the oligo

Biotin_D_Read2 and *red_oligo* (Table S1) in Hybridization buffer (5x SSC buffer (ThermoFisher 15557036), 5 mM

EDTA, 0.05% Tween20) for 15 minutes at 60 °C, subsequently the temperature was lowered to 40 °C for another 10

minutes. The *red_oligo* is conjugated to Alexa647 and binds a sparse subset of "fiducial" sequences on the chip, whose

fluorescence signal is used for initial image registration to the sequenced tiles.

Measuring melt curves on the chip

The Cy3 labeled *fluor_oligo* was annealed at 40˚C for 5 minutes, then the Black Hole Quencher (BHQ)-labeled oligo

quench_oligo (**Table S1**) using the same protocol. The chip was imaged after each step to ensure that hybridization

had occurred through increase or quenching of signal in the corresponding channel. The chip was then rinsed with

melt buffer (50 mM Na-HEPES pH 8.0, 25 mM NaCl, total Na^+ concentration about 88 mM at pH 8.0). We chose

HEPES buffer because it has one of the lowest sensitivities to temperature (-0.014 ΔpK_a/°C) among the most

commonly used buffers in biological experiments[31].

To quantify the melt curves, image station temperature was then lowered to 20˚C. For each temperature point, the

system was allowed to equilibrate to the new temperature (approximately 5 minutes) before refocusing and imaging.

The temperature was raised in 2.5˚C increments to a maximum temperature of 60˚C. Detailed protocol can be found

in XML imaging scripts in the supplementary information.

Processing sequencing data

Sequencing data from Illumina Miseq was processed to extract tile and coordinates of each sequenced cluster. Forward

and reverse paired-end reads were aligned using FLASH[32] with default settings. Consensus sequences from FLASH

were aligned to the reverse complement sequences of *fluor_oligo* and *quench_oligo* using a Needleman-Wunsch

alignment (nwalign3). For consensus sequences that successfully aligned to both with p-value <1e-3, evaluated as

described in[33], the variable region was extracted as the region between the two flanking regions and aligned to

sequences in the reference library. A reference sequence was assigned based on the best-scoring alignment with a p-

value < 1e-6. The resulting clusters were used for fluorescence quantification in the downstream data analysis.

Fluorescence data processing and image fitting

Fluorescent melt curve images were mapped to sequencing data from the Illumina Miseq. First, sequencing data was

processed to extract the tile and coordinates of each sequenced cluster. The coordinates were used to generate synthetic

images of clusters, which were registered to the fluorescent images iteratively to map the coordinates to the images at

sub-pixel resolution[34]. Once the identities of the clusters were determined, each cluster was fitted to a 2D normal

distribution to quantify its fluorescence[30].

Distance sensitivity curve

The repeat controls used to plot the distance sensitivity curve in Fig. 1d are repeats of varied lengths of the following

sequences: poly-A, T, AT, AAC, AAG, AAT, AC, ACC, AG, AGG, TC, TG, TTA, TTC, and TTG.

The theoretical curve fitted to the distance-to-fluorescence curve in Fig. 1d is $f(x) = \frac{1}{1+(a \cdot (x+b)^{-3})}$, where x is the

length in nt, $f(x)$ is the dimensionless fluorescence at length x, normalized to maximum fluorescence, b is a fitted

parameter in nt, which accounts for systemic distance offset, and a is another fitted parameter:

491
$$a = \frac{2\pi^{2/3}}{3} (K_{D,BHQ} \cdot N_A \cdot k^3)^{-1} \qquad (1)$$

492 where $K_{D,BHQ}$ is the dissociation constant for the fluor-quench pair, N_A is Avogadro's number, and $k =$

493 $0.64 \times 10^{-9} \, m \cdot nt^{-1}$ is a distance conversion factor from nt to physical distance for ssDNA. The fitted parameter

494 values are $a = 71.19$, $b = 0.08$, and $K_{D,BHQ} = 145.6 \, mol \cdot L^{-1}$. Using $\Delta G = RT \cdot \ln K_{D,BHQ}$, we have $dG_{37} =$

495 $3 \, kcal/mol$.

Fluorescence normalization

Size normalization

498 To reduce inter-cluster variation in fluorescence measurements, we normalized the Cy3 fluorescence during the melt

499 curve measurements by the total amount of ssDNA in that cluster, measured as Cy3 fluorescence right after

500 hybridization of *fluor_oligo* but before hybridization of *quench_oligo*. The normalizing factor Cy3 signal was clipped

501 to the 1st and 99th percentile of the total distribution.

Temperature dependence and photobleaching normalization

Across the timespan of the experiment, fluorescent signals are dependent on temperature and undergo photobleaching.

To correct for these factors, we further normalized the size-normalized signals of clusters to median signals of control

variants designed to have constant distances between the Cy3 fluorophore and the quench, including 1) "super stable

stem" variants with long GC stems that are not supposed to melt within the experimental temperature range and show

minimum fluorescence signals, and 2) "long repeat" controls that are at least 39 nt long, not supposed to form a hairpin,

and show maximum fluorescence signals. For both control variant groups, only the variants whose median signal

across clusters fall within the 5th and 95th percentile of all median signals at each temperature point within that control

variant group are used. Each variant is allowed to fall outside the percentile range at 1 temperature point at most. This

step filters out the variants not exhibiting the expected behavior. Afterwards, the median values at each temperature

point of both groups are taken and used to normalize the entire dataset. **(Fig. S1d)**

Direct melt curve fitting

514 The melt curve fitting pipeline was adapted from[34]. Briefly, the direct fitting process was carried out in the following

515 steps: 1) Directly fit normalized signals of single clusters; 2) Estimate the distribution of f_{max} and f_{min} from single

516 cluster fit results; 3) Refine the fit at the variant level with bootstrapping. The details are described below. The fitting

517 process is available as a snakemake[35] pipeline at https://github.com/keyuxi/array_analysis.

1) Single cluster fitting

Equation (2) was directly fitted to normalized signals of single clusters with minimal constraints, where T and T_m are temperatures in Kelvin, and $k_B = 0.0019872$ is the Boltzmann constant. For initial values, Tm was set to the temperature point where the normalized signal is closest to 0.5, ΔH to -40, f_{max} to 1, and f_{min} to 0. The python package *lmfit* (v1.0.3)[36] was used for least-squares fitting.

$$f[T] = f_{min} + \frac{f_{max} - f_{min}}{1 + exp(\frac{\Delta H}{k_B} \cdot \left(\frac{1}{T_m} - \frac{1}{T}\right))} \qquad (2)$$

2) Estimating distributions of f_{max} and f_{min}

We used the following empirically gated criteria to decide if a single cluster fit was successful: R^2 of the fit at least 0.5, fitted f_{min} between -1 and 2, the estimated standard error of the best-fitted parameter value $\sigma_{f_{min}}$ less than f_{min} + 1, fitted f_{max} between 0 and 3, $\sigma_{f_{max}}$ less than f_{max}, σ_{Tm} less than 10 K, and $\sigma_{\Delta H}$ less than 100 kcal/mol. A p value is calculated for each variant with a null hypothesis of 25% successful single cluster fits as in[34]. The single clusters are then aggregated to the variant level. Median values of the parameters of each variant were calculated. To select for variants used for estimation of f_{max}, the previously mentioned single cluster criteria were combined with the following extra criteria: 1) be a good fitter, defined as having a success p value less than 0.01; 2) be unfolded at high temperatures, defined as having a fitted f_{max} value of at least 0.5, and $p^{60°C}_{unfolded}$, the portion of unfolded molecules in the cluster calculated from the fitted parameters, of at least 0.975. Similarly, for f_{min} we select for variants with a p value less than 0.01, f_{min} less than 0.05, and $p^{20°C}_{unfolded}$ at most 0.025. We then fitted cluster-per-variant-dependent Gamma distributions to f_{max} and f_{min} as in[34].

3) Fit refinement

For each variant, we separately decided whether to enforce f_{max} and f_{min} distributions to correct for variants that never become fully unfolded or folded during the experiment. Specifically, we calculated the lower bound for f_{max} as $\mu_{f_{max}} - 10 \cdot \sigma_{f_{max}}$, where $\mu_{f_{max}}$ and $\sigma_{f_{max}}$ are the mean and standard deviation calculated from the Gamma distribution with the according number of clusters per variant from the previous step, and the upper bound for f_{min} as $\mu_{f_{min}} + \sigma_{f_{min}}$. Here, we set the margin for f_{max} larger to enforce global f_{max} less, as f_{max} is more variable between variants than f_{min} because of sequence-dependent effects. If the median signal at the highest temperature point is less than the lower bound for f_{max}, we enforce f_{max}; likewise, if the median signal at the lowest temperature point is greater than the upper bound for f_{min}, we enforce f_{min}.

We fitted each variant by bootstrapping the clusters within that variant. At each resampling step, we draw a f_{max} or f_{min} value from the global distribution if enforced or keep f_{max} or f_{min} floating otherwise, and fit **equation (2)** to the median signal of the drawn clusters, using the single cluster fit results as the initial parameters. Then, we calculate ΔG_{37} and ΔS for each resampling step:

<div align="right">549</div>

$$\Delta G[T] = \Delta H - T \cdot \Delta S = \Delta H \cdot \left(1 - \frac{T}{T_m}\right) \qquad (3)$$

<div align="right">550</div>

$$\Delta S = \frac{\Delta H}{Tm} \qquad (4)$$

where T is temperature in Kelvin. We repeated the process 100 times and obtained a 95% confidence interval and
standard deviation for each parameter, ΔH, Tm, ΔG$_{37}$ and ΔS.

ΔG Line fitting

We also fitted the data with an alternative line fitting to obtain heuristics for two-state behavior. We first re-normalize
the signal $f[T]$, previously normalized to initial fluorescence after hybridization as aforementioned, with fitted f_{max}
and f_{min}. We clip $f_{norm}[T] = (f[T] - f_{min})/(f_{max} - f_{min})$ between 0 and 1 with an $\epsilon = 0.01$ to avoid problems in
the next step. We transformed $f_{norm}[T]$ to $\Delta G[T]$ with

<div align="right">558</div>

$$\Delta G[T] = -k_B T ln\left(\frac{1}{f_{norm}[T]} - 1\right) \qquad (5)$$

559 Error propagation was calculated with the uncertainties python library. Finally, we fitted with a line with free
560 parameters ΔH and Tm:

<div align="right">561</div>

$$\Delta G[T] = \Delta H \cdot \left(1 - \frac{T}{T_m}\right) \qquad (6)$$

If the variant exhibits two-state behavior, $\Delta G[T]$ should fit well to a linear line. We first fit with RANSAC[37] to identify
outliers, then re-fit the inliers with ordinary least squares linear regression for the final parameters.

Filtering and combining data replicates

Each variant measured and fitted to curves in a replicate experiment was first filtered by the criteria: at least measured
in 5 clusters, bootstrapped error in ΔG$_{37}$ is less than 2 kcal/mol, error in Tm less than 25°C, error in ΔH less than 25
kcal/mol, and curve fitting RMSE less than 0.5. They were then compared to the line fitting results. To be labeled as
passing two-state criteria, the fitted ΔH value of a variant between the 2 methods need to agree within 20%, Tm within
2°C, the reduced χ^2 of the line fit below 1.5, and the number of inlier data points in line fitting greater than 10. The
replicates were then combined as described in[34]. Briefly, for each parameter (ΔH, Tm, ΔG_{37}, or ΔS), its measurements
X_i and bootstrapped error σ_i in each replicate, the combined error is $\sigma = \sqrt{\left(\sum_i \sigma_i^{-2}\right)^{-1}}$ and the combined parameter
$X = \sigma^2 \cdot \sum_i X_i/\sigma_i^2$, where the sum is over the replicates whose data are not missing for the given variant. The missing
values were ignored. We then apply a second filter for two-state behavior. A given variant needs to be labeled as
passing two-state criteria in at least 2 replicates to be kept. We combined 4 such replicate experiments, including 3
melting and 1 cooling curve measurements from 2 different chips (see table for detail). We apply a final filter on the

combined variants to only keep those whose Tm are between 25 and 55°C and fall within the dynamic range of the
system.

UV melting

Designing sequences for UV melting and quality control

We randomly selected hairpin sequences from each category of the Array Melt library for validations in **Fig. 2** and
designed random 16-mers that would fold into a hairpin with the secondary structure ((((((....)))))) for **Fig. S4d**,
filtering those with misfold secondary structures with NUPACK. DNA oligo samples were synthesized by Integrated
DNA Technologies (IDT) and were run through Ion Exchange High Pressure Liquid Chromatography (IE-HPLC)
remotely on Emerald Cloud Lab (ECL) with DNAPac PA200, 9x250 mm Semi-Prep column (Thermo Fisher
Scientific 063421), and the fractions were measured by a UV-Vis detector at 260 nm. Only the oligos with a single
peak were used for data analysis.

Running cloud lab UV melting experiments

UV melting experiments were run remotely on ECL. The Manifold data objects could be found at the data object links
in supplementary table 2. DNA oligos ordered from IDT were shipped to ECL and resuspended in high concentration
stock solutions. For hairpins, the DNA stock solutions are first snap frozen right before each experiment, being placed
at 90°C for 5 min and moved to 4°C for 30 min before diluted into working concentrations; for duplexes, the DNA
stock solutions are first diluted into working concentrations, then placed on 90°C for 5min and left annealing for 30
min to 1 hour. Melt curves are measured at varied concentrations, where the range is 3 to 12 uM for hairpins and 72
to 96 uM for duplexes. Hairpin data used for method validation in **Fig. 2** and model comparison in **Fig. S4d** were all
measured at 6 uM or 9 uM. Samples were measured on a Cary 3500 UV/Vis spectrophotometer, using micro scale
black walled UV quartz cuvettes with a pathlength of 10 mm. Samples were measured for 2 to 3 melt-cooling cycles
with a temperature ramp rate of 0.8°C and 15 min equilibrium time between each melt and cool sweep. Temperatures
were measured with an immersion probe. Data points were acquired every 1 °C at 260 nm from 5°C to 85 or 95 °C.

Analyzing UV melting curves

600 The UV melting curves were fitted to a modified version of **equation (2)** with an extra term to account for baseline
601 slope:

$$f[T] = f_{min} + a \cdot T + \frac{f_{max} - f_{min}}{1 + exp(\frac{\Delta H}{k_B} \cdot \left(\frac{1}{T_m} - \frac{1}{T}\right))} \qquad (7)$$

where f[T] is the absorbance, a is a free parameter for the slope, and the rest the same as in **equation (2)**. For each
individual melt or cooling curve, we first found the temperature where the derivative $\frac{df[T]}{T}$ is maximum, that is, $T_m^{diff} =$
$argmax_T \frac{df[T]}{T}$, as the initial value of T_m. We fixed the value of T_m when fitting **equation (7)** in the first round and

let the parameters a, f_{max}, f_{min}, and ΔH float. Afterwards, we use the best-fitted values from the previous round as

the initial values of the parameters and allow T_m to float in the second round of fitting. This two-step fitting process

empirically improved the fitting RMSE. We selected the curves with an RMSE less than 0.015, $\sigma_{\Delta H} < 10$ And $\sigma_{T_m} <$

5 and aggregated the fitted ΔH and Tm values of the melt and cooling curves by each sample tube with $X = \sigma^2 \cdot$

$\sum_i X_i/\sigma_i^2$ and $\sigma = \sqrt{\left(\sum_i \sigma_i^{-2}\right)^{-1}}$ as for the array experiments. We then calculated ΔG_{37} and ΔS from the aggregated

ΔH and T_m using equations (3) and (4), and used the uncertainty propagation equations $\sigma_{\Delta S} = -\Delta S \cdot$

$\sqrt{\left(\frac{\sigma_{\Delta H}}{\Delta H}\right)^2 + \left(\frac{\sigma_{T_m}}{T_m}\right)^2}$ and $\sigma_{\Delta G_T} = \sqrt{\sigma_{\Delta H}^2 + (T \cdot \sigma_{\Delta S})^2}$ to calculate the measurement uncertainties.

Salt correction

614   All salt corrections used the equation from[38]:

$$\frac{1}{T_m(2)} = \left(\frac{1}{T_m(1)} + (4.29 \cdot f(GC) - 3.95) \times 10^{-5} \cdot ln\frac{[Na^+]_2}{[Na^+]_1} + 9.4 \times 10^{-6}(ln^2[Na^+]_2 - ln^2[Na^+]_1) \quad (8)$$

616   where $T_m(1)$ and $T_m(2)$ are the melting temperatures before and after salt correction in Kelvin, $[Na^+]_1$ and $[Na^+]_2$

617   the corresponding sodium concentrations, and $f(GC)$ the fraction of GC in the sequence.

Variance analysis

Mismatch ΔΔG37 calculation

620   For each mismatch, we find its parent sequence variant, which is a hairpin with Watson-Crick pairing stem and can

621   derive into the mismatch by mutating a single nucleotide. The mismatches are categorized by the identities of those

622   nucleotide substitutions. We then calculated $\Delta\Delta G_{37}$ of mismatches by subtracting ΔG_{37} of the parent variant from that

623   of the mismatch variant. C>G single mismatches were excluded from the variance calculations because of insufficient

624   number of data points (only 2 variants).

Expected biological variance and model explained or introduced variance

For a dataset of n sequence variants with n corresponding $d\Delta$ values $y_1, \ldots, y_n$, measurement uncertainty values

$\sigma_1, \ldots, \sigma_n$, and nearest-neighbor model prediction values $\hat{y}_1, \ldots, \hat{y}_n$, we have total variance $\sigma^2_{tot} = \sum_{i=1}^n (y_i - \bar{y})^2/n$

and technical variance $\sigma^2_{tech} = \sum_{i=1}^n \sigma_i^2/n$. The expected biological variance $\sigma^2_{bio} = \sigma^2_{tot} - \sigma^2_{tech}$. Model

variance is the variance of prediction values $\sigma^2_{model} = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2/n$, and residual is $\sigma^2_{res} = \sum_{i=1}^n (y_i - \hat{y}_i)^2/n$.

From these we calculate variance explained $\sigma^2_{explained} = \sigma^2_{tot} - \sigma^2_{res}$. In the case where $\sigma^2_{res} > \sigma^2_{tot}$, we have

variance added by model $\sigma^2_{added} = -\sigma^2_{explained} = \sigma^2_{res} - \sigma^2_{tot}$.

Nucleotide importance score

633   We calculated nucleotide importance score (NIS) using $\Delta\Delta G_{37}$ for mismatches, and ΔG_{37} for triloop, tetraloop, and

634   bulges. For a dataset of n sequence variants with n corresponding energy values $y_1, \ldots, y_n$, either ΔG_{37} or $\Delta\Delta G_{37}$, we

635   group the n data points into four groups based on the identity of the nucleotide of interest. For each of the four groups,

636   we calculated the mean, $\overline{y_A}$, $\overline{y_T}$, $\overline{y_C}$, and $\overline{y_G}$. The grand mean $\overline{y_{tot}}$ is the mean of $y_1, \dots, y_n$. The NIS is simply the

637   between group sum of squares (SSB):

638
$$NIS = \sum_{i=\{A,T,C,G\}} (\overline{y_i} - \overline{y_{tot}})^2 \qquad (9)$$

639   *Pairwise interaction score*

640   For a pair of nucleotides of interest, M and N, we compare two linear models that predict the energy value Y, model

641   A with independence assumption and model B without. Model A is in the form of $M + N \sim Y$, and model B $MN \sim Y$.

642   The pairwise interaction score (PIS) is the difference of R^2 between the two linear models

643
$$PIS = R^2{}_{MN \sim Y} - R^2{}_{M+N \sim Y} \qquad (10)$$

644   **Calculations with NUPACK**

645   *Melting temperature*

646   Melting temperatures of DNA duplexes were calculated from the fraction of molecules unpaired. We first calculated

647   structure free energies of the paired state at temperature T, $\Delta G_{paired,T}$, then got the equilibrium constant

648
$$K = \frac{[dsDNA]}{[ssDNA]^2} = exp\left(-\beta \cdot \Delta G_{paired,T}\right) \qquad (11)$$

where $\beta = 1/RT$ is the thermodynamic beta, $[dsDNA]$ is the concentration of double-strand DNA, and [ssDNA] that

of the single-strand DNA. At melting temperature, $[dsDNA] = [ssDNA] = [DNA]/2$ for distinguishable strands and

$[dsDNA] = [DNA]/4$ for indistinguishable strands, where $[DNA]$ is the total concentration of DNA strand species.

This gives $K = \frac{2}{[DNA]}$ or $K = \frac{1}{[DNA]}$ for distinguishable and indistinguishable strands respectively. Solving this equation

with bisection gives us the melting temperature.

For DNA hairpins, melting temperature is independent of DNA strand concentration. The Boltzmann factor becomes:

655
$$e^{-\beta E} = exp\left(-\beta \cdot \Delta G_{folded}\right) \qquad (12)$$

656   T_m is then simply the temperature when $dG_{folded} = 0$. Therefore, we get structure free energies ΔG_1 and ΔG_2 at a

657   lower temperature T_1 and a higher one T_2, then calculate $\Delta S = -(\Delta G_2 - \Delta G_1)/(T_2 - T_1)$, $\Delta H = \Delta G_1 + T_1 \cdot \Delta S$, and

658    $T_m = \Delta H / \Delta S$.

659
$$T_m = \Delta H / \left(\frac{\Delta H - \Delta G_{37}}{T_{37}} - R \cdot \ln K\right) \qquad (13)$$

Data splitting for machine learning models

After filtering with the two-state criteria as described in "ΔG line fitting", we splitted the clean Array Melt into train, validation, and test sets, stratified by variant type. We first splitted 5% of the data as the test set (1438 variants), then splitted 5% of the remaining data as the validation set (1315 variants), and 95% of the rest as the training set (24979 variants). We splitted the external 384-duplex dataset and Oliveira et al. dataset to a 1:1 split of validation and test data. The cloud lab UV melting data were used for correction between Array Melt and bulk solution and were not used for model testing to prevent circularity.

Linear models

Weighted linear regression model

We used a singular value decomposition (SVD) solver to solve the linear regression solutions, taking measurement error of each sequence into account. For a training dataset of n sequence variants with n corresponding ΔH or ΔG_{37} values $y_1, \dots, y_n$, and measurement uncertainty values $\vec{\sigma} = [\sigma_1, \dots, \sigma_n]$, we have a n by $n_{feature}$ feature count matrix X derived from the sequences. We then did SVD on matrix $A = X/\vec{\sigma}$ and get U, S, V. Any diagonal value in S that is smaller than a relative numeric threshold 10^{-15} was set to zero. Then the estimated parameter values for the features are $V^T \cdot S_{inv} \cdot U^T \cdot \vec{b}$, and the estimated errors for the features are $\sqrt{\sum V^T \cdot S^2_{inv}}$, where $\vec{b} = \vec{y}/\vec{\sigma}$, S_{inv} is S but taking $\frac{1}{s}$ for all diagonal elements, and S^2_{inv} is S but taking $\frac{1}{s^2}$ for all elements.

The NUPACK-compatible nearest-neighbor model

We trained the improved nearest-neighbor model on the Array Melt training data set, containing 24,971 of the 27,732 sequences that passed quality control and two-state criteria, and tuned the model on a validation set of 1,315 sequences. We first adjusted salt concentration to 1M Na⁺ to be compatible with NUPACK and other softwares, and then corrected systematic errors by linearly shifting Tm and ΔG, such that the medians of each construct type match those of NUPACK predictions. The underlying assumption is that NUPACK predictions have minimal bias but could be improved by explaining more variance. Improved parameters for NUPACK were fitted with weighted linear regression. We allowed an extra intercept parameter to adjust mean absolute error. From sequence-structure pairs in the training set, we extracted features of stacks and loops. Stacks include nearest-neighbor Watson-Crick pairs and terminal stack penalty, as in NUPACK; loops include the following:

1) Hairpin loops, defined as a subsequence $\phi_{i:j}$ where $i \cdot j$ is the closing base pair. We extracted "hairpin_mismatch" features as in NUPACK, $\Delta G^{hairpinmm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}}$, and introduced an intermediate "hairpin_loop_mid" feature, $\Delta G^{hairpinmid}_{\phi_{[i+2:j-2]}}$ to account for sequence variations in the middle of hairpin loops for triloops and tetraloops. The final energy of hairpin is $\Delta G^{hairpin}_{\phi_{[i:j]}} = \Delta G^{hairpinsize}_n + \Delta G^{hairpinseq}_{\phi_{[i:j]}}$ as in[20], where

$$\Delta G^{hairpinseq}_{\phi_{[i:j]}} = \Delta G^{hairpinmm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}} + \Delta G^{hairpinmid}_{\phi_{[i+2:j-2]}} \left(+\Delta G^{terminalbp}_{\phi_j,\phi_i} \, if \, triloop \right) \qquad (14)$$

692   2) Interior loops, defined from 5' to 3' as two subsequences $\phi_{[i:d]}$ and $\phi_{[e:j]}$. Any other interior loop types were

693   defined as in[20], apart from the $\Delta G^{interiormm}_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}}$, which were fitted separately instead of shared with terminal

694   mismatches as in NUPACK, and that we introduced an intermediate parameter for the interaction between

695   the two base pairs surrounding 1x1 mismatches $\Delta G^{interiormmstacks}_{\phi_i,\phi_j,\phi_{i+2},\phi_{j-2}}$.

This results in 178 features for ΔH and ΔG each. Parameters for hairpin and interior loop sizes were taken from

NUPACK and clamped during fitting. Lookup tables of $\Delta G^{hairpinseq}_{\phi_{[i:j]}}$ and $\Delta G^{interiorseq}_{\phi_{[i:d]},\phi_{[e:j]}}$ for "interior_1_1",

"interior_1_2", and "interior_2_2" were calculated from the fitted parameters. All updated parameters were then

written to a NUPACK-compatible json file.

Afterwards, we obtained parameters for $\Delta G^{hairpinseq}_{\phi_{[i:j]}}$ for tetraloops and triloops directly from the '2nd flanking'

construct types of either tetraloop or triloop parts of the library (**Fig. 1c**). These variants covered all possible sequences

for hairpin loops of length 3 or 4, as well as the nearest-neighbor and next-nearest-neighbor closing pair sequences.

We first regressed out the next-nearest-neighbor closing pair of the loops from measured ΔH and ΔG values, then

aggregated the mean values for variants with identical hairpin loop and nearest-neighbor closing pair sequences. These

aggregated values have the correct ΔΔH and ΔΔG relationships between each other but need a constant to be added to

get the ΔH and ΔG values needed for the nearest-neighbor model. To solve this problem, we fitted two correction

values $\Delta\Delta H$ and $\Delta\Delta G$ in kcal/mol. We searched for values that minimize the equation

$f(\Delta\Delta H, \Delta\Delta G) = w_{Tm} \cdot RMSE(Tm) + w_{\Delta H} \cdot RMSE(\Delta H) + w_{\Delta G} \cdot RMSE(\Delta G) + \alpha \cdot (w_{\Delta H} \cdot \Delta\Delta H^2 + w_{\Delta G} \cdot \Delta\Delta G^2)$

(15)

where $w_{Tm} = 4, w_{\Delta H} = 0.3, w_{\Delta G} = 3.4$ and $\alpha = 0.0001$ are weights determined by validation on 100 sampled

variants in the training set. Calculating this equation is computationally expensive, as it involves directly calling

NUPACK in each iteration. We minimized it on 25 randomly sampled variants from the training set, repeating the

process 4 times on different subsets using the BFGS algorithm implemented in the 'scipy.optimize.minimize' function,

with parameters tol=1e-1 and maxiter=20. The resulting $\Delta\Delta H$ and $\Delta\Delta G$ values were mostly consistent between the

runs (**Fig. S4g**). We took the median values from the 4 runs ($\Delta\Delta H = -28 \; kcal/mol$, $\Delta\Delta G = -4 \; kcal/mol$) and

added them to the aggregated $\Delta\Delta H$ and $\Delta\Delta G$ values of hairpin loops.

Empirically, we found that model performance on the validation set was improved when we augmented the compact

178-parameter nearest-neighbor model with these curated hairpin loop parameters. This improvement might be due

to the dominantly overrepresentation of 'GAAA' hairpin loops, which led to highly skewed data distribution and

caused directly regressed hairpin loop parameters to be less accurate.

The richer parameter nearest-neighbor model

We trained the richer parameter model on the same Array Melt training data set as for the NUPACK-compatible

nearest-neighbor model. For feature extraction, we extracted nearest-neighbor Watson-Crick pairs and terminal stack

penalty as in NUPACK, and the following parameters for loops:

1)   Hairpin loops. For triloops and tetraloops, $\Delta G_{\phi_{[i:j]}}^{hairpinseq} = \Delta G_{\phi_{j-1},\phi_j,\phi_i,\phi_{i+1}}^{hairpinmm} + \Delta G_{\phi_{[i+1:j-1]}}^{hairpinloop}$, where

$\Delta G_{\phi_{[i+1:j-1]}}^{hairpinloop}$ is the full sequence of the hairpin loop, in contrast to only the middle of loops in the NUPACK-

compatible model. For hairpin loops longer than tetraloop, we applied a single loop sequence parameter

$\Delta G_{long}^{hairpinloop}$. The final energy of hairpin is also $\Delta G_{\phi_{[i:j]}}^{hairpin} = \Delta G_n^{hairpinsize} + \Delta G_{\phi_{[i:j]}}^{hairpinseq}$.

2)   Interior loops. For single and double bulges or single mismatches, the sequence-dependent parameter is fitted

for each sequence, $\Delta G_{\phi_{[i:d]},\phi_{[e:j]}}^{interiorseq}$. For double and triple mismatches, $\Delta G_{\phi_{[i:d]},\phi_{[e:j]}}^{interiorseq} = \Delta G_{\phi_i,\phi_j,\phi_{i+2},\phi_{j-2}}^{interiormmstacks} +$

$\Delta G_{\phi_{i+1},\phi_{d-1},\phi_{e+1},\phi_{j-1}}^{interiormm}$.

Deep learning models

Closest sequence lookup baseline model

We defined the distance between any two sequence-structure pairs by summing up the edit distance between the

sequences and the dot-bracket secondary structure representation. For every query sequence-structure pair, we find

the k closest sequence-structure pairs in the training set, whose distance-weighted sum becomes the predicted value

for the query using python package *scipy*[39]. We found the best value for k using the held-out Array Melt validation

data.

Graph neural network model

We implemented the graph neural network using python packages PyTorch and PyTorch Geometric[40]. The final model

has 4 graph transformer layers with a 125-vec node attribute at each layer, stacked with dropout of 0.01 at each layer,

a Set2Set module with 10 processing steps, and a 1 fully connected hidden layer with 128 channels and a dropout ratio

of 0.49. We normalized ΔH and Tm of the training data to fall within 0 and 1 and used the sum of the root mean square

error (RMSE) of both ΔH and Tm as the loss function and trained with Adam optimizer for 200 epoches. The model

has 287,136 parameters. The validation model registry was monitored and stored on weight and biases (wandb.ai).

Using the NUPACK-compatible model

The parameter file for the NUPACK-compatible nearest-neighbor model is available in the supplementary material

as 'NUPACK_compatible_NN_model.json'. To use this file, download and point to it when using the python

package NUPACK 4[41]:

my_model = nupack.Model(material='/path/to/parameter/file/NUPACK_compatible_NN_model.json', ...)

Code availability

Code for array data preprocessing from raw sequencing and imaging data plus curve fitting could be found at

https://github.com/GreenleafLab/array_analysis.

All subsequent analysis could be found at https://github.com/GreenleafLab/nnn_paper.

References

1. Kunkel, T. A. DNA Replication Fidelity. *J. Biol. Chem.* **279**, 16895–16898 (2004).

2. Rajski, S. R., Jackson, B. A. & Barton, J. K. DNA repair: models for damage and mismatch recognition. *Mutat.*

Res. Mol. Mech. Mutagen. **447**, 49–72 (2000).

3. Untergasser, A. *et al.* Primer3—new capabilities and interfaces. *Nucleic Acids Res.* **40**, e115 (2012).

4. Kibbe, W. A. OligoCalc: an online oligonucleotide properties calculator. *Nucleic Acids Res.* **35**, W43–W46 (2007).

5. Labun, K. *et al.* CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic Acids Res.* **47**, W171–W174 (2019).

6. Majikes, J. M. *et al.* Revealing thermodynamics of DNA origami folding via affine transformations. *Nucleic Acids Res.* **48**, 5268–5280 (2020).

7. Zadeh, J. N. *et al.* NUPACK: Analysis and design of nucleic acid systems. *J. Comput. Chem.* **32**, 170–173 (2011).

8. SantaLucia, J. & Hicks, D. The Thermodynamics of DNA Structural Motifs. *Annu. Rev. Biophys. Biomol. Struct.* **33**, 415–440 (2004).

9. SantaLucia, J. A unified view of polymer, dumbbell, and oligonucleotide DNA nearest-neighbor thermodynamics. *Proc. Natl. Acad. Sci.* **95**, 1460–1465 (1998).

10. SantaLucia, John, Allawi, H. T. & Seneviratne, P. A. Improved Nearest-Neighbor Parameters for Predicting DNA Duplex Stability. *Biochemistry* **35**, 3555–3562 (1996).

11. Zuker, M. On Finding All Suboptimal Foldings of an RNA Molecule. *Science* **244**, 48–52 (1989).

12. Schroeder, S. J. & Turner, D. H. Optical Melting Measurements of Nucleic Acid Thermodynamics. *Methods Enzymol.* **468**, 371–387 (2009).

13. Spink, C. H. Differential Scanning Calorimetry. in *Methods in Cell Biology* vol. 84 115–141 (Academic Press, 2008).

14. Hadiwikarta, W. W., Walter, J.-C., Hooyberghs, J. & Carlon, E. Probing hybridization parameters from microarray experiments: nearest-neighbor model and beyond. *Nucleic Acids Res.* **40**, e138 (2012).

15. Oliveira, L. M., Long, A. S., Brown, T., Fox, K. R. & Weber, G. Melting temperature measurement and mesoscopic evaluation of single, double and triple DNA mismatches. *Chem. Sci.* **11**, 8273–8287 (2020).

16. Bae, J. H., Fang, J. Z. & Zhang, D. Y. High-throughput methods for measuring DNA thermodynamics. *Nucleic Acids Res.* **48**, e89–e89 (2020).

17. You, Y., Tataurov, A. V. & Owczarzy, R. Measuring thermodynamic details of DNA hybridization using

fluorescence. *Biopolymers* **95**, 472–486 (2011).

18. Buenrostro, J. D. *et al.* Quantitative analysis of RNA-protein interactions on a massively parallel array reveals

biophysical and evolutionary landscapes. *Nat. Biotechnol.* **32**, 562–568 (2014).

19. Marklund, E., Ke, Y. & Greenleaf, W. J. High-throughput biochemistry in RNA sequence space: predicting

structure and function. *Nat. Rev. Genet.* **24**, 401–414 (2023).

20. Fornace, M. E., Porubsky, N. J. & Pierce, N. A. A Unified Dynamic Programming Framework for the Analysis

of Interacting Nucleic Acid Strands: Enhanced Models, Scalability, and Speed. *ACS Synth. Biol.* **9**, 2665–2678

(2020).

21. Tikhomirova, A., Beletskaya, I. V. & Chalikian, T. V. Stability of DNA Duplexes Containing GG, CC, AA, and

TT Mismatches. *Biochemistry* **45**, 10563–10571 (2006).

22. Khandelwal, G. & Bhyravabhotla, J. A Phenomenological Model for Predicting Melting Temperatures of DNA

Sequences. *PLOS ONE* **5**, e12433 (2010).

23. Yan, Z., Hamilton, W. L. & Blanchette, M. Graph neural representational learning of RNA secondary structures

for predicting RNA-protein interactions. *Bioinformatics* **36**, i276–i284 (2020).

24. Zhao, Q. *et al.* Review of machine learning methods for RNA secondary structure prediction. *PLOS Comput.*

Biol. **17**, e1009291 (2021).

25. Shi, Y. *et al.* Masked Label Prediction: Unified Message Passing Model for Semi-Supervised Classification.

Preprint at https://doi.org/10.48550/arXiv.2009.03509 (2021).

26. Veličković, P. *et al.* Graph Attention Networks. Preprint at https://doi.org/10.48550/arXiv.1710.10903 (2018).

27. Vinyals, O., Bengio, S. & Kudlur, M. Order Matters: Sequence to sequence for sets. Preprint at

http://arxiv.org/abs/1511.06391 (2016).

28. Spasic, A. *et al.* Improving RNA nearest neighbor parameters for helices by going beyond the two-state model.

Nucleic Acids Res. **46**, 4883–4892 (2018).

29. Boiko, D. A., MacKnight, R., Kline, B. & Gomes, G. Autonomous chemical research with large language

models. *Nature* **624**, 570–578 (2023).

30. She, R. *et al.* Comprehensive and quantitative mapping of RNA–protein interactions across a transcribed

eukaryotic genome. *Proc. Natl. Acad. Sci.* **114**, 3619–3624 (2017).

31. Good, N. E. *et al.* Hydrogen Ion Buffers for Biological Research*. *Biochemistry* **5**, 467–477 (1966).

32. Magoč, T. & Salzberg, S. L. FLASH: fast length adjustment of short reads to improve genome assemblies.

Bioinformatics **27**, 2957–2963 (2011).

33. Altschul, S. F. & Gish, W. [27] Local alignment statistics. in *Methods in Enzymology* vol. 266 460–480

(Academic Press, 1996).

34. Denny, S. K. *et al.* High-Throughput Investigation of Diverse Junction Elements in RNA Tertiary Folding. *Cell*

174, 377-390.e20 (2018).

35. Köster, J. & Rahmann, S. Snakemake—a scalable bioinformatics workflow engine. *Bioinformatics* **28**, 2520–

2522 (2012).

36. Newville, M., Stensitzki, T., Allen, D. B. & Ingargiola, A. LMFIT: Non-Linear Least-Square Minimization and

Curve-Fitting for Python. Zenodo https://doi.org/10.5281/zenodo.11813 (2014).

37. Fischler, M. A. & Bolles, R. C. Random sample consensus: a paradigm for model fitting with applications to

image analysis and automated cartography. *Commun. ACM* **24**, 381–395 (1981).

38. Owczarzy, R. *et al.* Effects of Sodium Ions on DNA Duplex Oligomers: Improved Predictions of Melting

Temperatures. *Biochemistry* **43**, 3537–3554 (2004).

39. Virtanen, P. *et al.* SciPy 1.0: fundamental algorithms for scientific computing in Python. *Nat. Methods* **17**, 261–

272 (2020).

40. Fey, M. & Lenssen, J. E. Fast Graph Representation Learning with PyTorch Geometric. Preprint at

https://doi.org/10.48550/arXiv.1903.02428 (2019).

41. Fornace, M. E. *et al. NUPACK: Analysis and Design of Nucleic Acid Structures, Devices, and Systems.*

https://chemrxiv.org/engage/chemrxiv/article-details/636c7089b588507d0045f283 (2022)

doi:10.26434/chemrxiv-2022-xv98l.

Supplementary Figures

SUPP FIGURE 1

Figure S1

a.   Number of measurements in a single Array Melt experiment.

b.   Pearson's R of ΔH, Tm, and ΔG_{37} between replicates.

c.   Controls used to normalize fluorescence across different temperatures, including always unfolded constructs

for f_{max} (maximum fluorescence) and folded constructs for f_{min} (minimum fluorescence). Some Watson-Crick

curves are randomly selected to visualize after the normalization, colored by their expected ΔG_{37} values

predicted by the nearest-neighbor model.

d.   Estimation of f_{max} and f_{min} after the initial round of curve fitting. Of all the fitted variants (blue), only a

fraction that passed quality control filters (orange) were used to estimate f_{max} and f_{min} parameters used for the

next round of refinement fitting. Uncertainties of f_{max} and f_{min} were corrected as a function of number of

clusters per variant by curve fitting (right).

SUPP FIGURE 2

849 **Figure S2**

850   a.   Schematic representation of Emerald Cloud Lab (ECL) workflow. Image from ECL with permission.

851   b.   UV melting measured Tm at different DNA hairpin oligo concentrations. Oligo used is WC68.

852   c.   Direct comparison of measured ΔG, ΔH and Tm between nearest-neighbor model prediction and UV melting

853   data. Each datapoint is one hairpin variant.

SUPP FIGURE 3

SUPP FIGURE 3

Figure S3

a. R^2 values between Array Melt data and nearest-neighbor model predictions for all single mismatches, and
for each mismatch type.
b. Relationship of biological variance to total variance in Array Melt data. Each dot is a group of variants with
identical mismatch type and adjacent nearest-neighbor base pairs as in **Fig. 3a**, colored by mismatched type.
The biological/total variance ratio is shown on the right.
c. Examples of when the nearest-neighbor model succeeds or fails to capture total variance in data. Each subplot
is calculated from a group of variants with identical mismatch type and adjacent base pairs. Green is the
histogram of Array Melt measurements, and purple is what is expected from the nearest-neighbor model
prediction with measurement uncertainty.
d. Comparison of nucleotide importance scores calculated from Array Melt data and nearest-neighbor model
predictions for Watson-Crick and single mismatch variants.
e. Comparison of nucleotide importance scores calculated from either $\Delta\Delta G37$ or ΔG_{37}, and on Array Melt data
or nearest-neighbor model predictions, for tetraloops and triloops.
f. Comparison of nucleotide importance scores calculated from Array Melt data and nearest-neighbor model
predictions of bulge variants.
g. Pairwise interaction scores calculated from nearest-neighbor model predictions for Watson-Crick and single
mismatch variants.
h. Pairwise interaction scores as a function of distance between the two nucleotides in the pair, for Watson-
Crick variants or single mismatch variants. Those for the latter are splitted by the distance from the anchor
point to the single mismatch.
i. Difference between pairwise interaction scores from Array Melt data and nearest-neighbor model prediction.
j. Comparison of nucleotide importance scores calculated from Array Melt data and nearest-neighbor model
predictions of triloop variants.

SUPP FIGURE 4

Figure S4

    a. Schematics of the NUPACK-compatible nearest-neighbor model.

    b. Difference between the new and original *dna04* single mismatch parameters.

    c. Comparison of two-state and ensemble model predictions from NUPACK against Array Melt measurements.

    d. Comparison of *dna04*, NUPACK-compatible, and richer parameterization nearest-neighbor model predictions with cloud lab UV melting measurements of randomly designed DNA hairpins.

    e. Performance of the original, NUPACK-compatible, and richer parameter nearest-neighbor models on literature UV melting data used to derive the original nearest-neighbor model.

    f. Performance of the original, NUPACK-compatible, and richer parameter nearest-neighbor model on Array Melt test set for Tm.

    g. Fitted ΔΔH and ΔΔG values for 'hairpin_tetraloop' and 'hairpin_triloop'. Each dot is from one individual optimization run.

SUPP FIGURE 5

Figure S5

a. Mean absolute error of GNNs with either Graph Attention Network or Graph Transformer Network architecture at the graph convolutional layer.

b. Mean absolute error of enthalpy on Array Melt data as a function of the number of graph convolution layers in the GNN.

c. Mean absolute error of melting temperature on Array Melt data as a function of percentage of training data used to fit the GNN.

d. Schematic of the baseline closest sequence lookup model.

e. GNN predicted ΔG_{37} vs. held out Array Melt, splitted by type of variant.

f. The objective function of GNN (RMSE of normalized ΔH and Tm), mean absolute error of Tm, and Pearson's R of Tm across the GNN training process.

g. UMAP of learned embeddings of test data, colored by their predicted Tm.

h. Benchmark of all models on the orthogonal Oliveira et al. dataset as in **Fig. 5** but plotted by model. Bars for each model are for no mismatch, one mismatch, two mismatches and three mismatches, respectively.

Supplementary Notes

Adjusted mean absolute error

Nearest-neighbor models inherently have parameterization degeneracy, meaning that adding a constant to one group of parameters and subtracting from another could lead to the same prediction. This degeneracy can result in a prediction bias, but its toll on prediction accuracy could be mitigated by applying a constant offset to align the predictions with the true values. This constant offset preserves the relative $\Delta\Delta G$ values but not the absolute ΔG values. For instance, in the Array Melt test set, NUPACK with the original parameter set has a ΔG_{37} bias of -0.71 kcal/mol, and our NUPACK-compatible model parameter set has a bias of -0.42 kcal/mol. After subtracting this bias to align the predictions with the true values, the MAE of ΔG_{37} was reduced from 1.07 to 0.87 kcal/mol for the original NUPACK model, and from 0.67 to 0.59 kcal/mol for our NUPACK-compatible model (**Table 1**). This adjustment demonstrated that, with the presence of this bias, the accuracy of predicted $\Delta\Delta G$ values between different sequence-structure pairs is generally higher than that of the predicted ΔG values compared to the measured ground truth. Despite this model bias, we argue that the accuracy of nearest-neighbor models is acceptable in practice, particularly for applications that primarily rely on relative $\Delta\Delta G$ values rather than absolute ΔG values. This is because many applications, such as using dynamic programming to find the minimum free energy structure, depend on the relative $\Delta\Delta G$ values between different candidate secondary structures.

Given this context, we find it reasonable to report both the raw and the adjusted MAE model metrics. The adjusted metrics are calculated by first centering the prediction values to match the mean of the true values, thereby providing a more informative estimate of the relative errors for a given model.